Visualization and Morphological Analysis of Individual Oligodendrocytes in the Mouse

White Matter

Yasuyuki Osanai[1,2,5], Sasikarn Looprasertkul[1,3,5], Batpurev Battulga[1], Reiji Yamazaki[1], Nobuhiko

Ohno[1,4].

[1]Department of Anatomy, Division of Histology and Cell Biology, School of Medicine, Jichi

Medical University, Shimotsuke, 329-0498, Japan. [2]Australian Regenerative Medicine Institute,

15 Innovation Walk, Monash University, Clayton, Victoria, 3800, Australia. [3]Section of Electron

Microscopy, Supportive Center for Brain Research, National Institute for Physiological Sciences,

Okazaki, Japan. [4]Division of Ultrastructural Research, National Institute for Physiological

Sciences, Okazaki, Japan. [5]These authors contribute equally.

Correspondence:

Yasuyuki Osanai, yasuyuki.osanai@jichi.ac.jp

Abstract

Myelin formation by oligodendrocytes is essential for the regulation of the conduction velocity

and proper brain function. To ensure accurate information processing in response to experiences

such as sensory stimuli and learning, oligodendrocytes adjust their number and morphology. In

addition, oligodendrocyte morphology changes with senescence and in the presence of

neurodegenerative diseases. Thus, visualizing oligodendrocytes and analyzing their morphology

is crucial for understanding how our brains change under such conditions. Herein, we describe

the methods for labeling and analyzing the morphologies of individual oligodendrocytes in mouse

white matter at the light microscopic level. PDGFRa-CreERT2:Tau-mGFP and PLP-

CreERT2:Tau-mGFP mice enable us to visualize and analyze later-born or early-born

oligodendrocyte morphology. In addition, sparse oligodendrocyte labeling with attenuated rabies

virus expressing GFP enables the visualization and morphological analysis of individual

oligodendrocytes in various brain white matter regions without the need for transgenic animals.

Furthermore, the combination with immunostaining in thick tissues enables the identification of

labeled oligodendrocytes and myelin sheaths, as well as their interactions with neuronal axons.

These methods are suitable for revealing how oligodendrocytes adapt their morphologies

depending on environmental stimuli or pathological conditions.

Keywords

oligodendrocyte, myelination, rabies virus, tissue clearing

Running head

Visualization and analysis of a single oligodendrocyte

1. Introduction

Oligodendrocytes, the myelin sheath-forming glial cells in the central nervous system, are critical for proper brain functions as they regulate the conduction velocity through myelin sheath formation and saltatory conduction and support neuronal metabolism. Physiological and pathological changes in brain functions are often accompanied by the structural regulation of oligodendrocytes and their myelin, which in turn modulates signal transduction in the neural circuits. Neuronal axons can be divided into multiple segments, including internode covered by myelin sheaths and nodes of Ranvier. The nodes of Ranvier are flanked by paranodes involving paranodal loops and axon-oligodendrocyte junctions [1]. Internodes are characterized by multilamellar compact myelin and inner and outer tongue cytoplasmic channels. Each oligodendrocyte, typically with a small cell body, extends multiple processes, with each connected to the outer tongue [2]. Characteristic oligodendrocyte morphology is developed through the differentiation of oligodendrocyte progenitor cells (OPCs), which express markers such as NG2 and PDGFRa and exhibit bipolar shapes, into mature myelinating oligodendrocytes,

which test positive for markers such as proteolipid protein (PLP) and myelin basic protein, extend

multiple processes, and form myelin sheaths.

Compared to other glial cells such as astrocytes and microglia, visualization of

oligodendrocytes is difficult because there are no reliable anti-bodies that label both their cell

bodies and processes (including myelin). Historically, silver carbonate methods or single-cell

injection of lucifer yellow have been used for visualizing the morphology of a single

oligodendrocyte [3, 4]. However, as silver carbonate staining requires special fixation and single-

cell lucifer yellow injection is technically difficult and makes it hard to obtain high sample

numbers, recent studies have focused on the use of transgenic animals and viral vectors [5-7].

PDGFRa-CreERT2:Tau-mGFP and PLP-CreERT:Tau-mGFP mice are transgenic lines

that label oligodendrocytes differentiated after tamoxifen administration or those generated before

tamoxifen administration, respectively [8, 9]. The PDGFRa-CreERT2 mouse expresses

tamoxifen-inducible Cre in OPCs, and the Tau-mGFP mouse possess the lox-STOP-lox

membrane-targeted GFP (mGFP) cassette downstream of the Tau (MAPT) promoter that is active

in neurons and mature oligodendrocytes. Tamoxifen-induced recombination removes the lox-

STOP-lox cassette from OPCs, and as they differentiate into oligodendrocytes, GFP expression

is driven by the Tau promoter. Thus, in PDGFRa-CreERT2:Tau-mGFP mice, we can label the

oligodendrocytes differentiated after the tamoxifen injection. In contrast, in PLP-CreERT mice,

tamoxifen-inducible Cre is expressed only in mature oligodendrocytes, since the PLP promoter is

active in mature oligodendrocytes but not in OPCs. Therefore, PLP-CreERT:Tau-mGFP mice

enable the labeling of oligodendrocytes differentiated before the tamoxifen injection. Thus, in

PLP-CreERT:Tau-mGFP mice, we can label early-born oligodendrocytes when we administer

tamoxifen in the early developmental period [9]. Therefore, these transgenic animals enable the

visualization of a temporarily distinct oligodendrocyte population.

Rabies virus and Semliki Forest virus vectors have been used to sparsely label

oligodendrocytes in the mouse white matter [10, 11]. We have previously analyzed the

interactions between oligodendrocytes and other cells, such as neurons, using a rabies virus vector

and immunostaining for thick (> 70μm) sections. Using this method, both single oligodendrocytes

and their myelinated axons can be labeled [9]. Thus, it is a powerful tool to analyze

oligodendrocyte-neuron interactions.

In this paper, we introduce methods to visualize and analyze single-cell oligodendrocyte

morphology at the light microscopic level, focusing on two labeling methods employing

transgenic mice and a viral vector. The methods using PDGFRa-CreERT2:Tau-mGFP and PLP-

CreERT:Tau-mGFP have been validated in the corpus callosum and cerebellum [9, 12], and

labeling using the rabies virus vector has been employed in multiple white matter regions,

including the corpus callosum, cerebellum, optic nerve and chiasm, and spinal cord [10]. Methods

utilizing these transgenic animals are potentially applicable to all brain regions, including the gray

matter. The rabies virus-mediated method is considered applicable not only to the white matter of

mice but also to that of other animals. These methods can be useful in examining various

morphological aspects of individual oligodendrocytes, such as the number of myelin sheaths, the

length and fiber diameter of each myelin sheath, height of each oligodendrocyte, and myelination

toward a particular subtype of axons.

2. Materials

2.1 Transgenic mice and reagents

1. PDGFRa-CreERT2 [13] or NG2-CreERT2 mice (Jackson, strain name: B6.Cg-Tg(Cspg4-

cre/Esr1*)BAkik/J, # 008538) [14].

2. PLP-CreERT mice (Jackson, strain name: B6.Cg-Tg(Plp1-cre/ERT)3Pop/J, # 005975) [15].

3. Tau-mGFP mice (Jackson, strain name: B6;129P2-Mapttm2Arbr/J, # 021162) [16].

4. Tamoxifen.

5. Corn oil.

6. Disposable syringes for small-volume injection.

7. 25G short length needles (10–16 mm).

2.2 Viral vector and stereotaxic injection

1. Rabies B19G-SADΔG-GFP (Salk Institute, viral vector core) or Rabies B19G-HepΔG-GFP [17] (*See* Note 1 for information on virus production and purchasing).

2. Apparatus for stereotaxic microinjection (For detailed procedures of mouse brain stereotaxic injection, please consult methodological papers on viral injection, such as Cetin et al., 2006) [18].

2.3 Perfusion and slicing of the brain

1. 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB) (pH 7.4).

2. Vibratome slicer.

3. Perfusion pump.

2.4 Thick tissue immunostaining and tissue clearing with RapiClear

Stock solutions without normal goat serum (NGS) or antibodies can be stored at room temperature (RT; 20–25°C) for over 1 year.

1. Penetration buffer: 2% Triton X-100 and 0.05% sodium azide in phosphate-buffered saline (PBS). Mix 2 g of Triton X-100 and 50 mg of sodium azide in 98 mL of PBS.

2. Blocking buffer: 10% NGS, 1% Triton X-100, 2.5% dimethyl sulfoxide (DMSO), and 0.1%

sodium azide in PBS. Mix 1 g of Triton X-100, and 2.5 g of DMSO in 87 mL of PBS. Add NGS

at one-tenth of the final volume to the solution before use.

3. Antibody dilution buffer: 1% NGS, 0.2% Triton X-100, 2.5% DMSO and 0.1% sodium azide

in PBS. Mix 2 mL of 10% Triton X-100 solution, 2.5 g DMSO, and 100 mg sodium azide in 96

mL of PBS. Add NGS at 1/100 of the final volume of the solution immediately before use.

4. Washing buffer: 3% NaCl and 0.2% Triton X-100 in PBS. Mix 3 g of NaCl, and 2 mL of 10%

Triton X-100 solution in 95 mL of PBS.

5. Antibodies: Mouse monoclonal anti-APC (CC1, 1/100, IgG2b, OP80, Merck) for labeling

mature oligodendrocytes, rabbit polyclonal anti-human Olig2 (1/100, 18953, Immuno-Biological

Laboratories America) or chicken polyclonal anti-Olig2 (1/500, OLIG2-0020, Aves) for labeling

oligodendrocyte lineage cells, rabbit polyclonal anti-Iba1 (1/400, 019-19741, Wako) for labeling

microglia, mouse monoclonal anti-GFAP (1/100, IgG1, G3893, Sigma-Aldrich) or mouse

polyclonal anti-S100B (1/100, ab34686, Abcam) for labeling astrocytes (*see* Note 2), rabbit

polyclonal anti-Caspr (1/500-1000, ab34151, Abcam) for labeling the paranodes of the myelin

sheath, rabbit polyclonal anti-NG2 (1/50, AB5320, Merck) for labeling OPCs, rabbit polyclonal

anti-calbindin (1/500, MSFR100390, Frontier Institute) for labeling Purkinje cells, chicken

polyclonal anti-neurofilament-H (1/500, NB300-217, Novusbio) for labeling neuronal axons, and

rat monoclonal anti-GFP (1/500, IgG2a, 04404-84, Nacalai Tesuque) for visualizing GFP.

6. RapiClear 1.52 (RC152001, SUNJin Lab, Taiwan)

7. Antifadant

3. Methods

3.1 Labeling oligodendrocytes using PDGFRa-CreERT2:Tau-mGFP and PLP-Cre:Tau-

mGFP mice

3.1.1 Tamoxifen administration and fixation

1. PDGFRa-CreERT2 and PLP-CreERT mice are crossed with Tau-mGFP mice to generate

PDGFRa-CreERT2 (either homozygous or heterozygous):Tau-mGFP (heterozygous) and

PLP-CreERT (either homozygous or heterozygous):Tau-mGFP (heterozygous) mice,

respectively.

2. Tamoxifen is dissolved in corn oil for one 1 h to make a tamoxifen stock solution (10 mg/mL).

During this process, the solution is incubated at 60 °C and vortexed every 10 min. Tamoxifen

stock solution can be aliquoted (0.5–1.5 mL) and stored for at least six months at −20°C. The

frozen stock solution is incubated at 60°C for 1 h before use, with gentle mixing by vortex

every 10 min during incubation. The thawed solution can be stored at 4°C for up to one week.

Before injection, the solution is incubated at 60°C for 10 min and vortexed thoroughly.

Ideally, the temperature of the solution be 37°C at the time of injection.

3. Tamoxifen is administered via intraperitoneal injection (50–100mg/kg) for two consecutive

days in mice younger than postnatal day 15 (P15). For mice older than P15, intraperitoneal

administration of 100 mg/kg tamoxifen for four consecutive days is recommended (*see* Notes

3, 4, and 5).

4. Mice are perfused with 4% PFA in 0.1 M PB at the desired postnatal age. Their brains are

then postfixed in the same fixative for 1 day at 4°C. After transferring the brains into PBS,

100 μm-thick brain slices are prepared using a vibratome slicer (*see* Note 6).

3.2 Labeling oligodendrocytes using attenuated rabies virus-GFP

The methods for stereotaxic microinjections are well established; therefore, they are not

described in detail here. For the detailed procedure, please refer to methodological papers on

stereotaxic injection, such as Cetin et al (2006).

1. Dilute the rabies virus stock with PBS to prepare a viral solution containing 5.0×10^3–5.0

$\times 10^4$ infectious units per microliter.

2. Inject 1 μL of the viral solution into the white matter. The following parameters are used

for injections into each brain region. In the adult mouse, for the corpus callosum 1.0 mm

posterior and 0.5–0.8 mm lateral to the bregma, at a depth of 1.0 mm from the pia, for the

181 optic chiasm, 0.2 mm posterior and 0.0 mm lateral to the bregma, at a depth of 5.0–5.3

182 mm from the pia, and for the white matter of cerebellar lobules IV/V, 6.45 mm posterior

183 and 1.00 mm lateral to the bregma, a depth of 0.5 mm.

184 3. Perfuse mice with 4% PFA in 0.1 M PB, 4–7 days after viral injection. The brains are then

185 postfixed in the same fixative for 1 day at 4°C. After transferring the brains into PBS, 100

186 µm-thick brain slices are prepared using a vibratome slicer.

188 **3.3 Immunostaining of thick brain sections using the Rapi clear method to analyze**

189 **oligodendrocyte morphology**

190 1. Select brain sections (50–200 µm) containing the target brain region and place 2–3

191 sections per well into a 24-well plate filled with PBS.

192 2. Incubate brain sections in the penetration buffer at RT for 24 h.

193 3. Incubate the brain sections in the blocking buffer for 1–2 days at 4°C,

194 4. The sections are incubated at 4°C for 3–4 days in the antibody dilution buffer (300-500

195 µL for 2–3 sections in 24 well plate) containing primary antibodies such as rat anti-GFP

196 (1/500), rabbit anti-Caspr (1/500-1000), rabbit anti-human Olig2 (1/100) or chicken anti-

197 Olig2 (1/500), and mouse anti-APC (CC1, 1/100).

198 5. Wash the sections in washing buffer three times for 1 h each at RT. Then, incubate the

specimens in washing buffer on an orbital shaker at 4℃ overnight.

6. Incubate the specimens with secondary antibodies in the antibody dilution buffer on an

orbital shaker at 4℃ for 2 days.

7. Wash the sections in washing buffer three times for 1 h each at RT. Then, keep the

specimens in the washing buffer on an orbital shaker at 4℃ overnight.

8. Stain with Hoechst (1/1500–2000) in PBS for 3 h at RT if needed.

9. If the tissue was stained with Hoechst, wash it with PBS three times for 1 h each on an

orbital shaker at RT. If not, with PBS twice for 10 min each.

10. For tissue clearing, place the specimen on a slide glass, add pre-warmed (37℃) RapiClear

solution with/without an antifade solution. After 20 min, applying the cover glass.

11. Apply the cover glass and seal its edges with a clear nail polish. Allow the slide to dry

overnight at RT (*see* Note 7).

3.4. Confocal microscopic imaging

Use a confocal microscope with Z-series sequential imaging function. A single

oligodendrocyte morphology can be captured by approximately 50–100 optical sections at 0.5

μm intervals, adjusted as needed to fully capture the dorsal-ventral extent of the cell bodies,

processes, and myelin sheaths of individual oligodendrocytes. We utilized a 60× objective lens

(N.A. 1.42) on the Dragonfly 200 high-speed confocal platform to achieve the high–

magnification image. High–resolution images are required for detailed morphological analyses

of individual oligodendrocytes and their myelin internodes.

1. The uppermost and lowermost individual oligodendrocytes and their myelin are set for

image acquisition, with a step size of 0.5 μm. GFP signal loss is used to define the

uppermost and lowermost limits of the confocal Z-stack imaging.

2. Images are captured and exported in the proprietary file format provided by the microscope

manufacturer. The subsequent analysis is performed using the Fiji software, which supports

most proprietary formats, such as .oif, .nd2, .czi, and .ims. These formats preserve z-plane

and scale information, facilitating the analysis of entire individual oligodendrocytes with

Fiji.

3.5. Analysis of oligodendrocyte morphology labeled using PDGFRa-CreERT2:Tau-mGFP

and PLP-Cre:Tau-mGFP mice

The oligodendrocyte in PDGFRa-CreERT2:Tau-mGFP (Fig. 1) and PLP-Cre:Tau-

mGFP mice (Fig. 2) are labeled via tamoxifen-induced Cre recombination at specific time points.

The oligodendrocyte cell membrane and myelin sheath membrane can be labeled by mGFP,

which enables the analysis of the oligodendrocyte's morphology. The age of mice, number of

administrations, and concentration of the tamoxifen are important for the analysis of a single

oligodendrocyte. If oligodendrocytes are densely labeled, it becomes difficult to analyze the

morphology of individual cells. Various morphological aspects can be analyzed in single

oligodendrocytes (*see* Note 8), such as the myelin sheath length, number of myelin sheaths

produced by a single oligodendrocyte [8, 19], fiber diameter of each myelin sheath [20], and

heights of single oligodendrocytes [12].

3.5.1 Measurement of the length and number of myelin sheaths formed by a single

oligodendrocyte

1. The number of sheaths and the myelin sheath length produced by a single oligodendrocyte

are quantified using the Simple Neurite Tracer (SNT) plugin

(https://imagej.net/plugins/snt/) [21], accessed via Fiji via Plugins > Neuroanatomy >

SNT....

2. In the SNT Startup Prompt window, select the directory containing the target file that

includes the z-stacked serial images of the GFP-labeled oligodendrocyte and its myelin

sheaths.

3. Since the images may appear with low intensity, adjust the brightness and contrast as needed

via Fiji > Image > Adjust > Brightness/Contrast….

4. Each individual myelin sheath is manually traced in the z-stacked serial images. The starting

point of each myelin sheath is defined at one tip of the myelin sheath (GFP-positive)

colocalized with the paranode marker (Caspr-positive, Fig. 1g–i), and subsequent points are

selected along the trajectory of the sheath across adjacent optical sections. Press either "Yes"

or the "Y" key to accept the path. For each pair of selected points, the A* search path-

finding algorithm auto-tracing within SNT is used to compute the optimal path based on

image intensity gradients. Each path segment is accepted by pressing the "Y" key. After

completing a full trace of one sheath, the path is finalized by selecting "Finish" or pressing

the "F" key.

5. Each completed myelin sheath tracing is stored as an individual path and can be verified

using SNT Path Manager (Fig. 1k, l).

6. After tracing all the myelin sheaths of a single oligodendrocyte, save the tracing data as

follows: File > Save Tracings… > Save As… (.traces file).

7. To determine the length of each myelin sheath and the total number of sheaths per cell,

export the path statistics via Analysis > Path Properties: Export CSV….

3.5.2 Height/ vertical territory measurement of an individual oligodendrocyte

1. Acquire z-stacked serial images of individual oligodendrocytes at a 0.5 μm step size,

ensuring that all myelin sheaths of the cell are included. Images should be saved in a file

format compatible with the Fiji software.

2. Import the z-stacked serial images into the Fiji software. The imported images display the

entire oligodendrocyte, with all its myelin sheaths.

3. Since the images may appear with low intensity, adjust the brightness and contrast

appropriate via Fiji > Image > Adjust > Brightness/Contrast….

4. Identify the outermost dorsal and ventral myelin processes of the oligodendrocyte using the

z-stack images.

5. Locate the uppermost dorsal and lowermost ventral myelin sheaths by tracing along their full

length and identifying the connection point between each myelin sheath and the

corresponding oligodendrocyte process.

6. Mark these connection points for further analysis.

7. Locate and mark the nucleus of each individual oligodendrocyte in the z-stack images.

8. Compress the confocal z-stack into a single-plane projection for linear distance measurement

via Image > Stacks > Z Project...

9. Measure the distance between the points where the outermost dorsal and ventral myelin

internodes connect to their process by drawing a straight line from the dorsal outermost

myelin sheath to the ventral outermost one. Ensure that the line passes through the nucleus

and is perpendicular to the connection points of the processes (Fig. 1j). This can be done

via Analyze > Measure.

3.5.3 Fiber diameter measurement of a single oligodendrocyte

1. Acquire z-stacked serial images of individual oligodendrocytes at a 0.5 μm step size,

ensuring the inclusion of all myelin sheaths of the cell. Images should be saved in a file

format compatible with the Fiji software.

2. Import the z-stacked serial images into the Fiji software. The imported images display the

entire oligodendrocyte, including all its myelin sheaths.

3. Since the images may appear with low intensity, adjust the brightness and contrast

appropriate via Fiji > Image > Adjust > Brightness/Contrast….

4. Select the myelin sheaths of each oligodendrocyte to measure their diameters. One approach

is to include: (i) the outermost myelin sheath on the dorsal side, (ii) the outermost sheath

on the ventral side, (iii) one internode located in the central region of the oligodendrocyte,

(iv) one internode between the central region and the outermost dorsal sheath, and (v) one

internode between the central region and the outermost ventral sheath, based on z-stack

images.

5. Compress the Z-stacked image to a single plane image. To determine where to measure the

diameter, draw a line parallel to the myelin internode of the selected myelin sheath and

measure the length of that line (Fig. 1m).

6. Since the fiber diameters can vary along the internode, myelin internodes are divided into

several segments for measurement. For example, the internodes are divided into six

segments using the following formula to determine the spacing along their total length:

Distance for measurement position (µm) = (Internode length (µm) − 2 µm) / 5

Place the first and last measurement points at a distance of 1 µm from each end of the

myelin internode because the internode's tip is a paranode area that is always larger than

the internode. The remaining four points are positioned equally between them based on

the calculated distance.

7. After the positions for the measurement are marked (Fig. 1m), draw a line from the upper

part to the lower part of that myelin area and measure its length (Fig. 1n) via Analyze >

Measure.

8. Collect data on the selected positions of each myelin sheath and calculate the average fiber

diameter.

9. Measure the fiber diameter of the selected internodes per oligodendrocyte—approximately

five internodes per oligodendrocyte. Then, calculate the average fiber diameter of each

oligodendrocyte.

3.6 Analysis of selective myelination toward a particular subtype of axons

We have previously demonstrated that oligodendrocytes that differentiated early in life

preferentially myelinate Purkinje cell axons in the cerebellar white matter [9]. By combining the

RapiClear method with oligodendrocyte labeling using transgenic mice or rabies virus, we are

able to investigate the selective myelination toward specific axon subtypes.

1. Label oligodendrocytes using transgenic mice or rabies virus. We labeled early-born

    oligodendrocytes in PLP-CreERT:Tau-mGFP mice by administering tamoxifen at P8 (100

    mg/kg) and perfusing the animals with 4% PFA in 0.1 M PB at P56 (Fig. 2).

2. Prepare 100 μm brain slices using a vibratome.

3. Performing thick-section immunostaining, RapiClear method, as described above, using

    markers for specific neuronal subtypes (e.g., calbindin for Purkinje cell axons) and the

    pan-neuronal axon marker neurofilament. GFP-labeled oligodendrocytes in transgenic

    animals should be enhanced with an anti-GFP antibody, whereas those labeled by rabies

    virus do not require immunostaining.

4. Acquire z-series confocal images of GFP-labeled oligodendrocyte/myelin and multiple

    subtypes of neuronal axons as described above.

5. Count the number of axons of each subtype myelinated by the GFP-labeled sheath using

Fiji. GFP-labeled processes ensheathing axons for more than 20 μm were defined as

myelin sheaths (*see* note 9).

6. Count the number of axons of each subtype not myelinated by GFP-labeled

oligodendrocyte/myelin within the height of the labeled oligodendrocytes (Fig. 2).

7. To determine whether GFP-labeled oligodendrocytes preferentially myelinate specific axon

subtypes, perform Fisher's exact test for individual confocal images using statistical

software (e.g., Prism, GraphPad). Input the following values:

A. Number of specific-subtype axons myelinated by GFP-labeled oligodendrocytes,

B. Number of specific-subtype axons not myelinated,

C. Number of non-specific-subtype axons myelinated,

D. Number of non-specific-subtype axons not myelinated.

Fisher's exact test is then performed to evaluate selective myelination.

3.7 Analysis of single oligodendrocytes morphology labeled by attenuated rabies virus

While the morphological analyses of individual oligodendrocytes in the sections labeled

with the rabies virus are mostly similar to those in PDGFRa-CreERT2:Tau-mGFP and PLP-

CreERT:Tau-mGFP mice, there are some differences. The GFP expression level in cells infected

by attenuated rabies virus encoding GFP (hereafter referred to as RV-GFP) is so strong that

immunostaining for GFP is unnecessary. Since RV-GFP can label cells other than

oligodendrocyte-lineage cells, we first need to check whether the cells labeled with RV-GFP are

oligodendrocytes. If the labeled cells exhibit morphological features of myelinating mature

oligodendrocytes and have GFP-positive ensheathment around neuronal axons, which are mostly

flanked by Caspr immunostaining, these cells are identified as oligodendrocytes.

In the white matter, myelinating mature oligodendrocytes labeled with RV-GFP extend

the processes corresponding to their myelin sheaths parallel to the direction of the neuronal axons.

Because RV-GFP labels the cytosol of myelin sheaths, the ensheathment of the GFP signal around

unstained axons can be clearly observed when the diameters of these axons are large (Fig. 3a, *see*

Note 9). Another distinguishing feature of RV-GFP-labeled oligodendrocytes is the swelling

observed at the paranodal regions (Fig. 3). Since the paranodes contain more cytoplasm than the

internodes, the paranodes appear more intensely labeled and expanded with GFP. The length of

each myelin sheath can be measured by identifying its paranodes as GFP-labeled swellings at

both ends of the sheath.

We can also use antibodies against oligodendrocytes/myelin markers. Notably, cell

body markers for oligodendrocytes (e.g., CC1) tend to be downregulated in RV-GFP (+) cells,

which might be attributed to the high-level GFP expression. To confirm whether GFP-labeled

cells form the myelin sheath, we recommend using an anti-Caspr antibody. If the ends of GFP-

positive processes are closely aligned with Caspr signals, this indicates that the processes

correspond to the myelin sheath flanked by the set of Caspr signals (Fig. 3). These morphological

features, combined with immunostaining, allow for reliable oligodendrocyte identification.

4. Notes

1. Salk Institute viral vector core provides a stock G-deleted rabies aliquot. Rabies virus vector

amplification techniques have been described in several studies [22, 23]. Rabies B19G-SADΔG-

GFP and rabies B19G-HepΔG viral vectors can be obtained from Vector Builder (Chicago, USA).

2. Immunostaining of membrane-located receptors is not always successful for the thick-tissue

immunostaining method. For example, immunostaining for PDGFRa is difficult with this method.

If you want to count the number of OPCs, we recommend quantifying CC1(−)/Olig2 cells [24] or

using immunostaining for NG2 [12]. In our experience, the anti-GFAP antibody exhibits reliable

performance in brain tissue, whereas the anti-S100B antibody lacks astrocyte specificity,

particularly in the corpus callosum. In contrast, in the optic nerve, anti-GFAP antibodies are

ineffective, but the anti-S100B antibody works well. To accurately determine the number of

astrocytes in the optic nerve, double immunostaining with anti-S100B and anti-Olig2 antibodies

is recommended, as both astrocytes and OPCs express S100B.

3. In our experience, 100 mg/kg tamoxifen administration for more than two days increases the

risk of mortality in mice younger than P15. Researchers should first confirm whether the mouse

strain they are using can tolerate two consecutive days of tamoxifen administration. If the

mortality rate is high, the number of tamoxifen administration days or its dosage should be

reduced.

4. In our experience, tamoxifen administration before P9 failed to label oligodendrocytes; instead

it labeled unidentified cells in PDGFRa-CreERT2:Tau-mGFP mice, possibly because PDGFRa

promoter activity is not yet restricted to OPC at such an early developmental stage.

5. Four days of consecutive injection of tamoxifen at 100 mg/kg helps maximize the

recombination rate. If a lower recombination rate is acceptable or if you wish to limit the number

of labeled oligodendrocytes, consider reducing the tamoxifen concentration or the number of

injection days. In PLP-CreERT:Tau-mGFP mice, oligodendrocytes in the white matter tend to be

densely labeled [9]. Therefore, to achieve sparse labeling and enable observation of single-cell

morphology, the tamoxifen concentration must be substantially reduced.

6. Brain slices can be stored for up to six months at 4°C in PBS containing 0.1% sodium azide or

stored for up to two years at −80°C after freezing in PBS containing 30% sucrose. Before freezing,

the slices are incubated in PBS containing 30% sucrose overnight. The GFP signal of Tau-mGFP

mouse tissue is weak and needs to be enhanced by immunostaining for GFP prior to imaging.

7. The immunostained sections embedded on slides can be stored at 4°C for up to one month. For

longer storage, the slides need to be stored at −20°C.

8. During the tracing process, it is important to distinguish the GFP-positive myelin sheaths of

the target oligodendrocyte from those of neighboring cells. Because adjacent oligodendrocytes

can also express GFP, their myelin sheaths may appear to be in close proximity, potentially

leading to misidentification. To avoid this issue, for the morphological analysis of single

oligodendrocytes, only cells that are isolated within the field of view (regions where a single GFP-

positive oligodendrocyte was clearly identifiable) are selected for tracing. Furthermore, only

myelin sheaths directly connected to the cell body through proximal processes of the targeted

oligodendrocyte are included in the analysis.

9. Myelination analyses performed on large-caliber axons, such as cerebellar white matter and the

optic nerve, are easier and more reliable compared to those performed on small-caliber axons,

such as the axons in the corpus callosum, because we can clearly observe GFP-positive processes

ensheathing neuronal axons (Figs. 2 and 3) [25].

References

1. Stadelmann C, Timmler S, Barrantes-Freer A and Simons M (2019) Myelin in the Central

Nervous System: Structure, Function, and Pathology. Physiol Rev 99(3):1381-1431

doi:10.1152/physrev.00031.2018

2. Osanai Y, Yamazaki R, Shinohara Y and Ohno N (2022) Heterogeneity and regulation of

oligodendrocyte morphology. Front Cell Dev Biol 10:1030486 doi:10.3389/fcell.2022.1030486

3. Pérez-Cerdá F, Sánchez-Gómez MV and Matute C (2015) Pío del Río Hortega and the

discovery of the oligodendrocytes. Front Neuroanat 9:92 doi:10.3389/fnana.2015.00092

4. Weruaga-Prieto E, Eggli P and Celio MR (1996) Topographic variations in rat brain

oligodendrocyte morphology elucidated by injection of Lucifer Yellow in fixed tissue slices. J

Neurocytol 25(1):19-31 doi:10.1007/bf02284783

5. Bacmeister CM et al. (2022) Motor learning drives dynamic patterns of intermittent myelination

on learning-activated axons. Nat Neurosci 25(10):1300-1313 doi:10.1038/s41593-022-01169-4

6. Osso LA, Rankin KA and Chan JR (2021) Experience-dependent myelination following stress

is mediated by the neuropeptide dynorphin. Neuron 109(22):3619-3632.e3615

doi:10.1016/j.neuron.2021.08.015

7. Summers BS et al. (2025) Demyelination Produces a Shift in the Population of Cortical Neurons

That Synapse with Callosal Oligodendrocyte Progenitor Cells. eNeuro 12(6)

doi:10.1523/eneuro.0113-25.2025

8. Young KM et al. (2013) Oligodendrocyte dynamics in the healthy adult CNS: evidence for

myelin remodeling. Neuron 77(5):873-885 doi:10.1016/j.neuron.2013.01.006

9. Battulga B et al. (2025) Axonal Selectivity of Myelination by Single Oligodendrocytes

Established During Development in Mouse Cerebellar White Matter. Glia 73(4):873-886

doi:10.1002/glia.24660

10. Osanai Y et al. (2017) Rabies virus-mediated oligodendrocyte labeling reveals a single

oligodendrocyte myelinates axons from distinct brain regions. Glia 65(1):93-105

doi:10.1002/glia.23076

11. Toth E, Rassul SM, Berry M and Fulton D (2021) A morphological analysis of activity-

dependent myelination and myelin injury in transitional oligodendrocytes. Sci Rep 11(1):9588

doi:10.1038/s41598-021-88887-0

12. Looprasertkul S, Yamazaki R, Osanai Y and Ohno N (2025) Reduced Accumulation Rate and

Morphological Changes of Newly Generated Myelinating Oligodendrocytes in the Corpus

Callosum of Aged Mice. Glia 73(11):2322-2334 doi:10.1002/glia.70070

13. Rivers LE et al. (2008) PDGFRA/NG2 glia generate myelinating oligodendrocytes and

piriform projection neurons in adult mice. Nat Neurosci 11(12):1392-1401 doi:10.1038/nn.2220

14. Zhu X et al. (2011) Age-dependent fate and lineage restriction of single NG2 cells.

Development 138(4):745-753 doi:10.1242/dev.047951

15. Doerflinger NH, Macklin WB and Popko B (2003) Inducible site-specific recombination in

myelinating cells. Genesis 35(1):63-72 doi:10.1002/gene.10154

16. Hippenmeyer S et al. (2005) A developmental switch in the response of DRG neurons to ETS

transcription factor signaling. PLoS Biol 3(5):e159 doi:10.1371/journal.pbio.0030159

17. Mori T and Morimoto K (2014) Rabies virus glycoprotein variants display different patterns

in rabies monosynaptic tracing. Front Neuroanat 7:47 doi:10.3389/fnana.2013.00047

18. Cetin A et al. (2006) Stereotaxic gene delivery in the rodent brain. Nat Protoc 1(6):3166-3173

doi:10.1038/nprot.2006.450

19. Pitman KA et al. (2020) The voltage-gated calcium channel CaV1.2 promotes adult

oligodendrocyte progenitor cell survival in the mouse corpus callosum but not motor cortex. Glia

68(2):376-392 doi:10.1002/glia.23723

20. Tanaka T et al. (2021) Large-scale electron microscopic volume imaging of interfascicular

oligodendrocytes in the mouse corpus callosum. Glia 69(10):2488-2502 doi:10.1002/glia.24055

21. Arshadi C et al. (2021) SNT: a unifying toolbox for quantification of neuronal anatomy. Nat

Methods 18(4):374-377 doi:10.1038/s41592-021-01105-7

22. Wickersham IR, Finke S, Conzelmann KK and Callaway EM (2007) Retrograde neuronal

tracing with a deletion-mutant rabies virus. Nat Methods 4(1):47-49 doi:10.1038/nmeth999

23. Osakada F and Callaway EM (2013) Design and generation of recombinant rabies virus vectors.

Nat Protoc 8(8):1583-1601 doi:10.1038/nprot.2013.094

24. Cooper JJM et al. (2023) Chronic demyelination of rabbit lesions is attributable to failed

oligodendrocyte progenitor cell repopulation. Glia 71(4):1018-1035 doi:10.1002/glia.24324

25. Osanai Y et al. (2018) Length of myelin internodes of individual oligodendrocytes is controlled

by microenvironment influenced by normal and input-deprived axonal activities in sensory

deprived mouse models. Glia 66(11):2514-2525 doi:10.1002/glia.23502

Figure legends

Fig 1. Visualization and morphological analysis of individual oligodendrocytes in the corpus callosum of PDGFRa-CreERT2:Tau-mGFP double transgenic mice.

(a-f) Representative z-stacked images of an oligodendrocyte in the corpus callosum of 10-week-old PDGFRa-CreERT2:Tau-mGFP mice that were injected tamoxifen two weeks before perfusion. The tissue was immunostained for GFP (green), CC1 (white), Caspr (red), counterstained with Hoechst (blue), and imaged using a confocal microscope. The rectangles in the left panels (a–c) are magnified in the right panels (d–f). (g–i) A higher-magnification z-stacked image of the area marked by the rectangle in d. The image shows a GFP-labeled myelin sheath (green, g and i) flanked by Caspr-positive paranodes (red, arrowheads, h and i). This method illustrates a single myelin sheath observable in the z-stacked serial images. (j) GFP-labeled oligodendrocyte (green, arrowhead) in a z-stacked image; double-headed arrow indicates cell height. (k–l) A z-stacked image shown in i with the yellow line added in the analysis with the SNT plugin in Fiji. The GFP-positive myelin sheath (green, k), flanked by Caspr-positive paranodes (red, k), is traced with a yellow line (k and l), representing the path used to measure the length of a single internode formed by an individual oligodendrocyte. This tracing was performed across z-stacked serial images. (m–n) Representative images showing fiber diameter measurement (axonal diameter + myelin thickness) of a single internode shown in g. A GFP-labeled myelin sheath of the individual oligodendrocyte was selected and traced along its full

length using a z-stacked serial image. A segmented line was then drawn on a single plane of the

z-stacked image and divided into five segments, starting 1 µm from the tip of each side (the white

line and its segment lines, m). Fiber diameters were measured at six points along each internode

(the arrowheads indicate measurement sites and double-headed arrows indicate span of the fiber

diameter, n).

Fig 2. Analysis of Purkinje cell selective myelination by early-born oligodendrocytes using

PLP-CreERT:Tau-mGFP double transgenic mouse.

Representative single-plane confocal image of cerebellar white matter of an 8-week-old PLP-

CreERT:Tau-mGFP double transgenic mouse treated with a single tamoxifen injection at P8. (a)

GFP-labeled early-born oligodendrocytes and their myelin sheaths (green). (b) Merged image

showing Purkinje cell axons (calbindin, red) and early-born oligodendrocytes (GFP, green). (c)

Neuronal axons (neurofilament, white), including Purkinje cell axons and other axon subtypes,

along with early-born oligodendrocytes (green). (d) Confocal image showing Purkinje cell axons

(red), neurofilament-positive axons (white), and early-born oligodendrocytes with their myelin

sheaths (green). Due to close proximity of labeled oligodendrocytes, individual cells cannot be

clearly distinguished. However, since individual myelin sheath can be identified, we enable to

assess whether early-born oligodendrocytes preferentially myelinate Purkinje cell axons (*see* note

5).

Fig 3. Oligodendrocytes labeled by RV-GFP.

(a–c) RV-GFP-labelled oligodendrocytes in the optic chiasm (a), cerebellar white matter (b), and

corpus callosum (c) of 8-week-old mice perfused 4 days after injection. (d–f). Double-

fluorescence images of the GFP and Caspr signals in the region outlined by the white square in

panels a–c. respectively. The distal ends of processes are co-localized with Caspr signals,

indicating that the processes are myelin sheaths.

GFP/CC1/Caspr/Hoechst

GFP/Calbindin/Neurofilament

20µm