The tree cricket ear is a highly phase sensitive biomechanical interferometer

Emine Celiker[1], Gregory Patrick Sutton[2], Natasha Mhatre[3]

[1]School of Engineering, University of Leicester, University Road, Leicester, LE1 7RH, UK,

[2]School of Life and Environmental Sciences, University of Lincoln, Joseph Banks Laboratories, Green Lane, Lincoln, LN6 7DL, UK,

[3]Department of Biology, Western University, London, ON, N6A 5B7, Canada.

Abstract

We show that tree crickets localize the sound by having an auditory system that first splits the soundwave into two parts, sending one wave to an Anterior tympanic membrane, and sending the second to a Posterior tympanic membrane. The membranes then carry the split waves to a common point, the Tracheal wall, where the re-assembly of the wave causes mechanical interference that is sensitive to microsecond time delays. This interference mechanically transduces the time delay into a simple lateral strain at the Tracheal wall. This principle, interferometry, is usually thought of in terms of light (Michaelson and Morley, 1887), and can measure extremely small time-delays. Tree crickets use the same principle as the light interferometer, except with sound, to measure extremely precise locations.

Finding a singing cricket in a living room is infuriatingly difficult. For while large vertebrates, like us, can determine the direction of a sound from several features that are altered between when the sound hits the left and right ears (Grothe et al,. 2010;), resolving this precisely requires the nervous system react to small (hundreds of microseconds) time differences between the ears (von Kriegstein et al., 2008). For crickets, using this mechanism is even more difficult because their small size would require their nervous system to differentiate between even more impossibly small (microsecond) time differences; and yet female crickets have no trouble locating singing males (Mhatre and Balakrishnan 2008), even when one ear is removed (Huber et al., 1984; Kohne et al., 1992). We will show that tree crickets (*Oecanthus henryi*) accomplish this task by using an elegant method of mechano-acoustic interferometry.

Tree crickets (in the suborder *ensifera*) (Figure 1a) possess ears located in the tibia of their forelegs (Michelsen et al., 1994; Mhatre et al., 2009). The ear in each leg has two tympana: an anterior tympanic membrane (ATM) and a posterior tympanic membrane (PTM). These tympana are backed by an air-filled tube called the acoustic trachea, with an opening, the acoustic spiracle, located on the thorax. Each tympana receives multiple inputs of the same sound: directly to the external side, and through the acoustic trachea (Michelsen et al., 1994; Michelsen, 1998). Both membranes attach ventrally to a common cuticular structure, the tracheal wall (TW), with sound waves being carried by both membranes to this common point. The tracheal wall is also where the sensory neurons are located (Field and Matheson, 1998; Mhatre et al., 2021). While the geometry of the ear does result in a directionally dependent intra-aural time delay (Pulver et al., 2022), this time delay (on the order of microseconds) is orders of magnitude smaller than the time delays detected by vertebrates. Larger crickets are thought to solve this problem using a four input pressure difference system (Larsen and Michelson, 1978),but for even smaller insects this requires novel solutions. Flies solve this by using an independently evolved intertympanal mechanical linker (Mason et al., 2001), but it is unclear how smaller crickets like tree crickets solve the problem of sound localization.

We have mechanically approximated the tree cricket ear with a three-element mechanical model: 1) an anterior tympanic membrane, which receives one sound input, 2) a posterior tympanic membrane, which receives a second sound input, 3) a tracheal wall, which is acted upon through the displacement of each membrane (Figure 1b,c). The specific mass, stiffness, and damping parameters used are detailed in supplementary material. This three-element model mechanically recreates the acoustic response of *O. henryi* tracheal wall (Figure 2).

The lumped element model shows that the key to this system is the individual membranes carrying the soundwaves to the common point, the tracheal wall, where the interference between the two soundwaves causes the strains at the tracheal wall to differ, depending on a phase difference between the sound waves hitting each membrane (Figure 2c,d and supplemental movie 1). A zero degree phase difference is transduced into vertical strains in the tracheal wall, a 180 degree phase difference into horizontal strains, and intermediate phase differences into elliptical strains (Figure 2 and supplemental movie 1). The two acoustic inputs resulting in elliptical movements have been observed previously in both insects and mammals (Stephen and Bennet-Clarke, 1982; Vavakou et al., 2024; Frost et al. 2025) with Bennet-Clark commenting "it would appear at first sight that the two drives will interact to reduce the sharpness of tuning of each other"; we show here that these ellipses are tied to the phase differences in the sound, and thus can be used to precisely determine the direction of the sound source. It is thus phase-dependent interference between the two sound waves is detected by the nervous system – making the tree cricket ear a biological example of a mechano-acoustic phase interferometer akin to that developed for light in (Michelson and Morley, 1887).

This bio-mechanical phase interferometer requires a small number of simple components: 1) two similarly-sized membranes transducing disparate vibrational inputs to a common point moving at similar levels, and 2) a system that allows for sound direction to create a phase difference between the soundwaves hitting each membrane and 3) mechanosensory neurons at that common point (the tracheal wall) that respond preferentially to a single direction of movement.

The first of these components is a feature common to many Ensiferan species and its functional significance was hitherto unappreciated. For instance bush crickets (fellow *ensiferans*), have a very similar ear morphology consisting of two membranes carrying the sound to a common point (Montealegre-Z and Robert, 2015). For component two, small phase differences between closely placed tympana may be created due to diffraction by the body of the animal and the substrate it stands on. Indeed, a directionally dependent phase difference in the sound hitting the ATM and PTM has been observed in bush crickets (Pulver et al., 2022). Such a difference is accentuated by the geometry of the tracheal branches carrying the sound to each membrane (Veitch et al., 2021). Moreover, similar elliptical movements in the tracheal wall have been observed in tree crickets (Mhatre et al, 2021) and also in bush crickets (Vavakou et al., 2024), strongly suggesting that this phase interferometry method is widespread among Ensiferans. Finally, it is also possible that mechanosensory neurons respond preferentially to certain axes of stimulation (Yarger and Fox, 2018; Chaiyasitdhi et al., 2025). The simple components required for such a system also leads the possibility that many hearing systems in small animals may be similar.

This simplicity provides a huge number of parameters that can then be further tuned to specialize different Ensiferan ears for different inputs. Membrane geometry (Mhatre et al., 2009; Rajarman et al., 2013), membrane mechanical properties (Vincent and Wegst, 2004), membrane anisotropy (Mhatre et al., 2009; Mhatre et al., 2012; Malkin et al., 2014), acoustic trachea geometry (Schmidt and Römer, 2013; Veitch et al., 2021; Celiker et al., 2022), fluid properties within the auditory vesicle (Veitch et al., 2021; Lomas et al., 2012), or mechanisms for active amplification

(Mhatre and Robert, 2013) could all be modified to specialize the interferometer for specific sound inputs.

Moreover, as this system is sensitive to the phase of the sound (as opposed to the time delay), the higher the frequency of the sound, the easier it is for the system to mechanically transduce the precise direction of the sound's source. This would have obvious utility in detecting bats (which have call frequencies in excess of 60kHz, Pulver et al., 2022) as well as detecting the song in ultrasonic crickets (Robillard et al., 2013).

We have trouble locating that singing cricket because of its tonal nature and our nervous system has difficulty resolving the minute time delays in such signals. Tree crickets, on the other hand, pinpoint the sound using mechano-acoustic phase interferometry, letting the tympanal membranes and the tracheal wall precisely tell them where a singing conspecific is.

Figure 1. The tree cricket ear anatomy. (a) The tree cricket *Oecanthus henryi*. (b) A cross-sectional sketch of the tree cricket *Oecanthus henryi* hearing organ near the tympanal end of the acoustic trachea. ATM = Anterior Tympanic Membrane, PTM = Posterior Tympanic Membrane, TO = Tympanal Organ. (c) The fully-coupled mechanical system representing the tracheal wall (TW) displacement. The tracheal wall (stiffness k_{TW}=20.62 N/m , damping c_{TW}=0.00047, and mass M_{TW}=2.8×10^{-10} kg) is acted upon by the anterior tympanic membrane (k_{ATM}=41.23 N/m , damping c_{ATM}=0.00047, and mass M_{ATM}=1.6×10^{-10} kg), and posterior tympanic membrane (k_{PTM}=41.23 N/m , damping c_{PTM}=0.00047, and mass M_{PTM}=1.4×10^{-10} kg). The angle between the tympana and tracheal wall $\theta = 60°$ (Mhatre et al., 2009). $F_1(t)$ and $F_2(t)$ represent the harmonic forces of equal magnitude and a frequency of 3 kHz acting on the tympana.

Figure 2. The displacement sensitivity of the tracheal wall in the xz-plane at the calling song frequency 3 kHz (at 24 °C) of *Oecanthus henryi*. Phase lag refers to the phase difference between the forces of same magnitude activating the anterior tympanic membrane (ATM), and the posterior tympanic membrane (PTM). (a) The displacement pattern of the tracheal wall when the ATM and PTM are acted on by forces *in-phase* (phase lag=0°) and *out-of-phase* (phase lag=180°) of equal magnitude (0.04 Pa), and (b) unequal magnitude. (c) The elliptic motion of the tracheal wall for large phase differences between the ATM and PTM. (d) The displacement magnitude of the tracheal wall in the x- and z- directions with respect to the phase lag of the forces acting on the ATM and PTM. (e) The elliptic motion of the trachea wall for small phase differences between the ATM and PTM.

References

Celiker, E., Woodrow, C., Rocha-Sánchez, A. Y., Chivers, B. D., Barrientos-Lozano, L., & Montealegre-Z, F. (2022). Beyond the exponential horn: a bush-cricket with ear canals which function as coupled resonators. *Royal Society Open Science*, **9**(10), 220532.

Chaiyasitdthi, A., Nowotny, M., Van der Heidjen, M., & Warren, B. (2025). Stretching of the insect mechanoreceptor evokes mechano-electrical transduction in auditory chordotonal neurons. *bioRxiv*, doi 10.1101/2025.06.27.662008v1.

Field, L. H., & Matheson, T. (1998). Chordotonal Organs of Insects. *Advances in Insect Physiology* 27, 1-56.

Frost, B. L., Strimbu, C. E., & Olson, E. S. (2025). Narrow elliptical motion at the outer hair cell-Deiters' cell junction explains disparate features of uniaxial displacement measurements. *Hearing Research*, 109189.

Grothe, B., Pecka, M., & McAlpine, D. (2010). Mechanisms of sound localization in mammals. *Physiological Reviews*, 90 (3) 983-1012.

Huber, F., Kleindienst, H. U., Weber, T., & Thorson, J. (1984). Auditory behaviour of the cricket. *Journal of Comparative Physiology A.*, 155, 725-738.

Kohne, R., Atkins, S., Stout, J., & Atkins, G. (1992). Enhanced calling song syllable period discrimination during one-eared phonotaxis by the female cricket (*Acheta domesticus*). *Journal of Comparative Physiology A.*, 170, 357-362.

Larsen, O. N., & Michelson, A. (1978). Biophysics of the Ensiferan Ear. *Journal of Comparative Physiology*, 123, 217-227.

Lomas, K. F., Greenwood, D. R., Windmill, J. F., Jackson, J. C., Corfield, J., & Parsons, S. (2012). Discovery of a lipid synthesising organ in the auditory system of an insect. *PLoS One*, **7**(12), e51486.

Malkin, R., McDonagh, T. R., Mhatre, N., Scott, T. S., & Robert, D. (2014). Energy localization and frequency analysis in the locust ear. *Journal of The Royal Society Interface*, **11**(90), 20130857.

Mason, A. C., Oshinsky, M. L., & Hoy R. R. (2001). Hyperacute directional hearing in a microscale auditory system. *Nature*, 410, 686-690.

Mhatre, N., & Balakrishnan, R.. (2008). Predicting acoustic orientation in complex real-world environments. *Journal of Experimental Biology*, 211(17) 2779-2785.

Mhatre, N., Montealegre-z, F., Balakrishnan, R., & Robert, D. (2009). Mechanical response of the tympanal membranes of the tree cricket *Oecanthus henryi*. *Journal of Comparative Physiology A*, **195**, 453-462.

Mhatre, N., Montealgre-Z, F., Balakrishnan, R., & Robert, D. (2012, April). Sound reception and radiation in a small insect. *In Acoustics 2012*.

Mhatre, N., & Robert, D. (2013). A tympanal insect ear exploits a critical oscillator for active amplification and tuning. *Current Biology*, **23**(19), 1952-1957.

Mhatre, N., Dewey, J. B., Quiñones, P. M., Mason, A., Applegate, B. E., & Oghalai, J. S. (2021). Reconstruction of sound driven, actively amplified and spontaneous motions within the tree cricket auditory organ. *bioRxiv*, 2021-11.

Michelsen, A., Popov, A. V., & Lewis, B. (1994). Physics of directional hearing in the cricket Gryllus bimaculatus. *Journal of Comparative Physiology A*, **175**, 153-164.

Michelsen, A. (1998). The tuned cricket. *Physiology*, **13**(1), 32-38.

Michelson, A. A. and Morley, A. A. (1887). *Philos. Mag. S.* **5** 24 No.151, 449-463.

Montealegre-Z, F., & Robert, D. (2015). Biomechanics of hearing in katydids. *Journal of Comparative Physiology A*, **201**(1), 5-18.

Pulver, C. A., Celiker, E., Woodrow, C., Geipel, I., Soulsbury, C. D., Cullen, D. A., Rogers, S. M., Veitch, D. & Montealegre-Z, F. (2022). Ear pinnae in a neotropical katydid (Orthoptera: Tettigoniidae) function as ultrasound guides for bat detection. *Elife*, **11**, e77628.

Rajaraman, K., Mhatre, N., Jain, M., Postles, M., Balakrishnan, R., & Robert, D. (2013). Low-pass filters and differential tympanal tuning in a paleotropical bushcricket with an unusually low frequency call. *Journal of Experimental Biology*, 216(5), 777-787.

Robillard, T., Montealegre-Z, F., Desutter-Grandcolas, L., Grandcolas, P., & Robert, D. (2013). Mechanisms of high-frequency song generation in brachypterous crickets and the role of ghost frequencies. *Journal of Experimental Biology*, **216**(11), 2001-2011.

Schmidt, A. K., & Römer, H. (2013). Diversity of acoustic tracheal system and its role for directional hearing in crickets. *Frontiers in Zoology*, **10**, 1-9.

Stephen, R. O., & Bennet-Clark, H. C. (1982). The anatomical and mechanical basis of stimulation and frequency analysis in the locust ear. *Journal of Experimental Biology*, **99**(1), 279-314.

Von Kriegstein, K., Griffiths, T. D., Thompson, S. K., & McAlpine, D. (2008). Responses to interaural time delay in human cortex. *Journal of Neurophysiology*, 100 (5) 2712-2718.

Vavakou, A., Scherberich, J., Nowotny, M., & Heijden, M. V. D. (2024, February). Elliptical sound-induced motion in the bushcricket hearing organ. In AIP Conference Proceedings (Vol. 3062, No. 1). AIP Publishing.

Veitch, D., Celiker, E., Aldridge, S., Pulver, C., Soulsbury, C. D., Jonsson, T., Woodrow, C., & Montealegre-Z, F. (2021). A narrow ear canal reduces sound velocity to create additional acoustic inputs in a microscale insect ear. *Proceedings of the National Academy of Sciences*, **118**(10), e2017281118.

Vincent, J. F., & Wegst, U. G. (2004). Design and mechanical properties of insect cuticle. *Arthropod structure & Development*, **33**(3), 187-199.

Yarger., A. M., & Fox, J. L. (2018). Single mechanosensory neurons encode lateral displacements using precise spike timing and threshholds. *Proceedings of the Royal Society B*, **285** (1887), DOI 10.1098/rspb.2018.1759.