ADHD-like traits reshape the balance between inhibitory control and predictive processes

Karolina Horváth [1*], Bianka Brezóczki [2,3,4*], Adrienn Holczer [1], Teodóra Vékony [1*], Dezső Németh [1,5,6*]

[1] Gran Canaria Cognitive Research Center, Atlántico Medio University, Las Palmas de Gran Canaria, Spain

[2] Doctoral School of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary

[3] Institute of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary

[4] Brain, Memory and Language Research Group, Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary

[5] BML-NAP Research Group, Institute of Psychology, Eötvös Loránd University and Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary

[6] Centre de Recherche en Neurosciences de Lyon CRNL U1028 UMR5292, INSERM, CRNS, Université Claude Bernard Lyon 1, 69500 Bron, France

* These authors contributed equally to this work

Correspondence: Correspondence should be addressed to Dezso Nemeth at dezso.nemeth [at]inserm.fr

Abstract

Adaptive behavior relies on a dynamic balance between flexible, goal-directed control and efficient, automatic processes. This equilibrium is often disrupted in Attention-Deficit/Hyperactivity Disorder (ADHD), a condition understood to exist along a continuum of traits in the general population. While ADHD is consistently linked to response inhibition deficits, its relationship with statistical learning (a mechanism for habit learning) and, crucially, the interaction between these functions, remains underexplored. Here, using a novel paradigm in a non-clinical sample of university students (n = 226), we investigated how ADHD-like traits modulate the interplay between response inhibition and statistical learning. We found that higher ADHD-like traits were associated with poorer response inhibition, confirming previous research. Importantly, we uncovered an antagonistic relationship between inhibition and statistical learning, where weaker inhibitory control typically led to enhanced learning of environmental regularities.

However, this learning advantage progressively diminished across the ADHD trait continuum, becoming markedly reduced in individuals with high symptom prevalence. These findings provide novel evidence that ADHD-like traits influence not only isolated neurocognitive processes but also their dynamic interaction, highlighting a spectrum-based mechanism that may underlie the transition from adaptive variability to maladaptive behavioral patterns. This work advocates for a dimensional approach to ADHD, emphasizing early detection and targeted interventions for individuals with varying levels of symptomatic expression, broadening support beyond traditional diagnostic boundaries.

Keywords: Attention Deficit/ Hyperactivity Disorder, statistical learning, cognitive control, cognitive tradeoff, spectrum-approach

Introduction

Navigating a complex and ever-changing world requires a dynamic interplay between flexible, goal-directed control and efficient, automatic processes. The capacity for goal-directed behavior allows for deliberate adaptation to novel demands, albeit at a significant cognitive cost (Grahek et al., 2023; Kurzban et al., 2013; Musslick & Cohen, 2021), while automatic processes enable rapid and efficient execution of well-learned actions at the expense of flexibility (Balleine & O'Doherty, 2010). An optimal balance between these two systems is fundamental for adaptive functioning. However, this equilibrium can be disrupted by structural or functional alterations within these neurocognitive systems (Gillan et al., 2015, 2016). Attention-Deficit/Hyperactivity Disorder (ADHD) is a prime example of such a disruption, characterized by enduring patterns of inattention and hyperactivity-impulsivity (American Psychiatric Association, 2022). A prevailing hypothesis posits that impaired goal-directed processes are a core neurocognitive deficit in ADHD (Barkley, 1997; Bush et al., 2005; Castellanos et al., 2006; Häger et al., 2021; Martel et al., 2007; Molitor et al., 2019; Natsheh et al., 2019; Seidman, 2006; Silverstein et al., 2020), which consequently perturbs the typical balance between goal-directed and automatic mechanisms (Hogarth et al., 2012). While the clinical presentation of ADHD is well-defined, mounting evidence suggests it is not a categorical disorder but rather exists on a continuum, with traits of inattention and hyperactivity-impulsivity distributed across the general population (Greven et al., 2016; Hengartner & Lehmann, 2017; Herrmann et al., 2009; Levy et al., 1997; T. Li et al., 2019; Paloyelis et al., 2009; Panagiotidi et al., 2018, 2019; Polner et al., 2015; Schippers et al., 2024; Vogel et al., 2018). This dimensional perspective provides a powerful framework to investigate how subtle variations in ADHD-like traits modulate the fundamental trade-off between cognitive control and automaticity in a non-clinical population.

To investigate this trade-off along the non-clinical ADHD spectrum, it is essential to examine the core cognitive functions involved. Goal-directed behavior depends on prefrontal cortex–related executive functions (Diamond, 2013; Friedman & Robbins, 2022; Miyake et al., 2000), such as response inhibition, which refers to the ability to suppress inappropriate or prepotent actions (Chambers et al., 2009; Wright et al., 2014). Prior research on isolated components of executive functioning has predominantly focused on response inhibition (Crosbie et al., 2013; Hadas et al., 2021; Hart et al., 2014; X. Li et al., 2025; Ma et al., 2012; Nigg, 2001; Nigg et al.,

2002; Silverstein et al., 2020; Wodka et al., 2007). A substantial body of evidence indicates that impairments in response inhibition are present in ADHD, both structurally and functionally (Bush et al., 2005; Castellanos & Tannock, 2002; Crosbie et al., 2013; Hadas et al., 2021; Hart et al., 2014; X. Li et al., 2025; Liston et al., 2011; Ma et al., 2012; Nigg, 2001; Nigg et al., 2002; Silverstein et al., 2020; Wodka et al., 2007). These alterations have been observed not only in clinical populations but also in individuals with non-clinical ADHD-like traits, where such traits are associated with reduced response inhibition performance and, consequently, greater impulsive behavior (Crosbie et al., 2013; Polner et al., 2015). In the context of balancing goal-directed and automatic processes, diminished inhibitory control may lead to a greater reliance on the automatic system, including mechanisms such as statistical learning (SL). SL is the unconscious acquisition of environmental regularities that support predictive processing and underlie the development of habits and skills (Aslin, 2017; Conway, 2020; Horváth et al., 2022; Theeuwes et al., 2022). While the majority of studies report largely intact SL in both clinical (Adi-Japha et al., 2011; Karatekin et al., 2009; Pedersen & Ohrmann, 2018; Takács et al., 2017; Vloet et al., 2010) (for exceptions, see [Barnes et al., 2010]), and non-clinical populations with ADHD-like traits (Parks & Stevenson, 2018), the interplay between inhibition and SL may still be significantly modulated by ADHD traits. We therefore posit that the interaction between these two distinct neurocognitive systems, impaired response inhibition and preserved statistical learning, represents a key mechanism underlying the behavioral profile of ADHD-like traits on a continuum, a relationship that has yet to be systematically investigated.

Accordingly, the core limitation of prior research is that these processes have primarily been examined in isolation, whereas real-world learning and adaptive behavior typically rely on the coordinated operation of multiple neurocognitive functions (Horváth et al., 2022; Pedraza, Farkas, et al., 2024). As such, this isolated approach is limited, since dysfunctions may emerge when these processes are forced to interact (Balleine & O'Doherty, 2010; Castellanos & Tannock, 2002). To date, only a single study in a clinical child population has investigated such interactions, revealing a bias toward automatic behavior in ADHD alongside disrupted goal-directed control (Natsheh et al., 2019). Importantly, ADHD in childhood and adulthood may differ at the neurocognitive level due to the maturation of prefrontal regions (Shaw et al., 2007). Therefore, it is crucial to examine the interactive relationship between SL processes and goal-directed versus automatic behavior.

Moreover, to better capture the underlying mechanisms of ADHD symptomatology, it is essential to move beyond traditional dichotomous clinical–control group comparisons. In the present study, we adopt a spectrum-based methodological approach to examine how individual differences in ADHD-like traits relate to neurocognitive processes, specifically the interplay between inhibitory control and SL. Based on existing literature, we hypothesized an antagonistic relationship between SL and inhibitory control, such that reduced inhibitory control may facilitate stronger statistical learning. Building on prior research, we further hypothesized that this relationship would be modulated by the prevalence of ADHD-like traits, with the interaction between SL and inhibitory control varying systematically across the trait spectrum rather than following a simple linear pattern.

Methods

Participants

We recruited three hundred and seventeen university students who participated in an online experiment and received course credit for their participation. All participants provided written informed consent, and the research protocol adhered to ethical guidelines established by the Declaration of Helsinki. All protocols were approved by the Eötvös Loránd University Research Ethics Board (2024/214). The dataset for this study is the same as that used by Vékony et al. (2025).

Several exclusion criteria were applied to ensure good data quality. Participants were excluded if they reported any neurological or psychiatric conditions ($n = 43$), were currently taking medications known to influence central nervous system functioning ($n = 8$), had consumed alcohol within three hours of testing ($n = 19$), or possessed prior familiarity with ASRT paradigms ($n = 10$). Incomplete data also resulted in exclusion, particularly for participants who did not complete the Adult ADHD Self-Report Scale (ASRS; $n = 25$) or who failed the attention test during the questionnaires ($n = 7$). To ensure participants were engaged, a statement was included in the questionnaires: "If you are paying attention right now, please select the 'Agree' option." Participants who did not follow this instruction were considered inattentive, and their data was excluded from the final analysis.

Task performance served as an additional screening criterion. Participants demonstrating accuracy rates below 80% (n = 11) were excluded to ensure adequate task engagement. Similarly, those requiring more than 50 minutes to complete the task (n = 7) or who restarted the procedure during testing (n = 3) were removed from the analyses. Following these exclusions, 91 participants were removed in total. Notably, some satisfied multiple exclusion criteria, resulting in a final sample of 226 participants (retention rate: 71.6%). The mean age of the participants was 21.42 ± 4.76 SD years (183 female, 42 male, and 1 non-specified). The sample of 226 participants showed a varied educational background: 178 participants (78.8%) had a high school education or lower, 41 (18.1%) had a bachelor's degree or equivalent, and 7 (3.1%) had a master's degree or equivalent.

The Cognitive Trade-off Task (CTT)

The Cognitive Trade-off Task (CTT) integrates the Alternating Serial Reaction Time Task (ASRT), a well-established visuomotor statistical learning paradigm (Farkas et al., 2023; Howard & Howard, 1997; Janacsek et al., 2012), with a Go/No-Go component, traditionally employed to assess response inhibition. In the present study, we adapted the ASRT task to simultaneously measure statistical learning and inhibitory control by incorporating a Go/No-Go component.

The experimental task was programmed in JavaScript using the jsPsych library v.6.1.0. (Vékony, 2021). Depending on the experimental condition (detailed below), either a dog or a cat head appeared in one of four positions arranged horizontally in the center of the screen (Figure 1A). The four circles were positioned in the middle of the screen, and participants received instructions to press the designated keyboard button ('S', 'F', 'J', or 'L' from left to right) as rapidly and accurately as possible when presented with go stimuli, corresponding to the stimulus position. Conversely, when presented with No-Go stimuli, participants were instructed to suppress their response and refrain from pressing any button.

When participants executed the correct button press on 'go' trials, the target stimulus disappeared immediately, followed by a 120 ms interstimulus interval before the presentation of the subsequent stimulus. Incorrect button presses on 'go' trials resulted in the stimulus remaining visible until participants selected the correct response. Notably, in 10 out of every 80 trials, either a cat head or a dog head stimulus functioned as the 'No-Go' signal, requiring participants to suppress any button

press. The No-Go stimuli remained displayed for 1000 ms before disappearing, regardless of the participants' responses.

Unbeknownst to participants, an 8-element sequence was continuously repeated throughout the experimental session. Within the eight-element sequence, sequence components alternated with random elements. Consequently, the presentation of a sequence stimulus was invariably followed by a random element, creating a consistent alternating pattern (e.g., $2 - r - 4 - r - 3 - r - 1 - r$, where 'r' represents a random location, while the numbers indicate the predetermined location of the sequence) (Figure 1B).

This alternating structure of the ASRT task led to certain three-trial combinations, known as triplets, occurring more frequently than others. Some triplets occur with a higher probability (high-probability triplets) than other triplets (low-probability triplets). Trial classification depends on the final trial within the triplet, independent of whether it constitutes a random trial. For example, in the sequence $2 - r - 4 - r - 3 - r - 1 - r$, high-probability triplets are, e.g., 2-r-4 and 4-r-3. Conversely, using the same sequence, low-probability triplets are, for example, 2-r-3 and 3-r-4. During the experiment, high-probability triplets were constructed either with two pattern trials and one random trial in the middle (50% of cases) or with two random trials and one pattern trial in the middle of the triplet (12.5% of trials). All other triplets were defined as low-probability triplets, which appeared in 37.5% of the trials. In total, 62.5% of trials functioned as the concluding element of high-probability triplets, while 37.5% of trials served as the final element of low-probability triplets (Figure 1C).

The experimental design utilized counterbalanced stimulus assignments to mitigate potential animal (cat or dog) preference effects. Half of the participants were assigned to respond to a dog head (go condition) while inhibiting responses to a cat head (No-Go condition), whereas the remaining participants received the opposite stimulus-response mapping. The appearance of 'No-Go' stimuli was randomized across blocks, appearing at both pattern and random positions within the alternating sequence. The structure also ensured that no two consecutive 'No-Go' stimuli were presented, and the initial two trials of each block were never 'No-Go' trials.

Participants completed 30 experimental blocks. Within each block, participants encountered the 8-element sequence 10 times, resulting in 80 stimuli per block. Each block contained 10 No-Go

stimuli and 70 Go stimuli, maintaining a consistent 12.5% No-Go trial ratio throughout the experiment.

Figure 1. The Cognitive Trade-off Task. (A) Participants completed 30 blocks, each containing 80 trials. In each trial, a 'go' stimulus appeared on the screen, and participants were instructed to press the corresponding button (S, F, J, or L) as quickly and accurately as possible. Ten percent of the trials were 'No-Go' stimuli, which required participants to withhold their response. A hidden structure was embedded in the task, where every second trial followed a predetermined, 8-element probabilistic sequence. **(B)** The alternating structure of the task led to certain three-trial combinations (triplets) occurring more frequently than others. These were categorized as high-probability triplets, while the less frequent combinations were considered low-probability triplets. The elements within these triplets were either pattern elements (indicated by a black background) or random elements (indicated by an orange background). **(C)** Due to the alternating structure of the task, high-probability triplets were formed in two ways. The most common configuration, occurring in 50% of the trials, consisted of two pattern elements and one random element in the middle. The second configuration, occurring on 12.5% of the trials, was formed by two random elements with a pattern element in the middle. In total, 62.5% of trials were categorized as high-probability triplets. The remaining 37.5% of trials were considered low-probability triplets.

Adult ADHD Self-Report Scale (ASRS)

ADHD tendencies were assessed using a self-report questionnaire, which enabled us to identify the prevalence of ADHD-like symptoms along the spectrum. The Adult ADHD Self-Report Scale (ASRS) was selected to provide a quantitative assessment of ADHD symptom prevalence within our non-clinical sample (Kessler et al., 2005). This psychometric instrument represents a widely validated screening tool developed specifically for identifying ADHD symptoms in adult populations (Kessler et al., 2005). The ASRS comprises 18 self-report items utilizing a 5-point Likert scale format (ranging from 0 = never to 5 = very often) to evaluate symptom frequency over the preceding six-month period. The maximum possible score on the ASRS questionnaire is 72 points. In our study, the highest score was 57 (Figure 2). This bifactorial structure aligns with the DSM-5 conceptualization of ADHD symptom presentation (American Psychiatric Association, 2022). The total ASRS score for each participant was calculated and used as a measure of ADHD-like symptoms in our analyses.

Figure 2. Distribution of the ASRS scores. The scores, which can range from 0 to 72, extended from a minimum of 6 to a maximum of 57 (M = 31.66, SD = 10.04). The score distribution exhibited a slight negative skewness (skewness = -0.013). The majority of participants scored in the mid-range of the scale.

Procedure

The experiment was hosted by the Gorilla Experiment Builder platform (https://gorilla.sc; (Anwyl-Irvine et al., 2020). Upon initial access, participants reviewed and provided informed consent

before proceeding to the experimental session. The testing session started with the CTT, which consisted of 30 blocks. Post-task questioning revealed that no participants could identify the hidden sequence or describe any underlying pattern, demonstrating that the task appropriately measured statistical learning (Horváth et al., 2022; Vékony et al., 2022). Additionally, demographic information was collected, including age, educational background, and other relevant participant characteristics. One week later, participants were asked to complete a battery of self-report questionnaires on Qualtrics, which included the ASRS.

Data preparation

All data processing, statistical computations, and visualization were performed using R version 4.4.2. (R Core Team, 2024). Prior to analysis, trials were categorized based on whether they served as the final element of a high-probability or low-probability triplet. For the calculation of statistical learning measures, 'No-Go' trials were excluded. Trials with response times below 100 ms or above 1000 ms were removed. Furthermore, trials falling below or beyond three standard deviations were excluded to control for extreme outlier responses. Repetitions (e.g., 3-3-3) and trills (e.g., 2-1-2) were excluded because participants typically demonstrate faster responses to these sequences due to reduced motor complexity rather than statistical learning effects (Soetens et al., 2004). Trials with incorrect responses were excluded.

No-Go performance score calculation

We measured response inhibition performance through 'No-Go' trials. Block wise accuracy on 'No-Go' trials was calculated as the proportion of trials where participants successfully inhibited their response. Correct inhibition was defined as refraining from pressing the corresponding key throughout the 1000 ms stimulus duration.

Statistical analysis

We applied linear mixed-effects modeling (LMM) using the *mixed* function from the *afex* package (Singmann et al., 2024). We initially implemented the maximal random effect structure. If convergence issues or singular fit warnings occurred, we first removed correlations between random variables. If they persisted, we removed interactions between variables. We retained the most complex models that converged adequately. Post-hoc analyses of simple means and simple

slopes were performed using estimated marginal means and estimated marginal trends calculated with the *emmeans* package (Lenth, 2018). Fixed effects were evaluated using Type III tests. For significant interactions involving continuous predictors, we divided continuous variables based on their mean values, with cut-points set at ±1 SD from the mean. This approach was applied when No-Go accuracy, Block, or ASRS scores were treated as continuous variables. Statistical significance was set at α = 0.05 for all tests. Data visualization was performed using the *ggplot2* package, with additional support from *ggplot* and *afex*.

We constructed two models to address our research questions. First, we examined how the level of ADHD-like traits related to response inhibition. Second, we investigated how ADHD-like traits and response inhibition interact to influence statistical learning. In our first model, the outcome variable was No-Go accuracy (measuring response inhibition; in percentage), calculated per participant for each block, with block (within-subject) and ASRS total score (between-subject) as predictor variables. Block and ASRS total scores were mean-centered. In our second model, the outcome variable was reaction time, calculated per participant for each Block and Triplet Type, with Block, Triplet type, and No-Go accuracy as within-subject predictors, and ASRS total score as a between-subject predictor. Block, No-Go accuracy, and ASRS total score were all mean-centered.

1. $Model_1$: *No-Go accuracy ~ Block x ASRS score + (Block | ID)*

2. $Model_2$: *Go RT ~ Block x Triplet type x ASRS score x No-Go accuracy + (Block | ID)*

Results

Higher ADHD-like traits are associated with decreased response inhibition.

We fitted a linear mixed-effects model with accuracy on No-Go trials as the outcome variable. This was operationalized as the mean percentage of correct 'No-Go' responses per block. Fixed effects included Block (centered), ASRS scores (centered), and their interaction. Random effects included participant ID and random slopes for block within participants, allowing individual variation in inhibitory control trajectories across time and estimating the correlation between participants' starting levels and their rate of change. The model revealed a significant main effect for both Block ($F_{(1, 224.61)}$ = 153.11, $p < .001$) and ASRS score ($F_{(1, 223.94)}$ = 8.59, $p = .004$),

indicating that response inhibition significantly differed across levels of both factors. Response inhibition was worse with higher levels of ASRS scores (b = -0.003, SE = 0.001) (Figure 3), and significantly decreased as the task progressed (b = -0.006, SE = 0.001). However, the interaction between ASRS scores and Block was not statistically significant ($F_{(1, 224.09)}$ = 0.44, p = .509), suggesting that the effect of ASRS score on response inhibition did not change with the progress of the task. These findings revealed that higher ADHD-like traits were associated with a decrease in response inhibition. Full results are presented in Supplementary Table 1, including b, SE b, 95% CI, t, df, and p values.

Figure 3. Relationship between ADHD-like symptoms and response inhibition. This figure illustrates the relationship between participants' self-reported ADHD-like symptoms and their response inhibition. The x-axis represents the ASRS scores, centered around the sample mean, while the y-axis shows No-Go accuracy in percentage. The graph shows a negative relationship, where higher ASRS scores were associated with a decrease in No-Go accuracy. This finding indicates that individuals with more self-reported ADHD-like traits tended to have poorer inhibitory control.

ADHD-like traits reshape the balance between inhibitory control and statistical learning.

We fitted a linear mixed-effects model with the blockwise mean reaction time to 'go' trials as the outcome variable. Fixed effects included a four-way interaction between Block (centered), Triplet Type, ASRS scores (centered), and No-Go accuracy (centered). Random effects included participant ID and random slopes for block within participants, allowing individual variation in learning trajectories across time. The model revealed a significant main effect of Block ($F_{(1, 228.50)}$

= 220.16, $p < .001$), indicating that reaction time to 'go' trials significantly decreases across levels of Block ($b = -1.806$, SE = 0.122). The model revealed a significant main effect of Triplet Type ($F_{(1, 13044.25)} = 83.34$, $p < .001$), indicating that reaction time to 'go' trials significantly differed between high and low probability triplets. Participants were faster to high-probability triplets ($b = -3.067$, SE = 0.336), indicating statistical learning. The interaction between Block and Triplet Type was not significant ($F_{(1, 13044.64)} = 2.57$, $p = .109$). The main effect of ASRS was also not significant ($F_{(1, 1224.74)} = 0.29$, $p = .594$). The main effect of No-Go performance was significant ($F_{(1, 13454.02)} = 801.40$, $p < .001$), indicating faster RT in blocks of weaker No-Go performance ($b = 55.63$, SE = 1.965). The interaction between No-Go Performance and Block was also significant ($F_{(1, 11556.93)} = 41.20$, $p < .001$). Simple slopes analyses were performed at the beginning, middle, and end points of the Block factor, revealing significant differences between time points in the slopes of No-Go performance (all $p < .001$). The largest difference in RTs based on No-Go accuracy performance was found at the beginning of the task. Participants with better response inhibition (higher No-Go accuracy) showed slower reaction times, particularly early in the task. This relationship diminished as the task progressed. Post-hoc comparison of estimated marginal trends indicated that the effect of No-Go Performance on 'go' reaction times weakened over time (early blocks: $b = 67.3$, $SE = 2.82$, $p < .001$; middle blocks: $b = 55.6$, $SE = 1.96$, $p < .001$; late blocks: $b = 43.8$, $SE = 2.54$, $p < .001$).

The interaction between No-Go Performance and Triplet Type was also significant ($F_{(1, 13049.04)} = 22.36$, $p < .001$). Simple effects analysis confirmed that the difference between high and low probability triplets varied significantly based on No-Go accuracy levels. At low No-Go accuracy, participants demonstrated the largest statistical learning effect ($b = -9.40$, $SE = 0.979$, $p < .001$). This effect remained significant but was diminished at moderate No-Go accuracy ($b = -6.16$, $SE = 0.672$, $p < .001$) and was smallest at high No-Go accuracy ($b = -2.92$, $SE = 0.940$, $p = 0.002$). Pairwise contrasts were conducted to investigate the differences in the statistical learning effect between No-Go accuracy levels. The analysis revealed significant differences across all comparisons. A significant decrease in the learning effect was observed when comparing the low and moderate No-Go accuracy levels ($b = -3.24$, $SE = 0.685$, $p < .0001$). The largest difference was found between the low and high No-Go accuracy levels, where the learning effect was significantly reduced ($b = -6.48$, $SE = 1.370$, $p < .001$). A significant difference was also present between the moderate and high No-Go accuracy levels, with the learning effect being lower in the

high accuracy level (b = -3.24, SE = 0.685, p < .001). These results indicate that the magnitude of the statistical learning effect diminishes as No-Go accuracy increases. The three-way interaction between No-Go performance, ASRS score, and Block was also significant ($F_{(1, 12400.85)}$ = 8.10, p = .004). However, the post hoc test did not reach significance (p > .476). The three-way interaction between No-Go performance, ASRS score, and Triplet Type was also significant ($F_{(1, 13046.16)}$ = 4.36, p = .037). The post hoc test revealed that for participants with low ASRS scores, statistical learning (i.e., the difference between high- and low-probability trials) was gradually worse at higher levels of No-Go performance [low vs. mean: b = –4.73, SE = 1.03, p < .001; low vs. high: b = –9.46, SE = 2.05, p < .001; mean vs. high: b = –4.73, SE = 1.03, p < .001]. For participants who had mean ASRS scores, statistical learning also gradually declined at higher levels of No-Go performance [low vs. mean: b = –6.47, SE = 1.37, p < .001; low vs. high: b = –6.47, SE = 1.37, p < .001; mean vs. high: b = –3.24, SE = 0.69, p < .001]. However, for participants with high ASRS scores, statistical learning did not vary based on No-Go performance: [low vs. mean: b = –1.74, SE = 0.95, p = .068; low vs. high: b = –6.47, SE = 1.37, p = .068; mean vs. high: b = –1.74, SE = 0.95, p = .068] (Figure 4). Full results are presented in Supplementary Table 2, including b, SE b, 95% CI, t, df, and p values.

Figure 4. The interaction between response inhibition, ADHD-like symptoms, and statistical learning. This figure illustrates the three-way interaction among No-Go accuracy (as a measure of response inhibition), statistical learning, and ASRS scores (as a measure of ADHD-like symptoms). The x-axis represents No-Go accuracy, while the y-axis shows Go reaction time in ms. The two lines represent reaction times for high-probability triplets (blue line) and low-probability triplets (yellow line). The graph shows how the relationship between response inhibition and statistical learning is moderated by ADHD-like symptom severity. For participants with low or mean ASRS scores, statistical learning declined as No-Go accuracy increased. In contrast, for participants with high ASRS scores, statistical learning remained relatively stable across levels of No-Go accuracy.

Discussion

Our study reveals how ADHD-like traits shape the dynamic balance between statistical learning and inhibitory control in a large non-clinical sample of young adults. Whereas prior research has typically examined these functions in isolation or in clinical groups, we employed a paradigm that concurrently probed both processes, enabling us to capture their interaction across the full spectrum of ADHD-like traits. We found that higher ADHD-like traits were reliably associated with reduced inhibitory control, confirming a hallmark deficit of ADHD even in individuals below diagnostic thresholds. Critically, we uncovered an antagonistic relationship between inhibition and statistical learning: weaker inhibitory control was linked to enhanced learning of environmental regularities. However, this compensatory advantage was not uniform. Instead, it was progressively attenuated across the ADHD trait continuum and weaker among individuals with high symptom severity. These findings provide evidence that ADHD-like traits modulate not only isolated neurocognitive processes but also their interplay, highlighting a spectrum-based mechanism that may underlie the transition from adaptive variability to maladaptive behavior.

Our initial findings, showing that higher ADHD-like symptoms were associated with poorer response inhibition, align with a substantial body of research on the neural underpinnings of ADHD (Cortese, 2012; Cortese et al., 2012; Guo et al., 2020). Deficits in inhibitory control are consistently found in individuals with ADHD (Nigg et al., 2002; Senkowski et al., 2024), while other research has shown that ADHD-like traits can also impair inhibition (Polner et al., 2015). Our study confirms this connection in a non-clinical sample, which highlights the need to assess neurocognitive functions across the full ADHD spectrum. This would enable the early detection of these declines, as inhibitory deficits can emerge even in individuals without clinically

significant symptoms. Importantly, we demonstrate that this deficit persists even in a demanding paradigm that simultaneously assesses inhibition and statistical learning, in contrast to previous studies that typically assessed inhibition in isolation (Nigg et al., 2002; Silverstein et al., 2020). This indicates that the inhibitory impairment in ADHD-like traits is robust enough to manifest under more ecologically valid conditions where multiple cognitive systems are engaged in parallel.

While extensive research has examined the relationship between ADHD and various cognitive processes in isolation (Crosbie et al., 2013; Hart et al., 2014; Polner et al., 2015; Wodka et al., 2007), this study addressed a critical gap by investigating their dynamic interaction and how non-clinical ADHD-like traits modulate this. Our findings reveal a strong association between response inhibition and statistical learning, with poorer inhibition corresponding to enhanced statistical learning. Crucially, we revealed that ADHD-like traits significantly influence this relationship, as the learning advantage gradually diminishes across the symptom spectrum and becomes markedly reduced in individuals with high ADHD-like traits. These results reinforce the competition hypothesis (Ambrus et al., 2020; Horváth et al., 2022; Janacsek et al., 2012; Nemeth et al., 2013; Pedraza, Farkas, et al., 2024; Pedraza, Vékony, et al., 2024), which posits an antagonistic relationship between executive functions and statistical learning. While individuals with low ADHD-like traits align with this common pattern, the antagonism weakens as symptoms escalate, evolving into an altered form of interaction. This suggests that non-clinical ADHD symptoms play a significant role in shaping the interplay between neurocognitive processes. We propose that the behavioral manifestations of ADHD may not stem from isolated impairments but rather from a disruption in the dynamic interaction among different cognitive systems, which may contribute to maladaptive behavioral patterns. Future research should systematically examine this interplay to fully elucidate the underlying mechanisms of ADHD symptoms.

This study also highlights the advantages of a spectrum-based methodological approach. Such an approach has already revealed associations with multiple neurocognitive processes in ADHD (Crosbie et al., 2013; Moses et al., 2022; Panagiotidi et al., 2018) and allows for the detection of psychiatric traits before they reach clinical thresholds. This is particularly relevant for ADHD, where diagnostic criteria require that symptoms appear before the age of 12 (American Psychiatric Association, 2022). Adopting a spectrum perspective makes it possible not only to identify symptoms that persist from childhood without further causing functional impairment but

also to uncover undiagnosed difficulties originating in childhood and to detect adult-onset ADHD (Moffitt et al., 2015). Furthermore, the spectrum approach has proven similarly useful in capturing alterations in the interplay between neurocognitive functions associated with other non-clinical traits (Brezóczki et al., 2025; Colzato et al., 2025). A key question for future research is whether these neurocognitive alterations are trait- or disorder-specific or instead reflect a transdiagnostic general psychopathology factor (Caspi et al., 2014; Gluschkoff et al., 2019) and whether they represent qualitative differences or merely quantitative shifts along the continuum from subclinical to clinical expression.

Another angle of interpretation emerges from research on mind wandering, the spontaneous shift of attention from external demands to internally generated thought (Killingsworth & Gilbert, 2010; Mooneyham & Schooler, 2013; Smallwood & Andrews-Hanna, 2013). Mind wandering, defined as the unintentional shift of attention, has recently been proposed as a potential mechanism underlying ADHD symptoms and associated functional impairments, given its strong correlations with core symptom domains such as inattention (Allan Cheyne et al., 2009; Seli et al., 2015), impulsivity (Allan Cheyne et al., 2009; Arabacı & Parris, 2018), and hyperactivity (Shaw & Giambra, 1993). Research on mind wandering in ADHD has examined both clinical populations and subclinical traits in non-clinical samples, including college students, demonstrating a positive association between mind wandering tendencies and symptom prevalence (Arabacı & Parris, 2018; Biederman et al., 2019; Helfer et al., 2021; Jonkman et al., 2017; Seli et al., 2015). Since attention deficits in ADHD are thought to result from dysregulation in brain networks, with spontaneous streams of thought (mind wandering) emerging as a consequence (Bozhilova et al., 2018; Helfer et al., 2021). This relationship is particularly noteworthy, as mind wandering can influence not only statistical learning (Simor et al., 2025; Vékony, Farkas, et al., 2025), but also the interaction between executive control processes, such as inhibition, and statistical learning (Vékony, Brezóczki, et al., 2025). We can speculate that mind wandering may act as a transdiagnostic factor in ADHD, carrying important implications for both prevention and intervention strategies. On the one hand, it can disrupt executive functioning (Jonkman et al., 2017; Kane et al., 2007); yet in return, it may foster creativity and constructive thought processes (Baird et al., 2012; Mooneyham & Schooler, 2013), and it may also facilitate statistical learning that supports the development of skills and habits (Simor et al., 2025; Vékony, Farkas, et al., 2025). This dual perspective indicates that, although mind wandering entails maladaptive aspects, its

adaptive potential could be strategically leveraged in therapeutic approaches to ADHD (Biederman et al., 2017, 2019; Seli et al., 2015), highlighting the need for future research to elucidate how ADHD and mind wandering jointly shape interactions among neurocognitive processes.

Future directions

To build on the current findings, future research should use a wider variety of tests to better understand the relationship between ADHD-like traits and cognitive processes. A logical next step involves incorporating additional diagnostic and symptom-based measures, such as the Conners' Adult ADHD Rating Scales (CAARS) or the Wender Utah Rating Scale (WURS) (Macey, 2003; Ward et al., 1993). These tools would allow researchers to more precisely evaluate ADHD symptoms, including those from childhood, and provide a richer context for the current findings. To develop a more comprehensive cognitive profile, future studies should also measure other core executive functions commonly affected by ADHD, including working memory and cognitive flexibility. Investigating the interplay among these functions and their relationship to statistical learning and inhibitory control is a crucial next step. Additionally, by comparing individuals with subclinical traits to those with formal clinical diagnosis, we can gain a more comprehensive understanding of how these cognitive functions differ across the entire spectrum of symptomatic expression. Finally, the theoretical framework proposed by our research has broader implications beyond ADHD. Future research could explore whether this trait-based approach to cognitive function applies to other disorders that show high comorbidity with ADHD. This would be a crucial step toward developing a new theoretical framework for diagnosis and intervention that moves beyond traditional diagnostic boundaries and is better equipped to support individuals with varying levels of symptomatic severity.

Conclusion

Our study demonstrates that ADHD-like traits influence not only inhibitory control and learning in isolation but also their dynamic interaction, revealing a cognitive trade-off that is systematically altered across the symptom spectrum. This finding underscores the importance of moving beyond binary case–control frameworks toward dimensional models that capture variability in the general population. Cognitive alterations can emerge even in individuals below diagnostic thresholds, with implications for everyday functioning and vulnerability to maladaptive outcomes. Recognizing

ADHD as a spectrum condition opens new opportunities for trait-based approaches that emphasize early detection and proactive intervention. Rather than waiting for symptoms to escalate into a formal diagnosis, clinicians could leverage these insights to design targeted strategies that strengthen both executive control and learning processes, as well as their interaction. Such an approach would broaden access to support, reduce stigma, and facilitate personalized psychoeducation. More broadly, our findings highlight the need for a new theoretical framework that situates ADHD within a continuum of cognitive functioning, with direct relevance for transdiagnostic psychiatry and precision mental health.

Acknowledgement

This work was supported by the French National Grant Agency (ANR-22-CPJ1-0042-01 and ANR-24-CE37-5807) (to D.N.); the National Brain Research Program project NAP2022-I-2/2022 (to D.N.) and the EKÖP-24 University Excellence Scholarship Program of the Ministry for Culture and Innovation from the Source of the National Research, Development and Innovation Fund (EKÖP-24-3-II-ELTE-1159 (to B.B).

Contributions

K.H.: Data Curation, Formal analysis, Writing – Original Draft, Writing – Review & Editing, Visualization; B.B.: Conceptualization, Software, Validation, Investigation, Data Curation, Formal analysis, Writing – Original Draft, Writing – Review & Editing,, Funding; A.H.: Formal analysis, Writing – Review & Editing, T.V.: Conceptualization, Software, Validation, Investigation, Data Curation, Formal analysis, Visualization, Resources, Writing – Review & Editing, Supervision; D.N.: Conceptualization, Software, Validation, Resources, Writing – Review & Editing, Supervision, Funding.

Declaration of interests

The authors declare no competing interests.

Data availability statement

Data and codes for analysis are available on the following link: https://osf.io/evm2k/

References

Adi-Japha, E., Fox, O., & Karni, A. (2011). Atypical acquisition and atypical expression of memory consolidation gains in a motor skill in young female adults with ADHD. *Research in Developmental Disabilities*, *32*(3), 1011–1020. https://doi.org/10.1016/j.ridd.2011.01.048

Allan Cheyne, J., Solman, G. J. F., Carriere, J. S. A., & Smilek, D. (2009). Anatomy of an error: A bidirectional state model of task engagement/disengagement and attention-related errors. *Cognition*, *111*(1), 98–113. https://doi.org/10.1016/j.cognition.2008.12.009

Ambrus, G. G., Vékony, T., Janacsek, K., Trimborn, A. B. C., Kovács, G., & Nemeth, D. (2020). When less is more: Enhanced statistical learning of non-adjacent dependencies after disruption of bilateral DLPFC. *Journal of Memory and Language*, *114*, 104144. https://doi.org/10.1016/j.jml.2020.104144

American Psychiatric Association. (2022). *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association Publishing. https://doi.org/10.1176/appi.books.9780890425787

Anwyl-Irvine, A. L., Massonnié, J., Flitton, A., Kirkham, N., & Evershed, J. K. (2020). Gorilla in our midst: An online behavioral experiment builder. *Behavior Research Methods*, *52*(1), 388–407. https://doi.org/10.3758/s13428-019-01237-x

Arabacı, G., & Parris, B. A. (2018). Probe-caught spontaneous and deliberate mind wandering in relation to self-reported inattentive, hyperactive and impulsive traits in adults. *Scientific Reports*, *8*(1), 4113. https://doi.org/10.1038/s41598-018-22390-x

Aslin, R. N. (2017). Statistical learning: a powerful mechanism that operates by mere exposure. *WIREs Cognitive Science*, *8*(1–2). https://doi.org/10.1002/wcs.1373

Baird, B., Smallwood, J., Mrazek, M. D., Kam, J. W. Y., Franklin, M. S., & Schooler, J. W. (2012). Inspired by Distraction. *Psychological Science*, *23*(10), 1117–1122. https://doi.org/10.1177/0956797612446024

Balleine, B. W., & O'Doherty, J. P. (2010). Human and Rodent Homologies in Action Control: Corticostriatal Determinants of Goal-Directed and Habitual Action. *Neuropsychopharmacology*, *35*(1), 48–69. https://doi.org/10.1038/npp.2009.131

Barkley, R. A. (1997). Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. *Psychological Bulletin*, *121*(1), 65–94. https://doi.org/10.1037/0033-2909.121.1.65

Barnes, K. A., Howard, J. H., Howard, D. V., Kenealy, L., & Vaidya, C. J. (2010). Two Forms of Implicit Learning in Childhood ADHD. *Developmental Neuropsychology*, *35*(5), 494–505. https://doi.org/10.1080/87565641.2010.494750

Biederman, J., Fitzgerald, M., Uchida, M., Spencer, T. J., Fried, R., Wicks, J., Saunders, A., & Faraone, S. V. (2017). Towards operationalising internal distractibility (Mind Wandering) in adults with ADHD. *Acta Neuropsychiatrica*, *29*(6), 330–336. https://doi.org/10.1017/neu.2016.70

Biederman, J., Lanier, J., DiSalvo, M., Noyes, E., Fried, R., Woodworth, K. Y., Biederman, I., & Faraone, S. V. (2019). Clinical correlates of mind wandering in adults with ADHD. *Journal of Psychiatric Research*, *117*, 15–23. https://doi.org/10.1016/j.jpsychires.2019.06.012

Bozhilova, N. S., Michelini, G., Kuntsi, J., & Asherson, P. (2018). Mind wandering perspective on attention-deficit/hyperactivity disorder. *Neuroscience & Biobehavioral Reviews*, *92*, 464–476. https://doi.org/10.1016/j.neubiorev.2018.07.010

Brezóczki, B., Vékony, T., Farkas, B. C., Hann, F., & Németh, D. (2025). *Obsessive-Compulsive Tendencies Shift the Balance Between Competitive Neurocognitive Processes*. https://doi.org/10.1101/2025.08.13.669948

Bush, G., Valera, E. M., & Seidman, L. J. (2005). Functional Neuroimaging of Attention-Deficit/Hyperactivity Disorder: A Review and Suggested Future Directions. *Biological Psychiatry*, *57*(11), 1273–1284. https://doi.org/10.1016/j.biopsych.2005.01.034

Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., Meier, M. H., Ramrakha, S., Shalev, I., Poulton, R., & Moffitt, T. E. (2014). The p Factor. *Clinical Psychological Science*, *2*(2), 119–137. https://doi.org/10.1177/2167702613497473

Castellanos, F. X., Sonuga-Barke, E. J. S., Milham, M. P., & Tannock, R. (2006). Characterizing cognition in ADHD: beyond executive dysfunction. *Trends in Cognitive Sciences*, *10*(3), 117–123. https://doi.org/10.1016/j.tics.2006.01.011

Castellanos, F. X., & Tannock, R. (2002). Neuroscience of attention-deficit/hyperactivity disorder: the search for endophenotypes. *Nature Reviews Neuroscience*, *3*(8), 617–628. https://doi.org/10.1038/nrn896

Chambers, C. D., Garavan, H., & Bellgrove, M. A. (2009). Insights into the neural basis of response inhibition from cognitive and clinical neuroscience. *Neuroscience & Biobehavioral Reviews*, *33*(5), 631–646. https://doi.org/10.1016/j.neubiorev.2008.08.016

Colzato, L., Zhang, H., Roessner, V., Beste, C., & Hommel, B. (2025). Non-bivalent psychopathology: Rethinking mental disorders through metacontrol. *Neuroscience & Biobehavioral Reviews*, *176*, 106297. https://doi.org/10.1016/j.neubiorev.2025.106297

Conway, C. M. (2020). How does the brain learn environmental structure? Ten core principles for understanding the neurocognitive mechanisms of statistical learning. *Neuroscience & Biobehavioral Reviews*, *112*, 279–299. https://doi.org/10.1016/j.neubiorev.2020.01.032

Cortese, S. (2012). The neurobiology and genetics of Attention-Deficit/Hyperactivity Disorder (ADHD): What every clinician should know. *European Journal of Paediatric Neurology*, *16*(5), 422–433. https://doi.org/10.1016/j.ejpn.2012.01.009

Cortese, S., Kelly, C., Chabernaud, C., Proal, E., Di Martino, A., Milham, M. P., & Castellanos, F. X. (2012). Toward Systems Neuroscience of ADHD: A Meta-Analysis of 55 fMRI Studies. *American Journal of Psychiatry*, *169*(10), 1038–1055. https://doi.org/10.1176/appi.ajp.2012.11101521

Crosbie, J., Arnold, P., Paterson, A., Swanson, J., Dupuis, A., Li, X., Shan, J., Goodale, T., Tam, C., Strug, L. J., & Schachar, R. J. (2013). Response Inhibition and ADHD Traits: Correlates and Heritability in a Community Sample. *Journal of Abnormal Child Psychology*, *41*(3), 497–507. https://doi.org/10.1007/s10802-012-9693-9

Diamond, A. (2013). Executive Functions. *Annual Review of Psychology*, *64*(1), 135–168. https://doi.org/10.1146/annurev-psych-113011-143750

Farkas, B. C., Krajcsi, A., Janacsek, K., & Nemeth, D. (2023). The complexity of measuring reliability in learning tasks: An illustration using the Alternating Serial Reaction Time Task. *Behavior Research Methods*, *56*(1), 301–317. https://doi.org/10.3758/s13428-022-02038-5

Friedman, N. P., & Robbins, T. W. (2022). The role of prefrontal cortex in cognitive control and executive function. *Neuropsychopharmacology*, *47*(1), 72–89. https://doi.org/10.1038/s41386-021-01132-0

Gillan, C. M., Kosinski, M., Whelan, R., Phelps, E. A., & Daw, N. D. (2016). Characterizing a psychiatric symptom dimension related to deficits in goal-directed control. *ELife*, *5*. https://doi.org/10.7554/eLife.11305

Gillan, C. M., Otto, A. R., Phelps, E. A., & Daw, N. D. (2015). Model-based learning protects against forming habits. *Cognitive, Affective, & Behavioral Neuroscience*, *15*(3), 523–536. https://doi.org/10.3758/s13415-015-0347-6

Gluschkoff, K., Jokela, M., & Rosenström, T. (2019). The General Psychopathology Factor: Structural Stability and Generalizability to Within-Individual Changes. *Frontiers in Psychiatry*, *10*. https://doi.org/10.3389/fpsyt.2019.00594

Grahek, I., Leng, X., Musslick, S., & Shenhav, A. (2023). *Control adjustment costs limit goal flexibility: Empirical evidence and a computational account*. https://doi.org/10.1101/2023.08.22.554296

Greven, C. U., Merwood, A., van der Meer, J. M. J., Haworth, C. M. A., Rommelse, N., & Buitelaar, J. K. (2016). The opposite end of the attention deficit hyperactivity disorder continuum: genetic and environmental aetiologies of extremely low <scp>ADHD</scp> traits. *Journal of Child Psychology and Psychiatry*, *57*(4), 523–531. https://doi.org/10.1111/jcpp.12475

Guo, X., Yao, D., Cao, Q., Liu, L., Zhao, Q., Li, H., Huang, F., Wang, Y., Qian, Q., Wang, Y., Calhoun, V. D., Johnstone, S. J., Sui, J., & Sun, L. (2020). Shared and distinct resting functional connectivity in children and adults with attention-deficit/hyperactivity disorder. *Translational Psychiatry*, *10*(1), 65. https://doi.org/10.1038/s41398-020-0740-y

Hadas, I., Hadar, A., Lazarovits, A., Daskalakis, Z. J., & Zangen, A. (2021). Right prefrontal activation predicts ADHD and its severity: A TMS-EEG study in young adults. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *111*, 110340. https://doi.org/10.1016/j.pnpbp.2021.110340

Häger, L. A., Åsberg Johnels, J., Kropotov, J. D., Weidle, B., Hollup, S., Zehentbauer, P. G., Gillberg, C., Billstedt, E., & Ogrim, G. (2021). Biomarker support for ADHD diagnosis based on Event Related Potentials and scores from an attention test. *Psychiatry Research*, *300*, 113879. https://doi.org/10.1016/j.psychres.2021.113879

Hart, H., Chantiluke, K., Cubillo, A. I., Smith, A. B., Simmons, A., Brammer, M. J., Marquand, A. F., & Rubia, K. (2014). Pattern classification of response inhibition in ADHD: Toward the development of neurobiological markers for ADHD. *Human Brain Mapping*, *35*(7), 3083–3094. https://doi.org/10.1002/hbm.22386

Helfer, B., Boxhoorn, S., Songa, J., Steel, C., Maltezos, S., & Asherson, P. (2021). Emotion recognition and mind wandering in adults with attention deficit hyperactivity disorder or autism spectrum disorder. *Journal of Psychiatric Research*, *134*, 89–96. https://doi.org/10.1016/j.jpsychires.2020.12.059

Hengartner, M. P., & Lehmann, S. N. (2017). Why Psychiatric Research Must Abandon Traditional Diagnostic Classification and Adopt a Fully Dimensional Scope: Two Solutions to a Persistent Problem. *Frontiers in Psychiatry*, *8*. https://doi.org/10.3389/fpsyt.2017.00101

Herrmann, M. J., Saathoff, C., Schreppel, T. J., Ehlis, A.-C., Scheuerpflug, P., Pauli, P., & Fallgatter, A. J. (2009). The effect of ADHD symptoms on performance monitoring in a non-clinical population. *Psychiatry Research*, *169*(2), 144–148. https://doi.org/10.1016/j.psychres.2008.06.015

Hogarth, L., Chase, H. W., & Baess, K. (2012). Impaired goal-directed behavioural control in human impulsivity. *Quarterly Journal of Experimental Psychology*, *65*(2), 305–316. https://doi.org/10.1080/17470218.2010.518242

Horváth, K., Nemeth, D., & Janacsek, K. (2022). Inhibitory control hinders habit change. *Scientific Reports*, *12*(1), 8338. https://doi.org/10.1038/s41598-022-11971-6

Howard, J. H., & Howard, D. V. (1997). Age differences in implicit learning of higher order dependencies in serial patterns. *Psychology and Aging*, *12*(4), 634–656. https://doi.org/10.1037/0882-7974.12.4.634

Janacsek, K., Fiser, J., & Nemeth, D. (2012). The best time to acquire new skills: age-related differences in implicit sequence learning across the human lifespan. *Developmental Science*, *15*(4), 496–505. https://doi.org/10.1111/j.1467-7687.2012.01150.x

Jonkman, L. M., Markus, C. Rob., Franklin, M. S., & van Dalfsen, J. H. (2017). Mind wandering during attention performance: Effects of ADHD-inattention symptomatology, negative mood, ruminative response style and working memory capacity. *PLOS ONE*, *12*(7), e0181213. https://doi.org/10.1371/journal.pone.0181213

Kane, M. J., Brown, L. H., McVay, J. C., Silvia, P. J., Myin-Germeys, I., & Kwapil, T. R. (2007). For Whom the Mind Wanders, and When. *Psychological Science*, *18*(7), 614–621. https://doi.org/10.1111/j.1467-9280.2007.01948.x

Karatekin, C., White, T., & Bingham, C. (2009). Incidental and intentional sequence learning in youth-onset psychosis and attention-deficit/hyperactivity disorder (ADHD). *Neuropsychology*, *23*(4), 445–459. https://doi.org/10.1037/a0015562

KESSLER, R. C., ADLER, L., AMES, M., DEMLER, O., FARAONE, S., HIRIPI, E., HOWES, M. J., JIN, R., SECNIK, K., SPENCER, T., USTUN, T. B., & WALTERS, E. E. (2005). The World Health Organization adult ADHD self-report scale (ASRS): a short screening scale for use in the general population. *Psychological Medicine*, *35*(2), 245–256. https://doi.org/10.1017/S0033291704002892

Killingsworth, M. A., & Gilbert, D. T. (2010). A Wandering Mind Is an Unhappy Mind. *Science*, *330*(6006), 932–932. https://doi.org/10.1126/science.1192439

Kurzban, R., Duckworth, A., Kable, J. W., & Myers, J. (2013). An opportunity cost model of subjective effort and task performance. *Behavioral and Brain Sciences*, *36*(6), 661–679. https://doi.org/10.1017/S0140525X12003196

Lenth, R. (n.d.). *emmeans: Estimated Marginal Means, aka Least-Squares Means [Computer software]*. https://CRAN.R-project.org/package=emmeans.

Levy, F., Hay, D. A., McStephen, M., Wood, C., & Waldman, I. (1997). Attention-Deficit Hyperactivity Disorder: A Category or a Continuum? Genetic Analysis of a Large-Scale Twin Study. *Journal of the American Academy of Child & Adolescent Psychiatry*, *36*(6), 737–744. https://doi.org/10.1097/00004583-199706000-00009

Li, T., Mota, N. R., Galesloot, T. E., Bralten, J., Buitelaar, J. K., IntHout, J., AriasVasquez, A., & Franke, B. (2019). ADHD symptoms in the adult general population are associated with factors linked to ADHD in adult patients. *European Neuropsychopharmacology*, *29*(10), 1117–1126. https://doi.org/10.1016/j.euroneuro.2019.07.136

Li, X., Liu, X., Jiang, Y., Cui, J., Ji, Y., Cheng, F., & Zhou, D. (2025). Identifying neuroimaging biomarkers of attention-deficit hyperactivity disorder (ADHD) from cortical hemodynamic responses during Go/NoGo task using machine learning approaches. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *140*, 111417. https://doi.org/10.1016/j.pnpbp.2025.111417

Liston, C., Cohen, M. M., Teslovich, T., Levenson, D., & Casey, B. J. (2011). Atypical Prefrontal Connectivity in Attention-Deficit/Hyperactivity Disorder: Pathway to Disease or Pathological End Point? *Biological Psychiatry*, *69*(12), 1168–1177. https://doi.org/10.1016/j.biopsych.2011.03.022

Ma, J., Lei, D., Jin, X., Du, X., Jiang, F., Li, F., Zhang, Y., & Shen, X. (2012). Compensatory brain activation in children with attention deficit/hyperactivity disorder during a simplified Go/No-go task. *Journal of Neural Transmission*, *119*(5), 613–619. https://doi.org/10.1007/s00702-011-0744-0

Macey, K. (2003). Conners' Adult ADHD Rating Scales (CAARS). By C.K. Conners, D. Erhardt, M.A. Sparrow. New York: Multihealth Systems, Inc., 1999. *Archives of Clinical Neuropsychology*, *18*(4), 431–437. https://doi.org/10.1016/S0887-6177(03)00021-0

Martel, M., Nikola, M., & Nigg, J. T. (2007). Executive Function in Adolescents With ADHD. *Journal of the American Academy of Child & Adolescent Psychiatry*, *46*(11), 1437–1444. https://doi.org/10.1097/chi.0b013e31814cf953

Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The Unity and Diversity of Executive Functions and Their Contributions to Complex "Frontal Lobe" Tasks: A Latent Variable Analysis. *Cognitive Psychology*, *41*(1), 49–100. https://doi.org/10.1006/cogp.1999.0734

Moffitt, T. E., Houts, R., Asherson, P., Belsky, D. W., Corcoran, D. L., Hammerle, M., Harrington, H., Hogan, S., Meier, M. H., Polanczyk, G. V., Poulton, R., Ramrakha, S., Sugden, K., Williams, B., Rohde, L. A., & Caspi, A. (2015). Is Adult ADHD a Childhood-Onset Neurodevelopmental Disorder? Evidence From a Four-Decade Longitudinal Cohort Study. *American Journal of Psychiatry*, *172*(10), 967–977. https://doi.org/10.1176/appi.ajp.2015.14101266

Molitor, S. J., Oddo, L. E., Eadeh, H.-M., & Langberg, J. M. (2019). Executive Function Deficits in Adolescents With ADHD: Untangling Possible Sources of Heterogeneity. *Journal of Emotional and Behavioral Disorders*, *27*(3), 165–177. https://doi.org/10.1177/1063426618763125

Mooneyham, B. W., & Schooler, J. W. (2013). The costs and benefits of mind-wandering: A review. *Canadian Journal of Experimental Psychology / Revue Canadienne de Psychologie Expérimentale*, *67*(1), 11–18. https://doi.org/10.1037/a0031569

Moses, M., Tiego, J., Demontis, D., Bragi Walters, G., Stefansson, H., Stefansson, K., Børglum, A. D., Arnatkeviciute, A., & Bellgrove, M. A. (2022). Working memory and reaction time variability mediate the relationship between polygenic risk and ADHD traits in a general population sample. *Molecular Psychiatry*, *27*(12), 5028–5037. https://doi.org/10.1038/s41380-022-01775-5

Musslick, S., & Cohen, J. D. (2021). Rationalizing constraints on the capacity for cognitive control. *Trends in Cognitive Sciences*, *25*(9), 757–775. https://doi.org/10.1016/j.tics.2021.06.001

Natsheh, J., Natsheh, A. Y., Dobryakova, E., DeLuca, J., Khalil, S., Ashhab, M., & Herzallah, M. M. (2019). Characterization of Goal-Directed and Habitual Behavior in Children with Attention-Deficit/Hyperactivity Disorder. *SSRN Electronic Journal*. https://doi.org/10.2139/ssrn.3396047

Nemeth, D., Janacsek, K., Polner, B., & Kovacs, Z. A. (2013). Boosting Human Learning by Hypnosis. *Cerebral Cortex*, *23*(4), 801–805. https://doi.org/10.1093/cercor/bhs068

Nigg, J. T. (2001). Is ADHD a disinhibitory disorder? *Psychological Bulletin*, *127*(5), 571–598. https://doi.org/10.1037/0033-2909.127.5.571

Nigg, J. T., Butler, K. M., Huang-Pollock, C. L., & Henderson, J. M. (2002). Inhibitory processes in adults with persistent childhood onset ADHD. *Journal of Consulting and Clinical Psychology*, *70*(1), 153–157. https://doi.org/10.1037/0022-006X.70.1.153

Paloyelis, Y., Asherson, P., & Kuntsi, J. (2009). Are ADHD Symptoms Associated With Delay Aversion or Choice Impulsivity? A General Population Study. *Journal of the American Academy of Child & Adolescent Psychiatry*, *48*(8), 837–846. https://doi.org/10.1097/CHI.0b013e3181ab8c97

Panagiotidi, M., Overton, P. G., & Stafford, T. (2018). The relationship between ADHD traits and sensory sensitivity in the general population. *Comprehensive Psychiatry*, *80*, 179–185. https://doi.org/10.1016/j.comppsych.2017.10.008

Panagiotidi, M., Overton, P. G., & Stafford, T. (2019). Co-Occurrence of ASD and ADHD Traits in an Adult Population. *Journal of Attention Disorders*, *23*(12), 1407–1415. https://doi.org/10.1177/1087054717720720

Parks, K. M. A., & Stevenson, R. A. (2018). Auditory and Visual Statistical Learning Are Not Related to ADHD Symptomatology: Evidence From a Research Domain Criteria (RDoC) Approach. *Frontiers in Psychology*, *9*. https://doi.org/10.3389/fpsyg.2018.02502

Pedersen, A., & Ohrmann, P. (2018). Impaired Behavioral Inhibition in Implicit Sequence Learning in Adult ADHD. *Journal of Attention Disorders*, *22*(3), 250–260. https://doi.org/10.1177/1087054712464392

Pedraza, F., Farkas, B. C., Vékony, T., Haesebaert, F., Phelipon, R., Mihalecz, I., Janacsek, K., Anders, R., Tillmann, B., Plancher, G., & Németh, D. (2024). Evidence for a competitive relationship between executive functions and statistical learning. *Npj Science of Learning*, *9*(1), 30. https://doi.org/10.1038/s41539-024-00243-9

Pedraza, F., Vékony, T., Farkas, B. C., Haesebaert, F., Phelipon, R., Mihalecz, I., Janacsek, K., Tillmann, B., Anders, R., Plancher, G., & Nemeth, D. (2024). *The interplay between executive functions and updating predictive representations*. https://doi.org/10.1101/2024.12.05.626969

Polner, B., Aichert, D., Macare, C., Costa, A., & Ettinger, U. (2015). Gently restless: association of ADHD-like traits with response inhibition and interference control. *European Archives of Psychiatry and Clinical Neuroscience*, *265*(8), 689–699. https://doi.org/10.1007/s00406-014-0531-7

R Core Team. (2024). *R: A language and environment for statistical computing (Version 4.4.2) [Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/*.

Schippers, L. M., Greven, C. U., & Hoogman, M. (2024). Associations between ADHD traits and self-reported strengths in the general population. *Comprehensive Psychiatry*, *130*, 152461. https://doi.org/10.1016/j.comppsych.2024.152461

Seidman, L. J. (2006). Neuropsychological functioning in people with ADHD across the lifespan. *Clinical Psychology Review*, *26*(4), 466–485. https://doi.org/10.1016/j.cpr.2006.01.004

Seli, P., Smallwood, J., Cheyne, J. A., & Smilek, D. (2015). On the relation of mind wandering and ADHD symptomatology. *Psychonomic Bulletin & Review*, *22*(3), 629–636. https://doi.org/10.3758/s13423-014-0793-0

Senkowski, D., Ziegler, T., Singh, M., Heinz, A., He, J., Silk, T., & Lorenz, R. C. (2024). Assessing Inhibitory Control Deficits in Adult ADHD: A Systematic Review and Meta-analysis of the Stop-signal Task. *Neuropsychology Review*, *34*(2), 548–567. https://doi.org/10.1007/s11065-023-09592-5

Shaw, G. A., & Giambra, L. (1993). Task-unrelated thoughts of college students diagnosed as hyperactive in childhood. *Developmental Neuropsychology*, *9*(1), 17–30. https://doi.org/10.1080/87565649309540541

Shaw, P., Eckstrand, K., Sharp, W., Blumenthal, J., Lerch, J. P., Greenstein, D., Clasen, L., Evans, A., Giedd, J., & Rapoport, J. L. (2007). Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proceedings of the National Academy of Sciences*, *104*(49), 19649–19654. https://doi.org/10.1073/pnas.0707741104

Silverstein, M. J., Faraone, S. V., Leon, T. L., Biederman, J., Spencer, T. J., & Adler, L. A. (2020). The Relationship Between Executive Function Deficits and *DSM*-5-Defined ADHD Symptoms. *Journal of Attention Disorders*, *24*(1), 41–51. https://doi.org/10.1177/1087054718804347

Simor, P., Vékony, T., Farkas, B. C., Szalárdy, O., Bogdány, T., Brezóczki, B., Csifcsák, G., & Németh, D. (2025). Mind Wandering during Implicit Learning Is Associated with Increased Periodic EEG Activity and Improved

Extraction of Hidden Probabilistic Patterns. *The Journal of Neuroscience*, *45*(19), e1421242025. https://doi.org/10.1523/JNEUROSCI.1421-24.2025

Singmann, H., Bolker, B., Højsgaard, S., Fox, J., Lawrence, M. A., & Mertens, U. (2024). *afex: Analysis of Factorial Experiments [Computer software]*.

Smallwood, J., & Andrews-Hanna, J. (2013). Not all minds that wander are lost: the importance of a balanced perspective on the mind-wandering state. *Frontiers in Psychology*, *4*. https://doi.org/10.3389/fpsyg.2013.00441

Soetens, E., Melis, A., & Notebaert, W. (2004). Sequence learning and sequential effects. *Psychological Research Psychologische Forschung*, *69*(1–2), 124–137. https://doi.org/10.1007/s00426-003-0163-4

Takács, Á., Shilon, Y., Janacsek, K., Kóbor, A., Tremblay, A., Németh, D., & Ullman, M. T. (2017). Procedural learning in Tourette syndrome, ADHD, and comorbid Tourette-ADHD: Evidence from a probabilistic sequence learning task. *Brain and Cognition*, *117*, 33–40. https://doi.org/10.1016/j.bandc.2017.06.009

Theeuwes, J., Bogaerts, L., & van Moorselaar, D. (2022). What to expect where and when: how statistical learning drives visual selection. *Trends in Cognitive Sciences*, *26*(10), 860–872. https://doi.org/10.1016/j.tics.2022.06.001

Vékony, T. (2021). *Alternating Serial Reaction Time Task created with jsPsych (Version 1.0.0)*.

Vékony, T., Ambrus, G. G., Janacsek, K., & Nemeth, D. (2022). Cautious or causal? Key implicit sequence learning paradigms should not be overlooked when assessing the role of DLPFC (Commentary on Prutean et al.). *Cortex*, *148*, 222–226. https://doi.org/10.1016/j.cortex.2021.10.001

Vékony, T., Brezóczki, B., Csifcsák, G., Németh, D., & Simor, P. (2025). *A functional trade-off between executive control and implicit statistical learning is dynamically gated by mind wandering*. https://doi.org/10.1101/2025.08.05.668618

Vékony, T., Farkas, B. C., Brezóczki, B., Mittner, M., Csifcsák, G., Simor, P., & Németh, D. (2025). Mind wandering enhances statistical learning. *IScience*, *28*(2), 111703. https://doi.org/10.1016/j.isci.2024.111703

Vloet, T. D., Marx, I., Kahraman-Lanzerath, B., Zepf, F. D., Herpertz-Dahlmann, B., & Konrad, K. (2010). Neurocognitive Performance in Children with ADHD and OCD. *Journal of Abnormal Child Psychology*, *38*(7), 961–969. https://doi.org/10.1007/s10802-010-9422-1

Vogel, S. W. N., ten Have, M., Bijlenga, D., de Graaf, R., Beekman, A. T. F., & Kooij, J. J. S. (2018). Distribution of ADHD symptoms, and associated comorbidity, exposure to risk factors and disability: Results from a general population study. *Psychiatry Research*, *267*, 256–265. https://doi.org/10.1016/j.psychres.2018.06.017

Ward, M. F., Wender, P. H., & Reimherr, F. W. (1993). The Wender Utah Rating Scale: an aid in the retrospective diagnosis of childhood attention deficit hyperactivity disorder [published erratum appears in Am J Psychiatry 1993 Aug;150(8):1280]. *American Journal of Psychiatry*, *150*(6), 885–890. https://doi.org/10.1176/ajp.150.6.885

Wodka, E. L., Mark Mahone, E., Blankner, J. G., Gidley Larson, J. C., Fotedar, S., Denckla, M. B., & Mostofsky, S. H. (2007). Evidence that response inhibition is a primary deficit in ADHD. *Journal of Clinical and Experimental Neuropsychology*, *29*(4), 345–356. https://doi.org/10.1080/13803390600678046

Wright, L., Lipszyc, J., Dupuis, A., Thayapararajah, S. W., & Schachar, R. (2014). Response inhibition and psychopathology: A meta-analysis of go/no-go task performance. *Journal of Abnormal Psychology*, *123*(2), 429–439. https://doi.org/10.1037/a0036295