Pseudouridine selects RNAs for extracellular transport

Alessandro Scacchetti[1†], Emily J. Shields[1], Lauren N. Reich[1], John F. Doherty[1], Julia A. Tasca[1], Grace E. Lee[2,3], Richard Lauman[1], Natali L. Chanaday[4], Benjamin A. Garcia[5], Colin C. Conine[2,3], and Roberto Bonasio[1†]

[1]Epigenetics Institute and Department of Cell and Developmental Biology; University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA.

[2]Departments of Genetics and Pediatrics; Epigenetics Institute; Institute of Regenerative Medicine; Center for Research on Reproduction and Women's Health; University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA.

[3]Division of Neonatology; Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA.

[4]Departments of Physiology and Neuroscience; University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA.

[5]Department of Biochemistry and Molecular Biophysics; Washington University School of Medicine, St. Louis, Missouri, United States

[†]To whom correspondence should be addressed: roberto@bonasiolab.org; metalsga@gmail.com

ABSTRACT

RNAs move through the extracellular space to transmit information between cells, including mammalian neurons, yet how specific RNAs are channeled into these extracellular routes is unknown. Using genome-wide CRISPR screening, proteomics, and high-sensitivity transcriptomics in a neuronal cell line, we identify domesticated retroviral proteins and RNA-modifying enzymes that regulate RNA loading into and transportation via extracellular vesicles. We show that the pseudouridine synthase PUS1 is a key determinant of RNA trafficking, and that its catalytic product in RNA, pseudouridine, is both necessary and sufficient for extracellular RNA export. We further show that myosin light chain 6 (MYL6) is a pseudouridine-binding protein required for secretion of synthetic and endogenous RNAs. These findings reveal a biochemical code linking chemical RNA modification to extracellular transport, and establish a framework to study the function of extracellular RNAs in the nervous system and beyond.

INTRODUCTION

Cells exchange information to coordinate their actions. In addition to hormones, neurotransmitters, and other small molecules, RNAs also cross the plasma membrane and traverse the extracellular space to exert regulatory functions across cellular, tissue, and sometimes organismal boundaries (Betti et al., 2021; Cai et al., 2018; Kaletsky et al., 2020; Maori et al., 2019; Molnar et al., 2010; Tassetto, Kunitomi, & Andino, 2017). In mammals, such intercellular transfer of RNA has been associated with diverse physiological and pathological processes, including brain function and disease (Budnik, Ruiz-Canada, & Wendler, 2016; Huo, Du, Li, Liu, & Xu, 2021; Marangon, Castro, & Lecca, 2022; Z. Y. Wang, Wen, Xu, Zhang, & Zhang, 2022), cancer (Moller & Lobb, 2020) and intergenerational epigenetic inheritance (Conine & Rando, 2022), but its mechanistic basis remain obscure. Here, we investigate how RNAs are selected for extracellular export, a necessary step toward understanding what type(s) of information they communicate between cells, and determining whether and how this process affect phenotypic, transcriptional, and epigenetic states.

Extracellular RNAs (exRNAs) are often found within extracellular vesicles (EVs), membrane-based particles of heterogenous size, origin, and composition (van Niel, D'Angelo, & Raposo, 2018; Welsh et al., 2024), produced by most cell types (Deatherage & Cookson, 2012), and capable of transferring their cargo between cells (O'Brien, Breyne, Ughetto, Laurent, & Breakefield, 2020). Other structures, including membrane-free ribonucleoprotein complexes (Murillo et al., 2019; Tosar & Cayota, 2020) and virus-like particles (Ashley et al., 2018; M'Angale et al., 2025; Moore et al., 2021; Pastuzyn et al., 2018; Segel et al., 2021), can also transport RNA through the extracellular environment, although these processes are even less well understood than those mediated by canonical EVs.

The exRNAs produced by a given cell type do not recapitulate its intracellular RNA content (Cha et al., 2015; Mittelbrunn et al., 2011; Shurtleff et al., 2017; Upton et al., 2021; Valadi et al., 2007), suggesting the existence of molecular pathways that facilitate (select) or prevent (filter) the release of specific RNAs in the extracellular environment. Mechanisms for exRNA selection have been proposed, including sequence-specific sorting mediated by RNA-binding proteins (Garcia-Martin et al., 2022; Leidal et al., 2020; Santangelo et al., 2016; Shurtleff, Temoche-Diaz, Karfilis, Ri, & Schekman, 2016; Shurtleff et al., 2017; Villarroya-Beltri et al., 2013) and selective processing of exRNAs after their release (Costa et al., 2023; Nechooshtan, Yunusov, Chang, & Gingeras, 2020; Tosar et al., 2020). To date, mechanistic studies have focused on extracellular transport mechanisms for miRNAs, which are more easily captured with traditional small RNA

sequencing protocols (Gustafsson et al., 2025; Scacchetti et al., 2024)**,** and largely utilized non-
neuronal cell lines. However, miRNAs are relatively scarce in EVs (Chevillet et al., 2014), and it
is becoming increasingly clear that neurons might have specialized strategies to select and
release exRNAs (Ashley et al., 2018; Bervoets et al., 2024; M'Angale et al., 2025; Pastuzyn et al.,
2018). Hence, the full spectrum of mechanisms that participate in exRNAs selection and transport,
especially in neurons, remain unknown.

Here, we describe a novel genetic screen that uncovered new regulators of exRNA export in a
neuronal cell line. These include domesticated retroviral proteins and RNA-modifying enzymes.
Among the latter, we show that the pseudouridine synthetase 1 (PUS1) and its catalytic product,
pseudouridine (Ψ), are necessary and sufficient for the selection and extracellular release of
certain exRNAs, including full-length tRNAs and snoRNAs.

RESULTS

Neuronal CADs produce RNA-containing EVs

To investigate the genetic and biochemical determinants of neuronal exRNA biogenesis in a
tractable model, we selected Cath.a-differentiated cells (CADs), an immortalized cell line that
resembles catecholaminergic neurons and is capable of forming neurite-like structures under
serum-free conditions (Mews et al., 2017; Qi, Wang, McMillian, & Chikaraishi, 1997; H. Wang &
Oxford, 2000). We isolated EVs from conditioned, serum-free medium from CAD cultures using
an established serial ultracentrifugation method (**Fig. 1A**) (Thery, Amigorena, Raposo, & Clayton,
2006). This relatively crude EV fraction consisted primarily of particles ranging from 90 to 150 nm
in diameter (median 123 nm, **Fig. 1B**), consistent with the expected size range of small EVs
(Welsh et al., 2024). This fraction was enriched for known EV markers (SDCBP, PDCD6IP/ALIX,
TSG101, CD9), and depleted for conventional control mitochondrial and ER proteins (COXB5 and
CANX) (**Fig. 1C**) (Kugeratski et al., 2021). Overall, 81% of the proteins enriched in our EV isolate
also appeared in a curated database of EV components (Vesiclepedia, **Fig. S1A**) (Chitti et al.,
2024), and gene ontology (GO) annotations for the proteins we identified in CAD EVs were
enriched for relevant terms, including "*intracellular vesicle*" and "*endosome*" (**Fig. 1D**). Consistent
with the neuronal identity of CADs, GO terms associated with neurons ("*synaptic membrane*",
"*axon*") were also enriched, as well as specific neuronal EV markers L1CAM and ATP1A3 (You
et al., 2023) (**Fig. 1C**, **Table S1**).

In contrast to the intercellular RNA composition of CADs, their EVs contained predominantly small
RNA species (< 500 nts; **Fig. 1E**), as previously observed in other neuronal EVs (Bellingham,
Coleman, & Hill, 2012). Because conventional, ligation-based protocols to sequence small RNAs
were predominantly developed to detect miRNAs (Crocker, Trigg, & Conine, 2022; Gustafsson
et al., 2025), we used LIDAR sequencing (Scacchetti et al., 2024), which captures a broader
range of RNAs regardless of their size and chemical modifications, to obtain a comprehensive
profile of the exRNAs produced by CADs. LIDAR revealed 2,854 RNAs that were significantly
(adjusted *p* value < 0.1) enriched in EVs compared to whole-cell extract (**Fig. 1F**). Compared to
the intracellular transcriptome, CAD exRNAs were enriched for nuclear tRNAs, Y-RNAs, and
some snRNAs, and depleted for mitochondrial tRNAs and miRNAs (**Fig. 1G; S1B**). Some classes
of snoRNAs such as the H/ACA box family were enriched in EVs, whereas others, such as C/D
box snoRNAs were more abundant inside the cells (**Fig. S1C**). Careful inspection of the reads
mapping to tRNA genes revealed the presence in EVs of a mixture of full-length tRNAs and
various types of tRNA-derived fragments (tDRs) (**Fig. S1D–E**).

Thus, CADs produce EVs that contain known EV-associated proteins, as well as neuron-specific
proteins, and these EVs carry a variety of exRNAs, including full-length tRNAs and their
fragments.

Transduced sgRNAs are loaded into EVs

The fact that CAD EVs are enriched for specific RNAs when compared to their intracellular content
suggests the presence of dedicated pathways ensuring that only certain transcripts are selected
to become exRNAs. Potential mechanisms for both active and passive regulation of RNA
secretion have been proposed (Tosar, Witwer, & Cayota, 2021), but very few examples,
seemingly dedicated to the export of specific miRNAs, have been demonstrated (Garcia-Martin
et al., 2022; Liu, Ma, & Schekman, 2021; Shurtleff et al., 2017; Villarroya-Beltri et al., 2013). It is
unclear whether these bespoke pathways also affect other RNA classes, which constitute the
majority of exRNAs (**Fig. S1B**).

In search for general regulators of exRNA biogenesis, we perform a genome-wide knockout
screen. In a typical CRISPR/Cas9 screen, a library of single-guide RNAs (sgRNAs) targeting
genes of interest (or all genes in the case of genome-wide screens) is transduced into cells that
are then selected for a phenotype, for example drug resistance. Enriched or depleted sgRNAs
provide information on genes that regulate that phenotype and are identified by sequencing the
encoding transgenes integrated into the genome of the selected cells, thus linking genotypes to

phenotypes. This approach cannot be used for phenotypes not physically connected to the cell of
origin, such as EV abundance and composition.

We made the serendipitous observation that transduced sgRNAs could be detected, albeit at a
low level, in the crude EV fraction from CADs (**Fig. S2A**), therefore coupling the sgRNA genotype
of the originating cell to phenotypes associated with the sgRNA-containing EVs (**Fig. 2A, S2B**).
We reasoned that knockout of genes required for EV production or for loading RNAs into EVs
would result in the disappearance of the corresponding sgRNAs from the EV fraction, allowing us
to perform a genome-wide screen to identify genes involved in these processes.

We envisioned that all targeted genes and the corresponding sgRNAs would fall within 4 possible
categories (**Fig. 2B**): 1) sgRNAs not affecting viability, EV production, or RNA export ("no effect")
would remain constant over time in both cells and EVs; 2) sgRNAs targeting genes promoting
survival or proliferation would be depleted over time in both intracellular and extracellular pools
("essential"); 3) positive regulators of EV production or RNA export would become less abundant
in the EV-associated pool over time, while possibly accumulating within cells ("positive"); and 4)
vice versa for negative regulators ("negative").

A genome-wide screen for EV regulators

We transduced CADs with the genome-wide Brie library, comprising ~80,000 sgRNAs targeting
~20,000 protein-coding genes (**Fig. 2A**) (Sanson et al., 2018). Two days after infection, we
harvested a fraction of the cells and EVs to quantify baseline sgRNA distribution. Eight days later
(day 10 after transduction), we harvested cells and EVs again, and prepared libraries to sequence
the sgRNAs, using a modified LIDAR protocol (**Fig. S2C**). We performed the screen twice,
obtaining two independent experimental replicates.

Deep sequencing detected > 95% of sgRNAs in all samples (**Fig. S2D**). As early as day 2 after
transduction, likely before most knockout effects would manifest, not all sgRNAs were equally
represented in EVs vs. cells (**Fig. S2E**), indicating that some sgRNAs are more likely to be loaded
into EVs than others. This suggests that these exogenous RNAs are likely subjected to the same
selection processes that act on endogenous RNAs, potentially influenced by differences in
sequence, structure, or both. Consistent with this, sgRNAs that were preferentially exported as
exRNAs were characterized by a higher frequency of uridines at the 5' (**Fig. S2F**), raising the
possibility that these uridines, or perhaps their chemical modification, might contribute to the
selection process.

We analyzed the relative changes in abundance of sgRNAs in cells and EVs over time and sorted
the sgRNAs in the four categories described above (**Fig. 2C**). As expected, most sgRNAs were
neither enriched nor depleted ("no effect"), while sgRNAs targeting reported essential genes, such
as *Pcna*, *Cdk1*, and *Rrm1* were depleted at day 10 compared to day 2, both in cells and in EVs
("essential"). Single-guide RNAs targeting tetraspanins, *Cd63*, *Tspan6*, and *Tspan15* were
depleted at day 10 in EVs, but not inside the cells, where they were either unchanged or enriched.
This indicates that these genes might function as positive regulators of EV biogenesis ("positive"),
in line with their known role in membrane trafficking (Toribio & Yanez-Mo, 2022). Conversely, we
observed an enrichment of sgRNAs in EVs, but not in cells, ("negative") for two subunits of ESCRT
complexes, *Tsg101* and *Hgs* (*Hrs*), consistent with recent reports suggesting that they may
function as negative regulators for release of some EV subtypes (Kunitake et al., 2024; Matsui,
Osaki, Hiragi, Sakamaki, & Fukuda, 2021). *Vps51*, encoding a subunit of the GARP complex
(Conibear, Cleck, & Stevens, 2003), was also classified by our screen as a negative regulator,
likely due to its role in mediating retrograde trafficking from early endosomes to the Golgi
apparatus (Reggiori, Wang, Stromhaug, Shintani, & Klionsky, 2003).

For each sgRNA, we calculated a "predicted" abundance in EVs compared to cells at day 10
based on the export efficiency measured at day 2 and then determined deviations from this
predicted value in the "observed" sgRNA abundances. Overall, we identified 1,275 and 1,228
genes targeted by significantly (p value < 0.05) depleted or enriched sgRNAs (**Fig. 2D; Table**
S2), and therefore likely to function as positive or negative regulators of exRNA
selection/transport, respectively. Overall, the candidate positive regulators in our screen were
enriched for GO terms related to known EV biogenesis routes, such as "*late endosome*",
"*multivesicular body*", and "*extracellular vesicle*" (**Fig. 2E**), and comprised various genes involved
in the regulation of EV production and release (*Tspan6*, *Cd63*, *Flot1*, *Smpd3*, *Vps37c*, *Chmp2a*,
Chmp5, and *Chmp7*) (**Fig. 2D**). Among these, *Sdcbp* (**Fig. 2D; S2G**) stood out because of its
demonstrated role in EV biogenesis (Baietti et al., 2012), established use as marker for these
structures (Kugeratski et al., 2021), and abundance in CAD EVs (**Fig. 1C**). To confirm its function
in neuronal EVs, we generated two independent *Sdcbp*$^{-/-}$ CAD cell clones (**Fig. S2H**). Loss of
SDCBP caused reduced particle numbers indicative of impaired EV production, consistent with
the effects observed in other cell types (Baietti et al., 2012) (**Fig. S2I–J**).

As our screen utilized extracellular sgRNAs directly as a readout, the identification of *Sdcbp*
suggests that sgRNAs are exported, at least in part, via EVs originating from a variation of the
ESCRT pathway (Baietti et al., 2012). However, several genes coding for domesticated retroviral

proteins (*Arc, Moap1, Nynrin, Pnma8b, Rtl5, Rtl8b*) (Henriques et al., 2024) were also among the

top candidates for positive regulators of sgRNA export (**Fig. 2F**). Encapsulation of RNAs in

capsid-like structures could therefore represent an alternative route for exRNA export, potentially

one reserved for neuronal cells (Al Othman et al., 2024; Ashley et al., 2018; Madigan et al., 2024;

Pastuzyn et al., 2018; Xu et al., 2024), given that several of these genes are preferentially

expressed in the nervous system (Henriques et al., 2024). Consistent with neuronal-biased

expression, all of these candidates except *Moap1* (for which our RNA-seq was not sufficiently

sensitive) were upregulated upon differentiation of CADs into their mature neuronal phenotype

(**Fig. S2K**) (Mews et al., 2017; Qi et al., 1997).

Our screen also identified several positive regulators with no obvious link to membrane trafficking.

We reasoned that some of these factors might be involved in RNA selection or loading into the

EVs and that, therefore, their depletion would cause the disappearance of the corresponding

sgRNAs over time. Amongst them, we noted several enzymes responsible for installing or

removing chemical modifications on RNA, including pseudouridine (*Pus1, Rpusd2*) (Rintala-

Dempsey & Kothe, 2017), wybutosine (*Tyw1*) (Sun et al., 2024), acp(3)uridine (acp^{3}U) (*Dtwd1*)

(Takakura, Ishiguro, Akichika, Miyauchi, & Suzuki, 2019), mannosyl-queuosine (*Gtdc1*) (Zhao et

al., 2023), inosine (*Adarb1*) (J. B. Li & Walkley, 2025) , methyl groups (*Mrm2*) (Rorbach et al.,

2014), and amino acids (*Vars1, Tars1, Tars2, Dtd2, Prorsd1*) (Kuncha et al., 2018; Ruan, Zhou,

Tan, & Wang, 2013; Rubio Gomez & Ibba, 2020) (**Fig. 2G**). More broadly, we identified 224 RNA-

binding proteins amongst the positive exRNA regulators via RBP2GO analysis (Caudron-Herger,

Jansen, Wassmer, & Diederichs, 2021) (**Table S3**), in line with our expectations that factors

involved in the selection and loading of RNAs into EVs would be revealed by our genetic screen.

In summary, we devised a pooled genome-wide strategy to link genetic perturbation to regulation

of exRNA export. We identified several factors that could be promoting or restricting exRNA

selection and transport, including several domesticated retroviral genes, suggesting an

alternative, potentially neuron-specific mode of RNA secretion. Our screen also uncovered a

variety of RNA modifiers and RNA-binding proteins that could be involved in marking and

selecting specific RNAs for extracellular trafficking, as demonstrated below for PUS1.

PUS1 shapes the neuronal exRNA population

Among the RNA-modifying enzymes identified as candidate positive regulators of exRNAs, we

chose to focus on the pseudouridine (Ψ) synthase PUS1 (Rintala-Dempsey & Kothe, 2017),

because it was among the top 100 most significant positive regulators (**Fig. 2D, G; Table S2**) and

because genetic evidence supports a role for Ψ in RNA mobility in *Arabidopsis thaliana* pollen
(Herridge et al., 2025).

To validate the results of the screen, we utilized an orthogonal, pooled CRISPR interference
approach (CRISPRi) (Kampmann, 2018), whereby we knocked down genes of interest by
repressing their transcription with dCas9-KRAB guided by sgRNAs different from to those used
for the primary screen (**Fig. S3A**). CRISPRi of *Pus1* in CADs suppressed the secretion of sgRNAs
significantly ($p = 0.01$) and to a larger extent than the positive control *Sdcbp* (**Fig. S3B**), confirming
its involvement in exRNA export.

To determine the contribution of PUS1 to the export of endogenous RNAs, we profiled intra- and
extra-cellular RNAs in wild type CADs and two independent *Pus1$^{-/-}$* clones (**Fig. 3A**). Loss of
PUS1 resulted in a significantly (adjusted p value < 0.1) altered profile of 560 exRNAs, skewed
more than 2-fold toward reduced export (390 reduced vs. 170 increased exRNAs; **Fig. 3B**).
Changes in exRNA abundance were not secondary to altered levels of the corresponding
transcripts inside the cells, as shown by a negligible overlap in the affected RNA subsets (**Fig.
3C; S3C–E**). The export of miRNAs, snoRNAs, and both mitochondrial and nuclear tRNAs was
significantly impaired in the absence of PUS1 compared to wild type (**Fig. 3D**). Further inspection
of tRNA-mapping reads revealed that full-length and 5' and 3' tRNA-derived fragments (tDRs;
Fig. 3E) were exported less efficiently into EVs in absence of PUS1, with the full-length transcripts
being most strongly affected (**Fig. 3F**).

We conclude that PUS1 is necessary for the secretion of certain RNAs, including miRNAs,
snoRNAs, and tRNAs, in CADs.

Pseudouridine installed by PUS1 is enriched in exRNAs

Based on our observation that PUS1 is required for the release of a subset of exRNAs, we
hypothesized that its catalytic product, Ψ, might guide certain RNAs to be incorporated into EVs,
facilitating their export into the extracellular space.

To determine if exRNAs contain Ψ , we incorporated the Ψ-bisulfite adduct production step from
BID-seq (Dai et al., 2023) into our high-sensitivity LIDAR approach (Scacchetti et al., 2024). This
allowed us to identify Ψ through the appearance of microdeletions in the sequencing reads (**Fig.
4A**), as demonstrated on a synthetic RNA (**Fig. S4A**) and within endogenous 18S rRNA, which
contains several known Ψ residues (**Fig. S4B**) (Dai et al., 2023). With this approach, we detected
1,111 Ψ sites on 466 distinct exRNAs from CADs (**Fig. S4C–D**). As expected, most Ψ sites were

within tRNAs both inside and outside the cells, but RNAs from other classes also contained this modification, (**Fig. 4B**), consistent with existing observations (Schwartz et al., 2014).

The majority of exRNAs with detectable Ψ also carried this modification inside the cells (**Fig. S4C**), in most cases at the same exact sites (**Fig. S4D**); however, we observed higher Ψ levels on exRNAs compared to cellular RNAs (**Fig. 4C**), suggesting an association of the mark with their extracellular export.

Next, we sought to determine whether Ψ was also enriched on physiologically relevant exRNAs *in vivo*. We isolated RNA from EVs purified from the cauda section of the mouse epididymis, (**Fig. 4D; S4E**), where RNAs are transferred to the developing sperm via specialized EVs (epididymosomes) in a process necessary for proper sperm function (Trigg & Conine, 2024) and implicated in intergenerational inheritance (Conine & Rando, 2022). Using bisulfite-modified LIDAR, we identified 1,976 Ψ sites on 846 cauda exRNAs. As observed in CADs, cauda exRNAs contained higher levels of Ψ compared to their intracellular counterparts (**Fig. 4E**), demonstrating that Ψ is enriched on exRNAs not only *in vitro* (see **Fig 4C**), but also *in vivo*.

Our genetic evidence (**Fig. 2–3**) and sequencing results (**Fig. 4C, E**) support the hypothesis that conversion of U to Ψ by PUS1 on facilitates the secretion of select RNAs. To test this hypothesis, we sought to determine whether the loss of PUS1-dependent Ψ in CADs would cause decreased export of specific exRNAs. Because Ψ can also be deposited by other enzymes, we first identified uridines specifically modified by PUS1, by comparing bisulfite-induced deletions in exRNAs from *Pus1*[-/-] vs. wild type CADs and found 479 PUS1-dependent Ψ sites on 270 transcripts, accounting for 43% of the total Ψ sites on exRNAs (**Fig. 4F**). Decreased Ψ levels on these PUS1 targets did not affect their abundance inside the cells compared to RNAs containing Ψ residues unaffected by PUS1 loss (**Fig. 4G**, left). However, the abundance of PUS1 targets in the EVs from *Pus1*[-/-] CADs was significantly decreased (**Fig. 4G**, right) ($p = 0.008$) suggesting that Ψ was required for their efficient export.

The majority of exRNAs carrying PUS1-dependent Ψ were full-length tRNAs (**Fig. S4F–G**), which were targeted by PUS1 specifically at position 27 and 28 of the anticodon stem (**Fig. S4H–J**), as previously reported in other cell types (Schwartz et al., 2014). Of the 8 full-length tRNAs that were significantly depleted in EVs from *Pus1*[-/-] CADs, all but 1 (Fisher's test $p = 0.006$) contained a PUS1-dependent Ψ at position 27 or 28 (**Fig. S4K**), as exemplified for *tRNA-Arg-CCG-3-1*, *tRNA-Arg-CCT-4-1, tRNA-Ile-AAT-1-1,* and *tRNA-Val-CAC-2-1* (**Fig. 4H–I**)

Not all Ψ sites on these extracellular tRNAs were PUS1 sensitive, as shown by the persistence

of Ψ signal at other positions in *Pus1$^{-/-}$* KOs (**Fig. S4I**), most notably base 54, which is known to

be modified by PUS10 (**Fig. S4J**) (Schwartz et al., 2014). This indicates that the role of Ψ in the

export of these exRNAs is context-dependent, and that sequence or structural features of the

surrounding RNA (or possibly additional RNA modifications) contribute to its function.

We observed a similar exRNA export defect in *Pus1$^{-/-}$*-derived EVs for snoRNAs (**Fig. S4L**),

although in this case only 6 out of 13 snoRNAs containing PUS1-sensitive sites displayed an

export defect in the knockout cells, as exemplified by *Snord59a* and *Snord87* (**Fig. 4J–K**).

Together, these observations support the conclusion that selective Ψ deposition by PUS1 on

specific RNAs is necessary to guarantee their efficient extracellular export.

Ψ stimulates RNA secretion through MYL6

Given that Ψ is enriched on exRNAs and that PUS1 facilitates their export into the extracellular

space (see **Fig. 3–4**), we wondered if Ψ is sufficient to induce RNA secretion. We *in vitro*

transcribed 102 bp-long RNAs replacing all U's with Ψ. We transfected these Ψ-containing RNA

into CADs and measured their appearance in the extracellular space by RT-qPCR after 24 hours

(**Fig. 5A**). RNA containing Ψ was exported ~4 times more efficiently compared to the U-containing

control, despite their levels being the same inside the cells (**Fig. 5B**). We tested additional RNAs,

differing in sequence at their 5' (**Table S5**) and for most of them (4 out of 6 sequences tested) the

presence of Ψ resulted in significantly increased abundance outside the cells, with a fifth one

(RNA #6) also showing a clear trend toward increased export (**Fig 5C**). Overall, these results

indicate that the presence of Ψ on an RNA is sufficient, in most cases, to stimulate its extracellular

export.

We hypothesized that one or more RNA-binding protein might recognize Ψ-marked RNAs to

channel them into secretory pathways. We incubated biotinylated RNA containing Ψ or only U's

as control with CAD whole-cell extracts, purified the RNA with streptavidin, and analyzed

associated proteins via mass spectrometry. 46 proteins bound specifically to Ψ-containing RNA

compared to the U-only control (**Fig. S5A; Table S4**)**,** among which the myosin light chain 6

(MYL6) was the most enriched Ψ-specific binder (**Fig. 5D; Table S4**). MYL6 is an essential light

chain component of the non-muscle myosin II complex (Brito & Sousa, 2020), and it was

previously reported as a putative non-canonical RNA-binding protein (Caudron-Herger et al.,

2019).

Consistent with a potential role in exRNA trafficking, MYL6 was enriched in the proteome from CAD EVs (**Fig. 1C; Fig. S5B**) and sgRNAs against it were depleted in our screen, although slightly below our significance threshold ($p = 0.053$) (**Fig. S5C**). We knocked down MYL6 in CADs (**Fig. S5D**) and found that the Ψ-dependent secretion of 2 out of 4 RNAs was substantially impaired, with a third one also demonstrating a clear trend toward reduction (**Fig. 5E**). To determine whether MYL6 was also involved in the Ψ-dependent export of endogenous exRNAs we performed LIDAR on intra- and extra-cellular RNAs after MYL6 knockdown. The abundance of intracellular RNAs was largely unaffected (**Fig. S5E**); however, the export of specifically Ψ-containing exRNAs was significantly impaired by MYL6 knockdown (**Fig. 5F–G).**

Our results demonstrate that the presence of Ψ—a modification enriched on exRNAs *in vitro* and *in vivo*—is sufficient to stimulate exRNA export, and that, in many cases, Ψ-mediated exRNA sorting is an active process mediated by the non-canonical RNA-binding protein MYL6.

DISCUSSION

In this study, we combined biochemical, genetic, and genomic approaches to reveal a role for PUS1 and MYL6 in guiding the extracellular traffic of RNAs via Ψ. We propose a model whereby PUS1 labels select RNAs with Ψ at defined sites, which are then bound by MYL6 to be packaged into EVs and secreted (**Fig. 5H**). In addition to PUS1 and MYL6, our work points to many new candidate regulators that might play important roles in selecting (RNA-modifying enzymes) and transporting (domesticated retroviral proteins) RNAs from neurons and other cell types into the extracellular space.

A genome-wide screen for EV and exRNA regulators

We developed a CRISPR/Cas9 pooled genetic screen strategy that linked extracellular phenotypes to cellular genotypes via direct sequencing of secreted sgRNAs (**Fig. 2; S2**). With this design, we could identify not only factors that regulate EV biogenesis in general, but also factors required for the selection, loading, and transport of exRNAs into EVs, and potentially other non-canonical RNA-containing extracellular particles.

Two other screens aimed at separately addressing one or the other process have been published to date. In one case, artificial tethering of sgRNAs though an EV-specific anchor (CD63-dCas9 or CD9-dCas9) was used to create a readout for EV production (Kunitake et al., 2024), which limited the screen to the discovery of factors regulating the biogenesis of these specific EV sub-populations containing CD63 or CD9, and, by design, bypassed the factors required to load RNA

into the EVs. In the other, barcoded miRNAs containing the EXO motif (Villarroya-Beltri et al.,
2013) were used as reporters for miRNA secretion, revealing the role of ARHGEF18 in regulating
the release of miRNA-containing EVs (Lu et al., 2018). In this latter case, only factors involved in
movement of this specific subset of exRNAs could be recovered.

While these two studies provided important insights into EVs and miRNA transport, respectively,
our unbiased screen cast a broader net for general regulator of all extracellular particles
containing exRNAs, as revealed by the extensive number of "hits" already known to take part in
membrane trafficking (**Fig. 2D–E**), and by the large number of candidates involved in RNA
biology, such as RNA-modifying enzymes (**Fig. 2G**), that have not been previously linked to RNA
secretion.

Importantly, ours was the first genetic screen for an EV phenotype to be conducted on a neuronal
cell line, which, in addition to its unique and broader design, likely contributed to our ability to
detect several candidates not previously associated with exRNA trafficking and that might be
restricted—in expression or function—to the nervous system.

Routes of exRNA secretion in neurons

While we designed our genetic screen with the goal of uncovering specific regulators of exRNA
selection and export, the strategy allowed us to simultaneously identify several factors involved
in the biogenesis and trafficking of EVs as "by-products". The observation that SDCBP was
identified as positive regulator (**Fig. S2G–J**) suggests that sgRNAs (and other RNAs) taking the
extracellular route may be loaded into EVs formed through a variation of the canonical ESCRT
pathway, in which SDCBP acts in concert with ALIX to regulate the production of endosomal
intraluminal vesicles (Baietti et al., 2012). This is further supported by the presence of several
components of the ESCRTIII complex (CHMP2A, CHMP5, CHMP7) amongst the list of candidate
positive regulators (**Fig. 2D**). We also recovered factors involved in alternative modes of exRNA
secretion in neuronal cells, such as intraluminal vesicles produced by endosomal membrane
curvatures induced by ceramide in lipid rafts (Dixson, Dawson, Di Vizio, & Weaver, 2023), since
both SMPD3 (nSmase2, responsible for ceramide synthesis) and FLOT1 (enriched in lipid rafts)
appeared amongst our positive regulators of exRNAs (**Table S2**).

In line with the neuronal identity of CADs, but somewhat unexpectedly, we detected synaptic
components in our EV preparation (**Fig. 1D; Table S1**) and some of them (e.g.: CXADR, GNAO1,
CNR1, ADAM10) were also identified as exRNA regulators in our screen (**Fig. 2E; Table S2**).
This suggests that in mammalian neurons, some EVs are produced at the presynaptic membrane,

as observed in *Drosophila* (Blanchette et al., 2022; Korkut et al., 2009), indicating the existence

of a dedicated, alternative route for neuronal exRNAs. Interestingly, fly and mammalian neurons

have been shown to package RNAs for secretion in structures resembling viral capsids originating

from domesticated retroviral proteins (Erlendsson et al., 2020; M'Angale et al., 2025; Pastuzyn et

al., 2018), whose intercellular exchanges appears important for synaptic function, as well as

learning and memory formation (Sullivan, Ravens, Walker, & Shepherd, 2025).

Among potentially neuron-specific regulators of exRNA trafficking, domesticated retroviral

proteins stand out as particularly intriguing, with six recovered in our screen (**Fig. 2F**). One of

these, ARC, is a GAG homolog that assembles into a virus-like particle that transports its own

mRNA across the plasma membrane in humans and flies (Al Othman et al., 2024; Ashley et al.,

2018; Pastuzyn et al., 2018). Several additional genes with homology to the retroviral GAG capsid

protein or other retroviral components are under purifying selection in mammalian genomes

(Henriques et al., 2024) and their expression is often high in the nervous system, hinting at the

tantalizing possibilities that brains have hijacked viral machinery for the purpose of RNA-based

communication.

Biochemical rules of exRNA selection

The most obvious class of candidate positive regulators of exRNAs revealed by our screen were

RNA-modifying enzymes (**Fig. 2G**). Among these, we chose to focus on PUS1 (see below);

however, not all pseudouridylated exRNAs depend on PUS1 for their secretion, and many

exRNAs do not contain Ψ. We suspect that other RNA modifying enzymes contribute—

independently or in concert with PUS1—to shape the exRNA population in neurons and other cell

types.

Among these other candidates, we note DTWD1, an enzyme responsible for acp^{3}U deposition on

tRNAs and possibly other substrates (Takakura et al., 2019). acp^{3}U is an attachment site for N-

glycans (Xie et al., 2024), and glycosylation of some RNAs seems to be important for their

secretion through EVs (Sharma, Jiao, Yang, Kwan, & Kiledjian, 2025), suggesting that glycans

could serve as alternative exRNA selection marks. Furthermore, we identified ALKBH6 (**Table

S2**), a poorly characterized protein predicted to have RNA demethylase activity (Ma et al., 2022),

as one of the top negative regulators (i.e.: factors that restrict exRNA export). A related RNA

demethylates ALKBH5 was recently shown to prevent RNA secretion via m6A removal (Chen et

al., 2025), and we speculate ALKBH6 may function similarly as a "filter" for exRNA export via

modulation of RNA methylation on yet unknown RNA substrates.

Our screen also uncovered hundreds of RNA-binding proteins that may regulate RNA secretion

without modifying them enzymatically (**Table S3**). Among these, we identified VAP-A, an

important regulator of exRNA export, previously reported to load RNAs into intraluminal vesicles

at contact sites between endoplasmic reticulum and multivesicular bodies (Barman et al., 2022).

We also identified ANXA7, a protein involved in regulating membrane fusion during exocytosis

(Caohuy & Pollard, 2002) and recently reported as an RNA binder (Tartaglia, Hollas, Havik,

Vedeler, & Pastore, 2025). Given its enrichment in CAD EVs (**Table S1**) and its reported

interaction with ESCRT-III (Sonder et al., 2019), we postulate it may additional function as an

RNA cargo loader during EV formation. On the other hand, we failed to identify some other RNA-

binding proteins previously shown to modulate exRNA export, such as ALYREF, FUS, ATG7/12,

SYNCRIP, YBX1, and HNRNPA2B1(Garcia-Martin et al., 2022; Leidal et al., 2020; Santangelo et

al., 2016; Shurtleff et al., 2016; Shurtleff et al., 2017; Villarroya-Beltri et al., 2013). This could be

due to ineffectiveness of sgRNAs, reporter RNA, or cell type used. In fact, we deem it likely that

different cell types utilize different sets of RNA-binding proteins and RNA modifications to guide

the secretion of different classes of RNAs.

A role for Ψ on exRNAs

Ψ is the most abundant RNA modification found in cells, and it is known for its roles in RNA

stability, translation, and splicing (Cerneckis, Cui, He, Yi, & Shi, 2022). However, the very strong

depletion of sgRNAs against PUS1 in our screen, as well as some indirect evidence in other

organisms motivated us to explore a potential role in exRNAs. In *Arabidopsis*, Ψ appears to

participate to the transfer of small RNAs from pollen to sperm cells (Herridge et al., 2025). In

human colorectal cancer cells, decreased Ψ levels caused by 5-fluorouracil treatment were linked

to alteration of extracellular small RNA profiles (Qu et al., 2024). In line with these reports, we

found Ψ to be 1) enriched in exRNAs both from CAD cell cultures and mouse tissue, 2) sufficient

for export of most synthetic RNAs tested, and 3) required for the release of endogenous RNA,

especially tRNAs and snoRNAs, into the extracellular space.

How does Ψ promote RNA secretion? We propose that the conversion of U to Ψ creates new

binding sites for specific proteins—Ψ readers. We identified MYL6, part of the non-muscle myosin

II complex, as a putative new Ψ reader and demonstrated its functional involvement in RNA export

of synthetic and endogenous exRNAs marked by Ψ **(Fig. 5)**. Intriguingly, other cytoskeletal

proteins, such as the actin-binding protein PFN1, have been reported to interact with Ψ (Wei et

al., 2025). RNAs containing Ψ might thus "hitchhike" cytoskeletal components to be loaded into

vesicles and eventually secreted, a pathway already proposed for other vesicular cargo such as

actin-associated proteins (Holliday, Faria, & Rody, 2019). It is worth noting, however, that not all

Ψ-containing synthetic and endogenous exRNAs were sensitive to MYL6 depletion **(Fig. 5G)**.

While this could be attributed to experimental limitations (such as incomplete knockdown), it is

also possible that other Ψ readers are involved, perhaps in an RNA sequence- or structure-

dependent fashion.

PUS1 is a conserved eukaryotic standalone pseudouridine synthetase which has been

extensively characterized in *S. cerevisiae* (Rintala-Dempsey & Kothe, 2017), where it is thought

to be involved in the export of specific tRNA from the nucleus to the cytosol (Grosshans, Lecointe,

Grosjean, Hurt, & Simos, 2001). In mammals, PUS1 localizes both to the nuclei and mitochondria

(Patton, Bykhovskaya, Mengesha, Bertolotto, & Fischel-Ghodsian, 2005). Our work suggests that

PUS1-mediated Ψ has an additional, fundamental function: guiding the export of exRNAs.

Knockout of PUS1 in neuronal CADs resulted in a significant loss of Ψ from many RNAs and,

importantly, in non-overlapping effects on cellular and extracellular RNA pools (**Fig. 3A–D**). PUS1

loss significantly decreased secretion of full-length tRNAs (**Fig. 3F**), likely due to the loss of PUS1-

dependent Ψ at position 27 and 28 in the anticodon stem. Interestingly, other Ψ sites remain

intact, indicating that modification at these other positions may not be sufficient for export (**Fig.**

4H–I) and that the export-related function of Ψ depends on sequence or structural context.

Conclusions and outlook

In conclusion, our screen in neuronal CADs discovered a new pathway for RNA secretion,

whereby PUS1 catalyzes the formation of Ψ on select RNAs, marking them for export via EVs,

through recognition by MYL6. The genetic identification and biochemical validation of this and

other mechanisms governing exRNA selection and transport will enable deeper mechanistic

investigation of their downstream biological functions, and, ultimately, help decode the language

of RNA-mediated cell-to-cell communication.

EXPERIMENTAL PROCEDURES

Cell culture and conditioned medium collection

CADs (kind gift from S. Berger) (Mews et al., 2017) were maintained in DMEM/F-12 medium (Gibco, Cat. No 11320033) supplemented with 10% ES-grade fetal bovine serum (Gibco, Cat. No 16141079), 1% L-Glutamine (Sigma-Aldrich, Cat. No G7513), and 0.5% penicillin/streptomycin solution (Sigma-Aldrich, Cat. No P0781). EV-depleted fetal bovine serum was generated by centrifugation at 100,000 g (28,500 rpm) on a SW-32Ti swinging bucket rotor (Beckman Coulter, Cat. No G369650) for 4 h at 4°C, followed by filtration through a 0.22 μm pore size filter (Sigma-Aldrich, Cat. No S2GPU10RE). For EV collection, cells were seeded at 3–15 million per 15-cm dish and washed twice with 1X PBS. Cells were incubated in serum-free medium (same as maintenance medium but without fetal bovine serum) for 48 h (proteomics) or for 24 h (all other analyses) before collecting EVs. For collection of EVs for pooled CRISPR screen (see below), cells were incubated in EV-depleted medium (same as maintenance medium but with EV-depleted fetal bovine serum) for 24 h. To obtain differentiated CADs, cells were maintained in serum-free medium for 5 days.

Extracellular vesicles (EV) isolation and nanoparticle tracking analysis

EV isolation was performed via differential centrifugation, as previously described (Thery et al., 2006) with minor modifications. Briefly, conditioned medium was centrifuged at 300 g for 5 min at 4°C, followed by centrifugation at 2,000 g for 10 min at 4°C and then 10,000 g for 30 min at 4°C. Then, the cleared conditioned medium was passed through a 0.22 μm pore size filter (Millipore-Sigma, Cat. No SCGP00525) and centrifuged at 100,000 g (28,500 rpm) on a SW-32Ti swinging bucket rotor for 2 h at 4°C. EV pellet was resuspended in 1X PBS and centrifuged again at 100,000 g (28,500 rpm) on a SW-32Ti swinging bucket rotor for 2 h at 4°C. The washed EV pellet was then resuspended in 100 μL of 1 X PBS and either used immediately or stored at -80°C.

For nanoparticle tracking analysis (NTA), EVs were diluted in deionized water and analyzed using the ZetaView PMX220 Twin (Patricle Metrix) with the following parameters: laser wavelength = 488 nm; filter wavelength = scatter; shutter = 100; sensitivity = 80. NTA was performed at the extracellular vesicle core at the University of Pennsylvania.

Animals

Male FVB/NJ (8–12 weeks old) were derived from breeders obtained from Jackson Laboratory (Strain #:001800) and maintained in-house. All animal procedures were in strict accordance with

the guidelines of the Children's Hospital of Philadelphia and the University of Pennsylvania

Institutional Animal Care and Use Committee regulations (CHOP IACUC Protocol #23-001364,

UPenn PSOM IACUC Protocol #806911).

Epidydimal tissue and EV collection

Mice were sacrificed by cervical dislocation, and epididymal tissue carefully dissected free of fat

and connective tissue. The cauda epididymis was separated and immersed in pre-warmed (37°C)

Biggers, Whitten, and Whittingham (BWW) medium (Cafe, Anderson, & Nixon, 2020). Seminal

fluid was gently expressed from the tissue. The tissue was removed from the media and flash

frozen in a microcentrifuge tube and kept at -80°C until processing for RNA isolation. The released

spermatozoa were incubated in BWW medium at 37 °C for 10 min to allow motile sperm to

disperse. The suspension was transferred to a fresh microcentrifuge tube and incubated for an

additional 10 min at 37 °C to enrich for motile sperm by swim-up. The upper fraction of the

suspension was carefully aspirated, leaving ~50 µL of residual medium behind. Spermatozoa

were pelleted by centrifugation (10,000 g, 5 min), and the supernatant was centrifuged twice at

10,000 g for 5 min to remove residual tissue and sperm, each time collecting only the supernatant.

A final spin at 10,000 g for 30 min at 4 °C was performed to further eliminate debris. The resulting

supernatant, containing EVs, was subjected to ultracentrifugation (Optima Max TL, Beckman

Coulter) at 120,000 g for 2 h at 4 °C. The pellet was washed in 1 mL cold PBS and ultracentrifuged

again at 120,000 g for 2 h at 4 °C. The final epididymosome pellet was resuspended in 30 µL cold

PBS, flash frozen in liquid nitrogen, and stored at -80 °C until RNA isolation.

Cells and EV proteomics

EVs and cells were resuspended in freshly prepared, ice-cold urea lysis buffer (50 mM Tris pH 8,

8M urea) and incubated for 1 h at 4°C. Lysates were then sonicated using Bioruptor (Diagenode)

with 5 cycles of 30s on / 30 s off and then cleared by centrifugation at 18,000 g for 10 min at 4°C.

Extract was quantified using the Micro BCA Protein Assay Kit (Thermo Scientific, Cat. No 23235)

and 50 µg were taken for further processing. Proteins were prepared using Protifi S-trap Micro

columns per manufacturer's instructions. Following S-trap elution, peptides were lyophilized,

resuspended in 0.1 % formic acid for further cleanup by desalting. Desalted peptides were

lyophilized and resuspended in 4% acetonitrile containing 0.1% formic acid for LC-MS analysis.

Mass spectra for peptides were obtained with a Dionex Ultimate3000 liquid chromatograph

(Thermo Fisher Scientific) coupled to a Thermo Fusion mass spectrometer (Thermo Fisher

Scientific). Peptides were loaded onto an in-house silica capillary column (75 µM inner diameter)

packed with C18 resin (ReproSil-Pur C18-AQ 2.4-µm resin; Dr. Maisch GmbH, Ammerbuch,

Germany). Peptides were eluted with a gradient of 2%–25% B (86 min) followed by a gradient of

25%–50% (32 min) and a hold at 98% B (15 min) (mobile phase A: 0.1% formic acid, mobile

phase B: 80% acetonitrile, 0.1% formic acid). Flow rate was 300 nL/min. Data were collected in

data-dependent acquisition mode. Full-scan MS settings were as follows: scan range, 300–1,100

(m/z; mass-to-charge ratio); resolution, 60,000; AGC, on; micro scan count, 1; Maximum IT, 54

ms. MS2 settings were as follows: resolution, 30,000; AGC, predicted; micro scan count, 1;

Maximum IT, 100 ms; fragmentation was enforced by higher-energy collisional dissociation with

normalized collision energy of 29; isolation width, 1.8 m/z. MS/MS spectra were processed using

MetaMorpheus version (https://github.com/smith-chem-wisc/MetaMorpheus) using a UniProt *Mus*

musculus reference proteome and common contaminants. The following search settings were

used: protease = trypsin; search for truncated proteins and proteolysis products = false; maximum

missed cleavages = 2; minimum peptide length = 7; maximum peptide length = 60; initiator

methionine behavior = variable; fixed modifications = carbamidomethyl on C, carbamidomethyl

on U; variable modifications = oxidation on M; max mods per peptide = 1; max modification

isoforms = 1,024; precursor mass tolerance = ±10.0000 PPM; product mass tolerance = ±0.0200

absolute; report PSM ambiguity = true. Results were filtered for score (> 5) and FDR (1 %).

RNA extraction

Cell pellets or purified EVs were resuspended in 1 mL of TriPure (Roche, Cat. No 11667165001)

and RNA was extracted following the recommended protocol. DNA was digested with Turbo

DNase-I (Invitrogen, Cat. No AM2238) at 37˚C for 30 min, RNA was re-purified using TriPure and

resuspended in modBTE (10mM Bis-Tris pH 6.7, 0.1 mM EDTA). For RNA extraction directly from

cleared conditioned medium, 0.3 M NaOAc (final) and 1 volume of phenol-chloroform-IAA

(Ambion, Cat. No AM9732) were added and RNA was purified using the recommended protocol.

DNA was digested and RNA was re-purified using TriPure as described above.

LIDAR, small RNA-seq, and mRNA-seq libraries

LIDAR libraries from 100–250 ng of cellular RNA or 1–50 ng of EV RNA were prepared as

described (Scacchetti et al., 2024) with some minor modification. LIDAR_RT_primer and

LIDAR_TSO_mix were diluted 1:3 for samples where RNA input was < 10 ng. Reverse

transcription conditions were changed to the following: 8°C for 10 min, 16°C for 10 min, 25°C for

10 min, 42°C for 80 min, 10 cycles at 50˚C for 2 min and 42˚C for 2 min, 85°C for 5 min. Lastly,

the amount of indexing primers was increased to 10 pmol. LIDAR libraries were analyzed on a

2% agarose gel and quantified using Qubit dsDNA Quantification Assay kit (Invitrogen, Cat. No
Q32854).

Small RNA libraries were prepared using the NEBnext small RNA library kit for Illumina (New
England Biolabs, Cat. No E7330S), following the standard protocol with the following parameters:
1) 1:2 dilution of 3'SR Adaptor, SR RT Primer, and 5'SR Adaptor, 2) 15 indexing PCR cycles, 3)
cleanup of final libraries with QIAgen MinElute PCR purification kit (QIAgen, Cat. No 28004),
followed by a second cleanup with 2X SPRI. Libraries were analyzed on 6% non-denaturing
polyacrylamide gels.

Conventional mRNA-seq libraries were prepared from 2 μg of total CAD RNA following the
protocol described in (Scacchetti et al., 2024).

Libraries were sequenced on Illumina NextSeq1000 or NextSeq2000 instruments.

Genome-wide pooled CRISPR-Cas9 screen

The mouse Brie sgRNA library (Doench et al., 2016) in the lentiCRISPR v2 and lentiGuide-Puro
backbones were obtained from Addgene (Cat. No 73632 and 73632). The library were amplified
in ElectroMAX Stbl4 competent cells (Invitrogen, Cat. No 11635018). To generate lentiviruses,
~100 million HEK293FT were transfected with 275 μg of amplified library together with 170 μg of
psPAX2 packaging plasmid (Addgene, Cat. No 12260) and 96 μg of pMDG.2 envelope plasmid
(Addgene, Cat. No 12259) using Lipofectamine 3000 Transfection Reagent (Invitrogen, Cat. No
L3000015). 18 h after transfection, the medium was changed to lentivirus_collection_medium
[DMEM high glucose (Gibco, Cat. No 11965126) supplemented with 10% ES-grade fetal bovine
serum (Gibco, Cat. No 16141079), 1% GlutaMAX (Gibco, Cat. No 35050061), 1% MEM non-
essential amino acid solution (Sigma-Aldrich, Cat. No M7145-100ML), 1% sodium pyruvate
(Millipore-Sigma, Cat. No S8636), and 1% bovine serum albumin (Sigma-Aldrich, Cat. No A1933).
Conditioned medium was collected 48 h later, filtered through a 0.45 μm filter (Millipore-Sigma,
Cat. No S2HVU11RE), and concentrated using Amicon Ultra centrifugal filters with 100 kDa
MWCO (Millipore-Sigma, Cat. No UFC910008). Concentrated lentiviruses were aliquoted and
stored at -80°C. Multiplicity of infection (MOI) in CADs was estimated as described (Shalem et
al., 2014). Briefly, CADs were infected with increasing amounts of lentivirus using 8 μg/mL
polybrene (final concentration). 24 h after infection, cells were divided equally into two new wells
and one of them was treated with 1 μg/mL puromycin (Invivogen, Cat. No ant-pr-1). 96 h later,
cells were counted, and the MOI was determined as the percentage of surviving cells. To test for
sgRNA presence in EV preparations, CADs were infected with the Brie lentiGuide-Puro (sgRNA

only) libraries at an MOI of 0.43 and ~1X coverage, selected with 1 µg/ml puromycin for 7 days,

and stably transduced cells were frozen in liquid N_2. RT-qPCR for sgRNA expression was

performed using Power SYBR Green RNA-to-C_T (Applied Biosystems, Cat. No 4389986) with the

sgRNA_1_qPCR primers (**Table S5**).

The genome-wide screen was performed in 2 biological replicates. 55 million CADs were

transduced on day 0 with the lentiviral library at an MOI of 0.21 using 8 µg/mL polybrene to result

in ~100-150X sgRNA coverage. 24 h after infection, cells were washed twice with PBS and EV-

depleted medium was added. 24 h later ("day 2" throughout the manuscript), EVs and half the

cells were collected as described above and resuspended in TriPure. The other half of the cells

was re-plated with maintenance media containing 1 µg/mL puromycin. Cells were selected with

puromycin for 7 days. Then, cells were washed twice with PBS and EV-depleted medium was

added. 24 h later ("day 10" throughout the manuscript), EVs and cells were collected once again

and resuspended in TriPure. RNA was extracted from all samples as described. For cell samples,

< 200 nts RNAs were enriched using Zymo RNA Clean & Concentrator – 25 (Zymo Research,

Cat. No. R1017) following the recommended protocol.

sgRNA-seq libraries were prepared using a modified LIDAR protocol starting from half of the total

amount of RNA obtained from cells and EVs. In the case of cells, multiple parallel libraries were

prepared to not exceed 1.5 µg of input RNA / library, which were then pooled before sequencing.

Briefly, RNAs was diluted to 2.8 µL and mixed with 0.4 µL of 10 µM LIDAR_sgRNA_RT_primer

(see **Table S5** for all oligonucleotide sequences). Samples were heated to 70°C for 2 min, then

cooled to 4°C (0.5°C/s). Next, 4.8 µL of RT_mix [25 mM Tris-HCl pH 8.3, 20 mM NaCl, 2.5 mM

$MgCl_2$, 8 mM DTT, 5% PEG-8000, 0.5 mM dNTPs, 1 mM GTP, 0.32 µL of 50 µM LIDAR_TSO

mix (Scacchetti et al., 2024) or SS3_TSO, 0.5 U/µL murine RNase inhibitor (New England Biolabs,

Cat. No M0314S), and 2U/µL Maxima H-minus RT (Thermo Scientific, Cat. No EP0752);

concentrations refer to a final volume of 8 µL] were added. Reverse transcription was performed

as follows: 42°C for 90 min, 10 cycles at 50°C for 2 min and 42°C for 2 min, 85°C for 5 min. cDNA

was pre-ampified by adding 12 µL of KAPA_mix [1X KAPA HiFi HotStart buffer Ready Mix (Roche,

Cat. No KK2601), 1 µM LIDAR_preamp_f, and 0.1 µM LIDAR_preamp_r (Scacchetti et al., 2024);

concentrations refer to a final volume of 20 µL] and by following these PCR cycling conditions:

denaturation (95°C for 3 min), 5 x (98°C for 20 s, 70°C for 30 s, 72°C for 30 s), final extension

(72°C for 5 min). Pre-amplified libraries were purified using 2.5X SPRI beads (Beckman Coulter,

Cat. No B23319) and eluted in 37 µL of TE buffer (10mM Tris-HCl pH 8, 1 mM EDTA). 0.5 µL of

each 10 µM custom Nextera indexing primers was added, followed by 12 µL of Q5_mix [1X Q5

buffer, 0.5 mM dNTPs, 0.02 U/µL Q5 High-Fidelity DNA Polymerase (New England Biolabs, Cat.

No M0491); concentrations refer to a final volume of 50 µL]. Samples were incubated with the

following PCR cycling conditions: denaturation (98°C for 30 s), 15 x (98°C for 10 s, 65°C for 20 s,

72°C for 20 s). Final libraries were purified using 2X SPRI beads (Beckman Coulter, Cat. No

B23319) and eluted in 45 µL of TE buffer (10 mM Tris-HCl pH 8, 1 mM EDTA). Libraries were

analyzed on a 2% agarose gel and quantified using Qubit dsDNA Quantification Assay kit

(Invitrogen, Cat. No Q32854). Libraries were sequenced on Illumina NextSeq500, NextSeq1000,

or NextSeq2000 instruments.

CRISPRi

CRISPRi sgRNAs targeting the promoter region of *Sdcbp*, *Pus1,* and control (see **Table S5** for

all oligonucleotide sequences) were cloned in Lenti-(BB)-EF1a-KRAB-dCas9-P2A-BlastR

(Addgene, Cat. No 118154) using Golden Gate. To generate lentiviruses, ~0.5 million HEK293FT

in 6-well plates were transfected with 1.5 µg of CRISPRi plasmid together with 1 µg of psPAX2

packaging plasmid (Addgene, Cat. No 12260) and 0.5 µg of pMDG.2 envelope plasmid (Addgene,

Cat. No 12259) using Lipofectamine 3000 Transfection Reagent (Invitrogen, Cat. No L3000015).

24 h after transfection, medium was changed to lentivirus_collection_medium. Conditioned

medium was collected 48 h later, and cell and debris were removed by centrifugation at 500 *g* for

5 min, followed by 1,250 *g* for 5 min. Cleared conditioned medium containing lentivirus was

concentrated to 50 µL using Amicon Ultra Centrifugal Filter with 100 kDa MWCO (Millipore-Sigma,

Cat. No UFC5100). To generate stable CRISPRi cells, 50 thousand CADs were transduced with

50 µL of lentiviruses using 8 µg/mL polybrene (final concentration). 48 h after infection, medium

was supplemented with with 20 µg/mL blasticidin (Invivogen, Cat. No ant-bl-1) and cells were

selected for 4 days. To generate the pool of different CRISPRi cells, approximately equal number

of cells from individual knockdowns were mixed and cells were expanded in the presence of 20

µg/mL blasticidin. Cells were washed twice with PBS and EV-depleted medium was added. 24 h

later, conditioned medium was collected and cleared at 300 *g* for 5 min at 4°C, followed by

centrifugation at 2,000 *g* for 10 min at 4°C, 10,000 *g* for 30 min at 4°C, and concentration using

Amicon Ultra Centrifugal Filter with 100 kDa MWCO (Millipore-Sigma, Cat. No UFC5100). RNA

was extracted from cells and concentrated conditioned medium as described above. sgRNA-seq

libraries were prepared as described for the genome-wide screen using LIDAR_TSO_mix during

the reverse transcription. Libraries were sequenced on Illumina NextSeq1000 or NextSeq2000

instruments.

Generation of *Sdcbp*[-/-] and *Pus1*[-/-] cells

sgRNAs targeting *Sdcbp* and *Pus1* (see **Table S5** for all oligonucleotide sequences) were cloned
in PX459 (Addgene, Cat. No 62988) using Golden Gate. 0.5 million CADs were transfected with
2.5 µg of the targeting plasmids (PX459 backbone) using Lipofectamine 3000 Transfection
Reagent (Invitrogen, Cat. No L3000015). 24 h after transfection, medium was supplemented with
2 µg/mL puromycin, and cells were selected for 48 h. Next, cells were seeded in 15-cm dishes at
500–1,000 cells per dish and incubated for 6 days without puromycin. Individual colonies
(corresponding to single clonal populations) were picked and transferred to 96-well plates. Clones
were expanded and tested by western blot to assess knock-out efficiency. Whole cell extracts
were prepared by lysis of cell pellets in RIPA buffer (10 mM Tris-HCl pH 8, 140 mM NaCl, 1 mM
EDTA, 0.1% Na-deoxycholate, 1% Triton-X-100, 0.1% SDS, 1X cOmplete EDTA-free Protease
Inhibitor), followed by sonication using Bioruptor (Diagenode) (30 s ON/OFF, 5 cycles) and
clarification by centrifugation at 18,000 g for 10 min at 4°C.The following antibody dilutions were
used for western blots: 1:1000 for αSDCBP (Abcam, Cat. No ab19903), 1:500 for αPUS1 (Abcam,
Cat. No ab203010), 1:1000 for αSUZ12 (Cell Signaling Technology, Cat. No 3737S), 1:1000 for
αTUBB (Abcam, Cat. No ab6046).

Export of synthetic sgRNAs

sgRNAs containing uridine or pseudouridine (Ψ) were synthesized by *in vitro* transcription. Briefly,
individual sgRNAs were PCR amplified to add a sense T7 promoter sequence (**Table S5**). RNA
was transcribed from ~500 ng of template DNA using HiScribe T7 High Yield RNA Synthesis Kit
(New England Biolabs, Cat. No E2040L) following the recommended protocol for short RNAs.
UTP was substituted with pseudo-UTP (Jena Bioscience, Cat. No NU-1139L) to generate Ψ-
containing sgRNAs. DNA was digested with Turbo DNase-I (Invitrogen, Cat. No AM2238) and
RNA was purified using TriPure (Roche, Cat. No 11667165001) as described above.

For assessing export with qPCR, 0.2 million CADs were transfected with 1 µg of UTP- or Ψ-
containing EGFP_1_sgRNA using Lipofectamine RNAiMAX Transfection Reagent (Invitrogen,
Cat. No 13778075). 6 h after transfection, cells were washed with serum-free medium, and then
incubated for additional 20 h in serum-free medium. Conditioned medium was collected and
cleared at 300 g for 5 min at 4°C, followed by centrifugation at 2,000 g for 10 min at 4°C and then
10,000 g for 30 min at 4°C. RNA was extracted from cells and cleared conditioned medium as
described above. RT-qPCR was performed using Power SYBR Green RNA-to-C$_T$ (Applied
Biosystems, Cat. No 4389986) with the EGFP_1_qPCR primers (**Table S5**). For assessing export

by sequencing, the same procedure was followed with some minor modifications. First, two

different sgRNA pools were created mixing 3 UTP-containing and 3 Ψ-containing sgRNAs (**Table**

S5). Second, 500 ng of sgRNA pools were used to transfect 0.3 million CADs. RNA was extracted

from cells and cleared conditioned medium as described above. sgRNA-seq libraries were

prepared using the same strategy as in the genome-wide screen. Libraries were sequenced on

Illumina NextSeq1000 or NextSeq2000 instruments.

Bisulfite-modified LIDAR

Bisulfite conversion and desulphonation of RNA was performed following the BID-seq protocol

(Dai et al., 2023). Briefly, input RNA (200 ng for cellular RNA and 10 ng for exRNAs) was diluted

in 8.5 μL of water, 45 μL of freshly prepared BS reagent (2.4 M Na_2SO_3 and 0.36 M $NaHSO_3$)

were added, and samples were incubated at 70°C for 3 h. Next, 75 μL of water, 270 μL of RNA

Binding Buffer (Zymo Research, Cat. No. R1013), and 400 μL of 100% EtOH were added, and

samples were transferred on a Zymo column (Zymo Research, Cat. No. R1013). After washing

with 200 μL of RNA wash buffer, samples were incubated in-column with 200 μL of RNA

desulphonation buffer (Zymo Research, Cat. No. R5001-3-40) for 1 h and 15 min at 25°C.

Samples were washed with twice with RNA wash buffer (400 μL and 700 μL), then eluted in 6 μL

of RNase-free water. Libraries were prepared starting from half of the eluate using the LIDAR

protocol described above with a few modifications: 1) initial RNA denaturation was conducted at

72°C for 3 min; 2) $MgCl_2$ was substituted with $MnCl_2$ in the RT_mix (keeping the same amounts);

3) reverse transcription conditions were changed to 8°C for 10 min, 16°C for 10 min, 25°C for 10

min, 42°C for 80 min, 10 cycles at 50˚C for 2 min and 42˚C for 2 min, 85°C for 5 min. Libraries

from untreated RNA input samples were prepared in parallel.

RNA affinity chromatography

Biotinylated sgRNAs containing U or Ψ were generated by *in vitro* transcription. Individual

sgRNAs were PCR amplified to add a sense T7 promoter sequence (**Table S5**). RNA was

transcribed from ~600 ng of template DNA (1:1 mix of EGFP_1sgRNA and SCR_1_sgRNA)

using HiScribe T7 High Yield RNA Synthesis Kit (New England Biolabs, Cat. No E2040L) following

the recommended protocol for short RNAs. In all reactions, ATP was substitute with a 10:1 mix of

ATP and N6-(6-Aminohexyl)-ATP-Biotin (Jena Bioscience, Cat. No NU-805-BIO). UTP was

substituted with pseudo-UTP (Jena Bioscience, Cat. No NU-1139L) to generate Ψ-containing

sgRNAs. DNA was digested with Turbo DNase-I (Invitrogen, Cat. No AM2238) and RNA was

purified using TriPure (Roche, Cat. No 11667165001) as described above.

To prepare protein extracts, approximately 50 million CADs were collected and washed with PBS.

Cell pellets were resuspended in 1 mL of ice-cold CHAPS lysis buffer [50 mM Tris-Cl pH = 7.5,

150 mM NaCl, 0.5% CHAPS, 0.5 mM DTT, 1 X cOmplete EDTA-free Protease Inhibitor Cocktail

(Roche, Cat. No 11873580001)] and cells were lysed via 2 cycles of freezing in dry ice/EtOH bath

followed by rapid thawing at 37°C. Extract was clarified by centrifugation at 18,000 g for 10 min

at 4°C. Protein amounts were quantified using Bio-Rad Protein Assay Dye Reagent Concentrate

(Bio-Rad, Cat. No 5000006).

50 µg of U- or Ψ-containing biotinylated sgRNAs were mixed with 1 mg of CAD protein extract in

CHAPS lysis buffer supplemented with 1.5 mM $MgCl_2$ and the samples were rotated for 1 h at

room temperature. As a control, RNA was omitted from 1 reaction ("beads only"). Next, 50 µL of

Dynabeads MyOne Streptavidin C1 beads (Invitrogen, Cat. No 65001), equilibrated in CHAPS

lysis buffer supplemented with 1.5 mM $MgCl_2$, were added and samples were rotated for 1 h at

4°C. Beads were washed 3 X with medium salt buffer (50 mM Tris-Cl pH = 7.5, 150 mM NaCl,

1.5 mM $MgCl_2$, 0.5 mM DTT, 1 X cOmplete EDTA-free Protease Inhibitor Cocktail) and 3 X with

high salt buffer (50 mM Tris-Cl pH = 7.5, 500 mM NaCl, 1.5 mM $MgCl_2$, 0.5 mM DTT, 1 X cOmplete

EDTA-free Protease Inhibitor Cocktail). Proteins were prepared using Protifi S-trap Micro columns

per manufacturer's instructions. Following S-trap elution, peptides were lyophilized, resuspended

in 0.1 % formic acid for further cleanup by desalting. Desalted peptides were lyophilized and

resuspended in 4% acetonitrile containing 0.1% formic acid for LC-MS analysis. Mass spectra for

peptides were obtained with a Dionex Ultimate3000 liquid chromatograph (Thermo Fisher

Scientific) coupled to a Q-Exactive HF mass spectrometer (Thermo Fisher Scientific). Peptides

were loaded onto an in-house silica capillary column (75 µM inner diameter) packed with C18

resin (ReproSil-Pur C18-AQ 2.4-µm resin; Dr. Maisch GmbH, Ammerbuch, Germany). Peptides

were eluted with a gradient of 2%–25% B (85 min) followed by a gradient of 25%–50% (30 min)

and a hold at 98% B (15 min) (mobile phase A: 0.1% formic acid, mobile phase B: 80%

acetonitrile, 0.1% formic acid). Flow rate was 300 nL/min. Data were collected in data-

independent acquisition (DIA) mode had the following settings: Full scan MS – Scan range, 385–

1015 m/z; AGC target, 1e6; Ion inject time, 60 ms; Resolution, 60,000; Isolation width, 630.

MS/MS scan – Resolution, 30,000; MS2 AGC target, 1e6, Ion inject time, 60 ms; Isolation width,

24; HCD energy, 30. The reference proteomes used for data processing were the Uniprot Mus

musculus reference proteome and a Cas9 FASTA (UniprotID: Q99ZW2). DDA spectra were

processed with ProteomeDiscoverer (version 2.4.1.15) and Sequest. False discovery rates (FDR)

were estimated using a decoy database search with strict and relaxed target FDRs set to 0.01

and 0.05, respectively. DIA spectra were processed with DIA-NN (version 1.9.2). Outputs were

filtered at 0.01 FDR, which were estimated based on in silico-generated spectral libraries

generated by DIA-NN.

***Myl6* knock-down**

siRNAs against *Myl6* and TYE 563 Control DsiRNA were obtained from Integrated DNA

Technologies (Cat. No mm.Ri.Myl6.13.2, and TriFECTa RNAi Kit). CADs were transfected with

siRNAs using Lipofectamine RNAiMAX using 15 pmol of siRNA per 0.3 million cells. Knockdown

efficiency was measured 72 h after transfection via RT-qPCR using Power SYBR Green RNA-to-

C_T (Applied Biosystems, Cat. No 4389986) with the Myl6_qPCR primers (**Table S5**).

DATA PROCESSING AND ANALYSIS

Proteomics data processing and analysis

Protein with measurable intensities only in cell or only in EV samples were excluded from the plot

in **Fig. 1C** and denoted as "Cell-exclusive" or "EV-exclusive" in **Table S1**. Raw intensities were

normalized using the `median_normalization` function or `proDA` package (Ahlmann-Eltze &

Anders, 2020). Differential abundance analysis was performed in R using the `limma` package as

previously described (Kammers, Cole, Tiengwe, & Ruczinski, 2015; Ritchie et al., 2015). Adjusted

p values (FDR) were calculated using the `p.adjust` function (method = 'fdr'*).* Gene ontology

analysis was performed using DAVID (Cellular component 5) (Huang, Sherman, & Lempicki,

2008; Sherman et al., 2022), including "EV-exclusive" proteins.

Processing, mapping, and read counting (LIDAR, BID-LIDAR, small RNA-seq, mRNA-seq)

Adapter trimming, UMI extraction, mapping to the *M. musculus* genome , deduplication and read

counting for LIDAR, BID-LIDAR, and small RNA-seq (NEB) was performed exactly as described

in (Scacchetti et al., 2024) with scripts available on GitHub (https://github.com/bonasio-lab). For

mRNA-seq libraries, paired-end reads were mapped directly using STAR (Dobin et al., 2013) with

the following parameters: `--peOverlapNbasesMin 10 --readMapNumber -1 --`

`clip3pNbases 0 --alignIntronMax 100000 --outSAMtype BAM Unsorted`. Counts

per gene were calculated in R using summarizeOverlaps function from GenomicAlignments.

RNAs with 0 counts in all wild type samples (cellular or exRNAs) were excluded. Raw counts

were batch corrected using `Combat_seq` (Zhang, Parmigiani, & Johnson, 2020).

Processing, mapping and, read counting (sgRNA libraries)

Adapter trimming on single-end reads was performed using `Cutadapt` (Martin, 2011) with the

following parameters: `-a gttttagagcta -m 6`. UMI extraction was performed using the

`process_files_single.sh` script available on GitHub (https://github.com/bonasio-lab) with

the following parameters: 8bp (libraries constructed using SS3_TSO) or LIDAR (libraries

constructed using LIDAR_TSO_mix). Reads were further trimmed with `Cutadapt` with the

following parameter: `-u 1`. Reads were mapped to the reference sgRNA Brie library or the sgRNA

sequences used in "Export of synthetic sgRNAs" using `bowtie2` (Langmead & Salzberg, 2012)

using standard parameters but specifying `--no-rc`. Aligned reads were deduplicated using

`umi_tools` (Smith, Heger, & Sudbery, 2017) with the following parameter: `--method unique`.

Reads shorter than 15 bp were filtered out using `bedtools` (Quinlan, 2014). Read counting was

performed from bam files using MAGeCK (W. Li et al., 2014) with the following parameters:

`count`. Raw counts were batch corrected using `Combat_seq` (Zhang et al., 2020).

CRISPR screen analysis

Batch corrected raw counts were normalized using MAGeCK with the following parameter: `count`

`-norm-method total`. Differential sgRNA abundance between cell and EVs at day 2 was

calculated with DESeq2 (Love, Huber, & Anders, 2014) with the following parameter: `fitType`

`= "local"`.

To identify candidates, first sgRNAs targeting genes not expressed in CADs (RPKM <1) were

excluded. Then, to account for individual sgRNA export efficiency as well as changes in sgRNA

representation in cells at day 10 due to fitness effects, we calculated a "predicted" sgRNA

abundance in day 10 EVs as follows:

$$i_{EV(day10)predicted} = average(\frac{i_{EV(day2)observed}}{i_{Cells(day2)observed}}) \times i_{Cells(day10)observed}$$

Where i is the sgRNA read frequency in the indicated samples. Differences in EV sgRNA

predicted and observed abundance at day 10 were calculated using MAGeCK with the following

parameter: `test -paired`. GO analysis was performed using DAVID (cellular component, GO

level #5) (Huang et al., 2008; Sherman et al., 2022).

Differential RNA abundance analysis (LIDAR)

797 Differential abundance between wild type cellular and exRNAs, as well as comparisons between

798 wild type and *Pus1*[-/-] mutant CAD, was performed using DESeq2 (Love et al., 2014) with the

799   following parameter: `fitType = "local"`. RNAs with an adjusted *p* value < 0.1 were considered

800   differentially enriched/exported.

Analysis of Ψ with BID-LIDAR

Detection of genomic positions with Ψ was done by identifying positions with deletions induced

by BID sequencing compared to a non-bisulfite-induced input. First, reads were mapped using

the LIDAR analysis pipeline, as described ab An in-house script using `GenomicRanges`

(Lawrence   et   al.,   2013),   `BioStrings`

(https://bioconductor.org/packages/release/bioc/html/Biostrings.html), and `bedtools` (Quinlan,

2014) was used to detect deletions within reads, map putative Ψ sites to the genome reference,

and calculate the fraction of reads at each genomic position that contain a deletion. Only positions

with at least 5% of reads deleted were considered to harbor a deletion. Positions where deletion

occurred 2 X more frequently in BID than input samples were classified as Ψ sites. Position with

a BID deletion rate in wild type < 0.75X compared to *Pus1⁻ᐟ⁻* were classified as PUS1-sensitive Ψ

sites. These positions were then assigned to the transcripts that contained them to generate a list

of modified RNAs.

For tRNA analysis of BID-LIDAR, the position of canonical bases for each tRNA gene were

annotated using the strategy described in (Scacchetti et al., 2024). These positions were used to

identify tRNA containing a U at position 27, which served as candidates for containing a Ψ at this

position, and to visualize and quantify reads with deletions at each canonical base.

DATA AVAILABILITY

819   Sequencing data will be deposited in GEO upon restoration of NCBI services.

820   Proteomics data deposited at PRIDE: PXD069544 and PXD069598

821   Code will be available on GitHub

ACKNOWLEDGMENTS

823   The authors thank Andy J. Minn for technical advice, and Kenneth Zaret and the Bonasio lab for

824   feedback on the experiments and manuscript. R.B. acknowledges support from the NIH

825   (R35GM153281; DP1NS148061). C.C.C. acknowledges support from the NIH (R35GM151087)

826   This project was supported in part by a Penn Epigenetics Institute Pilot grant to R.B. and A.S.

AUTHOR CONTRIBUTIONS

Conceptualization: A.S. and R.B.; methodology: A.S., E.J.S., R.B. ; formal analyses: A.S., E.J.S., R.B.; investigation: A.S., E.J.S., L.N.R, J.F.D, J.T., G.E.L., R.L., N.C. ; data curation: A.S., E.J.S., R.B.; writing – original draft: A.S. and R.B.; writing – review & editing: all authors; visualization: A.S., E.J.S., R.B.; supervision: R.B., C.C.C., N.C., B.A.G.; project administration: A.S. and R.B.; funding acquisition: R.B.

DECLARATION OF INTEREST

The authors declare no competing interest.

REFERENCES

Ahlmann-Eltze, C., & Anders, S. (2020). proDA: Probabilistic Dropout Analysis for Identifying Differentially Abundant Proteins in Label-Free Mass Spectrometry. *bioRxiv*. doi:10.1101/661496

Al Othman, A., Bagrov, D., Rozenberg, J. M., Glazova, O., Skryabin, G., Tchevkina, E., . . . Durymanov, M. (2024). Retrovirus-like gag protein Arc/Arg3.1 is involved in extracellular-vesicle-mediated mRNA transfer between glioma cells. *Biochim Biophys Acta Gen Subj, 1868*(1), 130522. doi:10.1016/j.bbagen.2023.130522

Ashley, J., Cordy, B., Lucia, D., Fradkin, L. G., Budnik, V., & Thomson, T. (2018). Retrovirus-like Gag Protein Arc1 Binds RNA and Traffics across Synaptic Boutons. *Cell, 172*(1-2), 262-274 e211. doi:10.1016/j.cell.2017.12.022

Baietti, M. F., Zhang, Z., Mortier, E., Melchior, A., Degeest, G., Geeraerts, A., . . . David, G. (2012). Syndecan-syntenin-ALIX regulates the biogenesis of exosomes. *Nature cell biology, 14*(7), 677-685. doi:10.1038/ncb2502

Barman, B., Sung, B. H., Krystofiak, E., Ping, J., Ramirez, M., Millis, B., . . . Weaver, A. M. (2022). VAP-A and its binding partner CERT drive biogenesis of RNA-containing extracellular vesicles at ER membrane contact sites. *Dev Cell, 57*(8), 974-994 e978. doi:10.1016/j.devcel.2022.03.012

Bellingham, S. A., Coleman, B. M., & Hill, A. F. (2012). Small RNA deep sequencing reveals a distinct miRNA signature released in exosomes from prion-infected neuronal cells. *Nucleic Acids Res, 40*(21), 10937-10949. doi:10.1093/nar/gks832

Bervoets, S., Jacob, M. S., Devineni, A. V., Mahoney, B. D., Sullivan, K. R., Butts, A. R., . . . Caron, S. J. C. (2024). dArc1 controls sugar reward valuation in Drosophila melanogaster. *bioRxiv*. doi:10.1101/2024.11.04.621761

Betti, F., Ladera-Carmona, M. J., Weits, D. A., Ferri, G., Iacopino, S., Novi, G., . . . Perata, P.
(2021). Exogenous miRNAs induce post-transcriptional gene silencing in plants. *Nat*
Plants, 7(10), 1379-1388. doi:10.1038/s41477-021-01005-w

Blanchette, C. R., Scalera, A. L., Harris, K. P., Zhao, Z., Dresselhaus, E. C., Koles, K., . . . Rodal,
A. A. (2022). Local regulation of extracellular vesicle traffic by the synaptic endocytic
machinery. *J Cell Biol, 221*(5). doi:10.1083/jcb.202112094

Brito, C., & Sousa, S. (2020). Non-Muscle Myosin 2A (NM2A): Structure, Regulation and
Function. *Cells, 9*(7). doi:10.3390/cells9071590

Budnik, V., Ruiz-Canada, C., & Wendler, F. (2016). Extracellular vesicles round off
communication in the nervous system. *Nat Rev Neurosci, 17*(3), 160-172.
doi:10.1038/nrn.2015.29

Cafe, S. L., Anderson, A. L., & Nixon, B. (2020). In vitro Induction and Detection of Acrosomal
Exocytosis in Human Spermatozoa. *Bio Protoc, 10*(14), e3689.
doi:10.21769/BioProtoc.3689

Cai, Q., Qiao, L., Wang, M., He, B., Lin, F. M., Palmquist, J., . . . Jin, H. (2018). Plants send small
RNAs in extracellular vesicles to fungal pathogen to silence virulence genes. *Science,*
360(6393), 1126-1129. doi:10.1126/science.aar4142

Caohuy, H., & Pollard, H. B. (2002). Protein kinase C and guanosine triphosphate combine to
potentiate calcium-dependent membrane fusion driven by annexin 7. *J Biol Chem,*
277(28), 25217-25225. doi:10.1074/jbc.M202452200

Caudron-Herger, M., Jansen, R. E., Wassmer, E., & Diederichs, S. (2021). RBP2GO: a
comprehensive pan-species database on RNA-binding proteins, their interactions and
functions. *Nucleic Acids Res, 49*(D1), D425-D436. doi:10.1093/nar/gkaa1040

Caudron-Herger, M., Rusin, S. F., Adamo, M. E., Seiler, J., Schmid, V. K., Barreau, E., . . .
Diederichs, S. (2019). R-DeeP: Proteome-wide and Quantitative Identification of RNA-
Dependent Proteins by Density Gradient Ultracentrifugation. *Mol Cell, 75*(1), 184-199
e110. doi:10.1016/j.molcel.2019.04.018

Cerneckis, J., Cui, Q., He, C., Yi, C., & Shi, Y. (2022). Decoding pseudouridine: an emerging
target for therapeutic development. *Trends Pharmacol Sci, 43*(6), 522-535.
doi:10.1016/j.tips.2022.03.008

Cha, D. J., Franklin, J. L., Dou, Y., Liu, Q., Higginbotham, J. N., Demory Beckler, M., . . . Patton,
J. G. (2015). KRAS-dependent sorting of miRNA to exosomes. *Elife, 4*, e07197.
doi:10.7554/eLife.07197

Chen, Z., Zhou, Y., Xue, C., Zeng, L., Deng, S., Xu, Z., . . . Zheng, J. (2025). Psychological stress-
induced ALKBH5 deficiency promotes tumour innervation and pancreatic cancer via
extracellular vesicle transfer of RNA. *Nat Cell Biol, 27*(6), 1035-1047. doi:10.1038/s41556-
025-01667-0

Chevillet, J. R., Kang, Q., Ruf, I. K., Briggs, H. A., Vojtech, L. N., Hughes, S. M., . . . Tewari, M.
(2014). Quantitative and stoichiometric analysis of the microRNA content of exosomes.
Proc Natl Acad Sci U S A, 111(41), 14888-14893. doi:10.1073/pnas.1408301111

Chitti, S. V., Gummadi, S., Kang, T., Shahi, S., Marzan, A. L., Nedeva, C., . . . Mathivanan, S.
(2024). Vesiclepedia 2024: an extracellular vesicles and extracellular particles repository.
Nucleic Acids Res, 52(D1), D1694-D1698. doi:10.1093/nar/gkad1007

Conibear, E., Cleck, J. N., & Stevens, T. H. (2003). Vps51p mediates the association of the GARP
(Vps52/53/54) complex with the late Golgi t-SNARE Tlg1p. *Mol Biol Cell, 14*(4), 1610-
1623. doi:10.1091/mbc.e02-10-0654

Conine, C. C., & Rando, O. J. (2022). Soma-to-germline RNA communication. *Nat Rev Genet,*
23(2), 73-88. doi:10.1038/s41576-021-00412-1

Costa, B., Li Calzi, M., Castellano, M., Blanco, V., Cuevasanta, E., Litvan, I., . . . Tosar, J. P.
(2023). Nicked tRNAs are stable reservoirs of tRNA halves in cells and biofluids. *Proc Natl*
Acad Sci U S A, 120(4), e2216330120. doi:10.1073/pnas.2216330120

Crocker, O. J., Trigg, N. A., & Conine, C. C. (2022). Cloning and Sequencing Eukaryotic Small
RNAs. *Curr Protoc, 2*(8), e495. doi:10.1002/cpz1.495

Dai, Q., Zhang, L. S., Sun, H. L., Pajdzik, K., Yang, L., Ye, C., . . . He, C. (2023). Quantitative
sequencing using BID-seq uncovers abundant pseudouridines in mammalian mRNA at
base resolution. *Nat Biotechnol, 41*(3), 344-354. doi:10.1038/s41587-022-01505-w

Deatherage, B. L., & Cookson, B. T. (2012). Membrane vesicle release in bacteria, eukaryotes,
and archaea: a conserved yet underappreciated aspect of microbial life. *Infect Immun,*
80(6), 1948-1957. doi:10.1128/IAI.06014-11

Dixson, A. C., Dawson, T. R., Di Vizio, D., & Weaver, A. M. (2023). Context-specific regulation of
extracellular vesicle biogenesis and cargo selection. *Nat Rev Mol Cell Biol, 24*(7), 454-
476. doi:10.1038/s41580-023-00576-0

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R.
(2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21.
doi:10.1093/bioinformatics/bts635

Doench, J. G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E. W., Donovan, K. F., . . . Root, D.
E. (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of
CRISPR-Cas9. *Nat Biotechnol, 34*(2), 184-191. doi:10.1038/nbt.3437

Erlendsson, S., Morado, D. R., Cullen, H. B., Feschotte, C., Shepherd, J. D., & Briggs, J. A. G.
(2020). Structures of virus-like capsids formed by the Drosophila neuronal Arc proteins.
Nat Neurosci, 23(2), 172-175. doi:10.1038/s41593-019-0569-y

Garcia-Martin, R., Wang, G., Brandao, B. B., Zanotto, T. M., Shah, S., Kumar Patel, S., . . . Kahn,
C. R. (2022). MicroRNA sequence codes for small extracellular vesicle release and
cellular retention. *Nature, 601*(7893), 446-451. doi:10.1038/s41586-021-04234-3

Grosshans, H., Lecointe, F., Grosjean, H., Hurt, E., & Simos, G. (2001). Pus1p-dependent tRNA
pseudouridinylation becomes essential when tRNA biogenesis is compromised in yeast.
J Biol Chem, 276(49), 46333-46339. doi:10.1074/jbc.M107141200

Gustafsson, H. T., Ferguson, L., Galan, C., Yu, T., Upton, H., Kaymak, E., . . . Rando, O. J. (2025).
Deep sequencing of yeast and mouse tRNAs and tRNA fragments using OTTR. *Elife, 14*.
doi:10.7554/eLife.77616

Henriques, W. S., Young, J. M., Nemudryi, A., Nemudraia, A., Wiedenheft, B., & Malik, H. S.
(2024). The Diverse Evolutionary Histories of Domesticated Metaviral Capsid Genes in
Mammals. *Mol Biol Evol, 41*(4). doi:10.1093/molbev/msae061

Herridge, R. P., Dolata, J., Migliori, V., de Santis Alves, C., Borges, F., Schorn, A. J., . . .
Martienssen, R. A. (2025). Pseudouridine guides germline small RNA transport and
epigenetic inheritance. *Nat Struct Mol Biol, 32*(2), 277-286. doi:10.1038/s41594-024-
01392-6

Holliday, L. S., Faria, L. P., & Rody, W. J., Jr. (2019). Actin and Actin-Associated Proteins in
Extracellular Vesicles Shed by Osteoclasts. *Int J Mol Sci, 21*(1).
doi:10.3390/ijms21010158

Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2008). Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nature Protocols, 4*(1), 44-57.
doi:10.1038/nprot.2008.211

Huo, L., Du, X., Li, X., Liu, S., & Xu, Y. (2021). The Emerging Role of Neural Cell-Derived
Exosomes in Intercellular Communication in Health and Neurodegenerative Diseases.
Front Neurosci, 15, 738442. doi:10.3389/fnins.2021.738442

Kaletsky, R., Moore, R. S., Vrla, G. D., Parsons, L. R., Gitai, Z., & Murphy, C. T. (2020). C. elegans
interprets bacterial non-coding RNAs to learn pathogenic avoidance. *Nature, 586*(7829),
445-451. doi:10.1038/s41586-020-2699-5

Kammers, K., Cole, R. N., Tiengwe, C., & Ruczinski, I. (2015). Detecting Significant Changes in
Protein Abundance. *EuPA Open Proteom, 7*, 11-19. doi:10.1016/j.euprot.2015.02.002

Kampmann, M. (2018). CRISPRi and CRISPRa Screens in Mammalian Cells for Precision
Biology and Medicine. *ACS Chem Biol, 13*(2), 406-416. doi:10.1021/acschembio.7b00657

Korkut, C., Ataman, B., Ramachandran, P., Ashley, J., Barria, R., Gherbesi, N., & Budnik, V.
(2009). Trans-synaptic transmission of vesicular Wnt signals through Evi/Wntless. *Cell,
139*(2), 393-404. doi:10.1016/j.cell.2009.07.051

Kugeratski, F. G., Hodge, K., Lilla, S., McAndrews, K. M., Zhou, X., Hwang, R. F., . . . Kalluri, R.
(2021). Quantitative proteomics identifies the core proteome of exosomes with syntenin-
1 as the highest abundant protein and a putative universal biomarker. *Nat Cell Biol, 23*(6),
631-641. doi:10.1038/s41556-021-00693-y

Kuncha, S. K., Mazeed, M., Singh, R., Kattula, B., Routh, S. B., & Sankaranarayanan, R. (2018).
A chiral selectivity relaxed paralog of DTD for proofreading tRNA mischarging in Animalia.
Nat Commun, 9(1), 511. doi:10.1038/s41467-017-02204-w

Kunitake, K., Mizuno, T., Hattori, K., Oneyama, C., Kamiya, M., Ota, S., . . . Kojima, R. (2024).
Barcoding of small extracellular vesicles with CRISPR-gRNA enables comprehensive,
subpopulation-specific analysis of their biogenesis and release regulators. *Nat Commun,*
15(1), 9777. doi:10.1038/s41467-024-53736-x

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods,*
9(4), 357-359. doi:10.1038/nmeth.1923

Lawrence, M., Huber, W., Pages, H., Aboyoun, P., Carlson, M., Gentleman, R., . . . Carey, V. J.
(2013). Software for computing and annotating genomic ranges. *PLoS Comput Biol, 9*(8),
e1003118. doi:10.1371/journal.pcbi.1003118

Leidal, A. M., Huang, H. H., Marsh, T., Solvik, T., Zhang, D., Ye, J., . . . Debnath, J. (2020). The
LC3-conjugation machinery specifies the loading of RNA-binding proteins into
extracellular vesicles. *Nat Cell Biol, 22*(2), 187-199. doi:10.1038/s41556-019-0450-y

Li, J. B., & Walkley, C. R. (2025). Leveraging genetics to understand ADAR1-mediated RNA
editing in health and disease. *Nat Rev Genet, 26*(8), 532-546. doi:10.1038/s41576-025-
00830-5

Li, W., Xu, H., Xiao, T., Cong, L., Love, M. I., Zhang, F., . . . Liu, X. S. (2014). MAGeCK enables
robust identification of essential genes from genome-scale CRISPR/Cas9 knockout
screens. *Genome Biol, 15*(12), 554. doi:10.1186/s13059-014-0554-4

Liu, X. M., Ma, L., & Schekman, R. (2021). Selective sorting of microRNAs into exosomes by
phase-separated YBX1 condensates. *Elife, 10*. doi:10.7554/eLife.71982

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-
0550-8

Lu, A., Wawro, P., Morgens, D. W., Portela, F., Bassik, M. C., & Pfeffer, S. R. (2018). Genome-
wide interrogation of extracellular vesicle biology using barcoded miRNAs. *Elife, 7*.
doi:10.7554/eLife.41460

M'Angale, P. G., Lemieux, A., Liu, Y., Wang, S., Zinter, M., Alegre, G., . . . Thomson, T. (2025).
Capsid transfer of the retrotransposon Copia controls structural synaptic plasticity in
Drosophila. *PLoS Biol, 23*(2), e3002983. doi:10.1371/journal.pbio.3002983

Ma, L., Lu, H., Tian, Z., Yang, M., Ma, J., Shang, G., . . . Chen, Z. (2022). Structural insights into
the interactions and epigenetic functions of human nucleic acid repair protein ALKBH6. *J*
Biol Chem, 298(3), 101671. doi:10.1016/j.jbc.2022.101671

Madigan, V., Zhang, Y., Raghavan, R., Wilkinson, M. E., Faure, G., Puccio, E., . . . Zhang, F.
(2024). Human paraneoplastic antigen Ma2 (PNMA2) forms icosahedral capsids that can
be engineered for mRNA delivery. *Proc Natl Acad Sci U S A, 121*(11), e2307812120.
doi:10.1073/pnas.2307812120

Maori, E., Garbian, Y., Kunik, V., Mozes-Koch, R., Malka, O., Kalev, H., . . . Shafir, S. (2019). A
Transmissible RNA Pathway in Honey Bees. *Cell Rep, 27*(7), 1949-1959 e1946.
doi:10.1016/j.celrep.2019.04.073

Marangon, D., Castro, E. S. J. H., & Lecca, D. (2022). Neuronal and Glial Communication via
Non-Coding RNAs: Messages in Extracellular Vesicles. *Int J Mol Sci, 24*(1).
doi:10.3390/ijms24010470

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal, 17(1). doi:10.14806/ej.17.1.200

Matsui, T., Osaki, F., Hiragi, S., Sakamaki, Y., & Fukuda, M. (2021). ALIX and ceramide
differentially control polarized small extracellular vesicle release from epithelial cells.
EMBO Rep, 22(5), e51475. doi:10.15252/embr.202051475

Mews, P., Donahue, G., Drake, A. M., Luczak, V., Abel, T., & Berger, S. L. (2017). Acetyl-CoA
synthetase regulates histone acetylation and hippocampal memory. *Nature, 546*(7658),
381-386. doi:10.1038/nature22405

Mittelbrunn, M., Gutierrez-Vazquez, C., Villarroya-Beltri, C., Gonzalez, S., Sanchez-Cabo, F.,
Gonzalez, M. A., . . . Sanchez-Madrid, F. (2011). Unidirectional transfer of microRNA-
loaded exosomes from T cells to antigen-presenting cells. *Nat Commun, 2*, 282.
doi:10.1038/ncomms1285

Moller, A., & Lobb, R. J. (2020). The evolving translational potential of small extracellular vesicles
in cancer. *Nat Rev Cancer, 20*(12), 697-709. doi:10.1038/s41568-020-00299-w

Molnar, A., Melnyk, C. W., Bassett, A., Hardcastle, T. J., Dunn, R., & Baulcombe, D. C. (2010).
Small silencing RNAs in plants are mobile and direct epigenetic modification in recipient
cells. *Science, 328*(5980), 872-875. doi:10.1126/science.1187959

Moore, R. S., Kaletsky, R., Lesnik, C., Cota, V., Blackman, E., Parsons, L. R., . . . Murphy, C. T.
(2021). The role of the Cer1 transposon in horizontal transfer of transgenerational
memory. *Cell, 184*(18), 4697-4712 e4618. doi:10.1016/j.cell.2021.07.022

Murillo, O. D., Thistlethwaite, W., Rozowsky, J., Subramanian, S. L., Lucero, R., Shah, N., . . .
Milosavljevic, A. (2019). exRNA Atlas Analysis Reveals Distinct Extracellular RNA Cargo
Types and Their Carriers Present across Human Biofluids. *Cell, 177*(2), 463-477 e415.
doi:10.1016/j.cell.2019.02.018

Nechooshtan, G., Yunusov, D., Chang, K., & Gingeras, T. R. (2020). Processing by RNase 1
forms tRNA halves and distinct Y RNA fragments in the extracellular environment. *Nucleic
Acids Res, 48*(14), 8035-8049. doi:10.1093/nar/gkaa526

O'Brien, K., Breyne, K., Ughetto, S., Laurent, L. C., & Breakefield, X. O. (2020). RNA delivery by
extracellular vesicles in mammalian cells and its applications. *Nat Rev Mol Cell Biol,
21*(10), 585-606. doi:10.1038/s41580-020-0251-y

Pastuzyn, E. D., Day, C. E., Kearns, R. B., Kyrke-Smith, M., Taibi, A. V., McCormick, J., . . .
Shepherd, J. D. (2018). The Neuronal Gene Arc Encodes a Repurposed Retrotransposon
Gag Protein that Mediates Intercellular RNA Transfer. *Cell, 172*(1-2), 275-288 e218.
doi:10.1016/j.cell.2017.12.024

Patton, J. R., Bykhovskaya, Y., Mengesha, E., Bertolotto, C., & Fischel-Ghodsian, N. (2005).
Mitochondrial myopathy and sideroblastic anemia (MLASA): missense mutation in the

pseudouridine synthase 1 (PUS1) gene is associated with the loss of tRNA
pseudouridylation. *J Biol Chem, 280*(20), 19823-19828. doi:10.1074/jbc.M500216200

Qi, Y., Wang, J. K., McMillian, M., & Chikaraishi, D. M. (1997). Characterization of a CNS cell line,
CAD, in which morphological differentiation is initiated by serum deprivation. *J Neurosci,*
17(4), 1217-1225. doi:10.1523/JNEUROSCI.17-04-01217.1997

Qu, S., Nelson, H. M., Liu, X., Wang, Y., Semler, E. M., Michell, D. L., . . . Patton, J. G. (2024). 5-
Fluorouracil treatment represses pseudouridine-containing miRNA export into
extracellular vesicles. *J Extracell Biol, 3*(9), e70010. doi:10.1002/jex2.70010

Quinlan, A. R. (2014). BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Curr*
Protoc Bioinformatics, 47, 11 12 11-34. doi:10.1002/0471250953.bi1112s47

Reggiori, F., Wang, C. W., Stromhaug, P. E., Shintani, T., & Klionsky, D. J. (2003). Vps51 is part
of the yeast Vps fifty-three tethering complex essential for retrograde traffic from the early
endosome and Cvt vesicle completion. *J Biol Chem, 278*(7), 5009-5020.
doi:10.1074/jbc.M210436200

Rintala-Dempsey, A. C., & Kothe, U. (2017). Eukaryotic stand-alone pseudouridine synthases -
RNA modifying enzymes and emerging regulators of gene expression? *RNA Biol, 14*(9),
1185-1196. doi:10.1080/15476286.2016.1276150

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res, 43(7), e47. doi:10.1093/nar/gkv007

Rorbach, J., Boesch, P., Gammage, P. A., Nicholls, T. J., Pearce, S. F., Patel, D., . . . Minczuk,
M. (2014). MRM2 and MRM3 are involved in biogenesis of the large subunit of the
mitochondrial ribosome. *Mol Biol Cell, 25*(17), 2542-2555. doi:10.1091/mbc.E14-01-0014

Ruan, L. L., Zhou, X. L., Tan, M., & Wang, E. D. (2013). Human cytoplasmic ProX edits
mischarged tRNAPro with amino acid but not tRNA specificity. *Biochem J, 450*(1), 243-
252. doi:10.1042/BJ20121493

Rubio Gomez, M. A., & Ibba, M. (2020). Aminoacyl-tRNA synthetases. *RNA, 26*(8), 910-936.
doi:10.1261/rna.071720.119

Sanson, K. R., Hanna, R. E., Hegde, M., Donovan, K. F., Strand, C., Sullender, M. E., . . . Doench,
J. G. (2018). Optimized libraries for CRISPR-Cas9 genetic screens with multiple
modalities. *Nat Commun, 9*(1), 5416. doi:10.1038/s41467-018-07901-8

Santangelo, L., Giurato, G., Cicchini, C., Montaldo, C., Mancone, C., Tarallo, R., . . . Tripodi, M.
(2016). The RNA-Binding Protein SYNCRIP Is a Component of the Hepatocyte Exosomal
Machinery Controlling MicroRNA Sorting. *Cell Rep, 17*(3), 799-808.
doi:10.1016/j.celrep.2016.09.031

Scacchetti, A., Shields, E. J., Trigg, N. A., Lee, G. S., Wilusz, J. E., Conine, C. C., & Bonasio, R.
(2024). A ligation-independent sequencing method reveals tRNA-derived RNAs with
blocked 3' termini. *Mol Cell, 84*(19), 3843-3859 e3848. doi:10.1016/j.molcel.2024.07.008

Schwartz, S., Bernstein, D. A., Mumbach, M. R., Jovanovic, M., Herbst, R. H., Leon-Ricardo, B.
X., . . . Regev, A. (2014). Transcriptome-wide mapping reveals widespread dynamic-
regulated pseudouridylation of ncRNA and mRNA. *Cell, 159*(1), 148-162.
doi:10.1016/j.cell.2014.08.028

Segel, M., Lash, B., Song, J., Ladha, A., Liu, C. C., Jin, X., . . . Zhang, F. (2021). Mammalian
retrovirus-like protein PEG10 packages its own mRNA and can be pseudotyped for mRNA
delivery. *Science, 373*(6557), 882-889. doi:10.1126/science.abg6155

Shalem, O., Sanjana, N. E., Hartenian, E., Shi, X., Scott, D. A., Mikkelson, T., . . . Zhang, F.
(2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science,*
343(6166), 84-87. doi:10.1126/science.1247005

Sharma, S., Jiao, X., Yang, J., Kwan, K. Y., & Kiledjian, M. (2025). Extracellular exosomal RNAs
are glyco-modified. *Nature Cell Biology, 27*(6), 983-991. doi:10.1038/s41556-025-01682-
1

Sherman, B. T., Hao, M., Qiu, J., Jiao, X., Baseler, M. W., Lane, H. C., . . . Chang, W. (2022).
DAVID: a web server for functional enrichment analysis and functional annotation of gene
lists (2021 update). *Nucleic Acids Res, 50*(W1), W216-W221. doi:10.1093/nar/gkac194

Shurtleff, M. J., Temoche-Diaz, M. M., Karfilis, K. V., Ri, S., & Schekman, R. (2016). Y-box protein
1 is required to sort microRNAs into exosomes in cells and in a cell-free reaction. *Elife, 5*.
doi:10.7554/eLife.19276

Shurtleff, M. J., Yao, J., Qin, Y., Nottingham, R. M., Temoche-Diaz, M. M., Schekman, R., &
Lambowitz, A. M. (2017). Broad role for YBX1 in defining the small noncoding RNA
composition of exosomes. *Proc Natl Acad Sci U S A, 114*(43), E8987-E8995.
doi:10.1073/pnas.1712108114

Smith, T., Heger, A., & Sudbery, I. (2017). UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. *Genome Res, 27*(3), 491-499.
doi:10.1101/gr.209601.116

Sonder, S. L., Boye, T. L., Tolle, R., Dengjel, J., Maeda, K., Jaattela, M., . . . Nylandsted, J. (2019).
Annexin A7 is required for ESCRT III-mediated plasma membrane repair. *Sci Rep, 9*(1),
6726. doi:10.1038/s41598-019-43143-4

Sullivan, K. R., Ravens, A., Walker, A. C., & Shepherd, J. D. (2025). "Arc - A viral vector of
memory and synaptic plasticity". *Curr Opin Neurobiol, 91*, 102979.
doi:10.1016/j.conb.2025.102979

Sun, C., Guo, R., Ye, X., Tang, S., Chen, M., Zhou, P., . . . Hu, H. (2024). Wybutosine
hypomodification of tRNAphe activates HERVK and impairs neuronal differentiation.
iScience, 27(5), 109748. doi:10.1016/j.isci.2024.109748

Takakura, M., Ishiguro, K., Akichika, S., Miyauchi, K., & Suzuki, T. (2019). Biogenesis and
functions of aminocarboxypropyluridine in tRNA. *Nat Commun, 10*(1), 5542.
doi:10.1038/s41467-019-13525-3

Tartaglia, G. G., Hollas, H., Havik, B., Vedeler, A., & Pastore, A. (2025). The RNA-Binding
Properties of Annexins. *J Mol Biol, 437*(4), 168933. doi:10.1016/j.jmb.2024.168933

Tassetto, M., Kunitomi, M., & Andino, R. (2017). Circulating Immune Cells Mediate a Systemic
RNAi-Based Adaptive Antiviral Response in Drosophila. *Cell, 169*(2), 314-325 e313.
doi:10.1016/j.cell.2017.03.033

Thery, C., Amigorena, S., Raposo, G., & Clayton, A. (2006). Isolation and characterization of
exosomes from cell culture supernatants and biological fluids. *Curr Protoc Cell Biol,*
Chapter 3, Unit 3 22. doi:10.1002/0471143030.cb0322s30

Toribio, V., & Yanez-Mo, M. (2022). Tetraspanins interweave EV secretion, endosomal network
dynamics and cellular metabolism. *Eur J Cell Biol, 101*(3), 151229.
doi:10.1016/j.ejcb.2022.151229

Tosar, J. P., & Cayota, A. (2020). Extracellular tRNAs and tRNA-derived fragments. *RNA Biol,*
17(8), 1149-1167. doi:10.1080/15476286.2020.1729584

Tosar, J. P., Segovia, M., Castellano, M., Gambaro, F., Akiyama, Y., Fagundez, P., . . . Cayota,
A. (2020). Fragmentation of extracellular ribosomes and tRNAs shapes the extracellular
RNAome. *Nucleic Acids Res, 48*(22), 12874-12888. doi:10.1093/nar/gkaa674

Tosar, J. P., Witwer, K., & Cayota, A. (2021). Revisiting Extracellular RNA Release, Processing,
and Function. *Trends Biochem Sci, 46*(6), 438-445. doi:10.1016/j.tibs.2020.12.008

Trigg, N. A., & Conine, C. C. (2024). Epididymal acquired sperm microRNAs modify post-
fertilization embryonic gene expression. *Cell Rep, 43*(9), 114698.
doi:10.1016/j.celrep.2024.114698

Upton, H. E., Ferguson, L., Temoche-Diaz, M. M., Liu, X. M., Pimentel, S. C., Ingolia, N. T., . . .
Collins, K. (2021). Low-bias ncRNA libraries using ordered two-template relay: Serial
template jumping by a modified retroelement reverse transcriptase. *Proc Natl Acad Sci U*
S A, 118(42). doi:10.1073/pnas.2107900118

Valadi, H., Ekstrom, K., Bossios, A., Sjostrand, M., Lee, J. J., & Lotvall, J. O. (2007). Exosome-
mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange
between cells. *Nature cell biology, 9*(6), 654-659. doi:10.1038/ncb1596

van Niel, G., D'Angelo, G., & Raposo, G. (2018). Shedding light on the cell biology of extracellular
vesicles. *Nat Rev Mol Cell Biol, 19*(4), 213-228. doi:10.1038/nrm.2017.125

Villarroya-Beltri, C., Gutierrez-Vazquez, C., Sanchez-Cabo, F., Perez-Hernandez, D., Vazquez,
J., Martin-Cofreces, N., . . . Sanchez-Madrid, F. (2013). Sumoylated hnRNPA2B1 controls
the sorting of miRNAs into exosomes through binding to specific motifs. *Nat Commun, 4*,
2980. doi:10.1038/ncomms3980

Wang, H., & Oxford, G. S. (2000). Voltage-dependent ion channels in CAD cells: A
catecholaminergic neuronal line that exhibits inducible differentiation. *J Neurophysiol,*
84(6), 2888-2895. doi:10.1152/jn.2000.84.6.2888

Wang, Z. Y., Wen, Z. J., Xu, H. M., Zhang, Y., & Zhang, Y. F. (2022). Exosomal noncoding RNAs
in central nervous system diseases: biological functions and potential clinical applications.
Front Mol Neurosci, 15, 1004221. doi:10.3389/fnmol.2022.1004221

Wei, S., Dai, X., Yuan, J., He, S., Shah, K., Guo, S., . . . Wang, Y. (2025). Quantitative Proteomics
Identifies Profilin-1 as a Pseudouridine-Binding Protein. *J Am Chem Soc, 147*(2), 1458-
1462. doi:10.1021/jacs.4c17659

Welsh, J. A., Goberdhan, D. C. I., O'Driscoll, L., Buzas, E. I., Blenkiron, C., Bussolati, B., . . .
Witwer, K. W. (2024). Minimal information for studies of extracellular vesicles
(MISEV2023): From basic to advanced approaches. *J Extracell Vesicles, 13*(2), e12404.
doi:10.1002/jev2.12404

Xie, Y., Chai, P., Till, N. A., Hemberger, H., Lebedenko, C. G., Porat, J., . . . Flynn, R. A. (2024).
The modified RNA base acp(3)U is an attachment site for N-glycans in glycoRNA. *Cell,*
187(19), 5228-5237 e5212. doi:10.1016/j.cell.2024.07.044

Xu, J., Erlendsson, S., Singh, M., Holling, G. A., Regier, M., Ibiricu, I., . . . Shepherd, J. D. (2024).
PNMA2 forms immunogenic non-enveloped virus-like capsids associated with
paraneoplastic neurological syndrome. *Cell, 187*(4), 831-845 e819.
doi:10.1016/j.cell.2024.01.009

You, Y., Zhang, Z., Sultana, N., Ericsson, M., Martens, Y. A., Sun, M., . . . Ikezu, T. (2023).
ATP1A3 as a target for isolating neuron-specific extracellular vesicles from human brain
and biofluids. *Sci Adv, 9*(37), eadi3647. doi:10.1126/sciadv.adi3647

Zhang, Y., Parmigiani, G., & Johnson, W. E. (2020). ComBat-seq: batch effect adjustment for
RNA-seq count data. *NAR Genom Bioinform, 2*(3), lqaa078. doi:10.1093/nargab/lqaa078

Zhao, X., Ma, D., Ishiguro, K., Saito, H., Akichika, S., Matsuzawa, I., . . . Suzuki, T. (2023).
Glycosylated queuosines in tRNAs optimize translational rate and post-embryonic growth.
Cell, 186(25), 5517-5535 e5524. doi:10.1016/j.cell.2023.10.026

FIGURE LEGENDS

Figure 1. CADs produce EVs enriched in tRNAs and snoRNAs

(A) Scheme of EV purification protocol. Conditioned medium was collected and subjected to

several round of centrifugation at increasing speeds to remove cells, cell debris, and large

vesicles. In the last step EVs were pelleted via ultracentrifugation at 100,000 x *g*.

(B) Nanoparticle tracking analysis of EVs from CADs. The plot shows the average (*n* = 2) number

of particles per mL on the *y* axis and their size on the *x* axis. The vertical dashed line indicates

the median size.

(C) Comparison of cellular and EV proteomes by mass spectrometry. Gray, significantly different

proteins (adjusted *p* < 0.1; *n* = 2); blue, EV markers; yellow, cellular markers.

(D) Top 9 enriched GO terms ("cellular component" aspect) associated with proteins significantly
enriched in EVs from (C).

(E) Gel electrophoresis (2% agarose-formaldehyde) comparing cellular *vs.* EV RNA. Individual
replicates were loaded in separate lanes.

(F) RNA content of CADs *vs.* EVs, measured by deep sequencing (LIDAR). Red, transcripts
significantly enriched in EVs [exRNAs; adjusted $p < 0.1$ and $\log_2$(EVs/cells) > 0; $n \geq 3$].

(G) Same as (F) with different classes of RNA highlighted by colors, as indicated.

Figure 2. A genome-wide screen identifies regulators of RNA secretion

(A) Scheme of the genetic screen. CADs were transduced with a lentiviral library containing
sgRNAs against most protein coding genes. Two (baseline) and 10 (readout) days later, cells and
EVs were harvested to measure sgRNA levels.

(B) Hypothetical gene categories with expected changes in sgRNA abundance over time in cells
and EVs. See text for explanation.

(C) Change in sgRNA abundance ($n = 4$ sgRNAs per gene), compared to their average
abundance in the corresponding sample (cells or EVs) at day 2. Individual biological replicates
(r1 and r2) are shown.

(D) Observed/predicted ratio of sgRNAs targeting putative positive (left) and negative (right)
regulators of RNA secretion, derived from MaGECK analysis. Significant hits (p value < 0.05) are
in red ($n = 2$). Candidates mentioned in the text are highlighted in blue.

(E) GO analysis (cellular component) of significant positive regulators from (D).

(F) Predicted (gray) vs. observed (red) sgRNA abundances in EVs ($\pm$ SEM), normalized to the
average predicted abundance (see methods), for retrovirus-derived proteins identified as
significant ($p < 0.05$) positive regulators from MaGECK analysis. Values for individual sgRNAs
are shown.

(G) Same as (F) but for RNA-modifying enzymes.

Figure 3. PUS1 facilitates the export of select exRNAs

(A) Western blot probing for PUS1, SDCBP, and β-TUB (control) in wild type (WT) and 2 different
Pus1$^{-/-}$ CAD clones.

(B) Differences in exRNA abundance between wild type and *Pus1*[-/-], as measured by LIDAR. Red,

transcripts showing significant difference ($n \geq 3$, adjusted p value < 0.1).

(C) Overlap between RNAs less abundant ("DOWN") in *Pus1*[-/-] cells compared to WT, either inside

the cells (grey, "cellular") or inside EVs (red, "extracellular").

(D) Ratios of RNA abundance in EVs and inside cells comparing wild type (blue) and *Pus1*[-/-] (gold)

CADs. Average and distributions are shown by the boxes and individual transcripts are shown in

points. RNAs were divided in the indicated classes and the p values were obtained with Student's

t tests. Data from ≥ 3 biological replicates.

(E) Scheme showing the different types of known tRNA-derived RNAs (tDRs).

(F) Changes in full-length tRNAs or tDRs between wild type and *Pus1*[-/-] in cells (grey) or EVs (red).

P values are from a Wilcoxon rank sum test. Data from ≥ 3 biological replicates.

Figure 4. PUS1-dependent Ψ selects RNA for secretion

(A) Scheme of the Ψ sequencing approach. Bisulfite-induced deletions were detected with a

modified LIDAR protocol which utilizes $MnCl_2$ during reverse transcription.

(B) Distribution of Ψ-modified RNAs among different classes in CADs and EVs. Data from 2

biological replicates.

(C) Ψ levels, inferred by the % of reads containing a deletion at the corresponding sites, for cellular

RNAs (gray) and exRNAs (red). P value is from a Wilcoxon rank sum test.

(D) Schematic representation of the mouse epididymis, highlighting the region of tissue and EVs

used for our analysis (cauda). The direction of sperm movement is indicated by the arrow.

(E) Same as (C) but for cauda tissue and corresponding EVs

(F) Levels of Ψ modification (% of total reads) at individual exRNA sites in wild type (x axis) vs

Pus1[-/-] (y axis) CAD EVs. PUS1-sensitive sites, where Ψ was reduced in *Pus1*[-/-] EVs by at least

25%, are in gold. Other Ψ sites are in gray. Data from 2 biological replicates.

(G) Ratio of the abundance of Ψ-containing RNAs in *Pus1*[-/-] CAD vs. WT cells (left) or EVs (right).

The RNAs were classified as PUS1-independent (gray, "others") and PUS1-sensitive (gold). Only

RNAs with at least one Ψ site detected in the EV fraction were considered. P values are from

Wilcoxon rank sum tests.

(H) % of molecules containing Ψ at the indicated bases according to bisulfite-LIDAR for four

example tRNAs in EVs from wild type (WT) and *Pus1*[-/-] CAD.

(I) Abundance ($\log_2$-converted, DEseq2 normalized counts) for the full-length tRNAs in (H) in wild

type (gray) or *Pus1*$^{-/-}$ (red) EVs. Asterisks indicate *p* value < 0.05 (calculated by DEseq2).

(J–K) Same as (H) and (I) but for the two indicated snoRNAs.

Figure 5. Ψ promotes exRNA export through MYL6

(A) Scheme of the sufficiency experiments shown in (B) and (C). CADs were transfected with

RNAs containing only Ψ or U at uridine positions for 6 hours. Their abundance inside and outside

cells was measured 24 hours later by RT-qPCR (B) or sequencing (C).

(B) Abundance of U- (gray) or Ψ-containing (yellow) RNAs 24 hours after transfection inside and

outside CADs 24 hours after transfection, measured by RT-qPCR. *P* values are from Student's *t*

tests.

(C) Export ratio (CPM_{cells} / CPM_{medium}) for U- (gray) or Ψ-containing (yellow) RNAs, 24 hours after

transfection into CADs. Six distinct sequences were tested, indicated by the numbers 1–6. The *p*

values are from Student's *t* tests.

(D) Comparison of protein abundance in the interactome of U- or Ψ-containing RNA, as measured

by mass spectrometry. Dark grey, significantly different proteins (adjusted *p* < 0.1; *n* = 2).

(E) Relative export ratio [export ratio(Ψ) / export ratio(U)] between U- and Ψ-containing RNAs in

the medium from CADs, treated with siRNAs against *Myl6* and control. The different RNA

sequences are identified by the same numbers as in (C). *P* values are from Student's *t* tests.

(F) Differences in exRNA abundance (from conditioned medium) between control and *Myl6*

knockdown (KD) as measured by LIDAR. Red, transcripts showing significant difference (*n* ≥ 5,

adjusted *p* value < 0.1).

(G) Ratio of the abundance of RNAs in *Myl6* knock down (KD) vs. control in cells (left) or

conditioned medium (right). The RNAs were classified as U-containing (gray, "others") and Ψ-

containing (blue). Only RNAs with at least one Ψ site detected in EVs were considered. *P* values

are from Wilcoxon rank sum tests.

(H) Proposed model for exRNA secretion. PUS1 marks specific RNAs with Ψ to select them as

exRNAs. Once deposited, Ψ provides a binding site for MYL6, which routes RNAs into EVs

formed via multivesicular bodies or by direct plasma membrane budding.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Additional analyses on exRNAs contained in EVs from CADs

(A) Overlap between the CAD EV proteome (see Fig. 1C) and proteins in the Vesiclepedia database (Kitti et al., 2024). The *p* value is from a hypergeometric test.

(B) RNA class distribution as % of transcripts significantly (adj. *p* value < 0.1) enriched in cells compared to EVs and *vice versa*.

(C) Distribution of average ($n \geq 3$) log-fold-changes between cells and EVs for C/D box and H/ACA box snoRNAs.

(D) Average ($n \geq 3$) insert size distribution of reads mapping to tRNAs in cells and EVs.

(E) Average ($n \geq 3$) tRNA and tDR distribution in cells and EVs.

Figure S2. Analyses and validations on the genome-wide screen for exRNAs

(A) RT-qPCR for sgRNA abundance (expressed as ratio to *Rpl32* RNA) in cells and EVs from sgRNA-expressing CAD and controls.

(B) Scheme of the logic underpinning the genome-wide screen. A fraction of sgRNAs expressed from a U6 promoter enter endogenous secretion pathways, for example by being loaded into intraluminal vesicles of multivesicular bodies (MVBs, derived from endosomal maturation) or directly by budding from the plasma membrane. This allows for the genotype of the originating cell to be determined via the sgRNA content of the EVs.

(C) Scheme of the LIDAR adaptation used to directly sequence the sgRNAs from the EVs.

(D) Number of sgRNAs that were detected (yellow) or not (blue) in cells and EV in each screen replicate. Individual samples are shown. Horizontal dashed line indicates the 95% of all sgRNAs included in the library.

(E) Comparison of the abundance of all individual sgRNAs inside cells vs. EVs at day 2 after transduction. Red, genes with significant difference ($n = 2$, adjusted *p* value < 0.1)

(F) Nucleotide representation at individual CRISPR RNA (crRNA) targeting sequence positions for all, preferentially exported, and preferentially retained sgRNAs. The bracket indicates the first 3 positions, enriched for Us in the exported sgRNAs.

(G) Average predicted (gray) and observed (red) values ($\pm$ SEM) for sgRNAs targeting *Sdcbp*. Circles represent values for the individual sgRNAs.

(H) Western blot for SDCBP and SUZ12 (control) in wild type and 2 different *Sdcbp*$^{-/-}$ CAD clones.

(I) Nanoparticle tracking analysis of EVs isolated from wild type and *Sdcbp*$^{-/-}$ CADs. The plot
shows the number of particles normalized to the number of starting million cells
[particles/(mL*million cells)] on the y axis and their size on the x axis.

(J) Average particles/(mL*million cells) ($\pm$ SEM), calculated as in (I), secreted by wild type and 2
different *Sdcbp*$^{-/-}$ CAD clones (*n* = 3). Individual replicates are shown.

(K) *Z* score-converted mRNA abundance (normalized DEseq2 counts) for the domesticated
retroviral genes showed in Fig. 2F, before ("CAD") and after ("dCAD") CAD cell differentiation into
a more mature neuronal form.

Figure S3. Validation of *Pus1* and analyses on exRNAs from *Pus1*$^{-/-}$ cells

(A) Scheme of secondary CRISPRi screen. Stable CADs, transduced with different CRISPRi
sgRNAs were mixed in equal numbers. Then, the abundance of sgRNAs in the pooled cells and
in the tissue culture medium (extracellular) was measured by LIDAR.

(B) Export rate of the indicated CRISPRi sgRNAs, calculated as a ratio of extracellular over
intercellular abundance. "Scr." denotes sgRNA scrambled control. *P* values are from Student's *t*
tests.

(C) Differences in cellular RNA abundance between wild type and *Pus1*$^{-/-}$, as measured by LIDAR.
Red, transcripts showing significant difference (*n* $\geq$ 3, adjusted *p* value < 0.1).

(D) Overlap between RNAs more abundant ("UP") in *Pus1*$^{-/-}$ cells compared to WT, either inside
the cells (gray, "cellular") or in EVs (red, "extracellular").

(E) Correlations in the change in RNA abundance between wild type and *Pus1*$^{-/-}$ CADs (*x axis*)
and EVs (*y* axis) for the RNAs significantly changed (adjusted p value < 0.1) in either set.
Spearman's correlation values are reported.

Figure S4. Validation of Ψ sequencing and analyses on exRNAs from *Pus1*$^{-/-}$ CADs

(A) LIDAR read densities (normalized to counts per millions) for individual nucleotides of an *in*
vitro-transcribed Ψ-containing RNA. Top, input; bottom, BID-treated

(B) Inferred % of Ψ based on % of deletion in LIDAR reads mapping to 18s rRNA from CAD RNA.
Top, input; bottom, BID-treated

(C–D) Overlap between Ψ-containing RNAs (C) or individual sites (D) identified in CADs and EVs.

P values are from hypergeometric tests.

(E) Denaturing agarose-formaldehyde (2%) electrophoresis of total tissue (cellular) and EV RNA

from cauda epididymis (individual replicates are shown).

(F) Distribution of transcripts containing PUS1-sensitive Ψ sites in CADs and EVs among the

indicated RNA classes. Data from 2 biological replicates.

(G) Distribution of reads containing PUS1-sensitive Ψ sites and mapping to tRNAs genes among

the indicated types of tRNA-derived species in cells and EVs.

(H) % of Ψ modification detected at each canonical tRNA position in wild type or *Pus1*[-/-] CAD EVs.

Data from the two different *Pus1*[-/-] clones are shown separately.

(I) % of Ψ at every canonical tRNA position (column) for every tRNA iso-acceptor containing a U

at position 27 (rows) in EVs from wild type (left) or *Pus1*[-/-] (right) CADs.

(J) Scheme of canonical Ψ sites on tRNAs. In blue, Ψ sites dependent on PUS1.

(K) Overlap between significantly reduced full-length tRNAs in *Pus1*[-/-] (red) and PUS1 full-length

tRNA targets (yellow) in EVs. *P* value is from a hypergeometric test

(L) Same as (K) but for snoRNAs.

For all panels data were from two biological replicates. In the case of *Pus1*[-/-] the two individual

knockout clones were used as combined replicates (n = 2 per clone).

Figure S5. Additional analyses for MYL6

(A) IP/beads values for U- and Ψ-RNAs interactomes measured by mass spectrometry. In red,

significantly different proteins (adjusted *p* < 0.1; *n* = 2).

(B) MYL6 protein abundance in CADs and EVs.

(C) Average predicted (gray) and observed (red) values (± SEM) for sgRNAs targeting *Myl6.*

Circles represent individual sgRNAs. The *p* value reported is from the MaGECK analysis.

(D) *Myl6* mRNA abundance relative to control (*Rpl32*). Bars show average ± SEM. Individual

replicates are shown.

(E) Differences in intracellular RNA abundance between control and *Myl6* knock down (KD) as

measured by LIDAR. Red, transcripts showing significant difference (*n* ≥ 5, adjusted *p* value <

0.1).

SUPPLEMENTAL TABLE LEGENDS

Table S1. Proteomics of CADs and EVs

Table S2. Genome-wide screen results

Table S3. RNA-binding proteins within positive regulators of exRNA biogenesis

Table S4. Proteomics results from U- and Ψ-containing RNA pulldowns

Table S5. Oligonucleotides used in this study

1376

Figure 1. CAD cells produce EVs enriched in tRNAs and snoRNAs

(A) Scheme of EV purification protocol. Conditioned medium was collected and subjected to several round of centrifugation at increasing speeds to remove cells, cell debris, and large vesicles. In the last step EVs were pelleted via ultracentrifugation at 100,000 x g.

(B) Nanoparticle tracking analysis of EVs from CADs. The plot shows the average (n = 2) number of particles per mL on the y axis and their size on the x axis. The vertical dashed line indicates the median size.

(C) Comparison of cellular and EV proteomes by mass spectrometry. Gray, significantly different proteins (adjusted p < 0.1; n = 2); blue, EV markers; yellow, cellular markers.

(D) Top 9 enriched GO terms ("cellular component" aspect) associated with proteins significantly enriched in EVs from (C).

(E) Gel electrophoresis (2% agarose-formaldehyde) comparing cellular vs. EV RNA. Individual replicates were loaded in separate lanes.

(F) RNA content of CADs vs. EVs, measured by deep sequencing (LIDAR). Red, transcripts significantly enriched in EVs [exRNAs; adjusted p < 0.1 and log2(EVs/cells) > 0; n ≥ 3].

(G) Same as (F) with different classes of RNA highlighted by colors, as indicated.

Figure 2. A genome-wide screen identifies regulators of RNA secretion

(A) Scheme of the genetic screen. CADs were transduced with a lentiviral library containing sgRNAs against most protein coding genes. Two (baseline) and 10 (readout) days later, cells and EVs were harvested to measure sgRNA levels.

(B) Hypothetical gene categories with expected changes in sgRNA abundance over time in cells and EVs. See text for explanation.

(C) Change in sgRNA abundance (n = 4 sgRNAs per gene), compared to their average abundance in the corresponding sample (cells or EVs) at day 2. Individual biological replicates (r1 and r2) are shown.

(D) Observed/predicted ratio of sgRNAs targeting putative positive (left) and negative (right) regulators of RNA secretion, derived from MaGECK analysis. Significant hits (p value < 0.05) are in red (n = 2). Candidates mentioned in the text are highlighted in blue.

(E) GO analysis (cellular component) of significant positive regulators from (D).

(F) Predicted (gray) vs. observed (red) sgRNA abundances in EVs (SEM), normalized to the average predicted abundance (see methods), for retrovirus-derived proteins identified as significant (p < 0.05) positive regulators from MaGECK analysis. Values for individual sgRNAs are shown.

(G) Same as (F) but for RNA-modifying enzymes.

Figure 3. PUS1 facilitates the export of select exRNAs

(A) Western blot probing for PUS1, SDCBP, and β-TUB (control) in wild type (WT) and 2 different Pus1-/- CAD clones.

(B) Differences in exRNA abundance between wild type and Pus1-/-, as measured by LIDAR. Red, transcripts showing significant difference (n ≥ 3, adjusted p value < 0.1).

(C) Overlap between RNAs less abundant ("DOWN") in Pus1-/- cells compared to WT, either inside the cells (grey, "cellular") or inside EVs (red, "extracellular").

(D) Ratios of RNA abundance in EVs and inside cells comparing wild type (blue) and Pus1-/- (gold) CADs. Average and distributions are shown by the boxes and individual transcripts are shown in points. RNAs were divided in the indicated classes and the p values were obtained with Student's t tests. Data from ≥ 3 biological replicates.

(E) Scheme showing the different types of known tRNA-derived RNAs (tDRs).

(F) Changes in full-length tRNAs or tDRs between wild type and Pus1-/- in cells (grey) or EVs (red). P values are from a Wilcoxon rank sum test. Data from ≥ 3 biological replicates

Figure 4. PUS1-dependent Ψ selects RNA for secretion

(A) Scheme of the Ψ sequencing approach. Bisulfite-induced deletions were detected with a modified LIDAR protocol which utilizes MnCl2 during reverse transcription.

(B) Distribution of Ψ-modified RNAs among different classes in CADs and EVs. Data from 2 biological replicates.

(C) Ψ levels, inferred by the % of reads containing a deletion at the corresponding sites, for cellular RNAs (gray) and exRNAs (red). P value is from a Wilcoxon rank sum test.

(D) Schematic representation of the mouse epididymis, highlighting the region of tissue and EVs used for our analysis (cauda). The direction of sperm movement is indicated by the arrow.

(E) Same as (C) but for cauda tissue and corresponding EVs

(F) Levels of Ψ modification (% of total reads) at individual exRNA sites in wild type (x axis) vs Pus1-/- (y axis) CAD EVs. PUS1-sensitive sites, where Ψ was reduced in Pus1-/- EVs by at least 25%, are in gold. Other Ψ sites are in gray. Data from 2 biological replicates.

(G) Ratio of the abundance of Ψ-containing RNAs in Pus1-/- CAD vs. WT cells (left) or EVs (right). The RNAs were classified as PUS1-independent (gray, "others") and PUS1-sensitive (gold). Only RNAs with at least one Ψ site detected in the EV fraction were considered. P values are from Wilcoxon rank sum tests.

(H) % of molecules containing Ψ at the indicated bases according to bisulfite-LIDAR for four example tRNAs in EVs from wild type (WT) and Pus1-/- CAD.

(I) Abundance (log2-converted, DEseq2 normalized counts) for the full-length tRNAs in (H) in wild type (gray) or Pus1-/- (red) EVs. Asterisks indicate p value < 0.05 (calculated by DEseq2).

(J–K) Same as (H) and (I) but for the two indicated snoRNAs.

Figure 5. Ψ promotes exRNA export through MYL6

(A) Scheme of the sufficiency experiments shown in (B) and (C). CADs were transfected with RNAs containing only Ψ or U at uridine positions for 6 hours. Their abundance inside and outside cells was measured 24 hours later by RT-qPCR (B) or sequencing (C).

(B) Abundance of U- (gray) or Ψ-containing (yellow) RNAs 24 hours after transfection inside and outside CADs 24 hours after transfection, measured by RT-qPCR. P values are from Student's t tests.

(C) Export ratio (CPM_{cells} / CPM_{medium}) for U- (gray) or Ψ-containing (yellow) RNAs, 24 hours after transfection into CADs. Six distinct sequences were tested, indicated by the numbers 1–6. The p values are from Student's t tests.

(D) Comparison of protein abundance in the interactome of U- or Ψ-containing RNA, as measured by mass spectrometry. Dark grey, significantly different proteins (adjusted $p < 0.1$; n = 2).

(E) Relative export ratio [export ratio(Ψ) / export ratio(U)] between U- and Ψ-containing RNAs in the medium from CADs, treated with siRNAs against Myl6 and control. The different RNA sequences are identified by the same numbers as in (C). P values are from Student's t tests.

(F) Differences in exRNA abundance (from conditioned medium) between control and Myl6 knockdown (KD) as measured by LIDAR. Red, transcripts showing significant difference (n ≥ 5, adjusted p value < 0.1).

(G) Ratio of the abundance of RNAs in Myl6 knock down (KD) vs. control in cells (left) or conditioned medium (right). The RNAs were classified as U-containing (gray, "others") and Ψ-containing (blue). Only RNAs with at least one Ψ site detected in EVs were considered. P values are from Wilcoxon rank sum tests.

(H) Proposed model for exRNA secretion. PUS1 marks specific RNAs with Ψ to select them as exRNAs. Once deposited, Ψ provides a binding site for MYL6, which routes RNAs into EVs formed via multivesicular bodies or by direct plasma membrane budding.

Figure S1. Additional analyses on exRNAs contained in EVs from CAD cells

(A) Overlap between the CAD EV proteome (see Fig. 1C) and proteins in the Vesiclepedia database (Kitti et al., 2024). The p value is from a hypergeometric test.

(B) RNA class distribution as % of transcripts significantly (adj. p value < 0.1) enriched in cells compared to EVs and vice versa.

(C) Distribution of average (n ≥ 3) log-fold-changes between cells and EVs for C/D box and H/ACA box snoRNAs.

(D) Average (n ≥ 3) insert size distribution of reads mapping to tRNAs in cells and EVs.

(E) Average (n ≥ 3) tRNA and tDR distribution in cells and EVs.

Figure S2. Analyses and validations on the genome-wide screen for exRNAs

(A) RT-qPCR for sgRNA abundance (expressed as ratio to Rpl32 RNA) in cells and EVs from sgRNA-expressing CAD and controls.

(B) Scheme of the logic underpinning the genome-wide screen. A fraction of sgRNAs expressed from a U6 promoter enter endogenous secretion pathways, for example by being loaded into intraluminal vesicles of multivesicular bodies (MVBs, derived from endosomal maturation) or directly by budding from the plasma membrane. This allows for the genotype of the originating cell to be determined via the sgRNA content of the EVs.

(C) Scheme of the LIDAR adaptation used to directly sequence the sgRNAs from the EVs.

(D) Number of sgRNAs that were detected (yellow) or not (blue) in cells and EV in each screen replicate. Individual samples are shown. Horizontal dashed line indicates the 95% of all sgRNAs included in the library.

(E) Comparison of the abundance of all individual sgRNAs inside cells vs. EVs at day 2 after transduction. Red, genes with significant difference (n = 2, adjusted p value < 0.1)

(F) Nucleotide representation at individual CRISPR RNA (crRNA) targeting sequence positions for all, preferentially exported, and preferentially retained sgRNAs. The bracket indicates the first 3 positions, enriched for Us in the exported sgRNAs.

(G) Average predicted (gray) and observed (red) values (SEM) for sgRNAs targeting Sdcbp. Circles represent values for the individual sgRNAs.

(H) Western blot for SDCBP and SUZ12 (control) in wild type and 2 different Sdcbp-/- CAD clones.

(I) Nanoparticle tracking analysis of EVs isolated from wild type and Sdcbp-/- CADs. The plot shows the number of particles normalized to the number of starting million cells [particles/(mL*million cells)] on the y axis and their size on the x axis.

(J) Average particles/(mL*million cells) (SEM), calculated as in (I), secreted by wild type and 2 different Sdcbp-/- CAD clones (n = 3). Individual replicates are shown.

(K) Z score-converted mRNA abundance (normalized DESeq2 counts) for the domesticated retroviral genes showed in Fig. 2F, before ("CAD") and after ("dCAD") CAD cell differentiation into a more mature neuronal form.

Figure S3. Validation of Pus1 and analyses on exRNAs from Pus1-/- cells

(A) Scheme of secondary CRISPRi screen. Stable CAD cells, transduced with different CRISPRi sgRNAs were mixed in equal numbers. Then, the abundance of sgRNAs in the pooled cells and in the tissue culture medium (extracellular) was measured by LIDAR.

(B) Export rate of the indicated CRISPRi sgRNAs, calculated as a ratio of extracellular over intercellular abundance. "Scr." denotes sgRNA scrambled control. P values are from Student's t tests.

(C) Differences in cellular RNA abundance between wild type and Pus1-/-, as measured by LIDAR. Red, transcripts showing significant difference (n ≥ 3, adjusted p value < 0.1).

(D) Overlap between RNAs more abundant ("UP") in Pus1-/- cells compared to WT, either inside the cells (gray, "cellular") or in EVs (red, "extracellular").

(E) Correlations in the change in RNA abundance between wild type and Pus1-/- CADs (x axis) and EVs (y axis) for the RNAs significantly changed (adjusted p value < 0.1) in either set. Spearman's correlation values are reported.

Figure S4. Validation of Ψ sequencing and analyses on exRNAs from Pus1-/- CADs

(A) LIDAR read densities (normalized to counts per millions) for individual nucleotides of an in vitro-transcribed Ψ-containing RNA. Top, input; bottom, BID-treated

(B) Inferred % of Ψ based on % of deletion in LIDAR reads mapping to 18s rRNA from CAD RNA. Top, input; bottom, BID-treated

(C–D) Overlap between Ψ-containing RNAs (C) or individual sites (D) identified in CADs and EVs. P values are from hypergeometric tests.

(E) Denaturing agarose-formaldehyde (2%) electrophoresis of total tissue (cellular) and EV RNA from cauda epididymis (individual replicates are shown).

(F) Distribution of transcripts containing PUS1-sensitive Ψ sites in CADs and EVs among the indicated RNA classes. Data from 2 biological replicates.

(G) Distribution of reads containing PUS1-sensitive Ψ sites and mapping to tRNAs genes among the indicated types of tRNA-derived species in cells and EVs.

(H) % of Ψ modification detected at each canonical tRNA position in wild type or Pus1-/- CAD EVs. Data from the two different Pus1-/- clones are shown separately.

(I) % of Ψ at every canonical tRNA position (column) for every tRNA iso-acceptor containing a U at position 27 (rows) in EVs from wild type (left) or Pus1-/- (right) CADs.

(J) Scheme of canonical Ψ sites on tRNAs. In blue, Ψ sites dependent on PUS1.

(K) Overlap between significantly reduced full-length tRNAs in Pus1-/- (red) and PUS1 full-length tRNA targets (yellow) in EVs. P value is from a hypergeometric test.

(L) Same as (K) but for snoRNAs.

For all panels data were from two biological replicates. In the case of Pus1-/- the two individual knockout clones were used as combined replicates (n = 2 per clone).

Figure S5. Additional analyses for MYL6

(A) IP/beads values for U- and Ψ-RNAs interactomes measured by mass spectrometry. In red, significantly different proteins (adjusted p < 0.1; n = 2).

(B) MYL6 protein abundance in CAD cells and EVs.

(C) Average predicted (gray) and observed (red) values (± SEM) for sgRNAs targeting Myl6. Circles represent individual sgRNAs. The p value reported is from the MaGECK analysis.

(D) Myl6 mRNA abundance relative to control (Rpl32). Bars show average   SEM. Individual replicates are shown.

(E) Differences in intracellular RNA abundance between control and Myl6 knock down (KD) as measured by LIDAR. Red, transcripts showing significant difference (n ≥ 5, adjusted p value < 0.1).