1   Neutrophils induce effective antibody responses to the pneumococcal conjugate vaccine by

2   inhibiting regulatory T cells

Essi Y. I. Tchalla, Anagha Betadpur, Andrew Y. Khalil, Elena Lopez, Manmeet Bhalla, Musea

Chang, Elizabeth A. Wohlfert and Elsa N. Bou Ghanem*

Author affiliations: Department of Microbiology and Immunology, University at Buffalo School of

Medicine, Buffalo, NY 14203, USA

*Address correspondence to: Elsa N. Bou Ghanem, 955 Main Street, Buffalo, NY, 14203.

Telephone: 716-829-2422. Email: elsaboug@buffalo.edu

Conflict-of-interest statement: The authors have declared that no conflict of interest exists.

Abstract

We recently found that neutrophils are required for production of protective antibodies in response to the pneumococcal conjugate vaccine (PCV), however, the mechanisms behind that are not known. Here, using a mouse model, we found that depletion of neutrophils at the time of vaccination in female mice led to aberrant T cell responses in the spleen resulting in increased regulatory T cells (Tregs) that was accompanied by a dysregulated cytokine environment. We found that following vaccination, neutrophils influx into secondary lymphoid organs and increase expression of T cell engaging/inhibiting markers. When we depleted Tregs in neutrophil deficient mice at the time of vaccination, the opsonic activity of the antibodies and the ability of the sera to protect naïve hosts against pneumococcal infection significantly increased compared to controls. Importantly, we examined responses following PCV administration in human female participants and found that PCV altered neutrophil phenotype and that *in vitro* coculture of neutrophils with PBMCs at one week following vaccination resulted in lower percentage and proliferation of FOXP3$^+$ T cells, showing the clinical relevance of this phenotype. This work reveals a new mechanism by which neutrophils ensure protective antibody responses to vaccination by suppressing Tregs.

Introduction

Neutrophils or polymorphonuclear cells (PMNs) are innate immune cells that are indispensable for the clearance of many infections and that have been traditionally studied as anti-microbial effector cells (1). However, in the past decade, PMNs have become increasingly appreciated for their ability to orchestrate adaptive immune responses (2, 3). In fact, they can affect B and T cells directly or via control of antigen presenting cells such as dendritic cells and macrophages (4-6). They can engage and stimulate T cells in response to antigens via expression of MHC-II, CD80, CD86 and/or CD11c (7-10) or inhibit T cells via production of IL-10, reactive oxygen species, arginase or by expressing PD-L1 or secreted protein acidic and rich in cysteine (SPARC) (11-15). The ectonucleotidases CD73 and CD39 have been associated with T cell inhibition via adenosine production and are also expressed on PMNs (16, 17). PMNs can also suppress B cell responses to protein antigens (18-20) or activate B cell responses to bacterial carbohydrates through secretion of B cell activating factor (BAFF), A proliferation-inducing ligand (APRIL), IL-21 or Pentraxin 3 (21, 22).

While most of the work examining PMN interaction with adaptive immune cells have used model antigens, a few studies have recently focused on the role of PMNs in vaccine responses (23). In an adenovirus-SIV vaccination model of Rhesus macaques, PMNs were found to migrate to the vaccine draining lymph nodes (vLNs) and acquire a B cell help phenotype as indicated by expression of BAFF, IL-21 and TNFα among others (23). PMNs were found to be required for effective immune responses to both tuberculosis and leishmaniasis live-attenuated vaccine models (24, 25). In the context of a clinically relevant vaccine, we showed that PMNs are required at the time of vaccination for induction of protective antibody responses to the licensed pneumococcal conjugate vaccine (PCV) Prevnar (26). In fact, depleting PMNs at the time of PCV vaccination led to a defect in antibody isotype switching to IgG2c and IgG3, a decline in antibody function, and decreased ability of the mice to control and survive a subsequent pneumococcal infection (26). However, the mechanisms by which PMNs control antibody responses to PCV

remain unclear. This is of high clinical relevance as PCV protects against infections caused by

the bacteria *Streptococcus pneumoniae* (pneumococcus), which remain a leading cause of

community acquired pneumonia globally, particularly in vulnerable populations such as older

adults (27). Understanding the mechanisms behind vaccine efficacy can lead to design of better

vaccines.

Conjugate vaccines such as PCV induce T follicular helper cells (Tfh) that aid the

production of antibodies by B cells (28, 29). In fact, PCV vaccination in humans induced the

formation of Tfhs, which was linked to enhanced antibody responses (30). Following vaccination

with a conjugate vaccine, germinal centers (GC) form and mature within secondary lymphoid

organs such as the lymph nodes (LNs) and spleen (31, 32). Inside GCs, polysaccharide-specific

B cells acquire conjugate antigens from the surface of follicular dendritic cells and process them

intracellularly for presentation to Tfhs (31-33). Interaction between the Tfhs and GC B cells

triggers B cell differentiation into antibody secreting cells and enhances antibody production, class

switching and affinity maturation (34, 35). These antibodies are required for efficient host

protection against infections as they facilitate binding, uptake and killing of bacteria by phagocytes

(36, 37). Antibody responses by B cells are also highly influenced by cytokines, which are

secreted by Tfhs as well as other T cell subsets (28, 38, 39).

A subset of T cells known to control immune responses are regulatory T cells (Tregs).

Tregs have suppressive capacity and play an essential role in homeostasis and immune tolerance

(40). In the context of vaccination, Tregs have been shown to limit protective immune responses

(41-46). In multiple HIV and cancer models of vaccination, Tregs were shown to dampen both

cellular and humoral immune responses, leading to viral replication or tumor progression (47, 48).

In a pneumococcal polysaccharides model of vaccination, blocking the Treg effector molecule

CTLA-4 enhanced antibody and cytokine responses to the polysaccharides (49). In humans,

higher Tregs in the PBMC of older adults were associated with lower CD4[+] T cell responses to

immunogenic pneumococcal proteins (50). A subset of Tregs that gain access to B cell follicles

are termed T follicular regulatory cells or Tfr (51, 52). Tfrs display phenotypic markers of T

follicular helper cells (Tfhs), in addition to their expression of FOPX3 (51, 53). They play a role in

humoral responses to antigens that is context dependent and have been shown to inhibit B cell

responses and accelerate GC contraction (54-56). Several studies have looked at the relationship

between Tregs and PMNs in models of inflammation and disease and found that Tregs modulate

PMN recruitment to the site of inflammation (57-59). PMNs in turn also promote Treg recruitment

in autoimmune and cancer models, and their depletion leads to a decrease in Tregs (60-63). The

interaction of PMNs and Tregs during vaccine responses (including PCV) has not been studied.

In this study, we tested the effect of PMNs on adaptive immune responses to PCV using

a preclinical murine model of vaccination as well as by examining responses in vaccinated human

participants. We found that following vaccination, PMNs acquire a T cell engaging and

suppressing phenotype and suppress Tregs, which is required for the production of protective

antibodies. This study reveals a new mechanism by which PMNs shape vaccine responses

through the control of Tregs and provides novel targets for the design of more effective vaccines

against bacterial infections.

Results

PMNs home into secondary lymphoid organs in response to PCV

In prior work, we showed that PMNs are required at the time of PCV administration for the induction of effective antibody response and optimal host protection against pneumococcal disease (26). There is a well-documented sex-based difference in responses to PCV where females elicit higher antibody responses and vaccine efficacy compared to males (64-66) and we recapitulated this in preclinical murine models (67). Therefore, to understand mechanisms of protection, we focused our studies on female hosts, that mount higher humoral and cellular responses to PCV and are better protected against subsequent pneumococcal infection compared to male hosts (67).

To better understand how PMNs control PCV efficacy, we first asked whether they are recruited to secondary lymphoid organs (spleen and vaccine draining lymph nodes (vLNs)) in response to vaccination. We looked at expression of migration markers CXCR4 and CCR7 on circulating PMNs 24 hours post vaccination as they are involved in PMN trafficking to the lymph nodes via the lymphatic system or high endothelial venules (Fig. S1 & Fig. 1A) (68, 69). We found that at baseline, circulating PMNs express relatively low levels of both CXCR4 and CCR7 and upon vaccination, expression of CCR7 increases (Fig. 1A). This higher CCR7 expression coincided with a decrease in the number of circulating PMNs 18 to 24 hours post vaccination and an influx of PMNs in the spleen and vLNs at those time points respectively (Fig. 1B). Following the peak of PMN influx into the spleen and vLNs, PMN numbers return to baseline by 48 hours post vaccination (Fig. 1B). These findings demonstrate that PMNs are rapidly recruited to secondary lymphoid organs in response to PCV.

PMNs do not affect B cell numbers in response to PCV

To start investigating how PMNs induce protective antibodies to PCV, we first examined changes to the B cell compartment. We depleted PMNs at the time of vaccination as presented

in Fig 2A and previously described (26) and investigated B cell responses in secondary lymphoid

organs on days 7, 14 and 21 post vaccination (Fig. S2A). Upon PMN depletion, there was no

change in the percentage and numbers of total B cells (B220$^+$) in the spleen compared to isotype

treated mice at any of the timepoints investigated (Fig. S3A). Similarly, no effect of PMN depletion

was observed on GC B cells (CD38^{lo}CD96$^+$) and plasmablasts (CD138$^+$) (Fig. S3B & S3C). In

prior work, we had observed a defect in isotype switching, opsonic and binding capacities of

antibodies in response to PCV in PMN deficient hosts (26), indicating a possible defect on somatic

hypermutation (SHM) and/or class switch recombination (CSR). Therefore, on days 7 post

vaccination, we assessed B cell expression of Activation Induced Cytidine Deaminase (AID), an

enzyme required for both SHM and CSR (70, 71) (Fig. S2B). For both total and GC B cells, there

was an increase in AID expression in mice following vaccination (Fig. S4). However, depleting

PMNs had no effect on the percentage of AID$^+$ B cells or the expression of AID by B cells (Fig.

S4). These findings suggest that PMNs are not directly acting on B cells during vaccination.

PMNs affect MHC-II expression by macrophages and dendritic cells in response to PCV

We then looked at the effect of PMNs on other immune cells important for vaccine

responses such as antigen presenting cells (APCs) (7, 72). For that, we depleted PMNs at the

time of vaccination (Fig. 2A) and analyzed the number of splenic macrophages (F4/80$^+$) and

dendritic cells (DC-CD11c$^+$) or their respective expression of antigen presenting molecules MHC-

II, CD80 and CD86 on days 7, 14 and 21 post vaccination (Fig. S5 & S6). Overall, we found that

depletion of PMNs had no effect on the percentages of both macrophages and DCs present in

the spleen (Fig. S6), except for a slight increase in macrophages in PMN-depleted mice compared

to PMN-sufficient mice at 21 post vaccination (Fig. S6A). A lower percentage of both

macrophages and DCs exprssed MHC-II in the spleen of PMN-depleted mice at day 14 post

vaccination compared to PMN-sufficient counterparts with no effect on CD80 and CD86

expression (Fig. S6). These findings suggest depletion of PMNs may impair antigen presentation.

Absence of PMNs at the time of vaccination alters the cytokine environment in the spleen

The cytokine environment in secondary lymphoid organs controls the quality of antibodies produced (73). Therefore, we next looked more broadly at the overall inflammatory environment in the spleen in the presence or absence of PMNs in vaccinated mice using a cytokine protein array. We found that overall, in the absence of PMNs at the time of vaccination, there was an increase in inflammatory mediators in the splenic milieu compared to when PMNs are present (Fig. S7). This increase in inflammation was seen at day 7 and resolved by day 14 through 21 post vaccination (Fig. S7), timepoints that coincide with return of PMNs after depletion (26). These data show that the absence of PMNs at the time of vaccination leads to an altered inflammatory response in the spleen, suggesting a regulatory role for PMNs during PCV.

We had observed a defect in isotype switching to IgG2 and IgG3 antibodies in the absence of PMNs (26), therefore, we directly tested whether PMNs controlled effective antibody response to PCV through IFNγ, as this cytokine is required for class switching to those isotypes (34, 74). We measured the percentage of IFNγ^+ PMNs in the spleen 18 hours post vaccination following *ex vivo* stimulation with PMA, heat-killed *S. pneumoniae* or PCV (Fig. S8). We found that more PMNs express IFNγ when stimulated with PMA, however, there was no change in the percentage of IFNγ^+ PMNs following vaccination and upon restimulation (Fig. S9A). Similarly, total IFNγ levels in the spleen within the first 28 days of vaccination were comparable between PMN-depleted and PMN-sufficient mice (Fig. S9B). Additionally, when we blocked IFNγ *in vivo* at the time of vaccination using an αIFNγ antibody following the same schedule as for the depletion of PMNs shown in Fig. 2A, there was no effect on bacterial burden in the lung and blood at 24 hours post infection (Fig. S9C) or on host survival (Fig. S9D) compared to the isotype-treated mice. To verify that the blocking antibody was functioning, we measured isotype switching to IgG2 which is controlled by IFNγ (34), and founded that blocking IFNγ led to an increase in IgG1 antibodies and

208   decrease in IgG2b and IgG2c (Fig. 9E). In summary, these data indicate that overall, the control

209   of pneumococcal antibody responses by PMNs is not mediated by IFNγ.

211   **PMNs acquire a T cell regulating phenotype upon entry into secondary lymphoid organs**

212   To better understand how PMNs control PCV efficacy, we then assessed how they

respond to vaccination. PMNs have been shown to acquire different phenotypes depending on

entry into organs/tissues and the composition of the tissue microenvironment (75, 76). Therefore,

we examined PMN phenotype within the first 24 hours post vaccination when they influx into

secondary lymphoid organs (Fig. 1B). As the cytokine responses in the spleen suggested that

PMNs play a regulatory role, we looked at the expression of various T cell engaging (MHC-II,

CD80, CD86 and Cd11c) and suppressing (PDL-1, CD73, CD39 and ARG1) markers on PMNs

(Fig. S1). In all sites investigated, overall, expression of both T cell engaging and suppressing

markers on PMNs did not change upon vaccination except for a decrease in PD-L1$^+$ PMNs in the

blood and spleen (Fig. 3 & 4). However, there were tissue specific differences (Fig. 5). Expression

of CD80, CD86, CD11c MHC-II and ARG1 were higher on PMNs in secondary lymphoid organs

compared to circulating PMNs in both naïve and vaccinated mice (Fig. 3, 4 and 5B). These data

are consistent with the literature showing that entry into tissues instructs PMNs to acquire organ

specific phenotypes (75, 76).

226   PMNs expressing MHC-II have been reported to express a regulatory phenotype (77).

Therefore, we assessed the expression of CD73, CD39, ARG1, PD-L1 and SPARC on MHC-II$^+$

versus MHC-II$^-$ PMNs (Fig. S1). In all tissues investigated, there are fewer MHC-II$^+$ PMNs

compared to MHC-II$^-$ PMNs (Fig. S10A). Those MHC-II$^+$ PMNs in the spleen expressed lower

ARG1, higher SPARC and PD-L1 and similar levels of CD73 and CD39 compared to their MHC-

II$^-$ PMNs in the spleen (Fig. S10B). Overall, all the above data show that upon influx into the

spleen and vLNs, PMNs acquire the ability to engage and suppress T cells following PCV

vaccination.

PMNs do not affect T helper cell responses to PCV

To test the effect of PMNs on T cell responses, we depleted PMNs at the time of vaccination (Fig 2A) and measured the percentages and numbers of T helper 1 (Th1) (CD4$^+$TCRβ$^+$FOXP3$^-$Tbet$^+$), Th2 (CD4$^+$TCRβ$^+$FOXP3$^-$GATA3$^+$), Th17 (CD4$^+$TCRβ$^+$FOXP3$^-$RORγt$^+$) and Tfh (CD4$^+$TCRβ$^+$CXCR5$^+$PD1$^+$) cells and in the spleen and vLNs, 7-, 14- and 21-days post vaccination (Fig. S11). Except for an increase in the percentage of Th1 cells in the spleen of PMN-depleted mice (Fig. S12A), we found that there was no difference in the percentages or numbers of Th2, Th17 and Tfhs between vaccinated PMN-depleted and PMN-sufficient mice at any the timepoints investigated in the spleen (Fig. S12B-D). As examining total CD4$^+$ T cells may mask changes within the smaller antigen experienced population, we also looked at antigen experienced cells by gating on CD49d$^+$CD11a$^+$ T cells (Fig. S11A) (78). Similarly, we found no effect of PMN depletion on antigen experienced Th1, Th2, Th17 and Tfhs in the spleen (Fig. S13).

As depleting PMNs had no effect on the amount of T helper cells (Fig. S12 & S13), we asked if their ability to produce cytokines was impaired. We examined cytokine responses *ex vivo* using PMA/Ionomycin as positive stimulant for cytokine production. Using flow cytometry, we measured expression of IFNγ, IL-4, IL-17, TGFβ and IL-10 by splenic CD44^{hi} activated CD4$^+$ T cells at day 14 post vaccination (Fig. S14). We used the protein Latency Associated Peptide (LAP) as a proxy for TGFβ production since it acts as a chaperone for TGFβ and their levels correlate (79, 80). The cytokine response was skewed towards production of IFNγ and LAP, both in the presence and absence of PMNs, as the percentage of CD4+TCRβ+ T cells expressing these cytokines was higher than the other cytokines investigated (Fig. S15). Overall, PMA/Ionomycin stimulation increased LAP, IFNγ and IL-4 expression by CD44^{hi} CD4$^+$ T cells above the no

stimulation condition (Fig. S15A-C). However, PMN depletion did not affect expression of any of

the cytokines by T cells in comparison to PMN-sufficient mice (Fig. S15).

PMNs inhibit Treg expansion in the spleen in response to PCV

We next assessed the effect of PMNs on regulatory T cells (Tregs) (CD4$^+$TCRβ$^+$FOXP3$^+$)

and T follicular regulatory cells (Tfrs) (CD4$^+$TCRβ$^+$CXCR5$^+$PD1$^+$FOXP3$^+$) following PCV (Fig. 6).

We saw an increase in the percentage of Tregs in the spleens of PMN-depleted mice compared

to PMN-sufficient mice at day 14 post vaccination (Fig. 6A). In PMN-depleted mice, there was

also an increase in the percentage and numbers of Tfrs at days 21 & 7 post vaccination

respectively (Fig. 6B). When looking at the antigen experienced compartment, there was an

increase in Tregs and Tfrs at day 14 post vaccination in the spleen of PMN-depleted mice

compared to PMN-sufficient mice (Fig. 6C & 6D). These data suggest that PMNs inhibit the

expansion of regulatory T cells in the spleen following PCV.

PMNs inhibit Treg expansion in response to PCV *in vitro*

To directly assess the effect of PMNs on Tregs, we used *in vitro* assays. For that, we

performed coculture assays of splenocytes isolated from day 1-vaccinated mice in the presence

or absence of PMNs using heat-killed bacteria or αCD3 as stimulants for 48 hours. PMNs were

removed from splenocytes using positive isolation magnetic sorting. Similar to what we observed

in vivo, in the absence of PMNs, there was an increase in the percentage of Tregs in response to

stimulation with heat-killed *S. pneumoniae* (Fig. 7). PMN removal did not alter other T helper cell

subsets (Fig. S16). This recapitulates the *in vivo* findings and suggests that PMNs control Tregs

response to bacterial polysaccharides.

Reducing Tregs in PMN-depleted hosts rescues antibody responses

Next, we tested whether the increase in regulatory T cells is responsible for the defective

antibody responses seen in the absence of PMNs at the time of vaccination (26). To investigate

this, we depleted Tregs using an αCD25 antibody (81) either alone or in conjunction with PMN

depletion using αLy6G (Fig. 2B). We achieved ~90% depletion of PMNs in the blood and ~40-

50% depletion of Tregs in both spleen and blood (Fig. S17A & S17B). Moreover, αCD25 and

αLy6G injections had no effect on percentages of Th1, Th2, Th17, Tfhs and B cells in the spleen

(Fig. S17C).

We then followed antibody levels upon Treg and/or PMN depletion in the blood of mice at

weeks 0, 2 & 4 post vaccination. We found an increase in total IgG against pneumococcal

polysaccharides in PMN-& Treg-depleted mice compared to PMN-depleted mice at 4 weeks post

vaccination (Fig. 8A). We next investigated the ability of antibodies in the sera to bind the surface

of the pneumococcus (Fig. S18). In all mouse groups, vaccination significantly increased binding

of IgG to the bacteria compared to baseline (Fig. 8B). As previously reported, sera from

vaccinated PMN-depleted mice had significantly lower IgG binding compared to sera from

vaccinated PMN-sufficient mice (Fig. 8B) (26). Interestingly, depletion of Tregs in PMN-depleted

mice rescued IgG binding in the sera to similar levels to that of PMN-sufficient vaccinated mice,

while depletion of Tregs alone had no effect (Fig. 8B). We also looked at the ability of antibodies

in the sera to promote opsonophagocytic killing of the pneumococcus by PMNs isolated from

naïve mice. As expected, there was an increase in bacterial killing by PMNs when sera from

vaccinated versus naïve mice were used as opsonins (Fig. 8C). We also found that sera from

PMN-depleted mice induced lower bacterial opsonophagocytic killing by PMNs compared to sera

from PMN-sufficient mice (Fig. 8C). Similar to above, depletion of Tregs in PMN-depleted mice

rescued the ability of the sera to induce opsonophagocytic bacterial killing to levels similar to

PMN-sufficient vaccinated mice, while depletion of Tregs alone did not affect opsonic capacity

(Fig. 8C).

To confirm these findings, we used a different model of Treg depletion for assessment of

antibody responses to PCV using diphtheria toxin (DT) mediated depletion (82). For that, we used

DT injections to deplete Tregs either alone or in conjunction with PMN depletion in DEREG mice

(83) (Fig. S19A). We achieved ~25% depletion of Tregs in the spleen (Fig. S19B). Treg depletion

in PMN-depleted DEREG mice increased their IgG levels 4 weeks post vaccination compared to

vehicle treated PMN-depleted mice (Fig. S19C). The above data suggest that the increase in

Tregs seen in PMN-depleted mice contributes to the defective antibody response observed in

these mice as reduction of the Tregs rescues the levels and function of the antibodies to similar

levels seen in PMN-sufficient vaccinated mice.

Reducing Tregs in PMN-depleted hosts rescues the protective capacity of the sera against

pneumococcal infection

Finally, we assessed the protectiveness of the sera from PMN- and/or Treg-depleted mice

in vivo against infection by adoptively transferring sera from each group into naïve mice. Recipient

mice were then infected 1hr later with 3×10^5 CFU of *S. pneumoniae* TIGR4 strain, assessed for

blood bacterial burden 48 hours post infection and monitored for survival. Recipients of sera from

vaccinated PMN-sufficient mice were able to control bacterial numbers better than the ones

receiving sera from vaccinated PMN-depleted mice with 50% and 87.5% of the mice becoming

bacteremic respectively (Fig. 9A). Receiving sera from PMN- & Treg-depleted mice rescued the

ability of the recipients to control infection with only 25% of the mice becoming bacteremic (Fig.

9A). Additionally, receiving sera from Treg-depleted mice did not affect control of blood bacterial

burden (Fig. 9A). Looking at survival, only 10% of recipients of sera from naïve mice survived the

infection compared to ~70% of the recipients of sera from vaccinated PMN-sufficient mice (Fig.

9B). Receiving sera from PMN-depleted mice resulted in only 30% survival, compared to ~81%

survival of mice that received sera from PMN- & Treg-depleted mice (Fig. 9B). The above data

suggest that reduction of Tregs in PMN-depleted mice rescues the protectiveness of the sera to

levels similar to PMN-sufficient vaccinated mice.

Vaccination of human participants alters the phenotype of circulating PMNs

As PMN responses vary significantly across species (84), we wanted to test the clinical

relevance of these findings. To do so we recruited unvaccinated adult female participants,

administered PCV to them, and monitored the phenotype of PMNs in the circulation over time in

response to vaccination by examining the expression of T cell engaging and inhibiting markers

using flow cytometry (Fig. S20). To test if PMNs respond to PCV and alter their phenotype, we

used the OMIQ software to perform opt-SNE analysis to visualize the data, and cluster them using

FLOWSOM. We found that PMNs separated into different clusters with varying levels of maturity

(based on CD10 expression, Fig. 10A and S21) with cluster 6 being highest followed by 3, 2, 4,

1, 9, 7, 5, and 8 the lowest. Clusters 6 (most mature) was the only cluster that expressed PDL-1

and had high levels of all the other markers, while cluster 8 (least mature), did not express any of

the markers we tested (Fig. 10A, 10C and S21). Upon vaccination, we observed a shift in the

abundance of circulating clusters where cluster 1, 2 and 6 decreased in abundance, cluster 3, 4

and 5 increased, while cluster 7 and 8 did not change (Fig. 10B). The clusters that decreased

from the circulation expressed the highest levels of CD39, ARG1 and HLADR (Fig. 10C) and may

be homing to the tissues as we observed in mice. When we examined the expression levels of

the various markers at week 1 following vaccination, we found that expression of HLADR, CD86,

CD39 and PDL-1 increased on all the clusters that expressed them, CD73 either increased or

decreased depending on the cluster, while ARG1 decreased (Fig. 10C). Overall, these findings

suggest that in response to PCV in humans, the phenotype of circulating PMNs changes, where

particular clusters increase expression of T cell engaging and suppressing markers and possibly

exit from the circulation into tissues.

PMNs from PCV immunized donors inhibit FOXP3$^+$ T cell expansion *in vitro*

Finally, we tested whether PMNs from vaccinated participants can suppress FOXP3$^+$ CD4$^+$ T cells. For that, at one week post vaccination, a timepoint where we observed phenotypic changes in PMNs in response to vaccination (Fig. 10), we performed an *in vitro* assay where PBMCs from each participant were cocultured with or without PMNs (from the same donor) and stimulated with PCV for 48 hours (Fig. S22). Similar to what we found in the preclinical model, the presence of PMNs resulted in reduction of FOXP3$^+$ T cells. Addition of PMNs to PBMC cultures lead to a decrease in the percentages of total and antigen experienced FOXP3$^+$ T cells in 67% of participants (Fig. 11A). We also observed a decrease in Ki67$^+$ antigen experienced FOXP3$^+$ T cells in the presence of PMNs in 100% of the participants (Fig. 11B), suggesting that PMNs were suppressing proliferation. These data suggest that PMNs inhibit FOXP3$^+$ T cell responses in humans following PCV administration.

Discussion

There are a growing number of studies investigating the effect of neutrophils on adaptive immune responses with a few of them focusing on vaccine responses (23-25). In this study, we report a new mechanism by which PMNs control antibody responses to the clinically relevant PCV by controlling Treg numbers. In fact, when neutrophils were depleted at the time of PCV vaccination, there was an expansion in the number of Tregs that resulted in a defect in protective antibody responses (26). Reduction of Tregs in PMN-deficient vaccinated mice boosted antibody number, function, and protective efficacy in the sera. This work demonstrates that PMNs drive production of functional antibodies against pneumococci in response to PCV by limiting the expansion of immune suppressive Tregs.

While the effect of PMNs on Tregs has not been investigated in the context of clinically relevant vaccines, several studies have looked at the relationship between Tregs and PMNs in models of inflammation and disease (57-63). Here we report that PMNs can inhibit Tregs, which is in contrast to prior work demonstrating that PMNs promote Tregs in models where high level of inflammation is induced (60, 61, 63). The exact way by which PMNs inhibit Tregs in response to PCV remains to be elucidated, however multiple mechanisms could be involved. Some possibilities could be via control of the transcription factor FOXP3, a master regulator of Treg induction and function (85, 86) or control of cytokines required for Treg differentiation and viability. In traumatic brain injury patients, release of serine proteases (neutrophil elastase and proteinase 3) by infiltrating PMNs cleaved cytokine receptors on the surface of T cells including IL-2Rα, the IL-2 receptor subunit that is important for IL-2 signaling in Tregs (87, 88). In addition, those proteases can cleave cytokines themselves including TGFβ and IL-2, leading to limited activation of T cell response (89, 90). In response to PCV, PMNs may similarly inhibit Treg induction by release of their proteases to deplete Treg-inducing cytokines and/or their respective receptors on the surface of Tregs. Weak antigenic TCR stimulation on naïve T cells in the presence of TGFβ can lead to Treg production via induction of FOXP3 (91, 92). We found that in the absence of

PMNs, a lower percentage of macrophages and DCs express MHC-II, which can weaken the TCR-MHC-II interaction by lowering antigen density on the surface of the antigen presenting cell (93). Therefore, the presence of PMNs at the time of vaccination can lead to Treg suppression by increasing the strength of APC interaction with T cells through high MHC-II expression. Signaling of retinoic acid (RA) in naïve T cells enhances TGFβ-mediated induction of Tregs (94) and RA boosts PMN anti-bacterial functions (95). It is possible that RA sequestration by PMNs limits its availability for Treg induction. PMNs could also be inhibiting Tregs through release of their granule component cathelicidin antimicrobial peptide, the human form of which (LL-37) was shown to induce Treg apoptosis *in vitro* (96). Alternatively, PMNs could be suppressing Tregs through the release of miRNA containing extracellular vesicles (EVs). PMN-derived EVs containing miRNAs have been shown to modulate responses in other cells including macrophages and endothelial cells (97-99). The absence of miRNA-146a has been shown to promote Treg expansion, and other miRNAs are known to control Treg induction (100, 101). PMNs, which express miRNA-146a among others (102, 103), could use such mechanism to limit Treg expansion. All these potential mechanisms can be avenues of future research for this work.

The findings here suggest that the expanded Tregs in PMN-depleted mice are detrimental and contribute to the defective antibody responses. In general, Tregs have been shown to negatively affect immune responses to vaccines. Depletion of Tregs using αCD25 in vaccination models against herpes simplex virus (HSV) and *Plasmodium berghei*, increased CD8$^+$ T cell responses to their respective vaccine epitopes and led to increase protection against subsequent challenge with the respective pathogens (43, 104). In humans, responses to an experimental vaccine of dendritic cells loaded with HIV peptides (105) found that patients with lower levels of HIV-specific Tregs displayed better T-effector cell responses following vaccination (48). In the context of *S. pneumoniae*, blocking of CTLA-4, an effector molecule on Tregs, enhanced levels of antibodies against pneumococcal polysaccharides in immunized mice and higher Tregs in older adults were associated with decreased proliferation of CD4$^+$ effector T cells in response to the

pneumococcal protein AliB (49, 50). Additionally, Tfrs, which we found are increased in PCV-

vaccinated PMN-depleted mice, have been shown to inhibit B cells responses and promote the

contraction of GC reactions (54, 56). The findings here, showing an improvement of vaccine

response upon depletion of the expanded Tregs in PMN-depleted mice, are in line with the

reported inhibitory role of Tregs in vaccine responses.

There have been several studies examining how Tregs and Tfrs inhibit B cell responses

(106). They can affect B cell number, differentiation, and antibody responses via release of

soluble mediators such as the neuropeptide neuritin, IL-10 and TGFβ or through contact

dependent mechanisms involving CTLA4, PD1 and granzymes (106-108). This suppression can

lead to a decrease in B cell expression of activation-induced deaminase (AID) (109). However, B

cell numbers and AID expression were not affected in this study. It is possible that the DNA

binding capacity or the enzymatic activity of AID are affected instead. AID activity is regulated by

its phosphorylation status which can either enhance or inhibit its enzymatic activity, depending on

the amino acid being phosphorylated (110, 111) by protein Kinase A (PKA) (112-115). It is

therefore possible for Tregs to be regulating AID phosphorylation in response to PCV, rather than

overall expression through modulation of cAMP as they have been shown to do in DCs and

effector T cells (116).

We found that the phenotypes observed in the preclinical model are also found in human

participants, highlighting the clinical relevance of this work. Upon vaccination of healthy human

donors, PMNs in the circulation changed phenotype suggesting they are responsive to

vaccination. Using peripheral blood collected from donors after PCV vaccination for *in vitro*

cocultures, addition of PMNs to PBMCs lead to a reduction in FOXP3$^+$ T cells and their

proliferation, recapitulating the Treg inhibiting property of PMNs seen in the mouse models.

However, further clinical studies are needed to assess the *in vivo* role of PMNs in protecting

human hosts against pneumococcal disease, including linking PMN phenotype to antibody

responses. Nevertheless, these findings can impact the design of improved pneumococcal

vaccine formulations that offer better efficacy, especially in the more vulnerable populations.

Despite their availability, pneumococcal vaccines do not mitigate the growing global burden of *S.*

pneumonia infections (27, 65, 117). This is due in part to the fact that the vaccines do not protect

all populations equally, where male donors and aged individuals mount lower responses

compared to female counterparts. Interestingly, Tregs are reported to be higher in males

compared to sex-matched females and higher in older adults compared to younger people (118-

120). In this work, we identified the role of the PMN-Treg axis in controlling antibody responses

in females that can be targeted for improvement of current vaccine formulations moving forward.

While further studies are needed to tease apart the specific mechanisms behind this inhibition of

Tregs by PMNs, the findings have important implications for development of future vaccine

strategies to improve efficacy to infections. Indeed, this PMN-Treg axis can be further studied in

the context of sex and age-driven differences in vaccine responses (121, 122) and can be targeted

to increase vaccination efficacy in the vulnerable older adults and male populations. The exact

mechanisms of Treg inhibition by PMNs need to be elucidated to pinpoint what pathways or

molecules to target specifically in the future.

Methods

Mice

Adult (10-12 weeks old) female C57BL/6 mice were purchased from Jackson Laboratories (Bar Harbor, Maine). Our study focused on female hosts, as there is a well-documented sex-based difference in responses to PCV where females elicit higher antibody responses and are better protected from subsequent infection compared to males (64, 67, 123). Therefore, we examined the mechanisms behind vaccine efficacy in the better protected hosts. Ongoing work in our lab is focused on sex-based difference in PMN responses in vaccinated hosts (67) which is beyond the scope of this study. Adult (10-16 weeks old) C57BL/6-Tg(Foxp3-DTR/EGFP)23.2Spar/Mmjax DEREG mice were a kind gift from Elizabeth Wohlfert and have been previously described in detail (82, 83). All mice were housed in a specific-pathogen-free environment at the University at Buffalo Jacobs School of Medicine and Biomedical Sciences animal facility.

Immunization and sera collection

Mice were vaccinated intramuscularly with 50ul of the Polysaccharide Conjugate Vaccine Prevnar 13 (Wyeth Pharmaceuticals) into the caudal thigh muscle. 10ul of blood from each mouse was collected from the tail at weeks 0, 2 and 4 following vaccination and used for assessment of antibody titers. Sera from each mouse group were collected via cardiac puncture 4 weeks following vaccination, pooled and aliquoted to assess antibody quantity and functions in subsequent assays.

Bacteria

Streptococcus pneumoniae TIGR4 strain (serotype 4) was a kind gift from Andrew Camilli. Bacteria were grown to mid-exponential phase in Todd–Hewitt broth (BD Biosciences) supplemented with Oxyrase at 37°C/5% carbon dioxide, aliquoted in growth media supplemented with 20% glycerol and saved at -80°C as previously described (26, 67). Prior to use, bacteria were

thawed on ice, washed, and enumerated by serial dilution in PBS followed by dribbled plating on

Tryptic Soy Agar (TSA) plates supplemented with 5% sheep blood (Northeast Laboratory

Services).

Neutrophil depletion

Mice were injected intraperitoneally with 50μg of the Ly6G-depleting antibody IA8 or isotype IgG

control following the same schedule as shown in Fig. 2A and previously described (26).

Regulatory T cell depletion

B6 mice were treated intraperitoneally with 300μg of the CD25-depleting antibody PC-61.5.3 or

isotype IgG1 control following the schedule shown in Fig. 2B. DEREG mice were treated

intraperitoneally with 1μg of diphtheria toxin (Millipore) or DMSO as vehicle control following the

schedule shown in Fig. S18A. Depletion efficiency was assessed by flow cytometry (Fig. S16 &

S18A).

Murine organ/tissue processing and flow cytometry

Blood, spleens and/or vaccine draining lymph nodes (vLNs- inguinal and popliteal) were

harvested from euthanized mice. Single cell suspensions from spleens and vLNs were prepared

in RPMI 1640 media supplemented with 10% Fetal Bovine Serum. Following red blood cells lysis

with an in-house hypotonic solution (8.29g ammonium chloride, 1g sodium bicarbonate, 0.038g

trypsin/ethylenediaminetetraacetic acid (EDTA) and 1L of water), the remaining cells were

enumerated, incubated with Fc block (2.4G2), Live/Dead dye (Invitrogen) and stained with

additional antibodies in separate panels.

For PMNs, cells were stained with Ly6G (1A8), CD11b (M1/70), CXCR4 (2B11), CCR7

(4B12), CD80 (16-10A1), CD86 (GL1), CD11c (N418), MHC-II (M5/114.15.2), SPARC (124413),

PD-L1 (MIH5), CD39 (24DMS1) and/or CD73 (TY/11.8) antibodies followed by permeabilization

with Cytofix/Cytoperm (BD) and intracellular staining with ARG1 (A1exF5).

For B cells, cells were stained with CD45R/B220 (RA3-6B2), CD138 (281-2), CD95

(SA367H8), and CD38 (90) antibodies followed by permeabilization with Fix/Perm buffer from the

FOXP3/Transcription factor staining buffer set (eBioscience) and intracellular staining with AID

(Polyclonal, Bioss).

For T cells, cells were stained with CD4 (RM4-5), TCRβ (H57-597), CD11a (M17/4),

CD49d (R1-2), PD1 (J43) and/or CXCR5 (SPRCL5) antibodies followed by permeabilization with

Fix/Perm buffer from the FOXP3/Transcription factor staining buffer set (eBioscience) and

intracellular staining with FOXP3 (FJK-16s), RORγT (931-378), Tbet (eBio4B10) and GATA3

(L580-823).

For intracellular cytokine staining in PMNs, cells were incubated in Brefeldin A in RPMI

1640 media supplemented with 10% Fetal Bovine Serum and 1% Pen/Strep and stimulated with

$9x10^6$ CFU/well of heat-killed *S. pneumoniae* TIGR4 strain, 10µl/well of PCV, 100nM of phorbol-

12-myristate 13-acetate (PMA) or media alone for 3 hours. Cells were then washed and stained

extracellularly with Fc block, Live/Dead dye, Ly6G, and CD11b antibodies followed by intracellular

staining with IFNγ (XMG1.2) after permeabilization with Cytofix/Cytoperm (BD). For intracellular

cytokine staining in T cells, cells were incubated in Brefeldin A (BD) in RPMI 1640 media

supplemented with 10% Fetal Bovine Serum and 1% Pen/Strep and stimulated with a mix of 81nM

of PMA and 1.34µM of Ionomycin (Cell activation cocktail, Biolegend) or media alone for 2 hours.

Cells were then washed and stained extracellularly with Fc block, Live/Dead dye, CD4, TCRβ,

CD44 (IM7) and LAP (TW7-16B4) antibodies followed by intracellular staining with IFNγ

(XMG1.2), IL-17 (931-378), IL-4 (11B11) and IL-10 (JES5-16E3) antibodies after permeabilization

with Fix/Perm buffer from the FOXP3/Transcription factor staining buffer set (eBioscience).

Following staining, cells were washed with FACS buffer (HBSS, 1% FBS and 0.1% sodium azide)

and fluorescence was assessed using a BD Fortessa. At least 20,000 cells in the live, CD4$^+$TCRβ^+

and B220$^+$ cells gates respectively were collected and analyzed using the FlowJo software.

Antibody Enzyme-Linked Immunosorbent Assay

Antibodies in sera were assessed using an ELISA assay as previously described (26, 124).

Briefly, Nunc-maxisorp plates (Thermo Scientific) were coated at 4°C overnight with type 4

Pneumococcal Polysaccharide (ATCC®) at 2µg/well. The next day, plates were washed in PBS

supplemented with 0.25% Tween (PBS/T) and sera preabsorbed with a pneumococcal cell wall

polysaccharide mixture (CWP-multi, Cederlane) were added to the plates. After a 2-hour

incubation, plates were washed and incubated with diluted horseradish peroxidase (HRP)

conjugated anti-mouse IgM (Invitrogen), IgG (Millipore Sigma) or IgG1, IgG2b, IgG2c or IgG3

(Southern Biotech). One hour later, plates were washed, TMB substrate (Thermo Scientific) was

added, and absorbance was read at 650nm using a Biotek microplate reader for 10 minutes at 1-

minute intervals. Antibody units were calculated as a percentage of standard sera included in

each assay. Standard sera were obtained from mice that were intranasally inoculated with *S.*

pneumoniae serotype 4 weekly for 3 weeks and immunized with PCV at week 4.

IFNγ Enzyme-Linked Immunosorbent Assay

583   Single cell suspension of splenocytes were prepared in 2ml of RPMI 1640 media supplemented

584   with 10% Fetal Bovine Serum. IFNγ concentrations were quantified using the Mouse IFNγ ELISA

585   kit (Invitrogen, KMC4021) per manufacturer's instructions.

IFNγ blocking *in vivo*

588   Mice were treated intraperitoneally with 200µg of the IFNγ-blocking antibody XMG1.2 or isotype

589   control following same schedule as neutrophil depletion shown in Fig. 2A.

Cytokine/Chemokine Array

Pan cytokines were assessed using the Proteome Profiler Mouse XL Cytokine Array kit (R&D, ARY028) following the manufacturer's instructions as previously described (26). Briefly, single cell suspension of splenocytes were prepared in 2ml of RPMI 1640 media supplemented with 10% Fetal Bovine Serum. Cell suspensions were aliquoted and saved at -80°C. Samples were thawed on ice prior to use, lysed with Triton-X (VWR, 0694) and quantified for protein concentrations using the Pierce BCA protein assay kit (Thermofisher, A55864) per the manufacturer's instructions. All samples were brought to the same protein concentrations and used for pan-cytokine assessment. Protein membranes were read using the BioRad ChemiDoc MP Imaging System and the intensity for each dot on the blots was quantified using the Quick Spots Image Analysis Software. The intensity values of each blot were blanked using their respective buffer only controls. Intensity values across multiple blots were normalized using the blot with the highest standard. Fold change of the values from vaccinated PMN-depleted mice over those of vaccinated PMN-sufficient mice were calculated and log transformed.

Splenocyte culture assays

For mice, single cell suspension of splenocytes were prepared in 2ml of RPMI 1640 media supplemented with 10% Fetal Bovine Serum one day post vaccination. To remove PMNs from the obtained suspensions, splenocytes were stained with a PE labeled anti-Ly6G antibody and treated using the EasySep™ Release Mouse PE Positive Selection Kit (StemCell) following manufacturer's instruction. One million cells per conditions were used and treated with $5x10^7$ CFU heat-killed *S. pneumoniae* or 1µl/ml αCD3 (positive control). Cells were cocultured for two days before assessing percentages of Tregs.

Antibody binding to the surface of *S. pneumoniae*

Sera collected 4-weeks post vaccination were used to assess antibody binding to *S. pneumoniae*

(TIGR4) as previously described (26, 67). After 30 minutes of incubation at 37°C, cells were

washed with FACS buffer and labeled with an APC-tagged anti-mouse IgG (polyclonal) antibody

for 30 minutes on ice in the dark. Cells were then washed, resuspended in FACS buffer and

fluorescence was assessed using a BD Fortessa. At least 10,000 cells were analyzed using the

FlowJo software.

Opsonophagocytic Killing Assay (OPH)

Neutrophils were isolated from the bone marrow of mice using density centrifugation with

Histopaque 1119 and Histopaque 1077 (Sigma) as previously described (26, 124, 125). Briefly,

Neutrophils were resuspended in HBSS (without Ca^{2+} or Mg^{2+}) media supplemented with 0.1%

gelatin and used in the assay. Sera collected 4-weeks post vaccination were incubated with *S.*

pneumoniae TIGR4 for 5 minutes and added to neutrophils in HBSS (with Ca^{2+} and Mg^{2+}).

Following a 40-minute incubation at 37°C, reactions were plated on TSA plates with 5% sheep

blood. Percent killing was calculated relative to a no neutrophils control for each sera condition.

Adoptive Transfer of Sera

Four weeks following immunization, naïve, vaccinated neutrophil- and Treg-sufficient, neutrophil-

depleted, neutrophils- and Treg-depleted and Treg-depleted were euthanized. Blood was

harvested via cardiac puncture; sera were obtained from the blood and pooled for each mouse

group. A volume of 250µL of sera was transferred intraperitoneally into naive recipients which

were infected 1hour later.

Animal Infections

Mice were infected intratracheally with 3×10^5 CFU of *S. pneumoniae* serotype 4 (TIGR4) using

the tongue pull method as previously described (26, 126). Mice were monitored for survival for a

total of 7 days post infection and assessed daily for bacteremia by plating blood collected in 50mM

EDTA on blood agar plates for enumeration of bacterial CFU.

Human participants

Healthy unvaccinated female participants (24-29 years old) were recruited under NCT06128915.

Individuals who were pregnant, had acute infections within the last 2 weeks, took medication 48

hours prior, had history of chronic infections, had conditions that altered immunity (autoimmunity,

on immune modulating drugs, history of cancer) were excluded from the study. At the first visit

(week 0), donors had their blood drawn then were administered the Polysaccharide Conjugate

Vaccine Prevnar 20 (Pfizer, Wyeth Pharmaceuticals) intramuscularly as per manufacturer's

instructions. At week 1 and 4 following vaccination, donors had their blood drawn for subsequent

PMN phenotyping. Blood from week 1 was also used for coculture assays. All blood was collected

into tubes containing sodium citrate and visits occurred in the morning (8AM-10AM) to control for

circadian variations in PMN phenotype (127). All enrolled participants signed approved informed

consent forms.

Human PMN isolation and flow cytometry

For participants, PMNs were isolated from peripheral blood of week 0, 1, 4 post vaccinated

donors using EasySep™ Direct Human Neutrophil Isolation Kit (StemCell) respectively following

manufacturers' instructions. For each donor and visit $1x10^5$ PMNs were added to each well and

then stained with CD16 (eBioCB16), CD14 (61D3), CD10 (MEM-78), HLA-DR (L243), CD86

(BU63), PDL1 (MIH1), CD73 (AD2), CD39 (124413), PD-L1 (MIH5), CD39 (eBioA1) and/or CD73

(TY/11.8) antibodies followed by permeabilization with Cytofix/Cytoperm (BD) and intracellular

staining with ARG1 (A1exF5). Following staining, cells were washed with FACS buffer and

fluorescence was assessed using a BD Fortessa. At least 5000 cells in the CD16+ gate were

collected and analyzed using the FlowJo software.

Human PMN FLOWSOM analysis

Analysis and data visualization of the PMN phenotype with and without vaccination over time was

completed using a software OMIQ (Dotmatics). Traditional gating was done to differentiate the

neutrophil population and all the markers. Dimensional reduction was next performed with

optimized parameters for T-distributed stochastic neighbor embedding which improves

visualization of large datasets by reducing it to 2D/3D map while preserving local structure.

Clustering into neutrophil subsets was done using Flow-cytometry specific self-organizing map

(FLOWSOM) algorithm. Clustered heat maps were made based on meta-cluster gating to

characterize the different PMN subsets based on the expression of different markers.

Human PMN/PBMC coculture assays

For participants, PBMCs and PMNs were isolated from peripheral blood of 1-week

vaccinated donors using the One-Step Polymorph (Accurate Chemical) and EasySep™ Direct

Human Neutrophil Isolation Kit (StemCell) respectively following manufacturers' instructions. For

each donor, $2x10^5$ PBMCs were incubated in (RPMI with 10% FBS) with or without addition of

$5x10^4$ PMNs and stimulated with 10µl of PCV. Cells were cocultured for two days before

assessing percentages and proliferation of T cells. At 48 hours post incubation, T cells were

stained with CD11a (HI111), PD1(EH12.1), CD4(S3.5), CD49d (9F10), CXCR5 (MU5UBEE),

CD25 (BC96) followed by permeabilization with Fix/Perm buffer from the FOXP3/Transcription

factor staining buffer set (eBioscience) and intracellular staining with FOXP3 (PCH101), RORγT

(AFJKS-9), Tbet (eB4B10), GATA3 (L580-823) and ki67 (B56). Following staining, cells were

washed with FACS buffer and data collected using a BD Fortessa. At least 10,000 cells in the live

CD4$^+$ gate were collected and analyzed using the FlowJo software.

Statistical Analysis

All statistical analysis was done using GraphPad Prism version 10 software. Data were checked

for normality distribution using the Shapiro-Wilk test. Significant differences were determined by

One-way analysis of variance followed Šídák's, Holm-Šídák's, Tukey's or Dunnett's multiple

comparisons test; Kruskal-Wallis One-way analysis of variance followed Dunn's multiple

comparisons; Mann-Whitney test or Student's t test; or one sample t-test as appropriate. Survival

was analyzed using the log-rank (Mantel-Cox) test. All p values <0.05 were considered significant.

Study Approvals

All work with mice was performed in accordance with the recommendations in the Guide for the

Care and Use of Laboratory Animals published by the National Institutes of Health. All procedures

were reviewed and approved by the University at Buffalo Institutional Animal Care and Use

Committee (IACUC), approval number MIC33018Y. All work with human donors was approved

by the University at Buffalo Human investigation Review Board (IRB), approval number

STUDY00007111.

Data Availability

All data presented within this manuscript are found in the supportive data values file and available

from the corresponding author upon request.

Author Contributions

EYIT conducted research, acquired data, analyzed data, and wrote the manuscript. AB

conducted research, acquired data, and analyzed data. AYK, EL, MB, MC conducted research

and acquired data. EAW provided materials. ENBG designed the research studies, wrote, and

edited the manuscript, and had responsibility for final content. All authors read and approved the

manuscript.

Funding Support

This work was supported by National Institute of Health grant R01 AG068568-01A1 to ENBG,

Mark Diamond Research Funds to EYIT, and an AHA predoctoral fellowship to AB. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health.

References

1.   Simmons SR, Bhalla M, Herring SE, Tchalla EYI, and Bou Ghanem EN. Older but

not wiser: The age-driven changes in neutrophil responses during pulmonary

infections. *Infect Immun.* 2021.

2.   Mantovani A, Cassatella MA, Costantini C, and Jaillon S. Neutrophils in the

activation and regulation of innate and adaptive immunity. *Nature reviews*

Immunology. 2011;11(8):519-31.

3.   Bert S, Nadkarni S, and Perretti M. Neutrophil-T cell crosstalk and the control of

the host inflammatory response. *Immunol Rev.* 2023;314(1):36-49.

4.   Li Y, Wang W, Yang F, Xu Y, Feng C, and Zhao Y. The regulatory roles of

neutrophils in adaptive immunity. *Cell Communication and Signaling.* 2019;17(1).

5.   Leliefeld PHC, Koenderman L, and Pillay J. How Neutrophils Shape Adaptive

Immune Responses. *Frontiers in Immunology.* 2015;6.

6.   Shafqat A, Khan JA, Alkachem AY, Sabur H, Alkattan K, Yaqinuddin A, et al. How

Neutrophils Shape the Immune Response: Reassessing Their Multifaceted Role

in Health and Disease. *Int J Mol Sci.* 2023;24(24).

7.   Abi Abdallah DS, Egan CE, Butcher BA, and Denkers EY. Mouse neutrophils are

professional antigen-presenting cells programmed to instruct Th1 and Th17 T-cell

differentiation. *International immunology.* 2011;23(5):317-26.

8.   Culshaw S, Millington OR, Brewer JM, and McInnes IB. Murine neutrophils present

Class II restricted antigen. *Immunol Lett.* 2008;118(1):49-54.

9.   Vono M, Lin A, Norrby-Teglund A, Koup RA, Liang F, and Lore K. Neutrophils

acquire the capacity for antigen presentation to memory CD4(+) T cells in vitro and

ex vivo. *Blood.* 2017;129(14):1991-2001.

10.   Matsushima H, Geng S, Lu R, Okamoto T, Yao Y, Mayuzumi N, et al. Neutrophil

771   differentiation into a unique hybrid population exhibiting dual phenotype and

772   functionality of neutrophils and dendritic cells. *Blood.* 2013;121(10):1677-89.

773   11.   de Kleijn S, Langereis JD, Leentjens J, Kox M, Netea MG, Koenderman L, et al.

774   IFN-γ-stimulated neutrophils suppress lymphocyte proliferation through

775   expression of PD-L1. *PLoS One.* 2013;8(8):e72249.

776   12.   Kasten KR, Muenzer JT, and Caldwell CC. Neutrophils are significant producers

777   of IL-10 during sepsis. *Biochemical and Biophysical Research Communications.*

778   2010;393(1):28-31.

779   13.   Pillay J, Kamp VM, Van Hoffen E, Visser T, Tak T, Lammers J-W, et al. A subset

780   of neutrophils in human systemic inflammation inhibits T cell responses through

781   Mac-1. *Journal of Clinical Investigation.* 2012;122(1):327-36.

782   14.   Schmielau J, and Finn OJ. Activated granulocytes and granulocyte-derived

783   hydrogen peroxide are the underlying mechanism of suppression of t-cell function

784   in advanced cancer patients. *Cancer research.* 2001;61(12):4756-60.

785   15.   Sangaletti S, Talarico G, Chiodoni C, Cappetti B, Botti L, Portararo P, et al. SPARC

786   Is a New Myeloid-Derived Suppressor Cell Marker Licensing Suppressive

787   Activities. *Front Immunol.* 2019;10:1369.

788   16.   Jiang X, Wu X, Xiao Y, Wang P, Zheng J, Wu X, et al. The ectonucleotidases CD39

789   and CD73 on T cells: The new pillar of hematological malignancy. *Front Immunol.*

790   2023;14:1110325.

791   17.   Allard B, Longhi MS, Robson SC, and Stagg J. The ectonucleotidases CD39 and

792   CD73: Novel checkpoint inhibitor targets. *Immunol Rev.* 2017;276(1):121-44.

18.   Jee J, Bonnegarde-Bernard A, Duverger A, Iwakura Y, Cormet-Boyaka E, Martin

TL, et al. Neutrophils negatively regulate induction of mucosal IgA responses after

sublingual immunization. *Mucosal Immunol.* 2015;8(4):735-45.

19.   Kamenyeva O, Boularan C, Kabat J, Cheung GYC, Cicala C, Yeh AJ, et al.

Neutrophil Recruitment to Lymph Nodes Limits Local Humoral Response to

Staphylococcus aureus. *PLoS pathogens.* 2015;11(4):e1004827.

20.   Bird AK, Chang M, Barnard J, Goldman BI, Meednu N, Rangel-Moreno J, et al.

Neutrophils Slow Disease Progression in Murine Lupus via Modulation of

Autoreactive Germinal Centers. *The Journal of Immunology.* 2017;199(2):458-66.

21.   Puga I, Cols M, Barra CM, He B, Cassis L, Gentile M, et al. B cell-helper

neutrophils stimulate the diversification and production of immunoglobulin in the

marginal zone of the spleen. *Nature immunology.* 2011;13(2):170-80.

22.   Chorny A, Casas-Recasens S, Sintes J, Shan M, Polentarutti N, Garcia-Escudero

R, et al. The soluble pattern recognition receptor PTX3 links humoral innate and

adaptive immune responses by helping marginal zone B cells. *The Journal of*

experimental medicine. 2016;213(10):2167-85.

23.   Musich T, Rahman MA, Mohanram V, Miller-Novak L, Demberg T, Venzon DJ, et

al. Neutrophil Vaccination Dynamics and Their Capacity To Mediate B Cell Help in

Rhesus Macaques. *Journal of immunology.* 2018;201(8):2287-302.

24.   Bhattacharya P, Dey R, Saxena A, Karmakar S, Ismail N, Gannavaram S, et al.

Essential Role of Neutrophils in the Protective Immune Response Induced by a

Live Attenuated Leishmania Vaccine. *Journal of immunology.* 2020;205(12):3333-

47.

25. Trentini MM, de Oliveira FM, Kipnis A, and Junqueira-Kipnis AP. The Role of
Neutrophils in the Induction of Specific Th1 and Th17 during Vaccination against
Tuberculosis. *Front Microbiol.* 2016;7:898.

26. Tchalla EYI, Bhalla M, Wohlfert EA, and Bou Ghanem EN. Neutrophils Are
Required During Immunization With the Pneumococcal Conjugate Vaccine for
Protective Antibody Responses and Host Defense Against Infection. *J Infect Dis.*
2020;222(8):1363-70.

27. CDC. Center for Disease Control and Prevention. Active Bacterial Core
Surveillance Report, Emerging Infections Program Network, Streptococcus
pneumoniae. 2022.

28. Vinuesa CG, Linterman MA, Yu D, and MacLennan IC. Follicular Helper T Cells.
Annual review of immunology. 2016;34:335-68.

29. Rappuoli R. Glycoconjugate vaccines: Principles and mechanisms. *Sci Transl*
Med. 2018;10(456).

30. Sterrett S, Peng BJ, Burton RL, LaFon DC, Westfall AO, Singh S, et al. Peripheral
CD4 T follicular cells induced by a conjugated pneumococcal vaccine correlate
with enhanced opsonophagocytic antibody responses in younger individuals.
Vaccine. 2020.

31. De Silva NS, and Klein U. Dynamics of B cells in germinal centres. *Nature reviews*
Immunology. 2015;15(3):137-48.

32. Pollard AJ, Perrett KP, and Beverley PC. Maintaining protection against invasive
bacteria with protein–polysaccharide conjugate vaccines. *Nature Reviews*
Immunology. 2009;9(3):213-20.

33.   Rappuoli R, De Gregorio E, and Costantino P. On the mechanisms of conjugate

840   vaccines. *Proceedings of the National Academy of Sciences.* 2019;116(1):14-6.

841   34.   Murphy KM, Weaver C, Mowat A, Berg L, Chaplin D, Janeway CA, et al. *Janeway's*

842   *immunobiology.* 2017.

843   35.   Rappuoli R, De Gregorio E, and Costantino P. On the mechanisms of conjugate

844   vaccines. *Proc Natl Acad Sci U S A.* 2019;116(1):14-6.

845   36.   Vidarsson G, Dekkers G, and Rispens T. IgG Subclasses and Allotypes: From

846   Structure to Effector Functions. *Frontiers in Immunology.* 2014;5(520).

847   37.   Forthal DN. Functions of Antibodies. *Microbiol Spectr.* 2014;2(4):1-17.

848   38.   Dong C. Cytokine Regulation and Function in T Cells. *Annual review of*

849   *immunology.* 2021;39(1):51-76.

850   39.   Vazquez MI, Catalan-Dibene J, and Zlotnik A. B cells responses and cytokine

851   production are regulated by their immune microenvironment. *Cytokine.*

852   2015;74(2):318-26.

853   40.   Shevyrev D, and Tereshchenko V. Treg Heterogeneity, Function, and

854   Homeostasis. *Front Immunol.* 2019;10:3100.

855   41.   Ndure J, and Flanagan KL. Targeting regulatory T cells to improve vaccine

856   immunogenicity in early life. *Front Microbiol.* 2014;5:477.

857   42.   Bayry J. Regulatory T cells as adjuvant target for enhancing the viral disease

858   vaccine efficacy. *Virusdisease.* 2014;25(1):18-25.

859   43.   Moore AC, Gallimore A, Draper SJ, Watkins KR, Gilbert SC, and Hill AV. Anti-

860   CD25 antibody enhancement of vaccine-induced immunogenicity: increased

durable cellular immunity with reduced immunodominance. *Journal of*

immunology. 2005;175(11):7264-73.

44. Pere H, Tanchot C, Bayry J, Terme M, Taieb J, Badoual C, et al. Comprehensive

analysis of current approaches to inhibit regulatory T cells in cancer.

Oncoimmunology. 2012;1(3):326-33.

45. Weingartner E, and Golding A. Direct control of B cells by Tregs: An opportunity

for long-term modulation of the humoral response. *Cell Immunol.* 2017;318:8-16.

46. Stepkowski S, Bekbolsynov D, Oenick J, Brar S, Mierzejewska B, Rees MA, et al.

The Major Role of T Regulatory Cells in the Efficiency of Vaccination in General

and Immunocompromised Populations: A Review. *Vaccines (Basel).* 2024;12(9).

47. Togashi Y, Shitara K, and Nishikawa H. Regulatory T cells in cancer

immunosuppression - implications for anticancer therapy. *Nat Rev Clin Oncol.*

2019;16(6):356-71.

48. Brezar V, Ruffin N, Richert L, Surenaud M, Lacabaratz C, Palucka K, et al.

Decreased HIV-specific T-regulatory responses are associated with effective DC-

vaccine induced immunity. *PLoS pathogens.* 2015;11(3):e1004752.

49. Boudewijns M, Jeurissen A, Wuyts M, Moens L, Boon L, Van Neerven JJ, et al.

Blockade of CTLA-4 (CD152) enhances the murine antibody response to

pneumococcal capsular polysaccharides. *J Leukoc Biol.* 2005;78(5):1060-9.

50. He SWJ, van de Garde MDB, Pieren DKJ, Poelen MCM, Voss F, Abdullah MR, et

al. Diminished Pneumococcal-Specific CD4(+) T-Cell Response is Associated

With Increased Regulatory T Cells at Older Age. *Front Aging.* 2021;2:746295.

51.   Maceiras AR, Fonseca VR, Agua-Doce A, and Graca L. T follicular regulatory cells

in mice and men. *Immunology.* 2017;152(1):25-35.

52.   Lu Y, and Craft J. T Follicular Regulatory Cells: Choreographers of Productive

Germinal Center Responses. *Front Immunol.* 2021;12:679909.

53.   Qi J, Liu C, Bai Z, Li X, and Yao G. T follicular helper cells and T follicular regulatory

cells in autoimmune diseases. *Front Immunol.* 2023;14:1178792.

54.   Sage PT, Ron-Harel N, Juneja VR, Sen DR, Maleri S, Sungnak W, et al.

Suppression by T(FR) cells leads to durable and selective inhibition of B cell

effector function. *Nature immunology.* 2016;17(12):1436-46.

55.   Xie MM, Chen Q, Liu H, Yang K, Koh B, Wu H, et al. T follicular regulatory cells

and IL-10 promote food antigen-specific IgE. *J Clin Invest.* 2020;130(7):3820-32.

56.   Jacobsen JT, Hu W, TB RC, Solem S, Galante A, Lin Z, et al. Expression of Foxp3

by T follicular helper cells in end-stage germinal centers. *Science.* 2021;373(6552).

57.   Himmel ME, Crome SQ, Ivison S, Piccirillo C, Steiner TS, and Levings MK. Human

CD4+ FOXP3+ regulatory T cells produce CXCL8 and recruit neutrophils.

European journal of immunology. 2011;41(2):306-12.

58.   Richards H, Williams A, Jones E, Hindley J, Godkin A, Simon AK, et al. Novel role

of regulatory T cells in limiting early neutrophil responses in skin. *Immunology.*

2010;131(4):583-92.

59.   Lewkowicz N, Klink M, Mycko MP, and Lewkowicz P. Neutrophil--CD4+CD25+ T

regulatory cell interactions: a possible new mechanism of infectious tolerance.

Immunobiology. 2013;218(4):455-64.

60.   Mishalian I, Bayuh R, Eruslanov E, Michaeli J, Levy L, Zolotarov L, et al.

906   Neutrophils recruit regulatory T-cells into tumors via secretion of CCL17--a new

907   mechanism of impaired antitumor immunity. *Int J Cancer.* 2014;135(5):1178-86.

908   61.   Perobelli SM, Mercadante AC, Galvani RG, Goncalves-Silva T, Alves AP, Pereira-

909   Neves A, et al. G-CSF-Induced Suppressor IL-10+ Neutrophils Promote

910   Regulatory T Cells That Inhibit Graft-Versus-Host Disease in a Long-Lasting and

911   Specific Way. *Journal of immunology.* 2016;197(9):3725-34.

912   62.   Nadkarni S, Smith J, Sferruzzi-Perri AN, Ledwozyw A, Kishore M, Haas R, et al.

913   Neutrophils induce proangiogenic T cells with a regulatory phenotype in

914   pregnancy. *Proc Natl Acad Sci U S A.* 2016;113(52):E8415-E24.

915   63.   Gao Y, Min K, Zhang Y, Su J, Greenwood M, and Gronert K. Female-Specific

916   Downregulation of Tissue Polymorphonuclear Neutrophils Drives Impaired

917   Regulatory T Cell and Amplified Effector T Cell Responses in Autoimmune Dry

918   Eye Disease. *Journal of immunology.* 2015;195(7):3086-99.

919   64.   Wagenvoort GH, Sanders EA, Vlaminckx BJ, de Melker HE, van der Ende A, and

920   Knol MJ. Sex differences in invasive pneumococcal disease and the impact of

921   pneumococcal conjugate vaccination in the Netherlands, 2004 to 2015. *Euro*

922   *Surveill.* 2017;22(10).

923   65.   Wiese AD, Grijalva CG, Zhu Y, Mitchel EF, Jr., and Griffin MR. Changes in

924   Childhood Pneumonia Hospitalizations by Race and Sex Associated with

925   Pneumococcal Conjugate Vaccines. *Emerg Infect Dis.* 2016;22(6).

926 66. Soneji S, and Metlay J. Mortality reductions for older adults differ by race/ethnicity

927   and gender since the introduction of adult and pediatric pneumococcal vaccines.

928   *Public Health Rep.* 2011;126(2):259-69.

929 67. Tchalla EYI, Betadpur A, Khalil AY, Bhalla M, and Bou Ghanem EN. Sex-based

930   difference in immune responses and efficacy of the pneumococcal conjugate

931   vaccine. *J Leukoc Biol.* 2024.

932 68. Beauvillain C, Cunin P, Doni A, Scotet M, Jaillon S, Loiry ML, et al. CCR7 is

933   involved in the migration of neutrophils to lymph nodes. *Blood.* 2011;117(4):1196-

934   204.

935 69. Hampton HR, Bailey J, Tomura M, Brink R, and Chtanova T. Microbe-dependent

936   lymphatic migration of neutrophils modulates lymphocyte proliferation in lymph

937   nodes. *Nat Commun.* 2015;6:7139.

938 70. Stavnezer J, Guikema JE, and Schrader CE. Mechanism and regulation of class

939   switch recombination. *Annual review of immunology.* 2008;26:261-92.

940 71. Di Noia JM, and Neuberger MS. Molecular mechanisms of antibody somatic

941   hypermutation. *Annu Rev Biochem.* 2007;76:1-22.

942 72. Morel C, Badell E, Abadie V, Robledo M, Setterblad N, Gluckman JC, et al.

943   Mycobacterium bovis BCG-infected neutrophils and dendritic cells cooperate to

944   induce specific T cell responses in humans and mice. *European journal of*

945   *immunology.* 2008;38(2):437-47.

946 73. Abbas AK, Burstein HJ, and Bogen SA. Determinants of helper T cell-dependent

947   antibody production. *Semin Immunol.* 1993;5(6):441-7.

948   74.   Karachunski PI, Ostlie NS, Monfardini C, and Conti-Fine BM. Absence of IFN-γ or

949   IL-12 Has Different Effects on Experimental Myasthenia Gravis in C57BL/6 Mice.

950   *The Journal of Immunology.* 2000;164(10):5236-44.

951   75.   Ballesteros I, Rubio-Ponce A, Genua M, Lusito E, Kwok I, Fernandez-Calvo G, et

952   al. Co-option of Neutrophil Fates by Tissue Environments. *Cell.* 2020;183(5):1282-

953   97 e18.

954   76.   Yang P, Li Y, Xie Y, and Liu Y. Different Faces for Different Places: Heterogeneity

955   of Neutrophil Phenotype and Function. *J Immunol Res.* 2019;2019:8016254.

956   77.   Rambault M, Doz-Deblauwe E, Le Vern Y, Carreras F, Cunha P, Germon P, et al.

957   Neutrophils Encompass a Regulatory Subset Suppressing T Cells in Apparently

958   Healthy Cattle and Mice. *Front Immunol.* 2021;12:625244.

959   78.   Christiaansen AF, Dixit UG, Coler RN, Marie Beckmann A, Reed SG, Winokur PL,

960   et al. CD11a and CD49d enhance the detection of antigen-specific T cells following

961   human vaccination. *Vaccine.* 2017;35(33):4255-61.

962   79.   Liu S, Ren J, and Ten Dijke P. Targeting TGFβ signal transduction for cancer

963   therapy. *Signal Transduction and Targeted Therapy.* 2021;6(1).

964   80.   Areström I, Zuber B, Bengtsson T, and Ahlborg N. Measurement of human latent

965   Transforming Growth Factor-β1 using a latency associated protein-reactive ELISA.

966   *J Immunol Methods.* 2012;379(1-2):23-9.

967   81.   Farkas AM, Marvel DM, and Finn OJ. Antigen choice determines vaccine-induced

968   generation of immunogenic versus tolerogenic dendritic cells that are marked by

969   differential expression of pancreatic enzymes. *Journal of immunology.*

970   2013;190(7):3319-27.

971   82.   Jin RM, Blair SJ, Warunek J, Heffner RR, Blader IJ, and Wohlfert EA. Regulatory

972   T Cells Promote Myositis and Muscle Damage in Toxoplasma gondii Infection.

973   *Journal of immunology.* 2017;198(1):352-62.

974   83.   Lahl K, Loddenkemper C, Drouin C, Freyer J, Arnason J, Eberl G, et al. Selective

975   depletion of Foxp3+ regulatory T cells induces a scurfy-like disease. *The Journal*

976   *of experimental medicine.* 2007;204(1):57-63.

977   84.   Nauseef WM. Human neutrophils ≠ murine neutrophils: Does it matter? *Immunol*

978   *Rev.* 2023;314(1):442-56.

979   85.   Fontenot JD, Gavin MA, and Rudensky AY. Foxp3 programs the development and

980   function of CD4+CD25+ regulatory T cells. *Nature immunology.* 2003;4(4):330-6.

981   86.   Ono M. Control of regulatory T-cell differentiation and function by T-cell receptor

982   signalling and Foxp3 transcription factor complexes. *Immunology.* 2020;160(1):24-

983   37.

984   87.   Bank U, Reinhold D, Schneemilch C, Kunz D, Synowitz HJ, and Ansorge S.

985   Selective proteolytic cleavage of IL-2 receptor and IL-6 receptor ligand binding

986   chains by neutrophil-derived serine proteases at foci of inflammation. *Journal of*

987   *interferon & cytokine research : the official journal of the International Society for*

988   *Interferon and Cytokine Research.* 1999;19(11):1277-87.

989   88.   Harris F, Berdugo YA, and Tree T. IL-2-based approaches to Treg enhancement.

990   *Clinical and experimental immunology.* 2023;211(2):149-63.

991   89.   Ariel A, Yavin EJ, Hershkoviz R, Avron A, Franitza S, Hardan I, et al. IL-2 induces

992   T cell adherence to extracellular matrix: inhibition of adherence and migration by

993   IL-2 peptides generated by leukocyte elastase. *Journal of immunology.*
994   1998;161(5):2465-72.

995 90. Fu Z, Akula S, Thorpe M, and Hellman L. Potent and Broad but not Unselective

996   Cleavage of Cytokines and Chemokines by Human Neutrophil Elastase and

997   Proteinase 3. *Int J Mol Sci.* 2020;21(2).

998 91. Chen W. TGF-beta Regulation of T Cells. *Annual review of immunology.*

999   2023;41:483-512.

1000 92. Kretschmer K, Apostolou I, Hawiger D, Khazaie K, Nussenzweig MC, and von

1001   Boehmer H. Inducing and expanding regulatory T cell populations by foreign

1002   antigen. *Nature immunology.* 2005;6(12):1219-27.

1003 93. Corse E, Gottschalk RA, and Allison JP. Strength of TCR-peptide/MHC

1004   interactions and in vivo T cell responses. *Journal of immunology.*

1005   2011;186(9):5039-45.

1006 94. Ohkura N, Hamaguchi M, and Sakaguchi S. FOXP3+ regulatory T cells: control of

1007   FOXP3 expression by pharmacological agents. *Trends Pharmacol Sci.*

1008   2011;32(3):158-66.

1009 95. Stream A, Corriden R, Dohrmann S, Gallo RL, Nizet V, and Anderson EL. The

1010   Effect of Retinoic Acid on Neutrophil Innate Immune Interactions With Cutaneous

1011   Bacterial Pathogens. *Infect Microbes Dis.* 2024;6(2):65-73.

1012 96. Mader JS, Ewen C, Hancock RE, and Bleackley RC. The human cathelicidin, LL-

1013   37, induces granzyme-mediated apoptosis in regulatory T cells. *J Immunother.*

1014   2011;34(3):229-35.

97. Glemain A, Neel M, Neel A, Andre-Gregoire G, Gavard J, Martinet B, et al.

Neutrophil-derived extracellular vesicles induce endothelial inflammation and

damage through the transfer of miRNAs. *J Autoimmun.* 2022;129:102826.

98. Gomez I, Ward B, Souilhol C, Recarti C, Ariaans M, Johnston J, et al. Neutrophil

microvesicles drive atherosclerosis by delivering miR-155 to atheroprone

endothelium. *Nat Commun.* 2020;11(1):214.

99. Jiao Y, Zhang T, Zhang C, Ji H, Tong X, Xia R, et al. Exosomal miR-30d-5p of

neutrophils induces M1 macrophage polarization and primes macrophage

pyroptosis in sepsis-related acute lung injury. *Crit Care.* 2021;25(1):356.

100. Lu J, Wang W, Li P, Wang X, Gao C, Zhang B, et al. MiR-146a regulates regulatory

T cells to suppress heart transplant rejection in mice. *Cell Death Discov.*

2021;7(1):165.

101. Kunze-Schumacher H, and Krueger A. The Role of MicroRNAs in Development

and Function of Regulatory T Cells - Lessons for a Better Understanding of

MicroRNA Biology. *Front Immunol.* 2020;11:2185.

102. Arroyo AB, Fernandez-Perez MP, Del Monte A, Aguila S, Mendez R, Hernandez-

Antolin R, et al. miR-146a is a pivotal regulator of neutrophil extracellular trap

formation promoting thrombosis. *Haematologica.* 2021;106(6):1636-46.

103. Garley M, Nowak K, and Jablonska E. Neutrophil microRNAs. *Biol Rev Camb*

Philos Soc. 2024;99(3):864-77.

104. Toka FN, Suvas S, and Rouse BT. CD4+ CD25+ T cells regulate vaccine-

generated primary and memory CD8+ T-cell responses against herpes simplex

virus type 1. *J Virol.* 2004;78(23):13082-9.

105. Levy Y, Thiebaut R, Montes M, Lacabaratz C, Sloan L, King B, et al. Dendritic cell-

based therapeutic vaccine elicits polyfunctional HIV-specific T-cell immunity

associated with control of viral load. *European journal of immunology.*

2014;44(9):2802-10.

106. Wang P, and Zheng SG. Regulatory T cells and B cells: implication on autoimmune

diseases. *Int J Clin Exp Pathol.* 2013;6(12):2668-74.

107. Gonzalez-Figueroa P, Roco JA, Papa I, Nunez Villacis L, Stanley M, Linterman

MA, et al. Follicular regulatory T cells produce neuritin to regulate B cells. *Cell.*

2021;184(7):1775-89 e19.

108. Tan D, Yin W, Guan F, Zeng W, Lee P, Candotti F, et al. B cell-T cell interplay in

immune regulation: A focus on follicular regulatory T and regulatory B cell

functions. *Front Cell Dev Biol.* 2022;10:991840.

109. Lim HW, Hillsamer P, Banham AH, and Kim CH. Cutting edge: direct suppression

of B cells by CD4+ CD25+ regulatory T cells. *Journal of immunology.*

2005;175(7):4180-3.

110. Park SR. Activation-induced Cytidine Deaminase in B Cell Immunity and Cancers.

Immune Netw. 2012;12(6):230-9.

111. McBride KM, Gazumyan A, Woo EM, Schwickert TA, Chait BT, and Nussenzweig

MC. Regulation of class switch recombination and somatic mutation by AID

phosphorylation. *The Journal of experimental medicine.* 2008;205(11):2585-94.

112. Chaudhuri J, Khuong C, and Alt FW. Replication protein A interacts with AID to

promote deamination of somatic hypermutation targets. *Nature.*

2004;430(7003):992-8.

113.   Basu U, Chaudhuri J, Alpert C, Dutt S, Ranganath S, Li G, et al. The AID antibody

diversification enzyme is regulated by protein kinase A phosphorylation. *Nature.*

2005;438(7067):508-11.

114.   Demorest ZL, Li M, and Harris RS. Phosphorylation directly regulates the intrinsic

DNA cytidine deaminase activity of activation-induced deaminase and

APOBEC3G protein. *J Biol Chem.* 2011;286(30):26568-75.

115.   Gazumyan A, Timachova K, Yuen G, Siden E, Di Virgilio M, Woo EM, et al. Amino-

terminal phosphorylation of activation-induced cytidine deaminase suppresses c-

myc/IgH translocation. *Mol Cell Biol.* 2011;31(3):442-9.

116.   Klein M, and Bopp T. Cyclic AMP Represents a Crucial Component of Treg Cell-

Mediated Immune Regulation. *Front Immunol.* 2016;7:315.

117.   Wiemken TL, Carrico RM, Klein SL, Jonsson CB, Peyrani P, Kelley RR, et al. The

effectiveness of the polysaccharide pneumococcal vaccine for the prevention of

hospitalizations due to Streptococcus pneumoniae community-acquired

pneumonia in the elderly differs between the sexes: results from the Community-

Acquired Pneumonia Organization (CAPO) international cohort study. *Vaccine.*

2014;32(19):2198-203.

118.   Robinson GA, Peng J, Peckham H, Butler G, Pineda-Torra I, Ciurtin C, et al.

Investigating sex differences in T regulatory cells from cisgender and transgender

healthy individuals and patients with autoimmune inflammatory disease: a cross-

sectional study. *Lancet Rheumatol.* 2022;4(10):e710-e24.

119. Afshan G, Afzal N, and Qureshi S. CD4+CD25(hi) regulatory T cells in healthy

males and females mediate gender difference in the prevalence of autoimmune

diseases. *Clin Lab.* 2012;58(5-6):567-71.

120. Raynor J, Lages CS, Shehata H, Hildeman DA, and Chougnet CA. Homeostasis

and function of regulatory T cells in aging. *Curr Opin Immunol.* 2012;24(4):482-7.

121. Flanagan KL, Fink AL, Plebanski M, and Klein SL. Sex and Gender Differences in

the Outcomes of Vaccination over the Life Course. *Annu Rev Cell Dev Biol.*

2017;33:577-99.

122. Pollard AJ, and Bijker EM. A guide to vaccinology: from basic principles to new

developments. *Nature Reviews Immunology.* 2021;21(2):83-100.

123. de St Maurice A, Schaffner W, Griffin MR, Halasa N, and Grijalva CG. Persistent

Sex Disparities in Invasive Pneumococcal Diseases in the Conjugate Vaccine Era.

J Infect Dis. 2016;214(5):792-7.

124. Bhalla M, Nayerhoda R, Tchalla EYI, Abamonte A, Park D, Simmons SR, et al.

Liposomal Encapsulation of Polysaccharides (LEPS) as an Effective Vaccine

Strategy to Protect Aged Hosts Against S. pneumoniae Infection. *Frontiers in*

Aging. 2021;2.

125. Bou Ghanem EN, Clark S, Roggensack SE, McIver SR, Alcaide P, Haydon PG, et

al. Extracellular Adenosine Protects against Streptococcus pneumoniae Lung

Infection by Regulating Pulmonary Neutrophil Recruitment. *PLoS Pathog.*

2015;11(8):e1005126.

126. Lenhard A, Joma BH, Siwapornchai N, Hakansson AP, Leong JM, and Bou

Ghanem EN. A Mouse Model for the Transition of Streptococcus pneumoniae from

Colonizer to Pathogen upon Viral Co-Infection Recapitulates Age-Exacerbated

Illness. *J Vis Exp.* 2022(187).

127.   Casanova-Acebes M, Pitaval C, Weiss LA, Nombela-Arrieta C, Chèvre R, N AG,

et al. Rhythmic modulation of the hematopoietic niche through neutrophil

clearance. *Cell.* 2013;153(5):1025-35.

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130 **Figures and legends**

A. Migration markers expression on circulating PMNs

B. PMN influx

1131

Figure 1: PMN migration to secondary lymphoid organs following vaccination. Adult

C57BL/6 female mice were injected with PCV or PBS. (A) Blood was collected from the mice 24

hours following vaccination and assessed using flow cytometry. Percentages (left) and gMFI

(right) of CCR7 and CXCR4 on PMNs (Ly6G$^+$CD11b$^+$) were determined. (B) At different

timepoints post vaccination, blood, spleen and vLNs were collected from all mice and PMN

(Ly6G$^+$CD11b$^+$) numbers was assessed using flow cytometry. (A) Data are pooled from 3

separate experiments and each dot represents an individual mouse with a total of n=5 naïve and

n=6 vaccinated mice. Graphs represent the mean +/- SD. (B) Data are pooled from 3 separate

experiments with a total of 6 mice per timepoint. Line graphs represent the mean +/- 95%

confidence interval. Abbreviations: gMFI = geometric Mean Fluorescence Intensity.

1142

A. PMN depletion

1143

1144   **Figure 2: PMN and Treg depletion models.** Adult C57BL/6 female mice were injected with PCV

1145   or PBS and depleted of their PMNs and/or Tregs or treated with isotype controls. (A) Depletion

1146   timeline of PMNs using an αLy6G antibody. (B) Depletion timeline of Tregs using an αCD25

1147   antibody.

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

T cell engaging markers

1159

Figure 3: PMN expression of T cell engaging markers following PCV vaccination. Adult

C57BL/6 female mice were injected with PCV or PBS. At 18- or 24 hours post vaccination,

blood, spleen and vLNs were collected and assessed using flow cytometry. Percentages (left)

and gMFI (right) of (A) CD80, (B) CD86, (C) MHC-II, and (D) CD11c on PMNs (Ly6G$^+$CD11b$^+$)

were determined. Data are pooled from 3 separate experiments and each dot represents an

individual mouse with a total of n=5 naïve and n=6 vaccinated mice. * denotes significant

differences between the indicated groups as determined by One-Way ANOVA followed by Šídák's

multiple comparisons test. * is $p \leq 0.05$, ** is $p \leq 0.01$, *** is $p \leq 0.001$, and **** is $p \leq 0.0001$. Graphs

represent the mean +/- SD. Abbreviations: gMFI = geometric Mean Fluorescence Intensity; MHC

= Major Histocompatibility Complex.

1170

1171

1172

1173

T cell suppressing markers

1174

Figure 4: PMN expression of T cell suppressing markers following PCV vaccination. Adult

C57BL/6 female mice were injected with PCV or PBS. At 18 hours post vaccination, blood, spleen

and vLNs were collected and assessed using flow cytometry. Percentages (left) and gMFI (right)

of (A) ARG1, (B) CD39, (C) CD73, and (D) PD-L1 on PMNs (Ly6G$^+$CD11b$^+$) were determined.

Data are pooled from 3 separate experiments and each dot represents an individual mouse with

a total of n=5 naïve and n=6 vaccinated mice. * denotes significant differences between the

indicated groups as determined by One-Way ANOVA followed by Šídák's multiple comparisons

test and # denotes significant differences between the indicated groups as determined by

Student's t test. * and # is $p{\leq}0.05$, ** is $p{\leq}0.01$, *** is $p{\leq}0.001$, and **** is $p{\leq}0.0001$. Graphs

represent the mean +/- SD. Abbreviations: gMFI = geometric Mean Fluorescence Intensity; ARG=

arginase.

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

PMN phenotype

A.

Blood　　　　　　　LNs　　　　　　　Spleen

● Vac　○ Naïve

B.

Naïve　　　　　　　　Vaccinated

○ Blood
○ Spleen
○ LNs

1198

Figure 5: PMN phenotype following PCV vaccination. Adult C57BL/6 female mice were

injected with PCV or PBS. (A). (B-C) At 18- or 24 hours post vaccination, blood, spleen and vLNs

were collected and assessed using flow cytometry. Percent expressions of MHC-II, CD80, CD86,

CD11c, PD-L1, ARG1, CD73 and CD39 on PMNs were determined. Data show percent

expression of markers on PMNs from naïve and vaccinated mice overlayed for each tissue/organ

(B) and PMNs in all organs/tissue overlayed for naïve and vaccinated mice (C). (A) (B-C) Data

are pooled from 3 separate experiments with a total of n=5 naïve and n=6 vaccinated mice and

are presented as radar plots generated using R studio.

1207

1208

Figure 6: Effect of PMN depletion on regulatory T cells. Adult C57BL/6 female mice were

injected with PCV or PBS and depleted of their PMNs or treated with an isotype control. Spleens

were collected from all mice and assessed using flow cytometry on days 7, 14 and 21 post

vaccination. (A-B) Total (CD4⁺TCRβ⁺) and (C-D) antigen experienced (CD11a⁺CD49d⁺) T cells

were gated on. Percentages (left) and numbers (right) of (A&C) regulatory T cells (FOXP3⁺) and

(B&D) T follicular regulatory T cells (CXCR5⁺PD1⁺ FOXP3⁺) were determined. Data are pooled

from 4 separate experiments and each dot represents an individual mouse with a total of n=6

naïve, n=6 vaccinated day 7, n=8-10 vaccinated day 14, and n=6 vaccinated day 21 mice per

group. * denotes significant differences between the indicated groups as determined by Mann-

Whitney test. * is $p≤0.05$, ** is $p≤0.01$, and *** is $p≤0.001$. Bar graphs represent the mean +/- SD.

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

Figure 7: Effect of PMN depletion on Tregs in mice *in vitro*. Adult C57BL/6 female mice were

injected with PCV or PBS and depleted of their PMNs or treated with an isotype control. On day

1 post vaccination, splenocytes were isolated and cultured in the presence or absence of PMNs

for 2 days with heat-killed *S. pneumoniae* or αCD3 as stimulants. Percentage of FOXP3[+] cells are

shown. Data are pooled from separate experiments. * denotes significant differences between

the indicated groups as determined by One-Way ANOVA followed by Šídák's multiple

comparisons test. * is $p≤0.05$ and ** is $p≤0.01$. Graphs represent the mean +/- SD.

Figure 8: Effect of Treg depletion on antibody responses. Adult C57BL/6 female mice were injected with PCV or PBS and depleted of their PMNs and/or Tregs (αCD25) or treated with an isotype control. (A) Production of IgM and total IgG against type 4 pneumococcal polysaccharides were assessed on weeks 0, 2 and 4 post vaccination. Sera collected 4-weeks post vaccination were used to assess (B) the binding efficacy of antibodies to *S. pneumoniae* using flow cytometry and (C) the ability of antibodies to induce opsonophagocytic killing by PMNs. (A) Data are presented as percentage of standard sera and are pooled from a total of 3 experiments with a total of n=9 mice per group per timepoint. * denotes significant differences between PMNs- & Tregs-depleted mice and group matching the symbol color as determined by Kruskal-Wallis test followed by Dunn's multiple comparisons test. (B) Data are from 5 separate experiments with

each dot representing the average of technical replicates. * denotes significant differences

between the indicated groups as determined by One-way ANOVA followed by Šídák's multiple

comparisons test and # denotes significant differences from naïve as determined by One-way

ANOVA followed by Holm-Šídák's multiple comparisons test. *** is $p{\leq}0.001$, and # is $p{\leq}0.001$.

Bar graphs represent the mean +/- SD. (C) Data are pooled from 3 separate experiments with

each dot representing the average of technical replicates from an individual mouse. All line graphs

represent the mean +/- 95% confidence interval and all bar graphs represent the mean +/- SD.

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

A.

B.

1286

Figure 9: Effect of Treg depletion on the protective efficacy of sera against pneumococcal

infection. Adult C57BL/6 female mice were injected with PCV or PBS and depleted of their PMNs

and/or Tregs (αCD25) or treated with an isotype control. Sera were collected from all mice 4

weeks post vaccination. Sera from each mouse group was adoptively transferred into a new set

of naïve mice which were subsequently infected with 3×10^5 CFU of *S. pneumoniae*. (A) Bacterial

burden in the blood was assessed 45 hours post infection. (B) All mice were monitored for survival

for 7 days following infection. (A) Data are pooled from 2 separate experiments and each dot

represents an individual mouse with a total of n=10 mice per group. * denotes significant

differences between the indicated groups as determined by Kruskal-Wallis test followed by Dunn's

multiple comparisons test. Bar graphs represent the mean +/- SD. (B) Data are pooled from 2

separate experiments with a total of n=10 mice per group. * denotes significant differences

between the indicated groups as determined by Mantel-Cox test. ** is $p \leq 0.01$, and ** is $p \leq 0.01$.

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

Figure 10: PMN phenotype following PCV vaccination in humans. Young (24-29 years old) female donors were vaccinated with PCV. Blood was collected prior (week 0) to and on weeks 1 & 4 post vaccination. PMNs were isolated from peripheral blood and assessed using flow cytometry. Expressions of HLA-DR, CD10, CD86, PD-L1, ARG1, CD73 and CD39 on PMNs were determined. (A) Clustering of investigated markers throughout the weeks, (B) cluster abundance

1326   throughout the weeks, and (C) heat map of markers throughout the different weeks and clusters

1327   were analyzed using FLOSOM via OMIQ. Data are pooled from six donors.

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

Figure 11: Human PMN and PBMC co-cultures following PCV vaccination. Young (24-29 years old) female donors were vaccinated with PCV. One week post vaccination, PBMCs and PMNs were isolated from peripheral blood. PBMCs were cultured in the presence or absence of matched donor PMNs for 2 days with PCV as a stimulant. (A) Percentage and (B) proliferation of total (left) and antigen experienced (right) FOXP3$^+$ T cells (CD4$^+$) are shown. Data are pooled from six donors and each dot represents a donor. Data are presented as fold change of PBMC divided by PBMC +PMNs conditions. The dashed lines represent a 20% difference in responses between the conditions. Graphs represent the mean, and # indicate significant differences from 1 as measured by a one-sample t-test.

1362