DRAFT

Dosage-sensitive *RBFOX2* autoregulation promotes cardiomyocyte differentiation through transcriptome maturation

Mengmeng Huang[1,2], Feria A. Ladha[1,2], Yunxia Wang[1,2], Michael A. Trembley[1,2], Hui-Min Yin[1,2], Rongbin Zheng[1,2], Maksymilian Prondzynski[1,2,†], Yashavi Tharani[1], Alexander A. Akerberg[1,2], Stefan Aigner[3], Brian A. Yee[3], Joshua Mayourin[1,2], Sarah U. Morton[2,4], Vassilios J. Bezzerides[1,2], William T. Pu[1,2], Gene W. Yeo[3], Kaifu Chen[1,2], C. Geoffrey Burns[1,2],* Caroline E. Burns[1,2]*

1 Department of Cardiology, Boston Children's Hospital, Boston, MA 02115, USA.

2 Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA

3 Department of Cellular and Molecular Medicine, University of California San Diego, La Jolla, CA 92093, USA.

4 Division of Newborn Medicine, Boston Children's Hospital, MA 02115, USA.

* Correspondence: Caroline.Burns@childrens.harvard.edu (C.E.B.);
Geoff.Burns@childrens.harvard.edu (C. G. B.)

Author Notes: †Present Address: Department of Biomedical Physiology and Kinesiology, Simon Fraser University, Burnaby, BC V5A 1S6, Canada; Cellular and Regenerative Medicine Centre, BC Children's Hospital Research Institute, Vancouver, BC V5Z 4H4, Canada

SUMMARY:

Haploinsufficiency of the RNA splicing regulator, *RBFOX2*, is linked to congenital heart disease (CHD), yet its pathogenic mechanisms remain unclear. Here, we demonstrate that RBFOX2 is essential for progressing cardiomyocyte (CM) differentiation by shifting exon usage profiles to more mature states in sarcomere, cytoskeletal, and focal adhesion genes, including ALPHA-ACTININ-2 (ACTN2). This maturation program is initiated by critical levels of RBFOX2 that facilitate self-splicing at mutually exclusive exons encoding early and late isoforms with distinct functional roles. In heterozygous CMs, self-splicing is disrupted, which skews isoform ratios and generates a dominant-negative caused by exon co-inclusion. Finally, we demonstrate that overexpression of correctly spliced ACTN2 rescues heterozygous, but not null, phenotypes by restoring contractility, which triggers a mechanosensing feedback loop involving upregulation of RBFOX2 from the wildtype allele and transcriptome maturation. Our data suggest that decreased RBFOX2 dosage and self-splicing impairs CM differentiation, contributing to CHD pathogenesis and heart failure susceptibility.

INTRODUCTION

Congenital heart disease (CHD) is among the most common birth defects, affecting ~1% of live births and accounting for nearly 40% of childhood morbidity from congenital anomalies[1,2]. Although advances in prenatal diagnosis, surgical palliation, and perioperative management have improved survival, ventricular dysfunction that progresses to heart failure (HF) remains a major cause of mortality across the life span[3,4].

This risk is particularly high in complex and heterogenous CHD subtypes such as hypoplastic left heart syndrome (HLHS) where left-sided structures, including the left ventricle, aorta, and valves, are variably underdeveloped[5]. Standard care involves three staged palliative surgeries to reroute the circulation for a single ventricle (SV) physiology, with the first two operations performed during the first year of life. In this first year, when the balance between the systemic and pulmonary circulations is unstable, nearly 34% of infants die or require a heart transplant[6–8]. The third surgical stage, the Fontan procedure, occurs at 2-4 years of age[9]. In these patients, the need for transplant remains high, primarily due ventricular dysfunction that progresses to HF[8,10]. In fact, fewer than half of people with HLHS remain transplant-free by 10-years of age[8,11–13]. These divergent trajectories in the HLHS population suggest that additional factors, including genetics, influence HF susceptibility in the setting of altered hemodynamics. Moreover, patients surviving with other forms of CHD also show this differential susceptibility to HF[14]. As such, new mechanistic insights into the genetic causes of CHD, and their impact on ventricular dysfunction and HF outcomes, are needed for improved risk stratification and targeted clinical intervention.

One gene associated with CHD, including HLHS, is *RBFOX2*[15,16], which encodes a highly conserved RNA binding protein with known roles in alterative splicing[17]. In a cohort of

11,555 probands, 74 individuals carried heterozygous *RBFOX2* predicted damaging variants across diverse CHD subtypes, including 10 with HLHS[16]. Although the overall number was too low to reach genome wide statistical significance, participants with HLHS were disproportionately enriched for protein-truncating or protein-damaging variants, suggesting a link between HLHS and *RBFOX2* haploinsufficiency.

As observed with many CHD genes, this dosage sensitivity appears unique to humans. In mice, loss of Rbfox2 in the *Nkx2-5* lineage causes embryonic lethality with hearts consisting of a common ventricle and outflow tract[18]. However, heterozygous animals are viable and phenotypically normal. We reported similar outcomes in zebrafish, where the function of RBFOX2 is split between two paralogs, *rbfox1l* and *rbfox2*[19]. Furthermore, despite their broad expression throughout the heart, myocardial-specific re-expression of Rbfox1l is sufficient to rescue heart structure and function in zebrafish double mutants, implicating defective myocardium as causal. While these cross-species comparisons established *RBFOX2* as an essential regulator of cardiac development, the species-specific differences in dosage sensitivity underscores the need for a human-based system to model the effects of *RBFOX2* haploinsufficiency, specifically in the cardiomyocyte (CM) lineage.

One strategy involves the use of genetically engineered human induced pluripotent stem cells (iPSCs) and their derivatives, including CMs[20]. Advantages include well-established protocols for CM differentiation and stage-specific profiling where specific cellular and molecular features have been defined along the iPSC-to-CM continuum that closely parallels fetal heart development, the time when HLHS arises[20,21]. Beyond differentiation, functional maturation can also be assessed *in vitro* using engineered heart tissues (EHTs) assembled from iPSC-CMs.

Here, we generated and characterized human iPSC-CMs and EHTs carrying heterozygous (het) or null RBFOX2 truncating alleles. Our data demonstrate that RBFOX2 is essential for progressing CM differentiation by shifting exon usage profiles, including those of *ALPHA-ACTININ-2 (ACTN2),* to more mature states. Moreover, we found that this process of transcriptome maturation is facilitated by critical levels of RBFOX2 that initiate self-splicing of RBFOX2 transcripts produced by the wildtype allele at mutually exclusive exons encoding early and late isoforms. In het CMs, this critical threshold is never met, leading to defective self-splicing that skews the early-to-late isoform ratio and generates protein-truncating isoforms caused by exon co-inclusion. Overexpression and rescue studies in iPSC-CMs, EHTs, and zebrafish suggest that the early isoform facilitates the initial stages of CM differentiation, the late isoform promotes CM maturation, while the aberrant isoform exerts potent dominant-negative activity. Finally, we demonstrate that overexpression of correctly spliced ACTN2 rescues het, but not null, phenotypes by restoring contractility, which triggers a mechanosensing feedback loop involving upregulation of the remaining RBFOX2 wildtype allele and transcriptome maturation. Collectively, our study offers a mechanistic framework for how RBFOX2 haploinsufficiency contributes to CHD pathogenesis and eventual susceptibility to HF. By extension, correcting RBFOX2 or ACTN2 expression or splicing might benefit people with CHD related to impaired RBFOX2 function.

RESULTS

Genetically engineered human iPSCs model RBFOX2 haploinsufficiency
To model *RBFOX2* deficiencies in human cardiomyocytes (CMs), we introduced deleterious mutations in exon 3, which is upstream of the RNA Recognition Motif (RRM; Figure 1A). We

isolated a heterozygous clone carrying a 1 base pair (bp) insertion in one allele (het cells) and a null clone with biallelic frameshifting mutations, including a 1 bp insertion and a 2 bp deletion (null cells). Both mutations shift the open reading frame and produce a premature stop codon. RBFOX2 deficiency did not alter stemness as iPSCs from each genotype similarly expressed the SSEA4 pluripotency marker by flow cytometry (Figure S1A)[22]. Using well-established small-molecule protocols (Figure S1B)[23,24], control, het, and null iPSCs were differentiated to the CM lineage and efficiency was assessed by cardiac Troponin T (cTnT) staining on day 21. The percentage of cTnT-expressing cells was comparable between genotypes (Figure S1C), which also showed dose-dependent reductions in RBFOX2 transcript and protein levels (Figures 1B and S1D). These data confirm that the genetic lesions reduce RBFOX2 as expected and show that RBFOX2 deficiency does not impair iPSC pluripotency or CM identity.

Molecular and cellular hallmarks have been defined along the directed differentiation continuum where iPSCs transition to mesoderm by day 2, CM progenitors by day 5, and early CMs by day 8 when spontaneous contractions begin. By day 15, CMs have reached a fetal-like morphological state where they are maintained in 2D culture or matured in 3D engineered heart tissues (EHTs) (Figure 1C)[20,21]. Under brightfield microscopy, we found that day 15 het CMs contract qualitatively similar to controls but with more sporadic activity (Video S1). In contrast, null CMs remained non-contractile (Video S1), mirroring the functional defects observed in *rbfox1l;rbfox2* zebrafish mutants[19]. Quantitatively, het CMs exhibited elevated spontaneous beat frequencies relative to controls (Figure 1D), which is consistent with a less mature electrophysiological phenotype[25].

During replating, we found that more het and null CMs were required to reach confluence, suggesting impaired cell growth, proliferation, or survival. Indeed, a greater

proportion of het and null CMs appeared smaller and rounder on day 15 than controls (Figure 1E). Quantitative analyses confirmed reduced CM areas (Figure 1F), and Vybrant adhesion assays demonstrated significantly impaired attachments to the matrix (Figure 1G). Although reduced CM proliferation has been linked to reduced left ventricular size in HLHS hearts[26], no genotype-specific alterations in EdU incorporation were observed (Figures 1H and S1E). Moreover, apoptotic rates assessed by TUNEL staining were also unchanged in het CMs but slightly elevated in null cells (Figures 1I and S1F). Together, these findings suggest that *RBFOX2* haploinsufficiency compromises CM growth and adhesion but not proliferation or apoptosis.

RBFOX2 promotes CM differentiation

We next evaluated additional hallmarks of CM differentiation, including myofibril structure, gene expression-based isoform switching, electrophysiology, and mitochondrial respiration. First, sarcomere alignment was assessed in day 21 CMs from each genotype using immunofluorescence for cTnT, which marks thin filaments, and ACTN2, which marks Z-discs (Figure 2A). In het CMs, myofibrils appeared less assembled with a 2-fold increase in the myofibrillar disarray index compared to controls (Figure 2B). Null CMs expressed cTnT and ACTN2 but lacked discernible myofibrils and Z-discs altogether (Figure 2A), similar to observations in zebrafish *rbfox1l;rbfox2* mutant hearts[19]. Transmission electron micrographs further supported these observations. Specifically, Z-discs appeared less defined in het CMs with significantly reduced heights (Figures 2C and 2D) but no differences in Z-disc to Z-disc lengths (Figure S2A).

As CM differentiate both *in vitro* and *in vivo*, their transcriptomes also progressively mature through differential gene expression and alternative splicing. In terms of gene expression,

immature CMs predominantly express MYH6, which is gradually replaced by MYH7 as the cells mature[20]. Accordingly, the MYH6:MYH7 ratio serves as a reliable proxy of CM differentiation status. On day 21, het and null CMs exhibited dose-dependent increases in MYH6:MYH7 ratios (Figure 2E), consistent with impaired differentiation. Calcium transients, recorded as changes in membrane potential in paced CMs, were also abnormal in both het and null cells, exhibiting prolonged durations (CTD75; Figure 2G) and slower downstroke velocities (Figure 2H). In null CMs, accelerated repolarization kinetics (Figure S3B), reflected by shortened action potential durations (APD75) were observed (Figure S2C). These changes in electrophysiology could reflect delays in CM differentiation or a fate switch from more ventricular to atrial phenotypes. Because we did not observe decreased expression of ventricular markers or increased expression of atrial markers (Figures S2D and S2E), we conclude that these alterations are caused by stalled differentiation rather than a fate switch.

We next assessed mitochondrial area, number, and respiration as dynamic changes have also been associated with CM differentiation and are perturbed in HLHS[27]. Despite no differences in mitochondrial number, we found 1.3- and 1.5-fold reductions in mitochondrial area in het and null CMs, respectively, in transmission electron micrographs (Figures S2F and S2G). Using the Seahorse assay to measure oxygen consumption rates (OCRs), we found dose-dependent respiratory defects, including reduced basal respiration, maximal respiration, and ATP production (Figures 2I and 2J). These findings also parallelled observations in *rbfox1l;rbfox2* mutant zebrafish, underscoring a conserved role for *RBFOX2* in maintaining mitochondrial function across species[19]. Collectively, these structural, molecular, electrophysiological, and metabolic abnormalities suggest that *RBFOX2* deficiency broadly impairs CM differentiation. Notably, these phenotypes are similar to those observed in iPSC-CMs derived from HLHS

patients with acute heart failure[27], further indicating that diverse genetic lesions that contribute to HLHS might converge on stalled CM differentiation.

Molecular profiling defines expression and splicing changes in RBFOX2 deficient CMs

To define the molecular mechanisms underlying the observed phenotypes, we performed bulk RNA sequencing (RNA-seq) on control, het, and null CMs on day 21 of differentiation. Principal component analysis showed genotype-dependent clustering, indicating significant transcriptional divergence (Figure 3A). In terms of differentially expressed genes (DEGs), we found that het and null CMs retained hallmarks of an immature transcriptional profile compared to controls[28] (Figure 3B). Specifically, het and null cells displayed dose-dependent elevations in fetal-to-adult isoform ratios in transcripts encoding structural genes, including *MYH6:MYH7*, *MYL7:MYL2*, and *TNNI1:TNNI3*. In addition, we observed dose-dependent increases in fetal ion channel expression (e.g., *HCN4*, *HCN1*, *CACNA1G*, *CACNA1H*) with reductions in the adult versions (e.g., *SCN5A*, *KCNJ2*, *CASQ2*, *PLN*).

Because CHD patients are haploinsufficient for RBFOX2, we performed Gene Ontology (GO) analysis on DEGs between het and WT samples. In the upregulated subset, we found enrichment of unfolded protein response (UPR) and endoplasmic reticulum (ER) stress terms (Figure 3C; Spreadsheet S1) that contained genes such as canonical UPR effectors (e.g., *HSPA5*(*GRP78/BiP*), *DDIT3*(*CHOP/GADD153*), *TRIB3*(*TRB3*), *CHAC1*), ER-associated degradation (ERAD) components (e.g., *HERPUD1***,** *DNAJB9*), and protein-folding catalysts/chaperones (e.g., *PDIA4***,** *HYOU1**(ORP150)***,** *HSPB8*)(Figure 3D; Spreadsheet S1)[29]. We also observed upregulation of *MANF,* which functions to protect cells from ER stress[29,30]. In addition, terms related to muscle system process were enriched that included additional stress-

responsive genes such as *ANKRD1*, *CCN2*, *TRPC3*, *HBEGF*, and *DES* (Figure 3D; Spreadsheet S1). Importantly, many of these transcripts have also been identified as upregulated in failing human hearts[31–34] and HLHS patient-derived iPSC-CMs[35], suggesting molecular parallels between RBFOX2 deficiency and HF.

In the downregulated subset, we found enrichment of extracellular matrix (ECM) organization and extracellular structural assembly (Figure 3C; Spreadsheet S1) with representative genes including the major collagens (e.g., *COL1A1*, *COL1A2*, *COL3A1*, *COL5A1*, *COL5A2*, *COL6A3*), laminin subunits (e.g., *LAMA1*, *LAMA5*, *LAMB1*), integrins (e.g., *ITGA2*, *ITGA11*, *ITGB4*), ECM glycoproteins (e.g., *FN1*, *TNC*, *DCN*, *LUM*), and junctional components (e.g., *GJA1*, *S100A10*) (Figure 3E). These transcriptional changes align with the observed defects in cell adhesion in het CMs (Figure 1G) and mirror previously described ECM remodeling deficiencies reported in *Rbfox2* null embryonic mouse hearts[18].

Because RBFOX2 is a well-established splicing regulator, we performed replicate Multivariate Analysis of Transcript Splicing (rMATS) to identify differential alternative splicing events (ASEs). From this analysis, we discovered 1,211 ASEs across 823 genes (Figure 3F; Spreadsheet S2). The most common event was skipped exon (SE), representing 76.1% of the total. GO analysis showed a high degree of overlap with downregulated DEGs, including terms like cell junction organization and cell-matrix adhesion (Figure 3G; Spreadsheet S2). Examples of mis-spliced transcripts include *ACTN1*, *ACTN2*, *KINDLIN2*, *PTK2*, *ITGA3*, *ITGA7*, *FN1*, and *PXN*. We also identified terms associated with CM differentiation such as heart development, myofibril differentiation, and actomyosin organization with mis-spliced transcripts including *TPM1*, *TTN*, *OBSCN*, *NEBL*, *FHOD3*, *MYH10*, *PDLIM5*, and *PDLIM7* (Figure 3G, Spreadsheet

S2). These data suggest that *RBFOX2*-sensitive exons are enriched in genes regulating sarcomere assembly, cytoskeletal architecture, and cell adhesion.

To determine which transcripts are directly bound by RBFOX2 in the CM lineage, we performed eCLIP-seq on day 15 WT CMs and identified 2,966 high-confidence peaks in 963 genes that were enriched for the RBFOX2 consensus binding sequence, UGCAUG ($p < 1 \times 10^{-237}$, Figure 3H). Consistent with previously published RBFOX2 eCLIP profiles from diverse cell types[36], over 70% of binding sites were found in introns with 57.1% in distal sequences (Figure 3H), previously defined as >500 base pairs (bp) away from an exon[36]. 15.1% were found in proximal introns (Figure 3H), defined as <500 bp from an exon[36]. A smaller fraction, 12.4%, were observed in 3′UTRs (Figure 3H), which can mediate alternative polyadenylation[37]. GO analysis revealed that RBFOX2-bound transcripts were significantly enriched for muscle-specific terms, such as muscle cell differentiation, heart contraction, and actin-mediated cell contraction (Figure 3I, Spreadsheet S3). Because the RBFOX2 binding profiles vary substantially between cell types, caution should be used when interpreting data from publicly available sources.

RBFOX2 directly regulates exon usage to mature the transcriptome

To distinguish direct from indirect splicing changes, we performed dataset integration between differential ASEs identified in het CMs and eCLIP+ RBFOX2 targets. Through this analysis, we observed significant overrepresentation of GO terms including muscle structure development, cell junction organization, muscle cell differentiation, and cell junction assembly (Figure 4A; Spreadsheet S4), suggesting that RBFOX2 directly regulates these splicing programs. By projecting these changes onto an annotated map of CM-CM and CM-ECM interactions, we

found alterations in genes that facilitate force transmission during systolic contraction[38] (Figure 4B). Specific genes affected encode sarcomere components (*TPM1*), Z-disc subunits (*PDLIM-5, ACTN2, ACTN4*), focal adhesion complex members (*ITGA7*, *KINDLIN2*), and intercellular junctional proteins (e.g., *ZO-1*).

By examining specific exons with altered inclusion level differences near RBFOX2 binding peaks, we found retention of exon usage profiles previously associated with relative immaturity[21,39]. Specifically, differential exon usage has been described between day 7 and day 14 iPSC-CMs[21] as well as fetal and adult mouse hearts[40]. For example, we found that despite showing similar levels of *PDLIM5* expression, het cells predominantly include the longer version of exon 5 (Figure 4C), which is more frequently found in day 7 than in day 14 iPSC-CMs[21]. In contrast, control cells almost always included the shorter exon (Figure 4C). The same exon 5 splicing defect was also observed in zebrafish and mouse *RBFOX2* null hearts[18,19]. Moreover, we found that RBFOX2 eCLIP peaks were enriched over the longer exon 5 sequence (Figure 4C), consistent with a direct role in splicing. Another example is *ACTN2* where both expression and splicing defects were observed (Figures 4C and 7A). In addition to 42% reductions in *ACTN2* expression levels, het cells more frequently skipped exon 6 (Figure 4C), which encodes the actin binding domain that is required for transcript stability and protein function in Z-disc and sarcomere assembly[39]. RBFOX2 eCLIP peaks were also found in the proximal intron downstream of exon 6 (Figure 4C), a location linked to exon inclusion[36]. Interestingly, we found similar fetal exon biases in *ZO-1*, *ITGA7*, *ACTN4*, *TPM1*, *CAPN3*, and *KINDLIN2* (Figures 4B and S3A), which we confirmed by qRT-PCR (Figure S3B).

Based on these observations, we hypothesized that RBFOX2 might facilitate more global transcriptome maturation through alternative splicing. To test this idea, we integrated our

datasets from day 21 control, het, and null CMs with publicly available day 7 and day 14 datasets[21] from the same parental strain. We found 95 ASEs in 68 independent genes that were enriched for GO terms involving muscle and cardiac cell development, sarcomere and cytoskeletal organization, cell junctions, adhesion and ECM assembly, and ion transport regulation (Figure 4D; Spreadsheet S5). Most strikingly, heatmaps revealed that our day 21 control CMs displayed more advanced splicing profiles than those observed on day 14, indicative of increased transcriptome maturation (Figure 4D). In contrast, *RBFOX2*-deficient cells failed to progress along this continuum with null CMs exhibiting less advanced profiles than those on day 7, and het CMs showing an intermediate phenotype (Figure 4D). Overall, these splicing abnormalities are consistent with dose-dependent requirements for RBFOX2 in promoting global transcriptome maturation, which is required to support progression of CM differentiation.

Based on these molecular profiles, we suspected that contractile function might be more significantly impaired in the context of 3D-tissue levels[41] than in monolayer culture as the matrix environment, which promotes cell-cell and cell-ECM attachments, is likely compromised[42]. We generated engineered heart tissues (EHTs) from WT, het, and null CMs using standard methods[43], cultured them for 28 days, and quantified metrics of contractility in paced samples. Unlike WT EHTs that showed robust force with typical contraction-relaxation kinetics, null EHTs were non-contractile (Figures 4E and 4 F, Video S2), reflecting their phenotype observed in 2D culture (Video S1). Het EHTs, however, appeared more functionally impaired than their 2D counterparts. Indeed, we found significant reductions in force generation with prolonged contraction-relaxation times compared to controls (Figures 4E and 4F; Video S2). Collectively, these data demonstrate that RBFOX2 deficiency disrupts transcriptome maturation, and in turn, CM differentiation and tissue-level function.

Primary defects in RBFOX2 self-splicing alter RBFOX2 isoform ratios in het CMs.

We next sought to identify initiating molecular events that might drive widespread mis-splicing. Among the eCLIP+ transcripts with differential ASEs was *RBFOX2* itself, which was mis-spliced in het CMs at previously described mutually exclusive exons encoding distinct C-terminal domains (Figure 5A)[40,44]. Inclusion of the 40 bp exon produces an amino acid sequence containing three tyrosine residues that was shown to promote assembly of the large assembly of splicing regulators (LASR), a higher-order protein complex that enhances splicing activity[44]. In contrast, inclusion of the 43 bp exon generates a sequence lacking these tyrosines, which prevents LASR formation. Both mouse and human CMs predominantly include the 40 bp exon during fetal life, which becomes almost completely replaced by the 43 bp exon in adulthood[40].

In keeping with stalled differentiation, het CMs preferentially included the 40 bp exon on day 21compared to controls, which favored the 43 bp exon (Figure 5A and 5B). Strikingly, het CMs also produced an aberrant isoform that included both exons (40+43; Figure 5B), which shifts the open reading frame, introduces a premature stop codon, and truncates the protein by 70 amino acids. This truncation eliminates the C-terminal splicing domain while preserving the RNA Recognition Motif (RRM), suggesting it might have dominant negative activity. Consistent with autoregulation, RBFOX2 eCLIP peaks were also enriched in the proximal intron upstream of the early exon (Figure 5B), a position associated with exon exclusion[44]. From these data, we conclude that exclusion of the early exon is sensitive to RBFOX2 dosage, which when reduced compromises transcript integrity once the late exon starts to be added.

To define the temporal dynamics of this splicing event, we quantified *RBFOX2* isoform ratios over a CM differentiation time course using fluorescent DNA fragment analysis (Figure

5C). In control cells, progressive declines in the percentages of early isoform expression were observed beginning on day 5 and continuing through day 21 of differentiation (Figure 5C). Concomitant increases in the late isoform were also seen, consistent with a temporal shift in RBFOX2 isoform expression. While this transition was significantly comprised in het CMs, perhaps the most profound difference was the continual rise in 40+43 isoforms (Figures 5C and 5D). Compared to WT CMs that expressed 2.6% and 5.4% on days 5 and 21, respectively, het CMs expressed 10.1% and 25.4%, representing a 3.9-4.7-fold increase over time. Moreover, the appearance of 40+43 isoforms coincided with RBFOX2 upregulation on day 5, which peaked on day 8 before subsequently declining. While a similar pattern was observed in het CMs, levels were consistently reduced by half (Figure 5E).

Overall, our data support a model where *RBFOX2* expression is upregulated on day 5, reaching the level required to reliably exclude the early 40 bp exon. In het CMs, this critical threshold is never met, resulting in sustained early exon inclusion. This initial defect becomes problematic with addition of the late exon, resulting in a progressive accumulation of 40+43 transcripts with potential dominant-negative activity. Although these abnormal transcripts were also detected in WT CMs, they were 4 to 5-fold less abundant than in het cells, where they constituted 25% of total *RBFOX2* mRNA. Thus, the primary defect caused by RBFOX2 haploinsufficiency is inadequate levels to consistently repress early exon inclusion, which leads to progressive accumulation of 40+43 isoforms that likely contribute to disease pathogenesis.

Functional assessments of RBFOX2 isoforms

To define the functions of each RBFOX2 isoform, we performed overexpression and rescue studies in WT, het, and null CMs. iPSCs were transduced with lentivirus to stably overexpress

epitope-tagged early, late, and 40+43 isoforms or the epitope tag alone (+FLAG). Immunoblot

analysis confirmed robust RBFOX2 isoform overexpression compared to baseline (Figure S6A).

Focusing on the early and late isoforms, we performed directed differentiation of iPSC

clones from each group and assessed sarcomere assembly and organization on day 21 by

immunofluorescence (Figures 6A and 6B). In WT CMs, overexpression of either early or late

isoforms had no effect, suggesting that high levels of either are non-toxic. Compared with the

+FLAG control, overexpression of the early isoform was sufficient to rescue the myofiber

disarray index in both het and null CMs, whereas the late isoform was only effective in het cells.

These findings suggest that the presence of the early isoform is essential to support the initial

stages of CM differentiation, which cannot be compensated for by the late isoform alone.

We next assessed contractile performance at the tissue level. In WT EHTs, early isoform

overexpression reduced force generation by 1.44-fold, while late isoform overexpression

augmented it by 1.64-fold. These changes were accompanied by alterations in contraction and

relaxation times (Figures 6C and S5A). In het EHTs, both isoforms restored contractile force to

control levels and normalized contraction–relaxation kinetics (Figures 6C and S5A). However, in

null EHTs, only the early isoform was sufficient for rescue (Figures 6C and S5A). Consistent

with our findings from 2D culture, these observations support distinct requirements for each

RBFOX2 isoform, with the early version supporting CM differentiation and the later version

promoting maturation. Consistently, the late isoform cannot substitute for the early isoform,

which is essential for CM differentiation.

Because 40+43 isoforms are predicted to have dominant-negative activity, we tested this

directly through overexpression in WT iPSCs (Figure S4B). As described above, we evaluated

sarcomere assembly by immunofluorescence on day 21 and contractility in EHTs on day 28.

Strikingly, 40+43-overexpressing CMs failed to form sarcomeres altogether, phenocopying the myofibril assembly defects observed in null CMs (Figure 6D). Similarly, EHTs were non-contractile (Figure 6E). These data demonstrate potent dominant-negative functions for 40+43 isoforms and suggest that its progressive accumulation in RBFOX2 haploinsufficient hearts might contribute to CHD.

To determine whether these isoform-specific roles are conserved *in vivo*, we performed rescue studies in zebrafish *rbfox*1l;*rbfox2* double knock-out embryos (DKO) by injecting mRNA encoding early, late, or 40+43 human RBFOX2 isoforms. Consistent with our previous findings, early isoform overexpression largely suppressed pericardial edema and restored ventricular morphology in mutant animals (Figure 6F)[19]. By contrast, late or aberrant isoform overexpression failed to rescue either phenotype (Figure 6F). Quantification of fractional area change (%FAC) confirmed that only the early RBFOX2 isoform, but not the late or 40+43 versions, restored contractility in DKO hearts (Figure 6G). Together, these findings support a division of labor between the early and late RBFOX2 isoforms during CM differentiation and functional maturation, respectively, and suggest that the 40+43 isoform is pathogenic.

ACTN2 overexpression rescues RBFOX2 haploinsufficiency through a positive feedback loop
RBFOX2 deficiency disrupts the expression and splicing of genes connecting the sarcomere to the ECM, including *ACTN2* (Figures 4B and 4C)[45]. To perform these functions, ACTN2 requires its actin-binding domain (ABD), encoded by exon 6. We found that exon 6 inclusion depends on RBFOX2, which binds downstream of the exon in a position known to promote inclusion (Figure 4C)[36]. Furthermore, addition of exon 6 begins on day 5 of CM differentiation[39], coinciding with *RBFOX2* upregulation (Figure 5E). Importantly, *ACTN2* transcripts lacking exon 6 are highly

unstable, likely due to a frameshift that introduces a premature stop codon and triggers non-sense mediated decay (NMD)[39]. Consistent with this mechanism, qPCR analysis confirmed dose-dependent reductions in *ACTN2* mRNA levels in het and null CMs (Figure 7A). Together, these findings suggest that overexpression of exon 6-containing ACTN2 isoforms in het and null CMs might rescue sarcomere organization and contractility in 2D culture and EHTs.

We transduced WT, het, and null iPSCs with lentivirus to stably overexpress epitope-tagged exon 6-containing ACTN2 (+ACTN2) or the epitope tag alone (+FLAG). Immunoblot analysis confirmed robust ACTN2 overexpression compared to baseline (Figure S4C). In both het and null CMs overexpressing ACTN2, sarcomere assembly appeared more organized on day 21 by immunofluorescence compared to the FLAG tag alone (Figure 7B). Quantifications of the myofibrillar disarray index confirmed rescue to WT levels in het CMs overexpressing ACTN2 (Figure 7C). However, rescue was only partial in null CMs + ACTN2, reaching levels comparable to het CMs + FLAG. In line with these findings, contractility was restored by ACTN2 in het CMs but showed only partial improvement in null cells.

In WT EHTs, ACTN2 overexpression augmented contractile force and contraction-relaxation kinetics, indicative of increased maturation (Figures 7D and S5B). Consistent with our findings in 2D culture, ACTN2 overexpression rescued contractile force and contraction-relaxation times in het EHTs, but not null. Remarkably, these findings demonstrate that ACTN2 overexpression can bypass RBFOX2 haploinsufficiency, but not full deletion, to restore CM differentiation and function. Moreover, this discrepancy between genotypes suggested that Z-disc stabilization and contractile function might initiate a positive feedback loop that matures the transcriptome through RBFOX2 upregulation, which would only be possible in the presence of a WT allele.

Indeed, ACTN2 restored total RBFOX2 expression to WT levels in het CMs (Figure 7E). We also found restoration of more appropriate isoform ratios in het cells with lower early, higher late, and decreased 40+43 RBFOX2 transcripts (Figure 7F). Because RBFOX2 levels and isoform ratios were rescued by ACTN2 overexpression, we assessed whether the more global RBFOX2-dependent maturation program might also be restored. Indeed, fetal-to-adult isoform ratios for MYH6:MYH7 and MYL7:MYL2 were rescued back to WT levels in het CMs (Figures 7G and 7H) and exon usage for PDLIM5 and TPM1 were also restored (Figures 7I and 7J). These results demonstrate that ACTN2 overexpression can bypass RBFOX2 haploinsufficiency to rescue CM differentiation and function by initiating a positive feedback loop that matures the transcriptome through RBFOX2 upregulation. By extension, these findings offer new therapeutic opportunities to stave off HF in RBFOX2 haploinsufficient hearts.

DISCUSSION

Here, we present a human model of RBFOX2 haploinsufficiency and identify molecular defects in RBFOX2 expression and self-splicing as initiating events that disrupt transcriptome maturation, which compromises CM differentiation and adhesion. These cellular defects were so severe that het and null CMs could not survive standard freeze-thaw protocols or assemble into embryoid bodies (EBs) under typical bioreactor conditions. Similar CM differentiation and adhesion defects that ultimately impair heart development have been reported in null mice when RBFOX2 is deleted at the CM progenitor stage (*Nkx2-5* lineage)[18]. By contrast, deletion of *RBFOX2* after CM differentiation has occurred (*Mlc2v* lineage) causes no developmental abnormalities[46]. Together, these data suggest that RBFOX2 is essential during the progenitor-to-

CM transition, but dispensable for subsequent steps of cardiac morphogenesis once a critical level of differentiation is achieved.

Defects in CM differentiation have been increasingly recognized as a common phenotype across genetically heterogeneous HLHS cases[35,47,48]. Molecular profiling of infant HLHS myocardium revealed retention of a fetal-like transcriptional program[35]. Likewise, single cell RNA-sequencing demonstrated distinct clustering of day 14 iPSC-CMs from control and HLHS individuals, with HLHS cells enriched in progenitor populations and depleted in mature ones[35]. Similar findings have been reported in iPSC-CMs derived from independent HLHS probands[27,47]. Because impaired CM differentiation is consistently observed across genotypes in both 2D and 3D *in vitro* systems, these data collectively suggest that HLHS is not merely a consequence of altered hemodynamics but arises from intrinsic myocardial defects.

We also found that contractility was sensitive to RBFOX2 dosage, again mirroring the functional abnormalities observed in iPSC-CMs derived from diverse HLHS genetic backgrounds[27,35,47]. On a molecular level, our RNA-seq analysis showed upregulation of UPR, ERAD, and fetal gene programs, consistent with finding in HLHS patient-derived iPSC-CMs[27,35], and adult HF samples [49–53]. In mice, decreased RBFOX2 expression was also observed in response to pressure overload-induced heart failure, which also associated with retention of target fetal exons[46,54]. Furthermore, CM-specific deletion of RBFOX2 leads to dilated cardiomyopathy followed by HF and lethality[46], providing *in vivo* relevance to the functional deficiencies observed *in vitro*. Although HF outcomes in patients with HLHS caused by RBFOX2 haploinsufficiency remain unknown, these findings suggest that these individuals may be at elevated risk for HF and likely express a less mature transcriptome.

Particularly intriguing are the potentially distinct mechanisms underlying stalled CM differentiation and transcriptome maturation in RBFOX2 null versus het CMs. In null CMs, exon usage fails to shift toward more mature patterns due to the complete absence of RBFOX2 protein. In contrast, het CMs present a more complex scenario in which overall expression is reduced by half, and isoform ratios, derived from the wildtype allele, are skewed with a substantial fraction encoding a truncated protein with dominant negative activity.

Comparable defects resulting from altered *RBFOX2* levels and isoform production have also been linked to cardiac phenotypes in other disease contexts. For example, diabetic patients frequently develop cardiovascular complications, including cardiomyopathy[55]. In both a mouse model of type 1 diabetes and in patients with type 2 diabetes, elevated glucose levels increase *RBFOX2* expression in the heart, promoting skipping of exon 6[56]. Exclusion of this exon, which encodes half of the RRM, generates a dominant negative isoform that causes mis-splicing of RBFOX2 target RNAs prior to the onset of cardiac phenotypes[56–58].

Similarly, patients with myotonic dystrophy type 1 (DM-1) often develop cardiac conduction delays and ventricular arrhythmias[59–61]. This phenotype has been linked to overexpression of the early *RBFOX2* isoform in both DM-1 patient hearts and a corresponding mouse model[40]. Mice engineered to overexpress or exclusively express wildtype levels of the early *RBFOX2* isoform live to adulthood but develop cardiac conduction defects that recapitulate the human disease phenotypes.

In the context of CHD, determining whether altered RBFOX2 expression levels, isoform imbalance, or dominant negative effects contributes most to the observed phenotypes will require more rigorous genetic testing. For example, it will be informative to assess how CM differentiation and contractility are affected in iPSCs engineered to only express wildtype levels

of either the early or late isoform, or to express half-doses of the early isoform in the absence of the dominant negative protein. Even in the absence of such data, the identification of variants in four HLHS probands that are predicted to truncate RBFOX2 within the same region as the dominant negative isoform supports the idea that this aberrant protein product is pathogenic[15,16]. It will also be important to determine whether missense variants that decrease RBFOX2 activity also lead to gradual accumulations of 40+43 isoforms, ultimately compromising CM function.

Remarkably, we discovered that the phenotypes caused by *RBFOX2* haploinsufficiency can be bypassed by ACTN2 overexpression. Prior work has demonstrated that sarcomeres initially assemble at protocostameres, sites of CM-ECM adhesion that resemble focal adhesions in terms of their molecular makeup[62]. ACTN2 acts a critical link between myosin and protocostameres to transmit initial mechanical forces that promote Z-disc and sarcomere maturation[62–65].

In summary, our study identifies *RBFOX2* isoform balance as a dosage-sensitive regulator that integrates post-transcriptional control with cardiomyocyte maturation. By linking splicing dysregulation to structural and functional failure, we establish a mechanism by which haploinsufficiency drives CHD-associated HF. Importantly, our findings highlight isoform correction and maturation-targeted strategies as potential avenues to reduce HF risk, offering a framework for precision interventions in this vulnerable patient population.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Caroline E. Burns (Caroline.Burns@childrens.harvard.edu) and C. Geoffrey Burns (Geoff.Burns@childrens.harvard.edu)

Materials availability

Requests for resources and reagents are available from the lead contact upon request.

Data and code availability

- The Bulk RNA-seq and eCLIP-seq datasets generated and/or analyzed during this study have been deposited in the Gene Expression Omnibus (GEO) under accession numbers GEO: XXXX and GEO: XXXX, respectively, and are publicly available as of the date of publication.
- This paper does not report original code.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

We thank the Boston Children's Hospital Aquatics Facility for zebrafish care. Sanger sequencing and DNA fragment analysis were performed at the Massachusetts General Hospital (MGH) Computational and Integrative Biology DNA Core. Transmission Electron Microscope imaging was performed at the Harvard Medical School (HMS) Electron Microscopy Core. We thank Christopher S. Chen (Boston University) for providing the pRRL-mApple–α-actinin-2 plasmid.

Funding information: M.H. was supported by an American Heart Association (AHA) postdoctoral fellowship (21POST834347) and a National Institutes of Health (NIH) Pathway to Independence Award (K99HL169927-01A1). M.P. and W.T.P. were supported by funding from Additional Ventures. G.W.Y was supported by an NIH R01 (HG0046569) and NIH U24 (HG009889). This work was supported by NIH R35 (HL135831), a Peer-Reviewed Medical Research Program (PRMRP) Department of Defense (DoD) award (W81XWH-20-1-0165), and

AUTHOR CONTRIBUTIONS

M.H. designed, performed, and interpreted the experiments. F.L. assisted with lentivirus production, rescue experiments, and data quantification. Y.W., R.Z., and K.C. performed bioinformatics analyses and dataset integration. M.P., Y.T., and W.T.P. provided training to M.H. for iPSC and EHT work. J.M. provided code for EHT contractility analysis. M.P. and A.A.A. prepared eCLIP samples from 3D bioreactor WT cardiomyocytes and S.A., B.A.Y., and G.W.Y. performed eCLIP-seq experiments and bioinformatic analyses. M.T., H.Y., and V.J.B. contributed to data analysis and discussion. S.U.M. provided information on RBFOX2 variants prior to publication. C.G.B. and C.E.B. co-directed the study, secured funding, designed experiments, and interpreted data. M.H. generated the figures. M.H. and C.E.B. wrote the manuscript with input from all authors.

DECLARATION OF INTERESTS

G.W.Y. is a cofounder, member of the board of directors, scientific advisory board member, equity holder and paid consultant for Eclipse BioInnovations. G.W.Y.'s interests have been reviewed and approved by the University of California San Diego in accordance with its conflict-of-interest policies.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
RBFOX2 Antibody	Bethyl Laboratories	Cat# A300-864A; RRID: AB_609476
GAPDH Monoclonal Antibody, Clone 14C10	Cell Signaling	Cat# 2118T; RRID: AB_561053
Cardiac Troponin T (CT3, anti-cTnT)	DSHB	Cat# CT3; RRID: AB_528495
Alpha Actinin antibody [EA-53] (Anti-ACTN2)	Abcam	Cat# ab9465; RRID: AB_307264
SSEA-4 antibody, FITC	Miltenyi Biotec	Cat# 130-098-371
Cardiac Troponin T, FITC	Miltenyi Biotec	Cat# 130-119-575
IgG-FITC	Miltenyi Biotec	Cat# 130-104-611
Goat anti-mouse IgG1 secondary antibody, Alexa 488	Invitrogen	Cat# A-21121; RRID: AB_2535764
Goat anti-mouse IgG2a secondary antibody, Alexa 555	Invitrogen	Cat# A-21137; RRID: AB_2535776
Anti-rabbit IgG, HRP-linked Antibody	Cell Signaling	Cat# 7074; RRID:AB_2099233
Anti-mouse IgG, HRP-linked Antibody	Cell Signaling	Cat# 7076; RRID:AB_330924
Rabbit IgG Isotype Control	Thermo Fisher Scientific	*Cat# 02-6102; RRID: AB_2532938*
Chemicals, peptides, and recombinant proteins		
Essential 8 Medium	Gibco	Cat# A1517001
RPMI 1640 Medium, GlutaMAX Supplement	Gibco	Cat# 61870127
RPMI 1640 Medium, minus glucose	Gibco	Cat# 11879020
50X B27 (minus insulin) supplement	Gibco	Cat# A1895601
Geltrex LDEV-free stem cell-qualified reduced-growth factor basement membrane matrix	Gibco	Cat# A1413302
Penicillin-Streptomycin	Gibco	Cat# 15140122
Versene Solution	Gibco	Cat# 1540066
ACCUTASE Cell Detachment Solution	STEMCELL	Cat# 07920
Doxycycline Hydrochloride	Sigma-Aldrich	Cat# D3072
Y-27632 dihydrochloride	Tocris	Cat# 1254
CHIR99021	STEMCELL	Cat# 72054
IWR-1-endo	STEMCELL	Cat# 72564
IWP 2	Tocris	Cat# 3533
Sodium L-lactate	Sigma-Aldrich	Cat# L7022
Insulin solution human	Sigma-Aldrich	Cat# I9278
Matrigel hESC-Qualified Matrix, LDEV-free	Corning	Cat# 354277
Thrombin from bovine plasma	Sigma-Aldrich	Cat# T7513-100UN
Aprotinin	Sigma-Aldrich	Cat# A1153-25MG
Fibrinogen from bovine plasma	Sigma-Aldrich	Cat# F8630-1G
DMEM, high glucose, pyruvate	Gibco	Cat# 11995065
OptiMEM	Gibco	Cat# 31985070

Fetal Bovine Serum (FBS)	R&D Systems	Cat# S11150
Puromycin	Boston BioProducts	Cat# ABT-440
Polyethylenimine (PEI 100000)	Polysciences	Cat# 25414-2
DAPI	Sigma-Aldrich	Cat# D9542
Hoechst 33342	Invitrogen	Cat# H3570
RIPA buffer	Thermo Fisher Scientific	Cat# 89900
Halt Protease Inhibitor Cocktail	Thermo Fisher Scientific	Cat# 78430
Pierce Lane Marker Reducing Sample Buffer	Thermo Fisher Scientific	Cat# 39000
SuperSignal West Pico PLUS Chemiluminescent Substrate	Thermo Fisher Scientific	Cat# 34577
TRIzol	Invitrogen	Cat# 15596-026
Paraformaldehyde	Electron Microscopy Sciences	Cat# 15710
Triton X-100	Sigma-Aldrich	Cat# T8787
BSA	Sigma-Aldrich	Cat# A3059
Critical commercial assays		
Human Stem Cell Nucleofector Kit 1	Lonza	Cat# VPH-5012
MEGAshortscript T7 Transcription Kit	Invitrogen	Cat# AM1354
mMESSAGE mMACHINE SP6 Transcription Kit	Invitrogen	Cat# AM1340
Direct-zol RNA Microprep Kits	Zymo Research	Cat# R2062
SuperScript III Reverse Transcriptase	Invitrogen	Cat# 18080044
Fast SYBR Green PCR Master Mix	Applied Biosystems	Cat# 4385612
Phusion High-Fidelity PCR Master Mix with GC Buffer	Thermo Fisher Scientific	Cat# F532S
Pierce BCA Protein Assay Kits	Thermo Fisher Scientific	Cat# 23225
Trans-Blot Turbo RTA Mini 0.2 µm PVDF Transfer Kit	Bio-Rad	Cat# 1704272
BD Cytofix/Cytoperm Fixation/Permeabilization Kit	BD Biosciences	Cat# 554714
Propidium Iodide Flow Cytometry Kit	Abcam	Cat# ab139418
Vybrant Cell Adhesion Assay Kit	Invitrogen	Cat# V13181
Click-iT Plus EdU Cell Proliferation Kit for Imaging, Alexa Fluor 647 dye	Invitrogen	Cat# C10640
In Situ Cell Death Detection Kit, TMR red	Sigma-Aldrich	Cat# 12156792910
FluoVolt Membrane Potential Kit	Invitrogen	Cat# F10488
X-Rhod-1, AM, cell permeant	Invitrogen	Cat# X14210
Seahorse XF Cell Mito Stress Test Kit	Agilent	Cat# 103015-100
HiFi DNA Assembly Cloning Kit	NEB	Cat# E5520S
Lenti-X Concentrator	Takara Bio	Cat# 631231
Deposited data		

Bulk RNA sequencing data of human iPSC-CMs	This paper	GEO:
eCLIP sequencing data of human iPSC-CMs	This paper	GEO:
Experimental models: Cell lines		
Human: WTC-11 iPSCs	Coriell Institute	Cat# GM25256
Human: HEK293T cells	American Type Culture Collection	Cat# CRL-3216
Experimental models: Organisms/strains		
Zebrafish: *rbfox1[chb5]*	Huang et al., 2022	N/A
Zebrafish: *rbfox2[chb6]*	Huang et al., 2022	N/A
Zebrafish: *Tg(myl7:EGFP)[f1]*	Burns et al., 2005	N/A
Oligonucleotides		
guide RNA for human *RBFOX2*: GCGTACTTCCGTAGAGTGTC<u>AG</u> <u>G</u>	This paper	N/A
PCR primers for human *RBFOX2* Genotyping: Forward: 5'-GAGCCGACAACAACTCCTGA-3' Reverse: 5'-AGGAGGAGGGAGGACAATGT-3'	This paper	N/A
PCR primers for DNA fragment analysis of human *RBFOX2* 40- and 43-bp region Forward: 5'-6-FAM-GGTACCTCCAACAGCCATCC-3' Reverse: 5'-GTGTACACCCTGCCATAA-3'	This paper	N/A
qRT-PCR primers (See Table S1)	This paper	N/A
Recombinant DNA		
pSLIK TT RBFOX2	Addgene	Cat# 59770
psPAX2	Addgene	Cat# 12260
pMD2.G	Addgene	Cat# 12259
pLenti CMV GFP Puro (658-5)	Addgene	Cat# 17448
pRRL mApple-ACTN2	Zhang et al., 2021[66]	N/A
Software and algorithms		
Fiji (ImageJ)	Fiji	http://fiji.sc; RRID:SCR_002285
GraphPad Prism 9	GraphPad Software	http://www.graphpad.com/; RRID:SCR_002798
BioRender	Created with BioRender.com	http://biorender.com; RRID:SCR_018361
MUSCLEMOTION	Sala et al., 2017[67]	https://github.com/l-sala/MUSCLEMOTION?utm_source=chatgpt.com
Seahorse Wave Software 2.4.0	Agilent	http://www.agilent.com/en-us/products/cell-analysis-(seahorse)/software-download-for-wave-desktop; RRID:SCR_014526
CyteSeer Software	Vala Sciences	https://valasciences.com/; RRID:SCR_012232

STAR (v2.7.9a)	Dobin et al.[68]	http://code.google.com/p/rna-star/; RRID:SCR_004463
RSEM (v1.3.1)	Li et al.[69]	http://deweylab.biostat.wisc.edu/rsem/; RRID:SCR_000262
DESeq2 (v1.32.0)	Love et al.[70]	https://bioconductor.org/packages/release/bioc/html/DESeq2.html; RRID:SCR_015687
clusterProfiler (v4.12.6)	Yu et al.[71]	http://bioconductor.org/packages/release/bioc/html/clusterProfiler.html; RRID:SCR_016884
rMATS (v4.1.2)	Shen et al.[72]	https://rmats.sourceforge.io; RRID:SCR_023485
Skipper (v1.1.0)	Boyle et al.[73]	https://github.com/YeoLab/skipper; RRID:SCR_026260
skewer (v0.2.2)	Jiang et al.[74]	http://sourceforge.net/projects/skewer/; RRID:SCR_001151
UMICollapse (v1.0.0)	Liu et al.[75]	https://github.com/Daniel-Liu-c0deb0t/UMICollapse
HOMER (v5.0.1)	Heinz et al.[76]	http://homer.ucsd.edu/; RRID:SCR_010881
Integrative Genomics Viewer (IGV)	Broad Institute	https://www.broadinstitute.org/igv/; RRID:SCR_011793
Metascape	Zhou et al.[77]	http://metascape.org/gp/index.html#/main/step1; RRID:SCR_016620
GeneOverlap (v1.40.0)	*Bioconductor*	https://rdrr.io/bioc/GeneOverlap/man/GeneOverlap.html; RRID:SCR_018419
BEDTools (v2.31.1)	Quinlan et al.[78]	https://github.com/arq5x/bedtools2; RRID:SCR_006646
Other		
Falcon round-bottom tubes with cell strainer cap	Corning	Cat# 352235
4–20% Mini-PROTEAN TGX precast gels	Bio-Rad	Cat# 4561095
PVDF membranes	Bio-Rad	Cat# 1704272
96 well polystyrene cell culture microplates, black	Greiner Bio-One	Cat# 655090
Seahorse XF96 cell culture microplates	Agilent	Cat# 103793-100
Silicon racks (transparent)	DNABIOS	Cat# C0001

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human iPSC derivation and culture

Human induced pluripotent stem cells (iPSCs) were derived from the WTC-11 background (Coriell Institute). Cells were maintained in Essential 8 Medium (Gibco) at 37°C in a humidified atmosphere with 5% CO_2 and passaged every 2-4 days using Versene (Thermo Fisher Scientific). Culture plates were pre-coated with 1% Geltrex matrix (Thermo Fisher Scientific) according to the manufacturer's instructions.

Zebrafish husbandry and strains

Zebrafish (Danio rerio) were maintained under standard conditions in accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) at Boston Children's Hospital, following the NIH Guide for the Care and Use of Laboratory Animals. The following strains were used: *rbfox1l*[chb5], *rbfox2*[chb6], and *Tg(myl7:EGFP)*[f1][19,79].

METHOD DETAILS

Human iPSC genome editing

CRISPR/Cas9 genome editing was performed to generate isogenic human iPSC lines. A guide RNA (gRNA) targeting *RBFOX2* (5′-GCGTACTTCCGTAGAGTGTC-3′) was synthesized using the MEGAshortscript T7 Transcription Kit (Thermo Fisher Scientific). WTC-11 iPSCs were genetically modified to express a doxycycline-inducible SpCas9 transgene in the AAVS1 (adeno-associated virus integration site 1) safe harbor locus as previously described[80,81]. To induce Cas9 expression, WTC-11 AAVS1[Cas9/+] hiPSCs were treated with doxycycline (2 μg/mL) for 16 h prior to transfection. Approximately 0.5×10^6 cells were then nucleofected with 5 μg gRNA using the Amaxa Nucleofector system (Lonza), and plated at low density for clonal isolation. Clonal genotyping of *RBFOX2* edits was performed by PCR (Forward: 5′-GAGCCGACAACAACTCCTGA-3′; Reverse: 5′-AGGAGGAGGGAGGACAATGT-3′), followed by Sanger and amplicon sequencing to identify heterozygous and null clones (n=4 independent clones per genotype).

Directed differentiation of hiPSCs into cardiomyocytes

Human iPSCs were differentiated into cardiomyocytes (iPSC-CMs) using a monolayer-based protocol with temporal Wnt signaling modulation, as previously described[24,82]. Cells were seeded onto 12-well plates and cultured for 48 h or until ~50% confluence. On day 0, cells were treated with RPMI (Thermo Fisher Scientific) supplemented with B27 minus insulin (Thermo Fisher Scientific) and 6.5–7 μM CHIR99021 (STEMCELL Technologies) for 48 h. On day 2, the medium was replaced with fresh RPMI/B27(–insulin) containing 5 μM IWR-1-endo (STEMCELL Technologies) and 5 μM IWP-2 (Tocris Bioscience) for 48 h. From days 4–10, cells were maintained in RPMI/B27(–insulin) with medium changes every 48 h. On day 10, metabolic selection was performed in glucose-free RPMI supplemented with 5 mM sodium lactate (Sigma-Aldrich) for 48 h. Cells were then recovered for at least 4 days in RPMI/B27(–insulin) prior to downstream assays. For dissociation, iPSC-CMs were incubated with Accutase (Thermo Fisher Scientific) at 37 °C for 10–15 min, resuspended in RPMI/B27 supplemented with 10 μM ROCK inhibitor (Y-27632; Tocris Bioscience).

Flow cytometry analysis

Pluripotency of hiPSCs and differentiation efficiency of hiPSC-CMs were assessed by flow cytometry. Briefly, hiPSCs and hiPSC-CMs were dissociated with Versene (5 min, 37 °C) or Accutase (10 min, 37 °C), respectively, and passed through a 70 μm cell strainer. Cells were washed and fixed using the Cytofix/Cytoperm Fixation/Permeabilization Kit (BD Biosciences) according to the manufacturer's instructions. Fixed cells were incubated overnight at 4 °C with FITC-conjugated antibodies against SSEA-4 (Miltenyi Biotec, 130-098-371), cardiac troponin T (cTnT; Miltenyi Biotec, 130-119-575), or an IgG isotype control (Miltenyi Biotec, 130-104-611), each diluted 1:100 in wash buffer. After two washes, cells were resuspended in wash buffer and transferred to Falcon tubes equipped with cell-strainer caps (Corning). Samples were kept on ice and analyzed on a BD LSRFortessa flow cytometer (BD Biosciences).

Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted using TRIzol reagent (Invitrogen) and purified with the Direct-zol Microprep Kit (Zymo Research). cDNA was synthesized with the SuperScript III First-Strand Synthesis System (Invitrogen) following the manufacturer's instructions. qRT-PCR was performed in triplicate on a QuantStudio 3 (Applied Biosystems) or CFX384 (Bio-Rad) instrument using the Fast SYBR Green Master Mix (Thermo Fisher Scientific). Relative mRNA expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method with GAPDH as an internal control. Isoform-specific primers were designed based on rMATS analysis to quantify alternative splicing events, whereas primers spanning constitutive exons were used to measure total transcript abundance. Primer sequences are listed in Supplementary Table S1.

Protein immunoblotting

Samples were lysed in RIPA buffer (Thermo Fisher Scientific) supplemented with Halt Protease Inhibitor Cocktail (Thermo Fisher Scientific). Protein concentrations were determined using the Pierce BCA Protein Assay Kit (Thermo Fisher Scientific), normalized, and reduced and denatured in sample buffer (Thermo Fisher Scientific). Equal amounts of protein were separated on 4–20% Mini-PROTEAN TGX precast gels (Bio-Rad), transferred to PVDF membranes (Bio-Rad) using the Trans-Blot Turbo Transfer System (Bio-Rad), and blocked in TBS-T (Tris-buffered saline with 0.1% Tween-20, pH 7.4) containing 5% BSA. Membranes were incubated overnight at 4 °C with primary antibodies, washed with TBS-T, and incubated with HRP-conjugated secondary antibodies (Cell Signaling, 7076 and 7074) for 1 h at room temperature. Signals were detected using SuperSignal West ECL substrate (Thermo Fisher Scientific) and imaged with a GE ImageQuant LAS-4000. Images were processed and quantified using Fiji (ImageJ). Primary antibodies used were: anti-RBFOX2 (Bethyl Laboratories, A300-864A), anti–α-actinin (Abcam, ab9465), and anti-GAPDH (Cell Signaling Technology, 2118T)

Analysis of contractility in 2D-cultured hiPSC-CMs

Contraction of 2D-cultured hiPSC-CMs in the original plates was recorded in bright-field mode (40x magnification) using a Keyence BZ-X700 microscope. Videos were captured at 33 frames per second to ensure sufficient temporal resolution for accurate

quantification of contractile dynamics. For quantitative analysis of spontaneous beat frequency, the acquired time-lapse images were processed using MUSCLEMOTION, as previously described[67].

Cell adhesion assay

Cell adhesion was quantified using the Vybrant Cell Adhesion Assay Kit (Thermo Fisher Scientific) according to the manufacturer's instructions. Briefly, flat-bottom 96-well plates (Greiner Bio-One) were pre-coated with 0.5% Geltrex matrix at 37 °C for 2 h. hiPSC-CMs were dissociated with Accutase (Thermo Fisher Scientific) for 10 min at 37 °C and labeled with 5 μM calcein AM (included in the kit) for 30 min at 37 °C. After two washes, cells were resuspended at 5×10^5 cells/mL, and 100 μL per well was seeded into the pre-coated plates and incubated for 30 min at 37 °C. Total fluorescence was measured using a FlexStation 3 Multi-Mode Microplate Reader (Molecular Devices). After gentle washing to remove nonadherent cells, fluorescence of adherent cells was recorded. Background wells contained medium only. The percentage of adhered cells was calculated as: Cell Adhesion (%) = [(Adherent fluorescence − Background) / Total fluorescence] × 100

Immunofluorescence, EdU staining, TUNEL staining and imaging

hiPSC-CMs were replated onto 1% Geltrex-coated 96-well plates (Greiner Bio-One) and allowed to recover for 4 days before fixation. Cells were fixed with 4% paraformaldehyde (Electron Microscopy Sciences) for 10 min, permeabilized in PBS containing 0.1% Triton X-100 for 5 min, and blocked in PBS containing 3% BSA. Cells were then incubated with primary antibodies overnight at 4 °C, followed by Alexa Fluor–conjugated secondary antibodies for 1 h at room temperature. Nuclei were counterstained with DAPI (1:1000; Sigma-Aldrich) for 5 min before imaging. EdU staining was performed using the Click-iT EdU Imaging Kit (Invitrogen) after a 2 h EdU pulse prior to fixation. TUNEL staining was performed using the In Situ Cell Death Detection Kit, TMR red (Sigma-Aldrich), followed by the same immunofluorescence procedures described above. Primary antibodies included Alpha-Actinin (ACTN2, 1:500; Abcam, ab9465) and cardiac Troponin T (cTnT, 1:500; DSHB, CT3). Secondary antibodies included goat anti-mouse IgG1 Alexa Fluor 488 (1:500; Invitrogen, A-21121), and goat anti-mouse IgG2a Alexa Fluor 555 (1:500; Invitrogen, A-21137). Images were acquired using an Olympus FV3000R resonant-scanning confocal microscope and processed with Fiji (ImageJ).

Morphological analysis of 2D-cultured hiPSC-CMs

For cell size analysis, hiPSC-CMs were seeded into 96-well plates for 24 h and imaged under bright field (100x magnification) using the EVOS M5000 Imaging System (Thermo Fisher Scientific). Cell area was quantified from confocal images using Fiji by manually tracing individual cell boundaries with the polygon selection tool. For myofibrillar disarray analysis, immunofluorescence staining for ACTN2 followed by confocal imaging (600x magnification) was performed as above. Myofibrillar disarray was quantified in Fiji by assessing the alignment consistency of sarcomeric Z-discs across multiple regions of interest, as previously described[83]. All analyses were performed in a blinded manner.

Transmission electron microscopy (TEM)

hiPSC-CMs were prepared and processed, and ultrathin sections were imaged at the Harvard Medical School Electron Microscopy Facility as previously described[19]. Briefly, cells were seeded onto Aclar coverslips in 24-well plates and allowed to recover for 4 days, then fixed for 1 h at room temperature in a 1:1 mixture of RPMI/B27(–insulin) and FGP fixative (2.5% paraformaldehyde, 5% glutaraldehyde, and 0.06% picric acid in 0.2 M cacodylate buffer). Samples were post-fixed in 1% osmium tetroxide/1.5% potassium ferrocyanide, stained with 1% uranyl acetate in maleate buffer (pH 5.2), dehydrated through a graded ethanol series, transitioned through propylene oxide prior to resin infiltration, and embedded in Taab 812 resin (Marivac-Canemo). Ultrathin sections (80 nm) were cut using a Leica Ultracut S ultramicrotome, post-stained with 0.2% lead citrate, and imaged using either a Philips Tecnai BioTwin Spirit or a JEOL 1200× transmission electron microscope.

Seahorse analyzer analysis of oxygen consumption rate (OCR)
Mitochondrial function in hiPSC-CMs was assessed using the Seahorse XF Cell Mito Stress Test Kit (Agilent) according to the manufacturer's instructions. hiPSC-CMs were seeded at 7.5×10^4 cells per well onto 1% Geltrex-coated Seahorse XF96 cell culture microplates (Agilent). Cells were maintained in RPMI/B27(–insulin) medium for 4 days to allow recovery prior to the assay. Oxygen consumption rate (OCR) was recorded using a Seahorse XF96 Extracellular Flux Analyzer, and data were analyzed with Wave software (version 2.4.0).

Calcium transient and membrane potential analysis
Calcium and membrane voltage imaging was recorded on a Vala Sciences Kinetic Image Cytometer (KIC) as previously described with some modifications[84]. Briefly, 1×10^5 hiPSC-CMs were seeded onto black-walled 96-well plates (Greiner Bio-One) and cultured for 4 days. Cells were incubated for 30 min at 37°C with a dye-loading solution prepared in Tyrode's buffer (137 mM NaCl, 2.7 mM KCl, 1 mM $MgCl_2$, 1.8 mM $CaCl_2$, 0.2 mM Na_2HPO_4, 12 mM $NaHCO_3$, 5.5 mM D-glucose, pH 7.4) containing X-Rhod-1 (1:2000, Thermo Fisher Scientific), FluoVolt membrane potential dye (1:1000) with PowerLoad (1:100, Thermo Fisher Scientific), and Hoechst 33342 (10 µg/mL, Invitrogen). After two washes with Tyrode's buffer, cells were incubated in fresh buffer for 10 min prior to imaging. Temperature-controlled recordings were acquired over 30 s using a three-phase protocol: 10 s of spontaneous activity, 10 s of pacing at 1 Hz and 15 V (via MyoPacer stimulator, IonOptix), and 10 s of post-pacing recovery. Line-scan images were processed using a custom ImageJ macro, and quantification of calcium transient kinetics and membrane potential dynamics was performed using CyteSeer software (Vala Sciences).

Generation and functional assessment of engineered heart tissues (EHTs)
EHTs were generated as previously described[43], with minor modifications. Briefly, 1×10^6 hiPSC-CMs were used per construct. Three days after casting, cells were transduced with an adenovirus encoding channelrhodopsin-2 fused to GFP (Ad-ChR2-GFP, a gift from Vassilios J Bezzerides)[85]. EHTs were maintained in RPMI/B27(–insulin), supplemented with 0.1% insulin (Sigma-Aldrich), 5 µg/mL aprotinin (Sigma-Aldrich), and 1% penicillin–streptomycin (Gibco), with medium changes every other day. EHTs were cultured in 24-well plates and kept in a stage-top incubator at 37°C with 5% CO_2. For

functional assessment, spontaneous and paced contractions were recorded starting on day 7, and quantitative analyses were performed on day 28. Optical pacing was applied using blue LEDs positioned above the wells, and contractions were recorded from below at 30 frames per second through a 561-nm long-pass filter (Semrock BLP02-561R-32) using an 8-mm f/1.4 lens (Thorlabs MVL8M1) mounted on a Basler acA1920 camera. Post deflection was tracked using the Multi-Template Matching plugin in Fiji (ImageJ), and twitch force was calculated by beam-bending analysis based on the known Young's modulus of the posts, as described elsewhere[86].

DNA fragment analysis

DNA fragment analysis was performed using cDNA synthesized as described above and Phusion DNA Polymerase (Thermo Fisher Scientific) with 6-FAM–labeled primers (listed in the Oligonucleotides table). PCR products were purified and analyzed at the MGH CCIB DNA Core Facility, where fluorescently labeled DNA fragments were separated on an ABI 3730xl Genetic Analyzer (Applied Biosystems). Fragment sizes and relative abundances were determined using GeneMapper v6.0 software (Applied Biosystems).

Plasmid cloning

For overexpression experiments, the mApple–ACTN2 fusion construct was PCR-amplified from the pRRL-mApple–α-actinin-2 plasmid (a gift from Christopher S. Chen)[66]. The early isoform of *RBFOX2* containing the 40-bp C-terminal exon was amplified from the pSLIK-TT-RBFOX2 plasmid (Addgene). To generate the late *RBFOX2* isoform (containing the 43-bp exon), the truncated isoform (containing both the 40-bp and 43-bp exons), and a 3×FLAG control construct, oligonucleotide pairs encoding the respective sequences were synthesized by Integrated DNA Technologies (IDT). All constructs were PCR-amplified and cloned into the pLenti-CMV backbone (Addgene) using the NEBuilder HiFi DNA Assembly Kit (New England Biolabs). Plasmids were propagated in NEB competent *E. coli* according to the manufacturer's instructions.

Lentivirus production

Lentivirus was produced by transfecting HEK293T cells (ATCC CRL-3216) seeded in 150-mm dishes with 20 mL DMEM (Thermo Fisher Scientific) supplemented with 10% FBS (R&D Systems). Cells were transfected with 28 µg of lentiviral transfer vector, 21 µg psPAX2 (Addgene), and 7 µg pMD2.G (Addgene) using polyethyleneimine (PEI; Polysciences) in Opti-MEM (Thermo Fisher Scientific) at a PEI:DNA ratio of 3:1 (w:w). Medium was replaced 24 h post-transfection, and virus-containing supernatant was harvested at 72 h. Lentivirus was concentrated using Lenti-X Concentrator (Takara) according to the manufacturer's instructions. To determine viral titers, hiPSCs were transduced with serial dilutions of the concentrated virus and selected with 1 µg/mL puromycin (Boston BioProducts). Transduction efficiency was quantified by counting puromycin-resistant colony-forming units.

Lentivirus overexpression studies

Lentiviral overexpression constructs were transduced into hiPSCs at a multiplicity of infection (MOI) >1. Medium was replaced the following day. For rescue overexpression experiments, transduced hiPSCs were selected for single-cell clones, and successful

integration of the transgene was confirmed by Sanger sequencing. Verified clones were subsequently expanded and differentiated into hiPSC-CMs as described above for downstream functional analyses.

Zebrafish genotyping and mRNA microinjection

Germline transmission of CRISPR/Cas9-induced *rbfox1l* and *rbfox2* deletions was detected using fluorescent PCR and DNA fragment analysis as previously described[19]. Microinjection of mRNA into zebrafish embryos was performed as described previously[19]. Human *RBFOX2* early (containing the 40-bp C-terminal exon), late (containing the 43-bp C-terminal exon), and truncated (containing both the 40-bp and 43-bp C-terminal exons) isoforms, were generated from constructs described in the plasmid cloning section. Each isoform was subcloned into the pCS2+ vector. Plasmids were linearized with *SacII* and used as templates for in vitro transcription using the mMESSAGE mMACHINE SP6 Transcription Kit (Thermo Fisher Scientific). Approximately 1 nL of mRNA (~100 pg) was microinjected into zebrafish embryos at the one-cell stage.

Bulk RNA-seq library preparation and sequencing

Total RNA was isolated and purified as described above. Biological replicates included three control, five *RBFOX2* heterozygous, and five null samples, each comprising $1–2 \times 10^6$ hiPSC-CMs from independent differentiations. Poly(A)-selected RNA-seq libraries were prepared using a standard Illumina mRNA library preparation workflow. Sequencing was performed on an Illumina NovaSeq 6000 platform (2×150 bp paired-end, single index), yielding approximately 350 million paired-end reads per lane.

Differential gene expression analysis

Raw FASTQ files were first processed for quality control and aligned to the reference genome (GENCODE v38) using STAR (v2.7.9a)[68]. Gene-level read counts in STAR outputs were obtained for each sample and imported into DESeq2 (v1.32.0) for differential expression analysis[70]. Significance was defined by an adjusted p-value (False Discovery Rate, FDR) < 0.05 and an absolute fold change > 1.5. Transcripts per million (TPM) were also calculated using RSEM (v1.3.1) to provide normalized expression measures for visualization and cross-sample comparison[69]. Gene Ontology (GO) enrichment analysis was performed using clusterProfiler (v4.12.6)[71].

Raw FASTQ files were first processed for quality control and aligned to the reference genome (GENCODE v38) using STAR (v2.7.9a). Gene-level read counts in STAR outputs were obtained for each sample and imported into DESeq2 (v1.32.0) for differential expression analysis. Significance was defined by an adjusted p-value (False Discovery Rate, FDR) < 0.05 and an absolute fold change > 1.5. Transcripts per million (TPM) were also calculated using RSEM (v1.3.1) to provide normalized expression measures for visualization and cross-sample comparison. Gene Ontology (GO) enrichment analysis was performed using clusterProfiler (v4.12.6).

Alternative splicing analysis

Alternative splicing event (ASE) analysis of bulk RNA-seq data was performed using rMATS tools (v4.1.2)[72], applying an FDR < 0.05 to identify the significant events. rMATS

detects five categories of ASEs, including skipped exon (SE), alternative 5′ splice site (A5SS), alternative 3′ splice site (A3SS), mutually exclusive exons (MXE), and retained intron (RI). Functional enrichment analysis of ASE-associated genes was performed as described above for GO term enrichment.

Integration analysis of alternative splicing across datasets
For integrative analysis of ASE between *RBFOX2* het and control samples and during human iPSC-CM differentiation (day 7 vs. day 14), bulk RNA-seq data from a published study were reanalyzed[21]. The dataset included hiPSC-CM samples collected at day 7 and day 14 (n = 3 biological replicates per time point). The same ASE analysis pipeline described above was applied to identify differential splicing events. Overlapping ASEs across datasets were defined based on shared genomic coordinates of regulated exons and inclusion level changes, and functional enrichment of the overlapping genes was analyzed for enriched GO terms as described above.

eCLIP-seq library preparation
Large-scale cultures (~1.5 × 10⁷ cells) of wild-type hiPSC-CMs (day 15) were harvested from an Eppendorf DASbox bioreactor as previously described[87]. eCLIP was performed following established protocols[88,89]. Cells were resuspended in minimal ice-cold PBS and UV-crosslinked on ice (254 nm, 400 mJ cm⁻²) to covalently link RNA–protein complexes. Crosslinked cells were scraped, pelleted (200 × g, 3 min), washed with PBS, flash-frozen on dry ice, and stored at −80 °C. Immunoprecipitations were carried out with anti-RBFOX2 antibody (Bethyl Laboratories, A300-864A) or IgG control (Thermo Fisher Scientific, 02-6102). RNA fragments bound to RBFOX2 were ligated to adapters containing random barcodes (N5/N10), reverse-transcribed, and PCR-amplified to generate sequencing libraries. Libraries were sequenced on an Illumina NovaSeq 6000 platform (single-end 50 bp). Two biological replicates were sequenced per condition.

eCLIP-seq analysis
Sequencing reads were processed using the Skipper pipeline[73]. Adapters were trimmed with skewer[74], reads were aligned to the human genome (hg38) using STAR[68], and PCR duplicates were removed with UMICollapse[90]. Crosslinking sites were identified by counting 5′ read ends within fixed-size tiled windows across annotated genic regions. Windows were stratified by GC content to estimate and correct for GC bias. Enrichment of RBFOX2 binding was determined by comparing immunoprecipitated reads with their size-matched input (SMinput) controls. Motif analysis was performed on significant peaks ($P < 0.001$ and >8-fold change) using HOMER[76], as previously described[89]. Genome browser tracks were visualized in IGV[91], and pathway enrichment analysis was performed using Metascape[77].

Overlap analysis between RBFOX2 binding and alternative splicing events
RBFOX2 eCLIP-seq peaks were determined from processed data and converted to BED format. Genes containing RBFOX2-bound regions were compared with differentially spliced genes using *GeneOverlap* (https://bioconductor.org/packages/GeneOverlap/). Overlap between RBFOX2 binding peaks and alternatively spliced exons or splice junctions (±250 nt) was assessed using *BEDTools*[78]. Fisher's exact test was used to

evaluate enrichment, and overlaps with an odds ratio > 1.5 and P < 0.001 were considered significant.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed using GraphPad Prism 9. The number of biological replicates, statistical tests, and exact P values are provided in the corresponding figure legends. Data are presented as mean ± standard deviation (SD) or as median with interquartile range, as indicated. Comparisons between two groups were performed using unpaired two-tailed t tests. Comparisons among three or more groups were analyzed by one-way ANOVA followed by Tukey's multiple comparisons test, and when two independent variables were assessed, two-way ANOVA with Bonferroni's post hoc test was applied. Sample sizes were not predetermined using statistical methods. Randomization and investigator blinding were not applied unless otherwise specified due to the nature of the study.

REFERENCES

1. Liu, Y., Chen, S., Zühlke, L., Black, G.C., Choy, M., Li, N., and Keavney, B.D. (2019). Global birth prevalence of congenital heart defects 1970–2017: updated systematic review and meta-analysis of 260 studies. Int. J. Epidemiology *48*, 455–463. https://doi.org/10.1093/ije/dyz009.

2. Lopez, K.N., Morris, S.A., Tejtel, S.K.S., Espaillat, A., and Salemi, J.L. (2020). US Mortality Attributable to Congenital Heart Disease Across the Lifespan From 1999 Through 2017 Exposes Persistent Racial/Ethnic Disparities. Circulation *142*, 1132–1147. https://doi.org/10.1161/circulationaha.120.046822.

3. Fahed, A.C., Roberts, A.E., Mital, S., and Lakdawala, N.K. (2014). Heart Failure in Congenital Heart Disease A Confluence of Acquired and Congenital. Hear. Fail. Clin. *10*, 219–227. https://doi.org/10.1016/j.hfc.2013.09.017.

4. Lewis, M.J., Hsieh, A., Qiao, L., Tan, R., Kazzi, B., Channing, A., Griffin, E.L., Jobanputra, V., Su, J., Shahryar, C., et al. (2023). Association of Predicted Damaging De Novo Variants on Ventricular Function in Individuals With Congenital Heart Disease. Circ.: Genom. Precis. Med. *16*, e003900. https://doi.org/10.1161/circgen.122.003900.

5. Grossfeld, P., Nie, S., Lin, L., Wang, L., and Anderson, R.H. (2019). Hypoplastic Left Heart Syndrome: A New Paradigm for an Old Disease? J. Cardiovasc. Dev. Dis. *6*, 10. https://doi.org/10.3390/jcdd6010010.

6. Ohye, R.G., Sleeper, L.A., Mahony, L., Newburger, J.W., Pearson, G.D., Lu, M., Goldberg, C.S., Tabbutt, S., Frommelt, P.C., Ghanayem, N.S., et al. (2010). Comparison of Shunt Types in the Norwood Procedure for Single-Ventricle Lesions. N. Engl. J. Med. *362*, 1980–1992. https://doi.org/10.1056/nejmoa0912461.

7. Butts, R.J., Savage, A.J., Atz, A.M., Heal, E.M., Burnette, A.L., Kavarana, M.M., Bradley, S.M., and Chowdhury, S.M. (2015). Validation of a Simple Score to Determine Risk of Early Rejection After Pediatric Heart Transplantation. JACC: Hear. Fail. *3*, 670–676. https://doi.org/10.1016/j.jchf.2015.04.014.

8. Garcia, A.M., Beatty, J.-T., and Nakano, S.J. (2020). Heart failure in single right ventricle congenital heart disease: physiological and molecular considerations. Am. J. Physiol.-Hear. Circ. Physiol. *318*, H947–H965. https://doi.org/10.1152/ajpheart.00518.2019.

9. Rychik, J., Atz, A.M., Celermajer, D.S., Deal, B.J., Gatzoulis, M.A., Gewillig, M.H., Hsia, T.-Y., Hsu, D.T., Kovacs, A.H., McCrindle, B.W., et al. Evaluation and Management of the Child and Adult With Fontan Circulation: A Scientific Statement From the American Heart Association. Circulation *140*, e234–e284. https://doi.org/10.1161/cir.0000000000000696.

10. McCormick, A.D., and Schumacher, K.R. (2019). Transplantation of the failing Fontan. Transl. Pediatr. *8*, 290–301. https://doi.org/10.21037/tp.2019.06.03.

11. Gaynor, J.W., Mahle, M., Ittenbach, R.F., Kaplinski, M., Lawrence, K.M., Hunt, M.L., Burnham, N.B., Fuller, S., Chen, J.M., Maeda, K., et al. (2025). Long-Term Survival and Patient-Reported Outcomes After Staged Reconstructive Surgery for Hypoplastic Left Heart Syndrome. J. Am. Coll. Cardiol. *85*, 2386–2398. https://doi.org/10.1016/j.jacc.2025.04.028.

12. Driscoll, D.J., Offord, K.P., Feldt, R.H., Schaff, H.V., Puga, F.J., and Danielson, G.K. (2018). Five- to fifteen-year follow-up after Fontan operation. Circulation *85*, 469–496. https://doi.org/10.1161/01.cir.85.2.469.

13. Gentles, T.L., Gauvreau, K., Mayer, J.E., Fishberger, S.B., Burnetta, J., Colan, S.D., Newburger, J.W., and Wernovsky, G. (1997). Functional outcome after the Fontan operation: Factors influencing late morbidity. J. Thorac. Cardiovasc. Surg. *114*, 392–403. https://doi.org/10.1016/s0022-5223(97)70184-3.

14. Hinton, R.B., and Ware, S.M. (2017). Heart Failure in Pediatric Patients With Congenital Heart Disease. Circ. Res. *120*, 978–994. https://doi.org/10.1161/circresaha.116.308996.

15. Homsy, J., Zaidi, S., Shen, Y., Ware, J.S., Samocha, K.E., Karczewski, K.J., DePalma, S.R., McKean, D., Wakimoto, H., Gorham, J., et al. (2015). De novo mutations in congenital heart disease with neurodevelopmental and other congenital anomalies. Science *350*, 1262–1266. https://doi.org/10.1126/science.aac9396.

16. Sierant, M.C., Jin, S.C., Bilguvar, K., Morton, S.U., Dong, W., Jiang, W., Lu, Z., Li, B., López-Giráldez, F., Tikhonova, I., et al. (2025). Genomic analysis of 11,555 probands identifies 60 dominant congenital heart disease genes. Proc. Natl. Acad. Sci. *122*, e2420343122. https://doi.org/10.1073/pnas.2420343122.

17. Underwood, J.G., Boutz, P.L., Dougherty, J.D., Stoilov, P., and Black, D.L. (2005). Homologues of the Caenorhabditis elegans Fox-1 Protein Are Neuronal Splicing Regulators in Mammals. Mol. Cell. Biol. *25*, 10005–10016. https://doi.org/10.1128/mcb.25.22.10005-10016.2005.

18. Verma, S.K., Deshmukh, V., Thatcher, K., Belanger, K.K., Rhyner, A.M., Meng, S., Holcomb, R.J., Bressan, M., Martin, J.F., Cooke, J.P., et al. (2022). RBFOX2 is required for establishing RNA regulatory networks essential for heart development. Nucleic Acids Res. *50*, 2270–2286. https://doi.org/10.1093/nar/gkac055.

19. Huang, M., Akerberg, A.A., Zhang, X., Yoon, H., Joshi, S., Hallinan, C., Nguyen, C., Pu, W.T., Haigis, M.C., Burns, C.G., et al. (2022). Intrinsic myocardial defects underlie an Rbfox-deficient zebrafish model of hypoplastic left heart syndrome. Nat. Commun. *13*, 5877. https://doi.org/10.1038/s41467-022-32982-x.

20. Guo, Y., and Pu, W.T. (2020). Cardiomyocyte Maturation: New Phase in Development. Circ. Res. *126*, 1086–1106. https://doi.org/10.1161/circresaha.119.315862.

21. Lau, E., Han, Y., Williams, D.R., Thomas, C.T., Shrestha, R., Wu, J.C., and Lam, M.P.Y. (2019). Splice-Junction-Based Mapping of Alternative Isoforms in the Human Proteome. Cell Rep. *29*, 3751-3765.e5. https://doi.org/10.1016/j.celrep.2019.11.026.

22. Saha, K., Mei, Y., Reisterer, C.M., Pyzocha, N.K., Yang, J., Muffat, J., Davies, M.C., Alexander, M.R., Langer, R., Anderson, D.G., et al. (2011). Surface-engineered substrates for improved human pluripotent stem cell culture under fully defined conditions. Proc. Natl. Acad. Sci. *108*, 18714–18719. https://doi.org/10.1073/pnas.1114854108.

23. Lian, X., Zhang, J., Azarin, S.M., Zhu, K., Hazeltine, L.B., Bao, X., Hsiao, C., Kamp, T.J., and Palecek, S.P. (2013). Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/β-catenin signaling under fully defined conditions. Nat. Protoc. *8*, 162–175. https://doi.org/10.1038/nprot.2012.150.

24. Lian, X., Hsiao, C., Wilson, G., Zhu, K., Hazeltine, L.B., Azarin, S.M., Raval, K.K., Zhang, J., Kamp, T.J., and Palecek, S.P. (2012). Robust cardiomyocyte differentiation from human pluripotent stem cells via temporal modulation of canonical Wnt signaling. Proc. Natl. Acad. Sci. *109*, E1848–E1857. https://doi.org/10.1073/pnas.1200250109.

25. Feyen, D.A.M., McKeithan, W.L., Bruyneel, A.A.N., Spiering, S., Hörmann, L., Ulmer, B., Zhang, H., Briganti, F., Schweizer, M., Hegyi, B., et al. (2020). Metabolic Maturation Media Improve Physiological Function of Human iPSC-Derived Cardiomyocytes. Cell Rep. *32*, 107925–107925. https://doi.org/10.1016/j.celrep.2020.107925.

26. Miao, Y., Tian, L., Martin, M., Paige, S.L., Galdos, F.X., Li, J., Klein, A., Zhang, H., Ma, N., Wei, Y., et al. (2020). Intrinsic Endocardial Defects Contribute to Hypoplastic Left Heart Syndrome. Cell Stem Cell *27*, 574-589.e8. https://doi.org/10.1016/j.stem.2020.07.015.

27. Xu, X., Jin, K., Bais, A.S., Zhu, W., Yagi, H., Feinstein, T.N., Nguyen, P.K., Criscione, J.D., Liu, X., Beutner, G., et al. (2022). Uncompensated mitochondrial oxidative stress underlies heart failure in an iPSC-derived model of congenital heart disease. Cell Stem Cell *29*, 840-855.e7. https://doi.org/10.1016/j.stem.2022.03.003.

28. Ronaldson-Bouchard, K., Ma, S.P., Yeager, K., Chen, T., Song, L., Sirabella, D., Morikawa, K., Teles, D., Yazawa, M., and Vunjak-Novakovic, G. (2018). Advanced maturation of human cardiac tissue grown from pluripotent stem cells. Nature *556*, 239–243. https://doi.org/10.1038/s41586-018-0016-3.

29. Ren, J., Bi, Y., Sowers, J.R., Hetz, C., and Zhang, Y. (2021). Endoplasmic reticulum stress and unfolded protein response in cardiovascular diseases. Nat. Rev. Cardiol. *18*, 499–521. https://doi.org/10.1038/s41569-021-00511-w.

30. Minamino, T., Komuro, I., and Kitakaze, M. (2010). Endoplasmic Reticulum Stress As a Therapeutic Target in Cardiovascular Disease. Circ. Res. *107*, 1071–1082. https://doi.org/10.1161/circresaha.110.227819.

31. Moulik, M., Vatta, M., Witt, S.H., Arola, A.M., Murphy, R.T., McKenna, W.J., Boriek, A.M., Oka, K., Labeit, S., Bowles, N.E., et al. (2009). ANKRD1, the Gene Encoding Cardiac Ankyrin Repeat Protein, Is a Novel Dilated Cardiomyopathy Gene. J. Am. Coll. Cardiol. *54*, 325–333. https://doi.org/10.1016/j.jacc.2009.02.076.

32. Dorn, L.E., Petrosino, J.M., Wright, P., and Accornero, F. (2018). CTGF/CCN2 is an autocrine regulator of cardiac fibrosis. J. Mol. Cell. Cardiol. *121*, 205–211. https://doi.org/10.1016/j.yjmcc.2018.07.130.

33. Wu, X., Eder, P., Chang, B., and Molkentin, J.D. (2010). TRPC channels are necessary mediators of pathologic cardiac hypertrophy. Proc. Natl. Acad. Sci. *107*, 7000–7005. https://doi.org/10.1073/pnas.1001825107.

34. Heling, A., Zimmermann, R., Kostin, S., Maeno, Y., Hein, S., Devaux, B., Bauer, E., Klövekorn, W.-P., Schlepper, M., Schaper, W., et al. (2000). Increased Expression of Cytoskeletal, Linkage, and Extracellular Proteins in Failing Human Myocardium. Circ. Res. *86*, 846–853. https://doi.org/10.1161/01.res.86.8.846.

35. Krane, M., Dreßen, M., Santamaria, G., My, I., Schneider, C.M., Dorn, T., Laue, S., Mastantuono, E., Berutti, R., Rawat, H., et al. (2021). Sequential Defects in Cardiac Lineage Commitment and Maturation Cause Hypoplastic Left Heart Syndrome. Circulation *144*, 1409–1428. https://doi.org/10.1161/circulationaha.121.056198.

36. Lovci, M.T., Ghanem, D., Marr, H., Arnold, J., Gee, S., Parra, M., Liang, T.Y., Stark, T.J., Gehman, L.T., Hoon, S., et al. (2013). Rbfox proteins regulate alternative mRNA splicing through evolutionarily conserved RNA bridges. Nat. Struct. Mol. Biol. *20*, 1434–1442. https://doi.org/10.1038/nsmb.2699.

37. Tian, B., and Manley, J.L. (2017). Alternative polyadenylation of mRNA precursors. Nat. Rev. Mol. Cell Biol. *18*, 18–30. https://doi.org/10.1038/nrm.2016.116.

38. Mezzano, V., Pellman, J., and Sheikh, F. (2014). Cell Junctions in the Specialized Conduction System of the Heart. Cell Commun. Adhes. *21*, 149–159. https://doi.org/10.3109/15419061.2014.905928.

39. Lu, S.H.-A., Lee, K.-Z., Hsu, P.W.-C., Su, L.-Y., Yeh, Y.-C., Pan, C.-Y., and Tsai, S.-Y. (2022). Alternative Splicing Mediated by RNA-Binding Protein RBM24 Facilitates Cardiac Myofibrillogenesis in a Differentiation Stage-Specific Manner. Circ. Res. *130*, 112–129. https://doi.org/10.1161/circresaha.121.320080.

40. Kalsotra, A., Xiao, X., Ward, A.J., Castle, J.C., Johnson, J.M., Burge, C.B., and Cooper, T.A. (2008). A postnatal switch of CELF and MBNL proteins reprograms alternative splicing in the

developing heart. Proc. Natl. Acad. Sci. *105*, 20333–20338. https://doi.org/10.1073/pnas.0809045105.

41. Lundy, S.D., Zhu, W.-Z., Regnier, M., and Laflamme, M.A. (2013). Structural and Functional Maturation of Cardiomyocytes Derived from Human Pluripotent Stem Cells. Stem Cells Dev. *22*, 1991–2002. https://doi.org/10.1089/scd.2012.0490.

42. Saraswathibhatla, A., Indana, D., and Chaudhuri, O. (2023). Cell–extracellular matrix mechanotransduction in 3D. Nat. Rev. Mol. Cell Biol. *24*, 495–516. https://doi.org/10.1038/s41580-023-00583-1.

43. Breckwoldt, K., Letuffe-Brenière, D., Mannhardt, I., Schulze, T., Ulmer, B., Werner, T., Benzin, A., Klampe, B., Reinsch, M.C., Laufer, S., et al. (2017). Differentiation of cardiomyocytes and generation of human engineered heart tissue. Nat. Protoc. *12*, 1177–1197. https://doi.org/10.1038/nprot.2017.033.

44. Ying, Y., Wang, X.-J., Vuong, C.K., Lin, C.-H., Damianov, A., and Black, D.L. (2017). Splicing Activation by Rbfox Requires Self-Aggregation through Its Tyrosine-Rich Domain. Cell *170*, 312-323.e10. https://doi.org/10.1016/j.cell.2017.06.022.

45. Ribeiro, E. de A., Pinotsis, N., Ghisleni, A., Salmazo, A., Konarev, P.V., Kostan, J., Sjöblom, B., Schreiner, C., Polyansky, A.A., Gkougkoulia, E.A., et al. (2014). The Structure and Regulation of Human Muscle α-Actinin. Cell *159*, 1447–1460. https://doi.org/10.1016/j.cell.2014.10.056.

46. Wei, C., Qiu, J., Zhou, Y., Xue, Y., Hu, J., Ouyang, K., Banerjee, I., Zhang, C., Chen, B., Li, H., et al. (2015). Repression of the Central Splicing Regulator RBFox2 Is Functionally Linked to Pressure Overload-Induced Heart Failure. Cell Rep. *10*, 1521–1533. https://doi.org/10.1016/j.celrep.2015.02.013.

47. Paige, S.L., Galdos, F.X., Lee, S., Chin, E.T., Ranjbarvaziri, S., Feyen, D.A.M., Darsha, A.K., Xu, S., Ryan, J.A., Beck, A.L., et al. (2020). Patient-Specific Induced Pluripotent Stem Cells Implicate Intrinsic Impaired Contractility in Hypoplastic Left Heart Syndrome. Circulation *142*, 1605–1608. https://doi.org/10.1161/circulationaha.119.045317.

48. Yang, C., Xu, Y., Yu, M., Lee, D., Alharti, S., Hellen, N., Shaik, N.A., Banaganapalli, B., Mohamoud, H.S.A., Elango, R., et al. (2017). Induced pluripotent stem cell modelling of HLHS underlines the contribution of dysfunctional NOTCH signalling to impaired cardiogenesis. Hum. Mol. Genet. *26*, 3031–3045. https://doi.org/10.1093/hmg/ddx140.

49. Gao, G., Xie, A., Zhang, J., Herman, A.M., Jeong, E.-M., Gu, L., Liu, M., Yang, K.-C., Kamp, T.J., and Dudley, S.C. (2013). Unfolded Protein Response Regulates Cardiac Sodium Current in Systolic Human Heart Failure. Circ.: Arrhythmia Electrophysiol. *6*, 1018–1024. https://doi.org/10.1161/circep.113.000274.

50. Blackwood, E.A., MacDonnell, L.F., Thuerauf, D.J., Bilal, A.S., Murray, V.B., Jr, K.C.B., Margulies, K.B., and Glembotski, C.C. (2023). Noncanonical Form of ERAD Regulates Cardiac Hypertrophy. Circulation *147*, 66–82. https://doi.org/10.1161/circulationaha.122.061557.

51. Groenendyk, J., Sreenivasaiah, P.K., Kim, D.H., Agellon, L.B., and Michalak, M. (2010). Biology of Endoplasmic Reticulum Stress in the Heart. Circ. Res. *107*, 1185–1197. https://doi.org/10.1161/circresaha.110.227033.

52. Taegtmeyer, H., Sen, S., and Vela, D. (2010). Return to the fetal gene program: a suggested metabolic link to gene expression in the heart. Ann. N. York Acad. Sci. *1188*, 191–198. https://doi.org/10.1111/j.1749-6632.2009.05100.x.

53. D'Antonio, M., Nguyen, J.P., Arthur, T.D., Matsui, H., Donovan, M.K.R., D'Antonio-Chronowska, A., and Frazer, K.A. (2022). In heart failure reactivation of RNA-binding proteins is associated with the expression of 1,523 fetal-specific isoforms. PLoS Comput. Biol. *18*, e1009918. https://doi.org/10.1371/journal.pcbi.1009918.

54. Gu, M., Zhao, Y., Wang, H., Cheng, W., Liu, J., Ouyang, K., and Wei, C. (2023). Suppression of RBFox2 by Multiple MiRNAs in Pressure Overload-Induced Heart Failure. Int. J. Mol. Sci. *24*, 1283. https://doi.org/10.3390/ijms24021283.

55. Trachanas, K., Sideris, S., Aggeli, C., Poulidakis, E., Gatzoulis, K., Tousoulis, D., and Kallikazaros, I. (2014). Diabetic cardiomyopathy: from pathophysiology to treatment. Hell. J. Cardiol. : HJC Hell. kardiologike epitheorese *55*, 411–421.

56. Nutter, C.A., Jaworski, E.A., Verma, S.K., Deshmukh, V., Wang, Q., Botvinnik, O.B., Lozano, M.J., Abass, I.J., Ijaz, T., Brasier, A.R., et al. (2016). Dysregulation of RBFOX2 Is an Early Event in Cardiac Pathogenesis of Diabetes. Cell Rep. *15*, 2200–2213. https://doi.org/10.1016/j.celrep.2016.05.002.

57. Damianov, A., Ying, Y., Lin, C.-H., Lee, J.-A., Tran, D., Vashisht, A.A., Bahrami-Samani, E., Xing, Y., Martin, K.C., Wohlschlegel, J.A., et al. (2016). Rbfox Proteins Regulate Splicing as Part of a Large Multiprotein Complex LASR. Cell *165*, 606–619. https://doi.org/10.1016/j.cell.2016.03.040.

58. Nakahata, S., and Kawamoto, S. (2005). Tissue-dependent isoforms of mammalian Fox-1 homologs are associated with tissue-specific splicing activities. Nucleic Acids Res. *33*, 2078–2089. https://doi.org/10.1093/nar/gki338.

59. Groh, W.J., Groh, M.R., Saha, C., Kincaid, J.C., Simmons, Z., Ciafaloni, E., Pourmand, R., Otten, R.F., Bhakta, D., Nair, G.V., et al. (2008). Electrocardiographic Abnormalities and Sudden Death in Myotonic Dystrophy Type 1. N. Engl. J. Med. *358*, 2688–2697. https://doi.org/10.1056/nejmoa062800.

60. Lazarus, A., Varin, J., Babuty, D., Anselme, F. rédéric, Coste, J., and Duboc, D. (2002). Long-term follow-up of arrhythmias in patients with myotonic dystrophy treated by pacing A

multicenter diagnostic pacemaker study. J. Am. Coll. Cardiol. *40*, 1645–1652.
https://doi.org/10.1016/s0735-1097(02)02339-2.

61. Tokgozoglu, L.S., Ashizawa, T., Pacifico, A., Armstrong, R.M., Epstein, H.F., and Zoghbi, W.A. (1995). Cardiac Involvement in a Large Kindred With Myotonic Dystrophy: Quantitative Assessment and Relation to Size of CTG Repeat Expansion. JAMA *274*, 813–819. https://doi.org/10.1001/jama.1995.03530100053034.

62. Chopra, A., Kutys, M.L., Zhang, K., Polacheck, W.J., Sheng, C.C., Luu, R.J., Eyckmans, J., Hinson, J.T., Seidman, J.G., Seidman, C.E., et al. (2018). Force Generation via β-Cardiac Myosin, Titin, and α-Actinin Drives Cardiac Sarcomere Assembly from Cell-Matrix Adhesions. Dev. Cell *44*, 87-96.e5. https://doi.org/10.1016/j.devcel.2017.12.012.

63. Geach, T.J., Hirst, E.M.A., and Zimmerman, L.B. (2015). Contractile activity is required for Z-disc sarcomere maturation in vivo. genesis *53*, 299–307. https://doi.org/10.1002/dvg.22851.

64. Majkut, S., Idema, T., Swift, J., Krieger, C., Liu, A., and Discher, D.E. (2013). Heart-Specific Stiffening in Early Embryos Parallels Matrix and Myosin Expression to Optimize Beating. Curr. Biol. *23*, 2434–2439. https://doi.org/10.1016/j.cub.2013.10.057.

65. Weitkunat, M., Kaya-Çopur, A., Grill, S.W., and Schnorrer, F. (2014). Tension and Force-Resistant Attachment Are Essential for Myofibrillogenesis in Drosophila Flight Muscle. Curr. Biol. *24*, 705–716. https://doi.org/10.1016/j.cub.2014.02.032.

66. Zhang, K., Cloonan, P.E., Sundaram, S., Liu, F., Das, S.L., Ewoldt, J.K., Bays, J.L., Tomp, S., Toepfer, C.N., Marsiglia, J.D.C., et al. (2021). Plakophilin-2 truncating variants impair cardiac contractility by disrupting sarcomere stability and organization. Sci. Adv. *7*, eabh3995. https://doi.org/10.1126/sciadv.abh3995.

67. Sala, L., Meer, B.J. van, Tertoolen, L.G.J., Bakkers, J., Bellin, M., Davis, R.P., Denning, C., Dieben, M.A.E., Eschenhagen, T., Giacomelli, E., et al. (2017). MUSCLEMOTION: A Versatile Open Software Tool to Quantify Cardiomyocyte and Cardiac Muscle Contraction In Vitro and In Vivo. Circ. Res. *122*, e5–e16. https://doi.org/10.1161/circresaha.117.312067.

68. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2012). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

69. Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinform. *12*, 323. https://doi.org/10.1186/1471-2105-12-323.

70. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-014-0550-8.

71. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. OMICS: A J. Integr. Biol. *16*, 284–287. https://doi.org/10.1089/omi.2011.0118.

72. Shen, S., Park, J.W., Lu, Z., Lin, L., Henry, M.D., Wu, Y.N., Zhou, Q., and Xing, Y. (2014). rMATS: Robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. Proc. Natl. Acad. Sci. *111*, E5593–E5601. https://doi.org/10.1073/pnas.1419161111.

73. Boyle, E.A., Her, H.-L., Mueller, J.R., Naritomi, J.T., Nguyen, G.G., and Yeo, G.W. (2023). Skipper analysis of eCLIP datasets enables sensitive detection of constrained translation factor binding sites. Cell Genom. *3*, 100317. https://doi.org/10.1016/j.xgen.2023.100317.

74. Jiang, H., Lei, R., Ding, S.-W., and Zhu, S. (2014). Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. BMC Bioinform. *15*, 182. https://doi.org/10.1186/1471-2105-15-182.

75. Boyle, E.A., Her, H.-L., Mueller, J.R., Naritomi, J.T., Nguyen, G.G., and Yeo, G.W. (2023). Skipper analysis of eCLIP datasets enables sensitive detection of constrained translation factor binding sites. Cell Genom. *3*, 100317. https://doi.org/10.1016/j.xgen.2023.100317.

76. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Mol. Cell *38*, 576–589. https://doi.org/10.1016/j.molcel.2010.05.004.

77. Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C., and Chanda, S.K. (2019). Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat. Commun. *10*, 1523. https://doi.org/10.1038/s41467-019-09234-6.

78. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics *26*, 841–842. https://doi.org/10.1093/bioinformatics/btq033.

79. Burns, C.G., Milan, D.J., Grande, E.J., Rottbauer, W., MacRae, C.A., and Fishman, M.C. (2005). High-throughput assay for small molecules that modulate zebrafish embryonic heart rate. Nat. Chem. Biol. *1*, 263–264. https://doi.org/10.1038/nchembio732.

80. Wang, G., Yang, L., Grishin, D., Rios, X., Ye, L.Y., Hu, Y., Li, K., Zhang, D., Church, G.M., and Pu, W.T. (2017). Efficient, footprint-free human iPSC genome editing by consolidation of Cas9/CRISPR and piggyBac technologies. Nat. Protoc. *12*, 88–103. https://doi.org/10.1038/nprot.2016.152.

81. Akerberg, A.A., Trembley, M., Butty, V., Schwertner, A., Zhao, L., Beerens, M., Liu, X., Mahamdeh, M., Yuan, S., Boyer, L., et al. (2022). RBPMS2 Is a Myocardial-Enriched Splicing Regulator Required for Cardiac Function. Circ. Res. *131*, 980–1000. https://doi.org/10.1161/circresaha.122.321728.

82. Lian, X., Zhang, J., Azarin, S.M., Zhu, K., Hazeltine, L.B., Bao, X., Hsiao, C., Kamp, T.J., and Palecek, S.P. (2013). Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/β-catenin signaling under fully defined conditions. Nat. Protoc. *8*, 162–175. https://doi.org/10.1038/nprot.2012.150.

83. Prondzynski, M., Lemoine, M.D., Zech, A.T., Horváth, A., Mauro, V.D., Koivumäki, J.T., Kresin, N., Busch, J., Krause, T., Krämer, E., et al. (2019). Disease modeling of a mutation in α-actinin 2 guides clinical therapy in hypertrophic cardiomyopathy. EMBO Mol. Med. *11*, e11115. https://doi.org/10.15252/emmm.201911115.

84. Bortolin, R.H., Nawar, F., Park, C., Trembley, M.A., Prondzynski, M., Sweat, M.E., Wang, P., Chen, J., Lu, F., Liou, C., et al. (2024). Antisense Oligonucleotide Therapy for Calmodulinopathy. Circulation *150*, 1199–1210. https://doi.org/10.1161/circulationaha.123.068111.

85. Yoshinaga, D., Craven, I., Feng, R., Prondzynski, M., Shani, K., Tharani, Y., Mayourian, J., Joseph, M., Walker, D., Bortolin, R.H., et al. (2025). Dysregulation of N-terminal acetylation causes cardiac arrhythmia and cardiomyopathy. Nat. Commun. *16*, 3604. https://doi.org/10.1038/s41467-025-58539-2.

86. Vandenburgh, H., Shansky, J., Benesch-Lee, F., Barbata, V., Reid, J., Thorrez, L., Valentini, R., and Crawford, G. (2008). Drug-screening platform based on the contractility of tissue-engineered muscle. Muscle Nerve *37*, 438–447. https://doi.org/10.1002/mus.20931.

87. Prondzynski, M., Berkson, P., Trembley, M.A., Tharani, Y., Shani, K., Bortolin, R.H., Sweat, M.E., Mayourian, J., Yucel, D., Cordoves, A.M., et al. (2024). Efficient and reproducible generation of human iPSC-derived cardiomyocytes and cardiac organoids in stirred suspension systems. Nat. Commun. *15*, 5929. https://doi.org/10.1038/s41467-024-50224-0.

88. Nostrand, E.L.V., Pratt, G.A., Shishkin, A.A., Gelboin-Burkhart, C., Fang, M.Y., Sundararaman, B., Blue, S.M., Nguyen, T.B., Surka, C., Elkins, K., et al. (2016). Robust transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP). Nat. Methods *13*, 508–514. https://doi.org/10.1038/nmeth.3810.

89. Lorenz, D.A., Her, H.-L., Shen, K.A., Rothamel, K., Hutt, K.R., Nojadera, A.C., Bruns, S.C., Manakov, S.A., Yee, B.A., Chapman, K.B., et al. (2023). Multiplexed transcriptome discovery of RNA-binding protein binding sites by antibody-barcode eCLIP. Nat. Methods *20*, 65–69. https://doi.org/10.1038/s41592-022-01708-8.

90. Liu, D. (2019). Algorithms for efficiently collapsing reads with Unique Molecular Identifiers. PeerJ *7*, e8275. https://doi.org/10.7717/peerj.8275.

91. Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26. https://doi.org/10.1038/nbt.1754.

Figure 1. *RBFOX2* deficiency impairs cardiomyocyte growth and adhesion

(A) Schematic of predicted protein products from *RBFOX2* heterozygous (het) and homozygous null alleles. Black asterisks mark premature stop codons. RRM, RNA-recognition motif.

(B) Schematic of human iPSC-directed cardiomyocyte (CM) differentiation with annotated stages: Early differentiation (days 0-7); Late differentiation (days 8-15); Maturation (3D EHT culture, days 0-28). EHT, engineered heart tissue.

(C) qRT-PCR analysis of *RBFOX2* transcript levels in day 21 CTRL, het, and null iPSC-CMs (n = 5).

(D) Spontaneous beating frequency of day 15 CTRL and het iPSC-CMs in 2D culture (n = 3).

(E) Brightfield images of replated day 15 CTRL, het, and null iPSC-CMs after 1 day (n = 3). White arrowheads indicate flattened cells; black arrowheads indicate rounded cells. Scale bars, 50 μm.

(F-I) Quantification of cell size (F; n = 3-4; each dot represents the average area of 35-55 CMs per replicate), cell adhesion (G; n = 3), percent EdU+ CM (H; n = 3; each dot represents one image count), and percent TUNEL+ CM (I; n = 3; each dot represents one image count) in day 15 CTRL, het, and null iPSC-CMs.

Data are presented as mean ± SD (C, D, and F–I). *n* indicates biological replicates. Statistical analyses were performed using one-way ANOVA with Tukey's post hoc test (C, F–I) or unpaired two-tailed *t*-test (D). Significance levels: *p < 0.05; **p < 0.01; ***p < **0.001; *******p* < **0.0001; *ns*, not significant.**

Figure 2. RBFOX2 promotes cardiomyocyte differentiation

(A, B) Immunofluorescence staining of day 21 CTRL, het, and null iPSC-CMs for cardiac Troponin T (cTnT, magenta), α-actinin-2 (ACTN2, green), and DAPI (blue) (A), and quantification of myofibrillar disarray (B) (n = 3; each dot represents one CM). Scale bars, 10 μm.

(C, D) Transmission electron micrographs of day 24 CTRL, het, and null iPSC-CMs (C), and quantification of Z-disc heights (n = 3; each dot represents one sarcomere). Yellow arrows highlight the Z-disc (n = 3). Scale bars, 500 nm.

(E) qRT-PCR analysis of *MYH6*:*MYH7* transcript ratios in day 21 CTRL, het, and null iPSC-CMs (n = 9-10).

(F-H) Representative calcium transient profiles under 1 Hz pacing (F) and quantification of calcium transient duration at 75% repolarization (CTD75; G), and downstroke velocity (H) in day 21 CTRL, het, and null iPSC-CMs (n = 3).

(I, J) Seahorse analysis of mitochondrial function. Profile of real-time oxygen consumption rate (OCR) in day 21 CTRL, het, and null iPSC-CMs (I), with quantification of basal respiration,

ATP production, and maximal respiration (J) (n = 3). FCCP, carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone.

Data are presented as mean ± SD (B, D, E, and J) or as median with interquartile range (G and H). n indicates biological replicates. Statistics are unpaired two-tailed t test (B and D) or one-way ANOVA with Tukey's post hoc test (E, G, H and J). Significance levels: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$; *ns*, not significant.

Figure 3. *RBFOX2* deficiency disrupts transcriptional and splicing programs.

(A) Principal component analysis (PCA) of bulk RNA-seq data from CTRL (n = 3), het (n = 5), and null (n = 5) day 21 iPSC-CMs.

(B) Heatmap of normalized expression from bulk RNA-seq showing immature (fetal ventricle-enriched) and mature (adult ventricle-enriched) structural, conduction, and calcium-handling transcriptional markers across CTRL (n = 3), het (n = 5), and null (n = 5) day21 iPSC-CMs. Values represent row Z-scores of log2(TPM + 1). TPM, transcripts per million. Yellow highlights structural gene or markers, and green highlights conduction, and calcium-handling genes.

(C) Bubble plot of enriched GO terms associated with downregulated and upregulated differentially expressed genes (DEGs) in het iPSC-CMs compared with CTRL. BP, biological process. Yellow highlights upregulated, and green highlights downregulated GO terms.

(D) Hockey stick plot of upregulated DEGs (blue) in het iPSC-CMs compared with CTRL. Representative genes (red) are highlighted (FDR < 0.05; |log2 fold change| > 0.5).

(E) Heatmap of extracellular matrix (ECM)-related GO terms showing differential expression across ECM structural components (collagens, laminins, elastic fibers, proteoglycans, glycoproteins), adhesion signaling, ECM remodeling factors, ECM-coagulation pathways, and other regulators between CTRL and het iPSC-CMs (86 genes). Values represent row Z-scores of log2(TPM + 1). TPM, transcripts per million.

(F) Distribution of alternative splicing events (ASEs) identified in het versus CTRL iPSC-CMs (FDR < 0.05; |inclusion level difference| > 0.1).

(G) GO terms enriched among 1,211 ASEs detected in 823 genes (het vs CTRL). Yellow highlights CM differentiation-related pathways, and green highlights junction-related pathways.

(H) Distribution of RBFOX2 eCLIP peaks across genomic features with HOMER motif analysis showing significant enrichment of the consensus UGCAUG motif (two-sided enrichment, $p <$ 1e-237; log2 fold change > 3; n = 2).

(I) GO terms enriched among 2,966 RBFOX2 eCLIP peaks in 963 genes. Yellow highlights CM differentiation-related pathways, and green highlights junction-related pathways.

Figure 4. RBFOX2 shifts exon usage to a more mature profile in target transcripts encoding sarcomere, cytoskeletal, and junctional components.

(A) Gene Ontology (GO) terms enriched among overlapping RBFOX2 eCLIP targets and alternative splicing events (ASEs) from het versus CTRL iPSC-CMs. Yellow highlights CM differentiation-related pathways, and green highlights junction-related pathways. BP, biological process.

(B) Schematic model of *RBFOX2*-dependent regulation of sarcomere, cytoskeletal, junctional, and adhesion networks through eCLIP+ ASE+ targets that coordinate force generation and transmission. CM, cardiomyocyte; ECM, extracellular matrix.

(C) Sashimi plots of splicing regulation at *PDLIM5* (A3SS) and *ACTN2* (SE) loci in CTRL, het, and null day 21 iPSC-CMs. Inclusion levels (IncLev) are indicated. Representative RBFOX2 eCLIP binding peaks are shown (IP) compared to input control. Numbers on tracks indicate normalized read counts. A3SS, alternative 3′ splice site; SE, skipped exon; ex, exon; early, early isoform; early, late isoform.

(D) Heatmap of overlapping ASEs (het vs CTRL and day 7 vs day 14 CMs from a published dataset) across day 7 (n = 3), day 14 (n = 3), CTRL (n = 3), het (n = 5), and null (n = 5) day 21 iPSC-CMs. GO biological process terms enriched among ASE-associated genes are shown at right.

(E) Brightfield images of engineered heart tissues (EHTs) generated from CTRL, het, and null iPSC-CMs at day 28 under 1 Hz pacing (n = 3). Scale bars: 1 mm.

(F) Functional assessment of EHT contractile performance (under 1 Hz pacing, day 28), including measurements of contractile force (mN), contraction time to 50% peak (C50, sec), and relaxation time to 90% peak (R90, sec) (n = 3; each dot represents one EHT).

Data in are presented as mean ± SD (F). *n* indicates biological replicates. Statistical analyses were performed using unpaired two-tailed *t*-test or one-way ANOVA with Tukey's post hoc test (F). Significance levels: ****$p < 0.0001$; *ns*, not significant.

Figure 5. Primary defects in RBFOX2 autoregulation alters RBFOX2 isoform ratios in het CMs.

(A) Schematic of *RBFOX2* mutually exclusive exons (MEXs) at the C-terminal domain, showing the early (40 bp) and late (43 bp) exons that define early versus late isoforms.

(B) Sashimi plots of *RBFOX2* exon inclusion in CTRL and het iPSC-CMs at day 21, with RBFOX2 eCLIP binding peaks near the regulated exons. Inclusion level differences (IncLev) are indicated. Numbers on tracks indicate normalized read counts.

(C) DNA fragment analysis of isoform distribution of early, late, and aberrant 40+43 *RBFOX2* transcripts in CTRL and het cohorts during iPSC-CM differentiation (days 0-21; n = 3). White dots indicate the increased levels of aberrant 40+43 isoform.

(D) Quantification of the percentage of aberrant 40+43 *RBFOX2* isoform in CTRL and het iPSC-CMs during differentiation (days 0-21; n = 3).

(E) qRT-PCR analysis of developmental expression dynamics of *RBFOX2* transcripts in CTRL and het cohorts across days 0-30 of differentiation, normalized to day 0 CTRL iPSCs (n = 3).

Data are presented as mean ± SD (C-E). n indicates biological replicates. Statistical analyses were performed using two-way ANOVA with Bonferroni's post hoc test (D) or multiple unpaired *t*-tests (E). Significance levels: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$; *ns*, not significant.

Figure 6. Isoform-specific roles for RBFOX2 in CM differentiation and maturation

(A, B) Immunofluorescence staining of day 15 WT, het, and null iPSC-CMs transduced with 3×FLAG control (+FLAG), Early, or Late *RBFOX2* isoforms, labeled with cTnT (magenta), labeled with ACTN2 (green), and DAPI (blue) (A), and quantification of myofibrillar disarray (B) (n = 3; each dot represents one CM). Scale bars, 10 μm.

(C) Quantification of EHT contractile force (under 1 Hz pacing, day 28) in WT, het, and null iPSC-CMs expressing FLAG, Early, or Late isoforms (n = 3; each dot represents one EHT).

(D) Immunofluorescence staining of WT iPSC-CMs transduced with FLAG control or aberrant 40+43 isoform, labeled with cTnT (magenta), ACTN2 (green), and DAPI (blue) (n = 3). Scale bar, 10 μm.

(E) Quantification of EHT contractile force (under 1 Hz pacing, day 28) in WT iPSC-CMs transduced with FLAG control or aberrant 40+43 isoform (n = 3; each dot represents one EHT).

(F, G) Brightfield images of 72 hpf zebrafish embryos and *Tg(myl7:GFP)* fluorescent overlays of the heart in CTRL, DKO, and DKO embryos injected with Early, Late, and 40+43 *RBFOX2* isoform mRNA (F), and quantification of zebrafish ventricular fractional area change (FAC%, G). Scale bar, 100 μm.

Data are presented as mean ± SD (B, C, E and G). n indicates biological replicates. Statistical analyses were performed using one-way ANOVA with Tukey's post hoc test (B, C, and G) or unpaired two-tailed *t*-test (E). Significance levels: **$p < 0.001$; ****$p < 0.0001$; *ns*, not significant.

Figure 7. ACTN2 restores contractility in het CMs and triggers a mechanosensitive feedback loop that reactivates RBFOX2 expression and transcriptome maturation.

(A) qRT-PCR analysis of *ACTN2* transcript levels in day 21 CTRL, het, and null iPSC-CMs (n = 8).

(B, C) Immunofluorescence staining of day 15 WT, het, and null iPSC-CMs transduced with 3×FLAG control (+FLAG) or ACTN2 late isoform (+ACTN2), labeled with cTnT (magenta), ACTN2 (green), and DAPI (blue) (B), and quantification of myofibrillar disarray (C) (n = 3; each dot represents one CM). Scale bar, 10 μm.

(D) Quantification of EHT contractile force (under 1 Hz pacing, day 28) in WT, het, and null iPSC-CMs transduced with FLAG or ACTN2 late isoform (n = 3; each dot represents one EHT).

(E) qRT-PCR analysis of *RBFOX2* transcript levels in day 21 WT and het iPSC-CMs transduced with FLAG or ACTN2 late isoform (n = 8).

(F) DNA fragment analysis of isoform percentages of early, late, and aberrant 40+43 *RBFOX2* transcripts in day 21 WT and het iPSC-CMs transduced with FLAG or ACTN2 late isoform (n = 8).

(G, H) qRT-PCR analysis of *MYH6*:*MYH7* transcript ratio (G) and *MYL7*:*MYL2* transcript ratio (H) in day 21 WT and het iPSC-CMs transduced with FLAG or ACTN2 late isoform (n = 7-8).

(I, J) qRT-PCR analysis of alternative splicing events *PDLIM5* exon 5 (A3SS), and *TPM1* exon 6 (SE). Isoform expression levels were normalized to total transcript levels in day 21 WT and het iPSC-CMs transduced with FLAG or ACTN2 late isoform (n = 7-8). A3SS, alternative 3′ splice site; SE, skipped exon.

Data are presented as mean ± SD (A, and C-J). *n* indicates biological replicates. Statistical analyses were performed using one-way ANOVA with Tukey's post hoc test. Significance levels: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$; *ns*, not significant.