1

# Two-Step Mechanism of Bruton's Tyrosine Kinase Membrane Recruitment and Activation

Rachel A. McAllister[1,2,3], Amy L. Stiegler[4], Keerthana Chari[1], Meera Chari[1], Moitrayee Bhattacharyya[3*], Kallol Gupta[1,2*]

*Co-corresponding author: Kallol Gupta, Moitrayee Bhattacharyya,

1) Nanobiology Institute, Yale University, West Haven, CT, USA.
2) Department of Cell Biology, Yale University School of Medicine, New Haven, CT, USA.
3) Department of Pharmacology, Yale University School of Medicine, New Haven, CT, USA.
4) Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, USA.

**Email:** kallol.gupta@yale.edu, moitrayee.bhattacharyya@yale.edu

**Author Contributions:** R.A.M., K.G. and M.B. designed the research and experiments. K.C. generated mutant plasmids, purified protein and analyzed data. R. A. M. performed all native mass spectrometry and kinase assays. M.C. analyzed native mass spectrometry data and generated graphs. R.A.M. performed the fluorescence polarization experiments under the guidance of A.L.S. R.A.M. and K.G wrote the manuscript with contributions from M.B.

**This PDF file includes:**

Main Text
Figures 1 to 5

**Abstract**

Peripheral membrane proteins (PMPs) are critical mediators of signaling cascades initiated via activation of cell-surface receptors. Their functions rely on their innate ability to interact with membranes dynamically in response to rapidly changing cellular conditions. This membrane recruitment may occur via high-affinity interactions with specific lipids or transient, low-affinity membrane interactions. These weak and dynamic interactions that are critical regulators of protein function are challenging to capture. Taking Bruton's Tyrosine Kinase (BTK), a non-receptor tyrosine kinase essential for B cell activation, as an example, we demonstrate a native mass spectrometry (nMS) platform to understand the recruitment of PMPs by directly studying it from lipid bilayers customized to target membranes. Our data demonstrates that BTK recognizes phosphatidylserine (PS) through sites distinct from phosphatidylinositol (3,4,5) phosphate ($PIP_3$) binding. We show PS-bound BTK retains $PIP_3$ binding simultaneously via high-affinity sites, exhibiting $PIP_3$-independent basal membrane recruitment of BTK. Biochemical assays show that this PS-mediated recruitment sensitizes BTK to $PIP_3$-mediated activation under near-physiological concentrations of $PIP_3$. Thus, we propose a two-step model for BTK membrane recruitment and activation. A low-affinity interaction with high-copy number PS enables plasma membrane recruitment of BTK and increases its membrane-bound concentration. Upon B-cell activation, this pre-recruited, membrane-bound BTK population localizes to $PIP_3$-rich domains through electrostatic gliding along the membrane driven by low-affinity PS and high-affinity $PIP_3$ binding. This indicates a cooperative mechanism where PS can amplify B-cell signaling through increased membrane-bound BTK concentration. Our

44  work demonstrates a general model of PH-domain-containing protein regulation by weak protein-
45  lipid interactions, which can be extended to many other PMPs.

47
48  **Main Text**

49
**Introduction**
Signaling pathways within the cell that initiate from organellar membrane receptors often rely on
downstream peripheral membrane proteins to propagate the signals into cellular actions. This
includes the MAPK pathway that regulates cell growth, B-cell receptor pathway that regulates
immune response, and mitochondrial fission-fusion dynamics that regulate mitochondrial health (1-
3). A significant subset of peripheral membrane proteins conditionally associates with membranes
via specific recognition of signaling lipids such as phosphatidylinositol (3,4,5) phosphate ($PIP_3$) (4-
7). Among the most well-studied peripheral membrane protein domains is the Pleckstrin Homology
(PH) domain, the 11$^{th}$ most common domain in the human proteome (8). A key PH-domain
containing peripheral membrane protein that plays an essential role in B Cell Receptor (BCR)
signaling is Bruton's tyrosine kinase (BTK) (9, 10). BTK is a Tec family kinase whose PH domain
has an adjoining unique zinc finger motif required for structural stability (Fig. 1A) (11, 12). Upon
antigen stimulation, the BCR undergoes clustering, leading to SYK/LYN-mediated phosphorylation
of the immunoreceptor tyrosine-based activation motifs (ITAMs) on the associated
immunoglobulins CD79a and CD79b (13, 14). This process initiates the downstream activation of
Phosphatidylinositol (3) Kinase (PI3K), which phosphorylates membrane lipid phosphatidylinositol
(4,5)-bisphosphate ($PIP_2$) to form phosphatidylinositol (3,4,5)-trisphosphate ($PIP_3$) at the plasma
membrane (15, 16). Previous studies have shown that $PIP_3$ binding leads to activation of BTK,
leading to auto-phosphorylation on Tyr223 (12, 17). In the cytoplasm, BTK exists as an
autoinhibited monomer with the SH2 and SH3 domains compact against the distal face of the
kinase (Fig. 1A) (18, 19). On the membrane surface, $PIP_3$ binding is suggested to directly compete
with one autoinhibitory conformation of the PHTH domain in the full-length kinase and stabilize
dimeric interfaces (20). This ability to relieve autoinhibition and stabilize BTK interactions is critical
for BTK *trans*-autophosphorylation within the cell (18, 21). Additionally, SYK or Src-family kinases,
parallelly activated downstream of the BCR, phosphorylate Tyr551 in the kinase domain activation
loop of BTK, further increasing its catalytic activity (13, 22). Activated BTK phosphorylates and
activates phospholipase-Cγ2 (PLCγ2), which cleaves $PIP_2$ to produce diacylglycerol (DAG) and
soluble secondary messenger inositol triphosphate ($IP_3$) (23, 24). $IP_3$ binds to the $IP_3$-receptor in
the endoplasmic reticulum (ER), releasing calcium and activating Nuclear Factor of Activated T cell
(NFAT) and Nuclear Factor-kappa B (NF-κB) pathways, leading to internalization of the antigen
and activation of the B Cell (25-27). Due to the central role of BTK in adaptive immunity, mutations
within its canonical $PIP_3$ binding pocket have been shown to cause X-linked agammaglobulinemia
(XLA), a severe immunodeficiency syndrome (Fig. 1B) (28, 29). A key step in the BCR activation
is the plasma membrane recruitment and activation of BTK (12, 21, 30, 31). While other modes of
activation in solution, such as activation via the inositol hexakisphosphate ($IP_6$) through a novel
binding site have been recently uncovered (18), the current mechanistic understanding is that $PIP_3$
alone is responsible for both the recruitment of BTK to the plasma membrane and its activation in
the membrane. The central hypothesis of this study is that BTK can simultaneously interact with
other abundant plasma membrane inner leaflet lipids, in addition to $PIP_3$, using binding sites distinct
from those used for $PIP_3$ binding. We propose that these likely weaker interactions promote $PIP_3$-
independent recruitment of BTK to the plasma membrane in an inactive state, increasing its local
concentration and enabling BTK to scan along the quasi-two-dimensional inner leaflet of the plasma
membrane for $PIP_3$. Upon generation of $PIP_3$, this membrane pre-localized pool enables a stronger
activation response, leading to a synergistic mechanism by which orthogonal weak affinity lipid
interactions can amplify $PIP_3$-mediated BTK activation and subsequent signaling. Indeed, while PH
domains were initially thought to primarily to recognize PIPs, in recent years studies have shown a
broad range of functions, including work on the isolated PH domains of GRP1, ASAP1, and PDK1
showing background electrostatic lipids induced higher membrane residence (32-35). These

studies all suggest that a broader mechanism of coincidence detection of negatively charged
phospholipids on the target membrane may play a significant role in recruiting peripheral membrane
proteins for activation. Establishing this dual-mode lipid interaction demands experimental avenues
that offer a dynamic detection range and molecular resolution to simultaneously detect both tightly
and weakly bound lipids and unambiguously determine their identities and stoichiometry.
Furthermore, it demands technical abilities to perform these analyses on formed target protein lipid
complexes directly from a membrane environment that can be customized to a target physiological
membrane. We have recently developed a native mass spectrometry (nMS) approach that enables
the study of integral membrane proteins from customizable proteoliposomes, allowing for a
quantitative understanding of their hierarchical organization and lipid binding directly from lipid
bilayers mimicking different organellar membrane contexts (36, 37). In the current work, we further
optimize and extend this platform to study how specific lipids regulate the recruitment of peripheral
membrane proteins to organellar membranes, taking BTK as the target protein. A key advantage
of our nMS is its unmatched molecular resolution to detect and identify lipids that are specifically
bound to target proteins—both individually or together—along with their binding stoichiometries,
directly from complex lipid bilayers (36, 37). The general sensitivity of nMS enables further
detection of even a minor fraction of protein bound to a lipid, allowing for application to a wide and
dynamic range of protein lipid interactions (38-42). Applying this lipid vesicle nMS platform, we
discovered that BTK can get recruited to the membrane surface through a weak affinity interaction
with phosphatidylserine (PS), independent of $PIP_3$. We further established that this low-affinity
interaction with PS, a highly abundant lipid in the plasma membrane inner leaflet, takes place via
sites independent of the canonical $PIP_3$ binding site or the recently proposed peripheral site. Using
*in vitro* kinase assays, we further show that this $PIP_3$ independent PS mediated recruitment leads
to an increase in the degree of BTK autophosphorylation in the presence of physiologically relevant
amount of $PIP_3$. This builds a two-step recruitment and activation mechanism of BTK, where $PIP_3$
and PS work in a synergistic manner. Beyond BTK, the work also presents a methodological
platform to understand membrane recruitment of peripheral membrane proteins with precise
molecular detail.

**Figure 1. Structural features, XLA mutations, and PS binding of BTK: (a)** Domain architecture
of BTK showing the N-terminal PH and Tec Homology (TH) domain which contains a zinc finger
motif, followed by an Src module composed of: an Src-homology 3 (SH3) domain, Src-homology 2
(SH2) domain, and a C-terminal kinase domain. Yellow circles denote major tyrosine
phosphorylation sites Tyr223 and Tyr551. The structural model shows the autoinhibited state, in
which the PHTH domain primarily folds against the kinase domain but can adopt a range of
conformations (18, 20, 43). **(b)** Percentage distribution of unique X-linked agammaglobulinemia
(XLA)-associated missense mutations mapped to each domain within BTK via *BTKbase* web-
repository (29). **(c)** Native mass spectra of the isolated PHTH domain **(top)** and in the presence of
18:1-18:1 phosphatidylserine (DOPS) **(bottom)**. Charge states are labeled and PS-bound peaks
are marked. Observance of PS-bound peaks to all charge state peaks confirms the binding of PS
to BTK. Spectra are normalized to 10% of major species in each spectra and repeated in triplicate.
**(d)** Native mass spectra of the full-length BTK in the presence of DOPS. Minor purification
impurities due to the presence of Heat Shock Protein 70 (HSP70) are annotated with a red HSP70.
Like the isolated PHTH domain, the full-length BTK also binds to PS. The spectra were repeated
in triplicate.

**Results**

Our first goal was to determine if BTK can bind to other plasma membrane lipids beyond PIP$_3$. To address this, at first, we utilized just the membrane binding PHTH domain of BTK and studied independent lipid binding in solution using nMS. By directly incubating the PHTH domain with different lipids independently and subjecting it to nMS, we found that the PHTH domain is capable of binding PS (Fig. 1C), a negatively charged phospholipid that is highly abundant on the plasma membrane. PS is primarily localized to the inner leaflet, accounting for ~1/5$^{th}$ of inner leaflet lipids (44-48). In contrast, no binding was detected for phosphatidylethanolamine (PE), another prominent plasma membrane lipid enriched on the inner leaflet, indicating the specificity of the protein for the anionic serine head group over the more zwitterionic PE head group (Fig. S1A-D) (44, 48). In the full-length autoinhibited conformation of the protein, the PH domain is sandwiched against the N-lobe of the kinase, yet explores a range of conformations, leading to minor heterogeneity in the auto-inhibited state (Fig. 1A) (18-20). Due to this potential occlusion of PHTH domain surfaces, we wanted to confirm that full-length BTK can also recognize PS. As shown in Figure 1D, full-length BTK also binds to PS, indicating that binding to PS is not abrogated by the PHTH domain conformation in the full-length protein. Finally, we aimed to demonstrate that this PS-BTK interaction can also occur in the presence of a bilayer, within the lipid-rich milieu that comprises the collective membrane environment. This is a critical consideration when studying protein-lipid interactions. While a target protein may bind a specific lipid in isolation, such interactions may not persist in the context of a complex physiologically relevant lipid bilayer(49). In these environments, other abundant lipids could potentially compete for binding, disrupt, or modulate the interaction (36, 37). Therefore, assessing lipid binding within native or native-like lipid mixtures is essential for accurately capturing biologically meaningful interactions.

To address this, we incorporated our recently published nMS method that enables the study of integral membrane protein-lipid interactions directly from customizable liposomes (36, 37). Here we adapted this approach to peripheral membrane proteins, where gentle ionization and in-source activation disrupts the vesicles enough to resolve direct protein-lipid complexes (Fig. 2A). First, from liposomes containing phosphatidylcholine (PC), PE and a minor amount PS – we observe a robust and exclusive PS binding peak (Fig. 2B). Next, we customized the lipid composition of the liposome to more closely resemble the lipid composition of the inner leaflet of the plasma membrane, the leaflet encountered by cytosolic peripheral membrane proteins. Typically, the inner leaflet is expected to be devoid of cholesterol and sphingomyelin, which primarily reside on the outer leaflet, and high PS (47, 48). As shown in Figure 2C, even in the plasma membrane inner leaflet-like liposome containing 7 different phospholipid species, PS binding is observed as the primary adduct. The ability to detect robust PS binding to BTK incubated with these custom lipid bilayers indicates the ability of PS in an inner leaflet-like bilayer to recruit BTK to the membrane surface.

To corroborate the nMS-based observations, we performed fluorescence polarization (FP) assays in solution. Fluorescent nitrobenzodiazole (NBD) – labeled PS, modified on the fatty acyl chain to preserve headgroup sterics, was kept at a fixed concentration while the PHTH domain concentration was varied. Excitation at 485 nm of polarized light and emission detection at 528 nm were used to monitor NBD fluorescence. In the absence of binding, the small, fast tumbling PS exhibits low polarization as the emitted light is largely depolarized due to rapid Brownian motions during the NBD excitation lifetime (50). Upon addition and binding to the bulkier PHTH domain the rotational speed is decreased and less light is depolarized. As shown (SI. Fig. 2A), this increases the polarization read-out, providing direct solution-phase evidence of BTK-PS binding. We further observe a PHTH-concentration-dependent increase in polarization, consistent with binding of the PHTH domain to PS (SI. Fig. 2B).

**Figure 2. Lipid binding to BTK directly from the plasma membrane inner leaflet-like**
**liposome. (a)** Flow chart of peripheral membrane protein proteo-liposome preparation. Lipids in
chloroform are dried under a nitrogen flow for 30 minutes, following a minimum of two hours under
vacuum, to remove residual organic solvent. Aqueous nMS buffer is added to the dried lipid film for
vortex rehydration prior to ten cycles of alternating liquid nitrogen and a 50 °C water bath, followed
by extrusion with 100 nm pore membranes using an Avestin Liposofast basic. The homogenous
vesicles are then incubated with protein prior to nMS, where the protein can be ejected from these
vesicles with bound lipid still attached. **(b)** Mass Spectra of full-length BTK in the presence of
PC/PE/PS vesicles. Lipid vesicle composition is shown on the right. As annotated, direct and
exclusive binding of PS to BTK could be observed from the liposome. Spectra were normalized to
10% of the apo peak and repeated in triplicate. **(c)** Mass Spectra of full-length BTK in the presence
of inner-leaflet plasma membrane-like vesicles. Lipid vesicle composition shown on the right.
Exclusive binding to PS to BTK was still observed, despite the presence of other inner-leaflet
anionic lipids at physiological ratios. Spectra were normalized to 10% of the apo peak and repeated
in triplicate. In both spectra, the * denotes a minor impurity of 241 Da addition that corresponds to
a covalent phosphogluconylation of BTK. A noted and established artifact of overexpression of
membrane-binding domains/proteins in E. coli (7).

Next, we asked whether the PS binding to BTK can occur independently of $PIP_3$ binding,
as both PS and $PIP_3$ are negatively charged lipids that may recognize similar positively charged
residues on proteins. To test this, we employed dual charge reversal mutations at the canonical
$PIP_3$ binding site formed by Lys 12, Arg 28, Asp 24 and Tyr 39 and performed nMS of this
R28C/N24D mutant BTK, which has been shown to abolish the $PIP_3$ binding to BTK (18, 28, 30).
nMS-based lipid binding studies on this mutant reveal that the mutant is still capable of binding to
PS (SI. Fig. 3A-C). We next tested a recently discovered secondary binding site for Inositol
Hexakisphosphate ($IP_6$), coordinated through Arg 52, Lys 36, Lys 49 and Tyr 40 (18).  nMS of
K49S/R52S mutant PHTH also shows no significant drop in the PS binding ability over the wild type
protein (SI. Fig. 4A-C). Together, these data indicate that BTK can bind to PS independent of the
specific $PIP_3$ binding sites.
To further establish the independent binding mechanism, we performed a series of dual
binding experiments involving full-length BTK, $PIP_3$, and PS. First, to unambiguously establish that
such dual binding can occur directly from a membrane environment, we performed nMS of BTK in
the presence of PS-containing liposomes, where we added $IP_4$, the functionally active headgroup
of $PIP_3$ (SI. Fig. 5). Prior work has established that $IP_4$ adopts the same binding pose as $PIP_3$ at the
canonical site (51). Additionally, at a steady concentration of BTK and PS, we increased $IP_4$
concentration, and the resultant solution was subjected to nMS analysis. The apo and lipid/$IP_4$ BTK
peaks were quantified via the area under the curve for the predominant charge state. As shown in
Figure 3A, and quantified in Figure 3B-C, the addition of $IP_4$ leads to a clear observation of $IP_4$-
bound BTK and critically, no significant loss in PS binding was observed. More importantly, as the
concentration of $IP_4$ in solution increases, a distinct peak at m/z 4574 arises, corresponding to the
mass of BTK bound simultaneously to $IP_4$ and PS. This indicates that PS-bound BTK can
simultaneously engage and bind the $PIP_3$ head-group. Finally, we confirmed the dual lipid-binding
capability of BTK by incubating PHTH domain with both PS and $PI(3,4,5)P_3$. nMS of this mixture
identified a characteristic peak that corresponds to the simultaneous $PIP_3$+PS bound state (Fig.
3D). This peak was absent in controls with BTK incubated with only PS or $PIP_3$ (SI Fig. 6). Together,
these data unambiguously establish that BTK can simultaneously bind PS while engaging with $PIP_3$
through the canonical binding site. Interestingly, the plasma membrane is composed of
approximately 10% PS, with the majority localized to the inner leaflet, constituting about 15-20
mol% of inner leaflet lipids (44, 46). For cytosolic peripheral membrane proteins like BTK, which
exclusively interact with the inner leaflet, this lower-affinity interaction with an abundant lipid like
PS can facilitate dynamic membrane exploration. Specifically, this PS-dependent interaction can
position BTK favorably on the membrane surface, allowing its PH domain to efficiently search for
and engage high-affinity activating ligands, such as $PIP_3$. Importantly, this PS-mediated basal
membrane association should also inherently elevate the local BTK concentration at the plasma
membrane independently of $PIP_3$ availability.

Figure 3. Ability of BTK to bind both PS and PIP3/IP4 simultaneously. (a) Expansion of charge
state 17 of nMS of full-length BTK after incubation with DOPS. Top to bottom: Increasing amounts
of inositol tetrakisphosphate (IP4) are added. BTK is kept at 2.5 μM, PS at 125 μM, and IP4
concentration (top to bottom: 0 μM, 0.625 μM, 1.25 μM, and 2.5 μM). Peak abundance is
normalized to 50% of the charge state 17 peak. As highlighted in a light fuchsia, all conditions
produce a PS-bound peak. Incubation with IP4 yields an IP4-bound peak highlighted in purple.
Additionally, increasing IP4 concentration yields a peak, highlighted in green, that corresponds to
the BTK+PS+IP4 species. (b) Plot of the peak area of the PS and IP4-bound peaks obtained from
(a), highlighted in red and purple, respectively. The area under the curve (AUC) was calculated for
each as a percentage of the apo charge state peak (mean ± sem, n = 3). Statistics calculated using
an unpaired, nonparametric t-test on Graphpad Prism. Statistical significance defined as $p > 0.05$,
ns; $p < 0.05$ as *, $p < 0.01$ as ** and $p < 0.001$ as ***. (c) Plot of the peak area of the peak at m/z =
4574 corresponding to dual IP4 and PS-bound BTK (highlighted in light green) (mean ± sem, n =3).
Statistics calculated using an unpaired, nonparametric t-test on Graphpad Prism. (d) Spectra of the
wild-type PHTH domain upon incubation with both 8:0 PIP3 and PS reveal a unique peak at m/z =
2445, which corresponds to the dual-bound species.

265   Given that BTK activation critically involves trans-autophosphorylation, this PS-mediated
266 increase in local BTK concentration on the plasma membrane is likely to facilitate
267 autophosphorylation and enhance BTK activation, particularly under conditions of limited $PIP_3$
268 availability. To assess the influence of PS-dependent membrane recruitment on BTK activation, we
269 utilized an *in vitro* BTK autophosphorylation assay (18). We set up BTK phosphorylation assays in
270 the presence of liposomes containing various concentrations of PS and $PIP_3$. Briefly, BTK was
271 incubated with custom liposomes before the introduction of ATP and $MgCl^{2+}$ to initiate reaction.
272 The reaction was stopped after 5 minutes by adding SDS buffer + EGTA and boiling the sample
273 (Fig. 4A). The extent of BTK autophosphorylation was measured and quantified by western blots
274 using BTK phosphoTyr223 antibodies and normalized with respect to total BTK concentration.
275 Previous work has established that $PIP_3$ is necessary and sufficient for activation of trans-
276 autophosphorylation of BTK on the membrane (28, 52). Aligning with this observation, PS and other
277 phosphatidylinositol-containing liposomes (PIP or $PIP_2$) failed to activate BTK by themselves in the
278 absence of $PIP_3$ (SI Fig. 7), affirming that $PIP_3$ is necessary for BTK activation. This aligns well with
279 previous mechanistic work that demonstrates how stable membrane binding of $PIP_3$ facilitates BTK
280 autophosphorylation (12, 30). Next, BTK autophosphorylation assays conducted with liposomes
281 containing high levels (5%) of $PIP_3$ showed autophosphorylation, although no additional influence
282 from increasing PS concentrations in the liposomes could be observed (Fig. 4b). However, under
283 10-fold lower $PIP_3$ concentration (1-0.5%), incremental increases in PS concentration led to a near-
284 linear enhancement of BTK autophosphorylation (Fig. 4C-D). Notably, the magnitude of this PS-
285 mediated amplification of autophosphorylation inversely correlated with $PIP_3$ abundance –
286 reduction in $PIP_3$ concentration heightened the impact of PS in the liposomes. Strikingly, at 0.5%
287 $PIP_3$, we note 2.77-fold increase in kinase autophosphorylation going from 0% PS to 20% PS (Fig.
288 4D). Based on our experimental data, we constructed a simple mathematical model to understand
289 how PS-mediated membrane recruitment can increase the rate of BTK autophosphorylation (See
290 Methods for details). For this, we leveraged a previously established framework and further
291 modified it to approximate how the effective concentration of BTK on the liposome surface would
292 change as a result of PS at low $PIP_3$ concentrations (18). This simple and approximate model
293 predicts approximately 1.97-fold increase in the autophosphorylation rate at 0.5% $PIP_3$ going from
294 0 to 20% PS (SI Fig. 8). This closely agrees with experimentally derived change in BTK
295 autophosphorylation, further establishing that increase in *trans*-autophosphorylation is a direct
296 representation of an increase in effective concentration in the presence of the liposomes. Together,
297 these results are in agreement with our nMS-based observations and a two-step model that low-
298 affinity interactions with abundant PS enhance local BTK concentrations at the membrane,
299 consequently promoting the ease and extent of BTK autophosphorylation under lower
300 concentrations of $PIP_3$.

302
**Figure 4. Influence of PS on the trans-*auto*phosphorylation of BTK. (a)** Schematic depicting
the kinase assay. Briefly, liposomes are created as in Fig 2A. After a 10-minute pre-incubation with
BTK, kinase reaction buffer is added and reaction proceeds for 5 minutes. The reaction was
stopped by the addition of Laemli w/ EGTA and boiling. For more details, see methods. All
conditions normalized to the mean of 0% PS for their respective $PIP_3$ concentrations and
representative westerns generated as a merge between colorimetric (ladder) and
chemiluminescence. **(b) (top)** Representative Coomassie stain of 5% $PIP_3$ liposomes to
demonstrate total Btk concentration in both the negative control in the absence of ATP and $MgCl_2$
as well as in the reaction conditions. **(middle)** Representative anti-phosphoTyr223 BTK western.
**(bottom)** Plot of the quantified and normalized phosphoTyr223/Total BTK ratio (mean ± sem, n
=4). No change in extent of phosphorylation observed despite changing PS concentrations.
Statistical significance defined as $p > 0.05$, ns; $p < 0.05$ as *, $p < 0.01$ as ** and $p < 0.001$ as ***. **(c)**
**(top)** Representative Coomassie stain of 1% $PIP_3$ liposomes to demonstrate total Btk concentration
in both the negative control in the absence of ATP and $MgCl_2$ as well as in the reaction conditions.
**(middle)** Representative anti-phosphoTyr223 BTK western. **(bottom)** Plot of the quantified and
normalized phosphoTyr223/Total BTK ratio (mean ± sem, n =3). Increases in relative
phosphorylation from 0%PS to 10% PS or 20% demonstrate an increase in BTK activation in the
background presence of anionic PS. **(d) (top)** Representative Coomassie stain of 0.5% $PIP_3$
liposomes to demonstrate total Btk concentration in both the negative control in the absence of
ATP and $MgCl_2$ as well as in the reaction conditions. **(middle)** Representative anti-phosphoTyr223
BTK western. **(bottom)** Plot of the quantified phosphoTyr223/Total BTK ratio (mean ± sem, n =4).
A 0.5% $PIP_3$ concentration, a stark increase in autophosphorylation level is observed upon
increasing the PS concentration, demonstrating a near linear increase in BTK activity with regards
to increase PS concentration.

**Discussion**

**Discussion**
Our findings reveal a novel interaction between BTK and PS, an abundant negatively
charged lipid predominantly localized on the inner leaflet of the plasma membrane. Importantly,
this interaction occurs at sites independent from the canonical $PIP_3$-binding pocket of BTK. This
allows BTK to simultaneously engage with PS and dynamically explore the membrane surface for
high-affinity interactions with $PIP_3$. We confirmed this through mutational analyses that selectively
disrupted $PIP_3$ binding without impairing PS interaction. Given the very low physiological
abundance of $PIP_3$ (<1 mol %) and its short life span, the observed low-affinity and high-copy
number PS interactions can substantially elevate local BTK concentration and facilitate rapid,
dynamic membrane scanning by the PHTH domain for $PIP_3$. The biological significance of this
electrostatic interaction is highlighted by our kinase assays, where it enhances membrane-localized
trans-autophosphorylation, thereby lowering the threshold of $PIP_3$ necessary to achieve robust BTK
activation in response to upstream B cell receptor (BCR) signaling events. Our liposome-based
kinase assay showed a 2.77-fold increase in the extent of BTK autophosphorylation. Further
mathematical modeling of the data corroborates with this change. Conceivably, at higher $PIP_3$
levels where BTK can be fully saturated on the high-affinity $PIP_3$, this effect of PS binding is
expected to plateau out. This is also supported in our kinase assay, where no effect of PS
concentration was observed at 5% PIP3 containing liposomes. At higher $PIP_3$ concentrations, the
excess of $PIP_3$, making up to one in 20 lipids, begins to occupy all BTK, leaving less free BTK to
recognize PS to recruit it on the surface, explaining how the difference in $PIP_3$ concentration alters
PS sensitivity.
An unresolved aspect of this study is the precise delineation of the binding sites for PS on
BTK. Despite extensive mutagenesis efforts, we were unable to localize specific residues whose
mutation disrupted PS binding. This finding raises the possibility that PS binding might not involve
distinct, well-defined sites but rather general electrostatic surface interactions. Surface charge
analysis further supports this hypothesis, demonstrating a large electropositive surface surrounding
the canonical $PIP_3$ site (Fig. 5A). Such an extensive electropositive surface is likely to facilitate a
generalized electrostatic mechanism, enabling PS-associated membrane bound BTK to remain
conformationally flexible, while gliding along the plasma membrane to present its canonical binding
site optimally for high-affinity $PIP_3$ interactions. This also guides the PHTH domain to have the
electrostatically positive surface surrounding the canonical $PIP_3$ site facing towards the membrane
surface (Fig. 5B). This orientation can facilitate faster search of the negatively charged inner leaflet
and have the PHTH domain already orientated with the area around the canonical site towards the
lipid head groups (Fig. 5B). An electrostatic gliding mechanism might allow BTK to rapidly scan the
PS-rich inner leaflet of the plasma membrane for $PIP_3$. Indeed, previous computational and FRET-
based studies on isolated PH domains of GRP1, a guanine nucleotide exchange factor, has
proposed electrostatic hopping that can facilitate $PIP_3$ binding (33). Notably, the nMS-based
workflow provides the analytical precision required to unravel such nuanced multi-modal
interactions directly from lipid membranes, as exemplified by our successful demonstration of dual
$PIP_3$ and PS binding to BTK. The autophosphorylation assay further illustrates how these weak-
affinity electrostatic interactions significantly reduce the threshold concentration of $PIP_3$ required to
achieve physiologically relevant kinase activation upon BCR engagement. Given that the PH
domain is one of the most widespread membrane-binding domains in the human proteome, found
in over 250 distinct proteins, understanding how cells rapidly recruit PH-domains proteins using
short -lived low abundant lipid such as $PIP_3$ presents an exciting mechanistic question (53). Our
work establishes a broadly applicable and robust analytical framework for investigating multi-modal
peripheral membrane protein-lipid interactions. The nMS approach described herein, using a highly
clinically-relevant peripheral membrane protein, offers unprecedented molecular resolution,
enabling the identification of novel protein-lipid interactions, and provides a powerful tool for
understanding the fundamental principles governing the selective recruitment of cytosolic effectors
to various cellular membranes.

Figure 5. (a) (top left) Electrostatic map of the PHTH domain of BTK (PDB ID: 4Y94) (18). Electrostatics calculated on a single PHTH monomer using Pymol APBS electrostatics. Canonical site highlighted in fuschia circle (top right) Same plot orientated 180° around the x axis showing lack of electropositive character on this face. (bottom left) Plot from top left oriented 90° around the y axis, with the canonical PIP$_3$ binding pocket highlighted in a fuschia circle. Large electrostatic positive character and pockets surround the canonical site. (bottom right) Plot from bottom left oriented 90° around the x axis highlighting another large electrostatic positive surface centered around the peripheral PIP$_3$ binding site. Canonical site highlighted at the edge of the structure with a fuschia arrow. Though the sites are two hotspots for binding PIP$_3$, many electropositive lysines and arginines are exposed on proximal loops to these sites as seen in bottom left and bottom right though have not been directly implicated in coordinating PIP$_3$ through the two defined sites. (b) Electrostatic map shown in (a) and model of membrane search mechanism, showing orientation of electropositive character surrounding the PIP$_3$ binding site gliding along membrane, orienting PHTH domain towards the anionic lipids. (c) Schematic depicting the proposed two-step binding of BTK on the inner leaflet of the plasma membrane. First, due to a low-affinity interaction with the highly abundant PS, BTK is recruited to the surface of the plasma membrane inner leaflet to dynamically explore for a high affinity activating ligand. Upon BCR stimulation, this initiates a cascade that activates two critical upstream proteins: PI3K and Src-family kinases. PI3K converts lipid phosphatidyl inositol (4,5) phosphate (PIP$_2$) into phosphatidyl inositol (3,4,5) phosphate (PIP$_3$). BTK is then activated by this high-affinity lipid and trans-autophosphorylates at Tyrosine 223 in the SH3 domain. Due to priming of BTK in the proximity of the membrane due to its recognition of PS, this activation can take place on a much faster scale. Src-family kinases, including LYN and FYN can phosphorylate Tyrosine 551 in the activation loop of BTK, increasing its catalytic activity. This fully active BTK can phosphorylate phospholipase C γ2, leading to cleavage of PIP$_2$ to diacylglycerol and IP$_3$, a secondary messenger leading to downstream BCR activation.

**Materials and Methods**

Purifications

Plasmids were transformed into *Escherichia Coli* BL21 (DE3) cells for protein expression. For full-length BTK, *E. Coli* cells also containing the expression constructs for tyrosine phosphatase (YopH) and GroEL/ES were generated and used. *Bos Taurus* BTK was used, as it retains > 98% sequence homology and retains identical lipid binding sites and can be expressed in high quantities in *E. Coli*. Cells were grown in Terrific Broth (TB) with 70 μg/ml Kanamycin (with an additional 50 ug/ml Spectinomycin and 50 ug/ml Chloramphenicol for full-length BTK) to an optical density of 1.6 at 37° C, at which point an equivalent amount of 4° C TB with 70 μg/ml Kanamycin (with an additional 50 ug/ml Spectinomycin and 50 ug/ml Chloramphenicol for full-length TB), 100 μM ZnCl2, and 1 mM IPTG was added to induce a 16-hour expression at 18 C. Cells were harvested at 4,000 x g for 30 min and cell pellets resuspended in Ni-NTA buffer A (25 mM Tris pH 8.5, 500 mM NaCl, 5% glycerol and 20 mM imidazole) with protease inhibitor cocktail and DNAse. Cells were lysed via sonication (40% amplitude, 20 sec on, 45 sec off, 5 min total) and debris centrifuged out at 200,000 x g for 45 minutes at 4° C. The supernatant was loaded onto an HisTrap 5 mL column using a peristaltic pump at 1 mL/min and washed with 10 Column volume (CV) Ni-NTA buffer A. The protein was eluted via a gradient going from 0% Ni NTA Elution buffer (25 mM Tris pH 8.5, 500 mM NaCl, 5% glycerol and 250 mM imidazole) to 100% over 5 CV increasing by 20% B for each CV. The eluted fractions containing protein were equilibrated into Desalting Buffer (25 mM Tris pH 8.0, 400 mM NaCl) on a HiPrep 26/10 desalting column to remove imidazole. The protein was incubated overnight with 500 μg His-tagged ULP1 protease at 4° C to cleave the His-SUMO tag leaving the untagged protein and the cleaved His-SUMO and His-ULP1. Cleaved protein was then flowed over a HisTrap 5 mL column equilibrated in Ni-NTA buffer A to remove ULP1 and the His-SUMO product and collected in the flow-through. The flowthrough containing untagged protein was purified further using size exclusion chromatography (SEC) into SEC buffer (25 mM Tris-HCl, pH 8.0, 400 mM NaCl, 5% glycerol, and 1.5 mM TCEP) on a HiLoad 16/600 Superdex 200 pg column prior to concentration, aliquoting and flash-freezing.

Preparation of samples for nativeMS

For native mass spectrometry of the purified proteins, proteins were buffer exchanged into 500 mM ammonium acetate, 1 mM DTT using Zeba desalting columns (Thermo Fisher Scientific) at room temperature. Protein concentration was determined after exchange into buffer. Protein ranges were kept between 1 and 5 μM and electrospray performed using in-house nano-emitter capillaries. These were generated by pulling borosilicate glass (O.D. – 1.2 mm, I.D. – 0.69 mm, 10 cm, Sutter Instruments) using a Flaming/Brown micropipette puller (Model P-100, Sutter Instruments) to a tip diameter within 4-5 μm. Platinum wire was inserted into the emitter and connected to the source for sample charging. nMS was performed In a Q Exactive UHMR using a Nanospray Flex ion source (Thermo Fisher Scientific). Mass spectrometry parameters were optimized for each sample, for instance, spray voltage was in the range between 0.9 – 1.5 kV, the capillary temperature was 150-250 °C, the resolving power of the MS was in the range between 3,125 – 6,250 the in-source desolvation was between 50V and 300V. For incubation with lipids in detergent, lipids from Avanti polar lipids were dried under nitrogen flow and vacuum prior to resuspension via vortex in 500 mM ammonium acetate + 1 mM DTT + 1.25x CMC OGNG. Protein and lipid were incubated for one hour prior to being subjected to nMS analysis as described above. PHTH domain interactions with lipid were kept at 4 μM protein to 200 μM lipid. For the titration, FL BTK is kept at 2.5 μM, PS at 125 μM, and IP$_4$ concentration at 0 μM, 0.625 μM, 1.25 μM, and 2.5 μM.

Fluorescence Polarization

Samples were prepared in four replicates for each condition using phosphatidylserine 18:1-12:0 labeled on the fatty-acyl chain with nitrobenzodiazole (NBD) from Avanti Polar Lipids. NBD-PS was dried under Nitrogen gas and vacuum to remove residual solvent before being resuspended in Polarization buffer (25 mM Tris pH 8.0, 400 mM NaCl, 1.5 mM TCEP, 5% glycerol

and 0.029% OGNG). NBD-PS at 25 nM was used in all conditions. OGNG kept at half the critical
micellar concentration to provide solubilization without forming larger micelles that could interfere
with solution dynamics. Increasing concentrations of the purified bovine wild-type PHTH domain
were added at intervals between 0 and 200 µM, incubated for 30 minutes and binding measured
by fluorescence polarization in black-welled 96-well plates. Final buffer conditions in all wells were
Polarization buffer and samples were measured using an Agilent BioTek Synergy H1. Linearly
polarized light at 485 nm was sent through to excite the NBD and the parallel and perpendicular
emitted light at 528 nm was measured, read at a 6.25 mm height. Filter settings were Excitation
485/20 and Emission 528/20. Polarization values were calculated using the instrument software
Gen5 v3.11 following the standard formula $P = (F_{\parallel} - F_{\perp})/(F_{\parallel} + F_{\perp})$. Briefly, this is the difference
between the emitted intensity parallel to the excitation light plane and perpendicular over the total
emitted intensity. Polarization values are reported in mP (milliPolarization units).

<u>Liposome Formation</u>
Lipids of appropriate concentrations were dried under nitrogen and vacuum for one hour before
being resuspended in 500 mM ammonium acetate supplemented with 1 mM DTT on vortex for one
hour. Texas Red DHPE as a minor component was included to monitor vesicle stability during the
extrusion process for nMS experiments. Lipid mixtures were then subjected to ten freeze-thaw
cycles and extruded using an Avestin liposofast extruder with 100 nm membranes to ensure
homogeneity. BTK was prepared in 500 mM ammonium acetate and 1 mM DTT as described
previously. BTK and respective liposomes were incubated at concentrations of 1.775 µM and 1.5
mM liposomes in 500 mM ammonium acetate and 1 mM DTT. For the detection of BTK in the
presence of both liposomes and IP4, 0.85 µM $IP_4$ was supplemented. $IP_4$ was diluted to a working
concentration in the same buffer prior to addition.

<u>Kinase Activity</u>
Liposomes were dried under nitrogen and vacuumed, resuspended in liposome
resuspension buffer (25 mM Tris-HCl pH 7.4, 100 mM NaCl) at 2 mM and subjected to ten free-
thaw cycles using a 50° C water bath and liquid nitrogen. These hydrated vesicles are then
extruded to ensure 100 nm homogeneity using an Avestin LiposoFast. Full-length BTK at 4 µM and
liposomes at 500 µM were prepared in kinase buffer (25 mM Tris pH 7.5, 150 mM NaCl, 5%
glycerol) and subjected to a ten-minute incubation at room temperature. At which, time point 0 was
taken. At time point 0, an equal volume of kinase reaction buffer (25 mM Tris pH 7.5, 150 mM NaCl,
5% glycerol, 20 mM MgCl2, 2 mM ATP, and 2mM sodium orthovanadate) or kinase negative control
buffer (25 mM Tris pH 7.5, 150 mM NaCl, 5% glycerol and 2mM sodium orthovanadate) was added
and the reaction proceeded for five minutes. All kinase and negative control reactions were stopped
via the addition of an equivalent volume of 2x Laemmli sample buffer (Bio-Rad) with 100 mM EGTA
and 5% β-mercaptoethanol and 15 minutes of 95 C boiling.
*Auto*-phosphorylation of BTK at phosphoTyr223 was measured using western blotting.
Briefly, samples were run on an SDS-PAGE gel, before transfer to a PVDF membrane using a
Tank transfer apparatus (BioRad). Membranes were blocked for one hour rocking at room
temperature in 4% Bovine Serum Albumin in Tris-buffered saline with tween (TBST). Overnight
incubation in a 4 C rocker with 1:2000 (1:500 for 5% PIP₃) phosphoTyr223 antibody in 1% BSA in
TBST was followed by five ten-minute TBST washes and secondary incubation with 1:5000
(1:10,000) goat anti-mouse in 1% BSA in TBST and a sequential five ten-minute TBST washes
prior to imaging with SuperSignal West Femto or Pico PLUS Maximum Sensitivity Substrate
(Thermo Fisher Scientific). For Coomassie staining, the gels were placed into fixation solution (40%
methanol, 10% acetic acid), microwaved for thirty seconds and placed on a room temperature
rocker for thirty minutes. Then, the gels were microwaved for thirty seconds in QC colloidal
Coomassie stain (BioRad) and placed on a room temperature rocker overnight before rinsing
several times with water for destaining. Both chemiluminescence and Coomassie blue staining
were imaged on a BioRad Chemidoc. Pre-stained ladder was run for molecular weight conformation
and imaged in the westerns. Analysis and integration of the bands was performed on Fiji.

<u>Mathematical modeling</u>
Mathematical model adopted from Qi et. al. (2015) (18). Previous estimates of effective
BTK concentration (C) on the membrane surface were calculated as follows:
Firstly, the active volume where BTK can operate within the lipid vesicle context, or the shell of the
vesicle containing BTK is calculated as V = A x L (SI Fig. 8A). In the model, A is the surface area
of the vesicle and L is the length of BTK, around 100 Å as in the BTK composite full-length structure.
From this, C can be calculated as $C = \frac{N_{Btk}}{V \, x \, N_a}$ where $N_{BTK}$ is the number of BTK on a vesicle and $N_a$
is Avogadro's number. $N_{BTK}$ is calculated from the molar fraction ($f$) of PIP$_3$ and the total number
of lipids in the vesicle ($N_0$). $N_0$ is given by A x $A_0$, the surface area of a lipid head group, at roughly
0.6 nm$^2$(54).
C can then be calculated as:

524
$$C = \frac{N}{V \, X \, N_0} = \frac{f \, X \, \frac{A}{A_0}}{A \, X \, L \, X \, N_a} = \frac{f}{A_0 X \, L \, X \, N_a}$$

525 From here, we can modify this equation for our kinase assays, to become

526
$$C = \frac{f_1 + f_2(P)}{A_0 \, X \, L \, X \, N_a}$$

Where $f_1$ represents molar fraction of PIP$_3$, $f_2$ represents molar fraction of PS and P represents the
percentage of PS occupied by BTK. We then estimate P (the fraction of PS bound) based on our
experimental data, and assuming the bulk of IP$_4$ (PIP$_3$ analog) is bound in the 1:1 stoichiometry as
in the mathematical model. In our experimental titration with 0.625 µM IP$_4$ we are using exactly
200-fold the concentration of PS as IP$_4$ and achieving roughly 2-fold (1.916) the area of bound BTK.
This approximates to around 1 in 100 PS is bound to BTK using the PIP$_3$ analog for control. Our
values become $f_1$ = 0.005, 0.01, 0.025 and 0.05 according to PIP$_3$ concentrations, $f_2$ = 0, 0.1 and
0.2 accordingly P=0.01 for 1 in 100 bound, $A_0$ = 60 Å$^2$ , $N_a$ = 6.03 x 10$^{23}$ mol$^{-1}$.
535
536

537 **Acknowledgments**
538
539 We would like to thank former members of the Kuriyan lab for providing us with the plasmids. The
540 Shah lab for the YopH construct. We would like to thank Titus Boggon and his lab for the assistance
541 with their plate reader for the fluorescence polarization assays. This work was supported by the
542 National Institutes of Health grant R01GM141192 (to K. G.), R35GM147095 (to M.B.) and
543 F31CA278383 (to R.M.).

544
545
546
547
548 **References**
549

1.     J. R. Molina, A. A. Adjei, The Ras/Raf/MAPK pathway. *J Thorac Oncol* **1**, 7-9 (2006).
2.     S. Pal Singh, F. Dammeijer, R. W. Hendriks, Role of Bruton's tyrosine kinase in B cells and
malignancies. *Mol Cancer* **17**, 57 (2018).
3.     Y. Adachi *et al.*, Coincident Phosphatidic Acid Interaction Restrains Drp1 in
Mitochondrial Division. *Mol Cell* **63**, 1034-1043 (2016).
4.     J. E. Harlan, P. J. Hajduk, H. S. Yoon, S. W. Fesik, Pleckstrin homology domains bind to
phosphatidylinositol-4,5-bisphosphate. *Nature* **371**, 168-170 (1994).
5.     M. A. Lemmon, K. M. Ferguson, R. O'Brien, P. B. Sigler, J. Schlessinger, Specific and high-
affinity binding of inositol phosphates to an isolated pleckstrin homology domain. *Proc
Natl Acad Sci U S A* **92**, 10472-10476 (1995).

6.    K. Moravcevic, C. L. Oxley, M. A. Lemmon, Conditional peripheral membrane proteins:
facing up to limited specificity. *Structure* **20**, 15-27 (2012).
7.    A. A. Alnaas *et al.*, Multivalent lipid targeting by the calcium-independent C2A domain of
synaptotagmin-like protein 4/granuphilin. *Journal of Biological Chemistry* **296** (2021).
8.    M. A. Lemmon, Pleckstrin homology (PH) domains and phosphoinositides. *Biochem Soc*
*Symp* 10.1042/BSS0740081, 81-93 (2007).
9.    Y. Kawakami *et al.*, Tyrosine phosphorylation and activation of Bruton tyrosine kinase
upon Fc epsilon RI cross-linking. *Mol Cell Biol* **14**, 5108-5113 (1994).
10.   K. Salim *et al.*, Distinct specificity in the recognition of phosphoinositides by the
pleckstrin homology domains of dynamin and Bruton's tyrosine kinase. *EMBO J* **15**,
6241-6250 (1996).
11.   M. Hyvonen, M. Saraste, Structure of the PH domain and Btk motif from Bruton's
tyrosine kinase: molecular explanations for X-linked agammaglobulinaemia. *EMBO J* **16**,
3396-3404 (1997).
12.   Q. Wang, Y. Pechersky, S. Sagawa, A. C. Pan, D. E. Shaw, Structural mechanism for
Bruton's tyrosine kinase activation at the cell membrane. *Proc Natl Acad Sci U S A* **116**,
9390-9399 (2019).
13.   S. A. Johnson *et al.*, Phosphorylated immunoreceptor signaling motifs (ITAMs) exhibit
unique abilities to bind and activate Lyn and Syk tyrosine kinases. *J Immunol* **155**, 4596-
4603 (1995).
14.   V. Rolli *et al.*, Amplification of B cell antigen receptor signaling by a Syk/ITAM positive
feedback loop. *Mol Cell* **10**, 1057-1069 (2002).
15.   E. H. Walker, O. Perisic, C. Ried, L. Stephens, R. L. Williams, Structural insights into
phosphoinositide 3-kinase catalysis and signalling. *Nature* **402**, 313-320 (1999).
16.   D. A. Fruman *et al.*, The PI3K Pathway in Human Disease. *Cell* **170**, 605-635 (2017).
17.   H. Park *et al.*, Regulation of Btk function by a major autophosphorylation site within the
SH3 domain. *Immunity* **4**, 515-525 (1996).
18.   Q. Wang *et al.*, Autoinhibition of Bruton's tyrosine kinase (Btk) and activation by soluble
inositol hexakisphosphate. *Elife* **4** (2015).
19.   L. E. Kueffer, R. E. Joseph, A. H. Andreotti, Reining in BTK: Interdomain Interactions and
Their Importance in the Regulatory Control of BTK. *Front Cell Dev Biol* **9**, 655489 (2021).
20.   D. Y. Lin *et al.*, Conformational heterogeneity of the BTK PHTH domain drives multiple
regulatory states. *Elife* **12** (2024).
21.   T. J. Eisen *et al.*, Conditional requirement for dimerization of the membrane-binding
module for BTK signaling in lymphocyte cell lines. *Sci Signal* **18**, eado1252 (2025).
22.   D. J. Rawlings *et al.*, Activation of BTK by a phosphorylation mechanism initiated by SRC
family kinases. *Science* **271**, 822-825 (1996).
23.   D. Watanabe *et al.*, Four tyrosine residues in phospholipase C-gamma 2, identified as
Btk-dependent phosphorylation sites, are required for B cell antigen receptor-coupled
calcium signaling. *J Biol Chem* **276**, 38595-38601 (2001).
24.   Y. J. Kim, F. Sekiya, B. Poulin, Y. S. Bae, S. G. Rhee, Mechanism of B-cell receptor-induced
phosphorylation and activation of phospholipase C-gamma2. *Mol Cell Biol* **24**, 9986-
9999 (2004).
25.   P. Antony *et al.*, B cell receptor directs the activation of NFAT and NF-kappaB via distinct
molecular mechanisms. *Exp Cell Res* **291**, 11-24 (2003).

26.    J. A. Burger, A. Wiestner, Targeting B cell receptor signalling in cancer: preclinical and
606           clinical advances. *Nat Rev Cancer* **18**, 148-167 (2018).
607    27.    H. Tang *et al.*, IP(3)R-mediated Ca(2+) signaling controls B cell proliferation through
608           metabolic reprogramming. *iScience* **25**, 104209 (2022).
609    28.    M. Fukuda, T. Kojima, H. Kabayama, K. Mikoshiba, Mutation of the pleckstrin homology
610           domain of Bruton's tyrosine kinase in immunodeficiency impaired inositol 1,3,4,5-
611           tetrakisphosphate binding capacity. *J Biol Chem* **271**, 30303-30306 (1996).
612    29.    G. C. P. Schaafsma *et al.*, BTKbase, Bruton Tyrosine Kinase Variant Database in X-Linked
613           Agammaglobulinemia: Looking Back and Ahead. *Hum Mutat* **2023**, 5797541 (2023).
614    30.    J. K. Chung *et al.*, Switch-like activation of Bruton's tyrosine kinase by membrane-
615           mediated dimerization. *Proc Natl Acad Sci U S A* **116**, 10798-10803 (2019).
616    31.    A. D. Fenton, J. K. Chung, Membrane-dependent assembly of Bruton's tyrosine kinase
617           mediated by the Proline-rich region and SH3 domain. *Protein Sci* **34**, e70213 (2025).
618    32.    J. A. Corbin, R. A. Dirkx, J. J. Falke, GRP1 pleckstrin homology domain: activation
619           parameters and novel search mechanism for rare target lipid. *Biochemistry* **43**, 16161-
620           16173 (2004).
621    33.    C. L. Lai *et al.*, Molecular mechanism of membrane binding of the GRP1 PH domain. *J
622           Mol Biol* **425**, 3073-3090 (2013).
623    34.    X. Jian *et al.*, Molecular Basis for Cooperative Binding of Anionic Phospholipids to the PH
624           Domain of the Arf GAP ASAP1. *Structure* **23**, 1977-1988 (2015).
625    35.    N. Lucas, W. Cho, Phosphatidylserine binding is essential for plasma membrane
626           recruitment and signaling function of 3-phosphoinositide-dependent kinase-1. *J Biol
627           Chem* **286**, 41265-41272 (2011).
628    36.    A. Panda, C. Brown, K. Gupta, Studying Membrane Protein-Lipid Specificity through
629           Direct Native Mass Spectrometric Analysis from Tunable Proteoliposomes. *J Am Soc
630           Mass Spectrom* **34**, 1917-1927 (2023).
631    37.    A. Panda *et al.*, Direct determination of oligomeric organization of integral membrane
632           proteins and lipids from intact customizable bilayer. *Nat Methods* **20**, 891-897 (2023).
633    38.    H. Y. Yen *et al.*, PtdIns(4,5)P(2) stabilizes active states of GPCRs and enhances selectivity
634           of G-protein coupling. *Nature* **559**, 423-427 (2018).
635    39.    S. Chen *et al.*, Capturing a rhodopsin receptor signalling cascade across a native
636           membrane. *Nature* **604**, 384-390 (2022).
637    40.    Y. Zhu *et al.*, Cupric Ions Selectively Modulate TRAAK-Phosphatidylserine Interactions. *J
638           Am Chem Soc* **144**, 7048-7053 (2022).
639    41.    H. M. Sanders, M. M. Kostelic, C. K. Zak, M. T. Marty, Lipids and EGCG Affect alpha-
640           Synuclein Association and Disruption of Nanodiscs. *Biochemistry* **61**, 1014-1021 (2022).
641    42.    J. Le, J. A. Loo, Native Top-Down Mass Spectrometry Characterization of Model Integral
642           Membrane Protein Bacteriorhodopsin. *J Am Soc Mass Spectrom* **36**, 961-968 (2025).
643    43.    R. E. Joseph, T. E. Wales, D. B. Fulton, J. R. Engen, A. H. Andreotti, Achieving a Graded
644           Immune Response: BTK Adopts a Range of Active/Inactive Conformations Dictated by
645           Multiple Interdomain Contacts. *Structure* **25**, 1481-1494 e1484 (2017).
646    44.    G. van Meer, D. R. Voelker, G. W. Feigenson, Membrane lipids: where they are and how
647           they behave. *Nat Rev Mol Cell Biol* **9**, 112-124 (2008).
648    45.    S. Nagata, J. Suzuki, K. Segawa, T. Fujii, Exposure of phosphatidylserine on the cell
649           surface. *Cell Death Differ* **23**, 952-961 (2016).

46.   J. G. Kay, G. D. Fairn, Distribution, dynamics and functional roles of phosphatidylserine
within the cell. *Cell Commun Signal* **17**, 126 (2019).
47.   W. M. Moore *et al.*, Leaflet-specific phospholipid imaging using genetically encoded
proximity sensors. *Nat Chem Biol* 10.1038/s41589-025-02021-z (2025).
48.   M. Doktorova *et al.*, Cell membranes sustain phospholipid imbalance via cholesterol
asymmetry. *Cell* 10.1016/j.cell.2025.02.034 (2025).
49.   S. Kumar *et al.*, Native Mass Spectrometry of Membrane Protein-Lipid Interactions in
Different Detergent Environments. *Anal Chem* **96**, 16768-16776 (2024).
50.   W. A. Lea, A. Simeonov, Fluorescence polarization assays in small molecule screening.
*Expert Opin Drug Discov* **6**, 17-32 (2011).
51.   D. J. Marcotte *et al.*, Structures of human Bruton's tyrosine kinase in active and inactive
conformations suggest a mechanism of activation for TEC family kinases. *Protein Sci* **19**,
429-439 (2010).
52.   K. Saito, A. M. Scharenberg, J. P. Kinet, Interaction between the Btk PH domain and
phosphatidylinositol-3,4,5-trisphosphate directly regulates Btk. *J Biol Chem* **276**, 16201-
16206 (2001).
53.   G. Powis, E. J. Meuillet, M. Indarte, G. Booher, L. Kirkpatrick, Pleckstrin Homology [PH]
domain, structure, mechanism, and contribution to human disease. *Biomed*
*Pharmacother* **165**, 115024 (2023).
54.   T. Gulik-Krzywicki, E. Rivas, V. Luzzati, Structure and polymorphism of lipids: x-ray
diffraction study of the system formed from beef heart mitochondria lipids and water. *J*
*Mol Biol* **27**, 303-322 (1967).

672
673