Evolutionarily conserved transcriptional regulators control monoaminergic neuron development

Clifton Lewis[1,2,11], Matthew Goulty[1,3], Aniela Wroblewska[1], Nicola Croxall[4], David Onion[4], Sue Robinson[5], Robert P. Zinzen[6], Jordi Solana[7,8], Charlambos P. Kyriacou[1], Ezio Rosato[1], Roberto Feuda[1,9,10]

[1] Neurogenetics Group, Division of Genetics and Genome Biology, School of Biology and Biomedical Sciences, College of Life Sciences, University of Leicester, Leicester LE1 7RH, UK

[2] Centre for Chromosome Biology, Biomedical Sciences Building, University of Galway, Galway, H91 W2TY, Ireland

[3] The Francis Crick Institute, London, UK

[4] Faculty of Medicine & Health Sciences, Queen's Medical Centre, Nottingham NG7 2UH, UK

[5] Division of Molecular and Cell Biology, College of Life Sciences, University of Leicester, Leicester LE1 7RH, UK

[6] Systems Biology Imaging Technology Platform, Max Delbrück Centre for Molecular Medicine (MDC-BIMSB), Berlin, Germany

[7] Living Systems Institute, University of Exeter, Exeter, UK

[8] Department of Biosciences, University of Exeter, Exeter, UK

[9] Department of Biology, Geology and Environmental Science, University of Bologna, Bologna, Italy

[10] Correspondence: rf190@leicester.ac.uk / roberto.feuda@unibo.it (R.F.)

[11] Correspondence: clifton.lewis@leicester.ac.uk / clifton.lewis@universityofgalway.ie (C.L.)

Abstract

To what extent conserved developmental programs specify homologous cell types is a central question in biology. Here, we address this by focusing on reconstructing monoaminergic neuron development in *Drosophila melanogaster* embryo using time-resolved single-cell genomics, spatial transcript mapping with hybridisation chain reaction, and targeted metabolomics. We uncover a regulatory landscape in which specific transcription factors are activated before biosynthetic enzymes, establishing a prospective temporal architecture for monoaminergic fate specification. Comparative analyses of developmental single-cell atlases from zebrafish and sea urchin indicate that components of this machinery are conserved across ~550 million years of bilaterian evolution with orthologous transcription factors showing similar temporal dynamics. Together, these findings point to a putatively conserved regulatory core that interfaces with other context-dependent transcription factors; this interplay accommodates monoaminergic multifunction and subtype diversity across distinct neuroanatomies.

Introduction

Understanding whether conserved or divergent developmental programs underlie the specification of homologous cell types is a central goal in evolutionary and developmental biology. We address this question by focusing on monoaminergic neurons, a class of cells that produce neuromodulators such as dopamine and serotonin. Monoaminergic neurons provide a powerful system to study evolution and development for several reasons. They are molecularly defined by the expression of key biosynthetic enzymes (Libersat and Pflueger, 2004), they are widespread across bilaterians, their monoamine products can be precisely quantified, and the genes underlying their biosynthetic pathways have been shown to derive from a common evolutionary origin (Parent, 1984; Yamamoto and Vernier, 2011; Goulty *et al.*, 2023).

Although these neurons are extremely rare (<1% of neurons in fly and mammalian brains), they play fundamental roles in processes ranging from motor control and circadian rhythms to learning and mood (Libersat and Pflueger, 2004; Flames and Hobert, 2011; Monti, 2011; Waddell, 2013; Zheng *et al.*, 2025). Furthermore, their dysfunction contributes to major human neurological and psychiatric disorders, including Parkinson's disease, depression, and schizophrenia (Kurian *et al.*, 2011; Volkow *et al.*, 2011; Grace, 2016).

The developmental specification of monoaminergic neurons requires the coordinated activity of multiple transcription factors (TFs) and other regulatory genes organised into hierarchical gene regulatory networks (GRNs) (Hobert, 2008; Flames and Hobert, 2009, 2011). Studies across vertebrates and invertebrates have identified TFs that are critical for establishing monoaminergic fate (Flames and Hobert, 2011). In vertebrates, for example, LIM-homeodomain proteins Lmx1a/b, the ETS factor Fev/Pet-1, and Nkx2.2 are essential for serotonergic neuron specification (Cheng *et al.*, 2003; Ding *et al.*, 2003; Hendricks *et al.*, 2003). In *Drosophila*, the zinc-finger proteins Eagle (eg) and Huckebein (hkb) play similar roles in serotonergic differentiation (Dittrich *et al.*, 1997). In *C. elegans*, dopaminergic fate is controlled by terminal selector TFs such as the ETS transcription factor, AST-1 (Flames and Hobert, 2009). Previous work from our lab established that monoaminergic cell identity is defined by a conserved set of transcription factors shared across bilaterian species, suggesting that a common regulatory basis for these neurons originated early in animal evolution (Goulty *et al.*, 2025). Yet these candidate-based studies provide only

a partial view of the regulatory landscape underlying monoaminergic specification and

their possible conservation. We therefore still lack a systems-level understanding of

how monoaminergic neurons develop. Addressing these questions is essential for

understanding whether homologous cell-types develop following a conserved

transcriptional program.

Here to address this open question, we combine time-resolved single-cell

transcriptomics, trajectory modelling, spatial transcript validation, and targeted

metabolomics in embryonic neurogenesis of *Drosophila melanogaster*, and test for

evolutionary conservation in zebrafish and sea urchin development. Our results reveal

the developmental program of monoaminergic neurons in the *Drosophila* embryo,

including key TFs—*fd59A*, *dmrt99B*, and *Lmx1a*—that pattern these lineages and

differences in maturation rates between subtypes. We also identified several TF

orthogroups conserved across *Drosophila*, sea urchin, and zebrafish. Comparison of

monoaminergic developmental trajectory between *Drosophila* and zebrafish revealed

a similar developmental cascade of TFs. Together, these findings support a putative

conserved temporal logic for monoaminergic fate acquisition and begin to explain how

developmental fate and function can align and/or diverge across species.

Results

A temporally resolved single-cell atlas captures neurogenesis in *Drosophila*. To

understand the development of monoaminergic neurons, we performed single-cell

RNA sequencing (scRNA-seq) on *Drosophila melanogaster* embryos collected at eight

partially overlapping time windows spanning 0 to 22 hours after egg laying (AEL), with

two biological replicates per window (**Figure 1A.**), covering the transition from

neuroectoderm to fully differentiated embryonic neurons. To enrich for progenitor and

early neural population, we used a SoxN-sfGFP transgenic line (VDRC #318062,

Sarov *et al.*, 2016) to isolate proneural cells and neuroblasts, and anti-Elav

immunostaining to enrich for differentiating neurons, followed by fluorescence-

activated cell sorting (FACS) (**Figure S1.**). Single cell RNA-seq libraries were

prepared on the 10X Genomics v3.1 platform and sequenced using Illumina NovaSeq

X Plus.

After standard quality control, we retained 57,476 high-quality cells, resolving

into transcriptionally distinct clusters. Cluster identities were assigned by mapping

cluster-specific marker genes—selected as the top differentially expressed features
with adjusted p-value < 0.05—to *in situ* expression patterns from the Berkeley
Drosophila Genome Project (BDGP) using an annotation pipeline adapted from
Calderon *et al.* (2022). This analysis revealed a clear temporal transition: early clusters
were enriched for maternal and early zygotic genes, whereas later stages featured
organised, differentiated tissues; including the brain primordium, embryonic brain, and
embryonic gut (**Figure 1B.** and **Table S1.**). Embedding all time points together
revealed a continuous developmental trajectory (**Figure 1C.**), capturing the
progressive emergence of neuronal identity.

Neurotransmitter subtype emergence reveals early co-expression and lineage
complexity. Although our dataset is enriched for neural progenitors and differentiated
neurons, cluster annotations revealed the presence of additional cell types (**Figure**
1C.). This likely reflects autofluorescence and/or putatively *SoxN*-driven GFP
expression in the developing gut, leading to non-specific capture during FACS
enrichment (**Figure S1.**).
To resolve distinct stages of neurogenesis, we used established markers - e.g.
dpn, *wor*, *mir*, *elav*, *nSyb* - to identify neuroblasts, ganglion mother cells (GMCs),
intermediate neural progenitors (INPs), and mature neurons in a population of ~40,000
cells (**Figure 2A.**, **Figure S3A.**) (Dillon *et al.*, 2022; Peng *et al.*, 2024). To capture the
full neurogenic continuum, spanning from neuroblasts to mature neurons, we
subsampled to ~10,000 high-quality cells, excluding disparate clusters (**Figure 2B.**,
Figure S3B.). These cells were distributed across all the developmental time points,
enabling the reconstruction of continuous differentiation trajectories (**Figure 2C.**).
Among these terminal-state mature neurons, we identified seven neurotransmitter-
defined populations: cholinergic (931 cells), GABAergic (816 cells), glutamatergic (173
cells), monoaminergic (77 cells), and three mixed populations co-expressing markers
of GABA and glutamate (100 cells), acetylcholine and GABA (556 cells), or serotonin
and GABA (388 cells), respectively (**Figure 2B.**).
Given that distinct TF combinations are known to contribute to neuronal subtype
identity, we next sought to identify TFs associated with each neurotransmitter-defined
population (**Figure 2E.**). Our differential TF expression analysis revealed that, with the
exception of neuroblasts/GMCs, GABA/glutamatergic neurons, and monoaminergic
neurons, most of these neuronal populations could not be unambiguously resolved

based on TF expression alone. This outcome suggests that TFs in these other
lineages may lack strict subtype specificity, that neuronal identity may arise from more
complex regulatory interactions beyond TF expression, or that the dataset may not
have sufficient depth to fully resolve these distinctions. Monoaminergic neurons,
however, showed co-expression of multiple TFs, including *fd59A*, *dmrt99B*, *Lmx1a*,
and *Vsx1* and *Vsx2* (**Figure 2E.**). In contrast, the mixed serotonergic/GABAergic
population was marked by the co-expression of *CG8301* and *br* (**Figure 2E.**). These
findings define candidate regulatory genes underlying monoaminergic fate.
150
RNA velocity reveals continuous differentiation trajectories. To investigate the
transcriptional dynamics underlying monoaminergic neuron differentiation, we applied
RNA velocity analysis using scVelo (La Manno *et al.*, 2018; Bergen *et al.*, 2020) and
modelled lineage trajectories with CellRank (v2) (Lange *et al.*, 2022; Weiler *et al.*,
2024), which integrates splicing kinetics (Velocity Kernel), transcriptomic similarity
(Connectivity Kernel), and CytoTRACE dynamics to infer directed developmental
transitions (CytoTRACE Kernel). We first applied scVelo to the full dataset. Velocity
vector fields projected onto the UMAP embedding revealed a strong correlation
between transcriptional dynamics and the temporal structure of the dataset (**Figure
S4**).

We then focused on the subclustered population spanning the full neurogenic
trajectory from neuroblasts to mature neurons (**Figure 2B–D**). Application of scVelo
and CellRank to this dataset recovered clear trajectories for each subtype and a
smooth latent-time ordering (**Figure S5A**). To interpret the underlying programs, we
identified velocity-inferred dynamical genes with the scVelo dynamical model and
ordered cells by latent time (**Figure S5B.**). We discretised latent time into 20 bins and,
for each bin, compiled the genes whose fitted expression/velocity profiles peaked
within that interval. Gene Ontology (GO) over-representation analyses (ORA) were
then performed using all expressed genes in the subset as background with
Benjamini–Hochberg correction. The enrichment landscape shows a coherent
progression: (i) early bins are dominated by cell-state remodelling and tissue-building
programs—RNA processing, cell cycle/proliferation, morphogenesis, and
organogenesis; (ii) intermediate bins transition toward neuronal differentiation, with
terms related to synapse formation and cell morphogenesis involved in neuron

differentiation; while (iii) late bins are enriched for synaptic and excitability modules,
including neuron recognition, axon guidance, and neuroactive ligand–receptor
interaction, consistent with the acquisition of functional neuronal properties. Together,
these GO trends align with the velocity-derived latent-time ordering and provide
functional support for a continuous maturation from progenitor states to functionally
specialised neurons (**Figure S5C.**).

To reconstruct the differentiation trajectory of monoaminergic neurons in
greater detail, we applied CellRank to define initial and terminal states and to model
transcriptional dynamics within this specific lineage (**Figure 3A**). Genes encoding key
enzymes required for monoaminergic neurotransmission—including *ple*, *Tdc1*, *Tdc2*,
Tbh, and *Vmat*—were strongly upregulated toward the terminal state, as expected,
marking the final stages of differentiation. With the exception of *Trhn* and *Ddc*, these
enzymes exhibited high specificity for the monoaminergic lineage (**Figure S6.**),
indicating subtype-restricted induction of the biosynthetic program. Similarly, top-
ranking TFs associated with monoaminergic identity—including *fd59A*, *dmrt99B*, and
Lmx1a—were also selectively expressed in the monoaminergic trajectory (**Figure 3B.**)
and peaked at approximately the same latent time as the biosynthetic genes (**Figure
3A.**). This coordinated expression suggests a tightly coupled regulatory program in
which TFs may directly activate terminal differentiation effectors. In contrast—and
consistent with their role as responders to monoamine signals—genes encoding
monoaminergic receptors showed broader expression across multiple neuronal
lineages (**Figure S7.**). This pattern suggests that receptor deployment is decoupled
from biosynthetic identity and instead reflects widespread postsynaptic
responsiveness to monoamines.
199
Spatial validation supports coordinated transcriptional activation and resolves
monoaminergic subtype identity. Although our scRNA-seq atlas resolves the major
neuronal lineages and transmitter classes, monoaminergic neurons are a rare
population in the *Drosophila* CNS (Busch *et al.*, 2009; Hartenstein *et al.*, 2017; Kasture
et al., 2018; Babski, Codianni and Bhandawat, 2024). As a result, it was difficult to
clearly separate specific monoaminergic subtypes from transcriptomes alone. To both
validate CellRank-inferred dynamics and add spatial specificity, we focused on two

subtypes – serotonergic and dopaminergic - and performed hybridisation chain
reaction (HCR) (Choi *et al.*, 2018)**.**

We assayed a panel that included most of the monoaminergic biosynthetic
enzymes, their receptors, and six transcription factors (**Figure 4.**, **Figure S8-9.**). TFs
were chosen based on differential expression within the monoaminergic lineage
(**Figure 2C.**) and prior evidence for roles in monoaminergic fate. Spatial profiling
showed that *fd59A*, *dmrt99B*, and *Lmx1a* are expressed in the CNS beginning at
stages 10–12 (~8 h AEL), preceding the onset of biosynthetic gene expression. In
contrast, monoaminergic enzymes were robustly expressed from stage 12 (~10 h
AEL) onward, consistent with the timing of neuronal specification. Notably, this TF
induction occurs slightly earlier than suggested by our latent-time analysis, which
predicted a more coordinated co-activation with the enzymes. Receptors generally
followed a similar onset, with most appearing from stage 12. A notable exception was
5-HT2A, which was detected as early as stage 1 and in non-neural tissues, consistent
with a recent study implicating it in early morphogenetic roles beyond
neurotransmission (Karki *et al.*, 2023)**.**

Together, these data mostly corroborate the temporal ordering inferred by
CellRank and, crucially, resolve subtype identity by localising expression to
serotonergic or dopaminergic domains. They further suggest that these TFs, *fd59A*,
dmrt99B, and *Lmx1a*, may act upstream to specifically initiate part of the
monoaminergic program, with effector enzymes and receptors switching on
subsequently.

Mass spectrometry reveals temporally distinct onset of monoamine production.
Our transcriptional and spatial analyses suggests that by stage 12 (~9–10 h AEL), the
Drosophila embryo expresses the biosynthetic enzymes and receptors required to
produce and respond to monoamines. However, whether these neurons are
functionally active at this stage remains unresolved. Previous studies have detected
serotonin and dopamine via immunohistochemistry only from stage 17 (~15 h AEL)
onwards (Lundell and Hirsh, 1994), raising questions about the relationship between
transcriptional commitment and neurotransmitter availability. To resolve this
dichotomy, we conducted targeted mass spectrometry across seven developmental
windows spanning 0–22 h AEL. At each time point, we analysed ≥6 biological

replicates (54 samples total) and quantified 19 analytes, including all canonical
monoamines (**Table S2.**) GABA, glutamate, and acetylcholine **(Figure S11.).**

Our results revealed distinct temporal profiles across monoaminergic subtypes
(**Figure 5.; Figure S10.**). Histamine and tyramine were constitutively present
throughout development, suggesting their production may occur independently of
canonical neurogenesis. In contrast, both dopamine and serotonin exhibited
temporally restricted accumulation, with sharp increases after stage 12—closely
aligning with the transcriptional induction of their biosynthetic enzymes. Although
dopamine and serotonin followed similar temporal trajectories, their relative levels
differed, likely reflecting variations in biosynthetic efficiency or substrate availability. It
is important to note that this effect may also reflect differences in the detection
capabilities for these molecules.

Together, these findings indicate that transcriptional programs broadly predict
the functional potential of monoaminergic neurons, but the timing and extent of
neurotransmitter production are modulated by subtype-specific biochemical properties
and/or regulatory constraints.

Conserved transcriptional dynamics underpin monoaminergic neuron
development in *Drosophila* and zebrafish. Monoaminergic neurons are widely
distributed across bilaterian animals, yet whether they arise from a conserved
developmental program remains unclear. Our findings in *Drosophila* suggest that a
specific set of TFs are activated during monoaminergic neuron differentiation,
providing an opportunity to test the evolutionary conservation of this regulatory
program. To address this question, we reanalysed published developmental single-
cell RNA-seq data from *Danio rerio* and sea urchin, *Strongylocentrotus purpuratus*.
For *Danio rerio*, we reprocessed a developmental time series spanning embryonic to
larval stages (10 hours post-fertilisation [hpf] 232 to 10 days post-fertilisation [dpf];
Lange *et al.*, 2024). For *S. purpuratus*, we merged two complementary datasets,
covering the 8-cell embryo through the 72 hpf larval stages (Foster, Oulhen and
Wessel, 2020; Perillo *et al.*, 2020). Central nervous system cell types were annotated
using orthologous marker genes applied previously in *Drosophila* (**Figure 2D.**) and
identified monoaminergic neuron populations based on expression of canonical
biosynthetic genes (**Figure S12A-B.**).

Applying the same differential-expression (DE) criteria used in fly (adjusted p <
0.05; |log$_2$FC| ≥ 0.25), we identified transcription factors (TFs) enriched in zebrafish
and sea urchin monoaminergic clusters analysed independently (**Figure 6A.**; **Table**
S3). We analysed clusters separately, rather than collapsing them into a single
monoaminergic group, to gather an inclusive list of TFs that define monoaminergic
identity across subtypes. To enable cross-species comparisons despite lineage-
specific duplications, genes were mapped to bilaterian-level eggNOG orthogroups.
Orthologs of key fly regulators were recovered in vertebrates, including *lmx1a* paralogs
(*lmx1a*, *lmx1al*, *lmx1ba*). In total, 15 TF orthogroups were differentially expressed in
both fly and zebrafish monoaminergic neurons, and 16 TF orthogroups were shared
between fly and sea urchin (**Figure 6C.**; **Table S4.**), defining a set of putative
conserved regulators. Notably, six TF orthogroups—*CG8301*, *Lmx1a*/*CG4328*, *Hr3*,
Eip78C, *net*, and *Pdp1*—were shared across all three species comparisons,
underscoring a potentially conserved core program for monoaminergic identity. The
partial overlap of the remaining TFs suggests that, while a conserved regulatory core
might exist, additional factors have diverged or been co-opted in lineage-specific
contexts

To test whether this conservation was unique to monoaminergic neurons, we
repeated the orthogroup-level analysis between *Drosophila* and zebrafish for other
neurotransmitter classes (Acetylcholine, GABA, Glutamate; **Figure S13.**). The number
of shared DE TF orthogroups was of similar magnitude across classes, despite vastly
different neuronal population sizes, supporting a model in which each transmitter
lineage may share a small, conserved set of regulators that is then diversified by
lineage- and species-specific additions.

We next asked whether not only TF identity but also their temporal activation is
conserved. Using the comparable time-resolved zebrafish dataset, we applied RNA
velocity and CellRank to reconstruct differentiation dynamics. Several orthologous
TFs, including *lmx1* paralogs, showed late upregulation along the zebrafish
monoaminergic trajectory, closely mirroring the dynamics observed in *Drosophila*
(**Figure 6B.**). Thus, some of the regulatory content and timing of activation are aligned
between insects and vertebrates, at least in these representative species.

Together, these findings support a model in which monoaminergic neurons
emerge from a potential conserved transcriptional program, with shared regulators,
such as *Lmx1a*, but critically has undergone significant species-specific elaborations.

Discussion

The extent to which homologous cell types are specified by conserved developmental and transcriptional programs remains a central question in evolutionary developmental biology. Monoaminergic neurons, which play fundamental roles in neurotransmission across bilaterians, offer a powerful model for addressing this issue. Previous comparative transcriptomic analyses have suggested that monoaminergic cell identity is defined by a shared set of transcription factors, indicating an origin in the bilaterian stem lineage (Goulty *et al.*, 2025). Yet, whether the developmental processes that give rise to these neurons are likewise conserved—or instead represent cases of convergent evolution—has remained unresolved. Additionally, by determining whether the developmental program underlying these neurons is shared across species, we further clarify the hypothesis of a conserved evolutionary origin. In this study, we characterise the transcriptional program governing monoaminergic neuron development in *Drosophila* embryos and reveal that key regulatory components are shared with vertebrate and deuterostome systems. Our findings suggest that monoaminergic neurons are specified by an evolutionarily conserved developmental framework that emerged early in bilaterian evolution.

Within *Drosophila*, we identify a regulatory program involving *fd59A*, *dmrt99B*, *Vsx1*, *Vsx2*, and *Lmx1a*. These TFs are selectively expressed in the monoaminergic lineage and become active around stage 12, coincident with the onset of biosynthetic enzyme expression (*ple*, *Tdc2*, *Tbh*, *Vmat*). Spatial validation by HCR confirmed the sequential activation of these genes and showed that TF induction precedes biosynthetic gene expression, supporting their role as primary regulators of monoaminergic fate. This organisation mirrors vertebrate serotonergic development, where Lmx1a/b and Pet-1/Fev initiate comparable programs (Ding *et al.*, 2003; Hendricks *et al.*, 2003; Flames and Hobert, 2011), pointing to deep evolutionary parallels.

Although biosynthetic gene expression initiates around stage 12, metabolomic profiling reveals pronounced differences in monoamine accumulation. Dopamine and serotonin both increase after stage 12, yet dopamine accumulates more rapidly—likely reflecting the higher catalytic efficiency of Dopa decarboxylase (Ddc) for L-3,4-dihydroxyphenylalanine (L-DOPA) than for 5-hydroxytryptophan (5-HTP), the respective intermediates in the dopamine and serotonin biosynthetic pathways

(Montioli and Borri Voltattorni, 2021). In contrast, octopamine remains undetectable
despite enzyme expression, suggesting either technical limitations or additional
biological constraints such as substrate availability, enzymatic kinetics, or post-
transcriptional control. These findings indicate that transcriptional activation alone is
insufficient for transmitter synthesis, pointing to additional regulatory layers that
modulate monoamine output. However, these interpretations remain constrained by
the detection sensitivity and chemical properties of the specific metabolites analysed.

Our comparative analysis of monoaminergic neuron development in
Drosophila, zebrafish, and sea urchins reveals a deeply conserved transcriptional
program, underscoring a shared regulatory logic that likely emerged early in bilaterian
evolution. The identification of six conserved TF orthogroups—*CG8301*,
Lmx1a/*CG4328*, *Hr3*, *Eip78C*, *net*, and *Pdp1*—establishes a core genetic program for
specifying monoaminergic identity. This finding supports the hypothesis that these
neurons represent potentially homologous cell types that originated early in bilaterian
evolution. Core regulatory modules specifying neuronal identity appear to have been
established in the bilaterian ancestor and maintained across deep evolutionary time
(Arendt, 2005; Arendt *et al.*, 2019; Rapti, 2022; Sachkova, Modepalli and Kittelmann,
2025). For instance, Lmx1a has a well-documented and critical role in the
development of dopaminergic neurons in vertebrates, where it regulates key effector
genes such as tyrosine hydroxylase (*Th*) and the dopamine transporter (*DAT*) to
ensure proper differentiation and maturation (Cai *et al.*, 2009; Doucet-Beaupré, Ang
and Lévesque, 2015). This role has not previously been demonstrated in non-
vertebrates, suggesting that *Lmx1a* may represent an ancient regulator of
monoaminergic fate specification—although convergent evolution of similar
transcriptional strategies cannot be excluded.

Importantly, this conservation is not absolute. We also observed substantial
regulatory divergence, with several TFs associated with monoaminergic differentiation
being unique to specific lineages. This pattern may reflect a dual-layered evolutionary
process—an ancient, conserved core program governing fate commitment,
complemented by lineage-specific regulators that refine neuronal subtype identity
(Hobert, 2008; Arendt *et al.*, 2019). Alternatively, the conserved TF set itself may
specify distinct monoaminergic identities across species, while divergent TFs act as
complementary components that integrate these core programs into broader gene-
regulatory networks, coordinating cross-talk and aligning specification with species-

specific neural architectures (Flames and Hobert, 2011; Rapti, 2022). While some
apparent differences may arise from technical factors (e.g., sequencing depth,
detection sensitivity, or annotation), others likely represent genuine regulatory
divergence. Given the substantial variation in neuroanatomical organisation among
insects, vertebrates, and echinoderms, such divergence may reflect adaptive
remodelling of the regulatory landscape to distinct neural architectures.

The temporal dynamics of conserved TFs suggest that these genes govern
monoaminergic fate commitment, whereas lineage-specific TFs modulate neuronal
subtype identity in a species-dependent manner. The hierarchical relationships
between conserved and lineage-specific TFs remain to be elucidated. Determining
whether these factors operate through feedforward or feedback regulatory loops—a
recurrent feature of GRN architecture that stabilises neuronal identity—will be crucial
(Flames and Hobert, 2011; Peter and Davidson, 2011). Understanding how conserved
GRNs are modified to generate lineage-specific phenotypes will be essential for
uncovering the general principles that govern cell-type diversification across evolution.

Our mass-spectrometry data indicate a partial decoupling between monoamine
abundance and biosynthetic enzyme expression. While enzyme kinetics may account
for some of the observed differences in transmitter accumulation, it remains unclear
whether such decoupling is a general feature of monoaminergic neurons across taxa.
Moreover, the functions of early monoamine signalling—potentially enabled by
precocious receptor expression (e.g., 5-HT2A)—during embryogenesis remain
unresolved.

Integrating transcriptomic, metabolomic, and spatial data across additional
bilaterian taxa will be important for testing the universality of these patterns.
Comparative perturbation experiments—targeting conserved TFs such as Lmx1a or
Hr3 in non-model systems—could determine whether shared transcriptional logic
indeed translates into conserved functional outcomes.

In conclusion, our study uncovers a conserved temporal program for
monoaminergic neuron specification, in which early transcription factor activation
orchestrates biosynthetic gene expression to align fate determination with functional
maturation. By linking transcriptional programs, transmitter synthesis, and receptor
deployment across species, our work provides a framework for understanding how
conserved and divergent GRNs generate neuromodulatory diversity and how neuronal
fate and function are developmentally coupled.

Methods

Fly husbandry

All *Drosophila melanogaster* stocks were maintained on standard cornmeal–yeast–agar medium at 18°C or 25°C, with relative humidity held at approximately 40%. We employed a transgenic SoxN–sfGFP line (VDRC #318062), which was reared continuously at 25°C (Sarov *et al.*, 2016).

Embryo collection

Adult flies were transferred into population cages fitted with agar plates prepared from mango juices, each supplemented with fresh yeast paste to boost fecundity. To eliminate circadian-driven fluctuations in gene expression, cages were kept in constant darkness (DD) beginning immediately upon transfer (Wijnen *et al.*, 2006). Following a 48-hour acclimation under DD, embryo collections was commenced, ensuring that all developmental stages progressed without light-induced transcriptional variability.

Embryos were collected over eight successive, overlapping time windows covering the full span of embryonic development; between 0-22 hours after egg laying (AEL). Collections were performed in biological duplicate to increase data reliability and resolution of transitional stages. Agar plates were replaced regularly to maintain optimal oviposition conditions, and collected embryos remained in the same DD incubator for precise aging.

Given the necessity of preserving RNA integrity for 10X Genomics single-cell RNA-sequencing, we avoided formaldehyde-based fixatives. Instead, we applied dithio-bis(succinimidyl propionate) (DSP; Sigma-Aldrich), a reversible cross-linker compatible with downstream library preparation. A 50 mg/mL DSP stock was made in DMSO, then diluted by adding 5 mL Dulbecco's PBS (DPBS; Gibco) to the 100 µL stock, followed by sterilisation through a 0.2 µm syringe filter. Collected embryos were fixed in this working solution, preserving stage-specific transcriptomes and enabling reliable reversal of cross-links prior to sequencing.

Embryo Harvesting, Dechorionation, and Fixation

Collected embryos were gently dislodged from agar plates by rinsing with distilled water and transferred into a 30 µm nylon mesh sieve using a fine paintbrush. Any residual yeast paste was washed away under a gentle stream of distilled water to

minimise contamination. Dechorionation was performed by submerging the embryos

in 50% bleach (Arco Essentials) for 3 minutes, then immediately rinsing them

thoroughly with distilled water until all bleach was removed.

For fixation, equal volumes of the DSP working solution and heptane (Fisher

Scientific) were combined in a glass scintillation vial. Dechorionated embryos were

added to this biphasic mixture and incubated on a gyroscopic rocker at room

temperature for 1 hour, ensuring gentle but continuous agitation. After fixation, the

lower aqueous phase was discarded, and cold methanol (VWR) was added to the

heptane layer to initiate devitellinisation. Embryos were vortexed for 2 minutes to

facilitate complete removal of the vitelline membrane, then allowed to settle to the

bottom of the vial. Finally, embryos were transferred to 1.5 mL Eppendorf tubes and

washed three times in fresh methanol. Fixed embryos were stored at –20°C until

further processing.

Immunofluorescence

All immunostaining was performed at 4°C under stringent RNase-free conditions:

Protector RNase Inhibitor (Roche, RNAINH-RO) was added to all buffers at 5 U/mL

during brief washes and 20 U/mL for all longer incubations, and both bovine serum

albumin (BSA; Gibco) and normal goat serum (NGS; Vector Laboratories) were heat-

inactivated at 60°C for 30 minutes to eliminate residual RNases. Prior to antibody

staining, DSP–methanol–fixed embryos were gradually rehydrated through an ethanol

series into PBT (PBS + 0.1% Triton X-100; Sigma-Aldrich T8787) and then washed

three times for 10 minutes each in PBT containing RNase inhibitor, with gentle rocking

to ensure complete removal of the methanol fixative. To block non-specific binding,

embryos were incubated overnight at 4°C in PBT supplemented with 1% heat-

inactivated BSA and 2% heat-inactivated NGS. The embryos were then transferred

into fresh blocking buffer (PBT + inhibitor, 1% BSA, 2% NGS) containing primary

antibodies: rabbit α-GFP (1:100; Invitrogen) to label SoxN–sfGFP$^+$ neurogenic cells,

and rat α-Elav (1:100; DSHB) to detect post-mitotic neurons. After overnight incubation

at 4°C with gentle agitation, excess primary antibody was removed by three 10-minute

washes in PBT. Secondary detection was performed by incubating embryos overnight

in the same blocking buffer supplemented with goat α-rabbit Alexa Fluor 488 (1:500;

Invitrogen), mouse α-rat PE (1:500; Invitrogen), and the nuclear counterstain DRAQ5

(4 µL/mL; Thermo Scientific). Finally, stained embryos were washed three times in
PBT (10 minutes each) and stored at 4 °C in PBT until mounting and imaging.
476

Microscopy validation and processing
Images were acquired using a Zeiss LSM 980 Airyscan 2 microscope in Airyscan
super-resolution mode, with the default settings applied for Airyscan image
processing. A Plan-Apochromat 10x/0.45 or Plan-Apochromat 20x/0.8 objective was
employed for image acquisition (University of Leicester Advanced Imaging Facility).
Post-processing was performed using a custom FIJI (Schindelin *et al.*, 2012) macro
for orienting the embryos along anatomical axes (Embryo Processing IF Script).
484

Single-cell dissociation
Embryos were transferred into an ice-cold 15 mL Dounce homogeniser in PBS
containing Protector RNase Inhibitor, and all subsequent steps were performed at 4
°C to preserve RNA integrity. After three gentle washes with cold Dulbecco's PBS
(DPBS), the supernatant was removed and replaced with PBS supplemented with
0.04% BSA (PBS/BSA). Embryos were then mechanically disrupted by 10–15 strokes
with the tight-fitting pestle, and the homogenate was collected into a fresh tube. To
maximise cell recovery, the homogeniser was rinsed with additional PBS/BSA, and
the combined suspension was centrifuged at 3,000 × g for 3 minutes. The supernatant
was discarded, and the pellet was resuspended in PBS/BSA.

To remove large debris, the resuspended cells were centrifuged at 40 × g for 3
minutes, after which the supernatant—containing single cells—was transferred to a
new tube and spun at 1,000 × g for 3 minutes. The resulting pellet was gently triturated
in PBS/BSA by passing the suspension through a 22-gauge needle with 20 strokes to
dissociate any remaining aggregates, then filtered through a 100 µm nylon mesh. A
final spin at 1,000 × g for 3 minutes pelleted the cleaned single-cell suspension, which
was resuspended in PBS/BSA and kept on ice prior to fluorescence-activated cell
sorting (FACS). Parallel control samples—unstained; stained with only primary or only
secondary antibodies; and combinations thereof with DRAQ5 nuclear stain—were
prepared and processed identically for gating and compensation during FACS
analysis.

FACS

Cells were sorted on a Beckman Coulter MoFlo Astrios cell sorter (Nottingham) equipped with a 100 µm nozzle, operating in "Purify" precision mode with a 1–2 drop envelope to maximise purity of collected fractions. Single-cell suspensions were maintained at 4°C throughout sorting by housing the entire flow chamber in a water-cooled environment, thereby minimising heat-induced stress and preserving cell integrity. Samples were deposited into a two-way tube holder pre-chilled on ice.

Gating strategies were defined using unstained and single-stained controls to eliminate autofluorescent events. We applied broad gates to capture both strongly and weakly fluorescent populations, acknowledging the continuum of marker expression. SoxN$^+$ and Elav$^+$ cells—representing early neurogenic and post-mitotic neuronal cohorts, respectively—were collected together. This inclusive approach ensured recovery of the full spectrum of developing neural cells.

Reverse cross-linking

After FACS, the cells were subjected to a reverse cross-linking step to remove the DSP fixative and prepare them for subsequent library preparation. The sorted cells were first centrifuged at 4°C at 1,000 x g for 3 minutes, and the supernatant was discarded. The cell pellet was then resuspended in PBS/BSA solution and incubated with 50 mM dithiothreitol (DTT, Sigma-Aldrich) at 37°C for 30 minutes to break the cross-links formed by DSP. Following incubation, the cells were centrifuged twice at 4°C at 1,000 x g for 3 minutes to thoroughly remove any residual DTT.

10X Genomics Library Preparation

The library preparation was conducted using the 10X Genomics Chromium Next GEM Single Cell 3' Reagent Kit v3.1 (10X Genomics). Prior to the preparation, cell concentration was assessed with a Luna FX7 Cell Counter, and cells were adjusted to a concentration of approximately 500-1500 cells/µl. Approximately 8000 cells per sample were used to generate single-cell gel beads in emulsion (GEMs), targeting approximately 5000 single-cell GEMs per sample. GEM emulsions were created with the 10X Chromium Controller, followed by complementary DNA (cDNA) synthesis and library preparation according to the manufacturer's protocol. cDNA amplification was performed for 12 cycles, and library amplification during the PCR indexing stage was similarly carried out for 12 cycles using the Dual Index Kit TT Set A (10X Genomics).

Quantification of cDNA was performed using Qubit dsDNA HS assays (Invitrogen) and D5000 HS ScreenTape (Agilent) on an Agilent Technologies TapeStation 4200, while library quantification was conducted with Qubit dsDNA HS assays and D1000 ScreenTape on the same platform. Libraries were prepared following the manufacturer's guidelines and were subsequently sequenced externally by Novogene on the Illumina NovaSeq X Plus platform, utilising 150 bp paired-end reads.

HCR Validation

Custom probe sets targeting mRNAs of key monoaminergic pathway enzymes, receptors, and TFs were designed from the longest isoform FlyBase annotations and synthesised by Molecular Instruments. Briefly, embryos were first equilibrated in hybridisation buffer (v3.1, Molecular Instruments) at 37°C for 30 min to block non-specific sites, then incubated overnight (12–16 h) at 37°C with 4 pmol of each probe set in fresh hybridisation buffer, following the manufacturer guidelines. Following this, unbound probes washed, and fluorophore-conjugated HCR hairpins (Molecular Instruments) were snap-cooled and applied to embryos in amplification buffer, then incubated for 12–16 h at room temperature in the dark. Post-amplification, embryos were washed to remove excess hairpins.

Finally, embryos were counterstained with DAPI (1:1000, Sigma-Aldrich) during the last wash, mounted in antifade medium (Vectashield; Vector Laboratories), and imaged by confocal microscopy and processed using a custom macro script in FIJI ([Embryo Processing HCR Script](#)).

Targeted Metabolomics

Embryos from seven consecutive developmental windows were rinsed thoroughly in distilled water to remove yeast, excess liquid discarded, weighed, and flash-frozen in liquid nitrogen. Samples were shipped on dry ice to the EMBL Metabolomics Core Facility (Heidelberg, Germany), where Bernhard Drotleff performed all downstream processing.

Each frozen sample (75 mg/mL) was extracted on dry ice by adding 80% acetonitrile (with internal standards and 1% formic acid), then homogenised using 1 mm zirconia beads in a bead-beater (FastPrep-24) at 6.0 m/s (5 × 30 s, 5 min rest). After incubation at –20 °C for 20 min, lysates were vortexed and centrifuged (15 000

× g, 10 min, 4 °C). Supernatants were transferred to glass vials and analysed within 2
h.

LC–MS/MS was carried out on an Agilent Infinity II BioLC coupled to a Sciex
QTRAP 6500+ (positive ESI). Chromatography used an Atlantis Premier BEH Z-HILIC
column (2.1 × 100 mm, 1.7 µm) at 40 °C, with mobile phases A (90:10 H_2O:ACN) and
B (98:2 ACN:H_2O), both buffered with 25 mM ammonium formate (pH 3.0). The 13
min gradient (0–1 min 100% B; 1–8 min to 60% B; 8–10 min 60% B; 10–10.5 min to
100% B; 10.5–13 min 100% B) ran at 0.3 mL/min, with 5 µL injections at 4 °C. Ion
source parameters were: curtain gas 40 psi, nebulizer gas 40 psi, heater gas 40 psi,
ion spray voltage 5 kV, source temperature 400 °C, medium collision gas. Data were
acquired in scheduled MRM mode (90 s windows, 600 ms cycle). Samples were
analysed in randomised order, with pooled QCs for system equilibration and every fifth
sample to monitor performance; a processed blank defined background. Raw peak
areas were extracted and processed in SciexOS. Data analysis and visualisation was
done in R using custom scripts (Metabolomics R Script).

scRNA-seq Analysis: Data alignment
Raw sequencing data were processed using Cell Ranger count (v9.0.1, 10X
Genomics). Reads were aligned to a custom *Drosophila melanogaster* reference
(Ensembl release BDGP6.46.110) into which the GFP transgene sequence had been
incorporated. Cell Ranger generated UMI-collapsed gene–cell count matrices for each
library.

scRNA-seq Analysis: Preprocessing and Quality Control
Downstream quality control and preprocessing were carried out in Seurat (v5.2.0; full
pipeline available at GitHub page). Briefly, cells with fewer than 800 or more than 100
000 UMIs, fewer than 200 or more than 8 000 detected genes, or mitochondrial
transcript content exceeding 8–10% were excluded. After log-normalisation and
identification of highly variable features, JackStraw analysis was used to select
significant principal components for dimensionality reduction. Uniform Manifold
Approximation and Projection (UMAP) was then performed on these PCs. To remove
artifactual multiplets, DoubletFinder (v2.0.6, McGinnis, Murrow and Gartner, 2019)
was applied with an expected doublet rate of 3.9%, consistent with 10X Genomics
loading recommendations.

scRNA-seq Analysis: Integration and Cluster Annotation

Biological replicates from each developmental time point were integrated using Harmony (v1.2.3, Korsunsky *et al.*, 2019) to correct batch effects while preserving biological variation. Clusters were initially defined in the integrated UMAP space and then annotated following a modified version of the Calderon *et al.* (2022) pipeline: marker genes for each cluster, selected as the top differentially expressed features with adjusted P < 0.05, were compared against curated spatial expression terms from the Berkeley Drosophila Genome Project (BDGP) using Fisher's exact test to identify over-represented *in situ* annotations (Tomancak *et al.*, 2002, 2007; Hammonds *et al.*, 2013; Calderon *et al.*, 2022). After Bonferroni correction, significant enrichment terms were manually reviewed and consolidated to assign each cluster to a defined neurogenic or neuronal identity. This combined data-driven and expert-curated approach ensured robust annotation of cell states across *Drosophila* embryonic neurogenesis.

To preserve the continuum of developmental variation, cells from all time points were merged without additional batch-correction; standard integration algorithms tended to over-correct and attenuate genuine developmental signals. The merged dataset was then reclustered in Seurat, and each resulting cluster was assigned an identity based on the modal annotation of its constituent cells from the individual time-point analyses. This majority-vote approach ensured that clusters reflected the predominant cell state across development.

scRNA-seq Analysis: Neuronal-Specific Annotation

We next focused on delineating specific neuronal lineages. Clusters corresponding to the progression from neuroblasts to mature neurons were reannotated using expression of subtype markers defined in Dillon *et al.* (2022) Peng *et al.* (2024). This approach yielded fine-grained, cell-type annotations that aligned with, and extended, the broader developmental ontology derived from BDGP enrichment annotation pipeline.

scRNA-seq Analysis: Trajectory Inference via RNA Velocity

To infer cellular dynamics and developmental trajectories, Cell Ranger BAM files were reprocessed with Velocyto (v0.17.17, La Manno *et al.*, 2018) to quantify spliced and

unspliced transcripts, producing Loom matrices. Next, the Seurat-annotated object
was converted to an AnnData structure using sccustomize (v3.0.1, Samuel Marsh *et*
al., 2024), and merged with the velocity matrices preserving metadata, embeddings,
and clustering labels. Downstream modeling was carried out in scVelo (v0.3.3, La
Manno *et al.*, 2018; Bergen *et al.*, 2020) using the dynamical model: data were first
filtered and normalised, moments were computed on the neighborhood graph, and
velocity vectors were estimated for each cell. Finally, velocity-stream embeddings
were projected onto the UMAP embedding to visualise inferred developmental
progressions.
650
scRNA-seq Analysis: Subclustering and Lineage Dynamic Analysis
To resolve finer neuronal subtypes, we isolated cells from the maturing neuronal
lineage and performed a second round of clustering in Seurat. Clusters were
reannotated based on the expression of established subtype marker genes, and dot
plots were generated to verify marker specificity. We then recalculated RNA-velocity
on this subset in scVelo (v0.3.3, La Manno *et al.*, 2018; Bergen *et al.*, 2020), re-
deriving the first- and second-order moments, velocity vectors, velocity-stream
embedding, and latent-time projections. These velocity maps were overlaid on the
UMAP to visualise intra-lineage transitions. Finally, CellRank was applied to the
subclustered dataset—integrating connectivity, velocity, and CytoTRACE kernels—to
infer both initial (progenitor) and terminal (mature neuron) fates and compute fate
probabilities within the neuronal lineage (Reuter *et al.*, 2018; Lange *et al.*, 2022;
Reuter, Klein and Lange, 2022; Weiler *et al.*, 2024).
664
scRNA-seq Analysis: Regulatory dynamics
To uncover TFs that drive subtype specification, we performed differential expression
analysis restricted to TFs annotated in the AnimalTFDB for *Drosophila melanogaster*
(Shen *et al.*, 2023). Candidate TFs with significant up-regulation in each neuronal
subtype were identified and ranked. Furthermore, leveraging the latent-time ordering
from scVelo, we modeled the dynamic expression of all genes along developmental
trajectories. We then refined this analysis using CellRank's multi-kernel framework—
combining connectivity, velocity, and CytoTRACE information—to pinpoint TFs whose
expression kinetics most strongly correlate with the latent time. This layered approach

highlighted key regulators of neuronal differentiation for subsequent validation using
HCR.
676
Comparative scRNAseq analysis: Dataset Acquisition and Preprocessing
To explore the evolutionary conservation of these regulators, we reanalysed a
published *Danio rerio* developmental single-cell RNA-seq dataset spanning 10
developmental stages, from 10 hours post-fertilisation (hpf) to 10 days post-fertilization
(dpf) (Lange *et al.*, 2024). Raw sequencing data were remapped using CellRanger
(v9.0.1, 10x Genomics) to the zebrafish reference genome (Danio_rerio.GRCz11.114,
Ensembl), ensuring retention of both spliced and unspliced reads. BAM files were
subsequently processed with Velocyto (v0.17.17, La Manno *et al.*, 2018) to generate
Loom files containing spliced/unspliced transcript counts.
686
Comparative scRNAseq analysis: Dataset Acquisition and Preprocessing
To explore the evolutionary conservation of these regulators, we reanalysed published
single-cell RNA-seq datasets from *Danio rerio* and *Strongylocentrotus purpuratus*. The
zebrafish atlas spans 10 developmental stages from 10 hours post-fertilisation (hpf) to
10 days post-fertilisation (dpf) (Lange *et al.*, 2024). The sea urchin dataset was
generated from combining two datasets, comprising 10 time points from the 8-cell
stage to the 72 hpf larval stage (Foster, Oulhen and Wessel, 2020; Perillo *et al.*, 2020).
For zebrafish, raw 10x Genomics sequencing data were reprocessed with Cell Ranger
(v9.0.1; 10x Genomics) against the Ensembl *D. rerio* reference
(Danio_rerio.GRCz11.114), retaining both spliced and unspliced reads. Resulting
BAM files were converted to spliced/unspliced count matrices using Velocyto
(v0.17.17, La Manno *et al.*, 2018), yielding loom files for downstream analysis. For sea
urchin, we used the author-provided count matrices and accompanying metadata
without remapping.
701
Comparative scRNAseq analysis: Data Integration and Annotation
Rather than reclustering the full datasets, we focused on neural lineages. For
zebrafish, central nervous system (CNS) populations were subset using the original
annotations provided in the source atlas (https://zebrahub.sf.czbiohub.org/data). For
sea urchin, we subset to neurogenic clusters based on the authors' labels.

Cross-species re-annotation was performed with orthologous marker genes
derived using eggNOG-based orthology mapping (described below), with the
Drosophila melanogaster subclustering marker set as the anchor. For sea urchin,
orthologues were called from the *S. purpuratus* proteome corresponding to the
reference genome used for mapping (GCF_000002235.4; Spur_4.2). Zebrafish
markers were referenced to Ensembl *D. rerio* GRCz11 (release 114). Monoaminergic
neuron populations were identified by expression of the orthologous canonical
biosynthetic enzymes and transporters.

Comparative scRNAseq analysis: Trajectory and Dynamic Expression Analyses
RNA velocity was estimated using scVelo (Bergen *et al.*, 2020) and lineage inference
was refined with CellRank (v2) (Weiler *et al.*, 2024) by integrating connectivity and
velocity information. Latent-time ordering was used to model dynamic gene expression
trajectories and identify TFs with temporal activation patterns correlating with neuronal
differentiation.

Comparative scRNAseq analysis: Differential Expression and Orthology
Mapping
Candidate TFs enriched in each neuronal subtype were identified using the same
thresholds applied to the *Drosophila* dataset (adjusted $p < 0.05$; $|log2FC| \geq 0.25$). For
cross-species comparison, we re-annotated the proteomes of zebrafish and
Drosophila with eggNOG (v5.0, Huerta-Cepas *et al.*, 2019), assigning genes to its
hierarchical orthogroup framework. We then conducted all comparative analyses at
the Bilateria-level orthogroups, which consolidate one-to-many and many-to-one
relationships, enabling robust mapping of putative orthologs and fair TF comparisons
across species.

Comparative scRNAseq analysis: Comparative and Class-specific Analyses
Shared and species-specific regulators were identified by comparing differentially
expressed TF orthogroups across *Drosophila* and zebrafish monoaminergic clusters.
To contextualise these findings, the orthogroup-level analysis was extended to other
neurotransmitter lineages (acetylcholine, GABA, glutamate).

Data availability

All data supporting this study are available on Zenodo. Processed FACS data are deposited at https://doi.org/10.5281/zenodo.17449471, targeted metabolomics data at https://doi.org/10.5281/zenodo.17449991, and processed single-cell RNA-seq data at https://doi.org/10.5281/zenodo.17441737. All datasets are released under the MIT License. Analysis pipelines and scripts are available at https://github.com/cliftonlewis/2025_Drosophila_scRNAseq_EmbryoNeurogenesis_ Monoamine.

Author Contributions

R.F. conceived the project. C.L. performed the majority of sample preparation for scRNA-seq, HCR, and targeted metabolomics, and carried out the bioinformatics analyses. M.G. provided guidance on the bioinformatics analyses. A.W. assisted with optimization of HCR protocols. N.C. and D.O. contributed to FACS sorting. S.R. supported the initial project setup. R.P.Z. provided the initial protocols for targeted scRNA-seq in *Drosophila* embryos. J.S. contributed to project development. E.R. and C.P.K. supervised and supported the project. R.F. was the primary supervisor. C.L. and R.F. wrote and edited the manuscript.

Acknowledgements

We gratefully acknowledge the support of the Royal Society through a University Research Fellowship (UF160226 and URF/R/221011) and a Research Grant (RGF\R1\181012) awarded to R.F. C.L. is supported by a BBSRC-MIBTP PhD Studentship (Grant number MIBTP2020: BB/T00746X/1). The authors wish to thank the technical and scientific staff of several core facilities for their invaluable support. We are especially grateful for the assistance received from the University of Leicester Advanced Imaging Facility (RRID: SCR_020967; BBSRC grant: BB/S019510/1) and specifically thank Dr. Kees Straatman for his expert microscopy advice. This research used the ALICE High Performance Computing Facility at the University of Leicester. We thank the Flow Cytometry Facility, particularly Dr. David Onion and Nicola Croxall, for their support with the Beckman Coulter MoFlo Astrios Flow Cytometer. We also thank the DEEP-SEQ Genomics Facility, specifically Dr. Christopher Moore and Dr.

Nadine Holmes, for their expert assistance with single-cell genomics, including 10X

Genomics support. Additionally, we are deeply appreciative of Dr. Bernhard Drotleff

from the EMBL Metabolomics Facility for running the targeted metabolomics

experiments. We would also like to thank Dr. Diego Calderon (University of California

San Francisco) for help with adapting the cell type annotation pipeline. Finally, we

would like to thank the Feuda Lab for their thoughtful proofreading and constructive

feedback on the manuscript.

References

Arendt, D. (2005) "Genes and homology in nervous system evolution: Comparing gene functions, expression patterns, and cell type molecular fingerprints," *Theory in Biosciences*, 124(2), pp. 185–197. Available at: https://doi.org/10.1016/j.thbio.2005.08.002.

Arendt, D. *et al.* (2019) "Evolution of neuronal types and families," *Current Opinion in Neurobiology*, 56, pp. 144–152. Available at: https://doi.org/10.1016/j.conb.2019.01.022.

Babski, H., Codianni, M. and Bhandawat, V. (2024) "Octopaminergic descending neurons in *Drosophila*: Connectivity, tonic activity and relation to locomotion," *Heliyon*, 10(9), p. e29952. Available at: https://doi.org/10.1016/j.heliyon.2024.e29952.

Bergen, V. *et al.* (2020) "Generalizing RNA velocity to transient cell states through dynamical modeling," *Nature Biotechnology*, 38(12), pp. 1408–1414. Available at: https://doi.org/10.1038/s41587-020-0591-3.

Busch, S. *et al.* (2009) "A map of octopaminergic neurons in the Drosophila brain," *The Journal of Comparative Neurology*, 513(6), pp. 643–667. Available at: https://doi.org/10.1002/cne.21966.

Cai, J. *et al.* (2009) "The role of Lmx1a in the differentiation of human embryonic stem cells into midbrain dopamine neurons in culture and after transplantation into a Parkinson's disease model," *Stem Cells (Dayton, Ohio)*, 27(1), pp. 220–229. Available at: https://doi.org/10.1634/stemcells.2008-0734.

Calderon, D. *et al.* (2022) "The continuum of *Drosophila* embryonic development at single-cell resolution," *Science*, 377(6606), p. eabn5800. Available at: https://doi.org/10.1126/science.abn5800.

Campos-Ortega, J.A. and Hartenstein, V. (1985) "Stages of Drosophila Embryogenesis," in Campos-Ortega, J. A. and Hartenstein, V., *The Embryonic Development of Drosophila melanogaster*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 9–84. Available at: https://doi.org/10.1007/978-3-662-02454-6_3.

Cheng, L. *et al.* (2003) "Lmx1b, Pet-1, and Nkx2.2 coordinately specify serotonergic neurotransmitter phenotype," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 23(31), pp. 9961–9967. Available at: https://doi.org/10.1523/JNEUROSCI.23-31-09961.2003.

Choi, H.M.T. *et al.* (2018) "Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust," *Development*, 145(12), p. dev165753. Available at: https://doi.org/10.1242/dev.165753.

Dillon, N. *et al.* (2022) "Single cell RNA-seq analysis reveals temporally-regulated and quiescence-regulated gene expression in Drosophila larval neuroblasts," *Neural Development*, 17(1), p. 7. Available at: https://doi.org/10.1186/s13064-022-00163-7.

Ding, Y.-Q. *et al.* (2003) "Lmx1b is essential for the development of serotonergic neurons," *Nature Neuroscience*, 6(9), pp. 933–938. Available at: https://doi.org/10.1038/nn1104.

Dittrich, R. *et al.* (1997) "The differentiation of the serotonergic neurons in the Drosophila ventral nerve cord depends on the combined function of the zinc finger proteins Eagle and Huckebein," *Development*, 124(13), pp. 2515–2525. Available at: https://doi.org/10.1242/dev.124.13.2515.

Doucet-Beaupré, H., Ang, S.-L. and Lévesque, M. (2015) "Cell fate determination, neuronal maintenance and disease state: The emerging role of transcription factors Lmx1a and Lmx1b," *FEBS Letters*, 589(24PartA), pp. 3727–3738. Available at: https://doi.org/10.1016/j.febslet.2015.10.020.

Flames, N. and Hobert, O. (2009) "Gene regulatory logic of dopamine neuron differentiation," *Nature*, 458(7240), pp. 885–889. Available at: https://doi.org/10.1038/nature07929.

Flames, N. and Hobert, O. (2011) "Transcriptional Control of the Terminal Fate of Monoaminergic Neurons," *Annual Review of Neuroscience*, 34(Volume 34, 2011), pp. 153–184. Available at: https://doi.org/10.1146/annurev-neuro-061010-113824.

Foster, S., Oulhen, N. and Wessel, G. (2020) "A single cell RNA sequencing resource for early sea urchin development," *Development*, 147(17), p. dev191528. Available at: https://doi.org/10.1242/dev.191528.

Goulty, M. *et al.* (2023) "The monoaminergic system is a bilaterian innovation," *Nature Communications*, 14(1), p. 3284. Available at: https://doi.org/10.1038/s41467-023-39030-2.

Goulty, M. *et al.* (2025) "Monoaminergic neurons share transcriptional identity across Bilaterian animals." bioRxiv, p. 2025.10.10.679534. Available at: https://doi.org/10.1101/2025.10.10.679534.

Grace, A.A. (2016) "Dysregulation of the dopamine system in the pathophysiology of schizophrenia and depression," *Nature Reviews. Neuroscience*, 17(8), pp. 524–532. Available at: https://doi.org/10.1038/nrn.2016.57.

Hammonds, A.S. *et al.* (2013) "Spatial expression of transcription factors in Drosophilaembryonic organ development," *Genome Biology*, 14(12), p. R140. Available at: https://doi.org/10.1186/gb-2013-14-12-r140.

Hartenstein, V. *et al.* (2017) "Developmental analysis of the dopamine-containing neurons of the Drosophila brain," *The Journal of comparative neurology*, 525(2), pp. 363–379. Available at: https://doi.org/10.1002/cne.24069.

Hendricks, T.J. *et al.* (2003) "Pet-1 ETS gene plays a critical role in 5-HT neuron development and is required for normal anxiety-like and aggressive behavior," *Neuron*, 37(2), pp. 233–247. Available at: https://doi.org/10.1016/s0896-6273(02)01167-4.

Hobert, O. (2008) "Regulatory logic of neuronal diversity: terminal selector genes
and selector motifs," *Proceedings of the National Academy of Sciences of the United*
States of America, 105(51), pp. 20067–20071. Available at:
https://doi.org/10.1073/pnas.0806070105.

Huerta-Cepas, J. *et al.* (2019) "eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502
viruses," *Nucleic Acids Research*, 47(D1), pp. D309–D314. Available at:
https://doi.org/10.1093/nar/gky1085.

Karki, S. *et al.* (2023) "Serotonin signaling regulates actomyosin contractility during
morphogenesis in evolutionarily divergent lineages," *Nature Communications*, 14(1),
p. 5547. Available at: https://doi.org/10.1038/s41467-023-41178-w.

Kasture, A.S. *et al.* (2018) "Big Lessons from Tiny Flies: Drosophila melanogaster as
a Model to Explore Dysfunction of Dopaminergic and Serotonergic Neurotransmitter
Systems," *International Journal of Molecular Sciences*, 19(6), p. 1788. Available at:
https://doi.org/10.3390/ijms19061788.

Korsunsky, I. *et al.* (2019) "Fast, sensitive and accurate integration of single-cell data
with Harmony," *Nature Methods*, 16(12), pp. 1289–1296. Available at:
https://doi.org/10.1038/s41592-019-0619-0.

Kurian, M.A. *et al.* (2011) "The monoamine neurotransmitter disorders: an expanding
range of neurological syndromes," *The Lancet. Neurology*, 10(8), pp. 721–733.
Available at: https://doi.org/10.1016/S1474-4422(11)70141-7.

La Manno, G. *et al.* (2018) "RNA velocity of single cells," *Nature*, 560(7719), pp.
494–498. Available at: https://doi.org/10.1038/s41586-018-0414-6.

Lange, M. *et al.* (2022) "CellRank for directed single-cell fate mapping," *Nature*
Methods, 19(2), pp. 159–170. Available at: https://doi.org/10.1038/s41592-021-
01346-6.

Lange, M. *et al.* (2024) "A multimodal zebrafish developmental atlas reveals the
state-transition dynamics of late-vertebrate pluripotent axial progenitors," *Cell*,
187(23), pp. 6742-6759.e17. Available at: https://doi.org/10.1016/j.cell.2024.09.047.

Libersat, F. and Pflueger, H.-J. (2004) "Monoamines and the Orchestration of
Behavior," *BioScience*, 54(1), pp. 17–25. Available at: https://doi.org/10.1641/0006-
3568(2004)054%255B0017:MATOOB%255D2.0.CO;2.

Lundell, M.J. and Hirsh, J. (1994) "Temporal and spatial development of serotonin
and dopamine neurons in the Drosophila CNS," *Developmental Biology*, 165(2), pp.
385–396. Available at: https://doi.org/10.1006/dbio.1994.1261.

McGinnis, C.S., Murrow, L.M. and Gartner, Z.J. (2019) "DoubletFinder: Doublet
Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors,"
Cell Systems, 8(4), pp. 329-337.e4. Available at:
https://doi.org/10.1016/j.cels.2019.03.003.

Monti, J.M. (2011) "Serotonin control of sleep-wake behavior," *Sleep Medicine*
Reviews, 15(4), pp. 269–281. Available at:
https://doi.org/10.1016/j.smrv.2010.11.003.

Montioli, R. and Borri Voltattorni, C. (2021) "Aromatic Amino Acid Decarboxylase
Deficiency: The Added Value of Biochemistry," *International Journal of Molecular*
Sciences, 22(6), p. 3146. Available at: https://doi.org/10.3390/ijms22063146.

Parent, A. (1984) "Functional Anatomy and Evolution of Monoaminergic Systems,"
American Zoologist, 24(3), pp. 783–790. Available at:
https://doi.org/10.1093/icb/24.3.783.

Peng, D. *et al.* (2024) "Organogenetic transcriptomes of the Drosophila embryo at
single cell resolution," *Development (Cambridge, England)*, 151(2), p. dev202097.
Available at: https://doi.org/10.1242/dev.202097.

Perillo, M. *et al.* (2020) "Regulation of dynamic pigment cell states at single-cell
resolution," *eLife*, 9, p. e60388. Available at: https://doi.org/10.7554/eLife.60388.

Peter, I.S. and Davidson, E.H. (2011) "Evolution of gene regulatory networks
controlling body plan development," *Cell*, 144(6), pp. 970–985. Available at:
https://doi.org/10.1016/j.cell.2011.02.017.

Rapti, G. (2022) "Open Frontiers in Neural Cell Type Investigations; Lessons From
Caenorhabditis elegans and Beyond, Toward a Multimodal Integration," *Frontiers in*
Neuroscience, 15. Available at: https://doi.org/10.3389/fnins.2021.787753.

Reuter, B. *et al.* (2018) "Generalized Markov State Modeling Method for
Nonequilibrium Biomolecular Dynamics: Exemplified on Amyloid β Conformational
Dynamics Driven by an Oscillating Electric Field," *Journal of Chemical Theory and*
Computation, 14(7), pp. 3579–3594. Available at:
https://doi.org/10.1021/acs.jctc.8b00079.

Reuter, B., Klein, M. and Lange, M. (2022) "pyGPCCA - python GPCCA: Generalized
Perron Cluster Cluster Analysis package to coarse-grain reversible and non-
reversible Markov state models." Zenodo. Available at:
https://doi.org/10.5281/zenodo.6914001.

Sachkova, M., Modepalli, V. and Kittelmann, M. (2025) "The Deep Evolutionary
Roots of the Nervous System," *Annual Review of Neuroscience*, 48(Volume 48,
2025), pp. 311–329. Available at: https://doi.org/10.1146/annurev-neuro-112723-
040945.

Samuel Marsh *et al.* (2024) "samuel-marsh/scCustomize: Version 3.0.1." Zenodo.
Available at: https://doi.org/10.5281/ZENODO.5706430.

Sarov, M. *et al.* (2016) "A genome-wide resource for the analysis of protein
localisation in Drosophila," *eLife*. Edited by H.J. Bellen, 5, p. e12068. Available at:
https://doi.org/10.7554/eLife.12068.

Schindelin, J. *et al.* (2012) "Fiji: an open-source platform for biological-image
analysis," *Nature Methods*, 9(7), pp. 676–682. Available at:
https://doi.org/10.1038/nmeth.2019.

Shen, W.-K. *et al.* (2023) "AnimalTFDB 4.0: a comprehensive animal transcription
factor database updated with variation and expression annotations," *Nucleic Acids*
Research, 51(D1), pp. D39–D45. Available at: https://doi.org/10.1093/nar/gkac907.

Tomancak, P. *et al.* (2002) "Systematic determination of patterns of gene expression
during Drosophila embryogenesis," *Genome Biology*, 3(12), p. research0088.1.
Available at: https://doi.org/10.1186/gb-2002-3-12-research0088.

Tomancak, P. *et al.* (2007) "Global analysis of patterns of gene expression during
Drosophilaembryogenesis," *Genome Biology*, 8(7), p. R145. Available at:
https://doi.org/10.1186/gb-2007-8-7-r145.

Volkow, N.D. *et al.* (2011) "Motivation deficit in ADHD is associated with dysfunction
of the dopamine reward pathway," *Molecular Psychiatry*, 16(11), pp. 1147–1154.
Available at: https://doi.org/10.1038/mp.2010.97.

Waddell, S. (2013) "Reinforcement signalling in Drosophila; dopamine does it all
after all," *Current Opinion in Neurobiology*, 23(3), pp. 324–329. Available at:
https://doi.org/10.1016/j.conb.2013.01.005.

Weiler, P. *et al.* (2024) "CellRank 2: unified fate mapping in multiview single-cell
data," *Nature Methods*, 21(7), pp. 1196–1205. Available at:
https://doi.org/10.1038/s41592-024-02303-9.

Wijnen, H. *et al.* (2006) "Control of Daily Transcript Oscillations in Drosophila by
Light and the Circadian Clock," *PLOS Genetics*, 2(3), p. e39. Available at:
https://doi.org/10.1371/journal.pgen.0020039.

Yamamoto, K. and Vernier, P. (2011) "The Evolution of Dopamine Systems in
Chordates," *Frontiers in Neuroanatomy*, 5. Available at:
https://doi.org/10.3389/fnana.2011.00021.

Zheng, Q. *et al.* (2025) "Bidirectional histone monoaminylation dynamics regulate
neural rhythmicity," *Nature*, 637(8047), pp. 974–982. Available at:
https://doi.org/10.1038/s41586-024-08371-3.

Figures

Figure 1. Time-resolved sampling strategy and developmental ontogeny annotation of targeted scRNA-seq across *Drosophila* embryogenesis.

A. Timeline of eight partially overlapping collection windows, spanning embryogenesis, from 0–22 hrs after egg laying (AEL) aligned to canonical embryonic stages (Bownes stages 1–17, (Campos-Ortega and Hartenstein, 1985)) and key morphogenetic landmarks (cleavage/blastoderm, gastrulation, germband elongation, germband retraction, dorsal closure, and differentiation). Major neurogenic milestones are indicated below the timeline, including neuroblast specification, diversification of sibling neuron fates, neuronal migration, neurite outgrowth, glial migration, and synaptogenesis. **B.** UMAPs for each time window showing single-cell transcriptomes coloured by developmental ontology. **C.** UMAP of the combined dataset with cells coloured by developmental ontology derived from *in situ*–guided annotation. Representative labels include maternal, cellular blastoderm, amnioserosa, dorsal ectoderm primordium, brain primordium, embryonic brain, and embryonic midgut.

Figure 2. Time-resolved single-cell landscape of embryonic neurogenesis in Drosophila

A. Neuronal annotation. UMAP of the full embryonic dataset with neural lineages highlighted and non-neuronal cells in grey. Colours denote major neural states: neuroblasts, neuroblasts/GMCs, neuroblasts/GMCs/immature neurons, intermediate neural progenitors (INPs), GMCs, new-born/immature neurons, mature neurons, and glia. **B. Neuronal subtype annotation.** Subclustered neurogenic atlas reannotated to subtypes including transmitter-class/subtype as well as neuronal developmental intermediates. Classes include acetylcholine, GABA, glutamate, monoamine, mixed identities (e.g., acetylcholine/GABA; GABA/glutamate), unknown mature neurons, and progenitor states (neuroblasts/GMCs; neuroblasts/GMCs/immature neurons; GMCs; new-born/immature neurons). **C. Timepoint.** Subclustered dataset coloured by collection window, showing temporal separation across the UMAP. **D. Markers.** Dot plot of selected marker genes across the neuronal states/subtypes in the subclustered dataset. Dot size indicates the fraction of cells expressing the gene; colour reflects average expression. **E. Differential transcription factors.** Dot plot of differential TFs across neuronal states/subtypes, highlighting regulators associated with progenitors, early differentiation, transmitter specification, and mature neurons.

A Monoaminergic Lineage - Dynamic Genes

Latent Time

B

1002

Figure 3. Monoaminergic lineage—dynamic gene programs and candidate transcription factors gene trends

A. Dynamic genes along latent time. Heatmap of CellRank-identified dynamical genes within the monoaminergic lineage. Cells are ordered by latent time (left→right; arrow) and expression is z-scaled per gene (yellow = higher expression). Many neurotransmitter-pathway genes and regulators turn on progressively toward late latent time as well as many of the candidate differentially expressed TFs identified

previously. **B. Monoaminergic transcription factors.** Smoothed expression trajectories across latent time for selected TFs (*fd59A*, *dmrt99B*, *CG4328*, *Lmx1a*, *Vsx1*, *Vsx2*). The monoaminergic lineage is highlighted in orange; other neuronal lineages (acetylcholine, GABA_1/2, glutamate) are shown for comparison. Curves show lineage-wise mean trends (y-axis, scaled expression) across latent time (x-axis), illustrating the preferential late-stage upregulation of these TFs in the monoaminergic lineage specifically.

1017

Figure 4. HCR validates the temporal onset and spatial organisation of serotonergic and dopaminergic programs in the embryo

HCR staining across four developmental windows (columns: stages 7–9, 10–12, 13–15, 16–17). Each image shows a whole-mount embryo, lateral view with anterior to the left. Scale bars 100 µm. **Serotonergic system**. HCR for *Trhn* (*tryptophan hydroxylase*, green), *SerT* (*serotonin transporter*, yellow) and *Vmat* (*vesicular monoamine transporter*, magenta), plus merges. No signal is detected at stages 7–9. From stage 10–12 onwards *Trhn*, *SerT* and *Vmat* appear in segmentally repeated CNS domains; marking serotonergic neurons as they mature. **Dopaminergic system**. HCR for *ple* (*Tyrosine hydroxylase*, green), *DAT* (*dopamine transporter*, yellow) and *Vmat* (magenta), plus merges. *ple* is first detected by stage 10–12, with *DAT* and *Vmat* also appearing by stage 10-12. *ple*/*DAT* signals co-localise with *Vmat* in the CNS. **Transcription factors**. HCR for *fd59A* or *dmrt99B* (orange) together with *Trhn* (green), plus merges. TF expression is evident in the CNS earlier (stages 7-9) and broadens by stages 10-12, preceding with enzyme onset, consistent with a putative upstream regulatory role in monoaminergic fate.

Figure 5. Targeted metabolomics across embryonic time reveals distinct onset of monoamine production

A–E. Concentrations of five monoamines measured by targeted LC–MS/MS across seven developmental windows (0–3, 3–6, 6–9, 9–12, 12–15, 15–18, 18–22 h AEL). Each dot is one biological replicate (≥6 per time point); colour encodes the time window. The black line connects the mean of replicates at each time. The y-axis shows $\log_{10}$(normalised peak area + 1). The red dashed line marks the empirical detection threshold for each analyte. **A. Serotonin** and **B. Dopamine:** low/near-baseline early, followed by a sharp rise beginning ~12–15 h, consistent with the transcriptional induction of biosynthetic enzymes. **C. Octopamine:** largely under the detection threshold. **D. Histamine** and **E. Tyramine**: detectable throughout development with gradual increases. Together, these profiles indicate that the functional availability of serotonin and dopamine lags their transcriptional commitment, whereas histamine and tyramine are present earlier and more continuously.

A

B

Monoaminergic TFs

Monoaminergic lineage

foxp4   lmx1ba   lmx1al   lmx1bb

C

Conserved TF Orthogroups

D. melanogaster	D. rerio
cic	cica
Nfl	nfia; nfixb
cbt	klf11b; klf11a
CG4328; Lmx1α	lmx1bb; lmx1al; lmx1ba
FoxP	foxp4
trh	NPAS3
Eip78C	nr1d1
net	atoh8
tj	mafa; mafb
Pdp1	hlfa
luna	klf6a

Monoamine

21 · 25 · 11 · 24 · 56

D. mel DE & Absent in D. rerio · D. mel DE & Present in D. rerio · DE_both · D. rerio DE & Present in D. mel · D. rerio DE & Absent in D. mel

Conserved TF Orthogroups

D. melanogaster	S. purpuratus
Vsx1;Vsx2	LOC575706
zfh1	LOC594820
CG4328;Lmx1α	LOC585635
Hr3	LOC578618
NFAT	LOC590859
Tusp	LOC591879
fd59A	LOC751845
Hr38	LOC581160
CG3104	LOC592680
Eip75B	LOC592902

Monoamine

23 · 24 · 10 · 36 · 11

D. mel DE & Absent in S. pur · D. mel DE & Present in S. pur · DE_both · S. pur DE & Present in D. mel · S. pur DE & Absent in D. mel

Figure 6. Cross-species conservation of monoaminergic TF programs

A. Differentially expressed TFs in *D. rerio* neurogenic subtypes. Dot plot of differentially expressed TFs across zebrafish neurogenic subtypes. Columns are grouped by transmitter/state classes (Acetylcholine, Glutamate, GABA, Serotonin, Dopamine, Monoamine; plus, developmental states such as Neuroblasts, GMCs, Immature/Mature neurons). Dot size shows the fraction of cells in the subtype expressing the TF; colour shows the z-scored average expression. **B. Monoaminergic TF dynamics.** Smoothed latent-time expression trajectories for TFs associated with monoaminergic fate (examples shown: *foxp4*, *lmx1ba*, *lmx1a1*, *lmx1a*). Curves are plotted for the major neuronal lineages; the monoaminergic lineage is highlighted in orange with a light-orange confidence band, illustrating late-latent-time up-regulation of these TFs along the monoaminergic trajectory. **C. Orthogroup-level conservation analysis.** Venn diagram summarising Bilateria eggNOG orthogroups containing TFs that are differentially expressed in monoaminergic clusters in *D. melanogaster* and *D. rerio* (criteria as in Methods: adjusted $p < 0.05$; $|\log2FC| \geq 0.25$). Left/right circles denote species-specific DE orthogroups; the intersection denotes orthogroups DE in both species. The table lists representative conserved TF orthogroups with the corresponding gene names in each species. Together, these panels show that a core set of TF orthogroups is shared across insects and vertebrates while each lineage also exhibits species-specific components of the monoaminergic program.