Developmental trajectories of cognitive traits in meerkats match socio-ecological demands

Tommaso Saccà[1,2], Elisa Protopapa[1,2], Adrian V. Jaeggi[3] & Sofia I. F. Forss[1,2]

[1]*Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland*

[2]*Kalahari Research Centre, Kuruman River Reserve, Northern Cape, South Africa*

[3]*Institute of Evolutionary Medicine, University of Zurich, Zurich, Switzerland*

Keywords: Heterochrony, life-history, ontogenetic patterns, inhibitory control, spatial memory and orientation, extractive foraging skills, individual variation.

Author contributions: T.S.: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Visualization, Writing – Original Draft. E.P.: Investigation, Data Curation, Writing – Review & Editing. A.V.J.: Methodology, Supervision, Writing – Review & Editing. S.I.F.F.: Conceptualization, Methodology, Supervision, Funding Acquisition, Writing – Review & Editing.

Abstract

Understanding cognitive development is fundamental for explaining variation in adult cognitive phenotypes, and thus the processes driving cognitive evolution. The ontogeny of cognitive traits is likely influenced by both population-wide pressures, such as ecological demands, and individual-specific factors, including early life experiences. To properly investigate cognitive variation, we must therefore identify species-level developmental trajectories and individual variation from normative ontogeny. We studied the ontogeny of three cognitive traits—inhibition control, spatial cognition, and physical problem-solving skills—in 28 wild meerkats (*Suricata suricatta*) from eight litters (seven groups). Our longitudinal study followed individuals from early life to nutritional independence and into sub-adulthood. We found that rates of development (learning curves) varied among traits, reflecting the timing of socio-ecological pressures in this species. Performance of cognitive traits did not correlate over ontogeny nor at specific time points, suggesting independent processes underlying each measured trait. While the development of inhibitory control appeared conserved, that of spatial cognition and physical problem-solving showed substantive individual differences. Furthermore, only physical problem-solving showed consistency in performances over time, emerging near nutritional independence, reflecting its ecological importance early in life. Future research should determine the drivers of individual developmental variation and how such differences translate into fitness consequences.

Introduction

Individual differences are increasingly relevant for explaining cognitive evolution (1). An important and underappreciated source of individual differences in cognition is developmental effects (2). Despite some recent work (3–5), this research is still in its infancy and studies investigating the ontogeny of cognitive traits remain especially sparse beyond primates and in wild conditions. Individual variation in cognitive developmental trajectories could facilitate adaptive responses when new selective pressures on existing phenotypes arise due to environmental variability (6–8). For example, given that advanced cognitive abilities may require larger brains, which are energetically costly (9) and thus trade-off with other functions such as reproduction (10,11), periods of food abundance may promote faster cognitive development while periods of food scarcity could select for slower developmental rates (12,13). When experienced during early life, ecological pressures such as diet (14), physical or social environment (3,15–17), have long lasting impacts on the adult cognitive phenotype. To explain how early-life pressures shape individual-level cognitive development, we need a clear account of population-level developmental trajectories. Crucially, the study of phylogenetic differences in developmental trajectories, known as heterochrony, has delivered important scientific advancements in our understanding of cognitive evolution (18–21). For example, comparative developmental studies of face recognition, gaze following, and goal-directed actions suggest that humans rely more heavily on social information than any of our closest primate relatives, with such differences emerging early in ontogeny (18,22). Furthermore, chimpanzees (*Pan troglodytes*) develop both spatial cognition and inhibitory control more rapidly than bonobos (*Pan paniscus*), which arguably reflects differences in their feeding ecology and social system (23,24).

Here we present a detailed, longitudinal investigation of early cognitive development in a wild social carnivorous mammal, the meerkat (*Suricata suricatta*). We studied meerkats throughout their ontogeny, from nutritional dependence to sub-adulthood, by experimentally measuring the developmental trajectories of three distinct physical cognitive skills: inhibitory control, spatial cognition, and extractive foraging represented through a physical problem-solving task (Figure 1). These cognitive traits reflect pressures experienced by meerkats in their ecological and social environment. Meerkats are highly social, cooperatively-breeding mammals in which adults and sub-adults contribute to raising offspring, defending against predators and outgroup conspecifics, and burrow maintenance (25). Cooperative breeding has been proposed to select for enhanced executive functions, including inhibitory control, because individuals must supress self-serving behaviours to engage in cooperative activities (26,27), but see (28). Moreover, meerkats live in environments characterised by high predation pressure and limited physical refuges (i.e., boltholes) from aerial predators (29). This makes the development of spatial cognition particularly important as immature individuals must learn proper, adult-like responses to alarm calls (30) and acquire adult-level knowledge of bolthole locations (31). Finally, the development of physical problem-solving is ecologically relevant because meerkats inhabit a complex foraging niche characterised by seasonality and habitat heterogeneity (32,33), and immature individuals rely on adults to learn foraging skills, define their food repertoire and acquire complex prey-handling skills (34–36). We

would also expect meerkat life-history to influence the cognitive performance of males and females. In meerkats, where reproduction is largely monopolised by dominant individuals (29), females are philopatric and can maximise fitness by attaining dominance within their birth group (37). Males in contrast must disperse to secure a dominance position, and as subordinates most of their reproductive success comes from prospecting and gaining extra-group copulations (29,38,39). Dispersal is risky due to predators and aggression from neighbouring groups (29), and its success likely involves the ability to navigate unfamiliar environments (spatial cognition), to efficiently locate food (physical problem-solving), and to prioritize effective decisions over prepotent responses (inhibition control) in this dangerous, highly arousing context. Overall, success in dispersal events requires effective decision making under limited cues and has been proposed to contribute to socially driven cognitive evolution (40).

Altogether, meerkat pups depend on adults to develop essential survival skills, such as foraging and anti-predator behaviours (41). Ontogenetic reliance on social inputs has been linked to predictions about domain-general cognition (i.e. correlated performance across cognitive traits). Extensive reliance on social learning has been argued to favour the emergence of human-like domain-general intelligence, by enhancing learning efficiency and capacity during development which translate across cognitive domains (the Cultural Intelligence Hypothesis) (42). On the other hand, strong reliance on social learning may reduce selection pressure on independent, generalized learning abilities, and avoid the associated energetic costs of large brains (43). In addition, it has been proposed that cooperative breeding systems present unique challenges that select for enhanced socio-cognitive abilities (the Cooperative Breeding Hypothesis) (26,44,45), as well as reducing energetic constrains on larger brains (46). Therefore, the social ecology and cooperative breeding system of meerkats provide a unique opportunity to evaluate hypotheses on the evolution of cognition in a non-primate mammalian species.

In this study we repeatedly assessed individual meerkat pups over multiple key developmental life stages (Figure 1). Our longitudinal approach enabled us to ask I) whether distinct cognitive traits develop in parallel within individuals, II) whether relationships among traits exist and whether these change over ontogeny, and lastly III) whether performance was consistent across developmental stages. Finally, we discuss population-level developmental trajectories for each trait in the context of species-specific socio-ecological demands.

Meerkat development and test timings

Birth · At burrow w/ babysitter(s) · 4 · Join group for foraging · 7th · 10th · 13th Nutritional Independence · 18th · 24th Sub-adulthood

Data collection · Week of age

Figure 1: Overview of meerkat early life development and data collection time points. Weeks of age are indicated as black ticks, testing weeks as orange rectangles, and key life history stages with longer ticks (see Methods for details).

Methods

Study population

We conducted our study at the Kalahari Research Centre, located in the Kuruman River Reserve (KRR) (26°59′ S, 21°50′ E) in the Kalahari Desert in South Africa. The KRR is a semi-arid savanna with distinct macro-habitats such as red sand dune and shrub grassland (33). Vegetation is sparse during the dry season and more abundant during the wet season (47). Rainfall averages 266mm per year, and temperatures are on average 30.3 °C during the wet season (maximum 44.2 °C) and 10.1 °C during the dry season (minimum -11.9 °C) (33). Here, we investigated 28 wild meerkats (12 females and 16 males), from seven groups and eight litters with average 3.5 pups (range 2 to 4) over two wet Seasons (December 2022 – May 2023, and November 2023 – May 2024). All groups and individuals included in our study were habituated to close observation.

The longitudinal developmental approach

For the first three weeks of life, meerkat pups stay below ground, after which they emerge and remain close to the natal burrow. During this time, they are always joined by usually at least one older group member to ensure their safety (48). At about four weeks of age, they start to leave the natal burrow and join their group foraging. They rely on adults for food acquisition until about 12 weeks of age, after which pups are considered nutritionally independent (49). They are considered sub-adult at 24 weeks of age, when they first start to participate in all cooperative activities, and adults at one year of age after reaching physiological reproductive maturity (50). We tested individuals on their cognitive performance at five key points during their development (see Figure 1). The earliest point was the 7[th] week of life, as this was when meerkat pups would first show interest in interacting with cognitive tests. At this time pups had left the safety of their natal burrow and were consistently joining their group foraging already for about 2-3 weeks. We re-tested all pups at 10, 13, 18 and 24 weeks to evenly space sampling effort around nutritional independence, with two measurements before, one during, and two after, with the final point corresponding to early sub-adulthood. Testing weeks were based on supposed date of birth and we allowed on average a two-day flexibility for testing individuals before and after the end of the formal week for the 7[th], 10[th] and 13[th] week, and four-day flexibility for the 18[th] and 24[th] week, to reduce logistical problems in data collection. At times, we could not test individuals due to external limitations or to individuals not being interested in the test apparatus. In other instances, we excluded single tests post hoc for not meeting sufficient test criteria. A complete list of deviations from planned data collection can be found in Supplementary Table S1. Differences in the number of presentations across developmental weeks were accounted for in the analysis.

Test paradigms of cognitive traits

Inhibition control task

To test for the ability to inhibit incorrect intuitive responses, we used a standardized inhibitory control task (sometimes referred to as the "cylinder test" (51)), which in our study consisted of a rectangular box with open ends containing a food reward in the middle. Prior to testing, at each developmental week, subjects were habituated to an opaque version of the test apparatus that was presented with the opaque barrier facing the subject. Before testing, we ensured that each subject had learnt to disregard the barrier and access the reward from the open sides, after which they were considered ready to be tested on their ability to recall the learned response and inhibit prepotent but incorrect responses. Detailed information about the habituation procedure can be found in Supplementary Text S2. During testing, we presented five times per developmental week to each subject a transparent version of the apparatus with the transparent barrier facing the subject (Figure 2). We considered a trial successful if the subject did not touch the transparent side before going around the open side and accessing the food reward.

Spatial cognition task

We evaluated spatial cognition using a four-armed maze with a central chamber and a starting box at the end of one arm. The starting box, the central chamber and all arms were fully opaque (Figure 2), while the top of the central chamber and the ends of each arm varied between data collection seasons (transparent in the first season and opaque in the second). This change was done to further minimize subjects' possible reliance on external cues that could not be kept constant between trials. Subjects were habituated to the apparatus each developmental week before testing. Additional details about habituation can be found in Supplementary Text S2. During testing, three exits were covered by plexiglass, while the fourth was always open and connected to the starting box. One of the covered exits could be opened, constituting the correct exit, while the others two were fastened shut. During testing, we assessed stress levels by monitoring characteristic lost calls (52) and other vocalizations. If subjects showed even mild signs of distress, judged from a combination of vocalizations and physical cues of arousal, we guided them to the correct exit by inserting our hand there and providing visual and auditory cues. We considered the first successful trial as the first trial where the individual successfully found the correct exit without external help. From then on, we tested each subject in three more trials. In those instances, if the subject became too stressed, we removed it from the test without guiding it to the correct exit and we considered the trial a mistrial.

The design of the spatial cognition task slightly differed between the first and second season of data collection. During the first season the plastic ends of the three exits and the top of the central chamber were made of transparent plexiglass. In addition, the correct exit for each subject could either be the same as the previous week or change. During the seventh week of the first study group, the main chamber of the maze was fully transparent. This data point was considered as the same to the datapoints from the first season, for sample size limitations on this design. During the second season the plastic end of the three exits were opaque, and the correct exit for each subject always changed between successive weeks, to avoid a purely associative learning effect.

Physical problem-solving task

To test the ability to solve physical problems to access a food reward, we presented an apparatus with four chambers, each containing a food reward inside (see Figure 2). Three chambers were covered by a different obstacle (henceforth: 'problem') to be removed or opened, with one chamber left unobstructed to maintain motivation to participate. Each problem could be solved with physical motions routinely used while foraging, specifically digging, poking, and pulling. Before each week of testing subjects were habituated to the test apparatus to avoid any potential effects linked to neophobia and approaching of novel human-made artefacts (53). Further details about habituation procedure and apparatus design can be found in Supplementary Text S2. During testing, we presented the test apparatus to each subject four times during each developmental week. We considered a problem solved if the subject had found a way to access and eat the food reward inside, even if we could not infer whether the action used was intended to solve the problem. We did not consider a chamber solved if the problem was opened by accident by an involuntary bump or any other sort of non-active solution.

Role of memory: To investigate whether the ontogeny of performance in the problem-solving task was primarily influenced by learning and memory from repeated exposure over development, we presented the apparatus to an additional six subjects (four males, two females) at ten weeks of age who had not encountered it before at seven weeks of age and recorded their performance. We hypothesised that, if the increase in performance was mainly due to memory and prior exposure, these six subjects would (a) perform similarly to subjects first presented with the apparatus at seven weeks of age, and (b) perform worse than those same subjects when the latter were re-presented with the apparatus at ten weeks of age.

Data scoring and statistical analyses

We recorded each presentation using video cameras (Hero 9 Black: GoPro Inc., 2020), and we scored cognitive task performances using the software Boris (54) and a predefined ethogram (Supplementary Table S3). For the spatial cognition test, to account for the effect of potential stress on performance, we noted as a binary variable for each trial if the subject produced any "lost call" vocalization during testing. For the physical problem-solving test, to account for motivational levels, which is known to influence cognitive performance (55), we scored the time the subject was interacting with or investigating the test apparatus. In addition, to account for differences in study design, we noted down whether the design was linked to the first or second season of testing. All analyses were conducted R v4.4.0 (56), using RStudio as an interface (57). For Bayesian hierarchical and multivariate model fitting we used the package BRMS (58). Throughout our analyses, we used weakly informative priors (intercepts: normal(0,2); fixed effects: normal(0,2); standard deviations of random effects: exponential(2); correlations of random effects: LKJ(2)) to aid model convergence and regularize parameter estimates, thus reducing type I errors (59). For all models, we confirmed proper model fitting by visual inspection of chain mixing, Rhat values for convergence (all = 1.00) and effective sample size measures (all > 1000).

Population-level developmental trajectories and individual variation

To investigate individual performance over development for each cognitive trait we fitted three

sets of Generalized Linear Mixed Models (GLMM) in a Bayesian framework, one for each cognitive task.

We added to each of the three models a random individual intercept and a random individual slope for the

effect of developmental week, treated as a continuous variable (henceforth: week). This allowed the

models to represent the developmental trend of each subject, accounting for both individual baseline

performance after adjusting for the effect of covariates (individual random intercept), and individual

differences in developmental rates (individual random slope for the effect of week on task performance).

These random effects were estimated as individual deviations (positive or negative) relative to,

respectively, the population-level performance (global intercept) and developmental trend (fixed effect of

week). This structure also enabled us to investigate to which extent performances could be attributed to

individual characteristics. To do so, we calculated the proportion of variance explained by the random

intercepts and slopes not explained by fixed effects using the function "variance_decomposition" from the

package Performance (60). This is the suggested approach when working with non-Guassian distribution

and complex random structures in Bayesian models fitted with BRMS package. This approach is

functionally equivalent to the widely used "adjusted intraclass correlation coefficient" (adjusted ICC).

Inhibition control - For the performance in the inhibitory control task we used as the response variable the

number of trials passed versus the number of total trials per week of testing, assuming a binomial

distribution, and we added as explanatory variable the effect of week and of sex.

Spatial cognition - For the performance in the spatial cognition task, we fitted two models, to investigate

competing variables linked to study design. First, we used the total amount of errors over the number of

trials per week of testing as a response variable, assuming a Poisson distribution, including as covariates

(a) the number of trials per week where the subject showed mild signs of distress, (b) week, (c) sex, and

(d) a binary variable for whether the correct exit was kept the same as for the previous week of testing.

Secondly, we fitted a model where all variables were equivalent, but we excluded the binary variable for

whether the correct exit changed from the previous week. Instead, we fitted a variable for study design,

distinguishing between first and second season. In both models, for the total amount of errors and

number of trials we excluded the performance in the first trial per week, since we could not assume the

subjects knew the solution and the performance was more due to chance than actual cognitive

processing.

Stress and season of testing - We investigated whether the physical design of the spatial

cognition test (i.e., ceiling and arm openings transparent during the first season and opaque during the

second season) influenced the level of stress experienced by subjects. We fitted a Generalized Linear

Model (GLM) with the number of trials per subject and week in which mild signs of stress were recorded

as the response variable, and season of testing included as an exploratory categorical predictor,

assuming a Poisson distribution for the response.

Physical problem-solving - For the performance in the physical problem-solving task we used the number

of problems solved over the number of problems available at each developmental timepoint as the

response variable, assuming a binomial distribution, and as explanatory variables we used (a) the

average time of exploration of the test apparatus for all trials during each week, (b) sex, and (c) week.

Role of memory - To compare the performance of subjects first presented with the problem-
solving task at ten weeks of age with that of subjects first presented at seven weeks and re-presented at
ten weeks, we fitted a GLMM with the proportion of problems solved out of those available as the
response, assuming binomial distribution. Explanatory variables included sex, the average time spent
investigating the apparatus across all trials at each developmental time point, and week as a factor with
three levels according to the experimental condition: (1) first presentation at seven weeks of age, (2) first
presentation at ten weeks of age, and (3) second presentation at ten weeks of age. We included
individual identity as a random factor.

Development of individual consistency in task performance

To investigate whether individual consistency changed over the course of early life, we used a
multivariate hierarchical modelling approach, using the brms package (61). For each cognitive test, we
fitted 5 Generalized Linear Models, one for each week of testing. We fitted the same set of fixed
response, explanatory variables and assumed distribution of the models used to investigate individual
performance over development, with the exclusion of week of age. For the random part, we allowed the
intercept calculation to vary for individual by fitting individual random intercepts. Then, for each cognitive
test, we used the multivariate hierarchical design to correlate individual intercepts calculated at each
week with each other. Random individual intercepts were calculated as deviation from the global
intercept, which for each model was the specific week investigated. Thus, we obtained within each week
a set of individual values that highlighted whether the subject performed above or below average, while
factoring out the covariates of interests that could confound cognitive performance and create spurious
correlations. Thus, comparing individual intercepts among weeks allowed us to investigate whether
subjects that performed above average at one point in time (e.g. week 10) were more likely to perform
above average at a future developmental stage (e.g. week 13).

Correlation among task performance of the three cognitive traits

Over the developmental period: To investigate whether performance in one cognitive task was related to
performance in another, we extended our hierarchical models into a multivariate framework. We refitted
the models described in the section "*Population-level developmental trajectories and individual variation*",
retaining for the spatial cognition task the version including study design as a predictor, as it
outperformed the alternative model with the binary 'correct exit changed' variable. This multivariate
approach allowed us to address two questions: (1) whether subjects who performed better than average
in one task also performed better in another, assessed by comparing individual intercepts across
tasks/models; and (2) whether subjects who improved more rapidly than average in one task showed
similar improvement in another, assessed by comparing individual slopes for the effect of week on
performance across tasks/models.

At specific developmental time points: To investigate whether relationships between performance in any
pair of cognitive tasks changed over the course of development, we again used a multivariate hierarchical
modelling approach. For each week of testing, we fitted three models (one for each cognitive task) using

the same specification (response, predictors, random effects, and assumed distributions) as described in

the section "*Development of individual consistency in task performance*". This framework allowed us to

assess, for each week, whether subjects performed above or below average after accounting for

confounding variables (individual intercepts). By comparing performance across tasks, we then

determined whether relationships between cognitive traits varied across early life.

Figure 2: Population-level developmental trajectories (second column) and individual ontogenetic variability (third column) for three cognitive tasks (first column). Vertical lines indicate nutritional independence and sub-adulthood. In all plots, age (developmental week) is on the x-axis and task performance on the y-axis. Population-level trajectories: black line shows the modelled population mean, green shading indicates the 95% confidence interval, and observed data are plotted in the background. Data points are semi-transparent, with darker shades indicating overlapping points, and jittered along the x-axis for clarity. Individual variability: lines show modelled trajectories for each individual, with shaded 95% confidence intervals; darker shading indicates overlapping intervals. For population-level spatial cognition, predictions average across seasons, and all population-level trends are averaged across sexes.

Results

321   We report model estimates as the mean with the 95% credible interval in square brackets. As a

322 measure of evidence, we calculated the proportion of posterior samples supporting the estimated effect

323 ($PP_{support}$). We encourage readers to treat $PP_{support}$ as a continuous measure of evidence without clear

boundaries, and to make their own decision on sufficient levels of credibility. However, to ease interpretation and discussion, we considered effects with $PP_{support} \geq 95\%$ as supported, and effects with $PP_{support}$ of between 70% and 95% as suggestive but inconclusive trends (62). We report the proportion of remaining variation (i.e., not explained by fixed effects) explained by random effects as the median with the 95% credible interval in square brackets. Although negative values for this proportion are numerically impossible, they can appear as artefacts of high variability in the posterior distribution (60). When the lower bound of the 95% credible interval extended below zero, we truncated it at zero and interpreted the effect as not being convincingly different from zero.

Population-level developmental trajectories and individual variation

Inhibition control – We found support for an increase in performance in the inhibitory control task over successive developmental week (mean effect of week = 0.09, 95% CI [0.05, 0.13], $PP_{support}$ = 100%, Figure 2). There was no evidence that males and females differed in their probability of passing the inhibitory control test (effect of males, relative to females = 0.08, [-0.55, 0.74], $PP_{support}$ = 60.6%).
Individual differences in learning curves (random slopes) accounted for 11% [0%, 45%] of the remaining variability that was not explained by fixed effects (henceforth: 'remaining variability'), while individual differences in baseline performance (random intercepts) accounted for 28% [0%, 56%], and the combination of the two explained 18% [0%, 44%]. Thus, we find little evidence for individual deviation from population-level developmental trends in this cognitive task. Individual baseline performance without the effect of covariates (random intercepts) and learning curves (random slopes) showed a non-supported trend towards a negative correlation (correlation = -0.46, [-0.95, 0.62], $PP_{support}$= 83.0%).

Spatial cognition – We found support for increasing performance in the spatial cognition task with successive developmental week, with subjects making less errors later in their development (week = -0.04, [-0.07, -0.01], $PP_{support}$ = 99.8%, Figure 2). Performance was also influenced by stress, with subjects making more errors when showing mild signs of stress (effect of stress = 0.36, [0.22, 0.50], $PP_{support}$ = 100%). We found no support for a difference between males and females in the average number of errors per trial (effect of male, relative to females = -0.06, [-0.64, 0.53], $PP_{support}$ = 58.1%). When the correct exit was kept the same from the previous testing week, we found a non-supported trend in favour of subjects making less errors when the exit was consistent among consecutive weeks (effect of consistent exit, relative to inconsistent = -0.09, [-0.38, 0.19], $PP_{support}$= 72.6%). Test design influenced performance, as we found that subjects made less errors during the second season (effect of second season, relative to first season = -1.01, [-1.73, -0.22], $PP_{support}$= 99.2%), while at the same time more likely to show mild signs of stress (second season, relative to first season = 1.25, [0.02, 2.76], $PP_{support}$= 97.6%). Accordingly, we used the model including the study design to calculate the proportion of remaining variance explained by random factors, to factor out possible spurious individual effects due to the two seasons.

Including individual learning curves (random slopes) in our model for the development of spatial cognition accounted for 72% [32%, 91%] of remaining variation, while including individual baseline performances (random intercepts) accounted for 80% [24%, 95%] of it. Including the combination of individual learning curves and baseline performance in the model accounted for 51% [0%, 81%] of remaining variability. Thus, we find strong support for individual differences in the development of

performance in the spatial cognition task in both baseline abilities and learning curves. The proportion of remaining variability was lower when random slopes and random intercepts were included together, possibly due to the covariance between the two factors (correlation = -0.69, [-0.92, -0.26], PP$_{support}$= 99.6%). This suggests that when analysed separately, each factor also captures part of the variability explained by the other. The negative covariance between random slopes and intercepts indicates that individuals with above-average baseline performance (i.e., making more errors after accounting for covariates) tended to show below-average rates of developmental change (i.e., a faster decrease in errors over time). Thus, subjects that performed worse at a baseline showed faster learning curves during development.

Physical problem-solving - The performance in the physical problem-solving task increased over ontogeny, with the effect of developmental week having a positive effect (week = 0.26, [0.17 - 0.36], PP$_{support}$= 100%, Figure 2). Subjects that spent more time exploring and interacting with the apparatus were more likely to solve the problems it posed (exploration time = 0.03, [0.01, 0.05], PP$_{support}$ = 100%). We found a suggestive trend for males being less likely to remove an obstacle than females, though we did not find full support for this finding (males, relative to females = -0.42, [-1.35, 0.51], PP$_{support}$ = 82.3%). Individual learning curves (random slopes) accounted for 60% [43%, 72%] of the remaining variability, while individual baseline performances (random intercepts) accounted for 44% [12%, 66%] of it. The combination of random slopes and intercepts accounted for 44% [21%, 63%] of remaining variability. Thus, we find moderate to medium to high levels of support for differences in both individual baseline performances and individual rates of development (learning curves). The decrease in variability explained when both random slopes and intercepts were included is possibly due to the two random factors covarying with each other (correlation = -0.65, [-0.88, -0.25], PP$_{support}$= 99.6%) and explaining in isolation part of the other factor's explained variability. In addition, this correlation highlights that subjects with poorer baseline performance showed faster learning curves during development.

Role of memory - Subjects who were presented with the physical problem-solving task for the first time at seven weeks of age performed worse than subjects who were presented with it for the first time at ten weeks of age (first presentation at seven weeks, relative to first presentation at ten weeks = -1.07, [-2.17, 0.02], PP$_{support}$ = 97.2%). We did not find support for a difference in performance between subjects who interacted with the apparatus for the first time at ten weeks and those that interacted with it for the second time at ten weeks, although there was a non-supported trend for the latter outperforming the former (second presentation at ten week, relative to first presentation at ten week = 0.35, [-0.75, 1.48], PP$_{support}$ = 73.4%).

Development of individual consistency in task performance

Detailed model outputs can be found in Supplementary Table S4.

Inhibitory control – We did not find support for individual performance at one developmental time point to predict performance at the subsequent time point (Figure 3). A non-supported positive trend was observed between week 7 and week 10 which disappeared in later comparisons. Likewise, performance at the onset of sub-adulthood (week 24) was not predicted by any earlier time point (Figure 3).

Spatial cognition – We did not find support for consistency in individual performance across developmental time points (Figure 3), although correlation estimates were consistently positive. A non-supported positive trend was observed between week 7 and week 10 and between week 13 and week 18, while support for the same correlation at week 10–13 was even weaker. Likewise, performance at the onset of sub-adulthood (week 24) could not be convincingly predicted by earlier time points.

Physical problem-solving – We found strong support for consistency in individual performance across consecutive developmental weeks, emerging early in ontogeny (Figure 3). A non-support positive trend was present between week 7 and week 10, which became supported in week comparisons from week 10 onwards. Reflecting this pattern, performance at the onset of sub-adulthood (week 24) was predicted by performance at week 18, 13 and 10, while at the earliest stage (week 7) the relationship remained a non-supported positive trend.

Correlation among task performance of the three cognitive traits over the developmental period

Because performance metrics were scaled in opposite directions (higher values indicated better performance in inhibitory control and physical problem-solving, while lower values indicated better performance in spatial cognition) we mirrored the correlation estimates involving spatial cognition. Accordingly, positive correlations between task performances consistently indicate that better performance in one task was associated with better performance in the other.

Inhibitory control & spatial cognition – We found a non-supported trend between the rates of development of the two traits, calculated as the random slope of the effect of week on performance for each subject, with faster learning curves in inhibitory control performance associated with slower learning curves in spatial cognition performance (correlation = -0.21, [-0.83, 0.43], $PP_{support}$ = 73.7%). When comparing individual baseline performances after removing the effects of covariates (random intercepts), we found another non- supported trend, with a better performance in inhibitory control associated with a worse performance in spatial cognition (correlation = -0.36, [–0.81, 0.13], $PP_{support}$ = 91%).

Spatial cognition & physical problem solving – No supported correlation was detected between developmental rates (correlation = -0.02, [–0.47, 0.41], $PP_{support}$ = 52.6%). At the baseline performance level, we observed a non- supported trend, with a better performance in the physical cognition task associated with a worse performance in the spatial cognition task (correlation = -0.27, [–0.70, 0.16], $PP_{support}$ = 88.4%).

Inhibitory control & physical problem solving – We found non-supported positive trends for both developmental rates (correlation = 0.21, [–0.43, 0.80], $PP_{support}$ = 75.7%) and baseline individual performance (correlation = 0.22, [–0.30, 0.72], $PP_{support}$ = 78.8%), with better inhibitory control performance associated with better physical problem-solving performance.

Correlation among task performance of the three cognitive traits at specific developmental time points

Similarly to the previous section, we mirrored the correlation estimates involving spatial cognition so that
positive correlations between task performances consistently indicate better performance in one task
being associated with better performance in another. Detailed model outputs can be found in
Supplementary Table S5.

Inhibitory control & spatial cognition – We found no supported associations between inhibitory control and
spatial cognition at any time point (Figure 3). A non-supported negative trend was present at 7 weeks of
age.

Spatial cognition & physical problem solving – Associations between spatial cognition and physical
problem solving also lacked support at all time points (Figure 3). Estimates were consistently positive
across development, with a non-supported positive trend at weeks 10 and 13.

Inhibitory control & physical problem solving – No supported associations or unsupported trends were
detected between inhibitory control and physical problem solving at any time point (Figure 3).

Figure 3: Summary of correlations between individual random intercepts among tasks, and between
tasks, over the developmental period. Full, red lines are positive, supported correlations, while dotted
lines are non-supported correlations.

Discussion

In this study, we investigated the developmental trajectories of three ecologically relevant
cognitive traits in wild meerkats. We found evidence that cognitive performance increases in meerkats
over ontogeny, spanning from early life to early sub-adulthood. Cognitive development was present for all
measured traits: inhibition control, spatial cognition and physical problem-solving skills. At the population
level, inhibitory control and spatial cognition (spatial orientation and memory) showed a linear increase in
performance and did not reach a plateau within the developmental timeframe investigated, while physical
problem-solving skills showed the fastest increase in performance (steepest learning curves) and

reached a plateau before subjects grew into sub-adulthood (Figure 2). We found substantial evidence for
individual variation in the development of both spatial cognition and physical problem-solving ability but
not in the development of inhibitory control, suggesting the latter to be under developmental constrain. Of
all cognitive traits, we found support for individual consistency in performance only in physical problem-
solving ability, which arose early and endured until the end of the developmental period studied. Contrary
to our expectations based on male dispersal and the cognitive requirements of interactions with outgroup
conspecifics, we did not find evidence for differences in cognitive performance between males and
females in any of the three traits investigated. Furthermore, although we expected cognitive performance
to positively covary in meerkats due to their extensive reliance on social learning during development
(42), we found no support for such covariance during early development, either in individual trajectories
or at discrete time points.

The timing of cognitive development in each trait can be linked to specific socio-ecological
demands in meerkats. Early in their ontogeny meerkats must reach sufficient independent foraging
competence, as social provisioning by adults stops around the twelfth week of age. This timing aligns with
our findings on the cognitive ability to solve extractive physical problems, where meerkats had a 76% (CI
95% [57%, 89%]) probability of solving a physical barrier assuming 30 seconds of exploration at 12
weeks of age. Despite an early onset of independent foraging ability (50), meerkats remain within their
family groups way beyond when their foraging competence is reached (25) and can rely on social
information and cues from their group members for spatial navigation and coordination for a longer
period. Before nutritional independence, pups can react to alarm calls by following the heading of nearby
older individuals to reach safety (30,41). Afterwards, decisions about group movement are mostly
influenced by older individuals (63). Accordingly, our findings suggest that the development of spatial
cognition is slower and lacks a clear plateau within the developmental timeframe we investigated (Figure
2), suggesting that this cognitive trait is less critical early in life. Since investment in neurological tissue is
expensive (9,11,44,64), partly relying on collective information may be an energetically efficient strategy
(65), especially early in life when demands on independent spatial memory are lower. Similarly, there are
lower demands for inhibitory control early in meerkat life, as juveniles are seen to contribute less to
cooperative activities than older individuals (25). Selection may even favour low inhibitory control early in
pup development, as adults react to signals of hunger and feed pups that beg more intensively and
persistently more often (66,67). However, these benefits progressively decline as pups grow, because
adults stop responding to their begging calls when they reach three to four months of age (68), and adults
may need to reach a minimum baseline of inhibition ability for effective cooperative activities. Accordingly,
we found that inhibitory control shows a faster development than spatial cognition but developed more
slowly than physical problem-solving and lacks a clear plateau before sub-adulthood. Thus, life-history
demands likely influence the developmental timings of inhibitory control, spatial cognition, and physical
problem solving in meerkats.
504

We found little individual variation in inhibition control for both baseline performance (without the
effect of sex and development) and developmental rates (learning curves). In addition, we did not find
evidence for a difference in performance between males and females. We hypothesized that the ability to

inhibit behaviour would be a cognitive underpinning of cooperative behaviours (26,45). Although
meerkats show consistent and repeatable differences in their contributions to cooperative behaviours
(69,70) and males' relative contributions differ from females' (25), we did not find evidence that
differences in the early development of inhibitory control could underlie these patterns. Instead, the
ontogeny of this cognitive trait seems developmentally canalized, with limited influence of environmental
or genetic factors (71). Inhibitory control is still likely to play a part in cooperative activities, because adult
meerkats need a minimum level of inhibitory control to perform cooperative activities by supressing self-
serving behaviours such as ingesting foraged prey, after which internal and physiological states are likely
to drive individual variation. Specifically, in meerkats, contributions to cooperative activities are linked to
body condition (72), foraging success (73), and individual necessities (e.g., gathering information for
future dispersal) (25), and are under hormonal control (74). The conserved early life development of
inhibitory control in meerkats could also result from low genetic variability underlying this trait, arising from
selection removing sub-optimal variance. However, although individual variation was not evident during
early development, it may arise later in response to ecological or social pressures. Future studies should
investigate if adult performance is repeatable and whether it is linked to enduring differences in
cooperative contributions.

In the spatial cognition task, individuals differed in both baseline performances (without the effect
of development, sex, apparatus design or stress), and rates of development (learning curves). In addition,
although males are the dispersing sex, they performed similarly to females. Individual differences in
baseline performances could be due to intrinsic cognitive characteristics or spatial orientation strategy.
Spatial orientation can be either based on position in space in relation to an individual's own position and
direction (egocentric orientation), or on external cues like landmarks (allocentric orientation) (75). Our
study design focused on egocentric orientation, as the apparatus was opaque and the orientation of the
correct exit in relation to any other perceivable external cue like magnetic field and group position
(through sound) was inconsistent among trials. We found that when the apparatus was even partially
transparent during the first season, meerkats were more likely to make errors, which was not explained
by stress levels nor other study design decisions. Therefore, a decrease in performance during the first
season is likely linked to the partial transparency of the apparatus, and consequently individual spatial
orientation strategy. In addition, the ability to inhibit intuitive responses could have also influenced
performance in a partially transparent apparatus, as subjects had to inhibit their attempts to go through a
transparent obstacle. However, we did not find that performance in the inhibition control predict
performance in the spatial maze in our dataset. Interestingly, individuals that were below average in
baseline performances (without the effect of development) had faster learning curves, suggesting that
there are minimum levels of cognitive functions that individuals are selected to reach at the onset of sub-
adulthood.

In the physical problem-solving task, we found individual differences in both baseline
performances (without the effect of development, sex and explorative effort) and rates of development.
Individual performances were also consistent from the tenth week until sub-adulthood, suggesting that

differences in physical problem-solving performance in meerkats arise early in development and are
enduring. Individuals who were presented the task for the first time at ten weeks of age outperformed
their peers who were presented it for the first time at seven weeks and were not substantially different
from those same peers when the latter were re-presented with it at ten weeks of age. This suggest that
the increase in performance we observe during early development is not exclusively due to memory and
previous test experience, but in fact likely linked to the development of brain functions and motor skills.
Although motor skills are usually seen as a confounding factors when investigating underlying cognitive
ability, recent work has questioned whether motor skills and cognitive abilities could be meaningfully
disentangled from each other (76). As such, a physical problem-solving test where motor skills are
involved is a meaningful assessment of cognitive ability. Our results can be linked to different meerkat
foraging strategies. Individuals either scan the ground and find preys on the surface or identify specific
places in the environment where they dig up to thirty centimetres for the chance of finding prey below
ground. Foraging strategy is known to change with age (33,77), and young individuals may also vary in
their tendency to engage in energetically costly foraging with uncertain returns but valuable learning
opportunities. These different strategies could reflect variation in learning curves in their physical
problem-solving abilities and be linked to socio-ecological conditions experienced during development.
Similarly to performance in the spatial cognition task, when individuals had a below average baseline
performances, they showed faster learning curves. Thus, selection may shape meerkats to reach a
minimal necessary level of physical cognition at the onset of sub-adulthood to handle ecological
challenges arising from independent foraging.

Individual performances across the three tasks were uncorrelated, both in their learning curves,
individuals' baseline abilities, or performance at any developmental time point right up to sub-adulthood,
suggesting that in meerkats inhibitory control, spatial cognition (navigation and memory) and problem-
solving ability are likely modular rather than domain-general adaptations (42). Their cognitive
underpinnings may therefore be independent of one another, and potentially subject to separate selective
pressures, at least during early development. Moreover, we found that meerkats' chances of successfully
passing a standardized inhibitory control test at the onset of sub-adulthood (model predicted performance
= 53.4%, 95% CI [37.8%, 68.1%]) were worse than 20 of 32 species investigated by MacLean and
colleagues (51). However, meerkat performance in inhibitory control may not be fully developed by this
point in their ontogeny. Altogether, our results from a cooperative mammal do not align with hypotheses
that link social learning (42) or cooperative care (26,44,45) to the evolution of general intelligence or
enhanced cognitive abilities in primates. As such, our study highlight how selective pressures may not be
operating uniformly and require fine scale developmental investigations including different animal taxa.

Conclusion
We present the first detailed longitudinal study on cognitive development in a carnivorous
mammal under natural conditions. We provide evidence that three cognitive traits (inhibition control,
spatial cognition, and physical problem-solving) emerge at different times of early development in
meerkats, likely because of distinct socio-ecological demands. By investigating individual deviations from

population-level trends, we found that inhibitory control follows a conserved developmental trajectory, while problem-solving ability and spatial cognition show inter-individual developmental variability and are likely to be influenced by socio-ecological or genetic factors. Cognitive performance across the three cognitive traits did not correlate within individuals, either at singular points in time or in their developmental trajectories, suggesting modular rather than domain-general cognitive underpinnings. These findings highlight the importance of studying cognitive evolution across different taxa, social systems and developmental timescales. Future studies should uncover the sources of the individual variation in developmental trajectories and their adaptive consequences.

Ethical statement

Acknowledgements

We thank the Swiss National Science Foundation for funding this work through the grant no. PZ00P3_202052 awarded to SF. We thank the Kalahari Research Trust and the Kalahari Meerkat Project for organisational support and access to the study population. We are grateful to neighbouring landowners for permission to use their land. We are particularly thankful to Marta Manser for discussion and feedback on the manuscript. We also thank Walter Jubber, Jordan Martin, Josh Arbon, and Emily Stott for their valuable input and support. Finally, we appreciate all members of the Kalahari Research Centre (staff, volunteers, managers, and researchers) for maintaining the research infrastructure, habituating and monitoring the meerkats, and providing professional and logistical support during fieldwork.

Declaration of interests

The authors declare no conflict of interest.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT (OpenAI, 2025) to improve grammar, clarity and quality of language. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Bibliography

1. Morand‑Ferron J, Cole EF, Quinn JL. Studying the evolutionary ecology of cognition in the wild: a review of practical and conceptual challenges. Biol Rev. 2016 May;91(2):367–89.

2. Thornton A, Lukas D. Individual variation in cognitive performance: developmental and evolutionary perspectives. Philos Trans R Soc B Biol Sci. 2012 Oct 5;367(1603):2773–83.

3. Ashton BJ, Ridley AR, Edwards EK, Thornton A. Cognitive performance is linked to group size and affects fitness in Australian magpies. Nature. 2018 Feb;554(7692):364–7.

4. Howard-Spink E, Matsuzawa T, Carvalho S, Hobaiter C, Almeida-Warren K, Gruber T, et al. Old age variably impacts chimpanzee engagement and efficiency in stone tool use. eLife. 2025 Jul 15;14.

5. Speechley EM, Ashton BJ, Thornton A, King SL, Simmons LW, Ridley AR. Social Interactions Are Related to Cognitive Development in Western Australian Magpie Fledglings. Ecol Evol. 2025 Nov;15(11):e72435.

6. Dingemanse NJ, Both C, Drent PJ, Tinbergen JM. Fitness consequences of avian personalities in a fluctuating environment. Proc R Soc Lond B Biol Sci. 2004 Apr 22;271(1541):847–52.

7. Liedtke J, Fromhage L. Modelling the evolution of cognitive styles. BMC Evol Biol. 2019 Dec;19(1):234.

8. Cockrem JF. Individual variation, personality, and the ability of animals to cope with climate change. Front Ecol Evol. 2022 Sep 2;10:897314.

9. Isler K, Van Schaik CP. The Expensive Brain: A framework for explaining evolutionary changes in brain size. J Hum Evol. 2009 Oct;57(4):392–400.

10. Soravia C, Ashton BJ, Thornton A, Ridley AR. General cognitive performance declines with female age and is negatively related to fledging success in a wild bird. Proc R Soc B Biol Sci. 2022 Dec 21;289(1989).

11. Kotrschal A, Rogell B, Bundsen A, Svensson B, Zajitschek S, Brännström I, et al. Artificial Selection on Relative Brain Size in the Guppy Reveals Costs and Benefits of Evolving a Larger Brain. Curr Biol. 2013 Jan;23(2):168–71.

12. Bell BA, Phan ML, Meillère A, Evans JK, Leitner S, Vicario DS, et al. Influence of early-life nutritional stress on songbird memory formation. Proc R Soc B Biol Sci. 2018 Sep 26;285(1887):20181270.

13. Fisher MO, Nager RG, Monaghan P. Compensatory Growth Impairs Adult Cognitive Performance. DeVoogd T, editor. PLoS Biol. 2006 Jul 18;4(8):e251.

14. Nyaradi A, Li J, Hickling S, Foster J, Oddy WH. The role of nutrition in children's neurocognitive development, from pregnancy through childhood. Front Hum Neurosci. 2013;7.

15. Speechley EM, Ashton BJ, Thornton A, King SL, Simmons LW, Woodiss-Field SL, et al. Aggressive interactions influence cognitive performance in Western Australian magpies. Proc R Soc B Biol Sci. 2024 Jun;291(2024).

16. Grimberg-Henrici CGE, Vermaak P, Elizabeth Bolhuis J, Nordquist RE, Van Der Staay FJ. Effects of environmental enrichment on cognitive performance of pigs in a spatial holeboard discrimination task. Anim Cogn. 2016 Mar;19(2):271–83.

17. Salvanes AGV, Moberg O, Ebbesson LOE, Nilsen TO, Jensen KH, Braithwaite VA. Environmental enrichment promotes neural plasticity and cognitive ability in fish. Proc R Soc B Biol Sci. 2013 Sep 22;280(1767):20131331.

18. Rosati AG, Wobber V, Hughes K, Santos LR. Comparative Developmental Psychology: How is Human Cognitive Development Unique? Evol Psychol. 2014 Apr 1;12(2):448–73.

19. Wobber V, Wrangham R, Hare B. Application of the heterochrony framework to the study of behavior and cognition. Commun Integr Biol. 2010 Jul;3(4):337–9.

20. Langer J. The Heterochronic Evolution of Primate Cognitive Development. Biol Theory. 2006 Mar;1(1):41–3.

21. Gomez J. Species comparative studies and cognitive development. Trends Cogn Sci. 2005 Mar;9(3):118–25.

22. Herrmann E, Call J, Hernàndez-Lloreda MV, Hare B, Tomasello M. Humans Have Evolved Specialized Skills of Social Cognition: The Cultural Intelligence Hypothesis. Science. 2007 Sep 7;317(5843):1360–6.

23. Rosati AG. Heterochrony in chimpanzee and bonobo spatial memory development. Am J Phys Anthropol. 2019 Jun;169(2):302–21.

24. Wobber V, Wrangham R, Hare B. Bonobos Exhibit Delayed Development of Social Behavior and Cognition Relative to Chimpanzees. Curr Biol. 2010 Feb;20(3):226–30.

25. Clutton-Brock T, Russell AF, Sharpe LL, Young AJ, Balmforth Z, McIlrath GM. Evolution and Development of Sex Differences in Cooperative Behavior in Meerkats. Science. 2002 Jul 12;297(5579):253–6.

26. Burkart JM, Van Schaik CP. Cognitive consequences of cooperative breeding in primates? Anim Cogn. 2010 Jan;13(1):1–19.

27. Burkart JM, Allon O, Amici F, Fichtel C, Finkenwirth C, Heschl A, et al. The evolutionary origin of human hyper-cooperation. Nat Commun. 2014 Aug 27;5(1).

28. Thornton A, McAuliffe K. Cognitive consequences of cooperative breeding? A critical appraisal. J Zool. 2015 Jan;295(1):12–22.

29. Clutton-Brock T, Manser M. Meerkats: Cooperative breeding in the Kalahari. In: Koenig WD, Dickinson JL, editors. Cooperative Breeding in Vertebrates. 1st ed. Cambridge University Press; 2016. p. 294–317.

30. Hollén LI, Manser MB. Ontogeny of alarm call responses in meerkats, Suricata suricatta: the roles of age, sex and nearby conspecifics. Anim Behav. 2006 Dec;72(6):1345–53.

31. Manser MB, Bell MB. Spatial representation of shelter locations in meerkats, Suricata suricatta. Anim Behav. 2004 Jul;68(1):151–7.

32. Doolan SP, Macdonald DW. Diet and foraging behaviour of group-living meerkats, Suricata suricatta, in the southern Kalahari. J Zool. 1996 Aug;239(4):697–716.

33. Jubber WR, Manser MB, Fuller A. Feeding through the ages: Revisiting the diet of meerkats. J Arid Environ. 2025 Mar;227:105331.

34. Thornton A. Social learning about novel foods in young meerkats. Anim Behav. 2008 Oct;76(4):1411–21.

35. Thornton A, Hodge SJ. The development of foraging microhabitat preferences in meerkats. Behav Ecol. 2009;20(1):103–10.

36. Thornton A, McAuliffe K. Teaching in Wild Meerkats. Science. 2006 Jul 14;313(5784):227–9.

37. Clutton-Brock TH, Brotherton PNM, Smith R, McIlrath GM, Kansky R, Gaynor D, et al. Infanticide and expulsion of females in a cooperative mammal. Proc R Soc Lond B Biol Sci. 1998 Dec 7;265(1412):2291–5.

38. Griffin AS, Pemberton JM, Brotherton PNM, McIlrath G, Gaynor D, Kansky R, et al. A genetic analysis of breeding success in the cooperative meerkat (Suricata suricatta). Behav Ecol. 2003 Jul 1;14(4):472–80.

39. Young AJ, Spong G, Clutton-Brock T. Subordinate male meerkats prospect for extra-group paternity: alternative reproductive tactics in a cooperative mammal. Proc R Soc B Biol Sci. 2007 Jul 7;274(1618):1603–9.

40. Ashton BJ, Kennedy P, Radford AN. Interactions with conspecific outsiders as drivers of cognitive evolution. Nat Commun. 2020 Oct 6;11(1):4937.

41. Thornton A, Clutton-Brock T. Social learning and the development of individual and group behaviour
in mammal societies. Philos Trans R Soc B Biol Sci. 2011 Apr 12;366(1567):978–87.

42. Burkart JM, Schubiger MN, Van Schaik CP. The evolution of general intelligence. Behav Brain Sci.
2017;40:e195.

43. Whiten A. When does cultural transmission favour or instead substitute for general intelligence?
Behav Brain Sci. 2017;40:e222.

44. Van Schaik CP, Song Z, Schuppli C, Drobniak SM, Heldstab SA, Griesser M. Extended parental
provisioning and variation in vertebrate brain sizes. PLOS Biol. 2023 Feb 28;21(2):e3002016.

45. Burkart JM, Hrdy SB, Van Schaik CP. Cooperative breeding and human cognitive evolution. Evol
Anthropol Issues News Rev. 2009 Sep;18(5):175–86.

46. Isler K, Van Schaik CP. Why are there so few smart mammals (but so many smart birds)? Biol Lett.
2009 Feb 23;5(1):125–9.

47. Cozzi G, Maag N, Börger L, Clutton-Brock TH, Ozgul A. Socially informed dispersal in a territorial
cooperative breeder. Street G, editor. J Anim Ecol. 2018 May;87(3):838–49.

48. Clutton-Brock TH, Gaynor D, Kansky R, MacColl ADC, McIlrath G, Chadwick P, et al. Costs of
cooperative behaviour in suricates (Suricata suricatta). Proc R Soc Lond B Biol Sci. 1998 Feb
7;265(1392):185–90.

49. Clutton-Brock TH, Maccoll A, Chadwick P, Gaynor D, Kansky R, Skinner JD. Reproduction and
survival of suricates (Suricata suricatta) in the southern Kalahari. Afr J Ecol. 1999 Mar;37(1):69–80.

50. Duncan C, Turner Z, Gaynor D, Thorley J, Vink T, Clutton-Brock T. The ontogeny of foraging in
meerkats, a cooperatively breeding mongoose. Anim Behav. 2025 Sep;227:123302.

51. MacLean EL, Hare B, Nunn CL, Addessi E, Amici F, Anderson RC, et al. The evolution of self-control.
Proc Natl Acad Sci. 2014 May 20;111(20).

52. Collier K, Townsend SW, Manser MB. Call concatenation in wild meerkats. Anim Behav. 2017
Dec;134:257–69.

53. Birchmeier K, Johnson-Ulrich L, Stein J, Forss S. The Role of Umwelt in Animal Curiosity: A Within
and Between Species Comparison of Novelty Exploration in Mongooses. Anim Behav Cogn. 2023
Nov 1;10(4):329–54.

54. Friard O, Gamba M. BORIS : a free, versatile open‐source event‐logging software for video/audio
coding and live observations. Fitzjohn R, editor. Methods Ecol Evol. 2016 Nov;7(11):1325–30.

55. Van Horik JO, Madden JR. A problem with problem solving: motivational traits, but not cognition,
predict success on novel operant foraging tasks. Anim Behav. 2016 Apr;114:189–98.

56. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation
for Statistical Computing; 2024. Available from: https://www.R-project.org/

57. Posit team. RStudio: Integrated Development Environment for R. Posit Software [Internet]. Boston,
MA: Posit Software, PBC; 2025. Available from: http://www.posit.co/

58. Bürkner PC. **brms** : An *R* Package for Bayesian Multilevel Models Using *Stan*. J Stat Softw.
2017;80(1).

59. Lemoine NP. Moving beyond noninformative priors: why and how to choose weakly informative priors
in Bayesian analyses. Oikos. 2019 Jul;128(7):912–28.

60. Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. performance: An R Package for
Assessment, Comparison and Testing of Statistical Models. J Open Source Softw. 2021 Apr
21;6(60):3139.

61. Bürkner PC. Bayesian Item Response Modeling in *R* with **brms** and *Stan*. J Stat Softw [Internet].
2021 [cited 2025 May 27];100(5). Available from: https://www.jstatsoft.org/v100/i05/

62. Farrar BG, Vernouillet A, Garcia-Pelegrin E, Legg EW, Brecht KF, Lambert PJ, et al. Reporting and
interpreting non-significant results in animal cognition research. PeerJ. 2023 Mar 9;11:e14963.

63. Averly B, Sridhar VH, Demartsev V, Gall G, Manser M, Strandburg-Peshkin A. Disentangling
influence over group speed and direction reveals multiple patterns of influence in moving meerkat
groups. Sci Rep. 2022 Aug 16;12(1):13844.

64. Navarrete A, Van Schaik CP, Isler K. Energetics and the evolution of human brain size. Nature. 2011
Dec 1;480(7375):91–3.

65. Berdahl A, Torney CJ, Ioannou CC, Faria JJ, Couzin ID. Emergent Sensing of Complex
Environments by Mobile Animal Groups. Science. 2013 Feb;339(6119):574–6.

66. Brotherton PNM, Clutton-Brock TH, O'Riain MJ, Gaynor D, Sharpe L, Kansky R, et al. Offspring food
allocation by parents and helpers in a cooperative mammal. Behav Ecol. 2001 Sep 1;12(5):590–9.

67. Manser MB, Madden JR, Kunc HP, English S, Clutton-Brock T. Signals of need in a cooperatively
breeding mammal with mobile offspring. Anim Behav. 2008 Dec;76(6):1805–13.

68. Madden JR, Kunc HJP, English S, Clutton-Brock TH. Why do meerkat pups stop begging? Anim
Behav. 2009 Jul;78(1):85–9.

69. English S, Nakagawa S, Clutton-Brock TH. Consistent individual differences in cooperative behaviour
in meerkats (*Suricata suricatta*). J Evol Biol. 2010 Aug;23(8):1597–604.

70. Carter AJ, English S, Clutton-Brock TH. Cooperative personalities and social niche specialization in
female meerkats. J Evol Biol. 2014 May;27(5):815–25.

71. Siegal ML, Bergman A. Waddington's canalization revisited: Developmental stability and evolution.
Proc Natl Acad Sci. 2002 Aug 6;99(16):10528–32.

72. Clutton-Brock TH, Brotherton PNM, O'Riain MJ, Griffin AS, Gaynor D, Sharpe L, et al. Individual
contributions to babysitting in a cooperative mongoose, *Suricata suricatta*. Proc R Soc Lond B Biol
Sci. 2000 Feb 7;267(1440):301–5.

73. Clutton-Brock TH, Brotherton PNM, O'Riain MJ, Griffin AS, Gaynor D, Kansky R, et al. Contributions
to cooperative rearing in meerkats. Anim Behav. 2001 Apr;61(4):705–10.

74. Dantzer B, Dubuc C, Goncalves IB, Cram DL, Bennett NC, Ganswindt A, et al. The development of
individual differences in cooperative behaviour: maternal glucocorticoid hormones alter helping
behaviour of offspring in wild meerkats. Philos Trans R Soc B Biol Sci. 2019 Apr
15;374(1770):20180117.

75. Bottini R, Doeller CF. Knowledge Across Reference Frames: Cognitive Maps and Image Spaces.
Trends Cogn Sci. 2020 Aug;24(8):606–19.

76. Barton R, Barrett L. Embodied cognitive evolution and the limits of convergence. Philos Trans R Soc
B Biol Sci. 2025 Jun 26;380(1929).

77. Duncan C, Turner Z, Gaynor D, Thorley J, Vink T, Clutton-Brock T. The ontogeny of foraging in
meerkats, a cooperatively breeding mongoose. Anim Behav. 2025 Sep;227:123302.

794