Structural mechanisms of pump assembly and drug transport in the AcrAB-TolC

efflux system

Running title: Cryo-EM structures of TolC-YbjP and TolC-YbjP-AcrABZ

Xiaofei Ge[1,2,*], Zhiwei Gu[1], Jiawei Wang[1,*]

[1]State Key Laboratory of Membrane Biology, Beijing Frontier Research Center for Biological

Structure, School of Life Sciences, Tsinghua University, Beijing 100084, P. R. China

[2]Health and Wellness, City University of Macau, Macau, 999078, P. R. China

*Correspondence should be addressed to: X.G. (gxf16@tsinghua.org.cn) or J.W.

(jwwang@tsinghua.edu.cn)

Abstract

Tripartite multidrug efflux pumps that span the cell envelope are essential for antibiotic

resistance in Gram-negative bacteria. Here, we report cryo-EM structures of two native

efflux complexes from *Escherichia coli*: a TolC–YbjP subcomplex at 3.56 Å resolution

and the complete TolC–YbjP–AcrABZ pump at 3.39 Å. Structural analysis reveals that

YbjP, a previously uncharacterized lipoprotein, binds TolC in a 3:3 stoichiometry,

bridging its protomers at the equatorial domain. Clear density of the mature YbjP's N-

terminal Cys19 indicates that YbjP is anchored to the outer membrane by an N-terminal

lipid moiety. Notably, YbjP remains bound as TolC undergoes AcrA-induced opening,

suggesting that this accessory protein accommodates the conformational change. The

AcrB trimer simultaneously presents three distinct conformational states (L, T, O),

capturing a complete transport cycle. These high-resolution structures provide insights

into the architecture and mechanism of clinically relevant efflux machinery, identifying

YbjP as a previously unrecognized structural component that contributes to TolC

positioning and may assist in its membrane localization—offering potential new targets

for therapeutic intervention.

Keywords

Introduction

Tripartite efflux pumps are essential for Gram-negative bacteria to expel a wide range of toxic compounds, including antibiotics, across their dual-membrane cell envelope[1-5]. Among these systems, the AcrAB–TolC complex of *Escherichia coli* is one of the most thoroughly studied[2,6]. It comprises the inner membrane RND (Resistance–Nodulation–Division) transporter AcrB, the periplasmic membrane fusion protein AcrA, and the outer membrane channel TolC[4,6-9]. AcrA, anchored to the inner membrane via N-terminal lipidation, bridges AcrB and TolC, mediating conformational signaling essential for TolC gating[10-12]. Structural studies have shown that TolC transitions from a closed to an open state in response to AcrAB assembly, a process crucial for efflux activation[13,14].

Despite significant progress in structurally characterizing the AcrAB–TolC system, fundamental questions remain regarding how TolC is positioned and retained in the outer membrane. Unlike homologous channels such as *Pseudomonas aeruginosa* OprM[15] and *E. coli* CusC[16], which possess covalently attached N-terminal lipids, TolC lacks a lipid anchor. This raises the possibility that *E. coli* may utilize accessory factors to localize and stabilize TolC during pump assembly. Furthermore, while prior studies using crystallography and low-to-medium resolution cryo-EM have revealed snapshots of the assembled pump[10-14,17], the limited resolution and conformational heterogeneity in those structures impeded accurate modeling of flexible regions and may have masked the presence of additional subunits.

Here, we address these knowledge gaps by determining two high-resolution cryo-EM structures of the TolC-based efflux machinery in *E. coli*. The first is a 3.56 Å structure of a TolC–YbjP subcomplex, and the second is a 3.39 Å structure of the fully

assembled TolC–YbjP–AcrABZ pump. These structures reveal a previously

uncharacterized outer membrane lipoprotein, YbjP, bound to TolC, and shed light on how

TolC is secured and transitions to the open state. Together, these findings enhance our

understanding of the architecture, assembly, and regulation of this clinically important

efflux pump.

Results

Structure of a TolC–YbjP closed-state complex

Native purification of *Escherichia coli* membranes yielded a stable assembly containing

the outer-membrane channel TolC. Cryo-EM analysis resolved the complex at 3.56 Å

resolution (Figure S1, Supplementary Table 1). TolC forms a homotrimer composed of a

β-barrel that spans the outer membrane and an elongated periplasmic α-helical domain

(Figures 1A-D). The periplasmic α-helical domain comprises an α-helical barrel extended

by two helix-turn-helix (HTH) motifs at its periplasmic extremity (TolC repeat 1: helices

H3-H4; repeat 2: helices H7-H8). These structural elements converge at an equatorial

domain (green in Figures 1C-D). The inward-bending HTH coiled-coils occlude the

periplasmic tunnel, maintaining TolC in a closed conformation—consistent with previous

crystal structures[10] (Figure 1C).

An unexpected density at the periplasmic face of TolC indicated the presence of

an additional component (wheat in Figures 1A-B). An initial backbone trace, followed by

a DALI search[18] tentatively matched Tai3, a periplasmic type IV immunity protein

associated with the T6SS amidase effector Tae3 (PDB ID: 4HZ9)[19] from *Ralstonia*

pickettii. However, discrepancies in side chain density and species origin indicated that

the match was likely incorrect. Systematic screening of the AlphaFold Database[20] using

CryoNet[21] identified *E. coli* YbjP (UniProt P75818) as the top candidate, with all side

chains matching the experimental density (Figure S2). YbjP features a globular domain

and a structured N-terminal loop stabilized by a Cys36-Cys144 disulfide bond (Figures

1E-F). The N-terminal loop of YbjP adopts an extended conformation parallel to the first

α-helix of TolC's α-helical barrel, forming substantial intermolecular interfaces. Buried

surface area (BSA) analysis reveals 787 Å² of contact with the primary TolC protomer

(green in Figure 1F), and 1037.4 Å² with the adjacent protomer (light green), suggesting

asymmetric binding energetics. The N-terminal density extends to Cys19, which

corresponds to the predicted signal peptide cleavage site, positioning the lipoprotein at

the periplasmic face of the outer membrane.

UniProt predictions indicate that YbjP is dually lipidated via N-palmitoyl and S-

diacylglycerol modifications, allowing its anchorage to the outer membrane. This

represents an elegant evolutionary solution, as while many Gram-negative TolC

homologs (*e.g.*, *Pseudomonas* OprM[15], *E. coli* CusC[16]) possess native lipid anchors,

E. coli TolC instead recruits YbjP as a dedicated membrane-tethering partner. The YbjP

globular domain nestles between adjacent TolC equational domains, forming a stable 3:3

complex (Figure 1B). The identification of YbjP, along with its defined membrane

anchoring and structural compatibility with TolC, suggests a new class of outer

membrane partners that may regulate the TolC-dependent pathways.

Structure of TolC-YbjP-AcrABZ complex

An improved 3.39 Å resolution structure of the endogenous AcrAB–TolC complex from

E. coli was determined, revealing well-resolved density across the entire assembly

(Figure S3, Supplementary Table 1). This structure allows for the accurate modeling of

side-chain interactions throughout the tripartite channel, surpassing earlier models in

completeness and resolution[14] (Figures S4, S5). YbjP remains bound to TolC, suggesting

that this lipoprotein is stably retained during pump assembly and functions as a molecular

clamp that anchors TolC to the outer membrane during AcrA and AcrB engagement

(Figures 2A-D).

The TolC–YbjP–AcrABZ complex adopts a funnel-like architecture spanning the

cell envelope, consistent with prior models of tripartite pumps[14] (Figure 2B). TolC,

tethered to YbjP, caps the outer-membrane end and docks onto a hexameric ring of AcrA

adaptors in the periplasm, which in turn surrounds the AcrBZ trimer embedded in the

inner membrane (Figures 2B, 2D). The full assembly is about 33 nm tall, matching the

distance between the two membranes. The complex was purified directly from *E. coli* as

a stoichiometric assembly, without the need for artificial fusion or crosslinking.

Cryo-EM density reveals six AcrA molecules beneath each TolC trimer (Figure

2D), yielding a 3:6:3 stoichiometry for TolC:AcrA:AcrB. AcrA therefore forms an

elongated hexamer that bridges TolC and AcrB. Three sets of interfaces stabilize the

pump: contacts between the AcrB trimer and the basal regions of AcrA, extensive AcrA–

AcrA interactions within the hexameric ring, and tip-to-tip junctions between the upper

AcrA α-helical hairpin and TolC (Figure 2D). While each AcrA protomer maintains the

characteristic four-domain architecture—comprising the α-helical hairpin, lipoyl, β-

barrel, and membrane-proximal domains—functional asymmetry is observed in their

interactions (Figure 2E). The trimeric TolC, which contains six HTH motifs due to

internal repeats (TolC repeat 1/2), engages the AcrA hexamer via quasi-equivalent

binding: adjacent AcrA and AcrA* protomers interact differentially with the intra- and

inter-protomer grooves of TolC, respectively (Figure 2F). The primary structural

difference between AcrA and AcrA* lies in the configurations of their membrane-

proximal (MP) domains (Figure 2E).

AcrB assembles as a homotrimer on the inner membrane, with each protomer

comprising three distinct domains: a funnel domain, a porter domain, and a

transmembrane domain (Figure 2D). The funnel domain consists of DN and DC

subdomains (Figure 2G). The porter domain is organized into four subdomains (PN1,

PN2, PC1, and PC2) arranged in a clockwise orientation when viewed from the

periplasmic side, with PN1 positioned closest to the central axis of the trimer. In the

hexameric AcrA assembly, the lipoyl and β-barrel domains form two stacked concentric

rings. While the lipoyl domains remain uninvolved in AcrB binding, the β-barrel domains

specifically engage the funnel domain of AcrB. Strikingly, the membrane-proximal (MP)

subdomains of AcrA exhibit asymmetric binding: AcrA-MP interacts with the DC and

PC1 subdomains of AcrB, whereas AcrA*-MP contacts PN1 and DN via an extended

loop, respectively (Figures 2G-H). This differential engagement of MP subdomains

establishes the structural basis for the observed structural divergence between AcrA and

AcrA* (Figure 2E).

Our density maps clearly resolve the small transmembrane protein AcrZ (49

amino acids) bound to each AcrB protomer (Figures 2C, 2D). AcrZ adopts a helical

structure that is embedded within a hydrophobic groove formed by the transmembrane

helices of AcrB at the predicted interaction site, forming an AcrB₃AcrZ₃ complex.

Although AcrZ is not essential for pump assembly, its consistent presence in natively

purified complexes and its role in stabilizing otherwise flexible regions—particularly the

transmembrane helices—suggest that AcrZ may serve an allosteric role in modulating

conformational dynamics for specific substrates[22]. Collectively, these interactions

illustrate the sophisticated assembly mechanism of the pump: AcrZ reinforces AcrB's

transmembrane domain, AcrB's funnel and porter domains anchor AcrA, and AcrA's α-

helical hairpins engage TolC. Notably, the holocomplex remained intact throughout

purification, indicating a high-affinity assembly between components, consistent with

prior *in vitro* reconstitution studies[23].

Structural rearrangements underlying TolC's closed-to-open transition

A comparative structural analysis of TolC in its closed and open states reveals a striking

iris-like dilation mechanism at the periplasmic entrance that facilitates transition to the

fully open conformation (Figures 3A-B). Throughout this conformational transition, the

transmembrane β-barrel and α-helical barrel domains maintain remarkable structural

rigidity, whereas the coiled-coil helices undergo dramatic rearrangement. These helices

pivot around the equatorial domain, undergoing a 27° rigid-body superhelical rotation

(Figure 3C), which constitutes a major structural rearrangement. In the closed state, a

complex network of interprotomer hydrogen bonds stabilizes the structure by constricting

the pore to its narrowest point at Asp396 (Figures 3D,3F). Upon disruption of this

network in the open state, the pore expands significantly, reaching a diameter of

approximately 2 nm (Figures 3E-F). Notably, YbjP remains stably association with TolC

in both conformational states – the closed resting state and the open activated state

(Figure 3C). This persistent interaction suggests that YbjP serves as a structural scaffold:

anchoring TolC in the outer membrane, accommodating conformational changes during

activation, and functionally compensating for TolC's lack of intrinsic lipidation. These

findings not only support existing models of allosteric pump activation[12,17], but also

suggest how *E. coli* might utilize YbjP to fulfill the membrane-anchoring role typically

provided by intrinsic lipid modifications in other bacterial TolC homologs, such as OprM

and CusC[24,25].

Mechanism of substrate transport in the AcrB module

The AcrAB-TolC efflux system actively transports substrates against their concentration

gradient, moving them from regions of low to high concentration. This energetically

unfavorable process is driven by proton motive force (PMF), with the AcrB trimer

harnessing the electrochemical potential of protons to power substrate translocation[26].

Structural and functional studies support an asymmetric rotary mechanism, wherein each

AcrB protomer sequentially transitions through three conformational states: L (loose), T

(tight), and O (open)[17,27]. This concerted conformational rotation – analogous to the

functional cycling of F_1F_0 ATP synthase during ATP synthesis – enables continuous

vectorial transport through the pump complex[28].

Our high-resolution cryo-EM structure of the endogenous TolC-YbjP-AcrABZ

complex captures the AcrB trimer in three distinct conformational states (L, T, O),

providing key mechanistic insights (Figure 4A). While the transmembrane domain

remains largely conserved between the L and T states (Figure 4B, upper panel)—

maintaining salt bridges between Lys940 (TM10) and Asp407/Asp408 (TM4), the T→O

transition entails a coordinated rotation of TM2, TM4, TM5, and TM6, disrupting these

interactions (Figure 4B, lower panel). Substrate entry occurs between PC1 and PC2 in the

L state (Figure 4C, left panel), followed by an 8 Å shift of PC2 toward PC1 during the L

→T transition (Figure 4D, lower panel and Figure S6), facilitating drug transfer from the

access pocket to the high-affinity deep binding pocket (Figure 4C, middle panel).

Remarkably, PC1 remains in contact with AcrA throughout the cycle (Figure S6). Proton

translocation through the transmembrane domain induces three coordinated events: (1) a

further 5 Å compaction between PC2 and PC1 (Figure 4D, T, O states in lower panel);

(2) inward rotation of PN2's cytoplasmic face (Figure 4D, T, O states in upper panel);

and (3) outward opening of PN1's periplasmic side—together expelling the substrate into

the central channel. This rotary mechanism, reminiscent of F_1F_0 ATP synthase, is

completed when the pump resets to the L state, initiating a new catalytic cycle.

Discussion

Based on our structural data, we propose a model for the assembly and function of the

AcrABZ–TolC multidrug efflux pump, emphasizing the anchoring and orienting role of

YbjP. As shown in Figure 5, the assembly begins with TolC priming. In the absence of

AcrAB, TolC is embedded in the outer membrane in a closed conformation, stabilized by

its interaction with YbjP. The lipid moiety of YbjP inserts into the inner leaflet of the

outer membrane, possibly interacting transiently with the peptidoglycan layer to help

position TolC within the periplasmic space. Such anchoring likely ensures optimal

orientation of TolC's periplasmic end for interaction with incoming AcrA.

Upon synthesis and membrane insertion, AcrB (complexed with AcrZ) and AcrA

form an inner membrane–periplasmic subcomplex[29,30]. A conserved cysteine residue in

AcrA, located near the inner membrane, may aid in anchoring or stabilizing the complex.

The complete tripartite pump likely assembles via stochastic encounters between the

AcrAB subcomplex and the TolC–YbjP complex in the periplasm. Stabilization likely

occurs through tip-to-tip interactions between the hairpin domains of AcrA and TolC[31].

YbjP may facilitate this docking by stabilizing TolC in a fixed, properly oriented

conformation, thereby increasing its local availability to AcrAB and lowering the energy

barrier for assembly. In this context, YbjP may act as a structural placeholder that guides

AcrA toward the TolC entrance. Although its exact function remains to be confirmed,

YbjP's consistent presence in our cryo-EM reconstructions and its predicted lipoprotein

features suggest an anchoring role. Based on our structural observations and UniProt

annotations, we propose that YbjP contributes to TolC stabilization in the outer

membrane and facilitates its spatial orientation for efficient pump assembly.

Once assembled, the AcrABZ–TolC complex becomes active for drug efflux,

allowing substrates such as antibiotics and dyes to enter AcrB from the periplasm and be

expelled through TolC[32,33]. At the core of this process is the conserved D407–D408

proton relay pair in AcrB's transmembrane domain, which functions as both the proton-

binding site and the mechanochemical coupling hub[34,35]. Protonation of these residues

initiates conformational changes that propagate from the transmembrane domain to the

porter domain via an allosteric mechanism.

The AcrB trimer operates via an asymmetric rotary cycle[17,27], with each protomer

sequentially transitioning through three distinct conformational states (L→T→O) to drive

peristaltic drug transport. This concerted conformational rotation – bearing striking

similarity to the catalytic cycle of F_1F_0 ATP synthase[28] – enables continuous, vectorial

substrate transport against concentration gradients, powered by proton translocation.

In conclusion, we identify YbjP as a previously unrecognized lipoprotein

associated with the AcrAB–TolC efflux system. Although its precise function requires

further validation, YbjP's structural positioning suggests a role in anchoring TolC and

promoting tripartite pump assembly—critical for bacterial resistance to antibiotics and

toxic compounds. Notably, our structure captures all three conformational states of the

AcrB trimer (L, T, O) within the context of the fully assembled, native AcrABZ–TolC

complex for the first time. While these states have been observed in isolated AcrB

previously, our data reveal subtle but functionally relevant conformational differences

that arise in the intact pump. This study exemplifies how high-resolution cryo-EM, in

combination with integrative modeling, can reveal previously uncharacterized protein

factors and enhance our understanding of complex membrane-spanning systems in

bacteria.

Method details

Cell preparation and membrane protein extraction

Escherichia coli C600 cells were cultured to mid-log phase and then infected with a high-

titer λ phage to increase cellular stress. Cells were harvested by centrifugation and

resuspended in buffer containing 25 mM Tris-HCl and 150 mM NaCl (pH 8.0).

Lysozyme was added to a final concentration of 10 mg/mL, followed by incubation at

37 °C for 2 hours to facilitate cell lysis. Following removal of cell debris by

centrifugation at 24,793 × g for 20 min, membrane proteins were extracted from the

supernatant using 2% n-dodecyl-β-D-maltoside (DDM). The membrane fraction was

subsequently subjected to size-exclusion chromatography using a SR6 Increase column.

Fractions around 12 mL, which were enriched for target proteins, were collected, diluted,

and used for negative staining. Transmission electron microscopy revealed the presence

of target particles.

Cryo-EM grid preparation

The sample was concentrated to 0.7 mg/mL. A volume of 4 µL was applied to a glow-

discharged Au Quantifoil R1.2/1.3 300 mesh grid (glow discharge: medium setting, 30 s).

To enhance the concentration of protein on the grid, sample loading was repeated ten

times: after each loading, excess solution was gently removed using a pipette tip, and a

fresh 4 µL aliquot was applied. The grid was then plunge-frozen in liquid ethane pre-

cooled with liquid nitrogen using a Vitrobot Mark IV (Thermo Fisher Scientific).

Blotting was performed for 3.5 s after a 30 s wait time, under 100% humidity at 8 °C.

Cryo-EM data collection

A total of 3452 movies were acquired using a 300 kV Titan Krios transmission electron

microscope (Thermo Fisher Scientific) equipped with a Gatan K3 Summit direct electron

detector. The calibrated pixel size was 0.675 Å. Each movie was recorded with a total

electron dose of ~50 e⁻/Å², and defocus values ranged from -1.2 to -1.8 μm. Data

collection was automated using the AutoEMation software[36].

Image processing

Motion correction and contrast transfer function (CTF) estimation were carried out in

CryoSPARC[37]. Poor-quality micrographs were manually excluded. 43,200 particles

corresponding to TolC were identified during 2D classification. Ab-initio reconstruction

and NU-refinement applied with C3 symmetry generated a reference model for further

picking. Template picking, *ab-initio* reconstruction in C1 symmetry, and NU-refinement

in C3 symmetry followed. The resulting map at 4.17 Å resolution was further improved

to 3.56 Å after reference-based motion correction (RBMC), CTF refinement, and NU-

refinement.

From the same dataset, TolC-AcrABZ particle selection was performed using

blob picking, followed by particle extraction and several rounds of 2D classification,

which yielded 19,396 high-quality particles. These were used for ab-initio reconstruction

and non-uniform (NU) refinement to generate an initial 3D model, which was

subsequently used for template picking. After template picking and further 2D

classification, a total of 35,092 particles were selected. Final refinement steps—including

NU-refinement, RBMC, and local CTF refinement—resulted in a 3D map at 3.23 Å

resolution. Since the AcrB subunit exhibits intrinsic structural asymmetry, C3 symmetry

was initially relaxed through symmetry expansion combined with 3D classification

without rotational/translational alignment. One predominant class was selected after

deduplication of symmetry-expanded particles. Local refinement in C1 symmetry better

resolved the asymmetric AcrB region. A subsequent masked 3D classification revealed a

particle subset containing distinct extra densities, which underwent local refinement with

re-applied C3 symmetry to enhance resolution of symmetric structural components.

Protein model building and structure refinement

The atomic models of TolC, AcrA, AcrB, AcrZ were rigid-body fitted into their

corresponding cryo-EM maps using UCSF Chimera[38]. To identify the extra densities

observed in both EM maps, a backbone model was manually traced. Protein identity was

assessed using the "Protein Searching without Sequence" function in CryoNet[21]. YbjP

(P75818) was ranked highest among *E. coli* proteins in the AlphaFold Database (AFDB,

strain 4364). Side-chain densities in the final EM maps were manually compared to the

predicted model, allowing confirmation of YbjP as the protein corresponding to the extra

density. Subsequent manual adjustments to the models were performed using COOT[39]. The

refinement of the protein structures was conducted using either PHENIX[40] or Refmac5[41].

Statistical details regarding the 3D reconstruction and model refinement processes are

provided in Supplementary Table S1. Visual representations of the structures were

generated using PyMol[42] or UCSF ChimeraX[43].

Acknowledgements

We are thankful to the Tsinghua University Branch of the China National Center for

Protein Sciences (Beijing) for their generous assistance with cryo-EM facility support

and computational resources on the Bio-Computing Platform cluster. Additionally, we

appreciate the valuable technical support from J. Lei, X. Li, and F. Yang. This work was

supported by the National Natural Science Foundation of China (Grant Nos. 32501078 to

X.G. and 32371254, 32171190 to J.W.).

Author contributions

343 X.G. and J.W. conceived the project. X.G. and Z.G. optimized the preparation of cryo-

344 grids, recorded the cryo-EM data, and processed this data. J.W. built the atomic models.

345 X.G. and J.W. wrote the manuscript. All authors read and approved the final version of

346 the manuscript.

Declaration of interests

349 The authors declare no competing interests.

Data and code availability

352 Cryo-EM maps and the associated structural coordinates have been respectively

353 deposited into the Electron Microscopy Data Bank (EMDB) and the Protein Data Bank

354 (PDB) under the following accession codes: EMD-64784 / 9V52 (TolC-YbjP); EMD-

355 64785 / 9V53 (TolC-YbjP-AcrABZ); and EMD-64787 / 9V55 (TolC-YbjP-AcrA local).

References

Poole, K. Efflux-mediated antimicrobial resistance. *J Antimicrob Chemother* **56**, 20–51 (2005). https://doi.org/10.1093/jac/dki171

Li, X. Z., Plésiat, P. & Nikaido, H. The challenge of efflux-mediated antibiotic resistance in Gram-negative bacteria. *Clin Microbiol Rev* **28**, 337–418 (2015). https://doi.org/10.1128/cmr.00117-14

Du, D., van Veen, H. W., Murakami, S., Pos, K. M. & Luisi, B. F. Structure, mechanism and cooperation of bacterial multidrug transporters. *Curr Opin Struct Biol* **33**, 76–91 (2015). https://doi.org/10.1016/j.sbi.2015.07.015

Du, D. J. *et al.* Multidrug efflux pumps: structure, function and regulation. *Nature Reviews Microbiology* **16**, 523–539 (2018). https://doi.org/10.1038/s41579-018-0048-6

Blair, J. M., Richmond, G. E. & Piddock, L. J. Multidrug efflux pumps in Gram-negative bacteria and their role in antibiotic resistance. *Future Microbiol* **9**, 1165–1177 (2014). https://doi.org/10.2217/fmb.14.66

Wright, M., Kaur, M., Thompson, L. K. & Cox, G. A historical perspective on the multifunctional outer membrane channel protein TolC in Escherichia coli. *NPJ Antimicrob Resist* **3**, 6 (2025). https://doi.org/10.1038/s44259-025-00078-3

Ma, D. *et al.* Genes acrA and acrB encode a stress-induced efflux system of Escherichia coli. *Mol Microbiol* **16**, 45–55 (1995). https://doi.org/10.1111/j.1365-2958.1995.tb02390.x

Nikaido, H. Multidrug resistance in bacteria. *Annu Rev Biochem* **78**, 119–146 (2009). https://doi.org/10.1146/annurev.biochem.78.082907.145923

Okusu, H., Ma, D. & Nikaido, H. AcrAB efflux pump plays a major role in the antibiotic resistance phenotype of Escherichia coli multiple-antibiotic-resistance (Mar) mutants. *J Bacteriol* **178**, 306–308 (1996). https://doi.org/10.1128/jb.178.1.306-308.1996

Koronakis, V., Sharff, A., Koronakis, E., Luisi, B. & Hughes, C. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature* **405**, 914–919 (2000). https://doi.org/10.1038/35016007

Mikolosko, J., Bobyk, K., Zgurskaya, H. I. & Ghosh, P. Conformational flexibility in the multidrug efflux system protein AcrA. *Structure* **14**, 577–587 (2006). https://doi.org/10.1016/j.str.2005.11.015

Murakami, S., Nakashima, R., Yamashita, E. & Yamaguchi, A. Crystal structure of bacterial multidrug efflux transporter AcrB. *Nature* **419**, 587–593 (2002). https://doi.org/10.1038/nature01050

Shi, X. *et al.* In situ structure and assembly of the multidrug efflux pump AcrAB-TolC. *Nat Commun* **10**, 2635 (2019). https://doi.org/10.1038/s41467-019-10512-6

Wang, Z. *et al.* An allosteric transport mechanism for the AcrAB-TolC multidrug efflux pump. *Elife* **6** (2017). https://doi.org/10.7554/eLife.24905

Phan, G. *et al.* Structural and dynamical insights into the opening mechanism of P. aeruginosa OprM channel. *Structure* **18**, 507–517 (2010). https://doi.org/10.1016/j.str.2010.01.018

Kulathila, R., Kulathila, R., Indic, M. & van den Berg, B. Crystal structure of Escherichia coli CusC, the outer membrane component of a heavy metal efflux pump. *PLoS One* **6**, e15610 (2011). https://doi.org/10.1371/journal.pone.0015610

Eicher, T. *et al.* Transport of drugs by the multidrug transporter AcrB involves an access and a deep binding pocket that are separated by a switch-loop. *Proc Natl Acad Sci U S A* **109**, 5687–5692 (2012). https://doi.org/10.1073/pnas.1114944109

Holm, L., Laiho, A., Törönen, P. & Salgado, M. DALI shines a light on remote homologs: One hundred discoveries. *Protein Sci* **32**, e4519 (2023). https://doi.org/10.1002/pro.4519

Dong, C. *et al.* Structural insights into the inhibition of type VI effector Tae3 by its immunity protein Tai3. *Biochem J* **454**, 59–68 (2013). https://doi.org/10.1042/bj20130193

Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439–d444 (2022). https://doi.org/10.1093/nar/gkab1061

Xu, K. *et al.* in *33rd AAAI Conference on Artificial Intelligence / 31st Innovative Applications of Artificial Intelligence Conference / 9th AAAI Symposium on Educational Advances in Artificial Intelligence.* 1230–1237 (2019).

Du, D. *et al.* Interactions of a Bacterial RND Transporter with a Transmembrane Small Protein in a Lipid Environment. *Structure* **28**, 625–634.e626 (2020). https://doi.org/10.1016/j.str.2020.03.013

Horne, J. *et al.* A lipoprotein partner for the Escherichia coli outer membrane protein TolC. *bioRxiv*, 2025.2003.2019.644130 (2025). https://doi.org/10.1101/2025.03.19.644130

Murakami, S., Nakashima, R., Yamashita, E., Matsumoto, T. & Yamaguchi, A. Crystal structures of a multidrug transporter reveal a functionally rotating mechanism. *Nature* **443**, 173–179 (2006). https://doi.org/10.1038/nature05076

Seeger, M. A. *et al.* Structural asymmetry of AcrB trimer suggests a peristaltic pump mechanism. *Science* **313**, 1295–1298 (2006). https://doi.org/10.1126/science.1131542

Delmar, J. A., Su, C. C. & Yu, E. W. in *Annual Review of Biophysics, Vol 43* Vol. 43 *Annual Review of Biophysics* (ed K. A. Dill) 93–117 (2014).

Yu, Z., Shi, X. & Wang, Z. in *Macromolecular Protein Complexes V: Structure and Function*   (eds J. Robin Harris & Jon Marles-Wright) 1–16 (Springer International Publishing, 2024).

Boyer, P. D. The ATP synthase - A splendid molecular machine. *Annual Review of Biochemistry* **66**, 717–749 (1997). https://doi.org/10.1146/annurev.biochem.66.1.717

Zgurskaya, H. I. & Nikaido, H. Cross-linked complex between oligomeric periplasmic lipoprotein AcrA and the inner-membrane-associated multidrug efflux pump AcrB from Escherichia coli. *Journal of Bacteriology* **182**, 4264–4267 (2000). https://doi.org/10.1128/jb.182.15.4264-4267.2000

Touzé, T. *et al.* Interactions underlying assembly of the Escherichia coli AcrAB-TolC multidrug efflux system. *Molecular Microbiology* **53**, 697–706 (2004). https://doi.org/10.1111/j.1365-2958.2004.04158.x

Szal, T. *et al.* Disassembly of the Escherichia coli AcrABZ-TolC efflux pump by ligand-mediated disruption of TolC-AcrA interfacial contacts. *bioRxiv*, 2025.2002.2018.638599 (2025). https://doi.org/10.1101/2025.02.18.638599

Zwama, M. *et al.* Multiple entry pathways within the efflux transporter AcrB contribute to multidrug recognition. *Nature Communications* **9** (2018). https://doi.org/10.1038/s41467-017-02493-1

Smith, B. L., Fernando, S. & King, M. D. Escherichia coli resistance mechanism AcrAB-TolC efflux pump interactions with commonly used antibiotics: a molecular dynamics study. *Scientific Reports* **14** (2024). https://doi.org/10.1038/s41598-024-52536-z

Su, C. C. *et al.* Conformation of the AcrB multidrug efflux pump in mutants of the putative proton relay pathway. *Journal of Bacteriology* **188**, 7290–7296 (2006). https://doi.org/10.1128/jb.00684-06

Yue, Z., Chen, W., Zgurskaya, H. I. & Shen, J. N. Constant pH Molecular Dynamics Reveals How Proton Release Drives the Conformational Transition of a Transmembrane Efflux Pump. *Journal of Chemical Theory and Computation* **13**, 6405–6414 (2017). https://doi.org/10.1021/acs.jctc.7b00874

Lei, J. & Frank, J. Automated acquisition of cryo-electron micrographs for single particle reconstruction on an FEI Tecnai electron microscope. *Journal of structural biology* **150**, 69–80 (2005). https://doi.org/10.1016/j.jsb.2005.01.002

Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017). https://doi.org/10.1038/nmeth.4169

Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and analysis. *Journal of computational chemistry* **25**, 1605–1612 (2004). https://doi.org/10.1002/jcc.20084

Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010). https://doi.org/10.1107/S0907444910007493

Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta crystallographica. Section D, Structural biology* **74**, 531–544 (2018). https://doi.org/10.1107/s2059798318006551

Brown, A. *et al.* Tools for macromolecular model building and refinement into electron cryo-microscopy reconstructions. *Acta Crystallogr D Biol Crystallogr* **71**, 136–153 (2015). https://doi.org/10.1107/s1399004714021683

DeLano, W. L. The PyMOL Molecular Graphics System. *on World Wide Web http://www.pymol.org* (2002).

Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**, e4792 (2023). https://doi.org/10.1002/pro.4792

Smart, O. S., Neduvelil, J. G., Wang, X., Wallace, B. A. & Sansom, M. S. HOLE: a program for the analysis of the pore dimensions of ion channel structural models. *J Mol Graph* **14**, 354–360, 376 (1996). https://doi.org/10.1016/s0263-7855(97)00009-x

Figures

Figure 1   Structure of the TolC–YbjP complex (closed state).

(A,B) Cryo-EM density map of the endogenous TolC–YbjP complex at 3.56 Å

resolution, shown in top and side view, spanning the outer membrane (OM) and

periplasmic space. TolC forms a homotrimeric "channel-tunnel" ~14 nm long, with its

12-stranded β-barrel embedded in the OM and periplasmic coiled-coils sealed at the tip.

YbjP (wheat) wraps around the equatorial domain of TolC, with continuous density

linking YbjP's N-terminal Cys19 into the OM micelle region.

(C,D) Top view (from the outer membrane) and side view of the TolC–YbjP cartoon model, showing three YbjP molecules arranged symmetrically around TolC's threefold axis.

(E,F) Each YbjP straddles the interface between two TolC protomers, occupying a groove on the TolC surface. TolC is shown as surface (each protomer in a different green color) and YbjP in wheat cartoon. YbjP's globular domain inserts between TolC protomers, while an N-terminal linker connects to the OM.

Figure 2   Architecture of the fully assembled TolC–YbjP–AcrABZ efflux pump.

(A,B) Top and side view of the cryo-EM density map (3.39 Å) of the endogenous TolC–YbjP–AcrABZ complex, with the inner membrane (IM) and outer membrane (OM) boundaries indicated (black bars). The map reveals the complete pump spanning 33 nm. Densities for TolC (green) are supposed to be in the OM region, AcrB (blue) in the IM region as a trimer, and six AcrA protomers (purple) forming an elongated barrel bridging TolC and AcrB. Three AcrZ protomers (pink) are seen at the periphery of the AcrB trimer in the IM region.

(C,D) Cartoon models of TolC–YbjP–AcrABZ in top view and side view. Three AcrB protomers (blue) form a trimeric base, surrounded by six AcrA molecules (purple) and

three AcrZ helices (pink). Each AcrB protomer consists of funnel, porter, and transmembrane domains.

 (E) Structural alignment of the two AcrA conformations (AcrA and AcrA*) reveals differences in the orientation of membrane proximal domain.

(F) Zoomed-in view of the TolC–AcrA interface, show the tight interaction between TolC's periplasmic helix-turn-helix motifs and AcrA's α-helical hairpin domain.

(G,H) Close-up views of the interface between AcrA (purple), AcrA* (light purple), and AcrB. Panel (g) shows the frontal cartoon representation; panel (h) displays a 60° rotated view, highlighting the intimate packing of AcrA with AcrB.

Figure 3 Conformational changes upon pump assembly: closed to open TolC transition with YbjP.

(A) Side view of TolC–YbjP complex. TolC is closed by coiled-coil helices.

(B) Side view of TolC–YbjP part in TolC–YbjP–AcrABZ complex. TolC is in open state.

(C) Comparison of TolC in the TolC–YbjP complex (closed state, forest green) and in the TolC–YbjP–AcrABZ pump (open state, pale green). Upon assembly with AcrABZ, these contacts are disrupted and the TolC helices tilt outward, enlarging the aperture. The OM β-barrel domain remains static. YbjP positions are consistent in two structures.

(D,E) Top views of TolC-YbjP in closed vs. open states. In the closed conformation, the coiled-coil helices bundle tightly, leaving a ~4 Å diameter sealed pore. In the open state (pale green), the helices are splayed apart, creating a ~20 Å diameter open channel.

(F) Quantitative comparison of pore radii in closed and open TolC, as computed using HOLE software[44].

Figure 4   Conformational cycling of AcrB monomers within the functional trimer.

 (A) Structural superimposition of the three AcrB protomers (L, T, O states) showing their sequential conformational transitions during the transport cycle.

(B) Conformational changes in the transmembrane domain associated with proton translocation.

(C) Substrate-binding pocket architecture in a single protomer, highlighting subdomains involved in drug recognition.

(D) Large-scale structural rearrangements in the porter domain that facilitate substrate translocation.

Figure 5   Proposed model of pump assembly and membrane anchoring.

Schematic illustration of the TolC–YbjP–AcrABZ efflux pump within the Gram-negative cell membrane. YbjP is hypothesized to associate with the outer membrane via its N-terminal Cys19, which may undergo lipidation *in vivo*. YbjP lies just beneath the outer membrane and above the peptidoglycan layer, according to prior electron tomography studies[13]. In the absence of AcrABZ, TolC adopts a closed state stabilized by inward-pointing periplasmic helices. AcrA, also likely lipid-anchored at its N-terminus, stabilizes the AcrBZ trimer near inner membrane. Upon engagement with AcrA's α-helical hairpin domain, TolC helices rotate outward in a left-handed (counterclockwise) manner to open the channel, and enables small-molecule substrates to be exported through the fully assembled complex.

Supplementary information for

Structural mechanisms of pump assembly and drug transport

in the AcrAB-TolC efflux system

Supplementary Figures S1 to S6

Supplementary Table S1

Supplementary Figures

Figure S1. Cryo-EM processing workflow for TolC-YbjP complex.

  A. Cryo-EM processing workflow.

  B. Local resolution distribution in final map[1].

  C. Gold-standard Fourier Shell Correlation curves for final reconstruction[2].

  D. Angular distribution plot for final map.

Figure S2. Local EM-density of YbjP in TolC-YbjP complex.

The region and some bulky residues are labelled. EM density threshold: 8σ.

Figure S3. Cryo-EM processing workflow for TolC-YbjP-AcrABZ complex.

A. Cryo-EM processing workflow.

B. Local resolution distribution in C1 final map[1].

C. Gold-standard Fourier Shell Correlation curves for C1 final reconstruction[2].

D. Angular distribution plot for C1 final map.

E. Local resolution distribution in C3 final map[1].

F. Gold-standard Fourier Shell Correlation curves for C3 final reconstruction[2].

G. Angular distribution plot for C3 final map.

Figure S4. Local EM-density of TolC and AcrA in TolC-YbjP-AcrABZ complex.

The region and some bulky residues are labelled. EM density threshold: 8σ.

Figure S5. Local EM-density of AcrB in TolC-YbjP-AcrABZ complex.

The region and some bulky residues are labelled. EM density threshold: 5σ.

Figure S6. Conformational transitions in AcrB's porter domain during the transport cycle.

Structural superposition of porter domain (subdomains PN1, PN2, PC1 and PC2) of monomers between L and T states and T and O states, revealing key conformational rearrangements.

Supplementary Tables

Table S1 Cryo-EM data collection, refinement and validation statistics

	TolC-YbjP	TolC-YbjP-AcrABZ	TolC-YbjP-AcrA (local)
Voltage (kV)	300		
Electron exposure (e$^-$/Å^2)	50		
Defocus range (μm)	-1.2～-1.8		
Pixel size (Å)	1.35		
Micrographs (no.)	3452		
Symmetry imposed	C3	C1	C3
Final particle images (no.)	63,022	31,000	17,043
Map resolution (Å) 0.143 FSC threshold	3.56	3.39	3.26
Map sharpening B factor (Å^2)	-134.5	-66.6	-74.7
CC (model vs. data)	0.7945	0.8727	0.8456
Model composition			
Chain count	6	18	12
Non-hydrogen atoms	13,539	53,464	29,061
Protein residues	1,752	7,037	3,816
B factors (Å^2)			
Proteins	40.66	99.69	115.5

R.m.s. deviations			
Bond lengths (Å)	0.006	0.003	0.006
Bond angles (°)	0.665	0.498	0.697
Validation			
MolProbity score	2.05	1.63	1.77
Clashscore	11.62	7.22	7.94
Poor rotamers (%)	0.41	0.12	0.13
Ramachandran plot			
Favored (%)	92.47	96.49	95.15
Allowed (%)	7.36	3.36	4.56
Disallowed (%)	0.17	0.16	0.29
PDB code	9V52	9V53	9V55
EMDB code	EMD-64784	EMD-64785	EMD-64787

References

1.   Kucukelbir A, Sigworth FJ, Tagare HD. Quantifying the local resolution of cryo-EM density maps. *Nat Methods* **11**, 63-65 (2014).

2.   Rosenthal PB, Henderson R. Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J Mol Biol* **333**, 721-745 (2003).