Structural mechanisms of pump assembly and drug transport in the AcrAB–TolC

efflux system

Running title: Cryo-EM structures of TolC–YbjP and TolC–YbjP–AcrABZ

Xiaofei Ge[1,2,*], Zhiwei Gu[1], Jiawei Wang[1,*]

[1]State Key Laboratory of Membrane Biology, Beijing Frontier Research Center for Biological

Structure, School of Life Sciences, Tsinghua University, Beijing 100084, P. R. China

[2]Health and Wellness, City University of Macau, Macau, 999078, P. R. China

*Correspondence should be addressed to: X.G. (gxf16@tsinghua.org.cn) or J.W.

(jwwang@tsinghua.edu.cn)

Abstract

Tripartite multidrug efflux pumps that span the cell envelope are essential for antibiotic

resistance in Gram-negative bacteria. Here, we report cryo-EM structures of two

endogenous efflux complexes from *Escherichia coli*: a TolC–YbjP subcomplex at 3.56 Å

resolution and the complete TolC–YbjP–AcrABZ pump at 3.39 Å. Structural analysis

reveals that YbjP, a previously uncharacterized lipoprotein, binds TolC in a 3:3

stoichiometry, bridging the TolC protomers at their equatorial domain. Clear density of

the mature YbjP's N-terminal Cys19 indicates that YbjP is anchored to the outer

membrane by an N-terminal lipid moiety. Notably, YbjP remains bound as TolC

undergoes AcrA-induced opening, suggesting that this accessory protein accommodates

the conformational change. The AcrB trimer simultaneously presents three distinct

conformational states (L, T, O), capturing a complete transport cycle. These high-

resolution structures provide insights into the architecture and mechanism of clinically

relevant efflux machinery, identifying YbjP as a previously unrecognized structural

component that contributes to TolC positioning and may assist in its membrane

localization.

Keywords

Multidrug efflux pump, TolC, YbjP, AcrA, AcrB, lipoprotein, antibiotic resistance,

Gram-negative

Introduction

Tripartite efflux pumps are essential for Gram-negative bacteria to expel a wide range of

toxic compounds, including antibiotics, across their dual-membrane cell envelope[1-5].

Among these systems, the AcrAB–TolC complex of *Escherichia coli* is one of the most

thoroughly studied[2,6]. It comprises the inner membrane RND (Resistance–Nodulation–

Division) transporter AcrB, the periplasmic membrane fusion protein AcrA, and the outer

membrane channel TolC[4,6-9]. AcrA, anchored to the inner membrane via N-terminal

lipidation, bridges AcrB and TolC, mediating conformational signaling essential for TolC

gating[10-12]. The assembling pathway of the tripartite pump involves the formation of an

AcrA–AcrB subcomplex, followed by the recruitment of TolC to form the complete

efflux apparatus[13]. In its isolated form, the periplasmic tip of TolC adopts a closed

conformation, stabilized by interactions between the α-helices[14-16]. Structural studies have

shown that TolC transitions from a closed to an open state in response to AcrAB

assembly, a process crucial for efflux activation[13,17].

Despite significant progress in structurally characterizing the AcrAB–TolC

system, fundamental questions remain regarding how TolC is positioned and retained in

the outer membrane. Unlike homologous channels such as *Pseudomonas aeruginosa*

OprM[18] and *E. coli* CusC[19], which possess covalently attached N-terminal lipids, TolC

lacks a lipid anchor. This raises the possibility that *E. coli* may utilize accessory factors to

localize and stabilize TolC during pump assembly. Furthermore, while prior studies using

crystallography and cryo-EM have revealed low-to-medium resolution snapshots of the

assembled pump[10-13,17,20], the limited resolution and conformational heterogeneity in

those structures impeded accurate modeling of flexible regions and may have masked the

presence of additional subunits.

Here, we address these knowledge gaps by determining two high-resolution cryo-

EM structures of the TolC-based efflux machinery in *E. coli*. The first is a 3.56 Å

structure of a TolC–YbjP subcomplex, and the second is a 3.39 Å structure of the fully

assembled TolC–YbjP–AcrABZ pump. These structures reveal a previously

uncharacterized outer membrane lipoprotein, YbjP, bound to TolC, and shed light on how

TolC is secured and transitions to the open state. Together, these findings enhance our

understanding of the architecture, assembly, and regulation of this clinically important

efflux pump.

Results

Structure of a TolC–YbjP closed-state complex

Endogenous purification of *E. coli* membranes yielded a stable assembly containing the

outer-membrane channel TolC (Figure S1A). The complex was purified directly from *E.*

coli as a stoichiometric assembly, without the need for artificial fusion or crosslinking.

Cryo-EM analysis resolved the complex at 3.56 Å resolution (Figures S1B-E,

Supplementary Table S1). TolC forms a homotrimer composed of a β-barrel that spans

the outer membrane and an elongated periplasmic α-helical domain (Figures 1A-D). The

periplasmic α-helical domain comprises an α-helical barrel extended by two helix-turn-

helix (HTH) motifs at its periplasmic extremity (TolC repeat 1: helices H3-H4; repeat 2:

helices H7-H8). These structural elements converge at an equatorial domain (Figures 1C-

D). The inward-bending HTH coiled-coils occlude the periplasmic tunnel, maintaining

TolC in a closed conformation—consistent with previous crystal structures[10] (Figure 1C).

An unexpected density at the periplasmic face of TolC indicated the presence of

an additional component (wheat in Figures 1A-B). An initial backbone trace, followed by

a DALI search[21] tentatively matched Tai3, a periplasmic type IV immunity protein

associated with the T6SS amidase effector Tae3 (PDB ID: 4HZ9)[22] from *Ralstonia*

pickettii. However, discrepancies in side chain density and species origin indicated that

the match was likely incorrect. Systematic screening of the AlphaFold Database[23] using

CryoNet[24] identified *E. coli* YbjP (UniProt P75818) as the top candidate, with all side

chains matching the experimental density (Figure S2). YbjP features a globular domain

and a structured N-terminal loop stabilized by a Cys36-Cys144 disulfide bond (Figures

1E-F). The N-terminal loop of YbjP adopts an extended conformation parallel to the

Helix 2 (residues 38-61) of TolC's α-helical barrel, forming substantial intermolecular

interfaces. Buried surface area (BSA) analysis reveals 787 Å² of contact with the primary

TolC protomer (green in Figure 1F), and 1037.4 Å² with the adjacent protomer (light

green), suggesting asymmetric binding energetics. The N-terminal density extends to

Cys19, which corresponds to the predicted signal peptide cleavage site, positioning the

lipoprotein at the periplasmic face of the outer membrane.

UniProt predictions indicate that YbjP is dually lipidated via N-palmitoyl and S-

diacylglycerol modifications, allowing its anchorage to the outer membrane. This

represents an elegant evolutionary solution, as while many Gram-negative TolC

homologs (*e.g.*, *P. aeruginosa* OprM[18], *E. coli* CusC[19], *et al.*) possess native lipid anchors

(Figure S3), *E. coli* TolC instead recruits YbjP as a dedicated membrane-tethering

partner. The YbjP globular domain nestles between adjacent TolC equational domains,

forming a stable 3:3 complex (Figure 1B). The identification of YbjP, along with its

defined membrane anchoring and structural compatibility with TolC, suggests a new

class of outer membrane partners that may regulate the TolC-dependent pathways.

Notably, the structure of TolC shows no conformational changes upon YbjP binding

when compared to the free, closed form of TolC[10].

Structure of TolC–YbjP–AcrABZ complex

An improved 3.39 Å resolution structure of the endogenous AcrAB–TolC complex from

E. coli was determined, revealing well-resolved density across the entire assembly

(Figure S4, Supplementary Table S1). This structure allows for the accurate modeling of

side-chain interactions throughout the tripartite channel, surpassing earlier models in

completeness and resolution[17] (Figures S5-6). YbjP remains bound to TolC[25], suggesting

that this lipoprotein is stably retained during pump assembly and functions as a molecular

clamp that anchors TolC to the outer membrane during AcrA and AcrB engagement

(Figures 2A-D).

The TolC–YbjP–AcrABZ complex adopts a funnel-like architecture spanning the

cell envelope, consistent with prior models of tripartite pumps[17] (Figure 2B). TolC,

tethered to YbjP, caps the outer-membrane end and docks onto a hexameric ring of AcrA

adaptors in the periplasm, which in turn surrounds the AcrBZ trimer embedded in the

inner membrane (Figures 2B, 2D). The full assembly is about 33 nm tall, comprising a 14

nm TolC channel, a 14 nm periplasmic portion of AcrAB, and a 5 nm transmembrane

portion of AcrBZ, matching the distance between the two membranes (Figure 2B).

Cryo-EM density reveals six AcrA molecules positioned beneath each TolC

trimer (Figure 2D), yielding a 3:6:3 stoichiometry for TolC:AcrA:AcrB. AcrA therefore

forms an elongated hexamer that bridges TolC and AcrB. This tripartite architecture is

stabilized by three distinct sets of interfaces: (i) contacts between the AcrB trimer and the

basal regions of AcrA, (ii) extensive AcrA–AcrA lateral interactions within the

hexameric ring, and (iii) tip-to-tip junctions formed between the upper AcrA α-helical

hairpin and the periplasmic entrance of TolC (Figure 2D). While each AcrA protomer

maintains the characteristic four-domain architecture—comprising the α-helical hairpin,

lipoyl, β-barrel, and membrane-proximal domains—functional asymmetry is observed in

their interactions (Figure 2E). The trimeric TolC, which contains six HTH motifs due to

internal repeats (TolC repeat 1/2), engages the AcrA hexamer via quasi-equivalent

binding: adjacent AcrA and AcrA* protomers interact differently with the intra- and

inter-protomer grooves of TolC, respectively (Figure 2F). The primary structural

difference between AcrA and AcrA* lies in the configurations of their membrane-

proximal (MP) domains (Figure 2E).

AcrB assembles as a homotrimer on the inner membrane, with each protomer

comprising three distinct domains: a funnel/docking domain, a porter/periplasmic

domain, and a transmembrane domain (Figure 2D). The funnel domain consists of two

subdomains, DN and DC (denoting the N- and C-terminal subdomains) (Figure 2G). The

porter domain is organized into four subdomains – PN1, PN2, PC1, and PC2 – named for

the N- and C-terminal subdomains. When viewed from the periplasmic side, these

subdomains are arranged in a clockwise orientation, with PN1 positioned closest to the

central axis of the trimer. In the hexameric AcrA assembly, the lipoyl and β-barrel

domains form two stacked concentric rings. While the lipoyl domains remain uninvolved

in AcrB binding, the β-barrel domains specifically engage the funnel domain of AcrB.

Strikingly, the membrane-proximal (MP) subdomains of AcrA exhibit asymmetric

binding: AcrA–MP interacts with the DC and PC1 subdomains of AcrB, whereas AcrA*–

MP contacts PN1 and DN via an extended loop, respectively (Figures 2G-H). This

differential engagement of MP subdomains establishes the structural basis for the

observed structural divergence between AcrA and AcrA* (Figure 2E).

Our density maps clearly resolve the small transmembrane protein AcrZ (49

amino acids) bound to each AcrB protomer (Figures 2B, 2D). AcrZ adopts a helical

structure that is embedded within a hydrophobic groove formed by the transmembrane

helices of AcrB at the predicted interaction site, forming an AcrB☐AcrZ☐ complex.

Although AcrZ is not essential for pump assembly, its consistent presence in

endogenously purified complexes and its role in stabilizing otherwise flexible regions—

particularly the transmembrane helices—suggest that AcrZ may serve an allosteric role in

modulating conformational dynamics for specific substrates[26]. Collectively, these

interactions illustrate the sophisticated assembly mechanism of the pump: AcrZ

reinforces AcrB's transmembrane domain, AcrB's funnel and porter domains anchor

AcrA, and AcrA's α-helical hairpins engage TolC. Notably, the holocomplex remained

intact throughout purification, indicating a high-affinity assembly between components,

consistent with prior *in vitro* reconstitution studies[25].

Structural rearrangements underlying TolC's closed-to-open transition

A comparative structural analysis of TolC in its closed and open states reveals a striking

iris-like dilation mechanism at the periplasmic entrance that facilitates transition to the

fully open conformation (Figures 3A-B)[27]. Throughout this conformational transition, the

transmembrane β-barrel and α-helical barrel domains maintain remarkable structural

rigidity, whereas the coiled-coil helices undergo dramatic rearrangement. These helices

pivot around the equatorial domain, undergoing a 27° rigid-body superhelical rotation

(Figure 3C), which constitutes a major structural rearrangement. In the closed state, a

complex network of interprotomer hydrogen bonds stabilizes the structure by constricting

the pore to its narrowest point at Asp396 (Figures 3D, 3F)[28,29]. Upon disruption of this

network in the open state, the pore expands significantly, reaching a diameter of

approximately 2 nm (Figures 3E-F). Notably, YbjP remains stably association with

TolC in both conformational states – the closed resting state and the open activated state

(Figure 3C). This persistent interaction suggests that YbjP serves as a structural scaffold:

anchoring TolC in the outer membrane, accommodating conformational changes during

activation, and functionally compensating for TolC's lack of intrinsic lipidation (Figure

S3). These findings not only support existing models of allosteric pump

activation[12,17,20,27,30], but also suggest how *E. coli* might utilize YbjP to fulfill the

membrane-anchoring role typically provided by intrinsic lipid modifications in other

bacterial TolC homologs, such as OprM and CusC[31,32].

Mechanism of substrate transport in the AcrB module

The AcrAB–TolC efflux system actively transports substrates against their concentration

gradient, moving them from regions of low to high concentration. This energetically

unfavorable process is driven by proton motive force (PMF), with the AcrB trimer

harnessing the electrochemical potential of protons to power substrate translocation[33].

Structural and functional studies support an asymmetric rotary mechanism[32], wherein

each AcrB protomer sequentially transitions through three conformational states: L

(loose), T (tight), and O (open)[20,34]. This concerted conformational rotation – analogous

to the functional cycling of F_1F_0 ATP synthase during ATP synthesis – enables

continuous vectorial transport through the pump complex[35].

Our high-resolution cryo-EM structure of the endogenous TolC–YbjP–AcrABZ

complex captures the AcrB trimer in three distinct conformational states (L, T, O),

providing key mechanistic insights (Figures 4A, S7). The detailed conformational

changes associated with this functional rotation mechanism are distinctly different from

those observed in isolated AcrB structures[36], even those determined in the native

membrane environment[37] (Figure S8).

The transmembrane domain of AcrB consists of N- and C-terminal

transmembrane repeats (N-TMR and C-TMR), encompassing TM1-TM6 and TM7-T12,

respectively (Figure 4B). Key proton relay residues – D407 and D408 on TM4, and K940

and R971 on TM10 – are located within these domains. While the transmembrane

domain remains largely conserved between the L and T states (Figure 4B, upper panel) –

maintaining salt bridges between Lys940 (TM10) and Asp407/Asp408 (TM4), the T→O

transition entails a coordinated rotation of TM2, TM4, TM5, and TM6, disrupting these

interactions (Figure 4B, lower panel).

In the L state, the PC1 and PC2 subdomains form a cleft that constitutes part of

the access pocket (AP). Substrate entry occurs between PC1 and PC2 in the L state

(Figure 4C, left panel). A conformational change to the T state is initiated by a random-

coil-to-helix transition in the upper part of TM8, coupled with an 8 Å shift of PC2 toward

PC1 (Figure 4D, middle panel and Figure S7). This displaces the switch loop that

separates the AP from the deep binding pocket (DBP), facilitating drug transfer from the

AP to the high-affinity DBP (Figure 4C, middle panel). Remarkably, PC1 remains static

and in contact with AcrA throughout the cycle (Figure S7). Transition to the O state is

triggered by the protonation of D407 and/or D408. This causes an inward-facing

movement of the N-TMR and C-TMR, bringing PC2 even closer to PC1 (Figure 4D,

right panel). Concurrently, PN1 and PN2 subdomains move closer together on the

transmembrane side, closing the DBP (Figure 4D, right panel). These collective

movements open a continuous tunnel that extends to the funnel domain, enabling

substrate extrusion (Figure 4C, right panel). The release of protons into the cytoplasm,

down their electrochemical gradient, subsequently restores the charge on D407 and/or

D408, resetting the transmembrane domain to its engaged state. This rotary mechanism,

reminiscent of F_1F_0 ATP synthase, is completed when the pump resets to the L state,

initiating a new catalytic cycle.

Discussion

Based on our structural data, we propose a model for the assembly and function of the

AcrAB–TolC multidrug efflux pump, emphasizing the anchoring and orienting role of

YbjP. As shown in Figure 5, the assembly begins with TolC priming. In the absence of

AcrAB, TolC is embedded in the outer membrane in a closed conformation, stabilized by

its interaction with YbjP. The lipid moiety of YbjP inserts into the inner leaflet of the

outer membrane, providing structural compensation for TolC's lack of intrinsic lipid

anchoring (Figure S3). Additionally, the periplasmic side of TolC may transiently interact

with the peptidoglycan layer, helping to position it within the periplasmic space. This

anchoring likely ensures optimal orientation of TolC's periplasmic end for interaction

with incoming AcrA.

Upon synthesis and membrane insertion, AcrB (complexed with AcrZ) and AcrA

form an inner membrane–periplasmic subcomplex[38,39]. A conserved cysteine residue in

AcrA, located near the inner membrane, may aid in anchoring or stabilizing the complex.

The complete tripartite pump likely assembles via stochastic encounters between the

endogenous AcrABZ subcomplex and the TolC–YbjP complex in the periplasm.

Stabilization likely occurs through tip-to-tip interactions between the hairpin domains of

AcrA and TolC[40]. YbjP may facilitate this docking by stabilizing TolC in a fixed,

properly oriented conformation at the outer membrane, thereby increasing its local

availability to the endogenous AcrABZ and lowering the energy barrier for tripartite

pump assembly, which is supported by the comparative structural analysis of TolC

homologs (Figure S3). In this context, YbjP may act as a structural placeholder that

guides AcrA toward the TolC entrance without directly modulating TolC's gating or

AcrABZ's conformational dynamics. YbjP's consistent presence in our cryo-EM

reconstructions of endogenous complexes and its predicted lipoprotein features together

suggest a specialized anchoring role for TolC (Figure S3). Based on our structural

observations and UniProt annotations, we propose that YbjP contributes to TolC

stabilization in the outer membrane and facilitates its spatial orientation for efficient

pump assembly.

269   Once assembled, the AcrABZ–TolC complex becomes active for drug efflux,

270 allowing substrates such as antibiotics and dyes to enter AcrB from the periplasm and be

271 expelled through TolC[41,42]. At the core of this process is the conserved D407–D408

272 proton relay pair in AcrB's transmembrane domain, which functions as both the proton-

273 binding site and the mechanochemical coupling hub[43,44]. Protonation of these residues

274 initiates conformational changes that propagate from the transmembrane domain to the

275 porter domain via an allosteric mechanism, which is refined in the context of the intact

276 endogenous pump complex.

277   The AcrB trimer operates via an asymmetric rotary cycle[20,34], with each protomer

278 sequentially transitioning through three distinct conformational states (L→T→O) to drive

279 peristaltic drug transport. This concerted conformational rotation – bearing striking

280 similarity to the catalytic cycle of F1F0 ATP synthase – enables continuous, vectorial

281 substrate transport against concentration gradients powered by proton translocation, and

282 our data reveal subtle yet functionally relevant conformational differences of this cycle in

283 the intact endogenous pump compared to isolated AcrB (Figure S8).

284   In conclusion, we identify YbjP as a previously unrecognized lipoprotein

285 associated with the AcrAB–TolC efflux system. Although its precise function requires

286 further validation, YbjP's structural positioning in the endogenous complexes suggests a

287 specialized role in anchoring TolC at the outer membrane and promoting tripartite pump

288 assembly (Figure S3), a process critical for bacterial multidrug resistance to antibiotics

289 and toxic compounds. Notably, our high-resolution cryo-EM structures capture all three

290 conformational states of the AcrB trimer (L, T, O) within the context of the fully

291 assembled, endogenous TolC–YbjP–AcrABZ complex for the first time. While these

states have been observed in isolated AcrB previously, our data reveal subtle yet

functionally relevant conformational differences that arise in the intact endogenous pump

(Figure S8), which are shaped by the native constraints of AcrA and AcrZ binding. This

study exemplifies how high-resolution cryo-EM, in combination with integrative

modeling, can reveal previously uncharacterized protein factors and enhance our

understanding of complex membrane-spanning systems in bacteria.

Method details

Cell preparation and membrane protein extraction

E. coli C600 cells were cultured to mid-log phase and then infected with a high-titer λ

phage to increase cellular stress. Cells were harvested by centrifugation and resuspended

in buffer containing 25 mM Tris-HCl and 150 mM NaCl (pH 8.0). Lysozyme was

added to a final concentration of 10 mg/mL, followed by incubation at 37 °C for

2 hours to facilitate cell lysis. Following removal of cell debris by centrifugation at

24,793 × g for 20 min, membrane proteins were extracted from the supernatant using 2%

n-dodecyl-β-D-maltoside (DDM). The membrane fraction was subsequently subjected to

size-exclusion chromatography using a SR6 Increase column. Fractions around 12 mL,

which were enriched for target proteins, were collected, diluted, and used for negative

staining. Transmission electron microscopy revealed the presence of target particles.

Cryo-EM grid preparation

The sample was concentrated to 0.7 mg/mL. A volume of 4 µL was applied to a glow-

discharged Au Quantifoil R1.2/1.3 300 mesh grid (glow discharge: medium setting,

30 s). To enhance the concentration of protein on the grid, sample loading was repeated

ten times: after each loading, excess solution was gently removed using a pipette tip, and

a fresh 4 µL aliquot was applied. The grid was then plunge-frozen in liquid ethane pre-

cooled with liquid nitrogen using a Vitrobot Mark IV (Thermo Fisher Scientific).

Blotting was performed for 3.5 s after a 30 s wait time, under 100% humidity at 8 °C.

Cryo-EM data collection

A total of 3452 movies were acquired using a 300 kV Titan Krios transmission electron

microscope (Thermo Fisher Scientific) equipped with a Gatan K3 Summit direct electron

detector. The calibrated pixel size was 0.675 Å. Each movie was recorded with a total

electron dose of ~50 e⁻/Å², and defocus values ranged from -1.2 to -1.8 μm. Data

collection was automated using the AutoEMation software[45].

Image processing

Motion correction and contrast transfer function (CTF) estimation were carried out in

CryoSPARC[46]. Poor-quality micrographs were manually excluded. 43,200 particles

corresponding to TolC were identified during 2D classification. Ab-initio reconstruction

and NU-refinement applied with C3 symmetry generated a reference model for further

picking. Template picking, *ab-initio* reconstruction in C1 symmetry, and NU-refinement

in C3 symmetry followed. The resulting map at 4.17 Å resolution was further improved

to 3.56 Å after reference-based motion correction (RBMC), CTF refinement, and NU-

refinement.

From the same dataset, TolC–AcrABZ particle selection was performed using

blob picking, followed by particle extraction and several rounds of 2D classification,

which yielded 19,396 high-quality particles. These were used for ab-initio reconstruction

and non-uniform (NU) refinement to generate an initial 3D model, which was

subsequently used for template picking. After template picking and further 2D

classification, a total of 35,092 particles were selected. Final refinement steps—including

NU-refinement, RBMC, and local CTF refinement – resulted in a 3D map at 3.23 Å

resolution. Since the AcrB subunit exhibits intrinsic structural asymmetry, C3 symmetry

was initially relaxed through symmetry expansion combined with 3D classification

without rotational/translational alignment. One predominant class was selected after

deduplication of symmetry-expanded particles. Local refinement in C1 symmetry better

resolved the asymmetric AcrB region. A subsequent masked 3D classification revealed a

particle subset containing distinct extra densities, which underwent local refinement with

re-applied C3 symmetry to enhance resolution of symmetric structural components.

Protein model building and structure refinement

The atomic models of TolC, AcrA, AcrB, AcrZ were rigid-body fitted into their

corresponding cryo-EM maps using UCSF Chimera[47]. To identify the extra densities

observed in both EM maps, a backbone model was manually traced. Protein identity was

assessed using the "Protein Searching without Sequence" function in CryoNet[24]. YbjP

(P75818) was ranked highest among *E. coli* proteins in the AlphaFold Database (AFDB,

strain 4364). Side-chain densities in the final EM maps were manually compared to the

predicted model, allowing confirmation of YbjP as the protein corresponding to the extra

density. Subsequent manual adjustments to the models were performed using COOT[48].

The refinement of the protein structures was conducted using either PHENIX[49] or

Refmac5[50]. Statistical details regarding the 3D reconstruction and model refinement

processes are provided in Supplementary Table S1. Visual representations of the

structures were generated using PyMol[51] or UCSF ChimeraX[52].

368   Acknowledgements

We are thankful to the Tsinghua University Branch of the China National Center for

Protein Sciences (Beijing) for their generous assistance with cryo-EM facility support

and computational resources on the Bio-Computing Platform cluster. Additionally, we

appreciate the valuable technical support from J. Lei, X. Li, and F. Yang. This work was

supported by the Science and Technology Development Fund, Macau SAR (FDCT)

(Grant No. 0112/2025/ITP2 to X.G.), the National Natural Science Foundation of China

(Grant Nos. 32501078 to X.G., and 32371254 to J.W.), and the Natural Science

Foundation of Beijing Municipality (Grant No. 5262009 to J.W.).

378   Author contributions

379   X.G. and J.W. conceived the project. X.G. and Z.G. optimized the preparation of cryo-

380   grids, recorded the cryo-EM data, and processed this data. J.W. built the atomic models.

381   X.G. and J.W. wrote the manuscript. All authors read and approved the final version of

382   the manuscript.

384   Declaration of interests

385   The authors declare no competing interests.

387   Data and code availability

388   Cryo-EM maps and the associated structural coordinates have been respectively

389   deposited into the Electron Microscopy Data Bank (EMDB) and the Protein Data Bank

390   (PDB) under the following accession codes: EMD-64784 / 9V52 (TolC–YbjP); EMD-

391   64785 / 9V53 (TolC–YbjP–AcrABZ); and EMD-64787 / 9V55 (TolC–YbjP–AcrA local)

Figures

Figure 1   Structure of the TolC–YbjP complex (closed state).

(A,B) Cryo-EM density map of the endogenous TolC–YbjP complex at 3.56 Å

resolution, shown in top and side view, spanning the outer membrane (OM) and

periplasmic space. TolC forms a homotrimeric "channel-tunnel" ~14 nm long,

with its 12-stranded β-barrel embedded in the OM and periplasmic coiled-coils

sealed at the tip. YbjP (wheat) wraps around the equatorial domain of TolC, with

continuous density linking YbjP's N-terminal Cys19 into the OM micelle region.

(C,D) Top view (from the outer membrane) and side view of the TolC–YbjP

cartoon model, showing three YbjP molecules arranged symmetrically around

TolC's threefold axis.

(E,F) Each YbjP straddles the interface between two TolC protomers, occupying

a groove on the TolC surface. TolC is shown as surface and cartoon (each

protomer in a different green color) and YbjP in wheat cartoon. YbjP's globular

domain inserts between TolC protomers, while an N-terminal linker connects to

the OM.

Figure 2   Architecture of the fully assembled TolC–YbjP–AcrABZ efflux pump.

(A,B) Top and side view of the cryo-EM density map (3.39 Å) of the endogenous

TolC–YbjP–AcrABZ complex, with the inner membrane (IM) and outer

membrane (OM) boundaries indicated (black bars). The map reveals the complete

pump spanning 33 nm. Densities for TolC (green) are supposed to be in the OM

region, AcrB (blue) in the IM region as a trimer, and six AcrA protomers (purple)

forming an elongated barrel bridging TolC and AcrB. Three AcrZ protomers

(pink) are seen at the periphery of the AcrB trimer in the IM region.

(C,D) Cartoon models of TolC–YbjP–AcrABZ in top view and side view. Three

AcrB protomers (blue) form a trimeric base, surrounded by six AcrA molecules

(purple) and three AcrZ helices (pink). Each AcrB protomer consists of funnel,

porter, and transmembrane domains.

(E) Structural alignment of the two AcrA conformations (AcrA and AcrA*)

reveals differences in the orientation of membrane proximal domain.

(F) Zoomed-in view of the TolC–AcrA interface, show the tight interaction

between TolC's periplasmic helix-turn-helix motifs and AcrA's α-helical hairpin

domain.

(G,H) Close-up views of the interface between AcrA (purple), AcrA* (light

purple), and AcrB. Panel (G) shows the frontal cartoon representation; panel (H)

displays a 60° rotated view, highlighting the intimate packing of AcrA with AcrB.

Figure 3   Conformational changes upon pump assembly: closed to open TolC transition.

(A) Side view of TolC–YbjP complex. TolC is closed by coiled-coil helices.

(B) Side view of TolC–YbjP part in TolC–YbjP–AcrABZ complex. TolC is in open state.

(C) Comparison of TolC in the TolC–YbjP complex (closed state, forest green) and in the TolC–YbjP–AcrABZ pump (open state, pale green). Upon assembly with AcrABZ, these contacts are disrupted and the TolC helices tilt outward, enlarging the aperture. The OM β-barrel domain remains static. YbjP positions are consistent in two structures.

(D,E) Top views of TolC–YbjP in closed *vs.* open states. In the closed conformation, the coiled-coil helices bundle tightly, leaving a ~4 Å diameter

sealed pore. In the open state (pale green), the helices are splayed apart, creating a

~20 Å diameter open channel.

(F) Quantitative comparison of pore radii in closed and open TolC, as computed

using HOLE software[53].

Figure 4   Conformational cycling of AcrB monomers within the functional

trimer.

 (A) Structural superimposition of the three AcrB protomers (L, T, O states)

showing their sequential conformational transitions during the transport cycle.

(B) Conformational changes in the transmembrane domain associated with proton translocation.

(C) Substrate-binding pocket architecture in a single protomer, highlighting subdomains involved in drug recognition.

(D) Large-scale structural rearrangements in the porter domain that facilitate substrate translocation.

Figure 5   Proposed model of pump assembly and membrane anchoring.
Schematic illustration of the TolC–YbjP–AcrABZ efflux pump within the Gram-negative cell membrane. YbjP is hypothesized to associate with the outer membrane via its N-terminal Cys19, which may undergo lipidation *in vivo*. YbjP lies just

beneath the outer membrane and above the peptidoglycan layer, according to prior

electron tomography studies[13]. In the absence of AcrABZ, TolC adopts a closed state

stabilized by inward-pointing periplasmic helices. AcrA, also likely lipid-anchored at

its N-terminus, stabilizes the AcrBZ trimer near inner membrane. Upon engagement

with AcrA's α-helical hairpin domain, TolC helices rotate outward in a left-handed

(counterclockwise) manner to open the channel, and enables small-molecule

substrates to be exported through the fully assembled complex.

Supplementary information for

Structural mechanisms of pump assembly and drug transport

in the AcrAB-TolC efflux system

Supplementary Figures S1 to S8

Supplementary Table S1

Supplementary Figures

Figure S1. Cryo-EM processing workflow for TolC-YbjP complex.

A. Schematic representation of the multi-step purification protocol employed to isolate the TolC-YbjP and TolC-YbjP-AcrABZ complexes.

B. Cryo-EM processing workflow.

C. Local resolution distribution in final map[54].

D. Gold-standard Fourier Shell Correlation curves for final reconstruction[55].

E. Angular distribution plot for final map.

Figure S2. Local EM-density of YbjP in TolC-YbjP complex.

The region and some bulky residues are labelled. EM density threshold: 8σ.

Figure S3. Structural comparison of the anchoring domains from six prototypical efflux pump outer membrane factors.

Cartoon representations of the protomer structures from *E. coli* TolC-YbjP (PDB: 9V52; this study), *K. pneumoniae* TOprJ1 (AlphaFold model/PDB: 9J3D)[56], *C. jejuni* CmeC (PDB: 4MT4)[57], *E. coli* CusC (PDB: 3PIK)[19,58], *P. aeruginosa* OprM (PDB: 3D5K)[18], and *V. cholera* VceC (AlphaFold model/PDB: 1YC9)[59] are shown. The conserved structural region responsible for anchoring the pump to the outer membrane is highlighted within the dashed box for each structure.

Figure S4. Cryo-EM processing workflow for TolC-YbjP-AcrABZ complex.

- **A.** Cryo-EM processing workflow.

- **B.** Local resolution distribution in C1 final map[54].

- **C.** Gold-standard Fourier Shell Correlation curves for C1 final reconstruction[55].

- **D.** Angular distribution plot for C1 final map.

- **E.** Local resolution distribution in C3 final map[54].

- **F.** Gold-standard Fourier Shell Correlation curves for C3 final reconstruction[55].

- **G.** Angular distribution plot for C3 final map.

Figure S5. Local EM-density of TolC and AcrA in TolC-YbjP-AcrABZ complex.

The region and some bulky residues are labelled. EM density threshold: 8σ.

Figure S6. Local EM-density of AcrB and AcrZ in TolC-YbjP-AcrABZ complex.

The region and some bulky residues are labelled. EM density threshold: 5σ.

Figure S7. Conformational transitions in AcrB's porter domain during the transport cycle.

Structural superposition of porter domain (subdomains PN1, PN2, PC1 and PC2) of monomers between L and T states and T and O states, revealing key conformational rearrangements.

A

3AOB-L   3AOC-L

C

4DX5-O

B

2DR6-T   4DX5-T   5NC5-T

D

9DXN-AcrB in native membrane

E

6TA5-OprM-MexA

Figure S8. Structural comparison of AcrB conformational states.

A. Comparison of the AcrB L state from the holocomplex with known L-state structures (PDB: 3AOB, 3AOC)[60]. Top panels show the funnel and porter domains (top view); bottom panels show a single protomer (side view).

B. Comparison of the AcrB T state with reference T-state structures (PDB: 2DR6, 4DX5, 5NC5)[17,20,31]. Panels show top views of the funnel and porter domains (top) and transmembrane domains (middle), and a side view of a protomer (bottom).

C. Detailed comparison of the AcrB O state with 4DX5[20], using the same three views as in (B).

D. Comparison of the AcrB structure from the holocomplex with an AcrB structure determined in the native membrane environment (PDB: 9DXN).

E. Comparison of the AcrB transmembrane domain with that of the homologous MexB (PDB: 6TA5)[30].

Supplementary Tables

Table S1   Cryo-EM data collection, refinement and validation statistics

	TolC-YbjP	TolC-YbjP-AcrABZ	TolC-YbjP-AcrA (local)
Voltage (kV)	300		
Electron exposure (e$^-$/Å^2)	50		
Defocus range (μm)	-1.2 ~ -1.8		
Pixel size (Å)	1.35		
Micrographs (no.)	3452		
Symmetry imposed	C3	C1	C3
Final particle images (no.)	63,022	31,000	17,043
Map resolution (Å) 0.143 FSC threshold	3.56	3.39	3.26
Map sharpening B factor (Å^2)	-134.5	-66.6	-74.7
CC (model vs. data)	0.7945	0.8727	0.8456
Model composition			
Chain count	6	18	12
Non-hydrogen atoms	13,539	53,464	29,061
Protein residues	1,752	7,037	3,816
B factors (Å^2)			
Proteins	40.66	99.69	115.5
R.m.s. deviations			
Bond lengths (Å)	0.006	0.003	0.006
Bond angles (°)	0.665	0.498	0.697
Validation			
MolProbity score	2.05	1.63	1.77
Clashscore	11.62	7.22	7.94
Poor rotamers (%)	0.41	0.12	0.13
Ramachandran plot			
Favored (%)	92.47	96.49	95.15
Allowed (%)	7.36	3.36	4.56
Disallowed (%)	0.17	0.16	0.29
PDB code	9V52	9V53	9V55
EMDB code	EMD-64784	EMD-64785	EMD-64787

References

1   Poole, K. Efflux-mediated antimicrobial resistance. *J Antimicrob Chemother* **56**, 20–51

         (2005). https://doi.org/10.1093/jac/dki171

2   Li, X. Z., Plésiat, P. & Nikaido, H. The challenge of efflux-mediated antibiotic resistance

         in Gram-negative bacteria. *Clin Microbiol Rev* **28**, 337–418 (2015).

         https://doi.org/10.1128/cmr.00117-14

3   Du, D., van Veen, H. W., Murakami, S., Pos, K. M. & Luisi, B. F. Structure, mechanism

         and cooperation of bacterial multidrug transporters. *Curr Opin Struct Biol* **33**, 76–91

         (2015). https://doi.org/10.1016/j.sbi.2015.07.015

4   Du, D. J. *et al.* Multidrug efflux pumps: structure, function and regulation. *Nature*

         Reviews Microbiology **16**, 523–539 (2018). https://doi.org/10.1038/s41579-018-0048-6

5   Blair, J. M., Richmond, G. E. & Piddock, L. J. Multidrug efflux pumps in Gram-negative

         bacteria and their role in antibiotic resistance. *Future Microbiol* **9**, 1165–1177 (2014).

         https://doi.org/10.2217/fmb.14.66

6   Wright, M., Kaur, M., Thompson, L. K. & Cox, G. A historical perspective on the

         multifunctional outer membrane channel protein TolC in Escherichia coli. *NPJ*

         Antimicrob Resist **3**, 6 (2025). https://doi.org/10.1038/s44259-025-00078-3

7   Ma, D. *et al.* Genes acrA and acrB encode a stress-induced efflux system of Escherichia

         coli. *Mol Microbiol* **16**, 45–55 (1995). https://doi.org/10.1111/j.1365-

         2958.1995.tb02390.x

8   Nikaido, H. Multidrug resistance in bacteria. *Annu Rev Biochem* **78**, 119–146 (2009).

         https://doi.org/10.1146/annurev.biochem.78.082907.145923

Okusu, H., Ma, D. & Nikaido, H. AcrAB efflux pump plays a major role in the antibiotic resistance phenotype of Escherichia coli multiple-antibiotic-resistance (Mar) mutants. *J Bacteriol* **178**, 306–308 (1996). https://doi.org/10.1128/jb.178.1.306-308.1996

Koronakis, V., Sharff, A., Koronakis, E., Luisi, B. & Hughes, C. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature* **405**, 914–919 (2000). https://doi.org/10.1038/35016007

Mikolosko, J., Bobyk, K., Zgurskaya, H. I. & Ghosh, P. Conformational flexibility in the multidrug efflux system protein AcrA. *Structure* **14**, 577–587 (2006). https://doi.org/10.1016/j.str.2005.11.015

Murakami, S., Nakashima, R., Yamashita, E. & Yamaguchi, A. Crystal structure of bacterial multidrug efflux transporter AcrB. *Nature* **419**, 587–593 (2002). https://doi.org/10.1038/nature01050

Shi, X. *et al.* In situ structure and assembly of the multidrug efflux pump AcrAB-TolC. *Nat Commun* **10**, 2635 (2019). https://doi.org/10.1038/s41467-019-10512-6

Pei, X. Y. *et al.* Structures of sequential open states in a symmetrical opening transition of the TolC exit duct. *Proc Natl Acad Sci U S A* **108**, 2112–2117 (2011). https://doi.org/10.1073/pnas.1012588108

Andersen, C. *et al.* Transition to the open state of the ToIC periplasmic tunnel entrance. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 11103–11108 (2002). https://doi.org/10.1073/pnas.162039399

Bavro, V. N. *et al.* Assembly and channel opening in a bacterial drug efflux machine. *Mol Cell* **30**, 114–121 (2008). https://doi.org/10.1016/j.molcel.2008.02.015

Wang, Z. *et al.* An allosteric transport mechanism for the AcrAB-TolC multidrug efflux pump. *Elife* **6** (2017). https://doi.org/10.7554/eLife.24905

Phan, G. *et al.* Structural and dynamical insights into the opening mechanism of P. aeruginosa OprM channel. *Structure* **18**, 507–517 (2010). https://doi.org/10.1016/j.str.2010.01.018

Kulathila, R., Kulathila, R., Indic, M. & van den Berg, B. Crystal structure of Escherichia coli CusC, the outer membrane component of a heavy metal efflux pump. *PLoS One* **6**, e15610 (2011). https://doi.org/10.1371/journal.pone.0015610

Eicher, T. *et al.* Transport of drugs by the multidrug transporter AcrB involves an access and a deep binding pocket that are separated by a switch-loop. *Proc Natl Acad Sci U S A* **109**, 5687–5692 (2012). https://doi.org/10.1073/pnas.1114944109

Holm, L., Laiho, A., Törönen, P. & Salgado, M. DALI shines a light on remote homologs: One hundred discoveries. *Protein Sci* **32**, e4519 (2023). https://doi.org/10.1002/pro.4519

Dong, C. *et al.* Structural insights into the inhibition of type VI effector Tae3 by its immunity protein Tai3. *Biochem J* **454**, 59–68 (2013). https://doi.org/10.1042/bj20130193

Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439–d444 (2022). https://doi.org/10.1093/nar/gkab1061

Xu, K. *et al.* in *33rd AAAI Conference on Artificial Intelligence / 31st Innovative Applications of Artificial Intelligence Conference / 9th AAAI Symposium on Educational Advances in Artificial Intelligence.* 1230–1237 (2019).

Horne, J. *et al.* A lipoprotein partner for the Escherichia coli outer membrane

       protein TolC. *bioRxiv*, 2025.2003.2019.644130 (2025).

       https://doi.org/10.1101/2025.03.19.644130

26   Du, D. *et al.* Interactions of a Bacterial RND Transporter with a Transmembrane Small

       Protein in a Lipid Environment. *Structure* **28**, 625–634.e626 (2020).

       https://doi.org/10.1016/j.str.2020.03.013

27   Alav, I. *et al.* Structure, Assembly, and Function of Tripartite Efflux and Type 1

       Secretion Systems in Gram-Negative Bacteria. *Chem Rev* **121**, 5479–5596 (2021).

       https://doi.org/10.1021/acs.chemrev.1c00055

28   Andersen, C., Koronakis, E., Hughes, C. & Koronakis, V. An aspartate ring at the TolC

       tunnel entrance determines ion selectivity and presents a target for blocking by large

       cations. *Mol Microbiol* **44**, 1131–1139 (2002). https://doi.org/10.1046/j.1365-

       2958.2002.02898.x

29   Marshall, R. L. & Bavro, V. N. Mutations in the TolC Periplasmic Domain Affect

       Substrate Specificity of the AcrAB-TolC Pump. *Front Mol Biosci* **7**, 166 (2020).

       https://doi.org/10.3389/fmolb.2020.00166

30   Glavier, M. *et al.* Antibiotic export by MexB multidrug efflux transporter is allosterically

       controlled by a MexA-OprM chaperone-like complex. *Nat Commun* **11**, 4948 (2020).

       https://doi.org/10.1038/s41467-020-18770-5

31   Murakami, S., Nakashima, R., Yamashita, E., Matsumoto, T. & Yamaguchi, A. Crystal

       structures of a multidrug transporter reveal a functionally rotating mechanism. *Nature*

       443, 173–179 (2006). https://doi.org/10.1038/nature05076

Seeger, M. A. *et al.* Structural asymmetry of AcrB trimer suggests a peristaltic pump mechanism. *Science* **313**, 1295–1298 (2006). https://doi.org/10.1126/science.1131542

Delmar, J. A., Su, C. C. & Yu, E. W. in *Annual Review of Biophysics, Vol 43* Vol. 43 *Annual Review of Biophysics* (ed K. A. Dill) 93–117 (2014).

Yu, Z., Shi, X. & Wang, Z. in *Macromolecular Protein Complexes V: Structure and Function*   (eds J. Robin Harris & Jon Marles-Wright) 1–16 (Springer International Publishing, 2024).

Boyer, P. D. The ATP synthase - A splendid molecular machine. *Annual Review of Biochemistry* **66**, 717–749 (1997). https://doi.org/10.1146/annurev.biochem.66.1.717

Lazarova, M. *et al.* Conformational plasticity across phylogenetic clusters of RND multidrug efflux pumps and its impact on substrate specificity. *Nat Commun* **16**, 11649 (2025). https://doi.org/10.1038/s41467-025-66751-3

Liu, H., Tse, C. M. & Dang, S. Capturing the native structure of membrane proteins using vesicles. *Proc Natl Acad Sci U S A* **122**, e2423407122 (2025). https://doi.org/10.1073/pnas.2423407122

Zgurskaya, H. I. & Nikaido, H. Cross-linked complex between oligomeric periplasmic lipoprotein AcrA and the inner-membrane-associated multidrug efflux pump AcrB from Escherichia coli. *Journal of Bacteriology* **182**, 4264–4267 (2000). https://doi.org/10.1128/jb.182.15.4264-4267.2000

Touzé, T. *et al.* Interactions underlying assembly of the Escherichia coli AcrAB-TolC multidrug efflux system. *Molecular Microbiology* **53**, 697–706 (2004). https://doi.org/10.1111/j.1365-2958.2004.04158.x

Szal, T. *et al.* Disassembly of the Escherichia coli AcrABZ-TolC efflux pump by ligand-mediated disruption of TolC-AcrA interfacial contacts. *bioRxiv*, 2025.2002.2018.638599 (2025). https://doi.org/10.1101/2025.02.18.638599

Zwama, M. *et al.* Multiple entry pathways within the efflux transporter AcrB contribute to multidrug recognition. *Nature Communications* **9** (2018). https://doi.org/10.1038/s41467-017-02493-1

Smith, B. L., Fernando, S. & King, M. D. Escherichia coli resistance mechanism AcrAB-TolC efflux pump interactions with commonly used antibiotics: a molecular dynamics study. *Scientific Reports* **14** (2024). https://doi.org/10.1038/s41598-024-52536-z

Su, C. C. *et al.* Conformation of the AcrB multidrug efflux pump in mutants of the putative proton relay pathway. *Journal of Bacteriology* **188**, 7290–7296 (2006). https://doi.org/10.1128/jb.00684-06

Yue, Z., Chen, W., Zgurskaya, H. I. & Shen, J. N. Constant pH Molecular Dynamics Reveals How Proton Release Drives the Conformational Transition of a Transmembrane Efflux Pump. *Journal of Chemical Theory and Computation* **13**, 6405–6414 (2017). https://doi.org/10.1021/acs.jctc.7b00874

Lei, J. & Frank, J. Automated acquisition of cryo-electron micrographs for single particle reconstruction on an FEI Tecnai electron microscope. *Journal of structural biology* **150**, 69–80 (2005). https://doi.org/10.1016/j.jsb.2005.01.002

Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017). https://doi.org/10.1038/nmeth.4169

Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605–1612 (2004). https://doi.org/10.1002/jcc.20084

Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010). https://doi.org/10.1107/S0907444910007493

Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531–544 (2018). https://doi.org/10.1107/S2059798318006551

Brown, A. *et al.* Tools for macromolecular model building and refinement into electron cryo-microscopy reconstructions. *Acta Crystallogr D Biol Crystallogr* **71**, 136–153 (2015). https://doi.org/10.1107/s1399004714021683

DeLano, W. L. The PyMOL Molecular Graphics System. *on World Wide Web http://www.pymol.org* (2002).

Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**, e4792 (2023). https://doi.org/10.1002/pro.4792

Smart, O. S., Neduvelil, J. G., Wang, X., Wallace, B. A. & Sansom, M. S. HOLE: a program for the analysis of the pore dimensions of ion channel structural models. *J Mol Graph* **14**, 354–360, 376 (1996). https://doi.org/10.1016/s0263-7855(97)00009-x

Kucukelbir, A., Sigworth, F. J. & Tagare, H. D. Quantifying the local resolution of cryo-EM density maps. *Nat Methods* **11**, 63–65 (2014). https://doi.org/10.1038/nmeth.2727

Rosenthal, P. B. & Henderson, R. Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J Mol Biol* **333**, 721–745 (2003). https://doi.org/10.1016/j.jmb.2003.07.013

Shi, Y. *et al.* Assembly and inhibition of transferable TMexCD1-TOprJ1 efflux pump. *Nat Commun* **16**, 10025 (2025). https://doi.org/10.1038/s41467-025-65038-x

Su, C. C. *et al.* Crystal structure of the Campylobacter jejuni CmeC outer membrane channel. *Protein Sci* **23**, 954–961 (2014). https://doi.org/10.1002/pro.2478

Su, C. C. *et al.* Crystal structure of the CusBA heavy-metal efflux complex of Escherichia coli. *Nature* **470**, 558–562 (2011). https://doi.org/10.1038/nature09743

Federici, L. *et al.* The crystal structure of the outer membrane protein VceC from the bacterial pathogen Vibrio cholerae at 1.8 A resolution. *J Biol Chem* **280**, 15307–15314 (2005). https://doi.org/10.1074/jbc.M500401200

Nakashima, R., Sakurai, K., Yamasaki, S., Nishino, K. & Yamaguchi, A. Structures of the multidrug exporter AcrB reveal a proximal multisite drug-binding pocket. *Nature* **480**, 565–U199 (2011). https://doi.org/10.1038/nature10641