Gene duplication dynamics and regulatory evolution shape the diversification of Asteraceae

Erika R. Moore-Pollard[1,2]*‡, Paige A. Ellestad[1]*, Brannan R. Cliver[3,4]*, Jakub Baczyński[5,6], Matthew D. Pollard[1,7,8], Zachary Meharg[9,10], Andrew C. Willoughby[11], Samantha Drewry[1], Alex Harkess[10], J. Mauricio Bonifacino[12], Zachary L. Nimchuk[11,13], John M. Burke[5,14], Daniel S. Jones[3,4], Jennifer R. Mandel[1,15]‡

[1]*Department of Biological Sciences, University of Memphis, 3700 Walker Ave, Memphis, Tennessee 38152, USA*

[2]*Department of Ecology and Evolutionary Biology, University of Connecticut, 75 N Eagleville Rd, Unit 3043, Storrs, CT, 06269, USA*

[3]*Department of Biological Sciences, Clemson University, 132 Long Hall Clemson, SC 29634, USA*

[4]*Department of Biological Sciences, Auburn University, 101 Rouse Life Sciences, Auburn, Alabama 36849, USA*

[5]*Department of Plant Biology, Miller Plant Sciences, University of Georgia, Athens, Georgia 30602, USA*

[6]*Institute of Evolutionary Biology, Faculty of Biology, Biological and Chemical Research Centre, University of Warsaw, Żwirki i Wigury 101, 02-089 Warsaw, Poland*

[7]*Department of Systems Biology, UMass Chan Medical School, 368 Plantation Street, Worcester, Massachusetts 01605, USA*

[8]*Department of Genomics and Computational Biology, UMass Chan Medical School, 368 Plantation Street, Worcester, Massachusetts 01605, USA*

[9]*Department of Crop, Soil, and Environmental Sciences, Auburn University, Funchess Hall, 350 S College St Suite 201, Auburn, AL 36849, USA*

[10]*HudsonAlpha Institute for Biotechnology, Huntsville, Alabama 35806, USA*

[11]*Department of Biology, University of North Carolina at Chapel Hill, 250 Bell Tower Dr Chapel Hill, NC 27708, USA*

[12]*Laboratorio de Botánica, Facultad de Agronomía, Universidad de la República, Av. Garzón 12 780, Montevideo C.P. 12900, Uruguay*

[13]*Curriculum in Genetics and Molecular Biology, University of North Carolina at Chapel Hill, 250 Bell Tower Dr Chapel Hill, NC 27599, USA*

[14]*The Plant Center, University of Georgia, Athens, Georgia 30602, USA*

[15]*ACRE: Institute for Agricultural and Conservation Research and Education, University of*

Memphis, Memphis, Tennessee 38152, USA

* These authors contributed equally to this work

‡ Corresponding authors

Authors' contact information:

‡ Erika R. Moore-Pollard, moore.erika.r@gmail.com, ORCID: 0000-0002-1182-9274

Paige A. Ellestad, pllestad@memphis.edu, 0000-0003-3930-3239

Brannan R. Cliver, bcliver@clemson.edu, 0009-0002-4183-2789

Jakub Baczyński, jjb67768@uga.edu, 0000-0001-5272-9053

Matthew D. Pollard, matthew.pollard95@gmail.com, 0000-0002-5171-1211

Zach Meharg, zmeharg@hudsonalpha.org, 0009-0004-7382-8527

Andrew C. Willoughby, andrew.willoughby@duke.edu, 0000-0003-0355-1214

Samantha Drewry, sgdrewry@memphis.edu, 0009-0002-0278-6804

Alex Harkess, aharkess@hudsonalpha.org, 0000-0002-2035-0871

J. Mauricio Bonifacino, bonifacinoj@fagro.edu.uy, 0000-0003-0377-3853

Zachary L. Nimchuk, zackn@email.unc.edu, 0000-0003-1826-0671

John M. Burke, jmburke@uga.edu, 0000-0002-1412-5539

Daniel S. Jones, dsj2@clemson.edu, 0000-0002-9241-1813

‡ Jennifer Mandel, jmandel@memphis.edu, ORCID: 0000-0003-3539-2991

Abstract

The flowering plant order Asterales exhibits a striking disparity in species richness, with >30,000

species in Asteraceae compared to <50 in its sister family Calyceraceae. To investigate the

genomic basis of this imbalance, we assembled three new chromosome-level genomes,

including the first for Calyceraceae, and re-annotated five additional Asterales genomes.

Comparative analyses revealed exceptionally high repeat content in both Asteraceae and

Calyceraceae, pervasive chromosomal rearrangements, and evidence for shared and lineage-

specific whole-genome duplications. In Asteraceae, tandem and dispersed duplications

disproportionately drove expansions of gene families linked to secondary metabolism and stress

response, while segmental duplicates bore signatures of adaptive selection for the regulation of

biosynthetic and metabolic processes. Selective pressures on regulators of flowering time, but

not on core flowering time genes, suggest an evolved balance between regulatory flexibility and

developmental constraint in floral diversification. Overall, these patterns suggest that, beyond

ancient polyploidy, small-scale duplications and selective fine-tuning of regulatory networks

underpinned the ecological versatility in Asteraceae, fueling its extraordinary diversification.

Keywords: Asterales, comparative genomics, Compositae, episodic positive selection, gene

family evolution, lineage-specific duplications, whole-genome duplication

Introduction

A recurring pattern in flowering plant evolution is the uneven distribution of species richness across the angiosperm tree of life, whereby species-poor lineages are often sister to highly diverse ones. While extinction and gaps in the fossil record may obscure some histories, the overall pattern remains clear[1]. Nowhere is this contrast more striking than in Asterales. The order includes ~35,800 species across 11 families, but >30,000 species belong to Asteraceae alone. Accounting for ~10% of all flowering plants and spanning nearly every continent and ecological niche (Fig. 1a), Asteraceae ranks among the most diverse and economically important plant families worldwide[2–8]. Over 2,000 of its members are used by humans for food, medicine, and ornamentals, yet it also ranks as the second most invasive plant family, exacting heavy ecological and economic costs[5,9]. In sharp contrast, its sister family, Calyceraceae, comprises fewer than 50 species confined to southern South America (Fig. 1a)[10–12]. This extreme disparity raises a central question: what genomic innovations fueled Asteraceae's extraordinary radiation?

Asteraceae boasts a long evolutionary history shaped by rapid radiation, polyploidization, hybridization, widespread ecological adaptation, and morphological innovation[3–7]. Its extensive diversification during the late Cretaceous has been largely attributed to enhanced reproductive success through pollinator attraction to its unique head-like inflorescence comprised of many individual florets, the capitulum[4,13,14]. Although Calyceraceae inflorescences also resemble capitula, important differences distinguish the two families, including the presence/absence of a terminal flower, fused/un-fused anthers, partially fused filaments attached to the corolla versus free filaments (Calyceraceae versus Asteraceae, respectively), and distinct mechanisms of pollen presentation[13,15,16]. Additionally, the production of diverse phytochemicals in Asteraceae facilitated ecological adaptation, particularly through defense responses[7]. These traits, coupled with a diversity of inflorescence types derived from distinct morphological and organizational modifications of the capitulum (Fig. 1b)[14], likely contributed to the disparity in species richness between the two families. The genomic mechanisms underlying their origin, however, have yet to be elucidated.

Acting across multiple scales, from individual genes to entire genomes, gene duplications have been a major driver of plant diversification by providing the raw genetic material for functional innovation[17]. Among these processes, whole-genome duplications (WGDs) are particularly pervasive throughout the evolutionary history of angiosperms and are widely recognized as a central force contributing to both their expansive diversity and widespread distribution[18–20]. The triplicate genome structure of Asteraceae suggests that the

family's diversification was catalyzed by two successive WGD events (i.e., paleohexaploidy) at
its core which was then followed by lineage-specific WGDs in some of its most species-rich
groups, e.g., tribe Cichorieae and the Heliantheae alliance[21–24]. Although limited by available
genomic data, evidence also suggests that the sister family Calyceraceae experienced the
same paleohexaploidy event[2–4]. This finding raises further questions about the role of
mechanisms beyond WGDs in plant diversification. Resolving the precise phylogenetic
placement of these events within Asterales is key to understanding the enormous evolutionary
success of Asteraceae.

Beyond WGDs, transposable elements (TEs) represent another dynamic force in plant
diversification, driving genomic change through chromosomal rearrangements and altering gene
expression via insertions or chromatin remodeling[25–27]. Asteraceae genomes, notable for their
exceptionally high repeat content dominated by TEs, highlight the likely role of TE activity in
promoting lineage-specific adaptations to diverse environments[28]. While both WGDs and TE-
mediated small-scale duplications have profoundly shaped plant evolution, their relative
contributions to diversification remain unresolved.

Despite the evolutionary and economic importance of Asteraceae, our understanding of
genome evolution in the family, and in the broader Asterales order, remains limited due to
sparse genomic and transcriptomic resources, beyond a handful of well-studied species (e.g.,
Helianthus [29] and *Lactuca* [24]). To date, assembled genomes are mostly concentrated to the
highly derived clades of Asteraceae with no representation of the early South American
lineages[30]. Similarly, the small sister family, Calyceraceae, lacks genomic data, leaving a major
gap in our understanding of the origin and early evolution of Asteraceae.

To investigate the genomic drivers of diversification across Asterales, we generated
three new chromosome-level assemblies of *Acicarpha procumbens* (the first Calyceraceae
genome), *Scaevola aemula* (Goodeniaceae), and *Platycodon grandiflorus* (Campanulaceae),
and re-annotated five publicly available genomes in Menyanthaceae (*Nymphoides indica*) and
Asteraceae (*Arctium lappa*, *Lactuca sativa*, *Chrysanthemum lavandulifolium*, and *Helianthus*
annuus) (Fig. 1c). In doing so, we address a central evolutionary question: How have WGDs
and small-scale gene duplications contributed to the origin and expansion of species diversity
within Asterales? Our analyses highlight lineage-specific patterns shaped by genome
duplications, chromosomal rearrangements, gene family dynamics, and signatures of selection
to provide a comparative framework for understanding the genomic basis of Asteraceae's
remarkable diversification.

Results

Extensive genome rearrangements yet similar composition in Asteraceae and Calyceraceae

We assembled chromosome-level genomes of *Platycodon*, *Scaevola*, and *Acicarpha*,

with *Scaevola* and *Acicarpha* assemblies being haplotype-resolved. The assemblies were highly

contiguous and complete based on BUSCO analysis, and comparable with the re-annotated

genomes (range = 96.6%–98%, mean = 97.6%; Supplementary Figs. 1 and 2, Supplementary

Tables 1 and 2). Genome rearrangements were widespread, with variations in chromosome

number indicating multiple chromosomal fission/fusion events across all Asterales taxa (Fig.

1c,d).

As expected, repeat content was high in Asteraceae (range = 80.9%–89.3%)[28], and

similarly elevated in *Scaevola* (Goodeniaceae, 82.1%) and *Acicarpha* (Calyceraceae).

Unexpectedly, *Acicarpha* exhibited the highest repeat proportion, reaching 90.1% (Fig. 1e,

Supplementary Table S2). In *Acicarpha*, the most abundant repeat type was *Ty1* (e.g., *copia*),

comprising 28.63% of the genome, though *Arctium* had the highest and *Nymphoides* had the

lowest *Ty1* content overall (33.82% and 10.84%, respectively). In contrast, *Helianthus* exhibited

the highest proportion of *Ty3* (e.g., *gypsy*) elements (45.4%), while *Chrysanthemum* had the

lowest (11.1%). The proportions of *Ty1*/*Ty3* retroelements show inverse patterns across the

phylogeny, with earlier-diverging species having higher percentages of *Ty1* and lower *Ty3* (e.g.,

Acicarpha), while more recently derived taxa show the opposite (e.g., *Helianthus*; Fig. 1e), a

pattern reported previously[28]. *Chrysanthemum* is the exception to this due to its low *Ty3*

percentage; however, it has the highest percentage of Unclassified LTRs, potentially causing a

decreased percentage in *Ty3* elements. Surprisingly, *Platycodon* and *Nymphoides* had an

abundance of DNA transposons (26.9% and 23.3%, respectively), which are not shared with the

rest of the species.

The proportions of duplicated gene types varied across Asterales genomes (Fig. 1d).

Segmental duplicates, likely derived from WGDs[32], were most common in *Acicarpha* (18.2%),

Arctium (19.1%)*,* and *Helianthus* (19.8%) and least in *Scaevola* (2.5%). In contrast, small-scale

duplicates (tandem, proximal, and dispersed), likely reflecting localized transposition or

recombination events[32], were most frequent in *Nymphoides* (12.9%) and showed modest

variation across other genomes (range = 7.6–10.8%, mean = 9.2%). Overall, *Acicarpha*'s

duplication frequencies fell within the range observed across Asteraceae (Fig. 1e).

Shared and lineage-specific WGD in Asteraceae and Calyceraceae

Asteraceae is hypothesized to have experienced one to multiple paleopolyploidization
events during its evolution[2–4,23,24], though the phylogenetic position of WGD events within the
family and related lineages remains uncertain. Syntenic gene depths corroborate the presence
of ancient WGDs for all taxa assessed, ranging from two (*Platycodon, Nymphoides,* and
Scaevola) to five (*Helianthus*), as compared to *Vitis vinifera* (an ideal reference due to its lack of
WGD since the gamma [γ] event shared by all eudicots[33]; Fig. 2a, Supplementary Table 3). In
Calyceraceae, we observed syntenic duplications and, to a lesser extent, triplications between
Acicarpha and *Scaevola* (Calyceraceae versus Goodeniaceae), and between *Acicarpha* and
Arctium (Calyceraceae versus Asteraceae), indicating lineage-specific duplications within
Asteraceae and Calyceraceae (Fig. 2b, Supplementary Table 4).
Comparing synonymous substitution rate (K_S) distributions across Asterales, we inferred
multiple duplications: an ancient WGD around K_S=2 (likely the γ event); a WGD shared by
Goodeniaceae, Calyceraecae, and Asteraceae; small-scale duplications in *Scaevola* and
Acicarpha; and separate WGDs in *Chrysanthemum* and *Helianthus* (Fig. 2c; Supplementary
Figs. 3 and 4). Alternatively, using K_S distributions adjusted to the rate of a focal taxon: three
small-scale duplications in *Nymphoides*; a WGD shared by Goodeniaceae, Calyceraceae, and
Asteraceae; two WGDs in *Scaevola*; two WGDs in *Acicarpha*; one WGD at the crown of
Asteraceae; one WGD in *Lactuca*; two in *Chrysanthemum*; and one in *Helianthus.* Prior to the
divergence of Asteraceae, both analyses indicated a WGD shared by Goodeniaceae,
Calyceraceae, and Asteraceae, but only the rate-adjusted method identified the Asteraceae-
specific WGD (Supplemental Figs. 5 and 6).
To further investigate WGDs, we phased the genomes and examined relationships
among duplicated subgenomes. Using phased sub-genomes that retained the most collinear
genes homologous to *Vitis* (Supplementary Datasets 1 and 2; Supplemental Figs. 7-13),
phylogenetic analyses generally revealed that *Acicarpha* and Asteraceae share at least one
duplication event, while each lineage also experienced independent duplications (Fig. 2d).
These findings were further supported by gene fractionation results, which indicated that
Acicarpha sub-genomes differ in age from those of Asteraceae, and are generally more recent,
although this trend was not always consistent (e.g., patterns of gene fractionation are distinct in
Lactuca but overlap in *Chrysanthemum*; Fig. 2e).
Given the variation in WGD signals across analyses, we performed hypothesis testing to
better place WGDs under various models. Under the critical model[34], which allows duplication
and loss rates to vary among branches, we found support for WGDs before and after the
Calyceraceae-Asteraceae split and lineage-specific events within *Acicarpha*, *Chrysanthemum*,

and *Helianthus*. Under the relaxed model[34], which allows rate heterogeneity among and within

branches, we found support for events in all tested lineages (Supplementary Fig.14). These

results indicate multiple WGD scenarios (Figs. 2f, 3), underscoring the difficulty of placing

Asteraceae WGDs. Some findings support earlier work that inferred two events: one shared

between Calyceraceae and Asteraceae and a second, independent duplication in each lineage

(Fig. 2f). However, other analyses support an older event predating the split of Goodeniaceae

and Calyceraceae+Asteraceae, followed by lineage-specific duplications in Asteraceae and

Calyceraceae (Fig. 2f), a pattern not previously reported.

Asteraceae gene family expansions comprised of mostly tandem duplicates

To assess how duplications contributed to Asteraceae diversification, we examined gene

orthogroup expansions across the family. Orthogroups that underwent substantial expansion at

the base of Asteraceae (n = 115; Supplementary Fig. 15) were significantly enriched in

processes related to stress response, signaling, and metabolic regulation, including cellular

response to amino acid stimulus, sesquiterpene biosynthesis, and cell surface receptor

signaling (Fig. 4a, Supplementary Fig. 16). Among the most expanded gene families were those

related to stress response and signaling (e.g. *GLP9*[35] and *WAKL2*[36]; Table 1), and to

biosynthesis of primary and secondary metabolites (e.g., *FAD2*, involved in fatty acid

synthesis[37]; and *CYP71B23*, involved in sesquiterpene synthesis and modification[38]; Table 1).

These were primarily tandem duplicates (30%), followed by dispersed (28%), proximal (23%),

and then segmental (18%) duplicates (Supplementary Fig. 17). Alternatively, *Acicarpha*

expansions were not enriched in macromolecule biosynthesis, pathways responding to

hormones like auxin and to environmental stimuli such as wounding or organic substances (Fig.

4b). Of these, multiple orthogroups included genes related to *CRK8*, a cysteine-rich receptor-

like kinase typically involved abiotic and biotic stress responses[39] (Table 1). Over half (62%) of

Acicarpha's expanded genes were dispersed duplicates, while 17% were tandem duplicates,

and only 4% were segmental duplicates (Supplementary Fig. 17).

The expansion of *FAD2* homologs in Asteraceae is hypothesized to have catalyzed

functional divergence[40] and supported the evolution of a unique N-C balance system in

conjunction with the loss of *PII,* a regulator of nitrogen-carbon assimilation[21]. The absence of *PII*

in *Acicarpha* and *Scaevola* suggests that this loss predates the Goodeniaceae split, clarifying

previous hypotheses about its timing. In contrast, *FAD2* expansion occurred only in Asteraceae

(Fig. 4c; Supplementary Dataset 3). Duplication patterns consisting of a few segmental genes,

with much larger proportions of proximal and tandem genes, mirrored previous research

suggesting that this expansion in Asteraceae arose via tandem duplications following genome
duplications[21] (Fig. 4c,d; Supplementary Fig. 18). This contrasts with Calyceraceae which has
fewer tandem and proximal duplicates (Fig. 4d). Also involved in metabolism, the expansion of
Asteraceae *CYP71B23* homologs involved a few segmental genes, but with higher proportions
of tandem, proximal, and dispersed genes (Fig. 4e,f; Supplementary Fig. 19). Overall, these
patterns indicate that gene family expansions, particularly those arising through tandem
duplications, have shaped metabolic and stress-response diversity in Asteraceae, linking
genome duplication and remodeling to the uneven diversification of Asterales.

Site-specific selection in Asteraceae acting on mostly segmental duplicates
Genes containing positively selected sites in all four Asteraceae taxa (n=586
orthogroups) were significantly enriched for the regulation of biosynthetic and metabolic
processes (Supplementary Fig. 20). Of these genes, segmental and dispersed duplicates
comprised 45% and 37%, respectively (Supplementary Fig. 17). In addition to the expansions in
the CYP superfamily—a large and functionally diverse gene family involved in secondary
metabolite synthesis, hormone regulation, and stress responses[41,42]—we observed positively
selected sites in all Asteraceae taxa for orthogroups related to *CYP96A1*, *CYP72A7*,
CYP706A4, *CYP706A6*, *CYP72A13*, and *CYP72A15* (Supplementary Dataset 3). Previous
studies also revealed strong lineage-specific diversification of CYP genes in Asteraceae,
indicating rapid evolutionary shifts, likely reflecting a central role in metabolic innovation[43].
To test for lineage-specific positive selection, we used the resulting 586 orthogroups with sites
under positive selection in Asteraceae species as input for branch model analyses. Of these, 47
showed evidence for episodic positive selection in an Asteraceae lineage (Supplementary
Dataset 3). Genes involved in the predominant biological processes associated with this gene
set (stress response and regulation of metabolic processes) include: *TMAC2*, a regulator of
abscisic acid and salt stress[44]; *WRKY27,* a transcription factor (TF) involved in plant defense
and floral development[45]; and *TCP5*, a TF that controls thermomorphogenesis[46], regulates
ethylene-mediated petal development and growth[47], and impacts leaf margin patterning[48] (Table
1). Other class II TCP TFs, *TCP2* and *TCP24*, also exhibited episodic positive selection in
Asteraceae. These genes regulate key developmental transitions and have been linked to floral
diversification in sunflower[49].

Lineage-specific selection in Asteraceae supports fine-tuning of flowering time
To focus on genes and pathways involved in floral innovation in Asteraceae, we
performed branch model selection analyses on a curated set of 335 flowering-associated

orthogroups (Supplementary Table 5). We identified 27 orthogroups with episodic positive

selection specific to all Asteraceae taxa (Supplementary Dataset 3) Of these, several

orthogroups contained genes involved in related pathways (photoperiod, autonomous, and

gibberellin) that regulate flowering time (Fig. 5). These genes function to directly or indirectly

suppress or activate *SOC1*, a floral integrator[50] that activates floral meristem identity genes,

such as *LFY* (a floral meristem identity transcriptional regulator that promotes flowering

transitions[51]) and *AP1* (a floral meristem identity MADS box gene[52]). Genes under selection in

Asteraceae were: *GI*, which promotes flowering via *FT* through the photoperiod pathway[53,54];

FVE, which promotes flowering by suppressing *FLC* in the autonomous pathway[55,56]; *CGA1* and

GNC, which delay flowering by suppressing *SOC1* in the gibberellin pathway[57,58]; *FD*, which

interacts with *FT* to activate *SOC1*; and *AGL24,* which integrates with *SOC1* to link floral signals

and promote flowering[59] (Fig. 5, Table 1). Of these, episodic positive selection was only

identified in *Acicarpha* homologs of *FD* and *CGA1*. Overall, selection on multiple genes across

interconnected pathways likely fine-tuned flowering time regulation in Asteraceae.

Discussion

While the role of WGD in driving gene expansion, diversification, and selection is evident

across our analyses, the number and timing of WGD events within Asterales remain difficult to

resolve despite having high-quality, contiguous genome assemblies. Contrasting results from

different approaches (Figs. 2f, 3) highlight the inherent challenges in detecting ancient WGDs,

particularly within large and diverse plant families. This complexity is compounded by processes

such as biased gene retention and loss, genome rearrangements, chromosomal fission and

fusion, incomplete lineage sorting, and hybridization, including allopolyploidy[60,61]. Synteny-

based methods, while powerful, often yield incomplete reconstructions of ancient events,

especially across distantly related taxa. Similarly, K_S-based approaches face limitations: signals

from ancient WGDs may be obscured by subsequent small-scale duplications and losses,

attenuated with time, or confounded with speciation signals, especially in cases of allopolyploidy

where increased variation in K_S values blurs inference[62].

Combining multiple forms of inference, our results support multiple hypotheses on WGD

placement in Asterales. Triplicate syntenic gene depths echo previous studies indicating a

shared paleohexaploidy event between Asteraceae and Calyceraceae [2–4] (Fig. 3a). However,

this inference was refined through sub-genome phasing and hypothesis testing to indicate that

Asteraceae and Calyceraceae shared a WGD, then each experienced their own (Fig. 3b). Some

K_S distribution and gene family expansion results suggest that the core WGD is shared not only

by Asteraceae and Calyceraceae, but also Goodeniaceae (Fig. 3c). However, this finding is

unlikely given the comparative syntenic gene depths and is probably an artefact of unaccounted

biological processes, such as small-scale duplications and biased gene retention[62].

Nonetheless, it is evident that both large-scale and small-scale duplications played a

formative role in the divergence and diversification of Asterales. While WGDs account for most

positively selected genes in Asteraceae, duplications originating from transposable element

activity (i.e., tandem, proximal, and dispersed duplicates) explained most gene family

expansions (Supplementary Fig. 17). A biased retention of segmental duplicates from

polyploidy, particularly those supporting complex processes and protein interactions, has been

observed in other systems, and the high proportion of segmental duplicates in Asteraceae and

Acicarpha may reflect this[32,63,64]. Rather than being lost or pseudogenized, retained duplicate

genes likely underwent subfunctionalization, neofunctionalization, or dosage amplification to

increase adaptive fitness. These findings highlight the substantial roles tandem duplication and

transposition, alongside WGD, have in generating genomic novelty[65].

In Asteraceae, post-duplication outcomes may have enhanced ecologically beneficial

processes such as secondary metabolite biosynthesis and cellular responses to environmental

stimuli, which are enriched in expanded and positively selected gene families (Supplementary

Fig. 20). Given Asteraceae's rich array of secondary compounds used in defense, attraction,

and environmental interaction[66], the lineage-specific retention and expansion genes involved in

metabolite production and regulation likely reflect specialization in metabolite biosynthesis.

Secondary chemistry contributes to key ecological interactions, including herbivore deterrence,

pathogen resistance, and pollinator attraction, and its diversification may have enhanced

environmental responsiveness and niche breadth[66]. Conversely, in *Acicarpha* the limited

expansion of metabolite biosynthesis genes may have limited their geographic range through

outcompetition. These patterns of gene family expansion, retention, and selection thus support

the broader view that chemical diversification, driven by underlying genomic innovation, has

been a major driver of Asteraceae's evolutionary success[5].

Advanced reproductive systems and strategies have also played a key role in

Asteraceae's evolutionary and ecological success. In a capitulum, flowering time and individual

floret development is precise and highly coordinated[13,14]. Our findings suggest that selective

pressures on duplicated genes and transposon activity may have fine-tuned gene regulation

associated with flowering time, conferring a key reproductive advantage in Asteraceae. Among

positively selected flowering time genes, patterns of evolution varied. For example, *FD*

homologs exhibited a two-step episode of positive selection: first on Asteraceae/*Acicarpha*

dispersed duplicates, then again on Asteraceae and *Acicarpha* duplicates individually. The *FVE*

homologs suggest that Asteraceae and *Acicarpha* segmental duplicates were derived from a

shared WGD, followed by positive selection on only one Asteraceae lineage. The patterns of *FD*

and *FVE* evolution reflect a shared history between Asteraceae and *Acicarpha*. Conversely, *GI*

and *CGA1* patterns reflect an Asteraceae-specific WGD. *GI* homologs showed two episodes of

diversifying selection in Asteraceae segmental duplicates, likely indicating sub-functionalization

or dosage compensation (Fig. 5). Among *CGA1* homologs, however, episodic positive selection

occurred only along one of the segmentally duplicated Asteraceae lineages. While likely

originating from transposon activity, rather than WGD, the dispersed duplicates of *Acicarpha*

CGA1 homologs exhibited a similar pattern of selection on one duplicated lineage. These

findings underscore distinct mechanisms of functional divergence and the fate of duplicates

between Asteraceae and *Acicarpha* that may have conferred unique adaptive advantages.

Furthermore, the absence of positive selection on core integrators (e.g. *SOC1*) and

meristem identity genes (e.g., *LFY* and *AP1*) suggests that strong purifying selection preserves

central flowering functions, while upstream regulators offer an evolutionary interface for

adjusting developmental timing. This pattern suggests that adaptive evolution in Asteraceae has

primarily targeted regulatory modifications within the flowering gene network, enabling flexibility

without disrupting deeply conserved functions. These patterns imply lineage-specific fine-tuning

of flowering time, likely in response to environmental heterogeneity. Because some genes in

traditional plant model systems, e.g., *FLC* in *Arabidopsis* (Supplementary Fig. 21), are lineage-

specific, these species may not serve as broadly representative models for flowering time

regulation. Instead, groups like Asteraceae and other diverse clades offer additional insights into

the evolutionary dynamics of these pathways.

Genome duplication has influenced not only speciation, but also the phenotypic and

ecological breadth of plants, shaping habitat use, life history strategies, competitive interactions,

responses to herbivores and pathogens, and pollinator relationships[67–72]. In both Asteraceae

and Calyceraceae, ancient large- and small-scale duplications left a strong genomic imprint, yet

distinct evolutionary pressures acting on duplicated genes, particularly those involved in

metabolite biosynthesis and flowering time regulation, drove the emergence of key ecological

traits and ultimately enabled the extraordinary diversification of Asteraceae.

Materials and Methods

<u>Taxon selection</u>

For this study, we generated de novo chromosome-level assemblies of *Acicarpha*

procumbens Less. (Calyceraceae)*,* hereafter *Acicarpha*; *Scaevola aemula* R.Br.

(Goodeniaceae)*,* hereafter *Scaevola*; and *Platycodon grandiflorus* (Jacq.) A.DC. 'Komachi'

(Campanulaceae), hereafter *Platycodon*, using the materials and methods described below. We

included additional publicly available Asterales species for comparison: *Arctium lappa* L.[73]

(Asteraceae; NCBI: PRJNA764011)*,* hereafter *Arctium*; *Lactuca sativa* L.[24] (Asteraceae; NCBI:

PRJNA173551)*,* hereafter *Lactuca*; *Chrysanthemum lavandulifolium* (Fisch. ex Trautv.)

Makino[74]*,* hereafter *Chrysanthemum* (Asteraceae; NCBI: PRJNA681093); *Helianthus annuus*

L.[29,75,76] (Asteraceae; HA412-HO v2: https://sunflowergenome.org/), hereafter *Helianthus*; and

Nymphoides indica (L.) Kuntze[77] (Menyanthaceae; CNGB: CNP0002767)*,* hereafter

Nymphoides. Additionally, for broader evolutionary and functional comparisons, we included the

well-studied genomes: *Vitis vinifera* L. cv. Cabernet Sauvignon cl. 08 v 1.1[78] (Vitaceae) ,

hereafter *Vitis,* and *Arabidopsis thaliana* (L.) Heynh. (Brassicaceae; TAIR10;

https://v2.arabidopsis.org/), hereafter *Arabidopsis* (Supplementary Table 6).

We collected seeds of *Acicarpha* in Uruguay, purchased seeds of *Platycodon* through

Park Seed (https://parkseed.com/), and acquired cuttings of *Scaevola* through North Carolina

Farms Inc. (https://www.ncfarmsinc.com/). We grew all plants in greenhouses at the University

of Memphis (Memphis, Tennessee, USA) and Auburn University (Auburn, Alabama, USA). A

single individual of *Platycodon* was taken through two successive rounds of inbreeding prior to

use. *Acicarpha* and *Scaevola* individuals were not inbred before sequencing. Vouchers for each

specimen of *Acicarpha*, *Scaevola*, and *Platycodon* are available at the University of Memphis

herbarium (MEM) and Auburn University Herbarium (AUA) (see Supplementary Table 7 for

details).

Sequencing

For *Acicarpha*, we generated short- and long-read DNA, Omni-C, and RNA sequencing

data. For short reads, genomic DNA was extracted from ~25 mg of fresh leaf tissue from a

single individual ground with a Bead Mill 24 Homogenizer (ThermoFisher Scientific, Atlanta,

Georgia, USA) using the E.Z.N.A. SP Plant DNA Kit (Atlanta, Georgia, USA). High molecular

weight (HMW) DNA for long-read sequencing was isolated from 2.5–3 g of flash-frozen young

leaf tissue ground in liquid nitrogen with an autoclaved mortar and pestle using the TaKaRa

NucleoBond HMW DNA kit (Düren, Germany). RNA was extracted from young leaf (YL), shoot

apical meristem (SAM), and mature leaf (ML) tissues using the E.Z.N.A. Plant RNA kit (Omega

Bio-tek, cat #R6827-01) and treated with RNase-free DNase (Omega Bio-tek, cat #E1091).

Flash-frozen tissue was sent to the HudsonAlpha Institute for Biotechnology (Huntsville,

Alabama, USA) for Omni-C library preparation using the Dovetail Omni-C kit (Dovetail

Genomics, Inc.). The gDNA, Omni-C libraries, and the RNA were sequenced using an Illumina

NovaSeq 6000 with PE150 chemistry, the HMW DNA was sequenced on a PacBio Sequel II at

HudsonAlpha Institute for Biotechnology.

For *Scaevola*, short- and long-read DNA, Hi-C, and RNA data were generated using

equivalent tissue amounts and extraction kits as for *Acicarpha*. Hi-C libraries were prepared with

the Proximo Hi-C Kit-Plant (Phase Genomics, cat #KT3040) and sequenced at the University of

North Carolina's High Throughput Sequencing Facility (UNC HTSF) on a NovaSeq 6000 (PE-50

and PE-100). The gDNA and HMW DNA were sequenced on NovaSeq PE150 and PacBio

Sequel II sequencers, respectively, at HudsonAlpha Institute for Biotechnology.

For *Platycodon*, we generated short- and long-read DNA, Hi-C, and RNA sequencing

data. Genomic DNA was extracted from ~100 mg of homogenized tissue using a modified

CTAB protocol. Short-read libraries were prepared at UNC's HTSF with the KAPA Hyper DNA

kit and sequenced at BGI-Genomics on the DNBseq platform (PE100). Total RNA from YL and

SAM tissues of five individuals was extracted using the E.Z.N.A. Plant RNA kit (Omega Bio-tek,

#R6827-01), treated with RNase-free DNase (#E1091), and used for library preparation (KAPA

stranded mRNA-seq kit, UNC HTSF). Hi-C libraries were prepared with the Proximo Hi-C Kit-

Plant (Phase Genomics, #KT3040) following the v4.0 protocol (October 2020) and sequenced at

UNC HTSF on a NovaSeq 6000 (PE100). For long reads, HMW DNA was extracted using the

ONT "High molecular weight gDNA extraction from plant leaves" (CTAB–Genomic-tip) protocol,

purified with Amicon Ultra filters (#UFC510008), and size-selected using the Circulomics Short

Read Eliminator Kit (#SS-100-101-01, v2). ONT libraries were prepared with the Ligation

Sequencing Kit (#SQK-LSK109) following previously described modifications[124] and sequenced

on a MinION Mk1B (MIN-101B) using FLO-MIN106D flow cells (R9.4.1 pore) controlled by

MinKNOW v3.6.5. Two flow cells were used; each washed three times with the Flow Cell Wash

Kit (#EXP-WSH003) and reloaded with the same library.

We checked raw long-read and short-read sequence data quality using FastQC v.

0.11.9, a program designed to highlight potential issues with sequence data. Additionally, we

checked raw long-read data quality using PAUVRE[82] v. 0.2.3, which plots the length and quality

distribution of raw Oxford Nanopore (ONT) or PacBio reads (Fig. S19). We then trimmed raw

reads to filter out inadequate reads and remove adapters using fastp[83] v. 0.23.2.

<u>Genome assembly and scaffolding</u>

We assembled long-read data for *Scaevola* and *Acicarpha* using HifiASM[84] v. 0.19.5 and

v. 0.18.5, respectively, with Hi-C data for *Scaevola* and Omni-C data for *Acicarpha* integrated

into the assemblies, resulting in two haplotypes per species. We base-called the *Platycodon*

MinION data using the python package guppy v. 3.1.6 (ONT) with model R9.4.1_450bps in

high-accuracy mode. We then assembled the genome with a hybrid approach utilizing ONT

long-read and DNBseq (BGI) short-read genomic DNA sequencing as inputs. For the hybrid

assembly, we used MaSuRCA[85] v. 3.4.2, and polished with short-read data using bowtie2[86] v.

2.1.4 and pilon[87] v. 1.22.

We aligned Omni-C and Hi-C reads to the assembly using BWA[88,89] v. 0.7.17 and used

Samtools[90] v1.11 to convert the output sam file into a bam file, sorted the bam file, and created

an index for the assembly fasta file to prepare for scaffolding. We conducted scaffolding,

followed by manual curation of the *Scaevola*, *Acicarpha*, and *Platycodon* assemblies, using Yet

another Hi-C Scaffolding tool (YaHS)[91] v1.1 for *Acicarpha* and v. 1.2a.2 for *Platycodon* and

Scaevola, with the Omni-C data for *Acicarpha*, Hi-C data for the *Scaevola*, and *Platycodon* used

as input. We prepared the resulting assemblies for Juicebox[92] (v. 2.17.00 for *Acicarpha* and v.

2.15.00 for *Scaevola* and *Platycodon*) visualization using the Juicer Pre tools v1.1 included in

YaHS before inputting them into the Juicebox GUI for manual scaffolding and correction to

produce pseudochromosomes. We assessed genome assembly quality and completeness

using Benchmarking Universal Single-Copy Orthologs (BUSCO)[93] v. 5.4.7 with embryophyta

from OrthoDB (www.orthodb.org) as the reference lineage. We then generated assembly

statistics of the finalized assemblies using the publicly available python script

(https://github.com/lexnederbragt/sequencetools/blob/master/assemblathon_stats.py).

<u>Genome annotation</u>

We screened the *Acicarpha*, *Scaevola*, and *Platycodon* genomes for bacterial

contaminants using StandardDB from Kraken2[94] v. 2.0.8 and removed all contigs with positive

hits. Using the resulting files as input, we then used the Extensive de-novo TE Annotator

(EDTA)[95] pipeline v. 1.9.3 to run RepeatModeler2[96] and identify remaining TEs utilizing the

default parameters, except --overwrite, --sensitive, --anno, and --evaluate were set to '1'. Using

the decontaminated genome file and the resulting TE library file from EDTA, we then ran

RepeatMasker[97] v. 4.1.5 to soft mask the genomes. We annotated the soft masked genomes

with RNA sequences using Braker3[98] v. 3.0.3 and labelled functional gene annotations using

InterProScan[99] v. 5.66-98.0. We then used OMArk v. 0.3.0 (https://omark.omabrowser.org/,

accessed June 3, 2024) to assess the proteome quality of *Acicarpha*, *Scaevola*, and *Platycodon*

with Asterids (NCBI ID: 71274) as the ancestral clade.

To ensure consistency between annotations and comparative genomic analyses, we re-

annotated five publicly available genomes from *Nymphoides, Arctium, Chrysanthemum,*

Helianthus, Lactuca following the steps detailed above. We assessed annotation quality across

all eight taxa using the gff3_stats python script

(https://github.com/HuffordLab/GenomeQC/blob/master/scripts/gff3_stats.py) from

GenomeQC[100].

<u>Plastid and mitochondrial genome assembly</u>

We extracted chloroplast sequences from all trimmed paired end short read data using

the GetOrganelle toolkit[101] following the recommended parameters for embryophyte plant

plastomes but increasing the number of rounds and the disentangle time limit (-R 75 --

disentangle-time-limit 7200). For long read data, PacBio and ONT minion, we used ptGAUL to

extract chloroplast sequences[102]. We then annotated complete circular sequences using

GeSeq[103] and assessed collinearity between plastomes using the progressive alignment in

Mauve[104]. In addition, we assembled mitochondrial genomes of *Platycodon, Scaevola,* and

Acicarpha using OatK v. 1.0 (https://github.com/c-zhou/oatk).

<u>Genome structure</u>

We characterized genomic sequences' size, heterozygosity, and repetitiveness by k-mer

frequency analyses (k=21) using Jellyfish[105] and GenomeScope[106] 2.0. We assessed

heterozygous k-mer pairs, obtained with KMC3[107], using Smudgeplot[106] to estimate ploidy levels

and infer genomic complexity in terms of sub-genome composition. We identified lower (L) and

upper (U) end cut-off values, below and above which all k-mers were discarded as errors, using

k-mer coverage output from Genomescope 2.0, as recommended in the Smudgeplot

documentation (https://github.com/KamilSJaron/smudgeplot), where L = (kcov/2)-5.

<u>Gene duplications</u>

We investigated gene duplications of all eight target Asterales genomes, plus genomes

from *Vitis* and *Arabidopsis*, by assessing homology and synteny to identify duplicated genes

and genomic regions, then by analyzing K_S values to estimate the relative age of duplication

events. To quantify and classify types of gene duplications within each genome, we used the

"duplicate gene classifier" program in MCScanX[32]. Through this program, duplicated genes

were classified as: segmental, proximal, tandem, or dispersed, as defined by [32].

We utilized OrthoFinder v. 2.5.5[108] to identify conserved orthologs and gene duplications

across the ten genomes, using the -M msa parameter to generate a multiple species alignment.

We then identified the best evolutionary model and performed a maximum likelihood

phylogenetic analysis on the alignment using IQTree[109] v. 2.1.3 with 1,000 bootstraps to

generate a species tree. We used the orthologous gene groups identified by OrthoFinder for

downstream analyses to investigate evolutionary processes, however, we filtered the groups to

exclude multiple isoforms of the same gene by keeping only the longest isoform.

Syntenic gene depths

To investigate putative paleopolyploidy, we assessed synteny among the eight Asterales

genomes. We used the JCVI[110][111] v. 1.3.8 python function "compara catalog ortholog" with the

default c-score cutoff of 0.7 to identify syntenic regions, then generated genome-wide synteny

plots of the full eight genomes following phylogenomic placement. We evaluated pairwise

syntenic gene depth between all genomes using the "compara synteny depth" option. We then

used these inferred values as input for subsequent analyses to phase subgenomes and further

elucidate polyploid origins (detailed in "Phasing subgenomes" below).

Phasing subgenomes

We implemented the two most prominent software tools to identify and separate

subgenomes, SubPhaser[112] and Whole-Genome Duplication Integrated (WGDI[111]). However,

given that the tested taxa have not experienced recent polyploidization events and are currently

recognized as diploids, SubPhaser could not be used with our data as it is unable to accurately

phase genomes that experienced paleopolyploidy events. Instead, we used WGDI v. 0.6.5 to

further investigate polyploidization events across Asterales using *Vitis* as a reference. Steps to

investigate polyploidy followed the subgenome aware analyses available on GitHub

(https://github.com/SunPengChuan/wgdi-example/blob/main/phase_subgenomes.md). Multiple

chromosomes of *Vitis* were tested as a reference for phasing, though only the chromosome with

the largest gene retention and collinearity, chromosome 11, was used for further analyses. Age

estimates from K_S values (e.g., Fig. S23) were used to help separate subgenomes if they were

one the same chromosome or were not distinct. We determined phylogenomic relationships

among the subgenomes using WGDI's '-at' parameter to generate gene trees with IQTree and

then ASTRAL-III[113] v. 5.7.8 to generate a final subgenome coalescent tree. We also assessed

gene fractionation within each subgenome to determine the evolutionary age of each

subgenome using the 'r' parameter of WGDI. We then reanalyzed the resulting subgenomes

within Asteraceae, Calyceraceae, and Goodeniaceae in JCVI to investigate synteny across

these phased regions using methods detailed above.

<u>Accelerated gene evolution in *Acicarpha* and Asteraceae</u>

To identify major targets of accelerated evolution within conserved genomic regions, we

used Cactus[114–116] v2.5.1, to generate a reference-free alignment of the Asterales subgenomes

that are syntenic to chromosome 11 of *Vitis*. Using the inferred phylogenomic relationship

between subgenomes as the guide tree to inform hierarchical alignment, we performed three

separate tests for accelerated evolution at different positions in the subgenome tree: across all

Asteraceae and *Acicarpha* subgenomes (clade 1, Fig. S11), in the *Acicarpha*-specific

duplication event (clade 2, Fig. S11), and in *Acicarpha* within the Subgenome 1 clade (clade 3,

Fig. S11).

Specifically, we created a neutral model of sequence evolution based on ancestral

repetitive sequences[117]. Repetitive elements were identified within the most ancestral sequence

in the subgenome alignment using RepeatModeler and RepeatMasker[97]. A random sample of

100,000 sites from these ancestral repeats was then selected to generate a multiple alignment

file (MAF) of repetitive sites across all subgenomes. This MAF was used to construct the neutral

model with PHAST phyloFit[118,119], using default settings and the same phylogeny used to

generate the subgenome alignment (Fig. S11).

With this neutral model, we identified conserved regions within the Asterales

subgenomes using PHAST phastCons[118,119], with *Vitis* as the reference. This was performed

separately for each of the three clade-specific analyses, with conserved regions detected in the

outgroup subgenomes. For each analysis, we obtained a list of the most-conserved elements

and converted their log-odds scores to length-normalized log-odds ratios. Elements with

normalized log-odds ratios below the 30th percentile were excluded from further analyses[120].

This filtering step produced three distinct lists of conserved regions in outgroup subgenomes,

allowing us to assess accelerated evolution within our clades of interest. We then tested for GC-

biased gene conversion (gBGC) within our ingroups using PHAST phastBias[118] and considered

any region with a score >0.5 to be impacted by gBGC. We filtered our lists of conserved regions

to remove any elements that were impacted by gBGC.

We used the likelihood ratio test of PHAST phyloP[118,121] to test the conserved elements

for accelerated evolution within our clades of interest. We corrected for multiple hypothesis

testing using Storey's correction method[122] (FDR=0.05). This produced a separate list of

accelerated regions (ARs) for each of our clade-specific analyses. To assess the biological

impacts of the accelerated evolution detected in each of our analyses, we identified the genes

most likely to be impacted by each AR. We associated an AR with a gene if it overlapped an

exon. Exon identities and locations were obtained from annotations of the *Vitis* genome[78].

ARs can be distributed unevenly across the genome, often clustering into regions of high

density, known as AR hotspots. To detect these hotspots, we employed the permutation method

of Cahill et al.[123]. Specifically, we used a sliding window approach, counting ARs in 100 kb

windows along *Vitis* chromosome 11, with windows offset by 10 kb steps. We conducted 100

permutations, each time randomly designating N conserved elements as accelerated, where N

corresponds to the actual number of ARs on the chromosome. For each permutation, we

repeated the sliding window approach to determine the maximum simulated AR density in a 100

kb window. Regions were designated as AR hotspots if their true AR density exceeded the 95th

percentile of maximum AR densities observed across all permutations. This approach prevents

regions with high conserved element density from being falsely identified as hotspots. We

identified separate hotspots for each of our clade-specific analyses.

<u>K_S distributions</u>

To investigate the influence of major genome duplication events and positive selection

on Asterales genomes, we assessed the non-synonymous (K_a) and synonymous (K_S)

substitution rates between gene pairs. Peaks in K_S value distributions, indicating the presence

of duplicated genes, were used to infer histories of WGD. For homologous gene pairs, we

estimated K_S values using an established pipeline[124] in which orthologs were aligned using

MUSCLE[125], then converted into nucleotides using PAL2NAL[126], then K_S values were calculated

between gene pairs using the F3X4 model in codeml within PAML[127]. We plotted K_S values of

paralogs and pairwise orthologous gene pairs in R[128,129] with kernel density estimations using

geom_density() from ggplot2[130]. Additionally, we used output from these analyses to identify

signatures of positive selection ($K_a/K_S > 1$), neutral selection ($K_a/K_S \approx 1$), and purifying selection

($K_a/K_S < 1$) of paralogous gene pairs. To more accurately compare estimations of divergence

derived from orthologous K_S distributions with estimations of WGD derived from paralogous K_S

distributions, we then re-scaled ortholog distributions to the paralog scale of each focal species

using *ksrates*[131]. For these analyses, *Platycodon* was used as the outgroup, therefore, WGD

events were not assessed for this lineage.

<u>WGD hyphothesis testing</u>

Directly following previously published methods and scripts[132,133]

(https://gitlab.com/barker-lab/ajb_special_issue_2024.git), we tested predicted WGD events,

hypothesized by previous literature and identified by *ksrates*, using WHALE[34]. WHALE is a tool

designed to test WGD hypotheses by analyzing the relative support for different evolutionary

scenarios. It uses gene tree topologies and considers the expected patterns of gene retention

and loss following WGD events. By applying a likelihood-based framework, WHALE evaluates

whether observed gene tree discordance is best explained by WGD or alternative models,

making it a valuable approach for investigating genome evolution. Given results from *ksrates*

and primary literature, we tested nine total WGD events split into three separate analyses: one

based solely off findings from previous studies, the second analysis used only the results of

ksrates, and the third analysis combined results of the first two analyses (Fig. S15).

<u>Evolution of gene family size</u>

To investigate the evolution of gene family sizes in Asterales, we reconstructed ancestral

states of the orthologous groups identified by OrthoFinder by employing the fast algorithm[134]

through the 'ace' function in the *phytools*[135] R package. From the reconstructed ancestral states,

we applied targeted filters to classify orthogroups as: 1) specific to *Acicarpha*, 2) lost in

Acicarpha, 3) specific to Asteraceae (evolved in the ancestral branch of Asteraceae and absent

from other lineages), 4) lost in the last common ancestor of Asteraceae, 5) specific to both

Asteraceae and *Acicarpha*, and 6) lost in the last common ancestor of both Asteraceae and

Acicarpha. Filtering was based on a Bayes factor threshold of 10, indicating that the likelihood of

an orthogroup being present or absent at a given node was ten times higher than the alternative

scenario.

Next, we used CAFE[136] v. 5.0 to further investigate the dynamics of orthogroup

expansion and contraction. Since CAFE requires an ultrametric tree, we used TreePL[137,138] to

calibrate the phylogeny reconstructed by OrthoFinder. We selected four calibration points based

on previous work[6], constraining the minimum ages of ancestral nodes for the following groups:

Asteraceae and Calyceraceae (*Tubulifloridites lillei* Type A[139]; 72.1 Mya), all Asteraceae

included in the study (*Raiguenrayun cura*[140]; 47.5 Mya), *Helianthus* and *Lactuca* (*Cichorium*

intybus type pollen[141]; 22 Mya), and *Helianthus* and *Chrysanthemum* (*Artemisia* type pollen[142];

31 Mya). The root age was constrained between 121 and 126 Mya using a secondary

calibration point for Pentapetalae[143] (Fig. S24). The optimal smoothing parameter was

determined through cross-validation, testing 20 values across a range of magnitudes (from 1e-

10 to 1e+09) to ensure the best fit.

The tested CAFE models varied in the number of gamma parameters (ranging from 1 to

3), reflecting assumed differences in expansion and contraction rates across orthogroups.

Orthogroups identified by the best-fitting model as significantly expanded or contracted (Viterbi

≤ 0.05) in Asteraceae, Acicarpha, and both Asteraceae and Acicarpha were retained for

downstream analyses, in addition to those identified as lost or specific to these groups. Genes

from significantly expanded orthogroups were then blasted to the *Arabidopsis thaliana*

database, TAIR10, using BLASTP[144] to determine closest gene similarity and trees were then

visualized in R.

<u>Positive selection</u>

We further investigated positive selection across Asterales genomes, focusing on

Asteraceae, using site-specific and branch-specific methods. For all taxa, we assessed

orthology and subsequent positive selection of single-copy orthologous groups using site-

specific methods within the AlexandrusPS pipeline[145]. For the orthogroups which were found to

exhibit positive selection for genes pertaining to all four Asteraceae taxa and no others, we

analyzed further by using branch-specific methods to determine if the found signatures of

positive selection originated from taxon-specific or family-specific lineages. Additionally, we

assessed a subset of orthogroups relating to floral development for lineage-specific positive

selection using branch-specific methods. These floral-related orthogroups were chosen based

on GO annotations for *Arabidopsis thaliana* genome (TAIR10: available at

https://v2.arabidopsis.org/) to include 18 terms related to meristem maintenance and

organization (GO:0010582, GO:0009933, GO:0009934, GO:0048507, GO:0010074,

GO:0019827), flower development (GO:0009908, GO:0010076, GO:0009910, GO:009911,

GO:0048438, GO:0010093, GO:2000032), vegetative shoot system/inflorescence development

(GO:0048367, GO:0010077, GO:0010492, GO:0090506,), and vegetative-reproductive phase

transition (GO:0048510). For both subsets of orthogroups, we generated codon alignments from

coding sequences by adding gaps for frameshifts and replacing stop codons with gaps using

MACSE[146]. Then we used the aBSREL (adaptive Branch-Site Random Effects Likelihood)

function in HyPhy[147] to test if positive selection has occurred on a proportion of branches. We

inspected results using HyPhy Vision (http://vision.hyphy.org/aBSREL), then visualized resulting

gene trees in R using Ape 5.0[148] and ggtree[149].

<u>GO enrichment</u>

For inferences on the functional role of genes identified in these analyses, we used the

basic local alignment search tool (BLAST[144]) to compare target proteins to a TAIR10 protein

database and identify matches based on an e-value threshold of 1e-10. For groups of gene

groups identified to be expanded or contracted or under positive selection, we performed gene

ontology (GO) enrichment analyses to summarize the significant GO terms for biological

processes[150]. We used the clusterProfiler[151] package in R, with annotations based on BLASTP

best-hit matches to the TAIR10 database. To account for multiple testing, p-values were

adjusted using the Benjamini–Hochberg false discovery rate (FDR) correction. Enrichment was

considered significant at an adjusted p-value (P-adjust) threshold of < 0.01.

Acknowledgements

We thank Jane Grimwood (HudsonAlpha), Eric Lyons (University of California, Berkley) for assistance and consultation on CoGe, and Eric Spangler and Kristian Skjervold of the University of Memphis High-Performance Computing (HPC) cluster.

Author contributions

ERMP, PE, JMB, DSJ, and JRM conceived the study. ERMP, PE, BRC, JB, MDP, and ZM performed data curation, formal analyses, and developed methodologies and software. ERMP, PE, JB, and JRM led the investigation, data validation, visualization, and drafting of the manuscript. ACW contributed to data visualization. MDP, SD, and ZM provided additional analytical and software support. AH, MB, ZN, JMB, DSJ, and JRM contributed to manuscript review and editing. MB and ZN provided resources, and ZN, JMB, DSJ, and JRM acquired funding and supervised the project. All authors read and approved of the final manuscript.

Data availability

Raw data and genome assembly data for *Acicarpha*, *Scaevola*, and *Platycodon* are uploaded to NCBI BioProject PRJNA1253301. Accession numbers of these newly developed and publicly available data can be found in Supplementary Table S6. Hardmasked, annotated genome assemblies for *Acicarpha, Scaevola,* and *Platycodon,* as well as the re-annotated assemblies for *Helianthus, Chrysanthemum, Lactuca, Arctium,* and *Nymphoides* were uploaded and made publicly accessible on the Comparative Genomics Platform[152] (id67898, id67904, id67903, id67900, id67899, id67901, id67867, id67902, respectively).

Funding

This work was supported by the National Science Foundation under grants IOS-2214472, DBI-2319905, and DEB-745197, and by the National Institutes of Health through an R35 MIRA award (R35GM119614).

References

695
1. Vamosi, J. C., Magallón, S., Mayrose, I., Otto, S. P. & Sauquet, H. Macroevolutionary
Patterns of Flowering Plant Speciation and Extinction. *Annu Rev Plant Biol* **69**, 685–706
(2018).
2. Barker, M. S. *et al.* Most Compositae (Asteraceae) are descendants of a paleohexaploid
and all share a paleotetraploid ancestor with the Calyceraceae. *Am J Bot* **103**, 1203–
1211 (2016).
3. Huang, C.-H. *et al.* Multiple Polyploidization Events across Asteraceae with Two Nested
Events in the Early History Revealed by Nuclear Phylogenomics. *Mol Biol Evol* **33**, 2820–
2835 (2016).
4. Zhang, C. *et al.* Phylotranscriptomic insights into Asteraceae diversity, polyploidy, and
morphological innovation. *J Integr Plant Biol* **63**, 1273–1293 (2021).
5. Palazzesi, L. *et al.* Asteraceae as a model system for evolutionary studies: from fossils to
genomes. *Botanical Journal of the Linnean Society* **200**, 143–164 (2022).
6. Mandel, J. R. *et al.* A fully resolved backbone phylogeny reveals numerous dispersals
and explosive diversifications throughout the history of Asteraceae. *Proc Natl Acad Sci U
S A* **116**, 14083–14088 (2019).
7. Panero, J. L. & Crozier, B. S. Macroevolutionary dynamics in the early diversification of
Asteraceae. *Mol Phylogenet Evol* **99**, 116–132 (2016).
8. Susanna, A. *et al.* The classification of the Compositae: A tribute to Vicki Ann Funk
(1947–2019). *Taxon* **69**, 807–814 (2020).
9. Diazgranados, M. *et al.* World Checklist of Useful Plant Species. *KNB Data Repository*
https://knb.ecoinformatics.org/view/doi:10.5063/F1CV4G34 (2020).
10. Lundberg, J. G. Asteraceae and relationships within Asterales. in (2009).
11. Brignone, N. F., Mazet, N., Pozner, R. & Denham, S. S. Calyceraceae: Unexpected
diversification pattern in the Southern Andes. *Perspect Plant Ecol Evol Syst* **60**, 125744
(2023).
12. Denham, S. S., Zavala-Gallo, L., Johnson, L. A. & Pozner, R. E. Insights into the
phylogeny and evolutionary history of Calyceraceae. *Taxon* **65**, 1328–1344 (2016).
13. Zhang, T. & Elomaa, P. Development and evolution of the Asteraceae capitulum. *New
Phytologist* **242**, 33–48 (2024).
14. Gurung, V., Muñoz-Gómez, S. & Jones, D. S. Putting heads together: Developmental
genetics of the Asteraceae capitulum. *Curr Opin Plant Biol* **81**, 102589 (2024).
15. Pozner, R., Zanotti, C. & Johnson, L. A. Evolutionary origin of the Asteraceae capitulum:
Insights from Calyceraceae. *Am J Bot* **99**, 1–13 (2012).
16. Teeri, T. H. *et al.* Reproductive meristem fates in Gerbera. *J Exp Bot* **57**, 3445–3455
(2006).
17. Flagel, L. E. & Wendel, J. F. Gene duplication and evolutionary novelty in plants. *New
Phytologist* **183**, 557–564 (2009).
18. Doyle, J. J. Polyploidy in Legumes. in *Polyploidy and Genome Evolution* 147–180
(Springer Berlin Heidelberg, Berlin, Heidelberg, 2012). doi:10.1007/978-3-642-31442-
1_9.

19.   Eric Schranz, M., Mohammadin, S. & Edger, P. P. Ancient whole genome duplications,
novelty and diversification: the WGD Radiation Lag-Time Model. *Curr Opin Plant Biol* **15**,
147–153 (2012).
20.   Soltis, D. E. *et al.* Polyploidy and angiosperm diversification. *Am J Bot* **96**, 336–348
(2009).
21.   Shen, F. *et al.* Comparative genomics reveals a unique nitrogen-carbon balance system
in Asteraceae. *Nat Commun* **14**, (2023).
22.   Kong, X. *et al.* Two-step model of paleohexaploidy, ancestral genome reshuffling and
plasticity of heat shock response in Asteraceae. *Hortic Res* **10**, (2023).
23.   Barker, M. S. *et al.* Multiple Paleopolyploidizations during the Evolution of the
Compositae Reveal Parallel Patterns of Duplicate Gene Retention after Millions of Years.
Mol Biol Evol **25**, 2445–2455 (2008).
24.   Reyes-Chin-Wo, S. *et al.* Genome assembly with in vitro proximity ligation data and
whole-genome triplication in lettuce. *Nat Commun* **8**, 14953 (2017).
25.   Oliver, K. R., McComb, J. A. & Greene, W. K. Transposable Elements: Powerful
Contributors to Angiosperm Evolution and Diversity. *Genome Biol Evol* **5**, 1886–1901
(2013).
26.   Fedoroff, N. Transposons and genome evolution in plants. *Proceedings of the National*
Academy of Sciences **97**, 7002–7007 (2000).
27.   Lippman, Z. *et al.* Role of transposable elements in heterochromatin and epigenetic
control. *Nature* **430**, 471–476 (2004).
28.   Staton, S. E. & Burke, J. M. Evolutionary transitions in the Asteraceae coincide with
marked shifts in transposable element abundance. *BMC Genomics* **16**, 623 (2015).
29.   Badouin, H. *et al.* The sunflower genome provides insights into oil metabolism, flowering
and Asterid evolution. *Nature* **546**, 148–152 (2017).
30.   Schwacke, R., Bolger, M. E. & Usadel, B. PubPlant: a continuously updated online
resource for sequenced and published plant genomes. Preprint at
https://doi.org/10.1101/2025.03.12.642823 (2025).
31.   Lundberg, J. & Bremer, K. A Phylogenetic Study of the Order Asterales Using One
Morphological and Three Molecular Data Sets. *Int J Plant Sci* **164**, 553–578 (2003).
32.   Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis of gene
synteny and collinearity. *Nucleic Acids Res* **40**, e49–e49 (2012).
33.   Jiao, Y. *et al.* A genome triplication associated with early diversification of the core
eudicots. *Genome Biol* **13**, R3 (2012).
34.   Zwaenepoel, A. & Van de Peer, Y. Inference of Ancient Whole-Genome Duplications and
the Evolution of Gene Duplication and Loss Rates. *Mol Biol Evol* **36**, 1384–1404 (2019).
35.   Jin, Y. *et al.* Functional divergence of GLP genes between G. barbadense and G.
hirsutum in response to Verticillium dahliae infection. *Genomics* **114**, 110470 (2022).
36.   Wang, Z. *et al.* Comparative genomics analysis of WAK/WAKL family in Rosaceae
identify candidate WAKs involved in the resistance to Botrytis cinerea. *BMC Genomics*
24, 337 (2023).
37.   Dar, A. A., Choudhury, A. R., Kancharla, P. K. & Arumugam, N. The FAD2 Gene in
Plants: Occurrence, Regulation, and Role. *Front Plant Sci* **8**, (2017).

38. Li, Y. & Wei, K. Comparative functional genomics analysis of cytochrome P450 gene
superfamily in wheat and maize. *BMC Plant Biol* **20**, 93 (2020).
39. Quezada, E.-H. *et al.* Cysteine-Rich Receptor-Like Kinase Gene Family Identification in
the Phaseolus Genome and Comparative Analysis of Their Expression Profiles Specific
to Mycorrhizal and Rhizobial Symbiosis. *Genes (Basel)* **10**, 59 (2019).
40. Feng, T. *et al.* FAD2 Gene Radiation and Positive Selection Contributed to Polyacetylene
Metabolism Evolution in Campanulids. *Plant Physiol* **181**, 714–728 (2019).
41. Pandian, B. A., Sathishraj, R., Djanaguiraman, M., Prasad, P. V. V. & Jugulam, M. Role
of Cytochrome P450 Enzymes in Plant Stress Response. *Antioxidants* **9**, 454 (2020).
42. Chakraborty, P., Biswas, A., Dey, S., Bhattacharjee, T. & Chakrabarty, S. Cytochrome
P450 Gene Families: Role in Plant Secondary Metabolites Production and Plant Defense.
J Xenobiot **13**, 402–423 (2023).
43. Chen, Y., Klinkhamer, P. G. L., Memelink, J. & Vrieling, K. Diversity and evolution of
cytochrome P450s of Jacobaea vulgaris and Jacobaea aquatica. *BMC Plant Biol* **20**, 342
(2020).
44. Alshareef, N. O. *et al.* Root remodeling mechanisms and salt tolerance trade-offs: The
roles of HKT1, TMAC2, and TIP2;2 in Arabidopsis. *PLoS Genet* **21**, e1011713 (2025).
45. Mukhtar, M. S., Deslandes, L., Auriac, M., Marco, Y. & Somssich, I. E. The Arabidopsis
transcription factor WRKY27 influences wilt disease symptom development caused by
Ralstonia solanacearum. *The Plant Journal* **56**, 935–947 (2008).
46. Han, X. *et al.* Arabidopsis Transcription Factor TCP5 Controls Plant
Thermomorphogenesis by Positively Regulating PIF4 Activity. *iScience* **15**, 611–622
(2019).
47. van Es, S. W. *et al.* Novel functions of the Arabidopsis transcription factor TCP5 in petal
development and ethylene biosynthesis. *The Plant Journal* **94**, 867–879 (2018).
48. Yu, H. *et al.* TCP5 controls leaf margin development by regulating KNOX and BEL-like
transcription factors in Arabidopsis. *J Exp Bot* **72**, 1809–1821 (2021).
49. Chapman, M. A., Leebens-Mack, J. H. & Burke, J. M. Positive Selection and Expression
Divergence Following Gene Duplication in the Sunflower CYCLOIDEA Gene Family. *Mol
Biol Evol* **25**, 1260–1273 (2008).
50. Lee, J. & Lee, I. Regulation and function of SOC1, a flowering pathway integrator. *J Exp
Bot* **61**, 2247–2254 (2010).
51. Blackman, B. K. *et al.* Contributions of Flowering Time Genes to Sunflower Domestication
and Improvement. *Genetics* **187**, 271–287 (2011).
52. Li, D. *et al.* Arabidopsis Class II TCP Transcription Factors Integrate with the FT–FD
Module to Control Flowering. *Plant Physiol* **181**, 97–111 (2019).
53. Sawa, M. & Kay, S. A. GIGANTEA directly activates Flowering Locus T in Arabidopsis
thaliana. *Proceedings of the National Academy of Sciences* **108**, 11698–11703 (2011).
54. Sawa, M., Nusinow, D. A., Kay, S. A. & Imaizumi, T. FKF1 and GIGANTEA Complex
Formation Is Required for Day-Length Measurement in Arabidopsis. *Science (1979)* **318**,
261–265 (2007).
55. Xie, H. *et al.* DNA Methylation of the Autonomous Pathway Is Associated with Flowering
Time Variations in Arabidopsis thaliana. *Int J Mol Sci* **25**, 7478 (2024).

56.   Simpson, G. G. The autonomous pathway: epigenetic and post-transcriptional gene
regulation in the control of Arabidopsis flowering time. *Curr Opin Plant Biol* **7**, 570–574
(2004).

57.   Richter, R., Bastakis, E. & Schwechheimer, C. Cross-Repressive Interactions between
SOC1 and the GATAs GNC and GNL/CGA1 in the Control of Greening, Cold Tolerance,
and Flowering Time in Arabidopsis . *Plant Physiol* **162**, 1992–2004 (2013).

58.   Bellucci, E. *et al.* Selection and adaptive introgression guided the complex evolutionary
history of the European common bean. *Nat Commun* **14**, 1908 (2023).

59.   Huang, N., Tien, H. & Yu, T. Arabidopsis leaf-expressed AGAMOUS-LIKE 24 mRNA
systemically specifies floral meristem differentiation. *New Phytologist* **241**, 504–515
(2024).

60.   Van de Peer, Y., Mizrachi, E. & Marchal, K. The evolutionary significance of polyploidy.
Nat Rev Genet **18**, 411–424 (2017).

61.   Soltis, P. S., Marchant, D. B., Van de Peer, Y. & Soltis, D. E. Polyploidy and genome
evolution in plants. *Curr Opin Genet Dev* **35**, 119–125 (2015).

62.   Dunn, T. & Sethuraman, A. Accurate Inference of the Polyploid Continuum Using
Forward-Time Simulations. *Mol Biol Evol* **41**, (2024).

63.   Freeling, M. Bias in Plant Gene Content Following Different Sorts of Duplication: Tandem,
Whole-Genome, Segmental, or by Transposition. *Annu Rev Plant Biol* **60**, 433–453
(2009).

64.   Freeling, M. & Thomas, B. C. Gene-balanced duplications, like tetraploidy, provide
predictable drive to increase morphological complexity. *Genome Res* **16**, 805–814
(2006).

65.   Qiao, X. *et al.* Different Modes of Gene Duplication Show Divergent Evolutionary Patterns
and Contribute Differently to the Expansion of Gene Families Involved in Important Fruit
Traits in Pear (Pyrus bretschneideri). *Front Plant Sci* **9**, (2018).

66.   *Systematics, Evolution, and Biogeography of Compositae*. (International Association for
Plant Taxonomy, Vienna, 2009).

67.   Ramsey, J. & Ramsey, T. S. Ecological studies of polyploidy in the 100 years following its
discovery. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**,
20130352 (2014).

68.   Strong, D. R. & Ayres, D. R. Ecological and Evolutionary Misadventures of *Spartina*.
Annu Rev Ecol Evol Syst **44**, 389–410 (2013).

69.   Martin, S. L. & Husband, B. C. Adaptation of diploid and tetraploid Chamerion
angustifolium to elevation but not local environment. *Evolution (N Y)* **67**, 1780–1791
(2013).

70.   Boalt, E., Arvanitis, L., Lehtilä, K. & Ehrlén, J. The association among herbivory
tolerance, ploidy level, and herbivory pressure in cardamine pratensis. *Evol Ecol* **24**,
1101–1113 (2010).

71.   Thompson, J. N. & Merg, K. F. Evolution of polyploidy and the diversification of plant-
pollinator interactions. *Ecology* **89**, 2197–2206 (2008).

72.   Oswald, B. P. & Nuismer, S. L. Neopolyploidy and pathogen resistance. *Proceedings of
the Royal Society B: Biological Sciences* **274**, 2393–2397 (2007).

73. Fan, W. *et al.* The genomes of chicory, endive, great burdock and yacon provide insights
into Asteraceae palaeo-polyploidization history and plant inulin production. *Mol Ecol*
Resour **22**, 3124–3140 (2022).

74. Wen, X. *et al.* The Chrysanthemum lavandulifolium genome and the molecular
mechanism underlying diverse capitulum types. *Hortic Res* **9**, (2022).

75. Todesco, M. *et al.* Massive haplotypes underlie ecotypic differentiation in sunflowers.
Nature **584**, 602–607 (2020).

76. Huang, K. *et al.* The genomics of linkage drag in inbred lines of sunflower. *Proceedings of*
the National Academy of Sciences **120**, (2023).

77. Yang, J.-S., Qian, Z.-H., Shi, T., Li, Z.-Z. & Chen, J.-M. Chromosome-level genome
assembly of the aquatic plant Nymphoides indica reveals transposable element bursts
and NBS-LRR gene family expansion shedding light on its invasiveness. *DNA Research*
29, (2022).

78. Minio, A., Cochetel, N., Vondras, A. M., Massonnet, M. & Cantu, D. Assembly of
complete diploid-phased chromosomes from draft genome sequences. *G3*
Genes|Genomes|Genetics **12**, (2022).

79. Doyle, J. J. & Doyle, J. L. A Rapid DNA Isolation Procedure for Small Quantities of Fresh
Leaf Tissue. *Phytochemical Bulletin* **19**, 11–15 (1987).

80. Vaillancourt, B. & Buell, C. R. High molecular weight DNA isolation method from diverse
plant species for use with Oxford Nanopore sequencing. Preprint at
https://doi.org/10.1101/783159 (2019).

81. Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data.
Preprint at (2010).

82. Schultz, D., Ebbert, M. & De Coster, W. PAUVRE. Preprint at (2019).

83. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

84. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170–175 (2021).

85. Zimin, A. V. *et al.* Hybrid assembly of the large and highly repetitive genome of Aegilops
tauschii, a progenitor of bread wheat, with the MaSuRCA mega-reads algorithm. *Genome*
Res **27**, 787–792 (2017).

86. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357–359 (2012).

87. Walker, B. J. *et al.* Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement. *PLoS One* **9**, e112963 (2014).

88. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
(2013).

89. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

90. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

91. Zhou, C., McCarthy, S. A. & Durbin, R. YaHS: yet another Hi-C scaffolding tool.
Bioinformatics **39**, (2023).

92. Durand, N. C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps with
Unlimited Zoom. *Cell Syst* **3**, 99–101 (2016).

93. Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M. BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol Biol Evol* **38**,
4647–4654 (2021).

94. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol **20**, 257 (2019).

95. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a
streamlined, comprehensive pipeline. *Genome Biol* **20**, 275 (2019).

96. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable
element families. *Proceedings of the National Academy of Sciences* **117**, 9451–9457
(2020).

97. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0. Preprint at
http://www.repeatmasker.org (2013).

98. Gabriel, L. *et al.* BRAKER3: Fully automated genome annotation using RNA-seq and
protein evidence with GeneMark-ETP, AUGUSTUS and TSEBRA. Preprint at
https://doi.org/10.1101/2023.06.10.544449 (2023).

99. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**, 1236–1240 (2014).

100. Manchanda, N. *et al.* GenomeQC: a quality assessment tool for genome assemblies and
gene structure annotations. *BMC Genomics* **21**, 193 (2020).

101. Jin, J. J. *et al.* GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of
organelle genomes. *Genome Biol* **21**, 1–31 (2020).

102. Zhou, W. *et al.* Plastid Genome Assembly Using Long-read data. *Mol Ecol Resour* **23**,
1442–1457 (2023).

103. Tillich, M. *et al.* GeSeq- versatile and accurate annotation of organelle genomes. *Nucleic
Acids Res* **45**, (2017).

104. Darling, A. C. E., Mau, B., Blattner, F. R. & Perna, N. T. Mauve: Multiple Alignment of
Conserved Genomic Sequence With Rearrangements. *Genome Res* **14**, 1394–1403
(2004).

105. Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. *Bioinformatics* **27**, 764–770 (2011).

106. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and
Smudgeplot for reference-free profiling of polyploid genomes. *Nat Commun* **11**, (2020).

107. Kokot, M., Dlugosz, M. & Deorowicz, S. KMC 3: counting and manipulating k-mer
statistics. *Bioinformatics* **33**, 2759–2761 (2017).

108. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**, (2019).

109. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530–1534 (2020).

110. Tang, H. *et al.* JCVI: A versatile toolkit for comparative genomics analysis. *iMeta* **3**,
(2024).

111. Sun, P. *et al.* WGDI: A user-friendly toolkit for evolutionary analyses of whole-genome
duplications and ancestral karyotypes. *Mol Plant* **15**, 1841–1851 (2022).

112. Jia, K. *et al.* Sub Phaser: a robust allopolyploid subgenome phasing method based on
subgenome-specific k-mers. *New Phytologist* **235**, 801–809 (2022).

113. Zhang, C., Rabiee, M., Sayyari, E. & Mirarab, S. ASTRAL-III: Polynomial time species
tree reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**, 15–30
(2018).

114. Zhang, R.-G., Shang, H.-Y., Jia, K.-H. & Ma, Y.-P. Subgenome phasing for complex
allopolyploidy: case-based benchmarking and recommendations. *Brief Bioinform* **25**,
(2023).

115. Paten, B. *et al.* Cactus: Algorithms for genome multiple sequence alignment. *Genome*
Res **21**, 1512–1528 (2011).

116. Armstrong, J. *et al.* Progressive Cactus is a multiple-genome aligner for the thousand-
genome era. *Nature* **587**, 246–251 (2020).

117. Sullivan, P. F. *et al.* Leveraging base-pair mammalian constraint to understand genetic
variation and human disease. *Science (1979)* **380**, (2023).

118. Hubisz, M. J., Pollard, K. S. & Siepel, A. PHAST and RPHAST: phylogenetic analysis
with space/time models. *Brief Bioinform* **12**, 41–51 (2011).

119. Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. *Genome Res* **15**, 1034–1050 (2005).

120. Keough, K. C. *et al.* Three-dimensional genome rewiring in loci with human accelerated
regions. *Science (1979)* **380**, (2023).

121. Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral
substitution rates on mammalian phylogenies. *Genome Res* **20**, 110–121 (2010).

122. Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proceedings*
of the National Academy of Sciences **100**, 9440–9445 (2003).

123. Cahill, J. A. *et al.* Positive selection in noncoding genomic regions of vocal learning birds
is associated with genes implicated in vocal learning and speech functions in humans.
Genome Res **31**, 2035–2049 (2021).

124. Yocca, A. *et al.* A chromosome-scale assembly for 'd'Anjou' pear. *G3: Genes, Genomes,*
Genetics **14**, (2024).

125. Edgar, R. C. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* **32**, 1792–1797 (2004).

126. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: Robust conversion of protein sequence
alignments into the corresponding codon alignments. *Nucleic Acids Res* **34**, (2006).

127. Yang, Z. *PAML: A Program Package for Phylogenetic Analysis by Maximum Likelihood*.
CABIOS APPLICATIONS NOTE vol. 13 http://mw511.biol.berkeley.edu/ziheng/paml.html.
(1997).

128. R Development Core Team (2008). R: A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0,
URL http://www.R-project.org. Preprint at https://stat.ethz.ch/pipermail/r-help/2008-
May/161481.html.

129. R Core Team. R: A language and environment for statistical computing. Preprint at
https://www.r-project.org/ (2017).

130. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, New York,
2016).

131. Sensalari, C., Maere, S. & Lohaus, R. ksrates: positioning whole-genome duplications
relative to speciation events in Ks distributions. *Bioinformatics* **38**, 530–532 (2022).

132. McKibben, M. T. W., Finch, G. & Barker, M. S. Species-tree topology impacts the
inference of ancient whole-genome duplications across the angiosperm phylogeny. *Am J*
Bot **111**, (2024).

133. Chen, H. *et al.* Revisiting ancient polyploidy in leptosporangiate ferns. *New Phytologist*
237, 1405–1417 (2023).

134. Pupko, T., Pe, I., Shamir, R. & Graur, D. A Fast Algorithm for Joint Reconstruction of
Ancestral Amino Acid Sequences. *Mol Biol Evol* **17**, 890–896 (2000).

135. Revell, L. J. phytools: An R package for phylogenetic comparative biology (and other
things). *Methods Ecol Evol* **3**, 217–223 (2012).

136. De Bie, T., Cristianini, N., Demuth, J. P. & Hahn, M. W. CAFE: a computational tool for
the study of gene family evolution. *Bioinformatics* **22**, 1269–1271 (2006).

137. Maurin, K. J. L. An empirical guide for producing a dated phylogeny with treePL in a
maximum likelihood framework. (2020).

138. Smith, S. A. & O'Meara, B. C. treePL: divergence time estimation using penalized
likelihood for large phylogenies. *Bioinformatics* **28**, 2689–2690 (2012).

139. Barreda, V. D. *et al.* Early evolution of the angiosperm clade Asteraceae in the
Cretaceous of Antarctica. *Proceedings of the National Academy of Sciences* **112**, 10989–
10994 (2015).

140. Barreda, V. D. *et al.* An extinct Eocene taxon of the daisy family (Asteraceae):
evolutionary, ecological and biogeographical implications. *Ann Bot* **109**, 127–134 (2012).

141. Hochuli, P. A. *Palynologische Untersuchungen Im Oligozän Und Untermiozän Der*
Zentralen Und Westlichen Paratethys. (Geologischen Institut der Eidg. Technischen
Hochschule und der Universtät Zürich, Zurich, 1978).

142. Beentje, H. J. & Caligari, P. D. S. *Compositae: Proceedings of the International*
Compositae Conference, Kew, 1994. (Royal Botanic Gardens, 1996).

143. Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L. & Hernández-Hernández, T. A
metacalibrated time-tree documents the early rise of flowering plant phylogenetic
diversity. *New Phytologist* **207**, 437–453 (2015).

144. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *J Mol Biol* **215**, 403–410 (1990).

145. Ceron-Noriega, A., Schoonenberg, V. A. C., Butter, F. & Levin, M. AlexandrusPS: A User-
Friendly Pipeline for the Automated Detection of Orthologous Gene Clusters and
Subsequent Positive Selection Analysis. *Genome Biol Evol* **15**, (2023).

146. Ranwez, V., Harispe, S., Delsuc, F. & Douzery, E. J. P. MACSE: Multiple Alignment of
Coding SEquences Accounting for Frameshifts and Stop Codons. *PLoS One* **6**, e22594
(2011).

147. Kosakovsky Pond, S. L. *et al.* HyPhy 2.5—A Customizable Platform for Evolutionary
Hypothesis Testing Using Phylogenies. *Mol Biol Evol* **37**, 295–299 (2020).

148. Paradis, E. & Schliep, K. Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2019).

149.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. ggtree: an r package for visualization
1040   and annotation of phylogenetic trees with their covariates and other associated data.
1041   *Methods Ecol Evol* **8**, 28–36 (2017).
1042   150.   Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nat Genet* **25**, 25–
1043   29 (2000).
1044   151.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing
1045   Biological Themes Among Gene Clusters. *OMICS* **16**, 284–287 (2012).
1046   152.   Lyons, E., Pedersen, B., Kane, J. & Freeling, M. The Value of Nonmodel Genomes and
1047   an Example Using SynMap Within CoGe to Dissect the Hexaploidy that Predates the
1048   Rosids. *Trop Plant Biol* **1**, 181–190 (2008).
1049   153.   GBIF.org User. Occurrence Download. Preprint at https://doi.org/10.15468/DL.RJ2PNP
1050   (2025).
1051   154.   GBIF.org User. Occurrence Download. Preprint at https://doi.org/10.15468/DL.BRQ5TF
1052   (2025).
1053   155.   GBIF.org User. Occurrence Download. Preprint at https://doi.org/10.15468/DL.66G87F
1054   (2025).
1055

1056 **Table 1.** Representative genes from analyses aiming to understand the genomic mechanisms underpinning diversification in

1057 Asterales. Gene symbols were obtained from BLASTP search results to a TAIR10 protein database. Functions were summarized

1058 from representative publications. Asterisk indicates that this gene was the best match for six different orthogroups. Complete results

1059 are listed in Supplementary Dataset S3.

Category	Gene	Description	Involved in
Top ten expanded orthogroups in Asteraceae	GLP9	germin-like protein 9	disease resistance (Jin et al., 2022)
	WAKL2	wall associated kinase-like 2	cellular communication, disease resistance (Wang et al., 2023)
	FAD2	fatty acid desaturase 2	fatty acid synthesis (Dar et al., 2017; Feng et al., 2019)
	CYP71B23	cytochrome P450, family 71, subfamily B, polypeptide 37	sisquiterpene synthesis and modification (Y. Li & Wei, 2020)
	AT1G43760	DNAse I-like superfamily protein	NA
	RPP13	NB-ARC domain-containing disease resistance protein	disease resistance (Yuan et al., 2025)
	WIP4	Wound-responsive family protein	root development (Diaz-Ramirez et al., 2022)
	HYR1	UDP-Glycosyltransferase superfamily protein	growth regulation (Tan et al., 2022)
	CYP72A15	cytochrome P450, family 72, subfamily A, polypeptide 15	merabolic resistance (Zhou et al., 2025)
	HTB9	Histone superfamily protein	chromatin remodeling (Liu et al., 2007)
Top ten expanded orthogroups in *Acicarpha*	NFU3	NFU domain protein 3	plant growth, development, and flowering (Nath et al., 2017)
	CRK8*	cysteine-rich RECEPTOR-like kinase	abiotic and biotic stress response (Quezada et al., 2019)
	LRR4	NB-ARC domain-containing disease resistance protein	disease resistance (Cao et al., 2024)
	AT2G01050	zinc ion binding; nucleic acid binding	NA
	SAUR2	SAUR-like auxin-responsive protein family	stress response (C. Li et al., 2024)
Lost in Goodeniaceae, Calyceraceae, and Asteraceae	GLB1(PII)	Nitrogen regulatory protein P-II homolog	nitrogen and metabolic regulation (Shen et al., 2023)
Positive Selection in Asteraceae	CYP72A15	cytochrome P450, family 72, subfamily A, polypeptide 15	secondary metabolism (Prall et al., 2016)
	TMAC2	AFP2 (ABI five-binding protein 2) family protein	stress response (Alshareef et al., 2025)
	TCP2,5,24	TEOSINTE BRANCHED 1, cycloidea and PCF transcription factor 2,5,24)	plant growth and development (D. Li et al., 2019; van Es et al., 2018; Yu et al., 2021)
	WRKY6,21,22,27, 31,39,42,57,74,75	WRKY DNA-binding proteins	stress response (Bakshi & Oelmüller, 2014)
	GI	gigantea protein	flowering time regulation (Sawa et al., 2007; Sawa & Kay, 2011)
	FVE	Transducin family protein / WD-40 repeat family protein	regulation of flowering time and cold response (Baek et al., 2008)
	CGA1	cytokinin-responsive gata factor 1	regulation of chloroplast development, stress response, and flowering time (Chiang et al., 2012; Richter et al., 2013)
	FD	Basic-leucine zipper (bZIP) transcription factor family protein	regulation of floral transition, plant development, and environmental response (Yang et al., 2025)
	AGL24	AGAMOUS-like 24	regulation of flowering time (Huang et al., 2024)

Table 1 References:

Alshareef, N. O., Melino, V. J., Saber, N., De Rosa, A., Rey, E., Wang, J. Y., AlBabili, S., Byrt, C., Tester, M. A., & Julkowska, M. M. (2025). Root remodeling mechanisms and salt tolerance trade-offs: The roles of HKT1, TMAC2, and TIP2;2 in Arabidopsis. PLOS Genetics, 21(6), e1011713. https://doi.org/10.1371/journal.pgen.1011713

Baek, I.-S., Park, H.-Y., You, M. K., Lee, J. H., & Kim, J.-K. (2008). Functional Conservation and Divergence of FVE Genes that Control Flowering Time and Cold Response in Rice and Arabidopsis. Molecules and Cells, 26(4), 368–372. https://doi.org/10.1016/S1016-8478(23)14010-6

Bakshi, M., & Oelmüller, R. (2014). WRKY transcription factors. Plant Signaling & Behavior, 9(2), e27700. https://doi.org/10.4161/psb.27700

Cao, Y., Mo, W., Li, Y., Xiong, Y., Wang, H., Zhang, Y., Lin, M., Zhang, L., & Li, X. (2024). Functional characterization of NBS-LRR genes reveals an NBS-LRR gene that mediates resistance against Fusarium wilt. BMC Biology, 22(1), 45. https://doi.org/10.1186/s12915-024-01836-x

Chiang, Y.-H., Zubo, Y. O., Tapken, W., Kim, H. J., Lavanway, A. M., Howard, L., Pilon, M., Kieber, J. J., & Schaller, G. E. (2012). Functional Characterization of the GATA Transcription Factors GNC and CGA1 Reveals Their Key Role in Chloroplast Development, Growth, and Division in Arabidopsis . Plant Physiology, 160(1), 332–348. https://doi.org/10.1104/pp.112.198705

Dar, A. A., Choudhury, A. R., Kancharla, P. K., & Arumugam, N. (2017). The FAD2 Gene in Plants: Occurrence, Regulation, and Role. Frontiers in Plant Science, 8. https://doi.org/10.3389/fpls.2017.01789

Diaz-Ramirez, D., Diaz-Garcia, U. S., Magdaleno-Garcia, G., Huep, G., Appelhagen, I., Sagasser, M., & Marsch-Martinez, N. (2022). Expression and Functional Analyses of the WIP Gene Family in Arabidopsis. Plants, 11(15), 2010. https://doi.org/10.3390/plants11152010

Feng, T., Yang, Y., Busta, L., Cahoon, E. B., Wang, H., & Lü, S. (2019). FAD2 Gene Radiation and Positive Selection Contributed to Polyacetylene Metabolism Evolution in Campanulids. Plant Physiology, 181(2), 714–728. https://doi.org/10.1104/pp.19.00800

Huang, N., Tien, H., & Yu, T. (2024). Arabidopsis leaf-expressed AGAMOUS-LIKE 24 mRNA systemically specifies floral meristem differentiation. New Phytologist, 241(1), 504–515. https://doi.org/10.1111/nph.19293

Jin, Y., Fan, L., Zhang, Y., Hu, W., Han, X., Yan, Q., Yang, J., Li, F., & Yang, Z. (2022). Functional divergence of GLP genes between G. barbadense and G. hirsutum in response to Verticillium dahliae infection. Genomics, 114(5), 110470. https://doi.org/10.1016/j.ygeno.2022.110470

Li, C., Jiang, C., Pan, X., Qi, Y., Zhao, W., Dong, H., & Liu, Q. (2024). OsSAUR2, a small auxin-up RNA gene, is crucial for arsenic tolerance and accumulation in rice. Environmental and Experimental Botany, 226, 105894. https://doi.org/10.1016/j.envexpbot.2024.105894

Li, D., Zhang, H., Mou, M., Chen, Y., Xiang, S., Chen, L., & Yu, D. (2019). Arabidopsis Class II TCP Transcription Factors Integrate with the FT–FD Module to Control Flowering. Plant Physiology, 181(1), 97–111. https://doi.org/10.1104/pp.19.00252

Li, Y., & Wei, K. (2020). Comparative functional genomics analysis of cytochrome P450 gene superfamily in wheat and maize. BMC Plant Biology, 20(1), 93. https://doi.org/10.1186/s12870-020-2288-7

Liu, Y., Koornneef, M., & Soppe, W. J. J. (2007). The Absence of Histone H2B Monoubiquitination in the Arabidopsis hub1 (rdo4) Mutant Reveals a Role for Chromatin Remodeling in Seed Dormancy. The Plant Cell, 19(2), 433–444. https://doi.org/10.1105/tpc.106.049221

Nath, K., O'Donnell, J. P., & Lu, Y. (2017). Chloroplastic iron-sulfur scaffold protein NFU3 is essential to overall plant fitness. Plant Signaling & Behavior, 12(2), e1282023. https://doi.org/10.1080/15592324.2017.1282023

Prall, W., Hendy, O., & Thornton, L. E. (2016). Utility of a Phylogenetic Perspective in Structural Analysis of CYP72A Enzymes from Flowering Plants. PLOS ONE, 11(9), e0163024. https://doi.org/10.1371/journal.pone.0163024

Quezada, E.-H., García, G.-X., Arthikala, M.-K., Melappa, G., Lara, M., & Nanjareddy, K. (2019). Cysteine-Rich Receptor-Like Kinase Gene Family Identification in the Phaseolus Genome and Comparative Analysis of Their Expression Profiles Specific to Mycorrhizal and Rhizobial Symbiosis. Genes, 10(1), 59. https://doi.org/10.3390/genes10010059

Richter, R., Bastakis, E., & Schwechheimer, C. (2013). Cross-Repressive Interactions between SOC1 and the GATAs GNC and GNL/CGA1 in the Control of Greening, Cold Tolerance, and Flowering Time in Arabidopsis . Plant Physiology, 162(4), 1992–2004. https://doi.org/10.1104/pp.113.219238

Sawa, M., & Kay, S. A. (2011). GIGANTEA directly activates Flowering Locus T in Arabidopsis thaliana. Proceedings of the National Academy of Sciences, 108(28), 11698–11703. https://doi.org/10.1073/pnas.1106771108

Sawa, M., Nusinow, D. A., Kay, S. A., & Imaizumi, T. (2007). FKF1 and GIGANTEA Complex Formation Is Required for Day-Length Measurement in Arabidopsis. Science, 318(5848), 261–265. https://doi.org/10.1126/science.1146994

Shen, F., Qin, Y., Wang, R., Huang, X., Wang, Y., Gao, T., He, J., Zhou, Y., Jiao, Y., Wei, J., Li, L., & Yang, X. (2023). Comparative genomics reveals a unique nitrogen-carbon balance system in Asteraceae. Nature Communications, 14(1). https://doi.org/10.1038/s41467-023-40002-9

Tan, Y., Yang, J., Jiang, Y., Wang, J., Liu, Y., Zhao, Y., Jin, B., Wang, X., Chen, T., Kang, L., Guo, J., Cui, G., Tang, J., & Huang, L. (2022). Functional Characterization of UDP-
Glycosyltransferases Involved in Anti-viral Lignan Glycosides Biosynthesis in Isatis indigotica. Frontiers in Plant Science, 13. https://doi.org/10.3389/fpls.2022.921815

van Es, S. W., Silveira, S. R., Rocha, D. I., Bimbo, A., Martinelli, A. P., Dornelas, M. C., Angenent, G. C., & Immink, R. G. H. (2018). Novel functions of the Arabidopsis transcription
factor TCP5 in petal development and ethylene biosynthesis. The Plant Journal, 94(5), 867–879. https://doi.org/10.1111/tpj.13904

Wang, Z., Ma, Y., Chen, M., Da, L., Su, Z., Zhang, Z., & Liu, X. (2023). Comparative genomics analysis of WAK/WAKL family in Rosaceae identify candidate WAKs involved in the
resistance to Botrytis cinerea. BMC Genomics, 24(1), 337. https://doi.org/10.1186/s12864-023-09371-9

Yang, H., Zhong, M., Liu, Y., Chen, L., Liu, B., Kong, F., Li, H., & Yue, L. (2025). Multifaceted roles of FD gene family in flowering, plant architecture, and adaptation. Frontiers in Plant
Science, 16. https://doi.org/10.3389/fpls.2025.1602756

Yu, H., Zhang, L., Wang, W., Tian, P., Wang, W., Wang, K., Gao, Z., Liu, S., Zhang, Y., Irish, V. F., & Huang, T. (2021). TCP5 controls leaf margin development by regulating KNOX
and BEL-like transcription factors in Arabidopsis. Journal of Experimental Botany, 72(5), 1809–1821. https://doi.org/10.1093/jxb/eraa569

Yuan, B., Li, C., Wang, Q., Yao, Q., Guo, X., Zhang, Y., & Wang, Z. (2025). Identification and functional characterization of the RPP13 gene family in potato (Solanum tuberosum L.)
for disease resistance. Frontiers in Plant Science, 15. https://doi.org/10.3389/fpls.2024.1515060

Zhou, F.-Y., Lu, H., Huang, X., Han, Y., Zhang, Y., Han, H., Tsykalchuk, L., Nyporko, A., & Yu, Q. (2025). Cytochrome CYP72A15 may play a role in metabolic resistance to
mesotrione in wild radish. Pesticide Biochemistry and Physiology, 210, 106380. https://doi.org/10.1016/j.pestbp.2025.106380

1117

Fig. 1. Characteristics of the eight focal Asterales species. **a)** from left to right, distribution of

Goodeniaceae (yellow), Calyceraceae (green), and Asteraceae (red), based on occurrence

points of preserved specimens accessed from the Global Biodiversity Information Facility[153–155].

b) Photos of some tested taxa. Colored box around image matches species colors in **c** and **d**.

From top to bottom: *Platycodon grandiflorus* (Campanulaceae); *Scaevola aemula*

(Goodeniaceae); *Acicarpha procumbens* (Calyceraceae); *Lactuca sativa* (Asteraceae);

Helianthus annuus (Asteraceae). Image of *Acicarpha procumbens* by D.S. Jones, remaining

images by A.C. Willoughby. **c)** Nuclear phylogenetic cladogram of the eight species generated

via the multiple-species alignment from OrthoFinder and IQTree. Values at nodes are bootstrap

support values. Tree to the bottom left shows branch lengths by nucleotide substitutions per

site. The four Asteraceae taxa are shaded gray. Values under each species name at the tip

label correspond to the estimated hapoloid chromosome number (x) and genome assembly size

in Gb, respectively. **d)** Synteny plots generated with JCVI ordered by phylogenetic placement of

the eight target taxa. Each bar is representative of a single chromosome, with circled numbers

above the bar indicating chromosome number. Gray lines between species' chromosomes

indicate a single syntenic gene. **e)** Barplots indicating the percentage of repeat elements (left)

and gene duplication type (right) across the genomes with percentage of the genome on the x-

axis and species on the y-axis. Colors correspond to the legends to the right. Taxon order

follows phylogenetic placement from **c.**

Fig. 2. Evidence of WGD events across the eight tested Asterales taxa. **a)** Pairwise gene depth ratios from JCVI between the eight Asterales species plus Vitis. The top right portion of the matrix indicates the gene depth ratio of the y-axis, while the bottom left portion is for the x-axis (e.g., *Scaevola:Acicarpha* is 1:3). Colors follow the legend in the bottom left of the panel. **b)** Syntenic regions between *Scaevola* (Goodeniacae), *Acicarpha* (Calyceraceae), and *Arctium* (Asteraceae) and as indicated by JCVI. Colored synteny ribbons are identified gene triplications between *Scaevola* to *Acicarpha* and follow those genes to *Arctium*. Bars represent chromosomes with circled numbers above the bars representing the chromosome number. **c)** K_S plot of paralogous gene pairs and orthologous gene pairs against Arctium for each species tested. Peaks closer to zero indicate a younger duplication event. Colors of lines correspond to the legend at the top right. **d)** Chromosome 11 subgenome phylogeny as determined by WGDI. Potential duplication events are indicated by clade, from Clades 1-3. Species names at the tips are colored according to subgenome number: blue: 1, green: 2, yellow: 3, pink: 4, orange: 5, purple: 6. Synteny plots at each tip were generated with JCVI and colored according to the synteny plots of Figure 1c. Local posterior probability (LPP) support values are at each node. **e)** Genome fractionation results from the subgenome analysis of chromosome 11 from WGDI for

Acicarpha and four Asteraceae subgenomes. Gene retention is on the y-axis and base pair

number on the x-axis. Colors in the graph correspond to the subgenome numbers which are

also represented as the tip label color in d. **f)** Summary of WGD results across this study.

Colored boxes indicate the findings, as indicated by the legend, that provide evidence for WGDs

on that branch. White boxes indicate no evidence was found for that analysis. Campanulaceae

has no boxes as WGDs were not directly tested for that family.

1160

Fig. 3. Possible WGD scenarios in Asterales. A colored pentagon indicates a WGD placement.

The scenarios are as follows: **a)** hypothesis for WGT at the crown of Asteraceae and

Calyceraceae, followed by unique WGD in *Chrysanthemum* and *Helianthus* lineages, supported

by previous literature, **b)** hypothesis for WGD at the crown of Asteraceae and Calyceraceae,

followed by WGD at the base of Asteraceae, then unique WGD in *Acicarpha, Chrysanthemum,*

and *Helianthus* lineages, supported by the critical model in WHALE, **c)** hypothesis for WGD at

the crown of Asteraceae, Calyceraceae, and Goodeniaceae, as well as unique WGD in

Nymphoides, Scaevola, Acicarpha, Lactuca, Chrysanthemum, and *Helianthus* lineages,

supported by the relaxed model in WHALE.

1170

Fig. 4. Genes underpinning Asteraceae diversification. GO enrichment of biological processes associated with expanded orthogroups in **a)** Asteraceae and **b)** *Acicarpha;* **c)** Expanded FAD2 gene family tree where tip colors represent the taxonomic family of each gene as indicated by the legend at the right and tip shapes represent the duplication type of each gene as indicated by the legend in d; **d)** Composition of each duplication type by taxonomic family for the FAD2 gene family; **e)** Expanded CYP1B23 gene family tree with same tip shape and color designations as c. **f)** Composition of each duplication type by taxonomic family for the CYP1B23 gene family.

1179

Fig. 5. Flowering time regulatory network involving: *CO, GI, FT, FD, FVE, FLD, FRI, FLC, CGA1, GNC, AGL24, SOC1, LFY,* and *AP1.* Colored gene boxes indicate orthogroups which experienced episodic positive selection in a lineage of Asteraceae homologs. Episodes of positive selection for only Asteraceae lineages are visualized as shaded blocks in phylogenetic trees and correspond to their respective gene color. Within the phylogenetic gene trees, additional episodes of positive selection are indicated as small black circles along nodes and tips, duplication types of genes are indicated by tip shapes, and plant families are indicated by color.