TET1 non-catalytic activity shapes the chromatin landscape that directs de novo

methylation establishment in the male germline

Prasasya, R. D.[1], Kim, J. J.[1], Liu, Z.[1], Kohli, R. M.[2,3], Bartolomei, M. S.[1,3]

[1]Department of Cell and Developmental Biology, Perelman School of Medicine, University of
Pennsylvania, Philadelphia, PA 19104, USA

[2]Department of Medicine, Perelman School of Medicine, University of Pennsylvania,
Philadelphia, PA 19104, USA

[3]Penn Epigenetics Institute, University of Pennsylvania, Philadelphia, PA, 19104, USA

*Correspondence: bartolom@pennmedicine.upenn.edu (MSB)

Abstract

In the germline, DNA methylation is globally erased in primordial germ cells (PGCs), enabling

establishment of sex-specific methylomes in prospermatogonia or oocytes. The catalytic activity

of TET1 is required for complete demethylation in PGCs, yet sperm from $Tet1^{-/-}$ mice display

methylation defects not explained by incomplete erasure. Instead, these defects arise from

abnormal de $novo$ methylation during development, coinciding with erosion of H3K4me3, a

chromatin modification that blocks DNMT3A/3L. Using a catalytically inactive $Tet1^{HxD}$ mouse

line, we demonstrate a non-catalytic role of TET1 in promoting H3K4me3 deposition and

protecting a subset of sperm hypomethylated regions from aberrant methylation in

prospermatogonia.

Introduction

DNA methylation is one of the most sexually dimorphic epigenetic marks in mammalian gametes (Stewart et al. 2016). Following specification from cells of the proximal epiblast, primordial germ cells (PGCs) undergo epigenetic reprogramming that involves both chromatin remodeling and genome-wide DNA methylation erasure (Saitou et al. 2005; Smith and Meissner 2013; Saitou et al. 2011). In the mouse developmental timeline, the PGC genome attains a hypomethylated state by embryonic day (E)13.5 (Seisenberger et al. 2012). From this common basal epigenetic state, the sperm genome becomes highly methylated (~90% overall methylation in the mouse) while the oocyte genome remains intermediate, with distinct hyper- and hypomethylated domains (Kobayashi et al. 2012; Hanna et al. 2018). In the male germline, acquisition of DNA methylation occurs in prospermatogonia between E15.5 and postnatal day 0. In contrast, *de novo* methylation occurs postnatally in oocytes when individual follicles are recruited for growth during each estrus cycle. Despite the distinct timing and the divergent resulting methylomes between the two sexes, *de novo* DNA methylation is facilitated by a common DNA methyltransferase (DNMT) complex: DNMT3A and its noncatalytic co-factor DNMT3L (DNMT3A/3L) (Kaneda et al. 2004; Kato et al. 2007; Bourc'his and Bestor 2004).

The mechanisms that target DNMT3A/3L to distinct loci in prospermatogonia versus the growing oocyte to generate sex-specific methylomes are still unclear. Biochemical studies suggest crosstalk between histone post-translational modifications (PTMs) and genome accessibility for DNMT3A/3L. For example, the PWWP domain of DNMT3A binds to di- or tri-methylated H3K36 (H3K36me2/me3) (Dhayalan et al. 2010). In the germline, H3K36me2 enriched domains are highly correlated with DNA methylation acquisition in prospermatogonia while transcription-coupled H3K36me3 deposition directs DNA methylation in the oocyte (Shirane et al. 2020; Chotalia et al. 2009; Stewart et al. 2015). In contrast, methylation of H3K4 (H3K4me1/2/3) repels the ADD domain of DNMT3A and DNMT3L, inhibiting DNA methylation activity (Zhang et al. 2010; Ooi et al. 2007). Consequently, H3K4me3 remodeling is required for

proper DNA methylation acquisition in the oocyte (Hanna et al. 2018). In the male germline,

DNA methylation acquisition occurs broadly, except at regions enriched for H3K4me2 at the

prospermatogonia stage (Singh et al. 2013) . Our previous data also confirmed the high

congruency of H3K4me3 enrichment in prospermatogonia with sperm hypomethylated regions

(HMRs) (Prasasya et al. 2024). To date, however, how the genome is patterned for

H3K36me2/3 or H3K4me1/2/3 prior to gametic *de novo* methylation establishment remains

unresolved.

In our previous investigation of the methylcytosine dioxygenase, Tet-eleven translocation

1 (TET1), in germline reprogramming, we described extensive methylation defects in *Tet1*[-/-]

sperm beyond the previously described limited numbers of TET1-dependent loci (i.e. imprinting

control regions -ICRs- and meiosis-associated gene promoters that require TET1 during PGC

epigenetic reprogramming) (Prasasya et al. 2024; Yamaguchi et al. 2012, 2013; Hill et al. 2018;

SanMiguel et al. 2018). TET enzymes iteratively oxidize 5-methylcytosine (5mC) to 5-

hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC), and 5-carboxylcytosine (5caC) (Ito et al.

2011). 5hmC is poorly recognized by maintenance DNMT1, while 5fC and 5caC can be excised

by thymine DNA glycosylase (TDG) resulting in an abasic site, activating the base excision

repair machinery to recover unmodified cytosine (Kohli and Zhang 2013). Using mouse models

harboring 5hmC-stalling (diminished activity in generating 5fC/5caC) TET1[V] and catalytically

inactive TET1[HxD], we established the necessity of TET1 iterative oxidative capability in germline

epigenetic reprogramming. Intriguingly, only a small portion of methylation defects observed in

Tet1[-/-] sperm can be attributed to the loss of TET1 catalytic activity. In fact, expression of full-

length, catalytically compromised TET1 (TET1[HxD]) was sufficient to rescue about 2/3 of

methylation defects observed in *Tet1*[-/-] sperm (Prasasya et al. 2024). In addition to the catalytic

core located within the C-terminus, TET proteins possess an extensive N-terminus with the

ability to interact with and recruit several histone modifiers such as OGT, Sin3a, and PRC2 *in*

vivo (Vella et al. 2013; Zhu et al. 2018; Wu et al. 2011; Chrysanthou et al. 2022). Here, we

demonstrate that the majority of hypermethylation in *Tet1*[-/-] sperm arises from abnormal *de novo*

methylation and coincides with diminished H3K4me3 at intergenic hypomethylated regions.

Importantly, catalytically inactive TET1 was sufficient to restore H3K4me3, establishing a non-

catalytic role for TET1 in shaping the chromatin landscape of the male germline.

Results and Discussion

We conducted global DNA methylation analysis of *Tet1*[-/-] and WT (*Tet1*[+/+]) PGCs at the

completion of reprogramming, E13.5, using the mouse Illumina Infinium Methylation BeadChip.

Consistent with previous reports, the *Tet1*[-/-] PGC genome was not globally hypermethylated

compared to WT (Supplemental Fig. 1A) (Yamaguchi et al. 2013; Hill et al. 2018). Aberrantly

methylated regions in *Tet1*[-/-] PGCs were almost exclusively hypermethylated and enriched for

ICRs (Figure 1A, Supplemental Fig. 1B, Supplemental Table S1A; 1049 hypermethylated

differentially methylated regions -DMRs-, 1 hypomethylated DMR; each DMR corresponds to a

single probe on the array). Gene ontology (GO) analysis revealed that hypermethylated DMRs

were enriched for genes involved in meiosis prophase I (Figure 1B), consistent with previous

findings by us and others showing that promoter activation requires proper demethylation (Hill et

al. 2018; Prasasya et al. 2024). We compared hypermethylated DMRs in *Tet1*[-/-] PGCs with

differentially expressed genes (DEGs) from our previously described RNA-seq (Figure 1C)

(Prasasya et al. 2024). Only DMRs within promoters of meiosis-associated genes corresponded

to downregulated expression, suggesting that the transcriptional consequences of TET1-

dependent hypermethylation outside of these meiosis-associated genes may manifest later in

germ cell development.

To assess whether incomplete DNA methylation erasure during PGC development

explains the hypermethylation defects previously reported in *Tet1*[-/-] sperm (Prasasya et al.

2024), we compared hypermethylated DMRs at the two stages (Figure 1D-E, Supplemental

Table S1B). 348 DMRs were hypermethylated in both *Tet1*[-/-] PGCs and sperm. As shown in the

distribution plot (Figure 1E, panel iii, Supplemental Table S1E), average methylation levels of

these regions were unchanged between sperm and PGCs for both genotypes, suggesting that

only a small fraction (6.18%) of $Tet1^{-/-}$ sperm hypermethylation results from incomplete erasure

in PGCs. In contrast, the majority of hypermethylated DMRs in $Tet1^{-/-}$ sperm showed no change

in $Tet1^{-/-}$PGCs compared to WT (Figure 1E, panel i-ii), indicating replication-coupled passive

dilution of DNA methylation was sufficient to reprogram these loci. Instead, these regions

abnormally acquired DNA methylation following successful erasure in $Tet1^{-/-}$ germ cells (Figure

1E, panel i-ii, Supplemental Table S1C-D).

When comparing WT PGCs and sperm, we noted a subset of regions where DMRs

typically gain methylation during normal male germline development; in $Tet1^{-/-}$, this acquisition

was exaggerated, resembling a hypermorphic phenotype (Figure 1E, panel i). In contrast, a

second subset of regions typically remained unmethylated in WT sperm but ectopically gained

methylation in $Tet1^{-/-}$ (Figure 1E, panel ii). Out of the 5283 $Tet1^{-/-}$ sperm DMRs that originated

from abnormal methylation acquisition, the majority belonged to the category of hypermorphic

gain (3667 hypermorphic gain, 1602 ectopic gain, 14 sperm DMRs did not pass filter in the PGC

data set). Finally, we identified 701 regions that required TET1 for reprogramming in PGCs but

ultimately became fully methylated in both $Tet1^{-/-}$ and WT sperm (e.g. paternally methylated

ICRs such as $H19$ or $IG-DMR$) or sperm HMRs that continued to demethylate as the germ cells

resumed mitosis in spermatogonial stem cells (SSCs) (Figure 1E, panel iv, Supplemental Table

S1F). Design group enrichment analysis showed that abnormal methylation gain with TET1 loss

occurred at enhancers and long noncoding RNA (lncRNA) promoters, whereas regions with

either incomplete erasure or exclusively hypermethylated in PGCs occurred mainly at ICRs

(Supplemental Fig. 1C–F). Overall, these findings suggest that TET1 not only supports

epigenetic reprogramming in PGCs but also regulates proper DNA methylation acquisition

during male germline development.

We hypothesized that abnormal gain of methylation observed in the *Tet1*[-/-] male

germline occurs concurrently with *de novo* methylation establishment and reflects changes in

the chromatin landscape in prospermatogonia. To test this, we conducted CUT&RUN for

H3K4me3, a histone PTM that inhibits DNMT3A/3L activity, in E17.5 prospermatogonia. We

chose E17.5 as a mid-point of global wave of *de novo* methylation that occurs between E15.5

and birth in the male germline (Kobayashi et al. 2013; Seisenberger et al. 2012; Singh et al.

2013). For both WT and *Tet1*[-/-], over 50% of identified peaks were located within distal

intergenic regions (Supplemental Fig. 2A). Most identified peaks showed overlap between *Tet1*[-

/-] and WT prospermatogonia, although *Tet1*[-/-] prospermatogonia had fewer consensus peaks

compared to WT (67205 vs 95643 peaks; consensus peaks were defined as peaks called in at

least 2 out of 3 biological replicates; Supplemental Fig. 2A, B, Supplemental Table S2A, B).

Averaged signals across genic regions or intergenic peaks revealed distinct H3K4me3 changes

in *Tet1*[-/-] prospermatogonia (Figure 2A). While *Tet1*[-/-] prospermatogonia showed higher overall

H3K4me3 near transcriptional start sites (TSSs), H3K4me3 was markedly lower at intergenic

peaks (Figure 2A). Differential enrichment (FDR ≤ 0.05, minimum 1.5-fold change vs WT)

identified 22904 regions that lost H3K4me3, mostly intergenic regions, in *Tet1*[-/-]

prospermatogonia (Supplemental Fig. 2D, Supplemental Table S3A). In contrast, 10971 regions

that gained H3K4me3 mapped predominantly within genic regions (Supplemental Fig. 2D,

Supplemental Table S3B). Notably, in prospermatogonia, intergenic H3K4me3 signal intensities

were comparable to those at TSSs (Figure 2A; Supplemental Fig. 2E-F), underscoring their

potential regulatory importance (Guenther et al. 2007) .

As we previously showed, H3K4me3 enrichment in prospermatogonia strongly

overlapped with sperm HMRs (Figure 2B, Supplemental Fig. 2C) (Prasasya et al. 2024). With

TET1 loss, we observed significantly less overlap (in absolute number of sperm HMRs) between

E17.5 H3K4me3 peaks and sperm HMRs (Figure 2B). This initial analysis suggested that fewer

sperm HMRs were protected by H3K4me3 in *Tet1*[-/-] prospermatogonia. To follow up, we

analyzed the relationship between *Tet1*[-/-] sperm DMRs and changes to H3K4me3 in *Tet1*[-/-]

prospermatogonia. With the caveat that the Infinium array provided methylation data for only a

representative subset of the genome, we found strong overlap between regions that lost

H3K4me3 in *Tet1*[-/-] prospermatogonia and hypermethylated DMRs in *Tet1*[-/-] sperm (Figure 2C).

As described above, the majority of *Tet1*[-/-] sperm hypermethylated DMRs originate from

abnormal methylation acquisition following successful DNA demethylation in E13.5 PGCs

(Figure 1E). Because methylated H3K4 protects the genome from targeting by *de novo* DNMTs,

we posit that diminished H3K4me3 in *Tet1*[-/-] prospermatogonia increases DNMT3A/3L access,

leading to aberrant methylation at these regions. Assessing H3K4me3 signals and *Tet1*[-/-] sperm

hypermethylated DMR subsets showed that those originating from abnormal methylation

(hypermorphic and ectopic) corresponded to regions with H3K4me3 erosion in

prospermatogonia, while those originating from incomplete methylation erasure did not show

changes in H3K4me3 signals (Figure 2D). DMRs specific to *Tet1*[-/-] PGCs also showed no

H3K4me3 changes at the prospermatogonia stage, suggesting a decoupling between the role of

TET1 in DNA demethylation during reprogramming and its role in patterning the chromatin for

de novo methylation acquisition (Figure 2D). Incidences of abnormal methylation gain that

coincide with H3K4me3 reduction in *Tet1*[-/-] prospermatogonia can be found in both genic and

intergenic regions (Figure 2E). ICRs such as those from *Nespas/Gnas* locus, where

hypermethylation originated from incomplete methylation erasure (Supplemental Fig. 1B), had

largely unchanged H3K4me3 in *Tet1*[-/-] prospermatogonia (Figure 2E).

H3K4me3 is highly enriched at TSSs of active genes in eukaryotic cells (Barski et al.

2007; Schneider et al. 2004). While there is strong evidence that H3K4me3 can serve as a

recognition signal for the transcription initiation complex, transcription can still occur without

H3K4me3 because the mark may not be instructive for gene expression genome wide (Lauberth

et al. 2013; Clouaire et al. 2012). We assessed the relationship between H3K4me3 changes

and gene expression in *Tet1*[-/-] prospermatogonia. While total RNA sequencing revealed many

differentially expressed genes in *Tet1*[-/-] prospermatogonia (DEGs: 2035 upregulated, 4772

downregulated; Supplemental Fig. 3A; Supplemental Table S4A), we did not observe any

directional correlations to changes in H3K4me3 (Figure 3A, Supplemental Table S4B). Perhaps

consistent with the known promiscuity or leakiness of gene expression in the male germline,

where up to 80% of all protein-coding genes can be detected at any given developmental stage,

DEGs are associated with biological processes at much later stages of spermatogenesis (e.g.

downregulation of motility-associated genes and upregulation of chromosome segregation

during meiosis-associated genes; Supplemental Fig. 3B, C) (Xia et al. 2020). Overall, our data

showed that H3K4me3 in prospermatogonia does not direct global gene expression.

To explore the regulatory potential of these altered regions across spermatogenesis, we

performed chromatin state discovery using ChromHMM. We built segmentation models from

publicly available histone PTMs, Pol2, and CTCF ChIP-seq datasets (Supplemental Fig. 3D,

Supplemental Table 5) spanning five stages of spermatogenesis: SSCs, differentiated

spermatogonia (SPG), pachytene spermatocyte (PS), round spermatid (RS), and sperm. We

assigned chromatin states based on Roadmap Epigenomics Consortium conventions

(Supplemental Fig. 3D) (Ernst and Kellis 2012) . Regions that gained H3K4me3 in *Tet1*[-/-]

prospermatogonia matched canonical TSS, while regions with depleted H3K4me3

corresponded to enhancer-like states (Figure 3B). Regions with reduced H3K4me3 in *Tet1*[-/-]

prospermatogonia were enriched for long terminal repeats (LTRs) and LINE elements

(Supplemental Fig. 3E). Motif enrichment analysis after repeat masking identified recognition

sites for methylation sensitive ETS and HOX families of transcription factors (Figure 3C)

(Stephens and Poon 2016; Huang et al. 2012) . Consistent with potentially serving as promoters

that are relevant for male germ line development (Supplemental Fig. 3F), regions that gained

H3K4me3 in *Tet1*[-/-] prospermatogonia contain motifs for DMRT6 binding, a germ cell-specific,

methylation sensitive transcription factor that coordinates the mitosis to meiosis transition

(Figure 3D) (Zhang et al. 2014).

Our findings support the hypothesis that non-catalytic domains of TET1 contribute to

H3K4me3 deposition, which protect intergenic sperm HMRs from *de novo* methylation

acquisition. Indeed, emerging evidence shows that non-catalytic domains of TET1 within the C-

terminus interact with chromatin modifiers in development and disease (Lian et al. 2016; Joshi

et al. 2022). We previously generated a mouse line with the *Tet1$^{H1654Y,D1656A(HXD)}$* mutation, which

produces a catalytically inactive but full-length TET1 protein (Ko et al. 2011; Chrysanthou et al.

2022; Caldwell et al. 2021; Prasasya et al. 2024). To test whether non-catalytic TET1

expression is sufficient to establish the chromatin landscape for proper male germline *de novo*

methylation, we analyzed *Tet1$^{HxD/HxD}$* prospermatogonia.

We conducted CUT&RUN for H3K4me3 in *Tet1$^{HxD/HxD}$* prospermatogonia at E17.5. In

support of our hypothesis, the majority of H3K4me3 changes seen in *Tet1$^{-/-}$* prospermatogonia

were absent in *Tet1$^{HxD/HxD}$* compared to WT (Figure 4A, Supplemental Fig. 4A, Supplemental

Table S2C). Of the regions that lost H3K4me3 in *Tet1$^{-/-}$* prospermatogonia, over 89% (20494 of

22904) were restored to WT levels in *Tet1$^{HxD/HxD}$* (Figure 4A, Supplemental Fig. 4A,

Supplemental Table S3C, D). The genomic distribution of H3K4me3 consensus peaks in

Tet1$^{HxD/HxD}$ prospermatogonia was comparable to *Tet1$^{-/-}$* and WT and consistently overlapped

with sperm HMRs (Supplemental Fig. 4B-C). At both TSS-centered and intergenic peaks,

H3K4me3 signals in *Tet1$^{HxD/HxD}$* prospermatogonia showed intermediate levels between WT and

in *Tet1$^{-/-}$* (i.e. increased around TSSs and decreased at intergenic peaks; Supplemental Fig. 4D-

E).

The limited number of regions where H3K4me3 was depleted in *Tet1$^{HxD/HxD}$*

prospermatogonia (compared to WT) remained associated with eventual hypermethylated

DMRs in *Tet1$^{HxD/HxD}$* sperm, although the overlap was significantly reduced compared to *Tet1$^{-/-}$*

germ cells (Figure 4B-C). Among 5631 hypermethylated DMRs identified in $Tet1^{-/-}$ sperm, 3710

(~2/3) were not significantly hypermethylated in $Tet1^{HxD/HxD}$ sperm (defined as rescued DMRs)

(Prasasya et al. 2024). These methylation defects cannot be fully attributed to loss of TET1

catalytic activity. Of note, regions that remained hypermethylated in $Tet1^{HxD/HxD}$ sperm are those

that require TET1-iterative oxidation capability such as ICRs and promoter of germline

responsive genes (Prasasya et al. 2024).

We examined how H3K4me3 enrichment in prospermatogonia relates to the rescue of

sperm hypermethylation defects in $Tet1^{HxD/HxD}$ compared to $Tet1^{-/-}$ sperm. To determine whether

restoration of H3K4me3 in $Tet1^{HxD/HxD}$ prospermatogonia underlies this rescue, we compared

H3K4me3 signal intensity across rescued and not-rescued DMRs in $Tet1^{HxD/HxD}$ and $Tet1^{-/-}$

(Figure 4D, Supplemental Table S3E-F). Overall, both DMR subsets exhibited higher H3K4me3

levels in $Tet1^{HxD/HxD}$ than in $Tet1^{-/-}$ prospermatogonia, consistent with a global restoration of this

mark in $Tet1^{HxD/HxD}$ prospermatogonia (Figure 4A, Supplemental Fig. 4A). To quantify the extent

of H3K4me3 restoration toward WT levels, we calculated the fold-change of H3K4me3 signal in

$Tet1^{HxD/HxD}$ relative to $Tet1^{-/-}$ prospermatogonia. The degree of restoration was significantly

greater across rescued DMRs than across those that remained hypermethylated (not-rescued;

Figure 4E), indicating that a threshold of partial recovery of H3K4me3 in $Tet1^{HxD/HxD}$

prospermatogonia correlates with absence of hypermethylation in $Tet1^{HxD/HxD}$ sperm.

To test this relationship more formally, we performed permutation based enrichment

analysis to assess whether rescued DMRs preferentially overlapped regions showing restored

H3K4me3 in $Tet1^{HxD/HxD}$ prospermatogonia (Gel et al. 2015). Both rescued and not-rescued

DMRs showed significant enrichment within H3K4me3 restored peaks in $Tet1^{HxD/HxD}$

prospermatogonia (Supplemental Fig. 4F-G), but the effect size (Z-score) was markedly higher

for rescued DMRs (Figure 4F). Together, these results indicate that non-catalytic full length

TET1^{HxD} is sufficient to enhance H3K4me3 deposition in prospermatogonia compared to the

absence of TET1, yet only regions surpassing a threshold of H3K4me3 enrichment achieve

protection from DNMT3A/3L-mediated *de novo* methylation (illustrated in Figure 4G).

In the male germline, hypermethylation is required, in part, to silence retroelements that

are toxic for meiosis (Bourc'his and Bestor 2004; Barau et al. 2016). Despite this, certain

regions of the mammalian sperm genome resist DNA methylation, including CpG-dense

promoters and younger, species-specific transposon subfamilies overlapping regulatory

elements (Shirane and Lorincz 2023; Molaro et al. 2011). Additionally, low methylation at CpG-

dense regions correlates with nucleosome retention during histone-to-protamine exchange, a

critical step male gamete maturation (Erkek et al. 2013) . Retained nucleosomes in sperm play

a crucial role in patterning chromatin of the early embryo, underscoring the importance of the

sperm epigenome beyond spermatogenesis (Fanourgakis et al. 2025). Thus, the integrity of

sperm HMRs is essential for both spermatogenesis and potentially embryo viability.

We previously demonstrated that most methylation defects from compromised TET1

activity occur within sperm HMRs (Prasasya et al. 2024). Iterative oxidation is required at ICRs

and meiosis-associated promoters, but these account for only a small fraction of *Tet1*[-/-] sperm

defects. In addition to its role in active demethylation, TET1 has been proposed protect against

DNA re-methylation during epigenetic reprogramming (Hill et al. 2018). However, DNMT3A/3L

activity is limited at E14.5, making this role less likely in PGCs. Our findings instead suggest a

role for TET1 in preventing abnormal methylation acquisition in prospermatogonia. Aberrant

DNA methylation in *Tet1*[-/-] sperm coincides with erosion of H3K4me3 in prospermatogonia, the

chromatin feature that protects and patterns sperm HMRs (Figure 4G). Consistently, the non-

catalytic function of TET1, as tested in *Tet1*[HxD/HxD] germline, is sufficient enhance H3K4me3

deposition and partially rescue DNA methylation (Prasasya et al. 2024). Analysis of H3K4me3

enrichment in *Tet1*[HxD/HxD] prospermatogonia suggests that there is likely a threshold of H3K4me3

that is needed to protect sperm HMRs from the activity of DNMT3A/3L during *de novo*

methylation as not all regions with H3K4me3 restoration corresponds to rescue of DNA

hypermethylation (Figure 4G).

TET proteins interact with chromatin modifiers in many contexts. In embryonic stem cells

(ESCs), the non-catalytic function of TET1 shapes bivalency (H3K4me3, H3K27me3) at

regulators of mesoderm and trophectoderm lineages by recruiting Polycomb repressive complex

2 (PRC2) and Sin3a (Chrysanthou et al. 2022) . TET1 is also a stable binding partner of *O-*

GlcNAc transferase (OGT) and is required for OGT-Sin3a-Hcf1 binding to chromatin (Vella et al.

2013; Williams et al. 2011). Hcf1 is a key component of the SET1/COMPASS histone

methyltransferase complex, a writer for H3K4me2/3. In hematopoiesis, the TET2/3-OGT

complex promotes Hcf1 O-GlcNAcylation, leading to SET1/COMPASS assembly, H3K4me3

deposition, and gene activation (Capotosti et al. 2011; Deplus et al. 2013). A similar mechanism

may apply in prospermatogonia, as single cell RNA-seq confirms *Ogt* and *Hcfc1* expression in

fetal and neonatal germ cells, coinciding with *de novo* methylation (Zhao et al. 2021).

It is also possible that in addition to promoting H3K4me3 deposition, TET1 may also

partially restrain DNA methylation by physically protecting its targets from DNMT3A/3L in

prospermatogonia. This is not likely to be the main mechanism of sperm HMRs protection

however, as TET1 and DNMT3A antagonism was previously found to be highly context

dependent within proximal promoters (Gu et al. 2018). Furthermore, in human ESCs, TET1 and

DNMT3A have been shown to co-bind, especially within DNMT3A preference binding regions

(Chao et al. 2022). Notably, TET1-dependent sperm HMRs are intergenic and acquire

enhancer-like chromatin states throughout spermatogenesis (Figure 2A, 3B). Top enriched

motifs in these regions correspond to ETS transcription factors. Significantly, the ETS family of

transcription factors are known to interact with and recruit histone acetyltransferases such as

CBP and p300, in further support of the potential of intergenic sperm HMRs as regulatory

elements (Yang et al. 1998; Goodman and Smolik 2000). We propose that protecting intergenic

sperm HMRs from methylation helps establish the enhancer landscape that regulates stage-

specific gene expression, particularly during the mitosis-to-meiosis transition (Maezawa et al.

2020). Indeed, *Tet1*[-/-] male mice reproductive defects are attributed to premature aging due to

reduced SSCs and progressively worse meiotic defects (Huang et al. 2020). While the

mechanism orchestrating chromatin patterning before *de novo* methylation remains unclear, our

findings reveal the non-catalytic function of TET1 as a key determinant in establishing sperm

HMRs.

Methods

Mouse models and germ cell collection

All experiments were approved by the Institutional Animal Care and Use Committee of the
University of Pennsylvania (protocol number: 804211). Mice were housed in polysulfone cages
within a pathogen-free facility under a 12 h light/dark cycle, with ad libitum access to water and
standard chow (Laboratory Autoclavable Rodent Diet 5010, LabDiet, St. Louis, MO, USA). *Tet1*
knockout mice (Dawlaty et al. 2013) (017358; B6;129S4-Tet1[tm1.1Jae]/J) and *Oct4-GFP* mice
(Lengner et al. 2007) (008214; B6;129S4-Pou5f1[tm2Jae]/J) were obtained from The Jackson
Laboratory and backcrossed for at least 10 generations to a C57BL/6J background (000664;
B6; The Jackson Laboratory). Catalytically inactive *Tet1*[HxD] mouse line was generated as
described in (Prasasya et al. 2024). The *Oct4-GFP* allele was maintained as homozygous in
Tet1 heterozygote breeders (*Tet1*[+/-]; *Oct4*[GFP/GFP] or *Tet1*[HxD/+]; *Oct4*[GFP/GFP]). Primers to determine
sex, *Tet1*[tm1.1Jae], *Tet1*[HxD], and *Oct4-GFP* alleles are listed in Supplemental Table S6.

Gonadal tissues from E13.5 and E17.5 embryos (post copulatory plug detection) were dissected
and digested into a single-cell suspension using 0.25% trypsin/EDTA for 10 minutes at 37 °C.
GFP[+] germ cells were isolated by flow cytometry. For bisulfite mutagenesis and RNA isolation,
germ cells were sorted into empty tubes and immediately snap-frozen for storage at -80 °C until
processing. For CUT&RUN, sorted cells were collected into chilled PGC medium (DMEM
supplemented with 15% FBS, 100 U/mL leukemia inhibitory factor, 5 µM forskolin, 1 ng/mL stem
cell factor). Following sorting, dimethyl sulfoxide (DMSO) was added to a final concentration of
10%, and the cells were slow frozen using Mr. Frosty until library preparation.

Genome-wide DNA methylation profiling using Infinium Mouse Methylation BeadChip

Snap frozen germ cell pellet was directly lysed and bisulfite mutagenized using EZ DNA
Methylation Direct Kit (Zymo Research) per manufacturer's instructions. Bisulfite mutagenized
(BSM) DNA was subjected to low-input whole-genome amplification workflow prior to
hybridization to Illumina BeadChip array as detailed in (Lee et al. 2024) (Workflow J). Briefly,
BSM DNA underwent 2 rounds of Klenow amplifications (NEB) using random hexamer primers
(NEB). For additional details on library preparation and data analysis, see the Supplemental
Material. Amplified DNA was cleaned using solid-phase reversible immobilization (SPRI) beads

(2x; Beckman Coulter) and concentration was measured with Qubit (Invitrogen).10 μL of BSM
DNA library was submitted for hybridization onto the Infinium Mouse Methylation-12v1-0
BeadChip (Illumina) at Children's Hospital of Philadelphia Center for Applied Genomics.

CUT&RUN library preparation

CUT&RUN experiments were performed on samples of 4,000 nuclei isolated per epitope from
cryopreserved germ cells as described above using CUTANA ChIC/CUT&RUN kit (EpiCypher).
Three biological replicates per genotype were prepared for H3K4me3 (1:100, Cell Signaling
Technology antibody 2729) and IgG (1:50, CUTANA Rabbit IgG negative control antibody). Cells
were rapidly thawed and counted, followed by nuclei extraction using NE buffer (20 mM
HEPES-KOH pH 7.9, 10mM KCl, 0.5 mM fresh spermidine, 0.1% Triton X-100, 20% glycerol
plus protease inhibitors). Nuclei were bound to concanavalin A-coated magnetic beads and
CUT&RUN was conducted per CUTANA product manual with modifications. CUT&RUN libraries
were prepared with NEBNext Ultra II DNA Library Prep Kit for Illumina (NEB). For additional
details on library preparation and data analysis, see the Supplemental Material.

Total RNAseq library preparation

Total RNA was extracted from 5000 snap frozen E17.5 germ cells isolated from a single embryo
using TRIzol reagent (Invitrogen). RNA integrity number and concentration were assessed using
the High Sensitivity RNA ScreenTape for TapeStation system (Agilent Technologies). Ribosomal
RNA-depleted total RNA libraries were generated using the SMART-Seq Total RNA Pico Input
Kit with UMIs with ZapR Mammalian (Takara Bio) from 5 ng input according to the
manufacturer's protocol. Libraries were indexed with NEBNext Multiplex Oligos for Illumina
Unique Dual Index Primer Pairs (NEB) and were amplified using SeqAmp DNA Polymerase
(Takara Bio). For additional details on library preparation and data analysis, see the
Supplemental Material.

Data availability

364 Data will be submitted to NIH Gene Expression Omnibus once US government resumes
365 operation. Raw, analyzed, and corresponding meta data are currently hosted on Box and
366 access will be granted upon request for review purposes.

Acknowledgement

368 This work was supported by National Institute of Health grant numbers R01GM146388 (MSB),

369 R01GM051279 (MSB), R01GM118501 (RMK), R01HG010646 (RMK), K99HD112543 (RDP).

Authors contribution

371 Experiments were designed by RDP with assistance and supervision from MSB and RMK. JJK

372 and RDP collected samples and performed all experiments. ZL performed initial quality check

373 and alignment for CUT&RUN and RNAseq experiments. Data and statistical analysis for

374   methylation array, CUT&RUN and RNAseq was performed by RDP. JJK illustrated the summary

375   figure. RDP wrote the manuscript, which was approved by all authors.

Figures and figure legends

Figure 1. The majority of methylation defects in *Tet1^-/-^* sperm do not result from

incomplete methylation erasure during epigenetic reprogramming.

A) Top enriched design groups based on the Illumina Infinium Mouse Methylation BeadChip

manifest for hypermethylated DMRs in *Tet1^-/-^* PGCs.

B) Heatmap of probes associated with the most enriched GO term among hypermethylated

DMRs in *Tet1^-/-^* PGCs.

C) Scatterplot showing overlap between hypermethylated DMRs and DEGs in *Tet1^-/-^* PGCs.

D) Comparison of hypermethylated probes (differentially methylated regions, DMRs: FDR ≤

0.05; >10% methylation difference) in E13.5 *Tet1^-/-^* PGCs (n=8) vs *Tet1^+/+^* (WT, n=8) and those

previously reported in *Tet1^-/-^* sperm (Prasasya et al. 2024).

E) Methylation levels in E13.5 PGCs (top) or sperm (bottom) for *Tet1^-/-^* (black) and *Tet1^+/+^* (blue).

DMRs are subsetted by origin of hypermethylation (i–iv). Average methylation levels are

denoted by dotted vertical lines.

Supplemental Figure 1. Additional analysis of methylation arrays for Tet1-/- vs Tet1+/+

PGCs.

A) Violin plots of average methylation values at each CpG probe with overall mean methylation

indicated (n=8 per group). Differentially methylated probes were identified using limma (FDR ≤

0.05, >10% difference).

B) ICRs are the most enriched design-group category among hypermethylated DMRs in *Tet1*$^{-/-}$

PGCs (15 of 21 array-represented ICRs are significantly hypermethylated). Boxplots show

average methylation of probes associated with hypermethylated ICRs.

C–F) Top enriched Infinium Mouse Methylation BeadChip design groups for different categories

of *Tet1*$^{-/-}$ sperm and PGC DMRs.

Figure 2. Abnormal de novo methylation acquisition is associated with H3K4me3 erosion

in _Tet1-/-_ male germ cells.

A) Metaplot of H3K4me3 signals over genic regions (top) or consensus intergenic peaks

(bottom).

B) Distribution of H3K4me3 peaks overlapping sperm HMRs compared to non-HMRs (**** p <

0.0001, Fisher's exact test).

C) Scatterplot of _Tet1-/-_ sperm DMRs and regions with differential H3K4me3 in _Tet1-/-_

prospermatogonia. H3K4me3 peaks overlapping with DMRs in _Tet1-/-_ sperm are highlighted in

black circles. Fisher's exact test for hypermethylated DMRs overlapping H3K4me3-depleted vs

other regions.

D) (Top) Metagene plots of H3K4me3 signals in _Tet1-/-_ and _Tet1+/+_ prospermatogonia

overlapping subsets of _Tet1-/-_ sperm hypermethylated DMRs (hypermorphic gain, ectopic gain,

incomplete erasure) or _Tet1-/-_ PGC hypermethylated DMRs. (Bottom) Boxplots of mean

H3K4me3 signals over 1 kb windows centered on subsets of hypermethylated DMRs (**** p <

0.0001, Wilcoxon signed-rank test).

E) Genome browser tracks for representative sperm hypermethylated DMRs from abnormal de

novo methylation at genic region (left) or intergenic region (middle). _Nespas_ is a representative

ICR that is hypermethylated due to incomplete erasure (right).

Supplemental Figure 2. Additional analysis of differentially enriched H3K4me3 signals in

Tet1⁻ᐟ⁻ **vs** *Tet1⁺ᐟ⁺* **prospermatogonia.**

A) Genomic compartmentalization of H3K4me3 consensus peaks (present in ≥2 of 3 replicates)

in $Tet1^{+/+}$ and $Tet1^{-/-}$ prospermatogonia.

B)) Distribution of H3K4me3 consensus peaks unique to or shared between $Tet1^{+/+}$ (n=3) and

$Tet1^{-/-}$ (n=3) E17.5 prospermatogonia detected by CUT&RUN.

C) Heatmaps of H3K4me3 signals in E17.5 $Tet1^{-/-}$ and $Tet1^{+/+}$ prospermatogonia overlapping

sperm hypomethylated regions (HMRs) defined by sperm WGBS (GSE56697; Wang et al.

2014). Plots represent averages of 3 biological replicates.

D) Genomic distribution of differentially enriched H3K4me3 peaks between $Tet1^{-/-}$ and $Tet1^{+/+}$

prospermatogonia (DiffBind, FDR ≤ 0.05, ≥1.5-fold change).

E-F) Genome browser tracks for representative regions showing H3K4me3 loss (E, intergenic)

or gain (F, promoter) in $Tet1^{-/-}$ prospermatogonia.

Figure 3. H3K4me3 enrichment in prospermatogonia may regulate later stages of

spermatogenesis

A) Scatter plot of significantly altered H3K4me3 peaks in $Tet1^{-/-}$ prospermatogonia (vs WT) at

gene promoters and their associated gene expression changes as measured by total RNA-seq.

Differentially expressed genes (FDR ≤ 0.05, ≥2-fold change) are highlighted with blue

(downregulated) or red (upregulated) dots.

B) Enrichment of H3K4me3 peaks lost or gained in $Tet1^{-/-}$ prospermatogonia across chromatin

states during spermatogenesis (SSC: spermatogonial stem cells; SPG: spermatogonia; PS:

pachytene spermatocyte; RS: round spermatid), using ChromHMM. Chromatin states

assignment is based on the ENCODE consortium projects and are defined in Supplemental Fig.

3D (Ernst and Kellis 2017)

C) Motif enrichment for regions with H3K4me3 loss after masking repetitive elements.

D) Motif enrichment for regions with H3K4me3 gain.

Supplemental Figure 3. Investigation into the potential role of H3K4me3 changes in *Tet1*[-/-]

prospermatogonia

A) MA-plot for differential expression analysis of total RNA-seq in *Tet1*[-/-] prospermatogonia vs WT (n=3, FDR ≤ 0.05, ≥2-fold change).

B) Top five enriched GO biological process terms for downregulated DEGs in *Tet1*[-/-] prospermatogonia.

C) Top five enriched GO biological process terms for upregulated DEGs in *Tet1*[-/-] prospermatogonia.

D) Chromatin states assigned by ChromHMM using public datasets for CTCF, Pol II, and histone modifications across spermatogonial stem cells (SSC), spermatogonia (SPG), pachytene spermatocytes (PS), round spermatids (RS), and sperm (datasets used are in supplemental table). Chromatin state assignment is based on Encode consortium convention (Ernst and Kellis 2017).

E) Classes of repetitive elements enriched in regions with H3K4me3 loss.

F) Top ten enriched GO biological process terms for regions with H3K4me3 gain.

Figure 4. The non-catalytic, full length TET1^{HxD} restores most H3K4me3 changes in *Tet1$^{-/-}$*

prospermatogonia.

A) Alluvial plot of differentially enriched H3K4me3 peaks in *Tet1$^{-/-}$* vs *Tet1$^{+/+}$* and their status in

Tet1$^{HxD/HxD}$ vs *Tet1$^{+/+}$*. Light brown ribbons indicate peaks altered in *Tet1$^{-/-}$* but not in *Tet1$^{HxD/HxD}$*.

B) Scatterplot of *Tet1$^{HxD/HxD}$* sperm DMRs and regions with differential H3K4me3 in *Tet1$^{HxD/HxD}$*

prospermatogonia. H3K4me3 peaks overlapping with DMRs in *Tet1$^{HxD/HxD}$* sperm are highlighted

in brown circles. Fisher's exact test for hypermethylated DMRs overlapping H3K4me3-depleted

vs other regions.

C) Proportion of regions with H3K4me3 loss in Tet1 mutants that overlap hypermethylated

sperm DMRs; two-sample proportion test.

D) Ridgeline plots showing the relative frequency distribution of DMRs that were significantly

hypermethylated in *Tet1$^{-/-}$* sperm but were not hypermethylated in *Tet1$^{HxD/HxD}$* sperm (left,

rescued DMRs) or stayed hypermethylated in *Tet1$^{HxD/HxD}$* sperm (right, not-rescued DMRs) and

the associated H3K4me3 signals found in *Tet1$^{+/+}$*, *Tet1$^{-/-}$*, or *Tet1$^{HxD/HxD}$* prospermatogonia. The

mean H3K4me3 signal for each genotype across the two subsets of sperm DMRs is indicated

with vertical dashed lines of matched colors (blue: *Tet1$^{+/+}$*, black: *Tet1$^{-/-}$*, or brown: *Tet1$^{HxD/HxD}$*)

E) Fold change of H3K4me3 signals in *Tet1$^{HxD/HxD}$* compared to *Tet1$^{-/-}$* calculated across two

subsets of sperm DMRs defined in D (**** p < 0.0001, Wilcoxon signed-rank test).

F) The strength of overlap between distinct subsets of DMRs (rescued in *Tet1$^{HxD/HxD}$* sperm or

not-rescued in *Tet1$^{HxD/HxD}$* sperm) and regions where H3K4me3 was restored to WT levels in

Tet1$^{HxD/HxD}$ prospermatogonia following observed depletion in *Tet1$^{-/-}$* prospermatogonia as

determined by permutation test (Z-score from 1000 permutations of randomly selected regions

consistent with Illumina Bead Chip array coverage).

G) With the absence of TET1, depletion of H3K4me3 at sperm HMRs (largely located at

intergenic regions) is associated with aberrant methylation acquisition in prospermatogonia (top

and middle panels). In the presence of full-length TET1^{HxD}, H3K4me3 deposition in

spermatogonia is globally increased but only a subset of intergenic sperm HMRs escaped

abnormal methylation acquisition in $Tet1^{HxD/HxD}$ sperm (bottom panel). Shaded regions indicate

relative H3K4me3 signals for each genotype, closed lollipops depict methylated CpGs.

Prasasya_Supplemental Fig 4

Supplemental Figure 4. Extended CUT&RUN analysis for H3K4me3 in *Tet1$^{HxD/HxD}$*

prospermatogonia.

A) Venn diagrams of differentially enriched peaks in *Tet1$^{-/-}$* vs *Tet1$^{+/+}$* and *Tet1$^{HxD/HxD}$* vs *Tet1$^{+/+}$*

prospermatogonia (DiffBind, FDR ≤ 0.05, ≥1.5-fold change).

B) Genomic compartmentalization of consensus peaks (≥2 of 3 replicates) in *Tet1$^{+/+}$*, *Tet1$^{HxD/HxD}$*,

and *Tet1$^{-/-}$* prospermatogonia.

C) Heatmaps of H3K4me3 signals in E17.5 *Tet1$^{HxD/HxD}$* prospermatogonia overlapping sperm

HMRs defined by WGBS (GSE56697; Wang et al. 2014). Plots represent averages of 3

replicates.

D–E) Metaplots of H3K4me3 signals over genic regions (D) or intergenic consensus peaks (E)

for *Tet1$^{+/+}$*, *Tet1$^{HxD/HxD}$*, and *Tet1$^{-/-}$* prospermatogonia.

F-G) The significance of enrichment of TET1^{HxD}-rescued (F) or TET1^{HxD}-not rescued (G) DMRs

within regions where H3K4me3 was restored to WT levels in *Tet1$^{HxD/HxD}$* prospermatogonia

following observed depletion in *Tet1$^{-/-}$* prospermatogonia. In the graph, the grey-shaded region

under the curve represents the distribution of the number of overlaps between 1000 randomized

DMRs (within the array coverage) and the fixed set of H3K4me3-restored regions, values above

the red line correspond to the area with a significance of $p < 0.05$. The solid green line

represents the observed number of overlaps between rescued or not-rescued DMRs and

H3K4me3-restored regions. Z-scores represent the strength or standardized magnitude of

enrichment of overlap.

Barau J, Teissandier A, Zamudio N, Roy S, Nalesso V, Hérault Y, Guillou F, Bourc'his D. 2016.
  The DNA methyltransferase DNMT3C protects male germ cells from transposon activity.
  Science **354**: 909–912.

Barski A, Cuddapah S, Cui K, Roh T-Y, Schones DE, Wang Z, Wei G, Chepelev I, Zhao K.
  2007. High-Resolution Profiling of Histone Methylations in the Human Genome. *Cell* **129**:
  823–837.

Bourc'his D, Bestor TH. 2004. Meiotic catastrophe and retrotransposon reactivation in male
  germ cells lacking Dnmt3L. *Nature* **431**: 96–99.

Caldwell BA, Liu MY, Prasasya RD, Wang T, DeNizio JE, Leu NA, Amoh NYA, Krapp C, Lan
  Y, Shields EJ, et al. 2021. Functionally distinct roles for TET-oxidized 5-methylcytosine
  bases in somatic reprogramming to pluripotency. *Mol Cell* **81**: 859-869.e8.

Capotosti F, Guernier S, Lammers F, Waridel P, Cai Y, Jin J, Conaway JW, Conaway RC, Herr
  W. 2011. O-GlcNAc Transferase Catalyzes Site-Specific Proteolysis of HCF-1. *Cell* **144**:
  376–388.

Chao L, Yang S, Li H, Long C, Xi Q, Zuo Y. 2022. Competitive binding of TET1 and
  DNMT3A/B cooperates the DNA methylation pattern in human embryonic stem cells.
  Biochim Biophys Acta (BBA) - Gene Regul Mech **1865**: 194861.

Chotalia M, Smallwood SA, Ruf N, Dawson C, Lucifero D, Frontera M, James K, Dean W,
  Kelsey G. 2009. Transcription is required for establishment of germline methylation marks at
  imprinted genes. *Genes Dev* **23**: 105–117.
  http://genesdev.cshlp.org/cgi/doi/10.1101/gad.495809.

Chrysanthou S, Tang Q, Lee J, Taylor SJ, Zhao Y, Steidl U, Zheng D, Dawlaty MM. 2022. The
  DNA dioxygenase Tet1 regulates H3K27 modification and embryonic stem cell biology
  independent of its catalytic activity. *Nucleic Acids Res* **50**: 3169–3189.

Clouaire T, Webb S, Skene P, Illingworth R, Kerr A, Andrews R, Lee J-H, Skalnik D, Bird A.
  2012. Cfp1 integrates both CpG content and gene activity for accurate H3K4me3 deposition
  in embryonic stem cells. *Genes Dev* **26**: 1714–1728.

Dawlaty MM, Breiling A, Le T, Raddatz G, Barrasa MI, Cheng AW, Gao Q, Powell BE, Li Z,
  Xu M, et al. 2013. Combined deficiency of Tet1 and Tet2 causes epigenetic abnormalities but
  is compatible with postnatal development. *Dev Cell* **24**: 310–323.
  http://linkinghub.elsevier.com/retrieve/pii/S1534580712005898.

Deplus R, Delatte B, Schwinn MK, Defrance M, Méndez J, Murphy N, Dawson MA, Volkmar
  M, Putmans P, Calonne E, et al. 2013. TET2 and TET3 regulate GlcNAcylation and H3K4

methylation through OGT and SET1/COMPASS. *EMBO J* **32**: 645–655.
http://emboj.embopress.org/cgi/doi/10.1038/emboj.2012.357.

Dhayalan A, Rajavelu A, Rathert P, Tamas R, Jurkowska RZ, Ragozin S, Jeltsch A. 2010. The
Dnmt3a PWWP Domain Reads Histone 3 Lysine 36 Trimethylation and Guides DNA
Methylation*. *J Biol Chem* **285**: 26114–26120.

Erkek S, Hisano M, Liang C-Y, Gill M, Murr R, Dieker J, Schübeler D, Vlag J van der, Stadler
MB, Peters AHFM. 2013. Molecular determinants of nucleosome retention at CpG-rich
sequences in mouse spermatozoa. *Nat Struct Mol Biol* **20**: 868–875.

Ernst J, Kellis M. 2017. Chromatin-state discovery and genome annotation with ChromHMM.
Nat Protoc **12**: 2478–2492.

Ernst J, Kellis M. 2012. ChromHMM: automating chromatin-state discovery and
characterization. *Nat Methods* **9**: 215–216.

Fanourgakis G, Gaspa-Toneu L, Komarov PA, Papasaikas P, Ozonov EA, Smallwood SA, Peters
AHFM. 2025. DNA methylation modulates nucleosome retention in sperm and H3K4
methylation deposition in early mouse embryos. *Nat Commun* **16**: 465.

Gel B, Díez-Villanueva A, Serra E, Buschbeck M, Peinado MA, Malinverni R. 2015. regioneR:
an R/Bioconductor package for the association analysis of genomic regions based on
permutation tests. *Bioinformatics* **32**: 289–291.

Goodman RH, Smolik S. 2000. CBP/p300 in cell growth, transformation, and development.
Genes Dev **14**: 1553–1577.

Gu T, Lin X, Cullen SM, Luo M, Jeong M, Estecio M, Shen J, Hardikar S, Sun D, Su J, et al.
2018. DNMT3A and TET1 cooperate to regulate promoter epigenetic landscapes in mouse
embryonic stem cells. *Genome Biol* **19**: 88.
https://genomebiology.biomedcentral.com/articles/10.1186/s13059-018-1464-7.

Guenther MG, Levine SS, Boyer LA, Jaenisch R, Young RA. 2007. A chromatin landmark and
transcription initiation at most promoters in human cells. *Cell* **130**: 77–88.

Hanna CW, Taudt A, Huang J, Gahurova L, Kranz A, Andrews S, Dean W, Stewart AF,
Colomé-Tatché M, Kelsey G. 2018. MLL2 conveys transcription-independent H3K4
trimethylation in oocytes. *Nat Struct Mol Biol* **25**: 73–82.
http://www.nature.com/articles/s41594-017-0013-5.

Hill PWS, Leitch HG, Requena CE, Sun Z, Amouroux R, Roman-Trufero M, Borkowska M,
Terragni J, Vaisvila R, Linnett S, et al. 2018. Epigenetic reprogramming enables the transition
from primordial germ cell to gonocyte. *Nature* **555**: 392–396.
http://www.nature.com/doifinder/10.1038/nature25964.

Huang G, Liu L, Wang H, Gou M, Gong P, Tian C, Deng W, Yang J, Zhou T-T, Xu G-L, et al. 2020. Tet1 Deficiency Leads to Premature Reproductive Aging by Reducing Spermatogonia Stem Cells and Germ Cell Differentiation. *iScience* **23**: 100908.

Huang Y, Sitwala K, Bronstein J, Sanders D, Dandekar M, Collins C, Robertson G, MacDonald J, Cezard T, Bilenky M, et al. 2012. Identification and characterization of Hoxa9 binding sites in hematopoietic cells. *Blood* **119**: 388–398.

Ito S, Shen L, Dai Q, Wu SC, Collins LB, Swenberg JA, He C, Zhang Y. 2011. Tet Proteins Can Convert 5-Methylcytosine to 5-Formylcytosine and 5-Carboxylcytosine. *Science* **333**: 1300–1303.

Joshi K, Liu S, S.J. PB, Zhang J. 2022. Mechanisms that regulate the activities of TET proteins. *Cell Mol Life Sci* **79**: 363.

Kaneda M, Okano M, Hata K, Sado T, Tsujimoto N, Li E, Sasaki H. 2004. Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature* **429**: 900–903.

Kato Y, Kaneda M, Hata K, Kumaki K, Hisano M, Kohara Y, Okano M, Li E, Nozaki M, Sasaki H. 2007. Role of the Dnmt3 family in de novo methylation of imprinted and repetitive sequences during male germ cell development in the mouse. *Hum Mol Genet* **16**: 2272–2280. https://academic.oup.com/hmg/article-lookup/doi/10.1093/hmg/ddm179.

Klobučar T, Kreibich E, Krueger F, Arez M, Pólvora-Brandão D, von Meyenn F, da Rocha ST, Eckersley-Maslin M. 2020. IMPLICON: an ultra-deep sequencing method to uncover DNA methylation at imprinted regions. *Nucleic Acids Res* **48**: gkaa567-.

Ko M, Bandukwala HS, An J, Lamperti ED, Thompson EC, Hastie R, Tsangaratou A, Rajewsky K, Koralov SB, Rao A. 2011. Ten-Eleven-Translocation 2 (TET2) negatively regulates homeostasis and differentiation of hematopoietic stem cells in mice. *Proc Natl Acad Sci USA* **108**: 14566–14571. http://www.pnas.org/cgi/doi/10.1073/pnas.1112317108.

Kobayashi H, Sakurai T, Imai M, Takahashi N, Fukuda A, Yayoi O, Sato S, Nakabayashi K, Hata K, Sotomaru Y, et al. 2012. Contribution of intragenic DNA methylation in mouse gametic DNA methylomes to establish oocyte-specific heritable marks. ed. W. Reik. *PLoS Genet* **8**: e1002440. https://dx.plos.org/10.1371/journal.pgen.1002440.

Kobayashi H, Sakurai T, Miura F, Imai M, Mochiduki K, Yanagisawa E, Sakashita A, Wakai T, Suzuki Y, Ito T, et al. 2013. High-resolution DNA methylome analysis of primordial germ cells identifies gender-specific reprogramming in mice. *Genome Res* **23**: 616–627. http://genome.cshlp.org/cgi/doi/10.1101/gr.148023.112.

Kohli RM, Zhang Y. 2013. TET enzymes, TDG and the dynamics of DNA demethylation. *Nature* **502**: 472–479. http://www.nature.com/doifinder/10.1038/nature12750.

Lauberth SM, Nakayama T, Wu X, Ferris AL, Tang Z, Hughes SH, Roeder RG. 2013. H3K4me3
Interactions with TAF3 Regulate Preinitiation Complex Assembly and Selective Gene
Activation. *Cell* **152**: 1021–1036.

Lee SM, Loo CE, Prasasya RD, Bartolomei MS, Kohli RM, Zhou W. 2024. Low-input and
single-cell methods for Infinium DNA methylation BeadChips. *Nucleic Acids Res* **52**: e38–
e38.

Lengner CJ, Camargo FD, Hochedlinger K, Welstead GG, Zaidi S, Gokhale S, Scholer HR,
Tomilin A, Jaenisch R. 2007. Oct4 expression is not required for mouse somatic stem cell
self-renewal. *Cell Stem Cell* **1**: 403–415.
https://linkinghub.elsevier.com/retrieve/pii/S1934590907000859.

Lian H, Li W-B, Jin W-L. 2016. The emerging insights into catalytic or non-catalytic roles of
TET proteins in tumors and neural development. *Oncotarget* **7**: 64512–64525.

Maezawa S, Sakashita A, Yukawa M, Chen X, Takahashi K, Alavattam KG, Nakata I, Weirauch
MT, Barski A, Namekawa SH. 2020. Super-enhancer switching drives a burst in gene
expression at the mitosis-to-meiosis transition. *Nat Struct Mol Biol* **27**: 978–988.

Molaro A, Hodges E, Fang F, Song Q, McCombie WR, Hannon GJ, Smith AD. 2011. Sperm
Methylation Profiles Reveal Features of Epigenetic Inheritance and Evolution in Primates.
Cell **146**: 1029–1041.

Ooi SKT, Qiu C, Bernstein E, Li K, Jia D, Yang Z, Erdjument-Bromage H, Tempst P, Lin S-P,
Allis CD, et al. 2007. DNMT3L connects unmethylated lysine 4 of histone H3 to de novo
methylation of DNA. *Nature* **448**: 714–717. http://www.nature.com/articles/nature05987.

Prasasya RD, Caldwell BA, Liu Z, Wu S, Leu NA, Fowler JM, Cincotta SA, Laird DJ, Kohli
RM, Bartolomei MS. 2024. Iterative oxidation by TET1 is required for reprogramming of
imprinting control regions and patterning of mouse sperm hypomethylated regions. *Dev Cell*
59: 1010-1027.e8.

Saitou M, Kagiwada S, Kurimoto K. 2011. Epigenetic reprogramming in mouse pre-implantation
development and primordial germ cells. *Development* **139**: 15–31.

Saitou M, Payer B, O'Carroll D, Ohinata Y, Surani MA. 2005. Blimp1 and the Emergence of the
Germ Line during Development in the Mouse. *Cell Cycle* **4**: 1736–1740.

SanMiguel JM, Abramowitz LK, Bartolomei MS. 2018. Imprinted gene dysregulation in a Tet1
null mouse model is stochastic and variable in the germline and offspring. *Development* **145**:
dev160622. http://dev.biologists.org/lookup/doi/10.1242/dev.160622.

Schneider R, Bannister AJ, Myers FA, Thorne AW, Crane-Robinson C, Kouzarides T. 2004.
Histone H3 lysine 4 methylation patterns in higher eukaryotic genes. *Nat Cell Biol* **6**: 73–77.

Seisenberger S, Andrews S, Krueger F, Arand J, Walter J, Santos F, Popp C, Thienpont B, Dean W, Reik W. 2012. The dynamics of genome-wide DNA methylation reprogramming in mouse primordial germ cells. *Mol Cell* **48**: 849–862. http://linkinghub.elsevier.com/retrieve/pii/S109727651200932X.

Shirane K, Lorincz M. 2023. Epigenetic Mechanisms Governing Female and Male Germline Development in Mammals. *Sex Dev* **16**: 365–387.

Shirane K, Miura F, Ito T, Lorincz MC. 2020. NSD1-deposited H3K36me2 directs de novo methylation in the mouse male germline and counteracts Polycomb-associated silencing. *Nat Genet* **52**: 1088–1098.

Singh P, Li AX, Tran DA, Oates N, Kang E-R, Wu X, Szabó PE. 2013. De novo DNA methylation in the male germ line occurs by default but is excluded at sites of H3K4 methylation. *Cell Rep* **4**: 205–219. https://linkinghub.elsevier.com/retrieve/pii/S2211124713002805.

Smith ZD, Meissner A. 2013. DNA methylation: roles in mammalian development. *Nat Rev Genet* **14**: 204–220.

Stephens DC, Poon GMK. 2016. Differential sensitivity to methylated DNA by ETS-family transcription factors is intrinsically encoded in their DNA-binding domains. *Nucleic Acids Res* **44**: 8671–8681.

Stewart KR, Veselovska L, Kelsey G. 2016. Establishment and functions of DNA methylation in the germline. *Epigenomics* **8**: 1399–1413. https://www.futuremedicine.com/doi/10.2217/epi-2016-0056.

Stewart KR, Veselovska L, Kim J, Huang J, Saadeh H, Tomizawa S-I, Smallwood SA, Chen T, Kelsey G. 2015. Dynamic changes in histone modifications precede de novo DNA methylation in oocytes. *Genes Dev* **29**: 2449–2462. http://genesdev.cshlp.org/lookup/doi/10.1101/gad.271353.115.

Vella P, Scelfo A, Jammula S, Chiacchiera F, Williams K, Cuomo A, Roberto A, Christensen J, Bonaldi T, Helin K, et al. 2013. Tet proteins connect the O-linked N-acetylglucosamine transferase Ogt to chromatin in embryonic stem cells. *Mol Cell* **49**: 645–656. https://linkinghub.elsevier.com/retrieve/pii/S1097276512010556.

Williams K, Christensen J, Pedersen MT, Johansen JV, Cloos PAC, Rappsilber J, Helin K. 2011. TET1 and hydroxymethylcytosine in transcription and DNA methylation fidelity. *Nature* **473**: 343–348. http://www.nature.com/articles/nature10066.

Wu H, D'Alessio AC, Ito S, Xia K, Wang Z, Cui K, Zhao K, Sun YE, Zhang Y. 2011. Dual functions of Tet1 in transcriptional regulation in mouse embryonic stem cells. *Nature* **473**: 389–393. http://www.nature.com/articles/nature09934.

Xia B, Yan Y, Baron M, Wagner F, Barkley D, Chiodin M, Kim SY, Keefe DL, Alukal JP, Boeke JD, et al. 2020. Widespread Transcriptional Scanning in the Testis Modulates Gene Evolution Rates. *Cell* **180**: 248-262.e21.

Yamaguchi S, Hong K, Liu R, Shen L, Inoue A, Diep D, Zhang K, Zhang Y. 2012. Tet1 controls meiosis by regulating meiotic gene expression. *Nature* **492**: 443–447. http://www.nature.com/articles/nature11709.

Yamaguchi S, Shen L, Liu Y, Sendler D, Zhang Y. 2013. Role of Tet1 in erasure of genomic imprinting. *Nature* **504**: 460–464. http://www.nature.com/articles/nature12805.

Yang C, Shapiro LH, Rivera M, Kumar A, Brindle PK. 1998. A Role for CREB Binding Protein and p300 Transcriptional Coactivators in Ets-1 Transactivation Functions. *Mol Cell Biol* **18**: 2218–2229.

Zhang T, Murphy MW, Gearhart MD, Bardwell VJ, Zarkower D. 2014. The mammalian Doublesex homolog DMRT6 coordinates the transition between mitotic and meiotic developmental programs during spermatogenesis. *Development* **141**: 3662–3671.

Zhang Y, Jurkowska R, Soeroes S, Rajavelu A, Dhayalan A, Bock I, Rathert P, Brandt O, Reinhardt R, Fischle W, et al. 2010. Chromatin methylation activity of Dnmt3a and Dnmt3a/3L is guided by interaction of the ADD domain with the histone H3 tail. *Nucleic Acids Res* **38**: 4246–4253.

Zhao J, Lu P, Wan C, Huang Y, Cui M, Yang X, Hu Y, Zheng Y, Dong J, Wang M, et al. 2021. Cell-fate transition and determination analysis of mouse male germ cells throughout development. *Nat Commun* **12**: 6839.

Zhu F, Zhu Q, Ye D, Zhang Q, Yang Y, Guo X, Liu Z, Jiapaer Z, Wan X, Wang G, et al. 2018. Sin3a–Tet1 interaction activates gene transcription and is required for embryonic stem cell pluripotency. *Nucleic Acids Res* **46**: 6026–6040.