Metabolic pathway analysis of an *Acinetobacter* strain capable of assimilating diverse
hydrocarbons

Shori Inoue, Shogo Yoshimoto and Katsutoshi Hori[#]

Department of Biomolecular Engineering, Graduate School of Engineering, Nagoya University, Furo-
cho, Chikusa-ku, Nagoya, Aichi 464-8603, Japan

Correspondence: Katsutoshi Hori

Department of Biomolecular Engineering, Graduate School of Engineering, Nagoya University, Furo-
cho, Chikusa-ku, Nagoya, Aichi 464-8603, Japan.

Tel: +81-52-789-3339

Email: khori@chembio.nagoya-u.ac.jp

15
16

ABSTRACT

The growing demand for sustainable bioproduction requires the development of robust microbial chassis with broad metabolic versatility and suitability for industrial applications. *Acinetobacter* sp. Tol 5 is a highly adhesive bacterium capable of utilizing various hydrocarbons, making it a promising chassis for immobilized whole-cell catalysis. In this study, we characterized the carbon metabolism of Tol 5 by reconstructing metabolic pathway maps from its genomic data and analyzing the transcriptomes of cells grown on ethanol, hexadecane, toluene, phenol, and L-phenylalanine. Genomic analysis revealed that Tol 5 possesses a wide range of metabolic pathways for both alkanes and aromatics, including five distinct aromatic degradation routes. Transcriptome analysis identified the specific carbon sources that regulate each degradation pathway. Most pathways were induced by their corresponding carbon sources with high specificity, whereas some were upregulated by multiple related substrates. In addition, Tol 5 appears to metabolize phenylalanine via the homogentisate pathway, which differs from the phenylacetate pathway reported in *Acinetobacter baumannii*. Genes involved in oxidative and osmotic stress resistance were also coordinately upregulated on hydrocarbon sources, indicating adaptation to these cellular stresses associated with carbon sources. These results provide a comprehensive view of the carbon metabolism of Tol 5 and highlight its potential as a robust microbial chassis for converting non-sugar feedstocks into valuable bioproducts, while revealing new aspects of metabolic versatility in the genus *Acinetobacter*.

IMPORTANCE

Members of the genus *Acinetobacter* are ubiquitous environmental bacteria, which have attracted increasing attention in ecological and biotechnological fields due to their broad metabolic versatility and strong tolerance to environmental stress. We characterized the carbon metabolism of the hydrocarbon-degrading strain Tol 5 and identified five aromatic degradation pathways, including a toluene dioxygenase pathway and a homogentisate pathway, which are rarely reported in this genus. These findings expand the known metabolic repertoire of *Acinetobacter* and reveal that Tol 5 possesses aromatic degradation capabilities comparable to those of *Pseudomonas putida* strains, which are widely used as chassis for aromatic compound bioconversions. Furthermore, since aromatic compound metabolism has been reported to be associated with virulence and antibiotic resistance in pathogenic *Acinetobacter* strains, these insights may contribute to future strategies for pathogen control as well as leading the rational design of microbial chassis for sustainable bioproduction.

INTRODUCTION

In recent years, there has been a global shift from petrochemical synthesis to more sustainable bioprocesses, driven by the need to reduce fossil resource consumption and lessen environmental impact. Microorganisms, especially bacteria, are considered useful hosts for bioproduction due to their rapid growth, ease of genetic manipulation, and capacity for engineering to improve the productivity of valuable compounds. *Escherichia coli* is one of the most widely used microbial chassis in bioproduction, owing to its well-characterized molecular biology and metabolic systems (1). Concurrently, environmental microorganisms capable of assimilating non-sugar carbon substrates, such as C1 compounds, aromatic hydrocarbons, and oils, are increasingly valued for converting underutilized resources, including petrochemical by-products, agricultural waste, lignocellulosic biomass, and exhaust gases into valuable products (2-4). Therefore, it is essential to explore and develop new microbial chassis that can metabolize diverse substrates, tolerate environmental conditions, and perform robustly in industrial bioprocesses.

Members of the genus *Acinetobacter* are ubiquitous and found in diverse environments, including soil, oceans, freshwater, sediments, activated sludge, and clinical settings (5, 6). The genus possesses extensive genomic plasticity, which contributes to the pathogenic potential of some species. However, this trait, combined with a high tolerance to environmental stress, has also made it a focus of great interest in environmental microbiology, where several species have shown potential in bioproduction and bioremediation. For example, *A. baylyi* ADP1 has been studied as a host for the valorization of aromatic compounds via the β-ketoadipate pathway (5-7). *A. venetianus* RAG-1, which has multiple alkane hydroxylase genes, exhibits a pronounced ability to degrade crude oil and has been employed in bioremediation applications (8). Furthermore, *A. junii* BP25 has been utilized for the production of polyhydroxyalkanoates (PHAs) from food waste-derived organic acids, while *A. venetianus* AMO1502 has been applied in the biosynthesis of bioemulsifiers (9, 10).

Acinetobacter sp. Tol 5, originally isolated as a toluene-degrading bacterium from an exhaust gas treatment reactor, can degrade various hydrocarbons, including benzene, toluene, and xylene (BTX) (11). In addition to metabolic versatility, Tol 5 exhibits rapid autoagglutination and high adhesiveness to various solid materials, such as plastics, glass, and metals (12). This adhesive phenotype is mediated by its fibrous cell surface protein AtaA, a member of the trimeric autotransporter adhesin (TAA) family proteins (13). Previous studies have revealed the functional domains and biophysical properties of AtaA (14-16). Based on these findings, a reversible cell immobilization method was proposed to enhance the efficiency and productivity of bioproduction processes (17). To demonstrate its utility, Tol 5 expressing heterologous biosynthetic genes was immobilized on porous carriers and successfully utilized in an unconventional gas-phase bioreaction process to produce a high-value-added monoterpenoid (18). Although these advances indicate that Tol 5 is a promising microbial chassis for

innovative bioproduction, its central metabolic network remains largely uncharacterized.

In this study, we characterized the carbon metabolism of Tol 5 by reconstructing metabolic pathways
from the complete genome sequence and performing transcriptomic analyses of Tol 5 cells grown on
ethanol, hexadecane, or aromatic compounds as the sole carbon source. The results showed that Tol 5
metabolizes a variety of non-sugar carbon sources and regulates the corresponding metabolic
pathways in response to the carbon source.

RESULTS

Central carbon metabolism

The complete genome of Tol 5 consists of two circular DNA components, a chromosome of
4,681,789 bp and a plasmid of 117,717 bp, collectively encoding 4,404 predicted coding sequences
(CDSs) (19). We first evaluated the functional annotation of each CDS using multiple databases,
including NCBI, UniProtKB/Swiss-Prot, KEGG, and EggNOG, as well as information from relevant
literature (Table S1). Tol 5 encodes all the enzymes for gluconeogenesis, pentose phosphate pathway,
tricarboxylic acid (TCA) cycle, glyoxylate shunt, purine and pyrimidine biosynthesis, β-oxidation, and
fatty acid biosynthesis (Fig. 1, Table S2). In contrast, glycolysis via the Embden–Meyerhof–Parnas
(EMP) pathway is incomplete due to the absence of hexokinase, phosphofructokinase, and pyruvate
kinase, which is a trait commonly observed among *Acinetobacter* species (20). While some
Acinetobacter strains, such as *A. baylyi* ADP1 and *A. baumannii*, metabolize glucose via nonspecific
pyrroloquinoline quinone-dependent aldose dehydrogenase and the Entner–Doudoroff (ED) pathway
(21-23), Tol 5 completely lacks these genes. Furthermore, *A. baumannii* ATCC 19606 metabolizes
pentoses such as L-arabinose, D-xylose, and D-ribose via a pathway encoded by *ara* genes that
converts them into α-ketoglutarate (24), but Tol 5 also lacks the *ara* genes. These results suggest that
Tol 5 has a very limited capacity for sugar utilization as carbon sources.

Many *Acinetobacter* strains utilize organic acids and fatty acids as carbon sources (23). Tol 5 possesses
three acetate assimilation routes, including the reversible acetate kinase-phosphotransacetylase
(AckA-Pta) pathway, the irreversible acetyl-CoA synthetase (ACS) pathway, and the reversible
pathway mediated by succinyl-CoA:coenzyme A transferase AarC (Fig. 1, Table S2). For fatty acid
degradation, multiple candidate genes corresponding to each step of β-oxidation were identified,
including two putative acyl-CoA ligases, twelve acyl-CoA dehydrogenases, four enoyl-CoA
hydratases, and three thiolases. Propionyl-CoA generated during the β-oxidation of odd-chain fatty
acids was predicted to be catabolized to succinate and pyruvate via the methylcitrate cycle, which
consists of 2-methylisocitrate lyase (PrpB), 2-methylcitrate synthase (PrpC), 2-methylcitrate

dehydratase (AcnD), 2-methylaconitate isomerase (PrpF), bifunctional aconitate hydratase/isomerase
(AcnA and/or AcnB), and 2-methylisocitrate lyase (PrpB). In addition, four extracellular lipase genes
were identified in Tol 5 (Table S1), all of which are tandemly arranged (*TOL5_03740–TOL5_03770*)
and share 44–52% amino acid sequence identities with the triacylglycerol lipase Lip reported in *P.*
aeruginosa PAO1 (25). Tol 5 further encodes key enzymes that redirect fatty acid oxidation
intermediates into carbon storage compounds. A bifunctional wax ester synthase/acyl-
CoA:diacylglycerol acyltransferase (WS/DGAT, AtfA) mediates the production of wax esters and
triacylglycerols, while polyhydroxyalkanoate synthase (PhaC) is responsible for the production of
PHAs.

Alkane metabolism

Under aerobic conditions, *n*-alkanes are first oxidized at their terminal methyl group to primary
alcohols by alkane hydroxylases, and these alcohols are then further oxidized to aldehydes and finally
to fatty acids (26). The Tol 5 chromosome encodes multiple alkane hydroxylases, including a
membrane-bound non-heme iron alkane monooxygenase AlkB, and three flavin-containing
monooxygenases encoded by *almA*, *ladA_1*, *ladA_2* (Table S2). Additionally, the plasmid of Tol 5
carries a gene encoding an alkane hydroxylase (*TOL5_43480*), whose amino acid sequence shares
99% identity with that of a cytochrome P450 monooxygenase of the CYP153 family found in
Acinetobacter sp. EB104 (27). It also includes genes encoding associated redox partners, ferredoxin
(*TOL5_43490*) and ferredoxin reductase (*TOL5_43470*). AlkB and CYP153 typically oxidize short-
and medium-chain *n*-alkanes (C16 or less), whereas AlmA and LadA act on long-chain *n*-alkanes (C20
or more) (28-30). The presence of enzymes targeting various *n*-alkane chain lengths suggests that Tol
5 can efficiently degrade a wide range of alkanes. The resulting primary alcohols are further oxidized
by alcohol dehydrogenases (ADHs) and aldehyde dehydrogenases (ALDHs). In Tol 5 genome, at least
six putative ADH genes and one ALDH gene were predicted to be involved in fatty alcohol and fatty
aldehyde oxidation (Table S2). Although the functions of most ADHs in Tol 5 remain unclear, the iron-
containing ADH encoded by *yiaY_1* shares 91% amino acid sequence identity with ADH4 from *A.*
baumannii ATCC 19606, which catalyzes the oxidation of ethanol, 1-propanol, and 1-butanol (31).
The putative ALDH encoded by *aldB* (AldB) shares 87% amino acid identity with a long-chain ALDH,
Ald1, reported in *Acinetobacter* sp. M-1 (46).

Aromatic compounds metabolism

Genes encoding multi-component oxygenases for aromatic compound oxidation, as well as enzymes

for successive reaction steps leading to central metabolites, are generally organized into operons and

regulated by specific transcriptional regulators (32). Based on the predicted operons involved in

aromatic compound degradation (Table S2, Fig. 2), we reconstructed the aromatic degradation

pathways in Tol 5, identifying five major aromatic degradation routes, TOD, PCA, CAT, PAA, and

HMG pathways (Fig. 3).

The *pca* genes encode the enzymes responsible for the degradation of protocatechuate via the PCA

pathway (Fig. 3), forming a gene cluster together with the aromatic compounds transport protein PcaK

and the transcriptional repressor PcaU (Fig. 2). The *cat* gene cluster encodes the enzymes responsible

for the degradation of catechol via the CAT pathway (Fig. 3) and the transcriptional activator CatM.

Protocatechuate and catechol are common intermediates derived from various aromatic compounds

through various upstream pathways. The PCA and CAT pathways funnel these intermediates into

acetyl-CoA and succinyl-CoA, serving as central routes for aromatic compound degradation that are

widely conserved in *Acinetobacter* species (21, 33-35). In Tol 5, the *pca* genes are located adjacent to

the *qui* genes (Fig. 2), which encode enzymes that convert quinate and shikimate to protocatechuate

(Fig. 3). The structure of this gene cluster is similar to that reported in *A. baylyi* ADP1 (36), and as

described for ADP1, the *pca* and *qui* genes are likely organized as a large single operon under the

regulator PcaU in Tol 5. Tol 5 also possesses the *pob* and *van* genes, which encode upstream pathways

leading to protocatechuate, along with their respective transcriptional regulators (PobR and VanR)

(Fig. 2 and 3). The *cat* genes are located adjacent to the *mph* gene cluster, which encodes the

components of the phenol monooxygenase complex MphKLMNOP, catechol 1,2-dioxygenase CatA

(encoded by *catA_2*), and the regulatory proteins MphX (repressor) and MphR (activator) (Fig. 2).

The structure of the *mph* operon in Tol 5 resembles that of *A. calcoaceticus* PHEA-2 (37), but is

distinguished by the presence of the additional *catA* gene (*catA_2*) located immediately downstream

in the same operon. In addition, Tol 5 possesses the *ben* and *iac* genes, which encode enzymes for the

upstream pathways that lead to catechol, along with their respective transcriptional regulators (BenM

and IacR) and transporters (BenE, BenK, and a putative transporter encoded by *TOL5_19290*) (Fig. 2

and 3). Notably, the *ben* genes in Tol 5 are located over 900 kbp away from the *cat* operon, in contrast

to *A. baylyi* ADP1, in which the *ben* genes are located adjacent to the *cat* genes to facilitate coordinated

regulation of gene expression (38). Despite its distinct genomic location, the *ben* operon in Tol 5

includes the third *catA* gene (encoded by *catA_4*), indicating that coordinated regulation between the

ben and *cat* genes is maintained.

The *tod* operon encodes the enzymes of the toluene dioxygenase (TOD) pathway (Fig. 3) and is located

between the *todS-todT* two-component regulatory system and the putative toluene transporter gene

fadL2, which is located in a separate operon (Fig. 2). In the TOD pathway, toluene is oxidized into 3-

methylcatechol and ultimately converted into acetate, pyruvate, and acetyl-CoA. Although rarely

found in other *Acinetobacter* species, the *tod* genes of Tol 5 share high similarity with those of *P.*
putida strains and play a central role in the degradation of toluene and benzene, thereby expanding the
aromatic-degrading capacity of the genus (39).

Furthermore, Tol 5 possesses the *paa* genes, which encode the enzymes of the phenylacetic acid (PAA)
pathway (Fig. 3), as well as the transcriptional repressor PaaX (Fig. 2). The PAA pathway degrades
phenylacetate via epoxidation of CoA thioesters. This pathway in *A. baumannii* strains is thought to
be primarily associated with phenylalanine degradation (40, 41). In addition, Tol 5 also possesses a
gene cluster encoding a putative homogentisate pathway (HMG pathway in Fig. 3), which includes 4-
hydroxyphenylpyruvate dioxygenase (HPD), an IclR-type transcriptional repressor (IclR),
maleylacetoacetate isomerase (MaiA), fumarylacetoacetase (FahA), 3-oxoacid CoA-transferase
subunits A and B (ScoA and ScoB), and an aromatic amino acid transporter (AroP), as well as a
putative homogentisate dioxygenase encoded by *TOL5_11140* (Fig. 2). Although the protein product
of *TOL5_11140* was annotated as a glyoxalase (Table S1), it shares 22.7% amino acid identity with
homogentisate 1,2-dioxygenase HmgA (B7VA96) and 18.8% amino acid identity with
metapyrocatechase XylE (Q04385) from *P. putida*. The HMG pathway in Tol 5 is likely involved in
tyrosine metabolism. However, because canonical pathways converting phenylalanine to
phenylacetate or to tyrosine were not identified, the phenylalanine degradation route in Tol 5 could
not be determined from the genomic information.

Substrate-dependent transcriptomic responses to lactate, ethanol, and hexadecane

The expression of genes involved in carbon catabolism is tightly regulated based on available carbon
sources, particularly for the degradation pathways of hydrocarbons and aromatic compounds that are
not directly linked to central metabolism (42). Therefore, we analyzed the transcriptomic responses of
Tol 5 to different carbon sources using RNA-seq (Table S3). Transcriptomes were compared between
Tol 5 cells grown on lactate and those grown on other carbon sources; differentially expressed genes
were defined by the thresholds of $|\log_2$ fold change (FC)$| > 1$, $\log_2$ counts per million (CPM) > 3, and
false discovery rate (FDR) < 0.05 (Fig. 4A and 5A, Table S4).

In ethanol culture, 17 genes were significantly upregulated, including *yiaY_1* and *aldB*, which encode
an iron-containing ADH and a putative ALDH, respectively (Table S5). Since *yiaY_1* was the only
ADH gene induced by ethanol, it is likely the primary ADH responsible for ethanol oxidation in Tol 5
(Fig. 4B). Although AldB shares 87% amino acid identity with Ald1 from *Acinetobacter* sp. M-1,
which has been characterized as a long-chain ALDH that is inactive toward short-chain substrates such
as acetaldehyde (43), its induction during growth on ethanol suggests that AldB functions differently
from Ald1 and catalyzes acetaldehyde oxidation in Tol 5. While Tol 5 possesses both the AckA-Pta

pathway and the ACS pathway for converting acetate into acetyl-CoA (Fig. 1), only the *pta* gene in

the AckA-Pta pathway was upregulated during growth on ethanol (Fig. 4B). Additionally, *aceA*, which

encodes isocitrate lyase of the glyoxylate shunt that bypasses the decarboxylation steps of the TCA

cycle and conserves carbon flux for gluconeogenesis (Fig. 1), was upregulated. A similar regulatory

pattern was observed in *A. oleivorans* DR1, where the AckA-Pta pathway and *aceA* were upregulated

during growth on acetate compared to succinate (44), suggesting that Tol 5 employs a similar strategy

for assimilating acetate when growing on ethanol.

In hexadecane culture, 515 genes were significantly upregulated (Table S5). Among the multiple

alkane hydroxylase genes in Tol 5, *alkB* and the CYP153 gene were specifically induced, indicating

their involvement in the initial hydroxylation of hexadecane (Fig. 4C). The genes encoding an iron-

containing ADH encoded by *yiaY_3* and a putative coniferyl aldehyde dehydrogenase encoded by *calB*

were also upregulated, suggesting their roles in oxidation of hexadecanol to hexadecanal and

subsequently to hexadecanoic acid. The genes involved in every step of the β-oxidation pathway were

upregulated. Among the multiple predicted enzymes for each step, only *fadD4* (one of the two

predicted acyl-CoA ligase genes) and *fadE*, *ydiO_3*, and *ydiO_5* (three of the twelve predicted acyl-

CoA dehydrogenase genes) were specifically induced (Fig. 4C).

Sixteen genes were downregulated in both ethanol and hexadecane cultures compared to lactate culture,

implying that these genes are specifically upregulated during growth on lactate (Table S5 and S6).

Among them, the *lldP*, *lldR*, *lldD_1*, and *dld* genes (Fig. 1), which form an operon similar to the

lldPRD operon involved in lactate metabolism reported in *A. baumannii,* were identified (45). The

aceE and *aceF* genes, which encode components of pyruvate dehydrogenase, were also upregulated

during growth on lactate. The *pca* and *qui* genes, responsible for quinate and shikimate utilization,

were downregulated in ethanol and hexadecane cultures, indicating catabolite repression similar to

that reported for *A. baylyi* ADP1, where these genes are repressed in the presence of preferred carbon

sources (33, 38).

Substrate-specific regulation of aromatics degradation pathways

The transcriptomes of Tol 5 cells grown on toluene, phenol, or L-phenylalanine were also compared

with those grown on lactate (Fig. 5A, Table S4). On these three carbon sources (toluene, phenol, and

L-phenylalanine), genes of the PCA pathway were downregulated, whereas the genes for other

aromatic compounds degradation pathways were upregulated, with each pathway most strongly

induced by its corresponding substrate (Fig. 5B and C, Table S5). In toluene culture, the TOD pathway

genes showed the highest induction among those of aromatic compounds degradation pathways (Fig.

5B). The *fadL2*, which is located adjacent to the *tod* operon and encodes a putative toluene transporter,

was also upregulated, although it was reported that its deletion did not affect growth on toluene (39).
In phenol culture, the *mph* genes and the CAT pathway genes were highly upregulated (Fig. 5B and
C). Additionally, the TOD pathway genes were significantly upregulated in phenol culture, with
expression levels of the genes comparable to those observed in toluene culture (Fig. 5B).

For L-phenylalanine, two degradation pathways, the PAA pathway and the HMG pathway, were
predicted in Tol 5 (Fig. 3). The transcriptome data showed that the upregulation of the HMG pathway
genes was marked in L-phenylalanine culture (Fig. 5B). Additionally, the *aco* operon, which encodes
enzymes for acetoacetate assimilation via acetoin, was upregulated, supporting the involvement of the
HMG pathway in phenylalanine metabolism, in which acetoacetate is generated as one of the final
metabolites (Fig. 5D and 5E, Table S5).

Aromatic substrates also influenced downstream central metabolic pathways (Fig. S1, Table S5).
Growth on these compounds triggered a differential response in aconitate hydratase isozymes, with
*acnA_2 b*eing upregulated, whereas *acnB* was downregulated. Concurrently, *pckG,* encoding the key
gluconeogenic enzyme phosphoenolpyruvate carboxykinase (46), was downregulated in all three
conditions, indicating a reduced carbon flux from the TCA cycle into gluconeogenesis.

Expression of intracellular stress resistance genes under different carbon sources

In addition to carbon metabolic pathways, genes involved in environmental stress resistance showed
significant transcriptional changes depending on the carbon sources (Fig. 6, Table S5). Notably,
several genes related to oxidative stress resistance were upregulated in Tol 5 cells grown on
hexadecane, toluene, phenol, or L-phenylalanine. The genes encoding monofunctional heme catalases
(*katA* and/or *katE*) were highly upregulated under all four conditions. The Mn/Fe superoxide
dismutase genes (*sodA_1* and/or *sodA_2*) were repressed in hexadecane and L-phenylalanine cultures
but upregulated in toluene and phenol cultures. The periplasmic Cu/Zn superoxide dismutase gene
(*sodC*) was upregulated in toluene, phenol, and L-phenylalanine cultures. The alkyl hydroperoxide
reductase genes (*ahpC_1* and *ahpC_2*), which play a role in peroxide detoxification, and the
methionine-(*S*)-sulfoxide reductase and methionine-(*R*)-sulfoxide reductase genes (*msrA* and *msrB*),
which repair oxidized methionine residues, were highly upregulated in L-phenylalanine culture.
Concurrently, the trehalose synthesis genes (*otsA* and *otsB*), associated with osmotic stress resistance
(47, 48), were also upregulated on all four alkane and aromatic carbon sources. The *otsB* gene*,* which
encodes trehalose-6-phosphate phosphatase, exhibited low expression levels (log$_2$ CPM) but showed
high upregulation (log$_2$ FC). These results indicate that Tol 5 activates mechanisms for oxidative and
osmotic stress resistance during growth on alkanes and aromatic compounds.

DISCUSSION

Members of the genus *Acinetobacter* inhabit diverse environments, including soil, oceans,
freshwater, sediments, activated sludge, and polluted sites (20, 49). Several *Acinetobacter* strains,
such as the *A. calcoaceticus–A. baumannii* complex, are also opportunistic nosocomial pathogens
that exhibit multidrug resistance (50). Consequently, most previous research on *Acinetobacter*
species has focused on antibiotic resistance, genomic plasticity, horizontal gene transfer, and
pathogenicity, whereas the metabolic capabilities of this genus remain comparatively underexplored
(20, 49). Although *A. baylyi* ADP1 is a widely used model strain and its 3.5 Mb genome has been
well characterized (21, 51), this genome is smaller than those of many other *Acinetobacter* species,
and thus represents only a subset of the metabolic potential present across the genus (20, 21). In this
study, we characterized the carbon metabolism of the highly adhesive hydrocarbon-degrading
Acinetobacter strain Tol 5, which has a 4.8 Mb genome (19). Genome and transcriptome analyses
revealed that Tol 5 has limited capability to metabolize sugars but possesses an extensive repertoire
of genes for alkane and aromatic metabolism and regulates them in response to available carbon
sources. In particular, this study identified the presence of the TOD and HMG pathways in
Acinetobacter species and demonstrated that the genes involved in these pathways are specifically
induced during growth on toluene and L-phenylalanine, respectively. Tol 5 also harbors the PAA
pathway, which in *A. baumannii* is primarily associated with phenylalanine degradation but was
weakly induced by L-phenylalanine in Tol 5. Because the PAA pathway has been implicated in the
degradation of other aromatics in *P. putida*, including phenylethylamine, phenylethanol, styrene, and
n-phenylalkanoates (52), it may also contribute to the degradation of multiple aromatic substrates in
Tol 5. These findings demonstrate that Tol 5 possesses a unique repertoire of aromatic degradation
pathways, combining routes broadly conserved in the genus *Acinetobacter* with additional pathways
(such as TOD and HMG) that are rare in this genus. This metabolic versatility is comparable to that
of *Pseudomonas putida* strains, widely used chassis for aromatic compound bioconversion. During
growth on hydrophobic carbon sources, such as hexadecane and aromatic compounds, oxidative and
osmotic stress resistance genes were also upregulated in Tol 5, indicating that these substrates
impose cellular stress and that Tol 5 expresses the corresponding stress response. These findings
provide new insights into the metabolic diversity of *Acinetobacter* species and demonstrate that Tol
5 can serve as a model with metabolic pathways distinct from those of previously well-characterized
species.

Acinetobacter species widely exhibit the capacity for alkane degradation, as more than 95% of
strains encode the alkane monooxygenases AlkB and AlmA (20). Oil-degrading strains such as *A.*
venetianus RAG-1 and *A. oleivorans* DR1 possess multiple AlkB homologs, enabling efficient

utilization of *n*-alkanes (8, 53). Although Tol 5 possesses only one *alkB* gene on the genome, it also

carries a plasmid-encoded cytochrome P450 monooxygenase of the CYP153 family. CYP153, which

is a distinct alkane hydroxylase targeting short- and medium-chain *n*-alkanes, is typically found in

alkane-degrading bacteria lacking AlkB but is relatively rare among *Acinetobacter* genomes (28, 54).

Both *alkB* and the CYP153 gene were induced in Tol 5 grown on hexadecane, indicating that Tol 5

employs both types of alkane hydroxylases to oxidize hexadecane efficiently (Fig. 4C and Table S5).

The expression of aromatic compound degradation pathways is generally regulated by pathway-

specific transcriptional regulators to prevent unnecessary synthesis of corresponding enzymes and

accumulation of toxic intermediates (55). In the TOD pathway, genes encoding all reaction steps for

toluene degradation, from the initial oxidation catalyzed by toluene dioxygenase to the generation of

pyruvate and acetyl-CoA, are clustered in a single *tod* operon (Fig. 2). This operon is regulated by

the TodS-TodT two-component system, in which the TodS senses a broad spectrum of aromatic

effectors, autophosphorylates, and activates transcription of the *tod* operon through phosphate

transfer to the transcriptional activator TodT (56, 57). Transcriptome analysis in Tol 5 revealed

distinct responses to different substrates: growth on phenol induced not only the *cat* and *mph* genes

(the latter including genes encoding phenol monooxygenase components) but also the *tod* genes,

whereas growth on toluene strongly induced only the *tod* operon (Fig. 5BC). Previous studies have

demonstrated that the toluene dioxygenase complex TodABC1C2, encoded by the *tod* operon,

oxidizes phenol to catechol and hydroquinone, and that a mutant of *P. putida* F1 lacking this enzyme

cannot grow on phenol as a sole carbon source (58, 59). Taken together, these findings suggest that

Tol 5 uses both toluene dioxygenase and phenol monooxygenase for phenol degradation.

The *cat* operon in Tol 5, which encodes the CAT pathway responsible for catechol degradation,

includes a catechol 1,2-dioxygenase gene *catA* (*catA_3*). Notably, homologs of the *catA* gene are

also present in both the *mph* and *ben* operons (Fig. 2). The presence of *catA* within the *mph* operon is

distinct from that in *A. calcoaceticus* PHEA-2 previously reported (37). Transcriptome analysis

revealed that this *catA* gene (*catA_2*) was induced with other *mph* genes at similar expression levels

by phenol (Fig. 5C), indicating that it is transcriptionally regulated as part of the same operon (Fig.

2). This genetic organization is analogous to the *ben* genes described in *P. putida* strains and *A.*

baylyi ADP1, where CatA encoded by the *ben* operon converts the catechol generated by benzoate

oxidation into *cis,cis*-muconate, and the *cis,cis*-muconate activates the downstream *cat* operon (60-

62). Such coordinated regulation prevents the accumulation of catechol, which is toxic at high

concentrations (63). Therefore, the organization of the *mph* operon as well as the *ben* operon in Tol 5

likely enables fine-tuned transcription during the degradation of phenol and probably benzoate,

representing an optimized gene organization for the degradation of aromatic compounds.

Carbon catabolic pathways are often subject to carbon catabolite repression (CCR) (42). In *A. baylyi*

ADP1, the catabolite repression control protein (Crc) represses the *pca* and *cat* operons (33, 38).

Succinate and acetate induce strong repression, whereas pyruvate and lactate cause relatively weak

repression. Tol 5 encodes a Crc homologue (TOL5_01610) that shares 89% amino acid identity with

ADP1 Crc, and both the *pca* and *cat* genes were repressed in Tol 5 during growth on ethanol, where

acetate is expected to accumulate (Fig. 5B, Table S5). This indicates that Tol 5 exhibits a similar

CCR. In addition, the *tod*, *paa*, and *hmg* genes showed a similar expression pattern, with lower

expression in lactate and ethanol cultures and higher expression under other carbon sources (Fig.

5B). This pattern may reflect CCR-like regulation that prioritizes the utilization of readily

metabolizable carbon sources, such as lactate and ethanol, over hydrocarbon degradation. Given that

disruption of CCR has been reported to markedly alter the utilization of aromatic substrates (64),

modulation of CCR should be considered a key strategy to control the expression of aromatic

compound metabolic pathways in Tol 5 for bioproduction applications.

Alkane and aromatic oxygenases catalyze the initial oxidation of their substrates by forming transient

radicals at the active site, which can lead to the leakage of reactive oxygen species (ROS) (65).

Consequently, ROS are considered to be by-products of alkane and aromatic hydroxylation, and cells

must deploy protective mechanisms to scavenge them. For instance, *A. calcoaceticus* grown on

hydrocarbons exhibits increased superoxide dismutase activity, while both *A. oleivorans* DR1 cultured

on triacontane and *Paraburkholderia xenovorans* p2-fldX1 grown on 4-hydroxyphenylacetate

upregulate multiple ROS scavengers (53, 65, 66). In Tol 5 grown on hexadecane or aromatic

compounds, oxidative stress resistance genes, including *sodA_2*, *sodC*, *katA*, *katE*, *msrA*, and *msrB*,

were similarly induced (Fig. 6). Furthermore, the expression of TCA cycle enzymes is also likely to

be replaced according to the intercellular oxidative stress. In *E. coli*, ROS-sensitive aconitate hydratase

AcnB functions as the primary aconitate hydratase during the growth phase, while ROS-tolerant

isozyme AcnA is induced under oxidative stress conditions to maintain the TCA cycle (67, 68). Thus,

the expression shift from *acnB* to *acnA_2* in the TCA cycle of Tol 5 grown on aromatic compounds

suggests that Tol 5 also replaces metabolic enzymes to maintain central metabolism in response to

intracellular oxidative stress (Fig. S1). In summary, Tol 5 responds to oxidative stress according to the

available carbon source, not only by activating stress resistance genes but also by replacing vulnerable

metabolic enzymes to sustain central metabolism under cellular stress.

In conclusion, Tol 5 possesses a diverse set of metabolic pathways for alkanes and aromatic

compounds, regulates these pathways according to available carbon sources, and activates stress

resistance genes. Owing to its strong adhesiveness and ability to metabolize alkanes and aromatic

compounds, Tol 5 represents a promising microbial chassis for bioprocesses utilizing non-sugar

substrates.

MATERIALS AND METHODS

Genome sequence data analysis

The complete genome sequence of Tol 5, consisting of a chromosome and a plasmid, was obtained from the NCBI database (accession: AP024708 and AP024709). All amino acid sequences of the CDSs were searched against the UniProtKB/Swiss-Prot database (as of August 2024) using BLASTP, and the best-scoring hit for each CDS was used for functional annotation. The sequences were further searched against the KEGG database using BlastKOALA and KEGG automatic annotation server (accession: T11084). For CDSs, that could not be annotated with the KEGG database, additional searches were performed against the EggNOG database using eggNOG-mapper. Based on the annotations obtained, a central carbon metabolic pathway map was reconstructed using KEGG mapper. Peripheral metabolic pathways, including those involved in the degradation of alkanes and aromatic compounds, were inferred by comparing operon organization with those previously characterized in closely related bacterial species. Operon structures were predicted using two tools: Operon-mapper (69), which infers operons based on intergenic distances and gene functions, and Rockhopper (70), which additionally incorporates transcriptomic data. Functional annotations, operon predictions, and metabolic pathways were manually curated based on information from relevant literature.

Growth conditions

Acinetobacter sp. Tol 5 was cultured in 20 mL of basal salt (BS) medium (11) supplemented with a single carbon source in 100-mL Erlenmeyer flasks. Each of DL-lactate, ethanol, hexadecane, toluene, phenol, and L-phenylalanine was supplied as the sole carbon source at a carbon equivalent concentration of 3.3×10^{-2} mol/L. Precultures were prepared by growing Tol 5 in lysogeny broth (LB) medium supplemented with the corresponding carbon source. The pre-culture was washed twice with an equal volume of BS medium and then inoculated into 20 mL of culture medium at a 1:100 ratio. Four rectangular pieces of polyurethane foam support (10 × 10 × 10 mm, CFH-20, 20 pores/25 mm; Inoac Corporation, Nagoya, Japan) were placed in each flask. The flasks were then capped with silicone stoppers for lactate, ethanol, and L-phenylalanine, or with Viton rubber stoppers for hexadecane, toluene, and phenol, and incubated at 28 °C with shaking at 115 rpm.

RNA sequencing

Tol 5 cells during growth were harvested by centrifugation (8,000 × g, 4°C, 3 min), and the cell

pellets were stored at -30°C. For each growth condition, three biological replicates were prepared.

Total RNA was extracted using the RNA Prep Kit (Cica Genus, Tokyo, Japan) according to the

manufacturer's protocol. Ribosomal RNA was removed using the NEBNext rRNA Depletion Kit

(Bacteria) (New England Biolabs, Ipswich, MA, USA), and cDNA libraries were generated with the

NEBNext Ultra II RNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA, USA).

The cDNA libraries were sequenced on the Illumina NextSeq550. Raw FASTQ reads were quality-

filtered using fastp (version 0.23.2) and mapped to the Tol 5 genome (accession: AP024708 and

AP024709) using Bowtie2 (version 2.5.1) with default parameters. Mapped reads were counted

using featureCounts (version 2.0.6).

Differential gene-expression analysis

Raw read counts of CDSs were analyzed in R using the edgeR package (version 3.42.4). Genes with

extremely low expression were removed using the filterByExpr function. Normalization was

performed using the trimmed mean of M-values (TMM) method implemented in the

calcNormFactors function. Differential gene-expression analysis was conducted using the quasi-

likelihood pipeline. A design matrix was constructed incorporating the cells grown on six different

carbon sources (n = 3), with the data from those grown on lactate set as the reference group.

Dispersion estimates were obtained using the estimateDisp function, and model fitting was

performed with the glmQLFit function. Differential expression between each condition and the

control condition (lactate condition) was assessed using the glmQLFTest function. Genes with FDR

< 0.05, $\log_2 FC > 1$, and $\log_2 CPM > 3$ were extracted as significantly differentially expressed genes.

ACKNOWLEDGMENTS

The authors wish to acknowledge the Center for Gene Research, Nagoya University, for technical

support with the RNA sequencing. The authors also thank Yuki Ohara for kindly discussing genome

sequence data analysis and reviewing the manuscript.

This research was supported by the Graduate Program of Transformative Chem-Bio Research at

Nagoya University supported by MEXT (WISE Program) to SI, the Japan Science and Technology

Agency (JST) SPRING (Grant Number 8 JPMJSP2125) to SI, the Japan Society for the Promotion

of Science (JSPS) KAKENHI (Grant Number JP24H00043) to KH and SY, and the GteX Program

Japan Grant number JPMJGX23B4 to KH.

AUTHOR CONTRIBUTIONS

Shori Inoue: Conceptualization, Formal analysis, Investigation, Data curation, Writing - Original Draft. **Shogo Yoshimoto:** Conceptualization, Writing - Review & Editing. **Katsutoshi Hori:** Conceptualization, Supervision, Project administration, Writing - Review & Editing.

DATA AVAILABILITY

RNA-seq data reported are available in the DDBJ Sequenced Read Archive under the accession numbers PRJDB35996.

REFERENCE

1. Pontrelli S, Chiu T-Y, Lan EI, Chen FYH, Chang P, Liao JC. 2018. *Escherichia coli* as a host for metabolic engineering. *Metabolic Engineering* 50:16-46.

2. Liew FE, Nogle R, Abdalla T, Rasor BJ, Canter C, Jensen RO, Wang L, Strutz J, Chirania P, De Tissera S, Mueller AP, Ruan Z, Gao A, Tran L, Engle NL, Bromley JC, Daniell J, Conrado R, Tschaplinski TJ, Giannone RJ, Hettich RL, Karim AS, Simpson SD, Brown SD, Leang C, Jewett MC, Köpke M. 2022. Carbon-negative production of acetone and isopropanol by gas fermentation at industrial pilot scale. *Nature Biotechnology* 40:335-344.

3. Liu H, Tao X, Ntakirutimana S, Liu Z-H, Li BZ, Yuan YJ. 2024. Engineering *Pseudomonas putida* for lignin bioconversion into *cis-cis* muconic acid. *Chemical Engineering Journal* 495:153375.

4. Martín-González J, Montero-Bullón J-F, Muñoz-Fernández G, Buey RM, Jiménez A. 2025. Valorization of waste cooking oil for bioproduction of industrially-relevant metabolites in *Ashbya gossypii*. *New Biotechnology* 88:32-38.

5. Biggs BW, Bedore SR, Arvay E, Huang S, Subramanian H, McIntyre EA, Duscent-Maitland CV, Neidle EL, Tyo Keith E J. 2020. Development of a genetic toolset for the highly engineerable and metabolically versatile *Acinetobacter baylyi* ADP1. *Nucleic Acids Research* 48:5169-5182.

6. Luo J, McIntyre Emily A, Bedore Stacy R, Santala V, Neidle Ellen L, Santala S. 2022. Characterization of highly ferulate-tolerant *Acinetobacter baylyi* ADP1 isolates by a rapid reverse engineering method. *Applied and Environmental Microbiology* 88:e01780-21.

7.   Kurnia K, Efimova E, Santala V, Santala S. 2024. Metabolic engineering of *Acinetobacter baylyi*
ADP1 for naringenin production. *Metabolic Engineering Communications* 19:e00249.

8.   Liu J, Zhao B, Lan Y, Ma T. 2021. Enhanced degradation of different crude oils by defined
engineered consortia of *Acinetobacter venetianus* RAG-1 mutants based on their alkane
metabolism. *Bioresource Technology* 327:124787.

9.   Sabapathy PC, Devaraj S, Parthipan A, Kathirvel P. 2019. Polyhydroxyalkanoate production
from statistically optimized media using rice mill effluent as sustainable substrate with an
analysis on the biopolymer's degradation potential. *International Journal of Biological*
Macromolecules 126:977-986.

10.   D'Almeida AP, de Azevedo DCS, Melo VMM, de Albuquerque TL, Rocha MVP. 2024.
Bioemulsifier production by *Acinetobacter venetianus* AMO1502: Potential for bioremediation
and environmentally friendly applications. *Marine Pollution Bulletin* 203:116436.

11.   Hori K, Yamashita S, Ishii Si, Kitagawa M, Tanji Y, Unno H. 2001. Isolation, Characterization
and application to off-gas treatment of toluene-degrading bacteria. *Journal of Chemical*
Engineering of Japan 34:1120-1126.

12.   Ishikawa M, Shigemori K, Suzuki A, Hori K. 2012. Evaluation of adhesiveness of *Acinetobacter*
sp. Tol 5 to abiotic surfaces. *Journal of Bioscience and Bioengineering* 113:719-725.

13.   Ishikawa M, Nakatani H, Hori K. 2012. AtaA, a new member of the trimeric autotransporter
adhesins from *Acinetobacter* sp. Tol 5 mediating high adhesiveness to various abiotic surfaces.
PLOS ONE 7:e48830.

14.   Yoshimoto S, Aoki S, Ohara Y, Ishikawa M, Suzuki A, Linke D, Lupas AN, Hori K. 2023.
Identification of the adhesive domain of AtaA from *Acinetobacter* sp. Tol 5 and its application in
immobilizing *Escherichia coli*. *Frontiers in Bioengineering and Biotechnology* 10:1095057.

15.   Yoshimoto S, Ishii S, Kawashiri A, Matsushita T, Linke D, Göttig S, Kempf VAJ, Takai M, Hori
K. 2024. Adhesion preference of the sticky bacterium *Acinetobacter* sp. Tol 5. *Frontiers in*
Bioengineering and Biotechnology 12:1342418.

16.   Inoue S, Yoshimoto S, Hori K. 2025. A new target of multiple lysine methylation in bacteria.
Journal of Bacteriology 207:e00325-24.

17.   Yoshimoto S, Ohara Y, Nakatani H, Hori K. 2017. Reversible bacterial immobilization based on
the salt-dependent adhesion of the bacterionanofiber protein AtaA. *Microbial Cell Factories*
16:123.

18.   Usami A, Ishikawa M, Hori K. 2020. Gas-phase bioproduction of a high-value-added
516   monoterpenoid (*E*)-geranic acid by metabolically engineered *Acinetobacter* sp. Tol 5. *Green*
517   *Chemistry* 22:1258-1268.

518   19.   Ishikawa M, Hori K. 2021. Complete genome sequence of the highly adhesive bacterium
519   *Acinetobacter* sp. strain Tol 5. *Microbiology Resource Announcements* 10:10.1128/mra.00567-
520   21.

521   20.   Zhao Y, Wei HM, Yuan JL, Xu L, Sun JQ. 2023. A comprehensive genomic analysis provides
522   insights on the high environmental adaptability of *Acinetobacter* strains. *Frontiers in*
523   *Microbiology* 14:1177951.

524   21.   Barbe V, Vallenet D, Fonknechten N, Kreimeyer A, Oztas S, Labarre L, Cruveiller S, Robert C,
525   Duprat S, Wincker P, Ornston LN, Weissenbach J, Marlière P, Cohen GN, Médigue C. 2004.
526   Unique features revealed by the genome sequence of *Acinetobacter* sp. ADP1, a versatile and
527   naturally transformation competent bacterium. *Nucleic Acids Research* 32:5766-5779.

528   22.   Kannisto MS, Mangayil RK, Shrivastava-Bhattacharya A, Pletschke BI, Karp MT, Santala VP.
529   2015. Metabolic engineering of *Acinetobacter baylyi* ADP1 for removal of *Clostridium*
530   *butyricum* growth inhibitors produced from lignocellulosic hydrolysates. *Biotechnology for*
531   *Biofuels* 8:198.

532   23.   Ren X, Palmer Lauren D. 2023. *Acinetobacter* metabolism in infection and antimicrobial
533   resistance. *Infection and Immunity* 91:e00433-22.

534   24.   Alberti L, König P, Zeidler S, Poehlein A, Daniel R, Averhoff B, Müller V. 2023. Identification
535   and characterization of a novel pathway for aldopentose degradation in *Acinetobacter*
536   *baumannii*. *Environmental Microbiology* 25:2416-2430.

537   25.   Wohlfarth S, Hoesche C, Strunk C, Winkler UK. 1992. Molecular genetics of the extracellular
538   lipase of *Pseudomonas aeruginosa* PAO1. *Microbiology* 138:1325-1335.

539   26.   Rojo F. 2009. Degradation of alkanes by bacteria. *Environmental Microbiology* 11:2477-2490.

540   27.   Maier T, Förster HH, Asperger O, Hahn U. 2001. Molecular characterization of the 56-kDa
541   CYP153 from *Acinetobacter* sp. EB104. *Biochemical and Biophysical Research*
542   *Communications* 286:652-658.

543   28.   Nie Y, Chi CQ, Fang H, Liang JL, Lu SL, Lai GL, Tang YQ, Wu XL. 2014. Diverse alkane
544   hydroxylase genes in microorganisms and environments. *Scientific Reports* 4:4968.

29.   Kong W, Zhao C, Gao X, Wang L, Tian Q, Liu Y, Xue S, Han Z, Chen F, Wang S. 2021.

Characterization and transcriptome analysis of a long-chain *n*-alkane-degrading strain

Acinetobacter pittii SW-1. *International Journal of Environmental Research and Public Health.*

18(12):6365.

30.   Chen S, Cao L, Lv T, Liu J, Gao G, Li M, Sun L, Tian W, Tian Y, Li G, Ma T. 2024. Regulation

mechanism of the long-chain *n*-alkane monooxygenase gene *almA* in *Acinetobacter venetianus*

RAG-1. *Applied and Environmental Microbiology* 91:e02050-24.

31.   Lin GH, Hsieh MC, Shu HY. 2021. Role of iron-containing alcohol dehydrogenases in

Acinetobacter baumannii ATCC 19606 stress resistance and virulence. *International Journal of*

Molecular Sciences 22(18):9921.

32.   Phale PS, Malhotra H, Shah BA. 2020. Chapter One - Degradation strategies and associated

regulatory mechanisms/features for aromatic compound metabolism in bacteria, p 1-65. *In* Gadd

GM, Sariaslani S (ed), *Advances in Applied Microbiology*, vol 112. Academic Press.

33.   Fischer R, Bleichrodt FS, Gerischer UC. 2008. Aromatic degradative pathways in *Acinetobacter*

baylyi underlie carbon catabolite repression. *Microbiology* 154:3095-3103.

34.   Stuani L, Lechaplais C, Salminen AV, Ségurens B, Durot M, Castelli V, Pinet A, Labadie K,

Cruveiller S, Weissenbach J, de Berardinis V, Salanoubat M, Perret A. 2014. Novel metabolic

features in *Acinetobacter baylyi* ADP1 revealed by a multiomics approach. *Metabolomics*

10:1223-1238.

35.   Breisch J, Huber LS, Kraiczy P, Hubloher J, Averhoff B. 2022. The ß-ketoadipate pathway of

Acinetobacter baumannii is involved in complement resistance and affects resistance against

aromatic antibiotics. *Environmental Microbiology Reports* 14:170-178.

36.   Dal S, Trautwein G, Gerischer U. 2005. Transcriptional organization of genes for

protocatechuate and quinate degradation from *Acinetobacter* sp. strain ADP1. *Applied and*

Environmental Microbiology 71:1025-1034.

37.   Yu H, Peng Z, Zhan Y, Wang J, Yan Y, Chen M, Lu W, Ping S, Zhang W, Zhao Z, Li S, Takeo M,

Lin M. 2011. Novel regulator MphX represses activation of phenol hydroxylase genes caused by

a XylR/DmpR-type regulator MphR in *Acinetobacter calcoaceticus*. *PLOS ONE* 6:e17350.

38.   Bleichrodt FS, Fischer R, Gerischer UC. 2010. The β-ketoadipate pathway of *Acinetobacter*

baylyi undergoes carbon catabolite repression, cross-regulation and vertical regulation, and is

affected by Crc. *Microbiology* 156:1313-1322.

39.   Yoshimoto S, Hattori M, Inoue S, Mori S, Ohara Y, Hori K. 2025. Identification of toluene

degradation genes in *Acinetobacter* sp. Tol 5. *Journal of Bioscience and Bioengineering*
140:284-289.

40.   Hooppaw Anna J, McGuffey Jenna C, Di Venanzio G, Ortiz-Marquez Juan C, Weber Brent S,
Lightly Tasia J, van Opijnen T, Scott Nichollas E, Cardona Silvia T, Feldman Mario F. 2022. The
phenylacetic acid catabolic pathway regulates antibiotic and oxidative stress responses in
Acinetobacter. *mBio* 13:e01863-21.

41.   Teufel R, Mascaraque V, Ismail W, Voss M, Perera J, Eisenreich W, Haehnel W, Fuchs G. 2010.
Bacterial phenylalanine and phenylacetate catabolic pathway revealed. *Proceedings of the*
National Academy of Sciences 107:14390-14395.

42.   Görke B, Stülke J. 2008. Carbon catabolite repression in bacteria: many ways to make the most
out of nutrients. *Nature Reviews Microbiology* 6:613-624.

43.   Ishige T, Tani A, Sakai Y, Kato N. 2000. Long-chain aldehyde dehydrogenase that participates in
n-alkane utilization and wax ester synthesis in *Acinetobacter* sp. strain M-1. *Applied and*
Environmental Microbiology 66:3481-3486.

44.   Park C, Shin B, Park W. 2019. Alternative fate of glyoxylate during acetate and hexadecane
metabolism in *Acinetobacter oleivorans* DR1. *Scientific Reports* 9:14402.

45.   Morris FC, Jiang Y, Fu Y, Kostoulias X, Murray GL, Yu Y, Peleg AY. 2024. Lactate metabolism
promotes in vivo fitness during *Acinetobacter baumannii* infection. *FEMS Microbiology Letters*
371:fnae032.

46.   Sauer U, Eikmanns BJ. 2005. The PEP—pyruvate—oxaloacetate node as the switch point for
carbon flux distribution in bacteria: We dedicate this paper to Rudolf K. Thauer, Director of the
Max-Planck-Institute for Terrestrial Microbiology in Marburg, Germany, on the occasion of his
65th birthday. *FEMS Microbiology Reviews* 29:765-794.

47.   Liu J, Wang D, Tong R, Ye S, Zhao Y, Wu J, Gan Y. 2025. Engineered rhizobia with trehalose-
producing genes enhance peanut growth under salinity stress. *Agronomy* 15(4):974.

48.   Zeidler S, Hubloher J, Schabacker K, Lamosa P, Santos H, Müller V. 2017. Trehalose, a
temperature- and salt-induced solute with implications in pathobiology of *Acinetobacter*
baumannii. *Environmental Microbiology* 19:5088-5099.

49.   Mateo-Estrada V, Graña-Miraglia L, López-Leal G, Castillo-Ramírez S. 2019. Phylogenomics
reveals clear cases of misclassification and genus-wide phylogenetic markers for *Acinetobacter*.
Genome Biology and Evolution 11:2531-2541.

50.   Vijayakumar S, Indranil B, and Veeraraghavan B. 2019. Accurate identification of clinically
609   important *Acinetobacter* Spp.: An update. *Future Science OA* 5:FSO395.

610   51.   Durot M, Le Fèvre F, de Berardinis V, Kreimeyer A, Vallenet D, Combe C, Smidtas S,
611   Salanoubat M, Weissenbach J, Schachter V. 2008. Iterative reconstruction of a global metabolic
612   model of *Acinetobacter baylyi* ADP1 using high-throughput growth phenotype and gene
613   essentiality data. *BMC Systems Biology* 2:85.

614   52.   Nogales J, García JL, Díaz E. 2017. Degradation of aromatic compounds in *Pseudomonas*: a
615   systems biology view, p 1-49. *In* Rojo F (ed), *Aerobic Utilization of Hydrocarbons, Oils and*
616   *Lipids*. Springer, Cham.

617   53.   Park C, Shin B, Jung J, Lee Y, Park W. 2017. Metabolic and stress responses of *Acinetobacter*
618   *oleivorans* DR1 during long-chain alkane degradation. *Microbial Biotechnology* 10:1809-1823.

619   54.   van Beilen Jan B, Funhoff Enrico G, van Loon A, Just A, Kaysser L, Bouza M, Holtackers R,
620   Röthlisberger M, Li Z, Witholt B. 2006. Cytochrome P450 alkane hydroxylases of the CYP153
621   family are common in alkane-degrading eubacteria lacking integral membrane alkane
622   hydroxylases. *Applied and Environmental Microbiology* 72:59-65.

623   55.   Shingler V. 2003. Integrated regulation in response to aromatic compounds: from signal sensing
624   to attractive behaviour. *Environmental Microbiology* 5:1226-1241.

625   56.   Lacal J, Busch A, Guazzaroni M-E, Krell T, Ramos JL. 2006. The TodS–TodT two-component
626   regulatory system recognizes a wide range of effectors and works with DNA-bending proteins.
627   *Proceedings of the National Academy of Sciences* 103:8191-8196.

628   57.   Busch A, Guazzaroni M-E, Lacal J, Ramos JL, Krell T. 2009. The sensor kinase TodS operates
629   by a multiple step phosphorelay mechanism involving two autokinase domains. *Journal of*
630   *Biological Chemistry* 284:10353-10360.

631   58.   Höring P, Rothschild-Mancinelli K, Sharma ND, Boyd DR, Allen CCR. 2016. Oxidative
632   biotransformations of phenol substrates catalysed by toluene dioxygenase: A molecular docking
633   study. *Journal of Molecular Catalysis B: Enzymatic* 134:396-406.

634   59.   Spain JC, Zylstra GJ, Blake CK, Gibson DT. 1989. Monohydroxylation of phenol and 2,5-
635   dichlorophenol by toluene dioxygenase in *Pseudomonas putida* F1. *Applied and Environmental*
636   *Microbiology* 55:2648-2652.

637   60.   Silva-Rocha R, de Lorenzo V. 2012. A GFP-*lacZ* bicistronic reporter system for promoter
638   analysis in environmental gram-negative bacteria. *PLOS ONE* 7:e34675.

61.   Cosper Nathaniel J, Collier Lauren S, Clark Todd J, Scott Robert A, Neidle Ellen L. 2000.
Mutations in *catB*, the gene encoding muconate cycloisomerase, activate transcription of the
distalben genes and contribute to a complex regulatory circuit in *Acinetobacter* sp. strain ADP1.
Journal of Bacteriology 182:7044-7052.

62.   Bundy BM, Collier LS, Hoover TR, Neidle EL. 2002. Synergistic transcriptional activation by
one regulatory protein in response to two metabolites. *Proceedings of the National Academy of*
Sciences 99:7693-7698.

63.   Jiménez JI, Pérez-Pantoja D, Chavarría M, Díaz E, de Lorenzo V. 2014. A second chromosomal
copy of the *catA* gene endows *Pseudomonas putida* mt-2 with an enzymatic safety valve for
excess of catechol. *Environmental Microbiology* 16:1767-1778.

64.   Werner AZ, Cordell WT, Lahive CW, Klein BC, Singer CA, Tan ECD, Ingraham MA, Ramirez
KJ, Kim DH, Pedersen JN, Johnson CW, Pfleger BF, Beckham GT, Salvachúa D. 2023. Lignin
conversion to β-ketoadipic acid by *Pseudomonas putida* via metabolic engineering and
bioprocess development. *Science Advances* 9:eadj0053.

65.   Sazykin I, Sazykina M, Khmelevtsova L, Khammami M, Karchava S, Zhuravleva M,
Kudeevskaya E. 2016. Expression of SOD and production of reactive oxygen species in
Acinetobacter calcoaceticus caused by hydrocarbon oxidation. *Annals of Microbiology* 66:1039-
1045.

66.   Rodríguez-Castro L, Durán RE, Méndez V, Dorochesi F, Zühlke D, Riedel K, Seeger M. 2024.
The long-chain flavodoxin FldX1 improves the biodegradation of 4-hydroxyphenylacetate and
3-hydroxyphenylacetate and counteracts the oxidative stress associated to aromatic catabolism in
Paraburkholderia xenovorans. *Biological Research* 57:12.

67.   Cunningham L, Gruer MJ, Guest JR. 1997. Transcriptional regulation of the aconitase genes
(*acnA* and *acnB*) of *Escherichia coli*. *Microbiology* 143:3795-3805.

68.   Jordan P, Tang Y, Bradbury AJ, Thomson AJ, JR. G. 1999. Biochemical and spectroscopic
characterization of *Escherichia coli* aconitases (AcnA and AcnB). *The Biochemical journal*
344:739-746.

69.   Taboada B, Estrada K, Ciria R, Merino E. 2018. Operon-mapper: a web server for precise operon
identification in bacterial and archaeal genomes. *Bioinformatics* 34:4118-4120.

70.   McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA, Vanderpool CK,
Tjaden B. 2013. Computational analysis of bacterial RNA-Seq data. *Nucleic Acids Research*
41:e140-e140.

671
672

Fig. 1. Prediction of the central metabolic pathway of *Acinetobacter* sp. Tol 5. Genes related to the

central metabolic pathway were analyzed and listed in Table S2. Dashed arrows indicate metabolic

reactions for which the corresponding genes are not found in Tol 5 genome. Abbreviations: CIT;

citrate, ICIT; isocitrate, AKG; α-ketoglutarate, SUC-CoA; succinyl-CoA, SUC; succinate, FUM;

fumarate, MAL; malate, OAA; oxaloacetate, PRPP; phosphoribosyl pyrophosphate, AH; alkane

hydroxylase, ADH; alcohol dehydrogenase, ALDH; aldehyde dehydrogenase.

Fig. 2. Genetic organization of genes involved in the degradation of aromatic compounds in
Acinetobacter sp. Tol 5. Predicted operon structures are illustrated based on analyses using Operon-
mapper and Rockhopper. Genes within the same operon are represented by arrows of the same color.
Proposed gene names or predicted functions are labeled above each arrow, and genomic coordinates
are indicated below.

690 **Fig. 3.** Predicted degradation pathways of aromatic compounds in *Acinetobacter* sp. Tol 5,

691 reconstructed based on the genes shown in Fig. 2 and Table S2. Question marks and dashed arrows

692 indicate steps for which the genes encoding the corresponding enzymes have not been identified in

693 Tol 5.

Fig. 4. Transcriptome analysis of *Acinetobacter* sp. Tol 5 grown on lactate, ethanol, and hexadecane.
(**A**) Volcano plots comparing gene expression between lactate and ethanol (left) or hexadecane
(right) as sole carbon sources (n = 3 biologically independent samples). Red and blue dots represent
significantly upregulated and downregulated genes, respectively (FDR < 0.05, |$\log_2$ FC| > 1). (**B**, **C**)
Proposed ethanol (**B**) and hexadecane (**C**) degradation pathways in Tol 5 based on transcriptome
analysis. Genes encoding enzymes significantly upregulated under each condition are shown in red.

Fig. 5. Differential gene expression of *Acinetobacter sp.* Tol 5 grown on various aromatics, ethanol, and hexadecane, using lactate as the reference condition. (**A**) Volcano plots comparing gene expression between lactate and toluene (left), phenol (center), or L-phenylalanine (right) as sole carbon sources (n = 3 biologically independent samples). Red and blue dots represent significantly upregulated and downregulated genes, respectively (FDR < 0.05, |$\log_2$ FC| > 1). (**B-C**) Heat maps showing the average $\log_2$ FC of the genes involved in the five major aromatic degradation pathways (*tod*, *pca*, *cat*, *paa*, and *hmg* genes) (**B**), the *mph* genes (**C**), and the *aco* operon (**D**). The color gradient indicates the magnitude of the $\log_2$ FC values, with the deepest red representing ≥ 10 and the deepest blue representing ≤ -10. Asterisks (*) denote differentially expressed genes defined by |$\log_2$ FC| > 1, $\log_2$ CPM > 3, and FDR < 0.05. (**E**) Proposed aromatic substrates degradation pathways in Tol 5 based on transcriptome data. Genes encoding enzymes significantly upregulated under each condition are shown in red.

Locus_tag	Function	Gene	Substrate				
			Eth	Hex	Tol	Phn	Phe
TOL5_17330	Fe-Mn family superoxide dismutase	*sodA_1*		*			*
TOL5_23820	Fe-Mn family superoxide dismutase	*sodA_2*		*	*	*	
TOL5_31810	Fe-Mn family superoxide dismutase	*sodB*					
TOL5_04020	Cu-Zn family superoxide dismutase	*sodC*			*	*	*
TOL5_04460	catalase	*katA*			*		*
TOL5_13540	catalase HPII	*katE*		*	*	*	*
TOL5_02070	alkyl hydroperoxide reductase	*ahpC_1*				*	*
TOL5_28070	alkyl hydroperoxide reductase	*ahpC_2*					*
TOL5_05100	peptide methionine sulfoxide reductase MsrA	*msrA*					*
TOL5_24000	methionine sulfoxide reductase B	*msrB*			*		*
TOL5_03100	thiol:disulfide interchange protein	*dsbC_1*					*
TOL5_08280	thiol:disulfide interchange protein	*dsbC_2*					
TOL5_42750	thiol:disulfide interchange protein	*dsbC_3*					
TOL5_13410	trehalose-6-phosphate synthase	*otsA*		*	*	*	*
TOL5_13420	trehalose-6-phosphate phosphatase	*otsB*					

log$_2$ fold change vs. lactate culture −5 0 5	**Eth**: Ethanol **Hex**: Hexadecane **Tol:** Toluene
	Phn: Phenol **Phe**: L-Phenylalanine

Fig. 6. Heat map showing the average log$_2$ FC of genes involved in oxidative stress and osmotic stress resistance. The color gradient indicates the magnitude of the log$_2$ FC values, with the deepest red representing ≥ 5 and the deepest blue representing ≤ -5. Asterisks (*) denote differentially expressed genes defined by $|\log_2 \text{FC}| > 1$, $\log_2 \text{CPM} > 3$, and FDR < 0.05.