A sleep promoting role of phototransduction in *Drosophila melanogaster*

Yu-Chien Hung[1], Mehran Akhtar[1], Nithish Sattoju[1], [#]Xinghua Li[1], [#]Steven Head[1], [#]Tobias Ollerenshaw[1,*], [#]Clelia

Siefer-Gaillardin[1,3], [#]Jashmine Arulchelvan[1,], Ben Warren[1,2], [⊠]Ko-Fan Chen[1]

1. Division of Genetics and Genome Biology, University of Leicester, Leicester, UK, LE1 7RH

2. School of Life Sciences, Keele University, Staffordshire, UK, ST5 5BG

3. Magistère Européen de Génétique, University Paris Cité, Paris 75013, France.

[#]These authors contribute equally

[*]Current addresses: **T.O.**: Mary Lyon Centre at MRC Harwell, Harwell Campus, Oxfordshire, UK, OX11 0RD.

[⊠]corresponding author: kc280@leicester.ac.uk

Abstract:

Daily sleep-wake cycle is a conserved behaviour defined by locomotion quiescence and enhanced

responsive threshold to sensory stimuli. Both circadian rhythm and a homeostatic process determine

the daily sleep profile, which is also regulated by environmental light, a major sensory input to

regulate circadian rhythm and alertness. With decades of investigation in *Drosophila*, the cellular

and circuital mechanism underlying light-mediated circadian synchronization are well-established,

yet the direct relationship between light/visual input and sleep remains unclear. To address this

knowledge gap, we have started an investigatory survey of sleep behaviour using classic mutant lines

to manipulate phototransduction and downstream neural transmission. We observed consistent day

sleep fragmentation in flies with mutations in multiple phototransduction components. We also

found hyperpolarised *Drosophila* photoreceptor resulted in shorter day sleep. We found a severe

reduction in locomotor speed in several visual mutants during normal waking time preventing

assessment of their sleep-linked immobility. In summary, we provide a rigorous quantification of

several phototransduction genes and reveal the key role of visual input to promote sleep.

1. Introduction

Sleep is vital for health. Disruptions in timing and/or amount of sleep has been linked to various metabolic and neurological diseases. Across animal kingdom, sleep is defined by behavioural quiescence and increased arousal threshold to environmental stimuli such as light or mechanic vibrations. Arousal threshold within a period of sleep varies in stages that are associated to specific electrophysiological signatures in electroencephalogram (EEG) and local field potentials (LFPs) in the brain (Lakhiani et al., 2023). With its versatile genetic toolkit and conserved behavioural and electrophysiological features of sleep (Jagannathan et al., 2024; Shafer and Keene, 2021), the fruit flies *Drosophila* has been used for identifying the biological basis and the conserved function of sleep for the past two decades: In parallel to murine investigations, studies in *Drosophila* have now demonstrated that neuronal redox status may form a cellular basis of sleep (Hartmann and Kempf, 2023) and sleep stages play roles in synaptic scaling that is important for learning and memory (Dissel et al., 2015; Gilestro et al., 2009; Tainton-Heap et al., 2021; Yap et al., 2017). Recent investigations in *Drosophila* also show that sleep is required for effective maintenance of the immune system, gut microbiome, and waste clearance (Alphen et al., 2021; Vaccaro et al., 2020; Vincent et al., 2022).

The onset and the duration of daily sleep are regulated by two intrinsic mechanisms: the circadian rhythm and homeostasis. The circadian rhythm defines the baseline sleep-wake level across 24 hours, whereas the homeostatic process maintains the overall level of sleep amount. In *Drosophila*, the neural circuits that generate circadian rhythm reside in the 242 so-called clock neurons (Reinhard et al., 2024) that express the core molecular clock mechanism, whereas the homeostatic control of sleep are distributed among neural domains including Fan-Shaped Body (FSB) and Ellipsoid Body (EB) (Jones et al., 2025; Shafer and Keene, 2021). The two mechanisms are likely coordinated as a recent study also demonstrated that clock neurons modulate the FSB-mediated sleep homeostat (Hsu et al., 2025).

Light is a prominent environmental stimulus to regulate both mechanisms in order to maintain animal's temporal niches with sufficient sleep. Such synchronisation between environmental light regime and the neural activity of clock neurons is called circadian photoentrainment. Similar to the mammalian system, visual inputs are part of photoentrainment in *Drosophila*. Visual system in the fly is comprise of peripheral light sensing neurons (photoreceptors in compound eyes, HB-eyelet and ocelli) and their downstream visual processing brain areas called the optic lobes and ventrolateral neuropils (Nériec and Desplan, 2016). Light signal is converted into membrane electrical depolarisation in the photoreceptors in compound eye and HB-eyelet by activation of cation channel

triggered by the *Rhodopsin* 1-6 (*Rh1-6*) linked phototransduction/GPCR signal cascades (Hardie,

2012). Activation of these two types of light-sensing neurons then release histamine or acetylcholine

to communicate with clock neurons either directly or indirectly via neural network in optic lobes

(Alejevski et al., 2019; Ogueta et al., 2020, 2018; Rieger et al., 2003; Schlichting et al., 2016; Tang et

al., 2022; Xiao et al., 2023). Apart from visual system, *Drosophila* molecular clock can also be

entrained by the cell autonomous light-sensing mechanisms mediated by *cryptochrome* (*cry*),

quasimodo (*qsm*) or *Rhodopsin 7* (*Rh7*) in clock neurons (Chen et al., 2011; Ni et al., 2017; Peschel et

al., 2009).

Light can directly regulate sleep by either promoting or supressing sleep via different neural

pathways in the murine system (Altimus et al., 2008; Pilorz et al., 2016). As a crepuscular species,

Drosophila sleep both in the day with the light and in the night in the darkness. While current

evidence indicate that distinct neurogenetic network may control day and night sleep (Chen et al.,

2019; Damulewicz et al., 2020; Hsu et al., 2025; Ishimoto et al., 2012; Lamaze et al., 2018; Sengupta

et al., 2019), visual experience in the light was recently found to promote sleep in *Drosophila*

(Kirszenblat et al., 2019), suggesting light and/or vision per se can serve as sleep homeostat drives,

similar to vertebrates (Horne and Walmsley, 1976; Korf et al., 2017; Miyamoto et al., 2003).

Despite growing research interests (Mazzotta et al., 2020), the mechanism underlying light-mediated

regulation of *Drosophila* sleep homeostat is far from clear: a previous study showed that synaptic

inhibition in *Drosophila* eye photoreceptor promote sleep (Damulewicz et al., 2020), yet the

optogenetic activation of all eye photoreceptors in another investigation did not result in sleep

suppression (Kirszenblat et al., 2019). Moreover, artificial activation of the direct downstream target

of eye photoreceptor, lamina monopolar neuron 2 (L2 LMC), resulted in drastic sleep loss (Muraro

and Ceriani, 2015). This finding is in contradiction to the increased sleep observed in flies with

reduced histamine (Oh et al., 2013), which is the main neurotransmitters released from

photoreceptors to inhibit L2 LMC neurons (Pantazis et al., 2008a). A recent systematic study of

mutants with defects in various light input pathway indicates that only *Rh7* is required for shade

sleep preference in the day, but the study did not investigate the baseline sleep level in these

mutants (Meyerhof et al., 2024).

In this study, we investigated the role of light input pathways in *Drosophila* sleep by focusing on the

role of visual system. Using the conventional infrared beam-based sleep assay (Chen et al., 2019;

Chiu et al., 2010), we surveyed the baseline sleep profiles for an array of fly lines with mutations in

genes involved in a) phototransduction, and the downstream histaminergic neurotransmission.

(Hardie, 2012) as well as hyperpolarising photoreceptors. Unlike night sleep, day sleep in *Drosophila*

is highly modifiable by sex, mating and nutrient status (Dove et al., 2017; Duhart et al., 2023; Isaac et

al., 2009; Wu et al., 2018). Typically, mated females can sleep as little as 150-200 minutes as

compare to 300-400 minutes in males in the 720 minutes of day time. In order to capture both sleep

increase and decrease upon genetic manipulation, we focused our investigation in virgin male flies.

By carefully examining sleep parameters and the phototransduction/neurotransmission defects in

electroretinogram (ERG), we identified a novel role of visual phototransduction in driving sleep.

2. Materials and methods

2.1 Fly husbandry

Drosophila strains are kept in the standard fly food (glucose 6.93% w/v, agar 0.5% w/v, cornmeal,

6.3% w/v, yeast, 4.38% w/v, propionic acid, 0.26% v/v, 0.2% v/v nipagin) at 25°C 12-hour light dark

cycle. The long-term stocks are kept in the same food and light regime at 18°C. The Canton S and

Oregon R wildtype strains are obtained from Kyriacou Lab (University of Leicester, UK), while w^{iso31}

(i.e. *iso*) control is obtained from Jepson lab (University College London, UK). Following visual

mutants, Gal4 and UAS transgenic are ordered from Bloomington stock centres (stock number): trp^1

(5692); trp^{MB} (23636); $trpl^{302}$ (31433); $trpl^{MB}$ (29134); $ninaE^8$ (2001); $ninaE^5$ (3531); $ninaE^{17}$ (5701);

$norpA^{P24}$ (9048); hdc^{MB} (25260); hdc^{JK910} (64203); ort^1 (1133); ort^{attP} (84559); *Rh1-Gal4* (8691); *gmr-*

Gal4 (1104); *UAS-kir2* (6595); *UAS-TeTxLC* (28837); *UAS-myr-mRFP* (7118). The 1st and 3rd

chromosomes of hdc^{JK910} were replace with those from w^{iso31} stock via following balancer crossings

with w^{iso31C};*Sco/CyO; MKRS/TM6B* and w^{iso31C};*Sco/CyO; +/+* (gifts from Jepson lab). All Gal4 and UAS

transgenic are brough into w^{iso31} background via five generation outcrossing following *[w+]* eye

colour marker cassette. For *Rh1-Gal4* outcrossing, we use *UAS-myr-mRFP/+; Rh1-Gal4/+* and follows

the eye RFP signal as visible marker under fluorescent scope for each generation. The final stock are

then established by balancer crossing with w^{iso31C};*Sco/CyO; MKRS/TM6B and then* w^{iso31C};*+/+;*

MKRS/TM6B for removing UAS-myr-mRFP.

2.2 Sleep recording and analysis

Prior sleep experiment, *Drosophila* strain or crossings are reared in a richer food recipe (wheat germ

1% w/v, soy 0.8% w/v, glucose/sucrose 3.3%/1.5% w/v, treacle 3% v/v, agar 1% w/v, cornmeal 1.5%

w/v, yeast 3.5% w/v, propionic acid 0.5% v/v, nipagin 0.1% v/v), as to keep the rearing condition

consistent with previous experimental conditions (Chen et al., 2019, 2011; Lamaze et al., 2018). Also,

we notice flies are generally less reproductively successful and motile in standard food. The sleep

recording are performed as described in (Chen et al., 2019): three to four days old virgin adult male

are transferred to typical behaviour tubes containing food (2% w/v agar and 4% w/v sucrose at one

end and cotton at the other end), and housed in typical Drosophila Activity Monitor (DAM,

Trikinetics Inc.) for 3 days of 12hours light-dark cycle at 25°C in a temperature and light controlled

incubator (MIR-254-PE, PHCbi). The lighting source is a day light fluorescent lamp (FL15D, 6500K)

with intensity between 1.2 and 6.7 w/m^2 depending on the distance to the lamp. The locomotion of

each fly is recorded as infrared beam crossing per minutes. The locomotion data for day 3 are

outputted into a previously customised Excel Calculator (Chen et al., 2019) to calculate all sleep

episodes (any period of immobility ≥ 5 minutes) along 24 hours and generate the following

parameters: **overall sleep profile**: minutes of sleep per 30 mins across 24 hours; **day sleep**: total

minutes in sleep during 12 hours of day; **night sleep**: total minutes in sleep during 12 hours of night;

day average sleep bout length: day sleep minutes divided by total numbers of sleep bout during 12

hours of day; **night average sleep bout length:** night sleep minutes divided by total numbers of sleep

bout during 12 hours of night; **day wake activity**: number of beam crosses divided by total wake

minutes during 12 hours of day**; night wake activity**: number of beam crosses divided by total wake

minutes during 12 hours of night.

2.3 Electroretinogram (ERG)

Three-to-five days old adult virgin male flies were inserted into a p1000 plastic pipette tip, so that its

head protruded from the narrow ending. The head was fixed using melted honey wax applied

between the thorax and head without blocking the mouthpart and eyes. The fly mounted pipette tip

was kept in the dark for 20 mins and then transferred into the ERG recording rig. Glass electrodes

(Borosilicate) with resistance of ~5 MO and filled with 3 M KCl were inserted into the compound eye

(recording) and head (reference). Each fly was then rest in the dark for 3 minutes before exposed to

four light intensities: 0.004, 0.02, 0.1 and 0.5 w/m^2 at 540nm LED green light source (M530L4-C1,

ThorLab). These intensities reside in the liner range of receptor potential response. We stimulated

with 1000 ms exposure to each light intensity and repeated two further times to get an average of

three presentations at each light intensity for each fly. The potential difference between the

electrodes was amplified 10x, and acquired by Neuroprobe Amplifier model 1600 and 8-channel

data acquisition interface DA-8 (A-M systems) using the SciWorks Data Acquisition & Analysis Suite.

2.4 Data display and statistics

`ggplot2` package in R is used to display sleep and ERG data. X-Y plots are used to display overall

sleep profile for each comparing genotypes. Boxplots with individual values are used to display sleep

parameters data. We calculated the average voltage changes of on transient, receptor potential and

off transient in the ERG profiles of each replicate for each fly (Figure 1H). The value for each fly is

then plotted against light intensity on logarithmic scales by `ggplot2`. Considering the variability in

sleep data (Chen et al., 2019; Lamaze et al., 2018), statistical significance are identified by Kruskal-
Wallis rank sum test for non-parametric multiple comparison (`kruskal.test`). Pairwise Wilcoxon
rank sum exact test with multiple comparison Benjamini-Hochberg correction
(`pairwise.wilcox.test`) is used to significant difference between pairwise genotype. ERG
voltage changes data are pooled among four intensities and used to identify genotypical difference
via Pairwise Wilcoxon rank sum exact test. Significance level is set at adj.p<0.05.   Effect size *r* is
calculated by `wilcox_effsize` with magnitude categories: small; moderate and large assigned. All
statistic comparisons, p values, effect size and sample size are listed in Table1 to Table 6 for the
corresponding Figure 1 to Figure 5.

3. Result

3.1 Defects in phototransduction genes caused short day sleep bout

To verify the role of visual phototransduction genes in sleep regulation, we selected eight mutant strains: *Rhodopsin 1/ninaE*, *Phospholipase C/norpA*, and the two cation channels *trp* and t*rpl*. We included both classic point mutations as well as more recently generated and characterised mutations using P-element insertion when available. To analyse the details of sleep-wake profiles under typical day of 12 hours light-12 hours dark, we examine the three conventional parameters for both day and night period: total sleep minutes, average sleep bout minutes and the waking activity.

3.1.1 *ninaE*

Upon exposure to light, 11-cis 3-hydroxy-retinal, the chromophore in the Rhodopsin protein undergoes photoisomerization, transforming Rhodopsin to Metarhodopsin and then trigger downstream GPCR signal cascade. Rhodopsin 1/ninaE protein is the main light-sensing GPCR in the outer photoreceptor R1-R6 of *Drosophila* compound eye. Reduction or loss of ninaE abolish or reduce block photoisomerization and therefore light induced photoreceptor potential and transients that is typically measurable for R1-R6 in ERG (Hardie, 2012).

Large part of visual tasks including navigation, phototaxis, optomotor and escapes required R1-R6 activity. We therefore first tested sleep profile of the class null allele *ninaE*[17] (O'Tousa et al., 1985). During the day, *ninaE*[17] mutants shows lower midday sleep peak in the sleep profile (Figure 1A) but the similar amount of total sleep minutes is found as compared to the white eye control, *w*[iso31] (here after *iso*, Figure 1Di). However, we found that the average sleep bout minutes in *ninaE*[17] is shorter than control, indicating a sleep fragmentation (Figure 1Bi). During the night, we observed an increased sleep amount (Figure 1Gi) while the average sleep bout minute is similar to control (Figure 1Ei). Since the sleep measurement in *Drosophila* depends on locomotion, the sleep increase can be falsely claimed by defect in locomotion. Therefore "waking activity", i.e., locomotor activity per waking minutes, was designed as a conventional measure to detangle such compound scenario (Andretic and Shaw, 2005; Harbison and Sehgal, 2008). We applied this measurement and found that waking activity are similar between *ninaE*[17] and control in both day and night (Figure 1Ci,Fi).

ninaE[17] flies contain an secondary *ebony*[1] mutation that impacts sleep-wake cycles (Rossi et al., 2015; Suh and Jackson, 2007). We therefore tested another two *ninaE* hypomorphs *ninaE*[5] and *ninaE*[8] (Scavarda et al., 1983). Both mutants show similar midday sleep peak reduction as seen in *ninaE*[17] flies (Figure 1A) This observation is consistent with significant reduced day sleep minutes in either mutant as compared to controls (Figure 1Dii-iii). Similarly, both mutants also show shorter

average sleep bout minutes in the day (Figure 1Bii-iii). Intriguingly, the loss of sleep in the two *ninaE*

hypomorphs coincide with loss of wake activity in the day (Figure 1Cii-iii). During the night, *ninaE*

hypomorphs both shows increased sleep (Figure 1Gii-iii) but only *ninaE^8* mutant shows significant

increase of the average sleep bout minutes as compared to control (Figure 1Eii-iii). However,

mutants severely reduce their wake activity (Figure 1Fii-iii with effect size *r* between 0.67 and 0.82,

Table 1). Therefore, the increased night sleep may simply reflect loss of locomotor activity. To

confirm *ninaE* mutants harbour phototransduction defects, we measured light-evoked receptor

potentials through Electroretinograms (ERG, Figure 1H). Consistent with previous reports, *ninaE^{17}*

and *ninaE^8* exhibit reduced light induction of receptor potential, on and off transients (Figure 1I-J)

whereas *ninaE^5* retain certain function (see normal receptor potential and off-transient in *ninaE^5*,

Figure 1J). Take results from the three mutants together, we conclude the reduction or loss of

Drosophila Rhodopsin 1 (ninaE) protein result in consist reduced day sleep bout length. This finding

indicates that phototransduction is required to drive day sleep pressure. However, recent studies

have found a non-visual function of ninaE (eg, Zanini et al., 2018). To verify the shorter day sleep

bouts is linked to phototransduction or just pleotropic function of ninaE protien, we further

examined the roles in sleep for three other proteins key for phototransduction.

3.1.2 *norpA*

Metarhodopsin triggers Gqα activation and enzymatic reaction of Phospholipase C (PLC) that

produce secondary messenger. norpA protein is the main PLC in the visual system, and the classic

mutant *norpA^{p24}* exhibit severe visual defects (Gao et al., 2008). The sleep profile of *norpA^{p24}* showed

decline of day sleep from the midday and a secondary dip in the midnight as compared to controls

(Figure 2A). The changes in sleep profile coincide with reduced sleep in the day (Figure 2D) and

shorter bout minutes in both day and night (Figure 2B, 2E). Similar to *ninaE^5* and *ninaE^8*, *norpA^{p24}*

mutant have low wake activity (Figure 2C, 2F, effect size 0.39/0.75, Table 2). We also confirm

abnormal ERG profiles in *norpA^{p24}* (Figure 2H).

3.1.3 *trp and trpl*

Activation of the two ion channels trp and trpl in the last step of phototransduction cause cation

influx resulting in depolarisation of photoreceptor. We therefore examine the sleep profile of four

mutant lines, two P-element insertion mediated mutants: *trp^{MB}*, *trpl^{MB}* (Chen et al., 2015) and two

classic null mutants: *trp^1*, and *trpl^{302}* (Cosens and Manning, 1969; Leung et al., 2000; Niemeyer et al.,

1996). Except for *trp^1*, these lines are white eyes as *ninaE* and *norpA* mutants as described above.

trp¹ mutant were first isolated from red eye wild type but the precise background is not known

(Cosens and Manning, 1969), so we compare its sleep profile to two widely used wild type in sleep

and vision research: *Canton S* and *Oregon R.* Significant reduction in day and night sleep amount and

average sleep bout length are shown as compared to either wild type flies (Figure 3Ai, 3Bi and 3Ei,

effect size: 0.29-0.54, Table 3). The wake activity of *trp¹* is comparable to *Canton S* but higher than

Oregon R, and has limited impact to sleep loss (Figure 3Ci and 3Fi). The sleep profile of *trp^MB* is

distinct from *trp¹, norpA* or *ninaE* mutants, with day sleep level similar to control but a significant

night sleep loss (Figure 3Aii, 3Gii). There are no significant changes in sleep bout minutes detected in

trp^MB flies (Figure 3Aii and 3Eii). Although, *trp^MB* flies sleep a similar amount during the day,

compared to control, their wake activity is significantly higher (Figure 3Cii).

This relative normal sleep-wake profile may relate to the less severe reduction in receptor potential

and light on/off transients in the ERG of *trp^MB* as compared to those of the *trp¹* null mutant (Figure

3Hi and 3Hii, effect size 0.11-0.83 vs 0.78-0.85, Table 3).

On the other hand, both *trpl³⁰²* and *trpl^MB* mutants exhibit day sleep profile similar to *ninaE* or *norpA*

mutants (Figure 3Aiii and 3Aiv), with significant reduced day sleep amount and average bout

minutes (Figure 3Biii, 3Biv, 3Diii and 3Div). Night sleep profiles and wake activities for the two *trpl*

mutants diverged: *trpl³⁰²* sleep more whereas *trpl^MB* sleep less (Figure 3Giii, 3Giv). No significant

changes in night sleep bout length were observed in both mutants as compared to the controls

(Figure 3Eiii and 3Eiv). *trpl³⁰²* mutants generally have low wake activity whereas *trpl^MB* maintain

relative normal wake locomotion as compared to the controls (Figure 3Ciii, 3Civ, 3Fiii and 3Fiv).

trpl³⁰² mutant is in *cn* and *bw* mutation background, which may affect the sleep. We therefore tested

w; trpl³⁰²/MB flies to reduce the impact of *cn bw* mutation. Similar to *trpl³⁰²* and *trpl^MB*, the *w;*

trpl³⁰²/MB transheterozygote mutant showed shorter day sleep and sleep bout length (Figure 3Biv). As

seen in previous reports (Chen et al., 2015; Leung et al., 2000) both mutants have limited impact on

ERG profile, *trpl³⁰²* showed large reduction of on/off transient (effect size 0.73/0.81, Table 4),

whereas *trpl^MB* maintain a normal ERG, possibly due to synaptic compensatory mechanism (Figure

3Hiii). Nevertheless, our data indicate the reduced trpl function and expression level cause sleep loss

in the day.

3.2 Divergent sleep function in histaminergic pathway

Drosophila photoreceptors can be divided into Rh1-expressing (R1-R6) and non-Rh1-exppressing

neurons (R7/R8). The R1-R6 photoreceptors form synapses onto lamina monopolar neurons (LMC),

while the R7/R8 send their output to medulla in the optic lobes. Histamine is the primary

neurotransmitters for Rh1-expessing photoreceptor. Therefore, we test if mutants with defects in

histamine synthesis or postsynaptic reception show altered sleep. Histidine decarboxylase (*hdc*) is

the main histamine synthesis enzyme (Pantazis et al., 2008b). We investigated the sleep profiles for

the two *hdc* mutants (*hdc^{JK910}* and *hdc^{MB}*) that were characterised previously and showed reduced

histamine levels in the photoreceptors (Burg et al., 1993; Stenesen et al., 2015).

As expected, both mutants selectively reduce on/off transients' levels, the components in ERG that

reflect synaptic transmission between photoreceptors and LMCs (Figure 4Hi). However, the sleep

profiles between the two mutants diverged: *hdc^{JK910}* showed increased day sleep and bout length,

whereas *hdc^{MB}* showed decreased day sleep and bout length (Figure 4Bi and 4Di). The wake activities

in the day are similar between *hdc* mutants and controls (Figure 4Ci). During the night, both *hdc*

mutants sleep less with shorter sleep bouts (Figure 4Ei and 4Gi). Intriguingly, both *hdc* mutants have

reduced wake activities compared to the controls at night (Figure 4Fi). Two histamine receptors ort

and HisCl1 are detected in *Drosophila* CNS, and they are both histamine gated chloride channels

(Pantazis et al., 2008b). Transcriptomic data in *Drosophila* visual system suggest that ort express

selectively in neurons postsynaptic to photoreceptors, whereas HisCl1 express almost exclusively in

photoreceptors as auto-receptors (Davis et al., 2020a). We therefore verify the sleep profile in ort

mutants to investigate the role of histamine signal downstream to photoreceptors.

First we monitored sleep profile of the classic *ort^1* mutant against *Canton S* as they share the same

genetic backgrounds (Inoue et al., 1989). A small but significant day time sleep loss are detected in

ort^1 flies (Figure 4Dii, effective size 0.27, Table 5). *ort^1* sleep for longer bouts during the day but have

higher day wake activity compared to *Canton S* (Figure 4Bii and 4Cii). In contrast to *hdc* mutants, no

night sleep phenotype is detected in *ort^1* (Figure 4Eii-4Gii). We also tested a recent CRISPR-

mediated null mutation *ort^{attp}* in the white eye background (Deng et al., 2019). The sleep profile of

ort^{attp} is distinct from *ort^1*, showing more sleep as compared to control (Figure 4Diii and 4Giii).

However, the wake activities of *ort^{attp}* are severely reduced (Figure 4Ciii and 4Fiii, effective size

0.55/0.60, Table 5) potentially compounding the observed sleep phenotype. Intriguingly, shorter day

sleep bouts and small reduction of night sleep are detected in *ort^{attp/+}* heterozygotes (Figure 4Biii and

4Giii), accompany improved wake activity (Figure 4Ciii and 4Fiii, wake activity *ort^{attp}* vs *ort^{attp/+}*, Table

5).

Both *ort* mutants exhibit ERG profile with loss of on/off transients, while *ort^1* also showed loss of

receptor potential (Figure 4Hii).

3.3. Hyperpolarisation of Rh1-expressing photoreceptor causes short day sleep bout

Our finding so far indicates that seven of the eight examined phototransduction mutants exhibit day

time sleep defects. However, there is divergence between sleep and ERG phenotypes in *trpl* and *trp*

mutants. Moreover, altered histaminergic pathway show divergent sleep defect in day or night.

Thus, we further test if direct inhibition of photoreceptor excitability causes sleep defects. At first,

we applied Gal-UAS system to block synaptic transmission from all photoreceptors by expressing

UAS-TeTxLC (tetanus toxin light chain) with gmr-Gal4 as in (Damulewicz et al., 2020). However, we

were not able to obtain viable adult flies even after maintaining the culture in 22°C to reduce the

known overexpression-related toxicity (Rezával et al., 2007; Sweeney et al., 1995). We next used

UAS-Kir2.1 (Baines et al., 2001; Nitabach et al., 2002) to hyperpolarise all or just Rh1-expressing

photoreceptors by gmr-Gal4 or Rh1-Gal4. Consistent with the finding for *ninaE* and *norpA* mutants,

gmr-Gal4>uas-Kir2.1 and *Rh1-Gal4>uas-Kir2.1* flies showed reduced day sleep as compared to

Gal4>+ or *+>uas-Kir2.1* controls (Figure 5Ai-ii, 5Di-ii), but only *Rh1-Gal4>uas-Kir2.1* flies exhibit

reduced sleep bout minutes in the day (Figure 5Bi-ii). No obvious changes in the analysed night sleep

parameters in both *gmr-Gal4>uas-Kir2.1* and *Rh1-Gal4>uas-Kir2.1* flies as compared to the controls

(Figure 5E-G). This is partly because the unexpected low sleep exhibited in *+>uas-Kir2.1* controls

(*gmr-Gal4>+* vs *+>uas-Kir2.1* and *Rh1-Gal4>+* vs *+>uas-Kir2.1*; effect size 0.62/0.74, Table 6). Kir2.1

expressing flies had no changes in wake activity (Figure 5C and 5F). Taken together, our data

indicates that phototransduction and the neuronal excitability in the *Drosophila* Rh1-expressing

photoreceptor promote day sleep.

4. Discussion

In this study, we surveyed ten mutants with either hyperpolarised photoreceptors or defects in phototransduction, and four mutants with disrupting histaminergic transmission, the main neurotransmitter used by *Drosophila photoreceptors*. We observed short- or fragmented-day sleep in majority of the phototransduction mutants as well as in mutants with hyperpolarised R1-R6 photoreceptors. This data indicates that R1-R6 photoreceptor neural activity is required for sleep homeostatic drive during daytime. In contrast, we found that reduced histamine production (*hdc* mutants) result in night time sleep loss, while reduced histaminergic reception by ort mutation cause short day sleep. Divergent sleep phenotypes between *hdc* and *ort* mutants may reflect a potential demarcation of day and night sleep neural network downstream of photoreceptors (see below). Moreover, *hdc* and *ort* expressions are detected outside photoreceptor or visual system, so we cannot rule out the sleep defect we observed is exclusively derived from visual system. Importantly, the mutants tested in this study don't loss their ability to sense light, as the non-visual pathway such as *cry*, *Rh7* and *qsm* remain intake. Nevertheless, we showed that reduced visual phototransduction or downstream neurotransmission cause sleep loss and/or fragmentation, indicating a clear role of *Drosophila* photoreceptor R1-R6 in sleep promoting.

Our finding is consistent with previous publication that a visually impaired *omb* mutant showed major sleep loss in both day and night (Kirszenblat et al., 2019) as well as the finding that activation of L2 LMC neurons cause sleep lost (Muraro and Ceriani, 2015).

However our data is in contrary to the finding that blocking neural transmission for R1-R6 via TeTxLC caused day sleep increase (Damulewicz et al., 2020). As mentioned above, we cannot produce viable adult flies that express TeTxLC in photoreceptors and instead we apply Kir2 channel to hyperpolarise photoreceptor. There may be two factors relate to these data discrepancy between us and Damulewicz et al (2020): Firstly, our Gal4 and UAS lines are outcrossed to the w^{iso31C} background, as a standard in the sleep research field, while such outcrossing were not described in Damulewicz et al. Secondly, vision is important for navigation, and the compromised vision in phototransduction mutants and flies with weaken photoreceptor outputs (i.e., *gmr-Gal4>UAS-kir2* or *gmr-Gal4>UAS-TeTxLC*) can result in reduced wake locomotor activities, which is a major compounding factor in assessing sleep increase. In fact, nine visual mutants in our survey (*ninaE^5*, *ninaE^8*, *norpA^{p24}*, *trp^1*, *trpl^{302}*, *hdc^{JK910}*, *hdc^{MB}*, *ort^{attp}* and *gmr-Gal4>uas-kir2*) showed significant reduced wake activity in either day and/or night. Consequentially, the reduced wake activity prevents us to conclude whether the increased night sleep seen in *ninaE^8*, *trpl^{302}* and *ort^{attp}* are genuine increase or a reflection of loss of locomotion during wakefulness. The wake activity was not analysed by Damulewicz et al (2020)

preventing a direct comparison between the two studies. More surprisingly, the sleep loss identified

in *hdc* and *ort* mutants is also opposite to previous finding indicating that histamine as pro-waking

factor (Oh et al., 2013). One major difference to Oh et al (2013) is that we only monitored male

sleeps in order to reach large range of sleep dynamics (see introduction). Importantly female flies

have very low levels of day sleep. Further investigation for female visual mutants will be required in

future to clarify if the role of visual input in sleep diverge between males and females. Nevertheless,

as shown in our data, three of the four mutants with histaminergic disruption show lower wake

activity, consistent with the arousal promoting role of histamine proposed by Oh et al (2013), yet

similar to Damulewicz et al (2020) Oh et al did not assess the impact of wake activity on sleep. Wake

activity is a classic but somehow ignored analytical tool in the *Drosophila* sleep research. We suggest

that this parameter should always be included for any locomotion-based sleep assay.

While uas-Kir2 has been used for previous sleep investigation, a *+>uas-Kir2* control were not

included (Shang et al., 2008). As mentioned, we observed an unexpected abnormal low night sleep

in *+>uas-Kir2* control flies which is likely due to leaking expression of Kir2 at other part of fly brain,

since the *gmr-Gal*4 and *UAS-Kir2* lines we used are isogenised to w^{iso31C} background. Nevertheless,

this limits our investigation to day sleep. Future investigation using gmr-LexA and LexAoP-Kir2 may

be required to verify this finding.

Unlike the isogenised Gal4 and UAS transgenic lines, the visual mutants come with various genetic

background including unknown *white* backgrounds and two wild type backgrounds. Due to the

complex mutagenic nature for these lines (point mutations or insertion) we were not able to bring all

these lines into common background. We therefore included in most cases at least two alleles and

most close background controls for sleep comparison, and we only interpreted the consistent

phenotype that presented across the alleles. Future investigation utilised the isogenised somatic

CRISPR knockout constructs will be required to rule out impact of background in this study.

As part of control experiments, we examined the ERG profiles for all the tested gene-specific

mutants. However, the severity of sleep loss phenotype does not directly correlate with ERG defects:

weakest *ninaE*[5] mutant (only on transient defected is detected) showed just as much day sleep

fragmentation as *ninaE*[8], *ninaE*[17] or *norpA*[p24]. Moreover, *trp*[1] has more dramatic defect in ERG as

compared to *trp*[MB]; while *trp*[1] showed reduce sleep in both day and night, *trp*[MB] only showed night

sleep loss. Intriguingly both *trpl*[302] and *trpl*[MB] exhibit sleep loss, despite that our ERG recordings

showing minor or no impact of *trpl* mutants on ERG profile, in agreement with previous studies. As

mentioned above these divergent sleep phenotypes may be partly due to the various genetic

background we did not control as well as the expression profile outside photoreceptors. When

combining with Gal4-UAS investigation (Figure 5), it is nevertheless clear that R1-R6 photoreceptor is

required for driving day sleep. Vision and light-affected sleep are previously shown to be mediated

both via circadian clock neurons (eg., LNvs and DN1) as well as ring neurons in EB (Lamaze et al.,

2018; Raccuglia et al., 2019; Shang et al., 2008; Singh et al., 2023). Various neural pathways have

been shown to connect photoreceptors to the wake promoting LNv: R1-R6 photoreceptor can

propagate the light signal via histamine-mediated inhibition of the downstream LMC neurons that

activates LNvs (Muraro and Ceriani, 2015); on the other hand, R8 has recently been shown to

communicate with LNv via the so-called AMA neurons and cholinergic signalling. Moreover, sleep

promoting signal from photoreceptors may also reach EB via Anterior Visual Pathway through MeTu

and TuBu neurons, upon which circadian clock neuron DN1 also converge (Guo et al., 2018; Lamaze

et al., 2018). It is currently unclear which of the above pathways are required for the photoreceptor

mediated day sleep promotion. Although the day sleep loss observed in ort^1, $ninaE$ mutants and $Rh1$-

$Gal4$>uas-$Kir2$ indicates that the day sleep depends on Rh1-positive R1-R6 histaminergic inhibition

on LMC neurons, we did not observe consistent day sleep loss among hdc mutants. Nevertheless,

hdc expression is wider than visual system(Oh et al., 2013). Moreover, loss of hdc activity does not

completely remove histamine production (eg., Stenesen et al., 2015) and is not equal to selective

blockage of ort-mediated histamine synaptic transmission. Five tested mutants also exhibit night

sleep loss (two hdc mutants, $norpA^{p24}$, trp^1 and trp^{MB}). One common feature of these five mutants is

that they may disrupt R7/R8 photoreceptors function in addition to R1-R6 photoreceptors.

Rhodopsin 3-expressing R7 was previously shown to selectively control night sleep (Damulewicz et

al., 2020). R8 was shown to synapse onto L1 LMC neurons and downstream motion sensing network

(Davis et al., 2020b). Consistently, a previous study suggest that a homeostatic night sleep drive can

be imposed by imbalanced activation of downstream targets in the LMC mediated motion sensing

network (Kirszenblat et al., 2019). Due to the low night sleep observed in +>uas-$Kir2$ flies, we cannot

assess the effect of photoreceptor inhibition in night sleep. It is possible the night sleep loss

observed in the five visual mutants are due to similar imbalance output from photoreceptor subsets.

In summary, our data indicates a clear sleep promoting role by $Drosophila$ photoreceptor. The

revelation of a potential neural network demarcation of day and night sleep require further

investigation beyond this report.

Acknowledgements

This study was funded by University of Leicester to M.A. and X.L. (F50/100 PhD studentships), N. S., S. H., T. O., and J. A (PGT and UG programme funding), an Erasmus + studentship to C. S.-G., a Royal Society University Fellowship to B. W., and a BBSRC New Investigator Scheme (BB/W014939/1) to K-F.C and Y-C. H.

Author Contributions

K-F.C – Conceptualization, Methodology, Investigation, Writing - Original Draft, Writing – Review & Editing, Visualization, Supervision, Project administration, Funding acquisition; Y.-C. H. – Investigation, Visualization, Writing – Review & Editing. B.W. – Methodology, Investigation, Writing – Review & Editing. M.A., X.L., N. S., S. H., T. O., J. A, and C. S.-G., – Investigation, Writing – Review & Editing.

Declaration Of Interests

The authors declare no competing interests.

 Figure captions

Figure 1. sleep profile, parameters and ERG quantification for *ninaE* mutants. A. Sleep traces of

average sleep minutes per 30 minutes over 12 hours light-dark cycle. ZT0: light on, ZT12: light off,

ZT24: end of night. Sleep traces of *iso* control and *ninaE* mutants are colour coded in the shades of

greys and reds. **B.** Average day sleep bout length (minutes), **C.** Average day wake activity (beam

breaks per waking minutes), **D.** Average day sleep (minutes), **E.** Average night sleep bout length

(minutes), **F.** Average night wake activity (beam breaks per waking minutes), and **G.** Average night

sleep (minutes) for *ninaE^{17}* (i), *ninaE^{8}* (ii) and *ninaE^{5}* (iii) compared to iso controls. Values for each

individual are plotted as filled circles except for those outside (B-G). **H.** An example ERG profile from

iso control with 1000 ms green light flash that evokes voltage changes of on-transient, receptor

potential and off-transient. **I.** Mean and standard deviation for voltage change (mV) in on transient,

receptor potential and off-transient of ERG upon four light intensity (0.004, 0.02, 0.1 and 0.5 w/m^2)

flash for *ninaE^{17}* and controls. **J.** Mean and standard deviation for voltage change (mV) in on

transient, receptor potential and off-transient of ERG upon four light intensity (0.004, 0.02, 0.1 and

0.5 w/m^2) 1000ms flash for *ninaE^{5}*, *ninaE^{8}* and controls. Plot scales are in either liner (A-G) and

logarithmic (I-J). Asterisk indicates statistical significance (p<0.05) between mutants and iso controls

by pairwise Wilcoxon rank sum exact test without (B-H) and with multiple comparison, Benjamini-

Hochberg correction (I, followed by Kruskal-Wallis rank sum test). See **Table 1** for sample size, p-

values and effect size for each comparison.

Figure 2. sleep profile, parameters and ERG quantification for *norpA* mutants. A. Sleep traces of

average sleep minutes per 30 minutes over 12 hours light-dark cycle. ZT0: light on, ZT12: light off,

ZT24: end of night. Sleep traces for *iso* control and *norpA^{p24}* mutants are colour coded in grey and

cyan. **B.** Average day sleep bout length (minutes), **C.** Average day wake activity (beam breaks per

waking minutes), **E.** Average night sleep bout length (minutes), **F.** Average night wake activity (beam

breaks per waking minutes), **G.** Average night sleep (minutes) for *norpA^{p24}* compared to iso controls.

Values for each individual are plotted as filled circles except for those outside (B-G). **H.** mean and

standard deviation for voltage change (mV) in on transient, receptor potential and off-transient of

ERG upon four light intensity (0.004, 0.02, 0.1 and 0.5 w/m^2) flash for *norpA^{p24}* and *iso* controls. Plot

scales are in either liner (A-G) and logarithmic (H). Asterisk indicates statistical significance (p<0.05)

between mutants and iso controls by pairwise Wilcoxon rank sum exact test. See **Table 2** for sample

size, p-values and effect size for each comparison.

Figure 3. sleep profile, parameters and ERG quantification for *trp* and *trpl* mutants. A. Sleep traces

of average sleep minutes per 30 minutes over 12 hours light-dark cycle. ZT0: light on, ZT12: light off,

ZT24: end of night. Sleep traces for *Canton S* (i), *Oregon R* (i) and *iso* controls (ii-iv) are coloured

coded in the shades of greys; *trp* (i-ii) and *trpl* (iii-iv) mutants are colour coded in shades of blue and

purples. **B.** Average day sleep bout length (minutes), **C.** Average day wake activity (beam breaks per

waking minutes), **D.** Average day sleep (minutes), **E.** Average night sleep bout length (minutes), **F.**

Average night wake activity (beam breaks per waking minutes), **G.** Average night sleep (minutes) for

trp^1 (i), *trp^{MB}* (ii), *trpl^{302}* (iii) and *trpl$^{302/MB}$* or *trpl^{MB}* (iv) compared to controls. Values for each

individual are plotted as filled circles except for those outside (B-G). **H.** Mean and standard deviation

for voltage change (mV) in on transient, receptor potential and off-transient of ERG upon four light

intensity (0.004, 0.02, 0.1 and 0.5 w/m^2) flash for *trp^1* (i), *trp^{MB}* (ii), *trpl^{302}* (iii) and *trpl^{MB}* (iii)

compared to controls. Plot scales are in either liner (A-G) and logarithmic (H). Asterisk indicates

statistical significance (p<0.05) between mutants and controls by pairwise Wilcoxon rank sum exact

test without (Aii-Gii, Aiii-Giii, Hi and Hii) and with multiple comparison, Benjamini-Hochberg

correction (Ai-Gi, Aiv-Giv, and Hiii followed by Kruskal-Wallis rank sum test). See **Table 3 and Table 4**

for sample size, p-values and effect size for each comparison.

Figure 4. sleep profile, parameters and ERG quantification for *hdc* and *ort* mutants. A. Sleep traces

of average sleep minutes per 30 minutes over 12 hours light-dark cycle. ZT0: light on, ZT12: light off,

ZT24: end of night. Sleep traces for *Canton S* (i) and *iso* controls (ii-iii) are coloured coded in the

shades of greys; *hdc* (i) and *ort* (ii-iii) mutants are colour coded in shades of indigos and greens. **B.**

Average day sleep bout length (minutes), **C.** Average day wake activity (beam breaks per waking

minutes), **D.** Average day sleep (minutes), **E.** Average night sleep bout length (minutes), **F.** Average

night wake activity (beam breaks per waking minutes), **G.** Average night sleep (minutes) for *hdc*

mutants (i), *ort^1* (ii) and *ort$^{attp/+}$* or *ort^{attp}* (iii) compared to controls. Values for each individual are

plotted as filled circles except for those outside (B-G). **H.** Mean and standard deviation for voltage

change (mV) in on transient, receptor potential and off-transient of ERG upon four light intensity

(0.004, 0.02, 0.1 and 0.5 w/m^2) flash for *hdc* (i) and *ort* (ii) mutants compared to controls. Plot scales

are in either liner (A-G) and logarithmic (H). Asterisk indicates statistical significance (p<0.05)

between mutants and controls by pairwise Wilcoxon rank sum exact test without (Aii-Gii) and with

multiple comparison, Benjamini-Hochberg correction (Ai-Hi, Aiii-Giii, and Hii followed by Kruskal-

Wallis rank sum test). See **Table 5** for sample size, p-values and effect size for each comparison.

Figure 5. sleep profile and parameters for flies with hyperpolarised photoreceptors. A. Sleep traces

of average sleep minutes per 30 minutes over 12 hours light-dark cycle. ZT0: light on, ZT12: light off,

ZT24: end of night. Sleep traces for *gmr-Gal4>+* (i), *Rh1-Gal4>+* (ii) and *+>UAS-Kir2* (i-ii) control flies

are coloured coded in the shades of greys; experimental mutants *gmr-Gal4> UAS-kir2* (i) and *Rh1-*

Gal4>UAS-kir2 (ii) are colour coded in light burgundy. **B.** Average day sleep bout length (minutes), **C.**

Average day wake activity (beam breaks per waking minutes), **D.** Average day sleep (minutes), **E.**

Average night sleep bout length (minutes), **F.** Average night wake activity (beam breaks per waking

minutes), **G.** Average night sleep (minutes) for *gmr-Gal4> UAS-kir2* (i) and *Rh1-Gal4>UAS-kir2* (ii)

compared to controls. Values for each individual are plotted as filled circles except for those outside

(B-G). Asterisk indicates statistical significance (p<0.05) between mutants and controls by pairwise

Wilcoxon rank sum exact test with multiple comparison, Benjamini-Hochberg correction (followed

by Kruskal-Wallis rank sum test). See **Table 6** for sample size, p-values and effect size for each

comparison.

Reference

Alejevski, F., Saint-Charles, A., Michard-Vanhee, C., Martin, B., Galant, S., Vasiliauskas, D., Rouyer, F.,
2019. The HisCl1 histamine receptor acts in photoreceptors to synchronize Drosophila
behavioral rhythms with light-dark cycles. Nature communications 10, 252.
https://doi.org/10.1038/s41467-018-08116-7

Alphen, B. van, Semenza, E.R., Yap, M., Swinderen, B. van, Allada, R., 2021. A deep sleep stage in
Drosophila with a functional role in waste clearance. Science Advances 7, eabc2999.
https://doi.org/10.1126/sciadv.abc2999

Altimus, C.M., Güler, A.D., Villa, K.L., McNeill, D.S., LeGates, T.A., Hattar, S., 2008. Rods-cones and
melanopsin detect light and dark to modulate sleep independent of image formation.
Proceedings of the National Academy of Sciences 105, 19998–20003.
https://doi.org/10.1073/pnas.0808312105

Andretic, R., Shaw, P.J., 2005. Essentials of Sleep Recordings in Drosophila: Moving Beyond Sleep
Time, in: Young, M.W. (Ed.), Methods in Enzymology, Circadian Rhythms. Academic Press,
pp. 759–772. https://doi.org/10.1016/S0076-6879(05)93040-1

Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T., Bate, M., 2001. Altered Electrical Properties in
DrosophilaNeurons Developing without Synaptic Transmission. J. Neurosci. 21, 1523–1531.
https://doi.org/10.1523/JNEUROSCI.21-05-01523.2001

Burg, M.G., Sarthy, P.V., Koliantz, G., Pak, W.L., 1993. Genetic and molecular identification of a
Drosophila histidine decarboxylase gene required in photoreceptor transmitter synthesis.
The EMBO Journal 12, 911–919. https://doi.org/10.1002/j.1460-2075.1993.tb05732.x

Chen, K.F., Lowe, S., Lamaze, A., Kratschmer, P., Jepson, J., 2019. Neurocalcin regulates nighttime
sleep and arousal in Drosophila. eLife 8. https://doi.org/10.7554/eLife.38114

Chen, K.F., Peschel, N., Zavodska, R., Sehadova, H., Stanewsky, R., 2011. QUASIMODO, a Novel GPI-
anchored zona pellucida protein involved in light input to the Drosophila circadian clock.
Current Biology 21, 719–729.

Chen, Z., Chen, H.-C., Montell, C., 2015. TRP and Rhodopsin Transport Depends on Dual XPORT ER
Chaperones Encoded by an Operon. Cell Reports 13, 573–584.
https://doi.org/10.1016/j.celrep.2015.09.018

Chiu, J.C., Low, K.H., Pike, D.H., Yildirim, E., Edery, I., 2010. Assaying Locomotor Activity to Study
Circadian Rhythms and Sleep Parameters in Drosophila. JoVE (Journal of Visualized
Experiments) e2157. https://doi.org/10.3791/2157

Cosens, D.J., Manning, A., 1969. Abnormal Electroretinogram from a Drosophila Mutant. Nature 224,
285–287. https://doi.org/10.1038/224285a0

Damulewicz, M., Ispizua, J.I., Ceriani, M.F., Pyza, E.M., 2020. Communication Among Photoreceptors

and the Central Clock Affects Sleep Profile. Front. Physiol. 11.

https://doi.org/10.3389/fphys.2020.00993

Davis, F.P., Nern, A., Picard, S., Reiser, M.B., Rubin, G.M., Eddy, S.R., Henry, G.L., 2020a. A genetic,

genomic, and computational resource for exploring neural circuit function. eLife 9, e50901.

https://doi.org/10.7554/eLife.50901

Davis, F.P., Nern, A., Picard, S., Reiser, M.B., Rubin, G.M., Eddy, S.R., Henry, G.L., 2020b. A genetic,

genomic, and computational resource for exploring neural circuit function. eLife 9, e50901.

https://doi.org/10.7554/eLife.50901

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang, W., Huang, J., Rao, Y.,

2019. Chemoconnectomics: Mapping Chemical Transmission in Drosophila. Neuron 101,

876-893.e4. https://doi.org/10.1016/j.neuron.2019.01.045

Dissel, S., Angadi, V., Kirszenblat, L., Suzuki, Y., Donlea, J., Klose, M., Koch, Z., English, D., Winsky-

Sommerer, R., van Swinderen, B., Shaw, P.J., 2015. Sleep Restores Behavioral Plasticity to

Drosophila Mutants. Current Biology 25, 1270–1281.

https://doi.org/10.1016/j.cub.2015.03.027

Dove, A.E., Cook, B.L., Irgebay, Z., Vecsey, C.G., 2017. Mechanisms of sleep plasticity due to sexual

experience in Drosophila melanogaster. Physiology & Behavior 180, 146–158.

https://doi.org/10.1016/j.physbeh.2017.08.020

Duhart, J.M., Buchler, J.R., Inami, S., Kennedy, K.J., Jenny, B.P., Afonso, D.J.S., Koh, K., 2023.

Modulation and neural correlates of postmating sleep plasticity in Drosophila females.

Current Biology 33, 2702-2716.e3. https://doi.org/10.1016/j.cub.2023.05.054

Gao, S., Takemura, S., Ting, C.-Y., Huang, S., Lu, Z., Luan, H., Rister, J., Thum, A.S., Yang, M., Hong, S.-

T., Wang, J.W., Odenwald, W.F., White, B.H., Meinertzhagen, I.A., Lee, C.-H., 2008. The

Neural Substrate of Spectral Preference in Drosophila. Neuron 60, 328–342.

https://doi.org/10.1016/j.neuron.2008.08.010

Gilestro, G.F., Tononi, G., Cirelli, C., 2009. Widespread Changes in Synaptic Markers as a Function of

Sleep and Wakefulness in Drosophila. Science 324, 109–112.

https://doi.org/10.1126/science.1166673

Guo, F., Holla, M., Díaz, M.M., Rosbash, M., 2018. A Circadian Output Circuit Controls Sleep-Wake

Arousal in Drosophila. Neuron 100, 624-635.e4.

https://doi.org/10.1016/j.neuron.2018.09.002

Harbison, S.T., Sehgal, A., 2008. Quantitative Genetic Analysis of Sleep in Drosophila melanogaster.

Genetics 178, 2341–2360. https://doi.org/10.1534/genetics.107.081232

Hardie, R.C., 2012. Phototransduction mechanisms in Drosophila microvillar photoreceptors. Wiley
Interdisciplinary Reviews: Membrane Transport and Signaling 1, 162–187.
https://doi.org/10.1002/wmts.20

Hartmann, C., Kempf, A., 2023. Mitochondrial control of sleep. Current Opinion in Neurobiology 81,
102733. https://doi.org/10.1016/j.conb.2023.102733

Horne, J.A., Walmsley, B., 1976. Daytime visual load and the effects upon human sleep.
Psychophysiology 13, 115–20.

Hsu, C.T., Guevara, C., Killiany, S.L., Shon, J., Dissel, S., Sehgal, A., 2025. Clock-dependent regulation
of a homeostatic sleep center maintains daytime sleep and evening activity. Current Biology
0. https://doi.org/10.1016/j.cub.2025.06.003

Inoue, H., Yoshioka, T., Hotta, Y., 1989. Diacylglycerol Kinase Defect in a *Drosophila* Retinal
Degeneration Mutant *rdgA*. Journal of Biological Chemistry 264, 5996–6000.
https://doi.org/10.1016/S0021-9258(18)83648-1

Isaac, R.E., Li, C., Leedale, A.E., Shirras, A.D., 2009. Drosophila male sex peptide inhibits siesta sleep
and promotes locomotor activity in the post-mated female. Proceedings of the Royal Society
B: Biological Sciences 277, 65–70. https://doi.org/10.1098/rspb.2009.1236

Ishimoto, H., Lark, A.R.S., Kitamoto, T., 2012. Factors that Differentially Affect Daytime and
Nighttime Sleep in Drosophila melanogaster. Front. Neurol. 3.
https://doi.org/10.3389/fneur.2012.00024

Jagannathan, S.R., Jeans, T., Van De Poll, M.N., van Swinderen, B., 2024. Multivariate classification of
multichannel long-term electrophysiology data identifies different sleep stages in fruit flies.
Science Advances 10, eadj4399. https://doi.org/10.1126/sciadv.adj4399

Jones, J.D., Holder, B.L., Montgomery, A.C., McAdams, C.V., He, E., Burns, A.E., Eiken, K.R., Vogt, A.,
Velarde, A.I., Elder, A.J., McEllin, J.A., Dissel, S., 2025. The dorsal fan-shaped body is a
neurochemically heterogeneous sleep-regulating center in Drosophila. PLOS Biology 23,
e3003014. https://doi.org/10.1371/journal.pbio.3003014

Kirszenblat, L., Yaun, R., van Swinderen, B., 2019. Visual experience drives sleep need in Drosophila.
Sleep 42. https://doi.org/10.1093/sleep/zsz102

Korf, E.M., Mölle, M., Born, J., Ngo, H.-V.V., 2017. Blindfolding during wakefulness causes decrease
in sleep slow wave activity. Physiological Reports 5, e13239.
https://doi.org/10.14814/phy2.13239

Lakhiani, R., Shanavas, S., Melnattur, K., 2023. Comparative biology of sleep in diverse animals. J Exp
Biol 226. https://doi.org/10.1242/jeb.245677

Lamaze, A., Krätschmer, P., Chen, K.-F., Lowe, S., Jepson, J.E.C., 2018. A Wake-Promoting Circadian
599   Output Circuit in Drosophila. Current Biology 28, 3098-3105.e3.
600   https://doi.org/10.1016/j.cub.2018.07.024

601   Leung, H.-T., Geng, C., Pak, W.L., 2000. Phenotypes of trpl Mutants and Interactions between the
602   Transient Receptor Potential (TRP) and TRP-Like Channels inDrosophila. J. Neurosci. 20,
603   6797–6803. https://doi.org/10.1523/JNEUROSCI.20-18-06797.2000

604   Mazzotta, G.M., Damulewicz, M., Cusumano, P., 2020. Better Sleep at Night: How Light Influences
605   Sleep in Drosophila. Front. Physiol. 11. https://doi.org/10.3389/fphys.2020.00997

606   Meyerhof, G.T., Easwaran, S., Bontempo, A.E., Montell, C., Montell, D.J., 2024. Altered circadian
607   rhythm, sleep, and *rhodopsin 7*–dependent shade preference during diapause in *Drosophila*
608   *melanogaster*. Proceedings of the National Academy of Sciences 121, e2400964121.
609   https://doi.org/10.1073/pnas.2400964121

610   Miyamoto, H., Katagiri, H., Hensch, T., 2003. Experience-dependent slow-wave sleep development.
611   Nat Neurosci 6, 553–554. https://doi.org/10.1038/nn1064

612   Muraro, N.I., Ceriani, M.F., 2015. Acetylcholine from visual circuits modulates the activity of arousal
613   neurons in Drosophila. Journal of Neuroscience 35, 16315–16327.

614   Nériec, N., Desplan, C., 2016. Chapter Fourteen - From the Eye to the Brain: Development of the
615   Drosophila Visual System, in: Wassarman, P.M. (Ed.), Current Topics in Developmental
616   Biology, Essays on Developmental Biology, Part A. Academic Press, pp. 247–271.
617   https://doi.org/10.1016/bs.ctdb.2015.11.032

618   Ni, J.D., Baik, L.S., Holmes, T.C., Montell, C., 2017. A rhodopsin in the brain functions in circadian
619   photoentrainment in Drosophila. Nature 545, 340–344.
620   https://doi.org/10.1038/nature22325

621   Niemeyer, B.A., Suzuki, E., Scott, K., Jalink, K., Zuker, C.S., 1996. The Drosophila Light-Activated
622   Conductance Is Composed of the Two Channels TRP and TRPL. Cell 85, 651–659.
623   https://doi.org/10.1016/S0092-8674(00)81232-5

624   Nitabach, M.N., Blau, J., Holmes, T.C., 2002. Electrical Silencing of Drosophila Pacemaker Neurons
625   Stops the Free-Running Circadian Clock. Cell 109, 485–495. https://doi.org/10.1016/S0092-
626   8674(02)00737-7

627   Ogueta, M., Hardie, R.C., Stanewsky, R., 2020. Light Sampling via Throttled Visual Phototransduction
628   Robustly Synchronizes the Drosophila Circadian Clock. Current Biology 30, 2551-2563.e3.
629   https://doi.org/10.1016/j.cub.2020.04.067

Ogueta, M., Hardie, R.C., Stanewsky, R., 2018. Non-canonical Phototransduction Mediates
Synchronization of the Drosophila melanogaster Circadian Clock and Retinal Light
Responses. Current Biology 28, 1725-1735.e3. https://doi.org/10.1016/j.cub.2018.04.016

Oh, Y., Jang, D., Sonn, J.Y., Choe, J., 2013. Histamine-HisCl1 Receptor Axis Regulates Wake-Promoting
Signals in Drosophila melanogaster. PLoS ONE 8, e68269.
https://doi.org/10.1371/journal.pone.0068269

O'Tousa, J.E., Baehr, W., Martin, R.L., Hirsh, J., Pak, W.L., Applebury, M.L., 1985. The Drosophila
ninaE gene encodes an opsin. Cell 40, 839–850. https://doi.org/10.1016/0092-
8674(85)90343-5

Pantazis, A., Segaran, A., Liu, C.-H., Nikolaev, A., Rister, J., Thum, A.S., Roeder, T., Semenov, E.,
Juusola, M., Hardie, R.C., 2008a. Distinct Roles for Two Histamine Receptors (hclA and hclB)
at the Drosophila Photoreceptor Synapse. J. Neurosci. 28, 7250–7259.
https://doi.org/10.1523/JNEUROSCI.1654-08.2008

Pantazis, A., Segaran, A., Liu, C.-H., Nikolaev, A., Rister, J., Thum, A.S., Roeder, T., Semenov, E.,
Juusola, M., Hardie, R.C., 2008b. Distinct Roles for Two Histamine Receptors (hclA and hclB)
at the Drosophila Photoreceptor Synapse. The Journal of Neuroscience 28, 7250–7259.
https://doi.org/10.1523/jneurosci.1654-08.2008

Peschel, N., Chen, K.F., Szabo, G., Stanewsky, R., 2009. Light-dependent interactions between the
Drosophila circadian clock factors cryptochrome, jetlag, and timeless. Current Biology 19,
241–247.

Pilorz, V., Tam, S.K.E., Hughes, S., Pothecary, C.A., Jagannath, A., Hankins, M.W., Bannerman, D.M.,
Lightman, S.L., Vyazovskiy, V.V., Nolan, P.M., Foster, R.G., Peirson, S.N., 2016. Melanopsin
Regulates Both Sleep-Promoting and Arousal-Promoting Responses to Light. PLOS Biology
14, e1002482. https://doi.org/10.1371/journal.pbio.1002482

Raccuglia, D., Huang, S., Ender, A., Heim, M.M., Laber, D., Suárez-Grimalt, R., Liotta, A., Sigrist, S.J.,
Geiger, J.R.P., Owald, D., 2019. Network-Specific Synchronization of Electrical Slow-Wave
Oscillations Regulates Sleep Drive in Drosophila. Current Biology.
https://doi.org/10.1016/j.cub.2019.08.070

Reinhard, N., Fukuda, A., Manoli, G., Derksen, E., Saito, A., Möller, G., Sekiguchi, M., Rieger, D.,
Helfrich-Förster, C., Yoshii, T., Zandawala, M., 2024. Synaptic connectome of the Drosophila
circadian clock. Nat Commun 15, 10392. https://doi.org/10.1038/s41467-024-54694-0

Rezával, C., Werbajh, S., Ceriani, M.F., 2007. Neuronal death in Drosophila triggered by GAL4
accumulation. European Journal of Neuroscience 25, 683–694.
https://doi.org/10.1111/j.1460-9568.2007.05317.x

Rieger, D., Stanewsky, R., Helfrich-Förster, C., 2003. Cryptochrome, Compound Eyes, Hofbauer-

665   Buchner Eyelets, and Ocelli Play Different Roles in the Entrainment and Masking Pathway of

666   the Locomotor Activity Rhythm in the Fruit Fly Drosophila Melanogaster. J Biol Rhythms 18,

667   377–391. https://doi.org/10.1177/0748730403256997

668   Rossi, F.A., Bochicchio, P.A., Quesada-Allué, L.A., Pérez, M.M., 2015. N-β-alanyldopamine

669   metabolism, locomotor activity and sleep in Drosophila melanogaster ebony and tan

670   mutants. Physiological Entomology 40, 166–174. https://doi.org/10.1111/phen.12100

671   Scavarda, N.J., O'Tousa, J., Pak, W.L., 1983. Drosophila locus with gene-dosage effects on rhodopsin.

672   Proceedings of the National Academy of Sciences 80, 4441–4445.

673   https://doi.org/10.1073/pnas.80.14.4441

674   Schlichting, M., Menegazzi, P., Lelito, K.R., Yao, Z., Buhl, E., Benetta, E.D., Bahle, A., Denike, J., Hodge,

675   J.J., Helfrich-Förster, C., Shafer, O.T., 2016. A Neural Network Underlying Circadian

676   Entrainment and Photoperiodic Adjustment of Sleep and Activity in Drosophila. J. Neurosci.

677   36, 9084–9096. https://doi.org/10.1523/JNEUROSCI.0992-16.2016

678   Sengupta, S., Crowe, L.B., You, S., Roberts, M.A., Jackson, F.R., 2019. A Secreted Ig-Domain Protein

679   Required in Both Astrocytes and Neurons for Regulation of Drosophila Night Sleep. Current

680   Biology 29, 2547-2554.e2. https://doi.org/10.1016/j.cub.2019.06.055

681   Shafer, O.T., Keene, A.C., 2021. The Regulation of Drosophila Sleep. Current Biology 31, R38–R49.

682   https://doi.org/10.1016/j.cub.2020.10.082

683   Shang, Y., Griffith, L.C., Rosbash, M., 2008. Light-arousal and circadian photoreception circuits

684   intersect at the large PDF cells of the Drosophila brain. Proceedings of the National Academy

685   of Sciences 105, 19587–19594. https://doi.org/10.1073/pnas.0809577105

686   Singh, P., Aleman, A., Omoto, J.J., Nguyen, B.-C., Kandimalla, P., Hartenstein, V., Donlea, J.M., 2023.

687   Examining Sleep Modulation by Drosophila Ellipsoid Body Neurons. eNeuro 10.

688   https://doi.org/10.1523/ENEURO.0281-23.2023

689   Stenesen, D., Moehlman, A.T., Krämer, H., 2015. The carcinine transporter CarT is required in

690   Drosophila photoreceptor neurons to sustain histamine recycling. eLife 4, e10972.

691   https://doi.org/10.7554/eLife.10972

692   Suh, J., Jackson, F.R., 2007. Drosophila Ebony Activity Is Required in Glia for the Circadian Regulation

693   of Locomotor Activity. Neuron 55, 435–447. https://doi.org/10.1016/j.neuron.2007.06.038

694   Sweeney, S.T., Broadie, K., Keane, J., Niemann, H., O'Kane, C.J., 1995. Targeted expression of tetanus

695   toxin light chain in Drosophila specifically eliminates synaptic transmission and causes

696   behavioral defects. Neuron 14, 341–351. https://doi.org/10.1016/0896-6273(95)90290-2

Tainton-Heap, L.A.L., Kirszenblat, L.C., Notaras, E.T., Grabowska, M.J., Jeans, R., Feng, K., Shaw, P.J.,
van Swinderen, B., 2021. A Paradoxical Kind of Sleep in Drosophila melanogaster. Current
Biology 31, 578-590.e6. https://doi.org/10.1016/j.cub.2020.10.081

Tang, M., Cao, L.-H., Yang, T., Ma, S.-X., Jing, B.-Y., Xiao, N., Xu, S., Leng, K.-R., Yang, D., Li, M.-T., Luo,
D.-G., 2022. An extra-clock ultradian brain oscillator sustains circadian timekeeping. Science
Advances 8, eabo5506. https://doi.org/10.1126/sciadv.abo5506

Vaccaro, A., Dor, Y.K., Nambara, K., Pollina, E.A., Lin, C., Greenberg, M.E., Rogulja, D., 2020. Sleep
Loss Can Cause Death through Accumulation of Reactive Oxygen Species in the Gut. Cell 0.
https://doi.org/10.1016/j.cell.2020.04.049

Vincent, C.M., Beckwith, E.J., Silva, C.J.S. da, Pearson, W.H., Kierdorf, K., Gilestro, G.F., Dionne, M.S.,
2022. Infection increases activity via Toll dependent and independent mechanisms in
Drosophila melanogaster. PLOS Pathogens 18, e1010826.
https://doi.org/10.1371/journal.ppat.1010826

Wu, B., Ma, L., Zhang, E., Du, J., Liu, S., Price, J., Li, S., Zhao, Z., 2018. Sexual dimorphism of sleep
regulated by juvenile hormone signaling in Drosophila. PLOS Genetics 14, e1007318.
https://doi.org/10.1371/journal.pgen.1007318

Xiao, N., Xu, S., Li, Z.-K., Tang, M., Mao, R., Yang, T., Ma, S.-X., Wang, P.-H., Li, M.-T., Sunilkumar, A.,
Rouyer, F., Cao, L.-H., Luo, D.-G., 2023. A single photoreceptor splits perception and
entrainment by cotransmission. Nature 623, 562–570. https://doi.org/10.1038/s41586-023-
06681-6

Yap, M.H.W., Grabowska, M.J., Rohrscheib, C., Jeans, R., Troup, M., Paulk, A.C., van Alphen, B., Shaw,
P.J., van Swinderen, B., 2017. Oscillatory brain activity in spontaneous and induced sleep
stages in flies. Nat Commun 8, 1815. https://doi.org/10.1038/s41467-017-02024-y

Zanini, D., Giraldo, D., Warren, B., Katana, R., Andrés, M., Reddy, S., Pauls, S., Schwedhelm-Domeyer,
N., Geurten, B.R.H., Göpfert, M.C., 2018. Proprioceptive Opsin Functions in Drosophila Larval
Locomotion. Neuron 98, 67-74.e4. https://doi.org/10.1016/j.neuron.2018.02.028

Table 1. Statistics of a. sleep and b. ERG parameters for *ninaE* mutants

a. sleep parameter	genotype comparison	p-value	effect size r	n1	n2	significant r magnitude
day sleep	*iso* vs *ninaE*^{17}	3.50E-01	0.09	60	52	
night sleep	*iso* vs *ninaE*^{17}	1.00E-07	0.51	60	51	large
day sleep bout length	*iso* vs *ninaE*^{17}	9.60E-06	0.42	60	52	moderate
night sleep bout length	*iso* vs *ninaE*^{17}	7.90E-01	0.03	44	35	
day wake activity	*iso* vs *ninaE*^{17}	5.70E-01	0.05	60	52	
night wake activity	*iso* vs *ninaE*^{17}	1.30E-01	0.14	60	51	
day sleep	*iso* vs *ninaE*^{5}	1.40E-03	0.40	32	31	moderate
night sleep	*iso* vs *ninaE*^{5}	3.20E-02	0.27	32	31	small
day sleep bout length	*iso* vs *ninaE*^{5}	5.30E-09	0.68	32	31	large
night sleep bout length	*iso* vs *ninaE*^{5}	7.10E-02	0.23	32	31	
day wake activity	*iso* vs *ninaE*^{5}	3.90E-13	0.79	32	31	large
night wake activity	*iso* vs *ninaE*^{5}	1.90E-10	0.72	32	31	large
day sleep	*iso* vs *ninaE*^{8}	9.30E-05	0.51	32	26	large
night sleep	*iso* vs *ninaE*^{8}	1.10E-08	0.75	32	26	large
day sleep bout length	*iso* vs *ninaE*^{8}	6.90E-09	0.76	32	26	large
night sleep bout length	*iso* vs *ninaE*^{8}	9.60E-02	0.22	32	26	
day wake activity	*iso* vs *ninaE*^{8}	1.40E-13	0.82	32	26	large
night wake activity	*iso* vs *ninaE*^{8}	2.90E-07	0.67	32	26	large

b.ERG parameter	genotype comparison	p-value	effect size r	n1	n2	significant r magnitude
receptor potential	*iso* vs *ninaE*^{17}	2.50E-06	0.71	24	12	large
on transient	*iso* vs *ninaE*^{17}	1.90E-08	0.78	24	12	large
off transient	*iso* vs *ninaE*^{17}	1.60E-09	0.81	24	12	large
receptor potential	*iso* vs *ninaE*^{5}	4.40E-01	0.17	12	12	
receptor potential	*iso* vs *ninaE*^{8}	4.40E-06	0.83	12	12	large
receptor potential	*ninaE*^{5} vs *ninaE*^{8}	4.40E-06	0.84	12	12	large
on transient	*iso* vs *ninaE*^{5}	1.70E-03	0.64	12	12	large
on transient	*iso* vs *ninaE*^{8}	4.40E-06	0.84	12	12	large
on transient	*ninaE*^{5} vs *ninaE*^{8}	5.97E-02	0.39	12	12	
off transient	*iso* vs *ninaE*^{5}	1.70E-02	0.48	12	12	moderate
off transient	*iso* vs *ninaE*^{8}	1.10E-06	0.85	12	12	large
off transient	*ninaE*^{5} vs *ninaE*^{8}	1.10E-06	0.85	12	12	large

760   **Table 2. Statistics of a. sleep and b. ERG parameters for *norpA* mutants**

a. sleep parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
day sleep bout length	*iso* vs *norpA*[p24]	9.40E-03	0.39	23	21	moderate
night sleep bout length	*iso* vs *norpA*[p24]	2.50E-06	0.66	23	21	large
day sleep	*iso* vs *norpA*[p24]	3.20E-02	0.32	23	21	moderate
night sleep	*iso* vs *norpA*[p24]	1.20E-01	0.23	23	21	
day wake activity	*iso* vs *norpA*[p24]	9.40E-03	0.39	23	21	moderate
night wake activity	*iso* vs *norpA*[p24]	1.70E-07	0.72	23	21	large
b.ERG parameter	**genotype comparison**	**p-value**	**effect size *r***	**n1**	**n2**	**significant *r* magnitude**
receptor potential	*iso* vs *norpA*[p24]	1.50E-06	0.85	16	16	large
on transient	*iso* vs *norpA*[p24]	3.30E-09	0.85	16	16	large
off transient	*iso* vs *norpA*[p24]	1.50E-06	0.85	16	16	large

773 Table 3. Statistics of a. sleep and b. ERG parameters for *trp* mutants

a. sleep parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
day sleep	*Canton S* vs *Oregon R*	5.33E-01	0.07	38	53	
day sleep	*Canton S* vs trp^{1}	1.50E-04	0.47	38	30	moderate
day sleep	*Oregon R* vs trp^{1}	2.90E-06	0.54	53	30	large
night sleep	*Canton S* vs *Oregon R*	1.20E-04	0.41	38	53	moderate
night sleep	*Canton S* vs trp^{1}	1.63E-02	0.29	38	30	small
night sleep	*Oregon R* vs trp^{1}	2.30E-07	0.59	53	30	large
day sleep bout length	*Canton S* vs *Oregon R*	8.66E-01	0.02	38	53	
day sleep bout length	*Canton S* vs trp^{1}	1.40E-02	0.32	38	30	moderate
day sleep bout length	*Oregon R* vs trp^{1}	1.20E-02	0.32	53	30	moderate
night sleep bout length	*Canton S* vs *Oregon R*	6.61E-01	0.05	38	53	
night sleep bout length	*Canton S* vs trp^{1}	5.20E-03	0.35	38	30	moderate
night sleep bout length	*Oregon R* vs trp^{1}	1.20E-03	0.39	53	30	moderate
day wake activity	*Canton S* vs *Oregon R*	8.20E-08	0.58	38	53	large
day wake activity	*Canton S* vs trp^{1}	7.70E-01	0.04	38	30	
day wake activity	*Oregon R* vs trp^{1}	2.90E-07	0.57	53	30	large
night wake activity	*Canton S* vs *Oregon R*	1.20E-07	0.58	38	53	large
night wake activity	*Canton S* vs trp^{1}	1.60E-02	0.31	38	30	moderate
night wake activity	*Oregon R* vs trp^{1}	6.70E-02	0.20	53	30	
day sleep	*iso* vs trp^{MB}	5.80E-01	0.09	16	24	
night sleep	*iso* vs trp^{MB}	3.70E-03	0.46	16	24	moderate
day sleep bout length	*iso* vs trp^{MB}	3.20E-01	0.16	16	24	
night sleep bout length	*iso* vs trp^{MB}	3.90E-01	0.14	16	24	
day wake activity	*iso* vs trp^{MB}	1.40E-03	0.49	16	24	moderate
night wake activity	*iso* vs trp^{MB}	2.90E-01	0.17	16	24	
b. ERG parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
receptor potential	*iso* vs trp^{1}	7.40E-07	0.85	12	12	large
on transient	*iso* vs trp^{1}	2.20E-05	0.78	12	12	large
off transient	*iso* vs trp^{1}	7.40E-07	0.85	12	12	large
receptor potential	*iso* vs trp^{MB}	4.40E-03	0.53	12	16	large
on transient	*iso* vs trp^{MB}	2.60E-07	0.83	12	16	large
off transient	*iso* vs trp^{MB}	5.70E-01	0.11	12	16	

808 Table 4. Statistics of a. sleep and b. ERG parameters for *trpl* mutants

a. sleep parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
day sleep	*iso* vs *trpl*^{302}	1.40E-02	0.32	30	30	moderate
night sleep	*iso* vs *trpl*^{302}	1.40E-06	0.62	30	30	large
day sleep bout length	*iso* vs *trpl*^{302}	1.40E-06	0.59	30	30	large
night sleep bout length	*iso* vs *trpl*^{302}	6.90E-01	0.05	30	28	
day wake activity	*iso* vs *trpl*^{302}	1.10E-08	0.68	30	30	large
night wake activity	*iso* vs *trpl*^{302}	9.70E-04	0.43	30	28	moderate
day sleep	*iso/iso* vs *trpl*^{302}/*trpl*^{MB}	3.70E-05	0.63	23	24	large
day sleep	*iso/iso* vs *trpl*^{MB}/*trpl*^{MB}	3.71E-02	0.32	23	21	moderate
day sleep	*trpl*^{302}/*trpl*^{MB} vs *trpl*^{MB}/*trpl*^{MB}	2.57E-02	0.35	24	21	moderate
night sleep	*iso/iso* vs *trpl*^{302}/*trpl*^{MB}	7.32E-02	0.28	24	22	
night sleep	*iso/iso* vs *trpl*^{MB}/*trpl*^{MB}	1.56E-02	0.39	24	23	moderate
night sleep	*trpl*^{302}/*trpl*^{MB} vs *trpl*^{MB}/*trpl*^{MB}	1.52E-01	0.22	22	23	
day sleep bout length	*iso/iso* vs *trpl*^{302}/*trpl*^{MB}	1.60E-04	0.59	23	23	large
day sleep bout length	*iso/iso* vs *trpl*^{MB}/*trpl*^{MB}	1.60E-04	0.58	23	21	large
day sleep bout length	*trpl*^{302}/*trpl*^{MB} vs *trpl*^{MB}/*trpl*^{MB}	8.69E-01	0.03	23	21	
night sleep bout length	*iso/iso* vs *trpl*^{302}/*trpl*^{MB}	2.20E-01	0.23	23	22	
night sleep bout length	*iso/iso* vs *trpl*^{MB}/*trpl*^{MB}	8.42E-02	0.31	23	21	
night sleep bout length	*trpl*^{302}/*trpl*^{MB} vs *trpl*^{MB}/*trpl*^{MB}	4.82E-01	0.13	22	21	
day wake activity	*iso/iso* vs *trpl*^{302}/*trpl*^{MB}	1.20E-04	0.58	23	23	large
day wake activity	*iso/iso* vs *trpl*^{MB}/*trpl*^{MB}	9.63E-01	0.01	23	21	
day wake activity	*trpl*^{302}/*trpl*^{MB} vs *trpl*^{MB}/*trpl*^{MB}	1.04E-03	0.50	23	21	large
night wake activity	*iso/iso* vs *trpl*^{302}/*trpl*^{MB}	2.91E-01	0.21	23	22	
night wake activity	*iso/iso* vs *trpl*^{MB}/*trpl*^{MB}	6.31E-01	0.12	23	21	
night wake activity	*trpl*^{302}/*trpl*^{MB} vs *trpl*^{MB}/*trpl*^{MB}	7.10E-01	0.07	22	21	
b. ERG parameter	**genotype comparison**	**p-value**	**effect size *r***	**n1**	**n2**	**significant *r* magnitude**
receptor potential	*iso* vs *trpl*^{302}	2.20E-01	0.26	24	12	
receptor potential	*iso* vs *trpl*^{MB}	8.60E-01	0.03	24	20	
receptor potential	*trpl*^{302} vs *trpl*^{MB}	2.20E-01	0.26	12	20	
on transient	*iso* vs *trpl*^{302}	2.40E-06	0.73	24	12	large
on transient	*iso* vs *trpl*^{MB}	1.50E-01	0.22	24	20	
on transient	*trpl*^{302} vs *trpl*^{MB}	1.00E-05	0.80	12	20	large
off transient	*iso* vs *trpl*^{302}	4.80E-09	0.81	24	12	large
off transient	*iso* vs *trpl*^{MB}	9.90E-01	0.00	24	20	
off transient	*trpl*^{302} vs *trpl*^{MB}	5.30E-08	0.81	12	20	large

845 Table 4. Statistics of a. sleep and b. ERG parameters for *hdc* mutants

a. sleep parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
day sleep	hdc^{MB} vs hdc^{JK910}	9.50E-09	0.81	16	24	large
day sleep	hdc^{MB} vs *iso*	6.10E-05	0.67	16	23	large
day sleep	hdc^{JK910} vs *iso*	7.30E-05	0.60	24	23	large
night sleep	hdc^{MB} vs hdc^{JK910}	1.70E-04	0.61	16	24	large
night sleep	hdc^{MB} vs *iso*	5.40E-08	0.78	16	23	large
night sleep	hdc^{JK910} vs *iso*	4.99E-03	0.42	24	23	moderate
day sleep bout length	hdc^{MB} vs hdc^{JK910}	5.10E-07	0.76	16	24	large
day sleep bout length	hdc^{MB} vs *iso*	4.90E-03	0.49	16	23	moderate
day sleep bout length	hdc^{JK910} vs *iso*	1.08E-01	0.25	24	23	
night sleep bout length	hdc^{MB} vs hdc^{JK910}	8.50E-01	0.03	16	24	
night sleep bout length	hdc^{MB} vs *iso*	3.30E-05	0.64	16	23	large
night sleep bout length	hdc^{JK910} vs *iso*	2.50E-06	0.66	24	23	large
day wake activity	hdc^{MB} vs hdc^{JK910}	7.10E-01	0.06	16	24	
day wake activity	hdc^{MB} vs *iso*	1.40E-01	0.27	16	23	
day wake activity	hdc^{JK910} vs *iso*	2.50E-01	0.19	24	23	
night wake activity	hdc^{MB} vs hdc^{JK910}	2.58E-02	0.36	16	24	moderate
night wake activity	hdc^{MB} vs *iso*	1.02E-02	0.43	16	23	moderate
night wake activity	hdc^{JK910} vs *iso*	1.50E-06	0.68	24	23	large
b. ERG parameter	**genotype comparison**	**p-value**	**effect size *r***	**n1**	**n2**	**significant *r* magnitude**
receptor potential	*iso* vs hdc^{JK910}	8.00E-01	0.24	12	12	
receptor potential	*iso* vs hdc^{MB}	4.00E-01	0.06	12	12	
receptor potential	hdc^{JK910} vs hdc^{MB}	3.80E-01	0.32	12	12	
on transient	*iso* vs hdc^{JK910}	4.20E-05	0.72	12	12	large
on transient	*iso* vs hdc^{MB}	2.20E-04	0.79	12	12	large
on transient	hdc^{JK910} vs hdc^{MB}	9.32E-01	0.02	12	12	
off transient	*iso* vs hdc^{JK910}	5.50E-05	0.85	12	12	large
off transient	*iso* vs hdc^{MB}	2.20E-06	0.85	12	12	large
off transient	hdc^{JK910} vs hdc^{MB}	4.80E-04	0.72	12	12	large

876 Table 5. Statistics of a. sleep and b. ERG parameters for *ort* mutants

a. sleep parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
day sleep	*Canton S* vs *ort^1*	5.20E-03	0.27	60	47	small
night sleep	*Canton S* vs *ort^1*	1.00E-01	0.16	60	47	
day sleep bout length	*Canton S* vs *ort^1*	2.60E-04	0.35	60	47	moderate
night sleep bout length	*Canton S* vs *ort^1*	7.30E-01	0.04	60	47	
day wake activity	*Canton S* vs *ort^1*	1.10E-08	0.55	60	47	large
night wake activity	*Canton S* vs *ort^1*	6.60E-01	0.03	60	47	
day sleep	*iso* vs *ort^{attp}*	4.10E-09	0.68	48	29	large
day sleep	*iso* vs *ort$^{attp/+}$*	8.30E-01	0.02	48	49	
day sleep	*ort^{attp}* vs *ort$^{attp/+}$*	4.50E-10	0.73	29	49	large
night sleep	*iso* vs *ort^{attp}*	6.10E-09	0.67	48	29	large
night sleep	*iso* vs *ort$^{attp/+}$*	1.50E-02	0.25	48	49	small
night sleep	*ort^{attp}* vs *ort$^{attp/+}$*	5.50E-10	0.72	29	49	large
day sleep bout length	*iso* vs *ort^{attp}*	6.20E-02	0.21	48	29	
day sleep bout length	*iso* vs *ort$^{attp/+}$*	1.80E-04	0.39	48	49	moderate
day sleep bout length	*ort^{attp}* vs *ort$^{attp/+}$*	9.00E-07	0.55	29	49	large
night sleep bout length	*iso* vs *ort^{attp}*	1.50E-02	0.32	48	29	moderate
night sleep bout length	*iso* vs *ort$^{attp/+}$*	7.50E-02	0.20	48	49	
night sleep bout length	*ort^{attp}* vs *ort$^{attp/+}$*	3.71E-01	0.10	29	49	
day wake activity	*iso* vs *ort^{attp}*	9.90E-07	0.55	48	29	large
day wake activity	*iso* vs *ort$^{attp/+}$*	3.00E-02	0.22	48	49	small
day wake activity	*ort^{attp}* vs *ort$^{attp/+}$*	6.70E-05	0.45	29	49	moderate
night wake activity	*iso* vs *ort^{attp}*	4.70E-08	0.60	48	29	large
night wake activity	*iso* vs *ort$^{attp/+}$*	4.00E-03	0.29	48	49	small
night wake activity	*ort^{attp}* vs *ort$^{attp/+}$*	3.00E-06	0.52	29	49	large

b. ERG parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
receptor potential	*iso* vs *ort^1*	2.70E-08	0.83	20	12	large
receptor potential	*iso* vs *ort^{attp}*	9.80E-01	0.01	20	12	
receptor potential	*ort^1* vs *ort^{attp}*	1.10E-06	0.85	12	12	large
on transient	*iso* vs *ort^1*	9.30E-08	0.81	20	12	large
on transient	*iso* vs *ort^{attp}*	9.30E-08	0.81	20	12	large
on transient	*ort^1* vs *ort^{attp}*	2.20E-01	0.26	12	12	
off transient	*iso* vs *ort^1*	2.70E-08	0.83	20	12	large
off transient	*iso* vs *ort^{attp}*	4.90E-06	0.83	20	12	large
off transient	*ort^1* vs *ort^{attp}*	7.90E-01	0.06	12	12	

Table 6. Statistics of sleep parameters for *gmrGal4>UAS-Kir2 and Rh1Gal4>UAS-Kir2* lines

sleep parameter	genotype comparison	p-value	effect size *r*	n1	n2	significant *r* magnitude
day sleep	*gmr-Gal4>+ vs gmr-Gal4>UAS-Kir2*	6.90E-03	0.36	32	32	moderate
day sleep	*gmr-Gal4>+ vs +>UAS-Kir2*	1.18E-01	0.20	32	32	
day sleep	*gmr-Gal4>UAS-Kir2 vs +>UAS-Kir2*	1.50E-04	0.51	32	32	large
night sleep	*gmr-Gal4>+ vs gmr-Gal4>UAS-Kir2*	3.90E-01	0.11	32	32	
night sleep	*gmr-Gal4>+ vs +>UAS-Kir2*	2.10E-06	0.62	32	32	large
night sleep	*gmr-Gal4>UAS-Kir2 vs +>UAS-Kir2*	3.80E-04	0.46	32	32	moderate
day sleep bout length	*gmr-Gal4>+ vs gmr-Gal4>UAS-Kir2*	8.80E-01	0.02	32	32	
day sleep bout length	*gmr-Gal4>+ vs +>UAS-Kir2*	6.00E-01	0.11	32	32	
day sleep bout length	*gmr-Gal4>UAS-Kir2 vs +>UAS-Kir2*	6.00E-01	0.12	32	32	
night sleep bout length	*gmr-Gal4>+ vs gmr-Gal4>UAS-Kir2*	2.70E-01	0.14	32	32	
night sleep bout length	*gmr-Gal4>+ vs +>UAS-Kir2*	1.14E-01	0.22	32	32	
night sleep bout length	*gmr-Gal4>UAS-Kir2 vs +>UAS-Kir2*	5.50E-02	0.30	32	32	
day wake activity	*gmr-Gal4>+ vs gmr-Gal4>UAS-Kir2*	9.73E-01	0.01	32	32	
day wake activity	*gmr-Gal4>+ vs +>UAS-Kir2*	7.10E-03	0.35	32	32	moderate
day wake activity	*gmr-Gal4>UAS-Kir2 vs +>UAS-Kir2*	7.10E-03	0.37	32	32	moderate
night wake activity	*gmr-Gal4>+ vs gmr-Gal4>UAS-Kir2*	1.10E-04	0.50	32	32	moderate
night wake activity	*gmr-Gal4>+ vs +>UAS-Kir2*	3.32E-02	0.29	32	32	small
night wake activity	*gmr-Gal4>UAS-Kir2 vs +>UAS-Kir2*	2.37E-01	0.15	32	32	
day sleep	*+>UAS-Kir2 vs Rh1-Gal4>+*	2.60E-02	0.29	28	31	small
day sleep	*+>UAS-Kir2 vs Rh1-Gal4>UAS-Kir2*	1.10E-02	0.36	28	30	moderate
day sleep	*Rh1-Gal4>+ vs Rh1-Gal4>UAS-Kir2*	2.00E-05	0.58	31	30	large
night sleep	*+>UAS-Kir2 vs Rh1-Gal4>+*	4.90E-08	0.74	28	31	large
night sleep	*+>UAS-Kir2 vs Rh1-Gal4>UAS-Kir2*	3.99E-03	0.38	28	30	moderate
night sleep	*Rh1-Gal4>+ vs Rh1-Gal4>UAS-Kir2*	9.40E-04	0.44	31	30	moderate
day sleep bout length	*+>UAS-Kir2 vs Rh1-Gal4>+*	8.74E-01	0.02	28	31	
day sleep bout length	*+>UAS-Kir2 vs Rh1-Gal4>UAS-Kir2*	6.60E-03	0.37	28	30	moderate
day sleep bout length	*Rh1-Gal4>+ vs Rh1-Gal4>UAS-Kir2*	1.50E-03	0.44	31	30	moderate
night sleep bout length	*+>UAS-Kir2 vs Rh1-Gal4>+*	6.70E-01	0.07	28	31	
night sleep bout length	*+>UAS-Kir2 vs Rh1-Gal4>UAS-Kir2*	4.40E-01	0.19	28	30	
night sleep bout length	*Rh1-Gal4>+ vs Rh1-Gal4>UAS-Kir2*	6.70E-01	0.06	31	30	
day wake activity	*+>UAS-Kir2 vs Rh1-Gal4>+*	3.10E-04	0.48	28	31	moderate
day wake activity	*+>UAS-Kir2 vs Rh1-Gal4>UAS-Kir2*	4.40E-06	0.63	28	30	large
day wake activity	*Rh1-Gal4>+ vs Rh1-Gal4>UAS-Kir2*	9.60E-01	0.01	31	30	
night wake activity	*+>UAS-Kir2 vs Rh1-Gal4>+*	4.80E-07	0.62	28	31	large
night wake activity	*+>UAS-Kir2 vs Rh1-Gal4>UAS-Kir2*	6.90E-08	0.68	28	30	large
night wake activity	*Rh1-Gal4>+ vs Rh1-Gal4>UAS-Kir2*	2.00E-01	0.17	31	30	