Title

A fast and tiny current as common generator of slow regular pacemaking in brain and heart

Authors

Arthur Fyon[1], Oleksandra Pavlova[2], Nick Schaar[3], Pietro Mesirca[4], Julien Brandoit[1], Sofian Ringlet[5], Alessio Franci[1,6], Matteo E. Mangoni[4], Jochen Roeper[3], Guillaume Drion[1†], Vincent Seutin[2*†] & Kevin Jehasse[1*†]

Affiliations

[1]Neuroengineering laboratory, Montefiore Institute, University of Liège, Belgium.

[2]Laboratory of Neurophysiology, GIGA Institute, University of Liège, Belgium.

[3]Institute of Neurophysiology, Neuroscience Center, Goethe University, Frankfurt am Main, Germany.

[4]Institut de Génomique Fonctionnelle, Université de Montpellier, CNRS, INSERM, France.

[5]Laboratory of Molecular Regulation of Neurogenesis, GIGA Institute, University of Liège, Belgium.

[6]WEL-T Department, WEL Research Institute, Belgium.

[†]denotes equal contribution

*Corresponding authors. Email: v.seutin@uliege.be – kevin.jehasse@uliege.be

Abstract

Slow pacemaking is a key physiological process in specific excitable cells, yet its underlying mechanisms remain debated. Here we identify a conserved, voltage-dependent pacemaker current that is essential for slow, regular firing in both midbrain dopaminergic neurons and sinoatrial node cells. Conductance-based models incorporating this current reproduce stable pacemaking, requiring a fast-activating, small-amplitude current. This is further confirmed by dynamic-clamp experiments in dopaminergic neurons. Replacing the pacemaker current in a model with a voltage-independent conductance such as the non-selective sodium leak channels fails to sustain slow rhythmicity, highlighting the necessity for an adequate voltage dependence. Our results suggest a novel and shared biophysical mechanism for slow pacemaking in neuronal and cardiac systems.

Introduction

Pacemaking activity is a fundamental property of a subset of excitable cells that underpins vital rhythmic functions across biological systems. Pacemaker cells can be classified, based on the firing frequency, as either fast or slow pacemakers. While the mechanism to sustain fast pacemaking is known (*1*), slow pacemaking remains controversial. It has been suggested that it originates from the balanced interplay between several depolarizing and hyperpolarizing currents (*2–5*). The main currents involved are due to hyperpolarization-activated cyclic nucleotide-gated (HCN sustaining I_H in neurons and I_F in cardiac cells) and voltage-gated Ca^{2+} (Ca_v) channels, on the one hand, and A-type fast inactivating K^+ channels (I_A), on the other hand. Yet, experimental data shows that blocking or altering any of these channels does not eradicate slow pacemaking, although both its frequency and regularity can be affected (*3, 6–9*). Another hypothesis is that the production of such a precise (coefficient of variation below 5%) and robust low frequency (1-5 Hz on average) pacemaking needs the reliable activation of a pacemaker current (I_{pace}) during the interspike interval (*1, 6, 10, 11*) . Yet the nature, dynamics, and origins of such an essential current have remained elusive to date.

Recently, we found that a compound previously shown to block gating pore currents, 1-(2,4-xylyl)guanidinium (XG), eradicates the pacemaking activity of rodent midbrain dopamine neurons (mDANs) (*6*). Using voltage ramps and pacemaker-clamp, we were able to isolate the tiny current blocked by XG, reminiscent of the elusive I_{pace}. Since slow pacemaking is such a fundamental behavior for life, we hypothesized that this mechanism is conserved across neuronal and cardiac pacemaker cells and species. We therefore investigated whether similar effects of XG would be observed in human mDANs and in rodent sinoatrial node (SAN) cells and found it to be the case. Dynamic-clamp experiments, in which electrophysiological recordings are coupled to exogenous current injections to either add or subtract conductances, confirmed that XG-sensitive I_{pace} is necessary and sufficient for slow pacemaking. Next, we used both a conductance-based model of a mDAN (*12*) and a reduced neuronal model and showed that the voltage dependence of I_{XG} is ideally suited to be the effector of slow pacemaking. Taken together, our data supports the hypothesis of a conserved mechanism for slow pacemaking, encompassing neuronal and cardiac cells.

Results

Evidence for a shared mechanism of slow pacemaking across species and systems

We had previously shown that pacemaking of rodent mDANs is blocked by XG and had isolated the underlying current (**Fig. 1A-C**) (*6*). We first asked whether pacemaking in human mDANs derived from two different induced pluripotent stem cells (hIPSCs) lines (see Methods) would be similarly affected by the drug. These neurons were identified as mDANs based on their spontaneous slow firing frequency, hyperpolarization by dopamine and, in some cases, the expression of tyrosine hydroxylase (**Fig. S1**; see Methods). Upon superfusion of XG, their spontaneous activity was silenced (2.28 ± 0.44 Hz in control vs 0.12 ± 0.05 in XG, n = 9, p = 0.0039, Wilcoxon test; **Fig. 1D, E**), similarly to what we had observed in rodent models. In particular, as in rodent mDANs, the silencing of the electrical activity by XG does not involve a hyperpolarization, contrary to what is seen with dopamine, which is known to open G-protein-coupled inwardly rectifying channels in these cells (**Fig. S1**). Moreover, we observed a current sensitive to XG in human mDANs which is similar the one observed in rat mDANs (**Fig. 1F**). Taken together, these results suggest that the mechanism of pacemaking is conserved across species in this cell type.

To investigate whether this finding could be extended to another cell type, we next explored the case of SAN cells of the heart, which are also slow (1-5 Hz), very regular, pacemakers. In addition, similar to mDANs, these cells express high levels of L-type $Ca_v1.3$ channels and HCN channels (*13*). Upon superfusion of XG, we observed a marked decrease of the pacemaking activity in SAN cells (from 166.5 ± 9.1 bpm in control to 24.5 ± 10.1 bpm in XG, n = 11, p = 0.001, Wilcoxon test ; **Fig. 1G, H**). The silencing effect of XG was not due to its interaction with K^+ or Ca_v channels, nor with HCN channels (**Fig. S2**), but due to the inhibition of a pacemaker current (named I_{pace} hereafter) that had similar characteristics to those previously observed in mDANs (**Fig. 1I, J**). Interestingly, I_{pace} starts operating around -50 mV and its amplitude increases up to ~-20 mV, a voltage more depolarized than the threshold of the action potential in both cell types. Taken together, this data is strongly suggestive of a shared mechanism for slow pacemaking in mDANs and in SAN cells.

Modeling the pacemaker current confirms its ability to sustain slow pacemaking

We used a well-established conductance-based model to study slow pacemaking (*12*). Since this model is set to pace steadily without a pacemaker current, it relies on shifting

the activation curve of Na$^+$ (g_{Na}) and L-type Ca^{2+} (g_{CaL}) conductances to artificially hyperpolarized potentials. We adjusted these activation curves to their measured physiological values in mDANs (*14*, *15*) (**Fig. S3C**), which led to a loss of spontaneous firing and a shift from type I to type II excitability (**Fig. S3B**). We then modeled the pacemaker conductance (g_{pace}) from the current-voltage curve of the XG-sensitive I_{pace} (**Fig. S3D**) and implemented it into the adjusted model (**Fig. S3E**). We were able to recover a pacemaking activity with a voltage range of activation of g_{pace} similar to what had been observed experimentally (**Fig. 2B**) (*6*), as well as to recover type 1 excitability (**Fig. S3B**). Currentscape analysis (*16*) revealed the crucial contribution of g_{pace} during pacemaking (**Fig. 2A**). We then measured the total current between two spikes and observed that the resulting current is driven by g_{pace} (**Fig. 2B**) and has a small amplitude similar to previous experimental observations in mDANs (*17*). Finally, we measured the charge transfer of each conductance during the interspike interval and the full period of the simulation (**Fig. 2C**). Interestingly, the transfer charge of g_{pace} (-190.76 fC) is smaller than the other conductance such as g_{CaL} (-1457.89 fC) or H-type conductance (g_H) (-2077.48 fC) during the interspike interval. Overall, these data highlight the essential role of the small conductance g_{pace} in slow pacemaking.

We next investigated the dynamical range of g_{pace} which is necessary to maintain a steady low frequency spontaneous activity (**Fig. 2D**). We explored both the density and activation time constant of g_{pace}, relative to the activation of Na$_v$ channels, and found that the model remains stable as long as the activation of g_{pace} is very fast, close to instantaneous. Moreover, g_{pace} can have a wide range of densities, making the phenomenon robust. To verify that these observations are not limited to the particular parameters used, we perfomed the same analysis using random parameters in which slow pacemaking can be observed (**Fig. S4**). In all cases, we observed similar requirements for the parameters of g_{pace}. Moreover, the addition of g_{pace} into the model increased ion channel degeneracy (**Fig. S5**), especially for g_{Na} and g_{CaL}, suggesting that it may normalize neuronal output in the face of biological variability (*18–21*).

To further explore the structure of this increased degeneracy, we performed principal component analysis on the pacemaking solutions of the adjusted model. The analysis revealed no preferred direction of variability, indicating that the contribution of g_{pace} does not impose a dominant constraint on the system but instead distributes flexibility across all conductances (**Fig. S5C**). This isotropic degeneracy suggests that slow

pacemaking remains robust even when the underlying ion channel expression varies
extensively, emphasizing the stabilizing role of g_{pace} in maintaining reliable activity.
Next, we used the dynamic-clamp approach to test how g_{pace} regulates slow
pacemaking in spontaneously active DANs from *substantia nigra pars compacta* (*SNc*)
(**Fig. 3A**). First, we injected several values of negative conductance and evaluated the
percentage of decrease in firing frequency (**Fig. 3B**). We observed a frequency
decrease of 50% at -0.06 nS and complete silencing at -0.3 nS (1.54 Hz (interquartile
range: 1.19 – 1.86 Hz) in ctrl vs. 0 Hz (interquartile range: 0) with g_{pace} = - 0,3 nS, n =
20, N = 3, p = <0,0001, Wilcoxon test ; **Fig 3B**). Injecting positive values for g_{pace}
increased the firing frequency as expected, leading to a depolarization block at the
highest value tested (above 0.3 nS, **Fig. 3C**). Finally, we preincubated *SNc* DANs with
XG to silence their activity and injected positive g_{pace} in order to test whether we are
able to rescue slow pacemaking (**Fig. 3D**). Overall, the rescue was successful in all
trials (n = 11) at a variable intensity of g_{pace} which is to be expected because of the
diversity in the mDAN population (*22*). The latter experiment demonstrates that our
modeling and experimental data are perfectly congruent.
152
Requirements for slow pacemaking
We next sought to understand how important g_{pace} is for slow pacemaking. We reduced
our conductance-based model to a minimal one, keeping only g_{Na} and g_{Kd} for spike
generation, g_{LNS} (leak conductance) to set the membrane at the desired potential.
Adding g_{pace} in this minimal configuration allows slow pacemaking similarly to our
complete model (**Fig. 4A**). In this model, slow pacemaking is still maintained with fast
activating g_{pace} over a wide range of densities, from 4.5 mS/mm² to 35.5 mS/mm² (**Fig.**
4B). Since our data shows that minimal configuration is sufficient to create slow
pacemaking, we investigate the role of the other conductances, especially A-type K$^+$
(g_{KA}), H-type (g_H) and Ca^{2+}-dependent K$^+$ (g_{KSK}) conductances. They have been shown
in slices to help maintaining the firing frequency and regularity constant during
membrane potential changes caused by excitatory and inhibitory inputs (*3, 17, 23*). We
introduced current noise in our minimal model and observed that the pacemaking
becomes irregular (**Fig. 4C**). In our complete model, the firing frequency is robust to
current injection, similarly to what have been observed by others (*3, 17, 23*). Moreover,
we also observed that these conductances, and especially g_{KSK}, help to reduce the
slope of the firing frequency – injected current relationship, which is expected.

We then challenged the requirement of a voltage-gated activation for g_{pace} in order to
obtain slow pacemaking. We replaced g_{pace} by a linear leak conductance similar to
NALCN channels (g_{Nalcn}), as they had been suggested to be the main drive of
pacemaking in mDANs (10, 24–26). g_{Nalcn} failed to sustain slow pacemaking (**Fig. S6**).
In fact, low or high density of g_{Nalcn} could not generate spontaneous firing (**Fig. S6B**),
while activity could be observed in a small window of density, going from type II
excitability to fast pacemaking type I excitability. Our data therefore supports the notion
that slow pacemaking requires a current with a specific voltage-dependent activation
during the interspike interval, not a linear one.

Discussion
Slow pacemaking is a unique property which is observed in a subset of cells that play
a major role in rhythmic biological processes. However, its exact mechanism is still
controversial. Recently, we found that XG inhibits the pacemaking of rodent mDANs
and were able to isolate the current blocked by this compound (6). Here we show that
this effect can be extended to human mDANs and to mouse SAN cells. Remarkably,
the XG-sensitive conductance or g_{pace} has very similar voltage-dependence in mDANS
and SAN cells, starting to operate at ~-50 mV and slowly increasing in amplitude
towards more depolarized potentials. In our minimal conductance-based model, we
show that such voltage dependence is able to generate slow pacemaking, whereas
addition of a linear conductance such as NALCN is unable to induce this firing pattern.
In our adjusted conductance-based model of a mDAN, we show that addition of g_{pace}
is able to induce biologically realistic pacemaking. The properties of g_{pace} allows to
extend ion channel degeneracy, which makes our model more robust to biological
variability (**Fig. S5**). In this work, we describe key properties that I_{pace} should have.
First, its activation should be fast. Second, its amplitude can vary within a relatively
large range, which guarantees robustness. In our minimal model, we show that the
largest conductances, such as g_{CaL}, g_{KA} or g_H are not required for slow pacemaking (3,
6), but that I_{pace} is equally able to generate this rhythm.
Dynamic-clamp experiments confirmed the importance of g_{pace} (**Fig. 3**). Moreover, the
range of conductances to either rescue slow pacemaking upon XG perfusion or to
silence the neurons was a small fraction of the cell conductance, around 3 nS
measured in mDANs (6). Overall, 0.3 nS was enough in both conditions, confirming
the fact that pacemaking relies on a tiny current. These data support our hypothesis

that the pacemaker channel should have a unitary conductance value in the fS range (6, 11). In fact, if g_{pace} were to have a single channel conductance similar to Na_v channels (around 20 pS), it would imply that slow pacemaking is sustained by at most 15 channels. Because of stochasticity, it will lead to very irregular firing in neurons and in the heart.

In mDANs, the role of several ion channels in pacemaking has been investigated over the years. L-type Ca_v channels were considered, since their pharmacological blockade with high concentrations of dihydropyridines silenced these neurons (27–29). However, these data were not observed using lower, pharmacologically relevant concentrations (6, 7, 30, 31), therefore suggesting an off-target effect of dihydropyridines at high concentrations. We also confirm that in this study, as our minimal model runs without a Ca_v conductance. These channels are now considered as a linear amplifier of the firing frequency in mDANs (7). NALCN channels were also considered as they are expressed in most rhythmic neurons (32). Further studies showed controversial results, in which NALCN gene deletion or pharmacological blockade suppressed (24–26, 33) or did not impact pacemaking activity (6, 25). Our simulations show that a linear depolarizing conductance is unlikely to sustain stable and slow pacemaking (**Fig. S6**).

On the other hand, SAN cell pacemaking has been linked to either Ca^{2+}-linked processes or HCN channel activity (34). In the Ca^{2+} clock theory, L-type $Ca_v1.3$ channels are responsible for the main entry of Ca^{2+}, which is then pumped by SERCA pumps in the sarcoplasmic reticulum or by the Na^+/Ca^{2+} exchanger NCX outside the cell. The genetic removal of $Ca_v1.3$ channels (35) or HCN channels (36) leads to arrhythmia, but not to a silenced activity. On the other hand, XG does silence pacemaking in these cells by blocking a current that is highly similar to the one found in mDANs. This observation clearly opens up a new hypothesis on cardiac pacemaking that is more consistent with previous experimental results than the current theories.

In conclusion, we shed new light on the controversial mechanism of slow pacemaking in excitable cells and, importantly, demonstrate a commonality across systems, an observation that is very novel and surprising.

Materials and Methods

Modeling

Programming language

The Julia programming language was used in this work (*37*). Numerical integration was

realized using *DifferentialEquations.jl*. Fitting was realized using *LsqFit.jl*. PCA was

realized using *MultivariateStats.jl*.

Equation 1:

$$C\dot{V} = -\bar{g}_{Na}\, m^3\, h\, h_s\, (V - E_{Na}) - \bar{g}_{CaL}\, l\, (V - E_{Ca}) - \bar{g}_{Kd}\, n^3\, (V - E_K)$$

$$- \bar{g}_{KA}\, p\, \left(\frac{q_1}{2} + \frac{q_2}{2}\right) (V - E_K) - \bar{g}_{KERG}\, o\, (V - E_K)$$

$$- \frac{\bar{g}_{KSK}}{1 + \left(\frac{0.00019}{[Ca^{+2}]}\right)^4}\, (V - E_K) - \bar{g}_H\, m_H^2\, (V - E_H) - g_{LCa}\, (V - E_{Ca})$$

$$- g_{LNS}\, (V - E_{LNS}) + \frac{100\, I_{app}}{\pi\, d\, L}$$

Here, the dot notation represents time derivative (in /ms), V is the membrane potential

(in mV), and C the membrane capacitance (in µF/cm²). The ionic currents are: I_{Na}

(transient sodium), I_{CaL} (L-type calcium), I_{Kd} (delayed rectifier potassium), I_{KA} (A-type

potassium), I_{KERG} (ether-à-go-go potassium), I_{KSK} (SK potassium), I_H

(hyperpolarization-activated current), I_{LCa} (calcium leak), and I_{LNS} (non-specific leak).

Reversal potentials are denoted by E. The intracellular calcium concentration $[Ca^{+2}]$

follows voltage-dependent kinetics as in (*13*). The soma geometry is defined by its

diameter d and length L (in µm). The applied current is I_{app} (in pA).

Maximal conductances are denoted by the bar notation (in mS/cm²), representing the

channel conductance when fully open. Gating variables follow first-order voltage-

dependent kinetics of the following form.

Equation 2:

$$\tau_x(V)\, \dot{x} = x_\infty(V) - x$$

Here, x is the gating variable, $\tau_x(V)$ its voltage-dependent time constant, $x_\infty(V)$ its

steady-state activation or inactivation function. Variables that were adjusted in this

work are specified in the following section.

Unless otherwise stated, the original model parameters were used (Table S1) (*13*).

$$\bar{g}_{Na} = 6;\ \bar{g}_{CaL} = 0.139;\ \bar{g}_{Kd} = 1.117;\ \bar{g}_{KA} = 1.68;\ \bar{g}_{KERG} = 0.13;$$

$$\bar{g}_{KSK} = 0.07;\ \bar{g}_H = 0.078;\ g_{LCa} = 0.00245;\ g_{LNS} = 0.28;$$

$$C = 1;\ E_{Na} = 60;\ E_K = -90;\ E_{Ca} = 50;\ E_H = -29;\ E_{LNS} = -65;\ d = 15;\ L = 25$$

Adjustment of the mDAN conductance-based model and fitting of the pacemaking
current

Before introducing the pacemaking current, activation of the transient sodium and L-
type calcium channel was realized. In the original model (*13*), the steady-state
activation functions were defined as follows.

Equation 3:

276
$$m_\infty(V) = \frac{1}{1 + e^{-\frac{V+30.09}{13.2}}}; \ l_\infty(V) = \frac{1}{1 + e^{-\frac{V+45}{7.5}}}$$

These sigmoidal functions have half-activation voltages of −30.09 mV and −45 mV,
respectively, which are substantially more hyperpolarized than values reported
experimentally. In the adjusted model, both activation curves were shifted to −10 mV.
The ultraslow inactivation of the transient sodium current (h_s) was disabled, as it was
an adjustment to generate slow pacemaking in the original model (*12*).

The pacemaking current was fitted to a previously published voltage–current
relationship (*3*) (**Fig. 1C**). The pacemaking current was modeled as a non-inactivating
current, which provided an accurate fit without unrealistic parameter estimates.

Equation 4:

288
$$\begin{cases} I_{pace} = \bar{g}_{pace} m_{pace} \left(V - E_{pace} \right) \\ \tau_{m,pace}(V) \, \dot{m}_{pace} = m_{pace,\infty}(V) - m_{pace} \\ m_{pace,\infty}(V) = \dfrac{1}{1 + e^{-\frac{V - V_{half}}{V_{slope}}}} \end{cases}$$

290   Because the fit was performed on steady-state voltage–current data, no biological
291   information was available on the time constant of pacemaking activation. We therefore
292   set

293   *Equation 5:*

294
$$\tau_{m,pace}(V) = \alpha \cdot \tau_m(V)$$

295   where $\tau_m(V)$ is the activation time constant of the transient sodium current.

297   Fitting yielded the following parameter values, given to one decimal place.

298
$$\bar{g}_{pace} = 6.5; \ V_{half} = -13.3; \ V_{slope} = 6; \ E_{pace} = 4.3$$

Since this pacemaking conductance corresponds to a specific neuron, it was varied throughout the study to test robustness rather than fixed at the fitted value. Unless otherwise stated, the following parameters were used in the adjusted model.

$$\bar{g}_{Na} = 25; \ \bar{g}_{CaL} = 1; \ \bar{g}_{Kd} = 10; \ \bar{g}_{KA} = 1.68; \ \bar{g}_{KERG} = 0.13; \ \bar{g}_{KSK} = 0.3;$$

$$\bar{g}_H = 0.078; \ g_{LCa} = 0.00245; \ g_{LNS} = 0.28; \ \bar{g}_{pace} = 5;$$

$$C = 1; E_{Na} = 60; E_K = -90; E_{Ca} = 50; E_H = -29; E_{LNS} = -65; d = 15; L = 25$$

Modeling of a NALCN current

In **Fig. 4C-E**, the NALCN conductance was modeled as a voltage-independent sodium current.

Equation 6:

$$I_{NALCN} = g_{NALCN} \ (V - E_{Na})$$

Degenerate population building

In **Fig. S5**, two degenerate populations (N=200) exhibiting pacemaking were generated for the original and adjusted models. The procedure was identical for both populations. All ionic conductances were randomly assigned using a uniform distribution between 0 and an upper bound value, corresponding to the ordinates of **Fig. S5B**. Each model was simulated and retained in the population if its firing frequency was between 1 and 5 Hz, if the spike peak exceeded 0 mV, and if the minimum membrane potential was between −90 and −60 mV.

<u>Cell Culture</u>

Cell lines

Two human induced pluripotent stem cell (hiPSC) lines were used in this study: GM23338 (gift from Prof. Ira Mercedes Espuny Camacho, GIGA Neurosciences, University of Liège) and a human TH-reporter iPSC line expressing mCherry (gift from Dr. Aleksandar Rakovic, Institute of Neurogenetics, University of Lübeck) (*38*). Prior to differentiation, cells were maintained for 2–3 consecutive passages in E8 medium using standard EDTA-based passaging when confluency reached 70–80%.

Differentiation to midbrain dopaminergic neurons

Differentiation was carried out using a slightly modified version of the protocol published by Stathakos et al. (2019) (*39*). Briefly, hiPSCs were cultured in E8 medium for up to 4 days before switching to neural induction medium consisting of N2B27 supplemented with 100 nM LDN193189, 10 µM SB431542, 300 ng/mL SHH, and 0.6 µM CHIR99021. This was considered neuralization day 0. Cells were passaged on polyornithine/laminin-coated plates at days 3 and 7 using Accutase at a density of 4 × 10^4 cells/cm² in induction medium with 10 µM Y27632 or 1:100 RevitaCell.

From day 11, the medium was replaced with progenitor expansion medium (N2B27 with 20 ng/mL BDNF, 20 ng/mL GDNF, 0.2 mM ascorbic acid, and 10 µM Y27632). Upon confluence, cells were passaged every 3-4 days in this medium. Terminal differentiation started on day 16 by seeding cells at 5,000 cells/cm² onto PO/laminin-coated coverslips in N2B27 supplemented with 20 ng/mL BDNF, 20 ng/mL GDNF, 0.2 mM ascorbic acid, 500 µM db-cAMP, 10 µM DAPT, and 10 µM Y27632. Cells were allowed to attach for ~1 hour before flooding wells with the same medium. Media was refreshed every 3 days with 50% volume exchange.

Tissue preparation

Animals

For dynamic-clamp experiments, adult male C57BL/6N (Charles River Laboratories) mice (8 to 14 weeks of age) were used. A maximum of 5 mice were housed per cage and single animal housing was avoided when possible. Animals were maintained on a 12-hour light-dark cycle and provided with water and food ad libitum. All experimental procedures involving mice were approved by the German Regional Council of Darmstadt.

For SAN cells experiments, the study is in accordance with the Guide for the Care and Use of Laboratory Animals published by the US National Institute of Health (NIH Publication No. 85-23, revised 1996) and European directives (2010/63/EU). All mice used were from the C57BL/6J background (Charles River and Janvier labs). Experimental procedures were approved by the Ethical Panel of the University of Montpellier and the French Ministry of Agriculture (protocol no: 2017010310594939). Animals were housed in the IGF animal facility with free access to food and water and were exposed to 12 h light/dark cycles in a thermostatically controlled room (21–22 °C) with 40−60% humidity.

Slice preparation for mouse midbrain dopamine neurons

Animals were anesthetized by intraperitoneal injection of ketamine (250 mg/kg; Ketaset, Zoetis) and medetomidine hydrochloride (2.5 mg/kg; Domitor, OrionPharma) before intracardial perfusion using ice-cold ACSF consisting of the following: 50 mM sucrose, 125 mM NaCl, 2.5 mM KCl, 25 mM $NaHCO_3$, 1.25 mM NaH_2PO_4, 2.5 mM glucose, 6 mM $MgCl_2$, 0.1 mM $CaCl_2$, and 2.96 mM kynurenic acid (Sigma-Aldrich), oxygenated with 95% O_2 and 5% CO_2. Rostral coronal midbrain slices (bregma, −2.92 to −3.16 mm) were sectioned into 250-µm slices using a vibrating blade microtome (VT1200s, Leica). Before the experiment, slices were kept at 37°C for 1 hour in oxygenated extracellular solution containing the following: 22.5 mM sucrose, 125 mM NaCl, 3.5 mM KCl, 25 mM $NaHCO_3$, 1.25 mM NaH_2PO_4, 2.5 mM glucose, 1.2 mM $MgCl_2$, and 1.2 mM $CaCl_2$.

Mouse sinoatrial node cell isolation

Sinoatrial node myocytes were isolated as previously described (*40*). Briefly, Sinoatrial tissue (SAN) strips were immersed into "low Ca^{2+}" Tyrode's solution containing (in mM): 140 NaCl, 5.4 KCl, 0.5 $MgCl_2$, 1.2 KH_2PO_4, 50 taurine, 5.5 D-glucose, 1 mg/mL BSA, and 5 Hepes (adjusted to pH 6.9 with NaOH) for 5 min. The tissue was then transferred into the low-Ca^{2+} solution containing Liberase TH (0.15 mg/mL, Roche) and Elastase (0.48 mg/mL, Worthington). Digestion was carried out for 15-20 min at 36 °C. SAN strips were then washed in "Kraftbrühe" (KB) solution containing 100 L-glutamic acid potassium salt, 10 L-Aspartic acid potassium salt, 25 KCl, 80 KOH, 10 KH_2PO_4, 2 $MgSO_4$, 5 Creatine, 0.5 EGTA, 20 D-glucose, 20 taurine, 1 mg/mL BSA, and 5 Hepes (adjusted to pH 7.2 with KOH). Single myocytes were isolated from the SAN tissue by manual agitation using a flame-forged Pasteur's pipette in KB solution at 36 °C. For recovering of pacemaker activity, Ca^{2+} was gradually reintroduced into the cells' storage solution to a final concentration of 1.8 mM. Cells were then stored at room temperature until use.

Electrophysiological Recordings

Midbrain dopamine neurons

For rat mDANs, the recordings were obtained from previous work (*6*)

For human mDANs, electrophysiological properties of differentiated neurons were assessed using both on-cell and whole-cell patch-clamp recordings performed at 30°C with a Multiclamp 700B amplifier and Clampex 11.1 software. The extracellular solution contained (in mM): 140 NaCl, 5 KCl , 2 CaCl$_2$,2 MgCl$_2$,15 HEPES, and 10 D-glucose; pH was adjusted to 7.3 with NaOH. The intracellular pipette (3-5 MΩ) solution contained (in mM): 135 K-gluconate, 10 NaCl, 0.5 CaCl$_2$, 15 HEPES, 2 Na$_2$ATP, and 5 EGTA; pH adjusted to 7.3 with KOH, osmolarity ~ 300 mosm. Putative *substantia nigra pars compacta* (*SNc*)-type mDANs were identified by similar characteristic electrophysiological features we used to identify rodent mDANs (*6, 41–43*), including spontaneous pacemaking activity (0.5–5 Hz), action potential duration > 1.35 ms (measured at half-amplitude), a prominent hyperpolarization-activated inward current (I_H), and membrane hyperpolarization > 5 mV in response to 100 µM dopamine from a holding potential of -60 mV. I_H was observed in many, but not all neurons sastifying these criteria. To test the inhibition of pacemaker activity, 600 µM of 1-(2,4-xylyl)guanidinium mesylate (XG) was applied to the bath solution. XG was produced as previously described (*6*).

For dynamic-clamp recordings in mouse *SNc* DANs, slices were placed in a heated recording chamber (37°C) that was superfused with the oxygenated extracellular solution with a flow rate of 2 to 4 ml/min. CNQX (20 µM), DL-AP5 (10 µM), gabazine (4 µM; SR95531), and were added to inhibit respectively excitatory and inhibitory synaptic transmission. Neurons were visualized using infrared differential interference contrast video microscopy with a digital camera (VX55, Till Photonics) connected to an upright microscope (Axioskop 2, FSplus, Zeiss). Patch pipettes were pulled from borosilicate glass (GC150TF-10, Harvard Apparatus, Holliston, MA, USA) using a temperature controlled, horizontal pipette puller (DMZ-Universal Puller, Zeitz). Patch pipettes (3 to 5 MΩ) were filled with a solution containing the following: 135 mM K-gluconate, 5 mM KCl, 10 mM HEPES, 0.1 mM EGTA, 5 mM MgCl$_2$, 0.075 mM CaCl$_2$, 5 mM Na adenosine 5′-triphosphate, 1 mM Li guanosine 5′-triphosphate, and 0.1% Neurobiotin, adjusted to a pH 7.35 with KOH. For pharmacology, slices were preincubated for at least 10 min with XG mesylate (2 mM) before recording. Recordings were performed using an EPC-10 patch-clamp amplifier (HEKA Elektronik) with a sampling rate of 20 kHz and a low-pass filter (Bessel, 5 kHz). For analysis, recordings were further digitally filtered at 1 KHz. A realtime Linux-based data acquisition program

(RTXI; http://rtxi.org) was used to inject online calculated currents (acquisition rates of 10 kHz) of the modelled XG-sensitive conductance according to the following equation:

Equation 7

$$\begin{cases} I_{pace} = \bar{g}_{pace} m_{pace} \left(V - E_{pace} \right) \\ \tau_{m,pace} \, \dot{m}_{pace} = m_{pace,\infty}(V) - m_{pace} \end{cases}$$

V is the membrane voltage in mV, E_{pace} the reversal potential in mV and g_{pace} is the conductance in nS, according to fit the fit in **Fig.2A**.

Mouse sinoatrial node cells

SAN cells were harvested in custom-made chambers with glass bottoms for cell attachment and superfused with Tyrode solution containing (in mM): 140 NaCl, 5.4 KCl, 1 MgCl$_2$, 1.8 CaCl$_2$, 5.5 D-glucose and 5 Hepes (adjusted to pH 7.4 with NaOH). warmed at 36°C before recording.

XG-sensitive, delayed-rectifier potassium ($I_{K,r}$) and "funny" (I_F) currents were recorded in extracellular Tyrode's solution under the standard whole-cell configuration whereas action potentials were measured by perforated patch clamp technique with escin (0.04 mg/mL, Sigma-Aldrich) also in extracellular Tyrode's solution. For I_F recordings, 2 mM BaCl$_2$ was added to Tyrode's solution to block $I_{K,1}$. Electrodes had a resistance of 3–4 MΩ when filled with a solution containing (in mM): 80 K-aspartate, 50 KCl, 1 MgCl$_2$, 2 CaCl$_2$, 5 EGTA, 5 HEPES, and 3 ATP-Na (adjusted to pH 7.2 with KOH). Seal resistances were in the range of 2–5 GΩ.

Superfusion of pre-warmed (36°C) experimental solutions was achieved by using a multi-MPRE8 heating pen (Cell Micro Controls) and data acquisition was performed using Multiclamp 700B patch-clamp amplifiers connected to a Digidata 1322A interface (Molecular Devices, Saint Jose, CA, USA).

Immunocytochemistry

The DA phenotype of neurons in the process of maturation was confirmed by immunocytochemistry as described earlier (*44*) with the use of tyrosine hydroxylase (TH) (Merck, AB152) and βIII tubulin (TUBB3)/ TUJ1 (BioLegend, MMS-435P) antibodies. Samples were imaged using an inverted confocal microscope (Leica SP5), with a resolution of 512 x 512 pixels, pixel size is 0.758 μm. A ratio of TH+/TUBB3+

was measured by counting the number of TH+ or TUBB3+ cells, in order to assess the maturation of the human mDAN culture (*44*).

Data analysis

Dynamic-clamp data was digitally filtered at 1 kHz and further exported as MAT-files using Fitmaster (Heka Elektronik). Offline data analysis was performed using custom-written scripts in MATLAB. Spike thresholds were determined using the first derivative of the recording where $dVm/dt \geq 10$ mV/ms. Spike thresholds, peaks and minima after hyperpolarization potential were visually verified for each cell. The g_{pace}-modulated firing frequency was averaged for the first three ISIs (frequency in Hz).

All statistical analysis were performed using GraphPad Prism 10.2.1. Non-parametric Wilcoxon's tests were performed and data are reported as mean ± standard error of the mean (SEM) and as median (interquartile range). Number of cells are denoted by n, number of animals used as N.

Supplementary Text

Specific parameter values used

In **Fig. 4A-B**, the non-specific leak conductance was reduced to 0.01 to lower the firing frequency.

In **Fig. S4**, ten sets of conductances were randomly chosen from the degenerate population with a fixed random seed for reproducibility.

In **Fig. S6**, a NALCN conductance of 0.012 mS/cm^2 was used, and values of 0.004 mS/cm^2 and 0.012 mS/cm^2 were used for the FI curves in **Fig. S6C**. The CaL conductance was reduced to 0.05 mS/cm^2 and the Kd conductance to 3 mS/cm^2 to enlarge the firing region with respect to the NALCN conductance and facilitate visualization.

References

1. B. P. Bean, Mechanisms of pacemaking in mammalian neurons. *J. Physiol.* **n/a** (2024).

2. E. Marder, J.-M. Goaillard, Variability, compensation and homeostasis in neuron and network function. *Nat. Rev. Neurosci.* **7**, 563–574 (2006).

3.   J. Amendola, A. Woodhouse, M. F. Martin-Eauclaire, J. M. Goaillard, Ca
2+/cAMP-sensitive covariation of I A and I H voltage dependences tunes
rebound firing in dopaminergic neurons. *J. Neurosci.* **32**, 2166–2181 (2012).

4.   Y. Amarillo, E. Zagha, G. Mato, B. Rudy, M. S. Nadal, The interplay of seven
subthreshold conductances controls the resting membrane potential and the
oscillatory behavior of thalamocortical neurons. *J. Neurophysiol.* **112**, 393–410
(2014).

5.   E. G. Lakatta, V. A. Maltsev, T. M. Vinogradova, A Coupled SYSTEM of
Intracellular Ca2+ Clocks and Surface Membrane Voltage Clocks Controls the
Timekeeping Mechanism of the Heart's Pacemaker. *Circ. Res.* **106**, 659–673
(2010).

6.   K. Jehasse, L. Massotte, S. Hartmann, R. Vitello, S. Ringlet, M. Vitello, H. C.
Chua, S. A. Pless, D. Engel, J.-F. Liégeois, B. Lakaye, J. Roeper, V. Seutin, The
gating pore blocker 1-(2,4-xylyl)guanidinium selectively inhibits pacemaking of
midbrain dopaminergic neurons. *Neuropharmacology* **197**, 108722 (2021).

7.   J. Shin, L. Kovacheva, D. Thomas, S. Stojanovic, C. J. Knowlton, J. Mankel, J.
Boehm, N. Farassat, C. Paladini, J. Striessnig, C. C. Canavier, G. Geisslinger, J.
Roeper, Cav1.3 calcium channels are full-range linear amplifiers of firing
frequencies in lateral DA SN neurons. *Sci. Adv.* **8**, eabm4560 (2022).

8.   R. E. BoSmith, I. Briggs, N. C. Sturgess, Inhibitory actions of ZENECA ZD7288
on whole-cell hyperpolarization activated inward current (If) in guinea-pig
dissociated sinoatrial node cells. *Br. J. Pharmacol.* **110**, 343–349 (1993).

9.   E. E. Verheijck, A. C. G. van Ginneken, R. Wilders, L. N. Bouman, Contribution
of L-type Ca2+current to electrical activity in sinoatrial nodal myocytes of rabbits.
Am. J. Physiol.-Heart Circ. Physiol. **276**, H1064–H1077 (1999).

10.   Z. M. Khaliq, B. P. Bean, Pacemaking in Dopaminergic Ventral Tegmental Area
Neurons: Depolarizing Drive from Background and Voltage-Dependent Sodium
Conductances. *J. Neurosci.* **30**, 7401–7413 (2010).

11. V. Seutin, K. Jehasse, G. Drion, An alternative mechanism for slow pacemaking.
J. Physiol. **n/a** (2024).

12. N. Yu, C. C. Canavier, A Mathematical Model of a Midbrain Dopamine Neuron
Identifies Two Slow Variables Likely Responsible for Bursts Evoked by SK
Channel Antagonists and Terminated by Depolarization Block. *J. Math.*
Neurosci. JMN **5**, 5 (2015).

13. M. E. Mangoni, J. Nargeot, Genesis and Regulation of the Heart Automaticity.
Physiol. Rev. **88**, 919–982 (2008).

14. V. Seutin, D. Engel, Differences in Na+ conductance density and Na+ channel
functional properties between dopamine and GABA neurons of the rat substantia
nigra. *J. Neurophysiol.* **103**, 3099–3114 (2010).

15. P. Durante, C. G. Cardenas, J. A. Whittaker, S. T. Kitai, R. S. Scroggs, Low-
Threshold L-type Calcium Channels in Rat Dopamine Neurons. *J. Neurophysiol.*
91, 1450–1454 (2004).

16. L. M. Alonso, E. Marder, Visualization of currents in neural models with similar
behavior and different conductance densities. *eLife* **8**, e42722 (2019).

17. Z. M. Khaliq, B. P. Bean, Dynamic, Nonlinear Feedback Regulation of Slow
Pacemaking by A-Type Potassium Current in Ventral Tegmental Area Neurons.
J. Neurosci. **28**, 10905–10917 (2008).

18. J.-M. Goaillard, E. Marder, Ion Channel Degeneracy, Variability, and Covariation
in Neuron and Circuit Resilience. *Annu. Rev. Neurosci.* **44**, 335–357 (2021).

19. A. Fyon, A. Franci, P. Sacré, G. Drion, Dimensionality reduction of neuronal
degeneracy reveals two interfering physiological mechanisms. *PNAS Nexus* **3**,
pgae415 (2024).

20. G. Drion, T. O'Leary, E. Marder, Ion channel degeneracy enables robust and
tunable neuronal firing rates. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E5361–E5370
(2015).

21. T. O'Leary, A. H. Williams, A. Franci, E. Marder, Cell Types, Network
Homeostasis, and Pathological Compensation from a Biologically Plausible Ion
Channel Expression Model. *Neuron* **82**, 809–821 (2014).

22. J. Roeper, Dissecting the diversity of midbrain dopamine neurons. *Trends*
Neurosci. **36**, 336–342 (2013).

23. M. H. Higgs, J. A. Jones, C. J. Wilson, M. J. Beckstead, SK and Kv4 Channels
Limit Spike Timing Perturbations in Pacemaking Dopamine Neurons. *eNeuro* **10**
(2023).

24. F. Philippart, Z. M. Khaliq, Gi/o protein-coupled receptors in dopamine neurons
inhibit the sodium leak channel NALCN. *eLife* **7** (2018).

25. D. E. Cobb-Lewis, L. Sansalone, Z. M. Khaliq, Contributions of the Sodium Leak
Channel NALCN to Pacemaking of Medial Ventral Tegmental Area and
Substantia Nigra Dopaminergic Neurons. *J. Neurosci.* **43**, 6841–6853 (2023).

26. K. B. Um, S. Hahn, S. W. Kim, Y. J. Lee, L. Birnbaumer, H. J. Kim, M. K. Park,
TRPC3 and NALCN channels drive pacemaking in substantia nigra
dopaminergic neurons. *eLife* **10**, e70920 (2021).

27. M. Puopolo, E. Raviola, B. P. Bean, Roles of Subthreshold Calcium Current and
Sodium Current in Spontaneous Firing of Mouse Midbrain Dopamine Neurons.
J. Neurosci. **27**, 645–656 (2007).

28. S. Nedergaard, J. A. Flatman, I. Engberg, Nifedipine- and omega-conotoxin-
sensitive Ca2+ conductances in guinea-pig substantia nigra pars compacta
neurones. *J. Physiol.* **466**, 727–747 (1993).

29. C. S. Chan, J. N. Guzman, E. Ilijic, J. N. Mercer, C. Rick, T. Tkatch, G. E.
Meredith, D. J. Surmeier, 'Rejuvenation' protects neurons in mouse models of
Parkinson's disease. *Nature* **447**, 1081–1086 (2007).

30. G. Drion, L. Massotte, R. Sepulchre, V. Seutin, How Modeling Can Reconcile
Apparently Discrepant Experimental Results: The Case of Pacemaking in
Dopaminergic Neurons. *PLOS Comput. Biol.* **7**, e1002050 (2011).

31. J. N. Guzman, J. Sánchez-Padilla, C. S. Chan, D. J. Surmeier, Robust Pacemaking in Substantia Nigra Dopaminergic Neurons. *J. Neurosci.* **29**, 11011–11019 (2009).

32. T. Z. Lu, Z.-P. Feng, NALCN: A Regulator of Pacemaker Activity. *Mol. Neurobiol.* **45**, 415–423 (2012).

33. J. Wang, M. Su, D. Zhang, L. Zhang, C. Niu, C. Li, S. You, Y. Sang, Y. Zhang, X. Du, H. Zhang, The cation channel mechanisms of subthreshold inward depolarizing currents in the mice VTA dopaminergic neurons and their roles in the chronic-stress-induced depression-like behavior. *eLife* **12**, RP88319 (2024).

34. E. G. Lakatta, D. DiFrancesco, What keeps us ticking: a funny current, a calcium clock, or both? *J. Mol. Cell. Cardiol.* **47**, 157–170 (2009).

35. A. G. Torrente, P. Mesirca, P. Neco, R. Rizzetto, S. Dubel, C. Barrere, M. Sinegger-Brauns, J. Striessnig, S. Richard, J. Nargeot, A. M. Gomez, M. E. Mangoni, L-type Cav1.3 channels regulate ryanodine receptor-dependent Ca2+ release during sino-atrial node pacemaker activity. *Cardiovasc. Res.* **109**, 451–461 (2016).

36. P. Mesirca, J. Alig, A. G. Torrente, J. C. Müller, L. Marger, A. Rollin, C. Marquilly, A. Vincent, S. Dubel, I. Bidaud, A. Fernandez, A. Seniuk, B. Engeland, J. Singh, L. Miquerol, H. Ehmke, T. Eschenhagen, J. Nargeot, K. Wickman, D. Isbrandt, M. E. Mangoni, Cardiac arrhythmia induced by genetic silencing of 'funny' (f) channels is rescued by GIRK4 inactivation. *Nat. Commun.* **5**, 4664 (2014).

37. J. Bezanson, A. Edelman, S. Karpinski, V. B. Shah, Julia: A Fresh Approach to Numerical Computing. *SIAM Rev.* **59**, 65–98 (2017).

38. A. Rakovic, D. Voß, F. Vulinovic, B. Meier, A.-K. Hellberg, C. Nau, C. Klein, E. Leipold, Electrophysiological Properties of Induced Pluripotent Stem Cell-Derived Midbrain Dopaminergic Neurons Correlate With Expression of Tyrosine Hydroxylase. *Front. Cell. Neurosci.* **16** (2022).

39. P. Stathakos, N. Jimenez-Moreno, L. Crompton, P. Nistor, M. A. Caldwell, J. D. Lane, "Imaging Autophagy in hiPSC-Derived Midbrain Dopaminergic Neuronal Cultures for Parkinson's Disease Research" in *Autophagy: Methods and Protocols*, N. Ktistakis, O. Florey, Eds. (Springer, New York, NY, 2019; https://doi.org/10.1007/978-1-4939-8873-0_17), pp. 257–280.

40. J. Louradour, O. Bortolotti, E. Torre, I. Bidaud, N. Lamb, A. Fernandez, J.-Y. Le Guennec, M. E. Mangoni, P. Mesirca, L-Type Cav1.3 Calcium Channels Are Required for Beta-Adrenergic Triggered Automaticity in Dormant Mouse Sinoatrial Pacemaker Cells. *Cells* **11**, 1114 (2022).

41. F. Philippart, G. Destreel, P. Merino-Sepúlveda, P. Henny, D. Engel, V. Seutin, Differential Somatic Ca2+ Channel Profile in Midbrain Dopaminergic Neurons. *J. Neurosci.* **36**, 7234–7245 (2016).

42. G. Destreel, V. Seutin, D. Engel, Subsaturation of the N-methyl-D-aspartate receptor glycine site allows the regulation of bursting activity in juvenile rat nigral dopamine neurons. *Eur. J. Neurosci.* **50**, 3454–3471 (2019).

43. S. Ringlet, Z. Motta, L. Vandries, V. Seutin, K. Jehasse, L. Caldinelli, L. Pollegioni, D. Engel, Glycine-gated extrasynaptic NMDARs activated during glutamate spillover drive burst firing in nigral dopamine neurons. *Prog. Neurobiol.* **249**, 102773 (2025).

44. L. Borgs, E. Peyre, P. Alix, K. Hanon, B. Grobarczyk, J. D. Godin, A. Purnelle, N. Krusy, P. Maquet, P. Lefebvre, V. Seutin, B. Malgrange, L. Nguyen, Dopaminergic neurons differentiating from LRRK2 G2019S induced pluripotent stem cells show early neuritic branching defects. *Sci. Rep.* **6**, 33377 (2016).

Acknowledgments

We thank our colleagues from the laboratories of Neuroengineering and Neurophysiology for their valuable input and help, especially L.Vandries for the histology of human mDANs. We thank our colleagues Pr. J-F Liégeois and Dr. R. Vitello for providing the mesylate salt of XG. We thank B. Fischer, J. Sonntag, S. Betz, G. Amrhein and T. Wulf for technical assistance in this project.

Funding

This work was supported by the Belgian Government through the Federal Public
Service (FPS) Policy and Support (BOSA) grant NEMODEI (NEuroMOrphic Design of
Embodied Intelligence) (AFy, JB, AFr, GD, KJ) ; Fonds National de la Recherche
Scientifique (FNRS, Belgium) grants ASP-REN40024838 (AFy), J.0148.19 and
T.0188.22 (VS). AFy is a Postdoctoral Researcher of FNRS. Liège University ;
"Strategic opportunity grant" R.DIVE 1142-J-P-A-02 (OP, VS) ; Fondation Léon
Frédéricq (KJ) ; Deutsche Forschungsgemeinschaft (DFG) - CRC1451 (JR).

Author contributions

Conceptualization: AFy, VS, GD & KJ; Methodology: AFy, SR, JB, AFr, KJ;
Investigation: AFy, OP, NS, PM, & KJ; Visualization: AFy & KJ; Funding acquisition:
VS, GD & KJ; Project administration: VS, GD & KJ; Supervision: JR, MM, VS, GD &
KJ; Writing – original draft: AFy & KJ; Writing – review & editing: all authors

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

All data and codes will be available upon publication.

Figure legend

Fig. 1. XG inhibits slow pacemaker cells. (**A**) Pacemaking in rat mDANs is inhibited upon (2 mM) XG application. (**B**) Summary data of the effect of the application of 2 mM XG on the firing frequency. XG abolished the firing (1.94 ± 0.18 Hz in control vs 0.02 ± 0.001 Hz in XG, n = 8, p = 0.0078, Wilcoxon test). (**C**) XG-sensitive current isolated from rat mDANs. Inward current at -60 mV is the result from rundown of T-type calcium channels that was observed in one cell. (**A-C**) were adapted from *Jehasse et al. 2021* (*6*). (**D**) Pacemaking in human mDANs is inhibited upon (600 μM) XG application. (**E**) Summary data of the effect of the application of 600 μM XG on the firing frequency (2.28 ± 0.44 Hz in control vs 0.12 ± 0.05 in XG, n = 9, p = 0.0039, Wilcoxon test). (**F**) XG-sensitive current isolated from human mDANs (**G**) Pacemaking in mouse SAN cells is inhibited upon (300 μM) XG application. (**H**) Summary data of the effect of the application of 600 μM XG on the firing frequency, in beats per minute (bpm). XG strongly reduces the firing (166.50 ± 9.11 bpm in control vs 24.55 ± 10.10 bpm in XG, n = 11, p = 0.001, Wilcoxon test). (**I**) XG-sensitive currents isolated from mouse SAN cells. (**J**) Overlay of XG-sensitive current from rat (green) and human mDANs (orange), and SAN cells (black) shows similarities in voltage activation and current intensity. Dashed lines correspond to 0 mV for voltage traces and 0 pA for current traces

Fig. 2. Fast activating pacemaker conductance allows slow pacemaking in a mDAN conductance-based model. (**A**) Currentscape analysis reveals that, although its contribution seems small in comparison to other conductances, g_{pace} is crucial to generate slow pacemaking. (**B**) Simulated pacemaking activity in our adjusted pacemaking model (green) with I_{pace} dynamic below in red, showing a small amplitude during the interspike interval as previously observed (*6*). Bottom panel corresponds to total current amplitude from our adjusted model with (green) and without (red) the addition of g_{pace}. The total current observed is similar to what has been calculated (*10*). (**C**) Total charge transfer of each conductance during the interspike interval (ISI) and the full period of the simulation. Charge transfer was calculated by measuring the area under the curve in (**A**). (**D**) Heatmap analysis showing the multiplicative time constant of g_{pace} activation relative to g_{Na} according to the conductance density in which our adjusted model generates slow pacemaking.

Fig. 3. The XG-sensitive conductance (g_{pace}) modulates pacemaking in *SNc* DANs. (**A**) *Ex vivo* pacemaking in a mouse *SNc* (orange shade) DAN. (**B**) Silencing of a *SNc* DAN upon injection of negative g_{pace} (upper panel). Summary data of the effect of negative g_{pace} on the firing frequency (lower panel). Negative g_{pace} abolished pacemaking at -0.3 nS (1,54 Hz (interquartile range: 1,193 - 1,864 Hz) in ctrl vs. 0 Hz (interquartile range: 0) with g_{pace} = - 0,3 nS, n = 20, N = 3, p = <0,0001, Wilcoxon test). (**C**) Excitation and overexcitation (depolarization block) of a *SNc* DAN upon positive g_{pace} (upper panel). Summary data of the effect of positive g_{pace} on the firing frequency (lower panel). g_{pace} increased the firing frequency (1,48 ± 0,66 Hz in ctrl vs. 2,26 ± 0,91 Hz with g_{pace} = 0,3 nS, n = 17, N =3, p = <0,0001, paired t test). (**D**) Rescue of pacemaking after preincubation with (2 mM) XG and complete cessation of pacemaking. Firing pattern and g_{pace} value upon rescue display broad large variability (right). Time frame of steady state g_{pace} (bold red line). Time frame of manual deactivation of g_{pace} (discontinuous red line).

Fig. 4. Assessment of the conductances contribution in slow pacemaking. (**A**) Currentscape analysis from a minimal model including g_{Na}, g_{Kd}, g_{LNS} and g_{pace} shows that it is possible to generate slow pacemaking, similarly to what we observed *ex vivo* in the absence of other conductances (*6*), and that g_{pace} contribution is maximal during the interspike interval. (**B**) Heatmap analysis showing the multiplicative time constant of g_{pace} activation relative to g_{Na} according to the conductance density in which our minimal model generates slow pacemaking. (**C**) Robustness of slow pacemaking in response to noise (in red). In the minimal model (blue), current noise leads to irregular spiking, while the complete model (teal), current noise injection does not alter pacemaking regularity (left panel). Right panel corresponds to the effect of increasing the standard deviation (SD) of noise current to the coefficient of variation (CV) of the interspike interval (ISI). The addition of g_H, g_{KSK} and g_{KA} increases the robustness of slow pacemaking. In order to activate g_{KSK}, g_{LCa} was added. (**D**) Effect of the other conductances on the firing frequency upon continuous current injection. Left panel shows example of the firing frequency of the minimal (blue) and complete (teal) upon current injection of 20 pA. Right panel shows the effect of the combination of several conductances on the FI relationship. The results related to g_{KSK} addition only are located behind the complete and combination of g_{KSK} and g_H data points.

Supplementary figures

Fig. S1. Identification of human mDANs. (**A**) Top panel shows that application of

dopamine on mDANs triggers silencing of pacemaking by hyperpolarization due the

activation of DA autoreceptors (*44*), as observed in rodent mDANs (*41*). Bottom panel

shows an example traces of I_H in human mDANs. (**B**) Evolution of the expression of

TH, showing differentiation of neurons into mature mDANs.

 Fig. S2. Slow pacemaking silenced by XG is not due to an off-target effect on the

 main voltage-gated channels of SAN cells. (**A**) XG does not interact with neuronal

 and cardiac HCN currents. (**B**) XG interacts neither with delayed rectifier nor A-type

 potassium channels. (**C**) XG does not decrease L-type and T-type calcium currents of

 SAN cells. Graphs of mDANs were adapted from *Jehasse et al. 2021* (*6*).

Fig. S3. Adjustment of current conductance-based model of a mDAN. (**A**) Simulated pacemaking and currentscape analysis from the model of *Yu et Canavier 2015* (*12*). (**B**) Blue dots represent the firing frequency relative to current injections from the model of Yu et Canavier, showing a type I excitability similar to in vitro mDANs. Grey dots represent the firing frequency relative to current injections from the adujsted model of *Yu et Canavier*, changing the excitability type from I to II. Adding g_{pace} to the adjusted model (green dots) shifts the excitability of the neuron from type II to type I. (**C**) Activation curves from Na$_v$ (top) and L-type Ca$_v$ (bottom) channels. Blue curves correspond to the parameters used in the model of *Yu et Canavier*, while grey curves corresponds to our adjustment to physiological values as observed in mDANs for Na$_v$ (*14*) and L-type Ca$_v$ channels (*15*). (**D**) Fitting curve (red) based on XG-sensitive current (green) to generate g_{pace}. (**E**) Circuit for conductance-based model with nonlinear conductances and the pacemaking conductance (in red).

754 **Fig. S4. Additional heatmaps generated from other random conductance values**

755 **for all channels.** Overall, we observe a wide range of densities from g_{pace} in order to

756 produce slow pacemaking, with fast kinetics.

Fig. S5. Degenerate populations and PCA analysis of the adjusted model. (**A**) Scatter plots of degenerate populations for the original (blue) and adjusted (green) models in selected conductance subspaces (g_{Na}, g_{CaL} and g_{Kd}), highlighting that our model allows more degeneracy. (**B**) Violin plots of all conductances for both populations. (**C**) Summary of PCA analysis on the adjusted degenerate population (green in **A** and **B**). From left to right: bar plot of explained variance for each component, scree plot, projection onto the PC1–PC2 plane, and scatter plot of PC1/PC2 loadings.

Fig. S6. Voltage-dependence of g_{pace} **is required to generate steady and slow pacemaking.** (**A**) Replacement of g_{pace} by a linear depolarizing conductance in our adjusted model such as g_{NALCN} does not result in slow pacemaking. (**B**) Firing frequency related to the density of g_{NALCN} shows the restricted range of g_{NALCN} density where pacemaking occurs. (**C**) Firing frequency – current relationship curve shows that low density of g_{NALCN} leads to type II excitability, which does not correspond to mDANs excitability, whereas higher density of g_{NALCN} leads to type I fast pacemaking, also not corresponding to mDANs slow firing frequency.