Post-infection pig and ferret antisera show similar antigenic profiles for human
influenza A(H1N1pdm09) viruses

Ruth Harvey[1*], Tiphaine Cayol[2*], Basudev Paudyal[2*], Alice Lilley[1], Christine Carr[1], Catherine
Hatton[2], Emily Briggs[2], Rodney S. Daniels[1], Samuel Richardson[2], Thomas P. Peacock[2], Nicola
Lewis[1#], Ian Brown[2#], John W. McCauley[1#], Elma Tchilian[2#]

[*]These authors contributed equally

[#]These authors share senior authorship.

[1]The Worldwide Influenza Centre, The Francis Crick Institute, London, UK

[2]The Pirbright Institute, Pirbright, UK

Abstract

Background: Monitoring antigenic drift in human influenza A viruses is essential for vaccine strain selection and ensuring protection against circulating strains. Antigenic drift is traditionally assessed using ferret antisera, which provide monospecific responses, and human vaccinee sera, which reflect exposure to multiple antigens. In this study we evaluated the pig as an alternative source of antisera to study antigenic drift compared to immune responses in ferrets and humans. We included seasonal influenza A(H1N1pdm09) human viruses that had shown different antigenic characteristics when using ferret or human antisera.

Methods: Pairs of pigs were inoculated with six human A(H1N1)pdm09 viruses circulating between 2019 and 2023, a period of marked antigenic drift. Pig and ferret antisera were analysed by hemagglutination inhibition (HI) and virus neutralization (VN) assays.

Results: Pigs were successfully infected with all strains, shedding virus and producing antibody responses, confirming their susceptibility to human influenza A viruses. Antigenic reactivity of pig antisera was qualitatively comparable to ferret antisera in both HI and VN assays, although maximum homologous antibody titres were significantly higher in ferrets. The antisera raised against viruses in circulation in 2019 and before, exemified by A/Guangdong-Maonan/SWL1536/2019, clade 5a.1, were clearly differentiated by both ferret and pig antisera from those in clade 5a.2 and its derivatives that became predominant.

Conclusions: Ferrets and pigs showed comparable responses and both distinguished clade 5a.1 from clade 5a.2. However, neither model recognised antigenically drifted variants from 2019–2022, including subclades 5a.2-C, 5a.2a-C.1/C.1.9, and .5a.2a.1-C.1.1/D, which were distinguishable using human post-vaccination antisera.

Introduction

Human seasonal Influenza A viruses continually evolve genetically and antigenically necessitating constant monitoring and frequent updating of seasonal influenza vaccine strains. The Global Influenza Surveillance and Response System (GISRS), coordinated through the World Health Organisation (WHO), monitors the genetic and antigenic relatedness of currently circulating viruses, and each year the WHO makes separate recommendations of virus components for inclusion in seasonal influenza vaccines for both the Southern and Northern Hemisphere (SH and NH) [1].

Underpinning the vaccine strain recommendations are phenotypic data characterising the antigenic properties of currently circulating influenza strains relative to previously recommended Candidate Vaccine Viruses (CVVs) and to other reference viruses, including viruses chosen to be representative of emerging genetic groups. Routinely the ferret is used as a small animal model for antibody responses to human seasonal influenza and reference antisera raised against CVVs and other key reference viruses and used throughout each influenza season in hemagglutination inhibition (HI) and virus neutralisation (VN) assays to assess the antigenic properties of contemporary influenza virus strains. These ferret antisera antigenic analyses are supplemented by human serological response analyses using pre- and post-vaccination serum panels.

Periodically disparities have been observed between the relative ability of ferret and human immunological responses to similarly discriminate amino acid substitutions in hemagglutinin (HA) resulting in antigenic evolution, particularly with the A(H1N1pdm09) human influenza viruses. For example, while the ferret antibody responses to 2019 CVVs aligned with the human responses when assessed by HI assays, there were discrepancies with CVVs from 2021 such as A/Sydney/5/2021 (Syd21), where human serology indicated significant antigenic drift commensurate with a need to update the vaccine strain whereas ferret antisera did not [2, 3].

Pigs are an important host for influenza A viruses and are readily infected by the same H1N1 and H3N2 virus subtypes as humans with periodic transmission in both directions [4]. They also serve as a significant biomedical model for studying human disease [5]. Pigs have a longer life span, and are genetically, immunologically, physiologically, and anatomically more like humans than small laboratory animals [6, 7]. Additionally, pigs exhibit a comparable distribution of sialic acid receptors in the respiratory tract to humans [8]. Indeed we have previously shown that pig immune responses following infection with A(H1N1pdm09) influenza virus were similar to those induced in humans [9] thereby providing a model of human influenza infection [10].

Here we tested whether the pig immunological response was able to discriminate
between human A(H1N1pdm09) influenza viruses that might be distinguished by either post-
infection ferret antisera or post-vaccination human sera.

Results

Virus infection of pigs and nasal shedding. The panel of study viruses selected were
representative of viruses in circulation between 2019 and 2023 (**Table 1**). Over this period
some groups of viruses were differentiated by both post-infection ferret antisera and post-
vaccination human sera while others had previously shown detectable differences in response
in human serology studies but not when tested against post-infection ferret antisera. Six
groups of two pigs each were infected intranasally with one of the study viruses possessing
the HA amino acid substitutions indicated (**Table 1**) and sampled as described in Materials
and Methods and illustrated in **Fig 1A.** All animals were productively infected as determined
by nasal shedding of virus. The infection kinetics showed that mean shedding following
infection with all viruses generally peaked at 4-days post-infection (DPI) (Syd21 infected pigs
peaked at 2-DPI) and had been cleared in all pigs by 7-DPI (**Fig. 1B**). Virus whole genome
sequencing of the input and 4-DPI nasal swabs showed all shed viruses to be unchanged at
the consensus level from the infecting viruses, indicating lack of genetic mutations associated
with pig infection in a single animal infection cycle (data not shown).

HI responses of ferret and pig antisera. Antibody titres in HI showed differences in the
magnitude of responses between ferret and pig antisera against the panel of study viruses
(**Table 2A**). Ferret antisera consistently exhibited high titres of 5120 against the homologous
viruses used for infection, indicating a strong antigen-specific response. In contrast, pig
antisera generally demonstrated lower homologous titres, ranging from 160 to 640, depending
on the specific virus and individual animal.

Cross-reactivity patterns were broadly similar between ferret and pig antisera. The
ferret antiserum raised against A/Guangdong-Maonan/SWL1536/2019 (GM19;subclade 5a.1-
B) showed good recognition of the homologous antigen but very limited recognition of viruses
of subclades 5a.2-C A/Wisconsin588/2019 (Wsc19), 5A.2a-C.1 (Syd21), 5A.2a-C.1.9
A/Lisboa/188/2023 (Lis23), 5a.2a.1-D A/Victoria/4897/2022 (Vic22) and 5a.2a.1-C.1.1
A/Wisconsin/67/2022 (Wsc22). Correspondingly, the antisera raised against Wsc19, Syd21,
Lis23, Vic22 and Wsc22 all showed high HI titres for the homologous viruses and poor
recognition of GM19. However, each of these antisera recognised viruses in subclades 5a.2-
C, 5a.2a-C.1, 5a.2a-C.1.9, 5a.2a.1-D and 5a.2a.1-C.1.1 at titres within 2-fold of the
homologous titres, most at titres equal to the homologous titre.

While there were some differences in the magnitude of HI antibody titres between
individual pigs using the same infecting virus, two-fold differences were not considered
significant since the antigenic profiles were consistent when comparing serum reactivity
between homologous and heterologous viruses for each pair of pigs. One exception was a 4-
fold variation in the homologous titres for Vic22 infected pigs (pigs 61 & 62) but titres to
heterologous viruses showed identical reactivity profiles (**Table 2A**). Like the ferret antiserum
raised against GM19, the two pig antisera raised against GM19 recognised the homologous
virus well, at HI titres of 320, but recognised the heterologous viruses less well, although pig
53 and 54 antisera recognised Wsc19, Syd21 and Lis22 at titres within 4-fold lower than the
homologous titres. In contrast, all antisera raised against the other five study viruses
recognised GM19 poorly, but showed good recognition of the homologous and heterologous
viruses, with a single exception for pig 66 (infected with Wsc22) antiserum which recognised
Wsc19 at a titre 4-fold reduced compared to the homologous titre.
123
VN of ferret and pig antisera. Ferret antisera raised against GM19 showed a similar clear
differentiation in VN assays between the homologous virus and the other five study viruses,
as seen by HI assay (**Table 2B**). Moreover, the antisera raised against the other five study
viruses all recognised GM19 poorly. Nevertheless, these antisera showed more variable
recognition of the heterologous viruses: those raised against Lis23 (5a.2a-C.1.9), Wsc22
(5a.2a.1-C.1.1) and Vic22 (5a.2a.1-D) all recognised Wsc19 (5a.2-C) at titres >4-fold reduced
compared to homologous titres, as was the case for Lis23 antiserum with Vic22. Reductions
of 2-4-fold were seen for Wis19 antiserum with Vic22 and Vic22 antiserum with Lis23.
Within pig antisera pairs homologous VN titres ranged from equivalence, for GM19, up
to 5-fold (Wsc22) for four pairs while for Vic22 (pigs 61 and 62) 21-fold difference was seen
(**Table 2B**). Neutralization titres were consistently higher in ferret antisera for all viruses except
Syd21 which had homologous titres of 2358 in ferret and 7680 or 2648 in pigs. Further, the
pig antisera showed an even greater variation in recognition patterns in the VN assay than
was seen in the HI assay. However, heterologous titres were inconsistent between pig pairs
in their profile of reactivity when directly comparing to homologous titres for all infecting
viruses.
Again, pig antisera differentiated GM19 (clade 5a.1) from the five viruses in clade 5a.2
and derivatives thereof, antisera raised against them recognising GM19 poorly. Some
neutralization of the heterologous viruses was seen by the antisera raised against GM19, five
of the ten heterologous titres being within 4-fold of the homologous titres, one within 2-fold. Of
the 40 heterologous titres associated with antisera raised against the other five study viruses,
seven were >4-fold reduced: five related to Vic22 with antisera raised against Wsc19 (pig 63,
with a low homologous titre), Syd21 (both pigs) and Lis23 (both pigs), and two to Wsc19 for

antisera raised against Lis23 (pig 54) and Wsc22 (pig 66). Notably, Wsc19 was also
recognized poorly by ferret antisera raised against Lis23 and Wsc22.

Overall, antisera raised in both pigs and ferrets against viruses in clades 5a.2, 5a.2a
and 5a.2a.1 recognised the 5a.1 virus, GM19, poorly in both HI and VN assays. The antiserum
raised against GM19 in ferrets recognised the 5a.2, 5a.2a and 5a.2a.1 viruses poorly in both
assays, but the antisera raised in pigs recognised some of the viruses from the heterologous
subclades at variable titres, many at titres within 4- to 8-fold of the titres for the homologous
antigen. Antisera in ferrets and pigs raised against 5a.2, 5a.2a and 5a.2a.1 viruses recognised
viruses in the same clades well, all at titres within 4-fold of the homologous titres, all but one
within 2-fold, in HI assays. There was more variation seen in the VN assay but there was no
observed differentiation between antigens in individual clades 5a.2, 5a.2a and 5a.2a.1.

Human Serology. Three serum panels taken from adult volunteers (18 to 64 years of age)
over three consecutive vaccination campaigns were analysed: 2020 ahead of the northern
hemisphere (NH) 2020/21 season, 2021 for the NH 2021/22 season, and 2022 for the NH
2022/23 season. Volunteers were immunised with inactivated vaccine antigens corresponding
to egg-propagated GM19 (clade .5a.1) in 2020, and cell culture-propagated Wsc19 (clade
5a.2) in 2021 and 2022. These antisera were used in HI assays with the homologous vaccine
viruses and cell culture-propagated Wsc19 for the 2020 panel, and cell culture-propagated
cultivars of Syd21 (subclade 5a.2a-C.1) and Wsc22 (subclade 5a.2a.1-C.1.1) for the 2021 and
2022 panels.

Non-parametric analysis of the paired post-vaccination HI titres using a Wilcoxon
matched-pairs signed rank test showed that the human sera differentiated GM19 from Wsc19
(P< 0.0001) in the 2020 vaccinees. The sera from the 2021 vaccinees clearly differentiated
Wsc22 from Wsc19 (P=0.0005) but not Syd21 (P=0.083), whilst the sera from the 2022
vaccinees differentiated both Syd21 (P=0.0009) and Wsc22 (P=0.0001) from Wsc19 (**Fig 2**).

Analysing individual serum responses showed that of the 22 volunteers vaccinated
with egg-propagated GM19 eight recognised Wsc19 at titres reduced 4-fold or more. Of the
29 volunteers vaccinated with cell culture-propagated Wsc19 in 2021, 12 (41%) recognised
Syd21 and 22 (76%) recognised Wsc22 at titres reduced 4-fold or more while of the 25
volunteers vaccinated in 2022, 9 (36%) recognised Syd21and 13 (45%) recognised Wsc22 at
titres reduced 4-fold or more.

These results indicate that, like post-infection ferret and swine antisera, post-
vaccination human sera (although with some degree of variation) are able to differentiate the
clade 5a.1 vaccine virus from Wsc19 (clade 5a.2). Post-vaccination serum panels as a whole,
also differentiate the vaccine virus Wsc19 from Wisc22 (subclade 5a.2a.1-C.1.1) and in one
of the two seasons differentiated Wsc19 from Syd21 (subclade 5a.2a-C.1) unlike the ferret

and pig antisera. These results with human sera are consistent with published results from
post-vaccination human serum panels used in vaccine strain recommendations for the
Southern Hemisphere 2023 and the Northern Hemisphere 2023/24 influenza seasons [2, 3].
Nevertheless, individual sera differed in their discrimination between viruses in the different
subclades.
189
Discussion
Post-infection ferret antisera and post-vaccination human sera are usually used to define the
antigenic characteristics of human influenza viruses as they evolve. However, for some recent
seasonal influenza viruses there has been a disparity in the recognition of viruses between
human and ferret antisera which has impacted on human vaccine strain selection, notably so
for A(H1N1pdm09) viruses [2, 11-13]. Therefore, this study aimed to evaluate the immune
response in pigs following infection with A(H1N1pdm09) human influenza viruses, to compare
their recognition of recent A(H1N1pdm09) viruses with those of ferret and human antisera,
and to investigate whether post-infection pig antisera can enhance the antigenic analysis of
human influenza A(H1N1pdm09) viruses.
The overall study findings revealed that pigs could readily be infected with seasonal
human A(H1N1pdm09) influenza viruses. This was not unexpected and confirmed that pigs
remain susceptible to contemporary human seasonal viruses descended from the 2009 swine
origin pandemic virus and support ongoing detections of virus spillover from humans to pigs
[14, 15]. Broadly, pig immune responses produced consistent antigenic profiles directly
comparable to those of ferrets; although, we obtained some altered characteristics when
comparing neutralization profiles for ferret and pig antisera between homologous and
heterologous viruses, but these were not correlated with subclade.
The post-infection antisera from ferrets and pigs and the post-vaccination human sera
distinguished GM19 (clade 5a.1) from Wsc19 (clade 5a.2). The HA of this pair had the
substitutions K130N, N156K, L161I, A187D, E189Q and V250A in HA1 and E179D in HA2
(**Table 1**), and four of these seven substitutions can be mapped to antigenic sites: Sa (K130N
and N156K) and Sb (A187D and E189Q). Five of these HA1 substitutions (not E189Q) were
conserved in the remaining four members of the virus panel. The antisera from both ferrets
and pigs did not differentiate Wsc19 from Syd21 (subclade 5a.2a-C.1) in HI or VN assays,
with Syd21 having HA1 substitutions K54Q, D94N, A186T, Q189E (reversion), T216A, E224A,
R259K and K308R (**Table 1**). These antisera also did not distinguish Vic22 (subclade 5a.2a.1-
D), Lis23 (subclade 5a.2a-C.1.9) and Wsc22 (subclade 5a.2a.1-C.1.1) from Wsc19, despite
viruses in these subclades differing in HA1 amino acid sequence at several sites including the
substitutions T120A, P137S, K142R, K169Q, D260E and T277A (**Table 1**). Importantly
antisera from both animal species did not clearly discriminate between Wsc19 and Wsc22 in

contrast to the post-vaccination cohorts of human sera in 2021 and 2022, which showed a reduction in human sera reacting to Wsc22 (**Fig 2** and [2, 3]). This pair of viruses differed at nine HA1 residues K54Q, P137S, K142R, A186T, Q189E, E224A, R259K, D260E and T277A. The altered recognition by the post-vaccination sera resulted in a recommendation to update the cell-based and egg-propagated vaccine strain components for the NH 2023/24 season [3].

Although the pattern of recognition of ferret and pig antisera were similar, direct comparisons of the immune responses showed that while the pairs of pigs in this study had comparable homologous and heterologous HI titres they were lower than those of ferret (approximately 16-fold across all viruses, except for GM19, a virus antigenically very distinct from the other viruses examined) which reduces the dynamic range of the assays. When homologous titres in VN assays are compared, generally pig antisera titres were lower than those of ferret, showing pig-specific reductions of 12- to 211-fold, for all but antisera raised against Syd21 (**Table 2B**).

The utility of pig and ferret antisera in evaluating responses to phenotypic evolution of swine influenza viruses has recently been compared [17]. In that study swine sera were produced following intranasal infection and subsequent intramuscular immunisation with the same virus dose formulated with adjuvant, whereas ferret sera were obtained after intranasal infection only. Despite these differences in immunisation/infection regimes, the breadth and relative magnitude of the antibody responses were equivalent in both hosts, as assessed by HI and VN.

We have previously demonstrated that monoclonal antibodies (mAbs) derived from A(H1N1pdm09)-infected pigs recognized the same two major immunodominant HA sites, Sa (residues 160 and 163) and Ca (residue130) [9]. In 2014, A(H1N1pdm09) viruses acquired antigenic drift through HA1 K163Q substitution which was not detected using ferret antisera [11, 18-20]. Whether mAbs generated from the infected pigs in this study would reveal the fine specificity of epitopes in Wsc19 and Wsc22 remains to be determined. By 2016, the dominant A(H1N1pdm09) viruses circulating had acquired a NQS glycosylation motif through HA1 S162N substitution, effectively masking the Sa antigenic site.

These results reaffirm the continued importance of the interface between human and pigs for ongoing influenza virus evolution and potential antigenic divergence whilst retaining interspecies transmissibility. The genetic data obtained from viruses prior to infection and those shed four days after infection showed retention of consistent human-derived virus genetic characteristics, supporting maintenance of similar viruses in pigs albeit without defining transmissibility between pigs. Given the high susceptibility of pigs to human A(H3N2) influenza viruses and the maintenance of such viruses many years after the antigenic variants have disappeared from humans [21], due to genetic drift, there is also merit in assessing

comparative data between ferrets and pigs to discriminate antigenic drift leading to A(H3N2) vaccine strain changes.

This is the first study directly comparing ferret and pigs as a host to define antigenic reactivity and variation in human A(H1N1pdm09) influenza viruses. We were able to show little variation qualitatively between the two animal hosts when using post-infection antisera to discriminate between viruses. Therefore, pigs could offer a qualified alternative for measuring antigenic variation in human influenza A viruses. This study also broadens the scope of pigs to serve as an animal model for human influenza for evaluating immune responses and vaccine studies as one of the milestones for improved influenza vaccine research [10].

Materials and Methods

Virus propagation. Viruses provided by the Francis Crick Institute (Table 1) were propagated in Madin Darby Canine Kidney (MDCK) cells. MDCK were cultured in T75 flasks with Dulbecco's Modified Eagle's Medium (DMEM; Sigma Aldrich), 10% fetal bovine serum (FBS; Gibco) and 1% Penicillin and Streptomycin (PS, Sigma Aldrich). Cells were inoculated with 2 ml of a seed virus diluted in DMEM+PS and incubated for 30 minutes at room temperature before adding 10 ml of DMEM+PS+2µg/ml (TPCK)-treated trypsin and incubated for two to three days at 37°C with 5% CO_2. The gene sequence of each virus stock was confirmed before use.

Influenza infection of pigs. Twelve 6-8 weeks old, outbred Landrace x Large White pigs (6 female and 6 male) were obtained from a commercial high-health status herd. The pigs were screened by ELISA for the absence of serum IgG titres against A(H1N1pdm09) and A(H3N2) virus. For each study virus two pigs, with one male and one female randomly divided per group, were infected. After a week of acclimatization the two pigs in each study group were inoculated intranasally using a mucosal atomization device (MAD, Medtree, UK) with 2 ml (1 ml per nostril) with either A/Guangdong Maonan/SWL1536/2019 (GM19; 1.8 x 10^5 PFU/pig) or A/Lisboa/188/2023 (Lis23; 5 x 10^6 PFU/pig) or A/Sydney/5/2021 (Syd21; 8 x 10^6 PFU/pig) or A/Victoria/4897/2022 (Vic22; 4 x 10^6 PFU/pig) or A/Wisconsin/588/2019 (Wsc19; 1 x 10^6 PFU/pig) or A/Wisconsin/67/2022 (Wsc22; 1.5 x 10^6 PFU/pig) (**Fig. 1A**). At 14 days post-infection (DPI), pigs were humanely euthanized with an overdose of pentobarbital sodium anaesthetic and clotted blood and bronchoalveolar lavage fluid (BAL) was collected to assess immune responses to the viruses.

Sampling of study pigs. Nasal swabs (NS) (two per nostril) were taken from pigs four days before infection, and 2-, 4- and 7-DPI to assess virus shedding. The swabs were placed into 2 ml of virus transport medium comprising tissue culture medium 199 (Sigma-Aldrich)

supplemented with 25 mM 4-(2-hydroxyethyl)-1 piperazineethanesulfonic acid (HEPES), 0.035% sodium bicarbonate, 0.5% bovine serum albumin, penicillin 100 IU/ml, streptomycin 100 µg/ml, and nystatin 0.25 µg/ml, vortexed and centrifuged to remove debris, then stored at −80°C for subsequent virus titration. Blood samples were collected in BD Vacutainer SST™ serum separation tubes at the start of the study (prior to challenge) and at 14-DPI. BAL was collected from the entire left lung with 0.1% BSA in PBS. BAL samples were centrifuged at 500 x g for 5 min and the supernatant was aliquoted and frozen for further analysis.

Plaque assay. Virus titres in nasal swabs were determined by plaque assay on MDCK cells as described previously [22]. Plaques were visualized by staining the monolayer with 0.1% (v/v) crystal violet containing 20% methanol (Merck) and expressed as plaque forming units (PFU) per ml of nasal swab eluate.

Virus whole-genome sequencing Viral RNA extraction and influenza virus whole genome sequencing was performed as previously described [23]. Raw fastq sequences were then imported into Geneious Prime software and, trimmed to remove primers and aligned to whole genome reference sequences (GM19 [EPI_ISL_377080], Syd21 [EPI_ISL_19193801], Wsc19 [EPI_ISL_404527], Lis23 [EPI_ISL_18788764], Wsc22 [EPI_ISL_15928538], Vic22 [EPI_ISL_19192400]).

Ferret antisera production. Post-infection ferret antisera were generated by intranasally inoculating ferrets with 500 µl (250 µl per nostril) of virus. Two weeks after inoculation sera were collected.

Human serology panels. Pre- and two-week post-vaccination human sera were collected over three consecutive years from healthy volunteers working at the Francis Crick Institute. Inclusion criteria included the receipt of a seasonal vaccine, being employed at the Francis Crick Institute and consenting to give a pre- and post-vaccination blood sample. No details of previous vaccination/infection history were obtained. Samples were collected in BD Vacutainer SSTII tubes and centrifuged for 10 minutes at 2000rpm to separate red and white blood cells from the serum. The serum supernatants were transferred into separate tubes and heat inactivated at 56°C for 30 minutes before storage at -20°C.

Hemagglutination Inhibition assay (HI). Ferret and pig antisera were heat inactivated at 56°C for 30 minutes and treated with receptor destroying enzyme (RDE) (Denka Co.). RDE-treated sera were diluted to a working dilution of 1:20 in PBS with 0.1% azide.

Hemagglutination and hemagglutinin inhibition (HI) assays were performed according to
standard methods using suspensions of turkey RBCs (0.75% v/v) [24]. Four hemagglutination
units were used in all HI assays.

Virus neutralisation (VN). Viruses were titrated to determine the dilution at which 50% of
cells within a well of a 96-well plate were infected, these dilutions were used in the VN assay
as previously described [25].

Conflict of interest. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author contribution. ET, IB, NL, RH, RD, JM designed the study and obtained the funding; ET, IB, TC, BP, CH, EB designed and performed the pig animal experiments; TC, BP, CH, EB, SR, processed samples; RH, TC, BP, AL, CC, TP, acquired, analysed and interpreted the data; ET, IB, JM, RH, RD wrote the manuscript. All authors have read and approved the manuscript.

Acknowledgments. We are grateful to the animal staff at the APHA for providing excellent animal care. This work was supported by the UKRI Biotechnology and Biological Sciences Research Council (BBSRC) IAA award BB/X511134/1 Research at Pirbright was funded by BBSRC via the Pirbright Institute's Strategic Programme Grants (ISPGs) [BBS/E/PI/230002A; BBS/E/PI/230002B], BBSRC National Bioscience Research Infrastructure: High Containment and Low Containment Services and Science Platforms grants [BBS/E/PI/23NB0004, BBS/E/PI/23NB0003]. This work was supported by core funding from the Francis Crick Institute from Cancer Research UK (grant no. CC1114), the UK Medical Research Council (grant no. CC1114), and the Wellcome Trust (grant no. CC1114). SR, TP and IB are supported by the Biotechnology and Biological Sciences Research Council (BBSRC)/DEFRA 'FluTrailMap' consortium [BB/Y007298/1] and the UK Medical Research Council/Department for Environment, Food and Rural Affairs (Defra, UK) FluTrailMap-One Health consortium [MR/Y03368X/1]. TC was supported by Gates Foundation grant OPP1215550 (Pirbright Livestock Antibody Hub).

Ethics statement: The pig studies were approved by the ethical review processes at APHA and the Pirbright Institute in accordance with the UK Government Animal (Scientific Procedures) Act 1986 under Project License PP2064443. Collection of human pre- and post-vaccination serum samples was approved by The Francis Crick Institute Internal Ethics Review Committee, registration number: 2019 FC5.

Figure 1. Experimental pig infection and analyses of samples taken.

(**A**) Schematic of the experimental design is shown, two pigs each were intranasally infected with one of the A(H1N1pdm09) viruses shown in **Table 1**. Serum and nasal swabs were collected at indicated times with necropsy at 14-DPI. (**B**) Virus loads in nasal swabs at -4-, 2, 4- and 7-DPI were determined by plaque assay. Each data point represents the mean of the two pigs inoculated with the same virus and error bars show the standard deviation of that mean.

Figure 2. Post-vaccination HI titres of three serum panels from healthy adults.

Serum panels from three consecutive northern hemisphere (NH) vaccination campaigns: 2020 for 2020/21 season (in blue), 2021 for 2021/22 season (in green), and 2022 for 2022/23 season (in red). Each panel was tested against the A(H1N1pdm09) virus component of the vaccine for the season and test viruses as shown. Data is plotted on a $\log_2$ scale.

Table 1. Influenza A(H1N1pdm09) viruses used in this study.

Virus Name	Passage History$	Clade	Subclade	Vaccination seasons*	HA amino acid substitutions compared to previous vaccine strain⁋	
					HA1	HA2
A/Guangdong-Maonan/SWL1536/2019	C2/M2	5a.1	B	NH 2020/1 (E)	N/A	N/A
A/Wisconsin588/2019	C2/S1/M4	5a.2	C	SH 2021 to NH 2022/3 (C)	K130N, N156K, L161I, A187D, E189Q, V250A¥	E179D
A/Sydney/5/5/2021	M3/M4	5a.2a	C.1	SH 2023 (E & C)	K54Q, D94N, A186T, *Q189E*, T216A, E224A, R259K, K308R	
A/Lisboa/188/2023	S1/M3	5a.2a	C.1.9	N/A (reference virus)	*N94D*, T120A, K169Q	K58R, I91V, *D179E*, R182K
A/Wisconsin /67/2022	M2	5a.2a.1	C.1.1	NH 2023/4 to NH2025/6 (C)	*N94D*, P137S, K142R, *A216T*, D260E, T277A	E29D, I91V, N124H
A/Victoria/4897/2022	S2/M3	5a.2a.1	D	NH 2023/4 to NH2025/6 (E)	*N94D*, P137S, K142R, D260E, T277A	E29D, I91V, N124H

$ E = egg, C = cell (specifics unknown), S = SIAT-1, M = MDCK.

* NH = Northern Hemisphere, SH = Southern Hemisphere, E = Egg-based vaccines, C = Cell culture- and recombinant-based vaccines.

⁋ Substitutions relate to the previous vaccine virus in the order shown in the table. Those substitutions in italics and underlined indicate amino acid reversions to match a previous vaccine strain. All HA sequences have been submitted to GISAID: GD19 EPI3592700, Wsc19 EPI3571883, Syd21 EPI3363292, Lis23 EPI3363420, Wsc22 EPI3251687, Vic22 EPI3362351.

¥ A/Wisconsin/588/2019 had A215A/T/I/V polymorphism

N/A = Not Applicable.

Table 2. Virus neutralizing activity of ferret and pig antisera assessed by HI and VN.

A. HI Assay □ <4-fold ▮ 4-fold ▮ 8-fold ▮ >8-fold/Not recognized by antiserum

Viruses	Clade	Subclade	Passage	Infecting Virus Ferret/Pig number	A/Guangdong-Maonan/SWL1536/2019 F09/20	Pig 52	Pig 53	A/Wisconsin/588/2019 F35/20	Pig 63	Pig 64	A/Sydney/5/2021 F46/22	Pig 56	Pig 57	A/Lisboa/188/2023 F09/24	Pig 54	Pig 55	A/Wisconsin/67/2022 F17/23	Pig 65	Pig 66	A/Victoria/4897/2022 F05/23	Pig 61	Pig 62
REFERENCE VIRUSES																						
A/Guangdong Maonan/SWL1536/2019	5a.1	B	C2/MDCK2		5120	320	320	160	<40	<40	80	<40	<40	40	<40	<40	<40	<40	<40	<40	<40	<40
A/Wisconsin/588/2019	5a.2	C	C2/SIAT1/MDCK4		320	160	80	5120	160	320	5120	640	640	2560	320	320	5120	80	80	5120	320	80
A/Sydney/5/2021	5a.2a	C.1	MDCK3/MDCK4		160	80	80	5120	160	160	5120	640	640	5120	160	160	2560	80	160	5120	320	80
A/Lisboa/188/2023	5a.2a	C.1.9	SIAT1/MDCK3		160	80	80	5120	160	160	5120	640	640	5120	320	320	2560	80	160	5120	640	160
A/Wisconsin/67/2022	5a.2a.1	C.1.1	MDCK2		160	<40	40	5120	80	160	5120	640	640	5120	320	320	5120	160	320	5120	640	160
A/Victoria/4897/2022	5a.2a.1	D	SIAT2/MDCK3		80	<40	40	5120	160	160	2560	640	640	2560	320	320	5120	160	320	5120	640	160

B. VN Assay □ ≤2-fold ▮ >2 ≤ 4-fold ▮ >4 ≤ 8-fold ▮ >8-fold/Not recognized by antiserum

Viruses	Clade	Subclade	Passage	Infecting Virus Ferret/Pig number	A/Guangdong-Maonan/SWL1536/2019 F09/20	Pig 52	Pig 53	A/Wisconsin/588/2019 F35/20	Pig 63	Pig 64	A/Sydney/5/2021 F47/22	Pig 56	Pig 57	A/Lisboa/188/2023 F09/24	Pig 54	Pig 55	A/Wisconsin/67/2022 F17/23	Pig 65	Pig 66	A/Victoria/4897/2022 F05/23	Pig 61	Pig 62
REFERENCE VIRUSES																						
A/Guangdong Maonan/SWL1536/2019	5a.1	B	C2/MDCK2		10240	240	226	93	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40
A/Wisconsin/588/2019	5a.2	C	C2/SIAT1/MDCK4		259	59	103	10240	96	286	1760	2080	1329	4800	352	320	3584	74	120	2560	404	51
A/Sydney/5/2021	5a.2a	C.1	MDCK3/MDCK4		52	66	83	27762	68	676	2358	7680	2648	26713	1062	1114	7680	160	749	15360	2358	240
A/Lisboa/188/2023	5a.2a	C.1.9	SIAT1/MDCK3		73	123	110	17723	61	845	2211	7680	4306	31403	2520	1254	7680	174	1061	7680	1721	308
A/Wisconsin/67/2022	5a.2a.1	C.1.1	MDCK2		43	<40	76	23757	40	411	2181	7680	1707	19144	671	1323	14758	188	873	75619	4189	413
A/Victoria/4897/2022	5a.2a.1	D	SIAT2/MDCK3		<40	<40	<40	4622	<40	94	2048	1858	579	5120	201	297	7680	77	243	19402	1020	92

Assays, HI (**A**) and VN (**B**), were performed as described in Materials and Methods and the results are 'heat-mapped' in accordance with the respective keys for **A** and **B**.

References:

1. World Health Organisation *Global Influenza Programme Recommendations for influenza vaccine composition*. 2025 [cited 2025 16th September]; Available from: https://www.who.int/teams/global-influenza-programme/vaccines/who-recommendations.
2. World Health Organisation, *Weekly Epidemiological Record*. Weekly Epidemiological Record, 2022. **97**(43): p. 537-566.
3. World Health Organisation, *Weekly Epidemiological Record*. Weekly Epidemiological Record, 2023. **98**(11): p. 99-110.
4. Anderson, T.K., et al., *Swine Influenza A Viruses and the Tangled Relationship with Humans*. Cold Spring Harb Perspect Med, 2021. **11**(3).
5. Judge, E.P., et al., *Anatomy and bronchoscopy of the porcine lung. A model for translational respiratory medicine*. Am J Respir Cell Mol Biol, 2014. **51**(3): p. 334-43.
6. Pabst, R., *The pig as a model for immunology research*. Cell Tissue Res, 2020. **380**(2): p. 287-304.
7. Walters, E.M., et al., *Swine models, genomic tools and services to enhance our understanding of human health and diseases*. Lab Anim (NY), 2017. **46**(4): p. 167-172.
8. Trebbien, R., L.E. Larsen, and B.M. Viuff, *Distribution of sialic acid receptors and influenza A virus of avian and swine origin in experimentally infected pigs*. Virol J, 2011. **8**: p. 434.
9. Holzer, B., et al., *Protective porcine influenza virus-specific monoclonal antibodies recognize similar haemagglutinin epitopes as humans*. PLoS Pathog, 2021. **17**(3): p. e1009330.
10. Ostrowsky, J.T., et al., *Accomplishments and challenges in developing improved influenza vaccines: An evaluation of three years of progress toward the milestones of the influenza vaccines research and development roadmap*. Vaccine, 2025. **61**: p. 127431.
11. World Health Organisation, *Weekly Epidemiological Record*. Weekly Epidemiological Record, 2016. **91**(41): p. 469-484.
12. World Health Organisation, *Weekly Epidemiological Record*. Weekly Epidemiological Record, 2019. **94**(12): p. 141-160.
13. World Health Organisation, *Weekly Epidemiological Record*. Weekly Epidemiological Record, 2020. **95**(12): p. 105-116.
14. Kim, S.C., et al., *Major Shift of Influenza A Virus of Swine (IAV-S) by Human-to-Swine Spillover of the 2009 Pandemic Virus in Korea*. Transbound Emerg Dis, 2024. **2024**: p. 6366170.
15. Markin, A., et al., *Reverse-zoonoses of 2009 H1N1 pandemic influenza A viruses and evolution in United States swine results in viruses with zoonotic potential*. PLoS Pathog, 2023. **19**(7): p. e1011476.
16. Linnik, J., et al., *Model-based inference of neutralizing antibody avidities against influenza virus*. PLoS Pathog, 2022. **18**(1): p. e1010243.
17. Coggon, A., et al., *Quantifying the zoonotic risk profile of European influenza A viruses in swine from 2010 to 2020 inclusive*. J Virol, 2025. **99**(7): p. e0030625.

18. Huang, K.Y., et al., *Focused antibody response to influenza linked to antigenic*
drift. J Clin Invest, 2015. **125**(7): p. 2631-45.
19. Linderman, S.L., et al., *Potential antigenic explanation for atypical H1N1*
infections among middle-aged adults during the 2013-2014 influenza season.
Proc Natl Acad Sci U S A, 2014. **111**(44): p. 15798-803.
20. McCauley;, J., et al., *Report prepared for the WHO annual consultation on the*
composition of influenza vaccine for the Southern Hemisphere 2016. 2015, The
Francis Crick Institute.
21. Brown, I.H., *The epidemiology and evolution of influenza viruses in pigs*. Vet
Microbiol, 2000. **74**(1-2): p. 29-46.
22. Edmans, M., et al., *Magnitude and Kinetics of T Cell and Antibody Responses*
During H1N1pdm09 Infection in Inbred Babraham Pigs and Outbred Pigs. Front
Immunol, 2020. **11**: p. 604913.
23. Dholakia, V., et al., *Polymerase mutations underlie early adaptation of H5N1*
influenza virus to dairy cattle and other mammals. bioRxiv, 2025: p.
2025.01.06.631435.
24. World Health Organisation, *Manual for the laboratory diagnosis and virological*
surveillance of influenza. 2011, World Health Organisation.
25. Lin, Y., et al., *Optimisation of a micro-neutralisation assay and its application in*
antigenic characterisation of influenza viruses. Influenza Other Respir Viruses,
2015. **9**(6): p. 331-340.