Growth of *Corynebacterium glutamicum* on 5-oxo-L-proline (pyroglutamate) as

carbon and nitrogen source requires the PxpR-controlled *pxpTABC* genes

Lea Sundermeyer[1], Christina Mack[1],and Michael Bott[1,2*]

[1]IBG-1: Biotechnology, Institute of Bio- and Geosciences, Forschungszentrum Jülich,

52425 Jülich, Germany

[2]Bioeconomy Science Center (BioSC), Forschungszentrum Jülich, 52425 Jülich,

Germany

*Corresponding author: Prof. Dr. Michael Bott, IBG-1: Biotechnology, Institute of Bio-

and Geosciences, Forschungszentrum Jülich, 52425 Jülich, Germany

Phone +49 2461 613294; Email m.bott@fz-juelich.de

ORCID identifiers: Lea Sundermeyer 0000-0001-6333-2289; Christina Mack 0009-

0004-4192-8697; Michael Bott 0000-0002-4701-8254

Running title: 5-oxoproline utilization by *C. glutamicum*

ABSTRACT

5-Oxo-L-proline (5-OP) is inevitably formed in all cells by spontaneous cyclization of L-glutamate, L-glutamine, or γ-glutamyl phosphate. Its use as a substrate by bacteria has rarely been described. Here, we show that the actinobacterial species *Corynebacterium glutamicum* can grow well in minimal medium with 5-OP as sole carbon and nitrogen source. We identified the *pxpTABC* gene cluster as being essential for growth on 5-OP. The *pxpT* gene encodes a secondary transporter of the APC superfamily which most likely catalyzes 5-OP uptake into the cell. The *pxpABC* genes encode a recently identified ATP-dependent 5-oxoprolinase that converts 5-OP into L-glutamate for further metabolism. Both PxpT and PxpABC were required for growth with 5-OP. Upstream and divergent to *pxpT*, the gene *pxpR* is located, encoding a GntR-type transcriptional regulator. Deletion of *pxpR* improved growth on 5-OP suggesting that PxpR acts as repressor of the *pxpTABC* operon. This function was further supported by reporter gene studies. Purified PxpR was shown by isothermal titration calorimetry (ITC) to bind 5-OP with a K_D of 726 ± 23 nM. L-proline, L-glutamate, L-glutamine, and L-aspartate were not bound under the conditions tested, suggesting that PxpR is a specific 5-OP biosensor.

IMPORTANCE

The utilization of 5-OP as a carbon and nitrogen source for bacteria has rarely been investigated in the past, although this metabolite is ubiquitous in cells due to its non-enzymatic formation from the amino acids L-glutamate and L-glutamine. In *Corynebacterium glutamicum* cells, but also in other bacteria such as *Escherichia coli*, L-glutamate is present at concentrations in the range of 100 mM, suggesting a substantial and continuous 5-OP formation. Our results demonstrate that 5-OP can

serve both as carbon and nitrogen source for *C. glutamicum* and presumably also for

other bacteria.

KEYWORDS

Corynebacterium glutamicum, 5-oxoproline, 5-oxoprolinase, transport, transcriptional

regulation, isothermal titration calorimetry, biosensor

INTRODUCTION

5-Oxo-L-proline (5-OP), also termed pyroglutamate, is the cyclic lactam of glutamate.

Its presence in all cells is inevitable due to the spontaneous cyclization of L-glutamate,

L-glutamine, and γ-glutamyl phosphate (1). In the case of L-glutamine, approximately

10% is converted to 5-OP per day at 37°C with ammonia as second product (2, 3). 5-

OP is also formed enzymatically in the degradation of glutathione within the γ-glutamyl

cycle (4) and by cleavage of N-terminal 5-OP residues of proteins by pyroglutamyl

peptidase (5, 6).

Earlier reports on 5-OP utilization and metabolism by bacteria were reviewed by

van der Werf and Meister (7). A systematic study on the effects of 5-OP on different

prokaryotic species revealed that growth of thermoacidophilic archaea and some

bacteria was severely inhibited by the presence of 5-OP and that the extent of inhibition

increased with increasing growth temperature and decreasing pH (8). For example, 15

mM 5-OP completely inhibited growth of *Sulfolobus solfataricus* at 78°C and pH 3 (9).

In contrast, growth of some organisms living at neutral pH and moderate temperatures

was not inhibited and sometimes even stimulated by 5-OP (8). In halophilic and

alkaliphilic methanotrophs, 5-OP can serve as osmoprotectant (10). For *Salmonella*

enterica serovar *typhimurium* it was reported that 5-OP cannot be used as nutrient

source for growth, but has an influence on cellular aggregation (11).

5-OP formed during degradation of glutathione in eukaryotes is known to be

converted to glutamate in an ATP-dependent reaction catalyzed by 5-oxoprolinase

(12). An ATP-independent 5-oxoprolinase was identified in *Alcaligenes faecalis* (13).

However, the equilibrium constant of the reaction, K_{eq} ([L-glutamate]/[5-oxo-L-proline])

was found to be ~0.035, showing that the formation of 5-OP from L-glutamate is highly

favored. Interestingly, sequencing of the corresponding gene revealed that the ATP-

independent 5-oxoprolinase contains a Sec signal peptide and therefore is presumably

active in the periplasm (14). In a recent study it was shown that of 984 analyzed

bacterial and archaeal genomes only 115 contain homologs of the eukaryotic-type 5-

oxoprolinase and only 10 had homologs of the enzyme of *A. faecalis* (15). The majority

of the analyzed genomes was found to encode another type of ATP-dependent 5-

oxoprolinase showing no sequence similarity to the eukaryotic-type enzyme. It is

composed of three subunits and the corresponding genes were named *pxpA*, *pxpB*

and *pxpC* for prokaryotic 5-oxoprolinase (15). This enzyme was originally identified in

Pseudomonas (16). It is assumed that PxpB and PxpC are responsible for ATP-

dependent phosphorylation of 5-OP, whereas PxpA catalyzes the decyclization to γ-

glutamyl phosphate and the subsequent dephosphorylation to L-glutamate (17, 18).

Studies with *Bacillus subtilis* revealed that the *pxpABC* genes are essential for

utilization of 5-OP as sole nitrogen source in minimal medium, but their absence also

had a negative effect on growth with ammonia as nitrogen source (15). This suggests

that the absence of 5-oxoprolinase impairs growth fitness of *B. subtilis* also in the

absence of exogeneous 5-OP. It was shown that *B. subtilis* mutants lacking *pxpA*,

pxpB, or *pxpC* accumulate 5-OP both within the cells and in the medium, whereas the

wild type does not (15). Therefore, the negative effect on growth observed for these

mutants might be due to 5-OP accumulation. 5-OP can be used as sole nitrogen source

by *B. subtilis*, but not by mutants lacking the 5-oxoprolinase genes (15). Recently, the

role of the *pxp* genes was explored in *Clostridioides difficile* (19). In contrast to *B.*

subtilis, the genes showed no effect on sporulation in *C. difficile*, but were necessary

for the growth-promoting effect of 5-OP observed for the wild type of this anaerobic

enteric pathogen. It was suggested that the *pxpAGBC* genes facilitate the use of 5-OP

as a carbon source, nitrogen source, or both a carbon and nitrogen source (19).

In this study, we analyzed the role of 5-OP in *Corynebacterium glutamicum*. This

actinobacterial species is a model organism in microbial biotechnology since it is the

most important industrial amino acid producer, with L-glutamate and L-lysine being

produced at a scale of several million tons per year (20-23). Also L-glutamine-

producing strains of *C. glutamicum* have been developed (24). Although both L-

glutamate and L-glutamine spontaneously cyclize to form 5-OP, the presence and fate

of this metabolite has never been studied in *C. glutamicum*. We show that

C. glutamicum is able to grow with 5-OP as sole carbon and nitrogen source and

delineate the molecular basis. 5-OP is taken up by the secondary transporter PxpT

and converted to L-glutamate by the 5-oxoprolinase PxpABC. The corresponding

operon *pxpTABC* is regulated by a GntR-type transcriptional regulator, which we

named PxpR and which specifically binds 5-OP with high affinity.

RESULTS

Growth of *C. glutamicum* with 5-OP as a sole carbon and nitrogen source

5-OP is an unavoidable metabolite in all cells formed by spontaneous cyclization of L-

glutamate, L-glutamine, or γ-glutamyl phosphate. Since the cytoplasmic concentration

of L-glutamate in *C. glutamicum* cells under standard conditions has been reported to

be in the range of 100-200 mM (25-28), it can be expected that 5-OP is constantly

formed. Nevertheless, no studies were performed so far on this metabolite, which

prompted us to analyze how *C. glutamicum* deals with 5-OP. Initially we tested if 5-OP

can serve as carbon and nitrogen source. As shown in Fig. 1, cultivation of the wild

type ATCC 13032 (WT) in modified CGXII minimal medium lacking the nitrogen

sources ammonium sulfate and urea and containing 12.9 g/L (100 mM) 5-OP enabled

growth, showing that 5-OP can serve as a sole carbon and nitrogen source. As a

control, standard CGXII medium with 20 g/L $(NH_4)_2SO_4$ and 5 g/L urea as nitrogen

sources and 20 g/L glucose as carbon source was used. The lag phase for growth on

5-OP was long if the cells were pre-grown in glucose minimal medium, but absent if

cells were pre-cultured with 5-OP, suggesting that the genes required for 5-OP

utilization are inducible (Fig. 1).

Genes required for growth of *C. glutamicum* on 5-OP

Inspection of the genome of *C. glutamicum* (29, 30) for genes encoding 5-oxoprolinase

revealed a cluster of three genes encoding proteins showing 31–43% amino acid

sequence identity to the PxpA, PxpB, and PxpC proteins of *B. subtilis* (15). These

genes were therefore named *pxpA* (cg1141)*, pxpB* (cg1140), and *pxpC* (cg1139).

Upstream of *pxpA*, the gene *cg1142* is located in the same orientation (Fig. 2) that

encodes a protein with 50% sequence identity to YcsG of *B. subtilis*, which is encoded

downstream of *pxpA* and was shown to be required for utilization of 5-OP as a nitrogen

source in *B. subtilis* (15). Based on the experiments reported below, cg1142 was

named *pxpT* for 5-OP transport.

The relevance of the *pxpABC* genes and of *pxpT* for 5-OP utilization was tested

by constructing and analyzing the *C. glutamicum* deletion mutants Δ*pxpABC*,

Δ*pxpTABC*, and Δ*pxpT.* All three mutants grew like the WT in standard CGXII minimal

medium with glucose (20 g/L) (Fig. 3A), but were unable to grow with 5-OP (12.9 g/L)

as sole carbon and nitrogen source (Fig. 3B), suggesting that PxpABC and PxpT are

essential for 5-OP utilization. For confirmation, the expression plasmids pPREx2-

pxpABC, pPREx2-*pxpTABC,* and pPREx2-*pxpT* were constructed and transformed

into the mutant strains. As shown in Fig. 3B, plasmid-based expression of *pxpABC* in

the mutant strain Δ*pxpABC* not only restored growth on 5-OP, but even improved

growth (reduced lag phase, increased growth rate) compared to the WT. Similarly,

plasmid-based expression of either *pxpT* alone or of *pxpTABC* recovered growth of the

Δ*pxpT* mutant on 5-OP, however, expression of *pxpTABC* enabled markedly better

growth than *pxpT* alone (Fig. 3C). Complementation of the Δ*pxpTABC* mutant with

plasmid-encoded *pxpTABC* recovered growth on 5-OP, whereas complementation

with *pxpABC* alone or *pxpT* alone did not (Fig. 3D). These results confirm that both

pxpT and *pxpABC* are required for growth of the Δ*pxpTABC* mutant on 5-OP. In some

cases, the Δ*pxpTABC* mutant transformed with pPREx2-*pxpTABC* did not grow on 5-

OP and plasmid resequencing showed that these strains lacked the transporter gene

and part of 5-oxoprolinase-encoding sequence. The reason for this effect is unknown

yet. We also tested whether pre-cultivation on 5-OP influences growth of the strain

Δ*pxpT* with pPREx2-*pxpT*. As shown in Fig. 3E, precultivation on 5-OP strongly

reduced the lag phase and increased the growth rate, suggesting that in this strain the

pxpABC genes remain inducible by 5-OP.

Regulation of the *pxpTABC* genes by PxpR

Upstream and divergently oriented to *pxpT*, a gene (cg1143) encoding a transcriptional

regulator of the GntR family (31) is located and was designated *pxpR* (Fig. 2). PxpR

consists of an N-terminal helix-turn-helix (HTH) domain (residues 11-69) and a C-

terminal ligand-binding domain (residues 79-205). Due to its genomic localization, an

involvement of PxpR in transcriptional regulation of the *pxpTABC* genes appeared

likely and was tested by the construction and analysis of a Δ*pxpR* deletion mutant. As

shown in Fig. 4, the absence of PxpR had no effect on growth on glucose, but strongly

improved growth on 5-OP by shortening the lag phase and increasing the growth rate.

Complementation of the Δ*pxpR* deletion mutant with plasmid pPREx2-pxpR-Strep

inhibited growth on 5-OP, and this inhibition became stronger when *pxpR* expression

was increased by IPTG addition. These results suggest a function of PxpR as a

repressor of *pxpTABC* whose absence leads to constitutive expression of the genes.

To confirm the repression of *pxpTABC* by PxpR, the *pxpT* promoter region (212

bp upstream of the *pxpT* start codon) was fused to the *venus* reporter gene, resulting

in plasmid pJC1-P$_{pxpT}$-Venus. The plasmid was introduced into the WT and the Δ*pxpR*

mutant and the two strains were cultivated in standard CGXII glucose medium. Both

strains showed the same growth behavior, but the specific Venus fluorescence (ratio

of fluorescence to backscatter) was 3-fold higher for the Δ*pxpR* mutant (0.177 ± 0.005)

than for the wild type (0.063 ± 0.018), supporting the role of PxpR as a repressor for

the *pxpT* promoter.

PxpR functions as a biosensor for 5-OP

Based on the results described above, it appeared likely that PxpR senses the 5-OP

concentration in the cytoplasm and that binding of 5-OP to PxpR triggers a

conformational change that leads to the derepression of the *pxpTABC* genes. To test

for 5-OP binding to PxpR, the protein modified by a carboxyterminal Strep-tag was

overproduced in *E. coli* BL21(DE3) and purified by StrepTactin affinity chromatography

followed by size-exclusion chromatography (Fig. 5). The purified protein was used for

binding studies using isothermal titration calorimetry, which measures binding

interactions by detecting the heat absorbed or released during a binding event (32,

33). Deconvolution of the binding isotherm gives the number of binding sites (n), the

enthalpy of binding (ΔH) and the equilibrium dissociation constant (K$_D$). From these

parameters, the Gibbs free energy (ΔG) and the entropy of binding (ΔS) can be

derived. The ITC experiments (Fig. 5 and Table S1) showed that PxpR binds 5-OP

with a K_D of 726 ± 23 nM (mean value of five determinations). Binding of 5-OP is an

exothermic event with a ΔH of -98.7 ± 1.0 kJ mol^{-1} and a loss of entropy (-

TΔS = 63.6 ± 1.1 kJ mol^{-1}). The binding stoichiometry was 0.324 ± 0.004 if each PxpR

monomer binds one molecule of 5-OP, suggesting that a fraction of the protein was in

a binding-incompetent state. Importantly, L-glutamate, L-aspartate, L-glutamine, and

L-proline did not bind to PxpR under the same experimental conditions, suggesting

that PxpR serves as a specific 5-OP biosensor.

DISCUSSION

In this study we analyzed the utilization of 5-OP by *C. glutamicum*. To the best of

our knowledge, no studies have been performed to date on the fate of this metabolite

in *C. glutamicum*, although it is well known that 5-OP is formed spontaneously from L-

glutamate, L-glutamine, and γ-glutamyl phosphate. The presence of 5-OP in

C. glutamicum cells was previously reported in a metabolomics study by GC-MS (34).

We showed that *C. glutamicum* can grow with 5-OP as the sole carbon and nitrogen

source and identified the genes required for this metabolic pathway and their

transcriptional regulation. The assumed pathways for 5-OP formation and utilization

are summarized in Fig. 6. Due to the widespread occurrence of the *pxpABC* genes

(15), it can be assumed that the capability to utilize 5-OP as a carbon and nitrogen

source is widespread in bacteria.

The *pxpT* gene encodes a secondary transporter of 436 amino acid residues with

11 predicted transmembrane helices. BLAST searches in the Transporter

Classification Database (35) showed that the proteins with the highest sequence

identity, such as YcsG of *B. subtilis*, belong to the NRAMP family of divalent metal ions

transporters (TCDB 2.A.55). This family is part of the large amino acid-polyamine-

organoCation (APC) superfamily of secondary transporters, which also includes

several families involved in amino acid transport (36). Although we cannot exclude a

function of PxpT in the transport of divalent metal ions, our data suggest that PxpT is

required for uptake of 5-OP by *C. glutamicum*. Deletion of *pxpT* prevented growth on

5-OP, which could be reversed by transformation of the Δ*pxpT* mutant with plasmid-

encoded *pxpT*. The homologous YcsG protein of *B. subtilis* is encoded downstream of

the *pxpA* gene and deletion of *ycsG* prevented the use of 5-OP as nitrogen source

(15), supporting a function of YcsG in 5-OP uptake. A homolog of PxpT is also encoded

in the recently analyzed *pxpAGBC* operon of *C. difficile* (PxpG, 46% amino acid

sequence identity to PxpT), but its function has not been studied yet (19). The

observation that lack of PxpT prevents 5-OP utilization suggests that the transporters

involved in L-proline uptake in *C. glutamicum*, PutP (37), ProP, and EctP (38) cannot

compensate for the absence of PxpT under the conditions used in our experiments. A

different type of 5-OP uptake system has been described for *Bordetella pertussis*,

which belongs to the tripartite ATP-independent periplasmic (TRAP) transport systems

(39). The crystal structures of two extracytoplasmic solute receptors DctP6 and DctP7

were solved in complex with 5-OP. The K_D of DctP7 to 5-OP was found to be 0.3 µM.

Although 5-OP utilization could not be demonstrated, the fact that the genes BP1887

and BP1891 for DctP6 and DcdP7, respectively, are located in the vicinity of a gene

(locus tag BP1985) encoding a eukaryotic-type 5-oxoprolinase supports the existence

of TRAP-type 5-OP uptake systems in some bacteria.

Similar to the *pxpT* gene, the *pxpABC* genes, encoding 5-oxoprolinase, were

essential for growth of *C. glutamicum* on 5-OP as sole carbon and nitrogen source.

Interestingly, growth of mutants lacking *pxpABC* or *pxpTABC* on glucose was not

impaired (Fig. 3A), although 5-OP can presumably no longer be converted to glutamate

by 5-oxoprolinase and therefore might accumulate within the cells. As L-proline is

known to serve as a compatible solute in *C. glutamicum* that can accumulate to very

high concentrations (40), also 5-OP may be tolerated at high concentrations. As

mentioned before, 5-OP serves as a compatible solute in halophilic and alkaliphilic

methanotrophic bacteria, which contain high concentrations under hyperosmotic stress

(10). In *Methylomicrobium alcaliphilum* cells grown at a NaCl concentration of 1 M the

concentration of 5-OP reached 0.4 M (41). Therefore, it seems possible that also other

bacteria like *C. glutamicum* can tolerate high 5-OP concentrations.

To our knowledge, regulation of 5-OP metabolism in bacteria has not been studied

before. We demonstrated that the *pxpTABC* genes of *C. glutamicum* are repressed by

the GntR-type transcriptional regulator PxpR. Deletion of *pxpR* had no influence on

growth with glucose, but strongly improved growth on 5-OP by reducing the lag phase

and increasing the growth rate. This phenotype of the Δ*pxpR* mutant could be reversed

by plasmid-encoded expression of *pxpR* whereby stronger expression led to stronger

growth inhibition (Fig. 4). The repression of the *pxpTABC* genes by PxpR was

supported by a plasmid-encoded reporter gene fusion of the *pxpT* promoter with *venus*,

which revealed a threefold higher expression level in the Δ*pxpR* mutant than in the WT

after growth on glucose.

Regarding the DNA binding site of PxpR, the RegPrecise database (42), which

predicts transcriptional regulons based on comparative genomics (42), anticipates a

DNA binding site with a 17 bp inverted repeat sequence motif shown in Fig. S1 for

PxpR (Cg1143). This motif is centered 65 bp upstream of the start codon of *pxpT* (Fig.

S2). A putative -10 region (TAACCT) for *pxpT* was identified 95.5 bp upstream of the

pxpT start codon, supporting a a repressor function of PxpR. The transcriptional start

site of *pxpR* was identified in a genome-wide study at the ATG start codon of pxpR

(43), indicating that *pxpR* is a leaderless gene (44). Due to the 143-bp distance of the

predicted PxpR binding site to the transcriptional start site of *pxpR*, autoregulation of

pxpR expression appears unlikely. Although 5-OP can serve as sole nitrogen source,

the *pxpTABC* genes were not found to be part of the nitrogen starvation stimulon, which

includes the regulon of the master regulator of the nitrogen starvation response, AmtR

(45-48). However, a binding site (TGTGCTATAGGACACA) for the cAMP-responsive

global regulator GlxR was identified in the intergenic region between *pxpR* and *pxpT*

close to the assumed -10 region of *pxpT* (49, 50), suggesting a negative influence on

pxpTABC expression.

The function as a repressor of *pxpTABC* suggested that PxpR senses the

cytoplasmic concentration of 5-OP and relieves repression of *pxpTABC* once the 5-OP

level reaches a certain level. In agreement with this suggestion we could show by ITC

that purified PxpR binds 5-OP with a K_D of 726 nM. This suggests that 5-OP

concentrations in the µM range are sufficient for derepression of the *pxpTABC* genes.

Since we could not detect binding of L-glutamate, L-aspartate, L-glutamine, and L-

proline under the tested conditions, PxpR is presumably a specific 5-OP sensor. This

offers the opportunity to develop various types of 5-OP biosensors enabling

cytoplasmic 5-OP detection, e.g. based on expression of a reporter gene for a

fluorescent protein under the control of the *pxpT* promoter (51) or FRET-based

biosensors (52).

MATERIALS AND METHODS

Bacterial strains, media, and culture conditions

All bacterial strains and plasmids used in this work are listed in Table 2. *Escherichia*

coli cells were cultivated at 37 °C in lysogeny broth (LB) (53) or terrific broth (TB)

(12 g L^{-1} tryptone, 24 g L^{-1} yeast extract, 4 mL glycerol, 12.54 g L^{-1} K_2HPO_4,

2.31 g L^{-1} KH_2PO_4; pH 7.0) or on LB agar plates (Carl Roth, Karlsruhe, Germany).

C. glutamicum strains were cultivated at 30°C in brain heart infusion medium (BHI;

Difco Laboratories, Detroit, USA) or in CGXII medium with 20 g L^{-1} glucose (54)

containing 30 mg L^{-1} 3,4-dihydroxybenzoate as an iron chelator or in modified CGXII

medium lacking glucose, ammonium sulfate and urea and containing 12.9 g L^{-1} 5-oxo-

L-proline as the nitrogen and carbon source. Solid media were prepared by adding

15 g L^{-1} agar to these media. To maintain plasmid stability, kanamycin was added at

concentrations of 25 mg L^{-1} (*C. glutamicum*) or 50 mg L^{-1} (*E. coli*).

Growth experiments in BioLector microcultivation systems

Growth experiments with *C. glutamicum* strains were routinely performed as

microscale cultivations with BioLector I or II instruments (Beckman Coulter, Brea,

USA). Growth in this system was measured online as backscattered light at 620 nm

(55). Cultivations were performed in a 48-well FlowerPlate (Beckman Coulter, Brea,

USA) by using the defined CGXII media described above. When strains carried

expression plasmids, and 25 µg mL^{-1} kanamycin were added to the medium.

Inoculations were individually prepared to reach an initial OD_{600} of 1.0. BioLector

cultivations were performed at 30°C, 1200 rpm and 85% humidity. The fluorescence

of the reporter protein Venus was measured with an eYFP filter (excitation 508 nm,

emission 532 nm).

Standard recombinant DNA methods and construction of deletion mutants

Standard methods such as PCR and plasmid restriction were carried out according to

established protocols (56). All oligonucleotides used are listed in Table 2. Plasmids

were constructed by ligating DNA fragments obtained by restriction digestion or PCR

or by Gibson assembly (57). Oligonucleotide synthesis and DNA sequencing were

performed by Eurofins Genomics (Ebersberg, Germany). Transformation of *E. coli* was

performed using a standard protocol (58) and *C. glutamicum* transformation was

performed by electroporation (59). *C. glutamicum* deletion mutants were constructed

by double homologous recombination using pK19mobsacB-based plasmids as

described previously (60). Oligonucleotides annealing upstream and downstream of

the deleted genes were used to confirm genomic deletions by colony-PCR.

PxpR overproduction and purification

For testing the binding of 5-OP and other metabolites to PxpR, the protein was

overproduced in *E. coli* BL21(DE3) using the expression plasmid pPREx6-*pxpR*-Strep

encoding a PxpR protein with a C-terminal Strep-tag-II. The strain was cultivated in TB

medium with 50 µg mL^{-1} l kanamycin at 37°C. At an OD_{600} of about 0.6, *pxpR*

expression was induced by addition of 0.5 mM IPTG and then the culture was further

incubated overnight at 18°C. Afterwards cells were harvested and stored at -80°C. For

purification of PxpR, cells were resuspended (4 g cell wet weight mL^{-1}) in Strep binding

buffer (100 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA, pH 8) supplemented with

cOMPLETE EDTA-free protease inhibitor cocktail (Roche, Basel, Switzerland) and

disrupted by one passage through a Multishot Cell Disruptor (Constant System,

Georgia, USA) at 20,000 psi. The cell extract was first subjected to a low-speed

centrifugation (5,000 *g*, 4°C, 20 min) and subsequent ultracentrifugation of the

supernatant (100,000 *g*, 4°C, 1 h). Supernatants of the ultracentrifugation were loaded

onto a StrepTrap HP column (GE Healthcare, Chicago, IL, USA) and, after washing,

the Strep-tagged protein was eluted using binding buffer containing 2.5 mM

desthiobiotin. The protein was further purified by size exclusion chromatography on a

Superdex 200 10/300 GL column (GE Healthcare, Chicago, IL, USA) equilibrated in

HEPES buffer (40 mM HEPES-NaOH, 100 mM NaCl, pH 7.4). Protein concentrations

were determined using a Colibri microvolume spectrometer (Berthold Detection

Systems GmbH, Pforzheim, Germany) and the molar extinction coefficient at 280 nm

of 22920 M^{-1} cm^{-1} predicted by the ProtParam tool (http://web.expasy.org/protparam/).

Isothermal titration calorimetry

Purified PxpR-Strep was dialyzed overnight in HEPES buffer (40 mM HEPES–NaOH,

pH 7.4, 100 mM NaCl). 20 mM stock solutions of 5-OP, L-glutamic acid, L-glutamine,

L-proline, and 2-oxoglutaric acid were prepared in dialysis buffer, and the pH was

adjusted to pH 7.4 using NaOH. ITC measurements were performed with a MicroCal

PEAQ-ITC instrument operated at 25°C. The protein concentration was 40 µM and the

ligand concentration was 200 µM. Prior to filling the measuring cell with 300 µL protein

solution, the cell was rinsed with dialysis buffer, and the syringe was filled with 75 µL

ligand solution. An ITC run was started with an initial injection of 0.4 µL followed by 36

injections of 1 µL each. In addition, control experiments with ligand solution titrated into

the dialysis buffer were performed. The data were analyzed using the MicroCal ITC

analysis software (Malvern Panalytical, Malvern, United Kingdom).

Acknowledgements

374 This work was financially supported by the DFG-ANR project MetActino (DFG BO

375 903/4-1, ANR-18-CE92-0003). The authors thank Tobias Georgi for initial experiments

376 and Elena von Helden for support in the preparation of the figures.

References

1. Kumar A, Bachhawat AK. 2012. Pyroglutamic acid: throwing light on a lightly studied metabolite.
Curr Sci 102:288-297.
2. Tritsch GL, Moore GE. 1962. Spontaneous decomposition of glutamine in cell culture media.
Exp Cell Res 28:360-364.
3. Ozturk SS, Palsson BO. 1990. Chemical decomposition of glutamine in cell culture media: effect
of media type, pH, and serum concentration. Biotechnol Prog 6:121-8.
4. Bachhawat AK, Yadav S. 2018. The glutathione cycle: glutathione metabolism beyond the γ-
glutamyl cycle. IUBMB Life 70:585-592.

5.   Awade AC, Cleuziat P, Gonzales T, Robert-Baudouy J. 1994. Pyrrolidone carboxyl peptidase
(Pcp): an enzyme that removes pyroglutamic acid (pGlu) from pGlu-peptides and pGlu-proteins.
Proteins 20:34-51.
6.   Cummins PM, O'Connor B. 1998. Pyroglutamyl peptidase: an overview of the three known
enzymatic forms. Biochim Biophys Acta 1429:1-17.
7.   van der Werf P, Meister A. 1975. Metabolic formation and utilization of 5-oxo-L-proline (L-
pyroglutamate, L-pyrrolidone carboxylate). Advances in Enzymology and Related Areas of
Molecular Biology 43:519-556.
8.   Park CB, Ryu DD, Lee SB. 2003. Inhibitory effect of L-pyroglutamate on extremophiles:
correlation with growth temperature and pH. FEMS Microbiol Lett 221:187-190.
9.   Park CB, Lee SB, Ryu DD. 2001. L-pyroglutamate spontaneously formed from L-glutamate
inhibits growth of the hyperthermophilic archaeon *Sulfolobus solfataricus*. Appl Environ
Microbiol 67:3650-3654.
10. Khmelenina VN, Kalyuzhnaya MG, Sakharovsky VG, Suzina NE, Trotsenko YA, Gottschalk G.
1999. Osmoadaptation in halophilic and alkaliphilic methanotrophs. Arch Microbiol 172:321-
329.
11. Stern E, Shterzer N, Mills E. 2022. Characterizing 5-oxoproline sensing pathways of *Salmonella*
enterica serovar *typhimurium*. Sci Rep 12:15975.
12. van der Werf P, Orlowski M, Meister A. 1971. Enzymatic conversion of 5-oxo-L-proline (L-
pyrrolidone carboxylate) to L-glutamate coupled with cleavage of adenosine triphosphate to
adenosine diphosphate, a reaction in the γ-glutamyl cycle. Proc Natl Acad Sci USA 68:2982-
2985.
13. Nishimura A, Ozaki Y, Oyama H, Shin T, Murao S. 1999. Purification and characterization of a
novel 5-oxoprolinase (without ATP-hydrolyzing activity) from *Alcaligenes faecalis* N-38A. Appl
Environ Microbiol 65:712-177.
14. Nishimura A, Oyama H, Hamada T, Nobuoka K, Shin T, Murao S, Oda K. 2000. Molecular
cloning, sequencing, and expression in *Escherichia coli* of the gene encoding a novel 5-
oxoprolinase without ATP-hydrolyzing activity from *Alcaligenes faecalis* N-38A. Appl Environ
Microbiol 66:3201-3205.
15. Niehaus TD, Elbadawi-Sidhu M, de Crecy-Lagard V, Fiehn O, Hanson AD. 2017. Discovery of
a widespread prokaryotic 5-oxoprolinase that was hiding in plain sight. J Biol Chem 292:16360-
16367.
16. van der Werf P, Meister A. 1974. Isolation of 5-oxoprolinase from a prokaryote. Biochem
Biophys Res Commun 56:90-96.
17. Li LY, Seddon AP, Meister A. 1988. Interaction of the protein components of 5-oxoprolinase.
Substrate-dependent enzyme complex formation. J Biol Chem 263:6495-6501.
18. Oke M, Oni O, Bello R, Samuel-Omoyajowo K, Senbadejo T. 2019. Structure-function
relationships of the 5-oxoprolinase subunit A: Guiding biological sciences students down the
path less traveled. Biochem Mol Biol Educ 47:620-631.
19. Lee CD, Rizvi A, Carter ZA, Edwards AN, McBride SM. 2025. The Pxp complex detoxifies 5-
oxoproline and promotes the growth of *Clostridioides difficile*. Mol Microbiol 124:66-76.
20. Eggeling L, Bott M (ed). 2005. Handbook of *Corynebacterium glutamicum*. CRC Press, Taylor
& Francis Group, Boca Raton, Florida, USA.
21. Eggeling L, Bott M. 2015. A giant market and a powerful metabolism: L-lysine provided by
Corynebacterium glutamicum. Appl Microbiol Biotechnol 99:3387-3394.
22. Becker J, Wittmann C. 2012. Bio-based production of chemicals, materials and fuels -
Corynebacterium glutamicum as versatile cell factory. Curr Opin Biotechnol 23:631-640.
23. Wendisch VF, Jorge JMP, Perez-Garcia F, Sgobba E. 2016. Updates on industrial production
of amino acids using *Corynebacterium glutamicum*. World J Microbiol Biotechnol 32:105.
24. Lv Q, Hu M, Tian L, Liu F, Wang Q, Xu M, Rao Z. 2021. Enhancing L-glutamine production in
Corynebacterium glutamicum by rational metabolic engineering combined with a two-stage pH
control strategy. Bioresour Technol 341:125799.
25. Gutmann M, Hoischen C, Krämer R. 1992. Carrier-mediated glutamate secretion by
Corynebacterium glutamicum under biotin limitation. Biochim Biophys Acta 1112:115-123.
26. Lange J, Münch E, Müller J, Busche T, Kalinowski J, Takors R, Blombach B. 2018. Deciphering
the adaptation of *Corynebacterium glutamicum* in transition from aerobiosis via microaerobiosis
to anaerobiosis. Genes (Basel) 9:297.
27. da Luz JA, Hans E, Frank D, Zeng AP. 2017. Analysis of intracellular metabolites of
Corynebacterium glutamicum at high cell density with automated sampling and filtration and
assessment of engineered enzymes for effective L-lysine production. Eng Life Sci 17:512-522.

28. Niebisch A, Kabus A, Schultz C, Weil B, Bott M. 2006. Corynebacterial protein kinase G controls
2-oxoglutarate dehydrogenase activity via the phosphorylation status of the OdhI protein. J Biol
Chem 281:12300-12307.
29. Kalinowski J, Bathe B, Bartels D, Bischoff N, Bott M, Burkovski A, Dusch N, Eggeling L,
Eikmanns BJ, Gaigalat L, Goesmann A, Hartmann M, Huthmacher K, Krämer R, Linke B,
McHardy AC, Meyer F, Möckel B, Pfefferle W, Pühler A, Rey DA, Rückert C, Rupp O, Sahm H,
Wendisch VF, Wiegrabe I, Tauch A. 2003. The complete *Corynebacterium glutamicum* ATCC
13032 genome sequence and its impact on the production of L-aspartate-derived amino acids
and vitamins. J Biotechnol 104:5-25.
30. Ikeda M, Nakagawa S. 2003. The *Corynebacterium glutamicum* genome: features and impacts
on biotechnological processes. Appl Microbiol Biotechnol 62:99-109.
31. Hoskisson PA, Rigali S. 2009. Chapter 1: Variation in form and function the helix-turn-helix
regulators of the GntR superfamily. Adv Appl Microbiol 69:1-22.
32. Wiseman T, Williston S, Brandts JF, Lin LN. 1989. Rapid measurement of binding constants
and heats of binding using a new titration calorimeter. Anal Biochem 179:131-137.
33. Ladbury JE, Chowdry BZ. 1996. Sensing the heat: the application of isothermal titration
calorimetry to thermodynamic studies of biomolecular interactions. Chem Biol 3:791-801.
34. Strelkov S, von Elstermann M, Schomburg D. 2004. Comprehensive analysis of metabolites in
Corynebacterium glutamicum by gas chromatography/mass spectrometry. Biol Chem 385:853-
861.
35. Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, Lam KJK,
Tian N, Russum S, Wang J, Medrano-Soto A. 2021. The Transporter Classification Database
(TCDB): 2021 update. Nucleic Acids Res 49:D461-D467.
36. Vastermark A, Wollwage S, Houle ME, Rio R, Saier MH, Jr. 2014. Expansion of the APC
superfamily of secondary carriers. Proteins 82:2797-2811.
37. Peter H, Bader A, Burkovski A, Lambert C, Krämer R. 1997. Isolation of the *putP* gene of
Corynebacterium glutamicum and characterization of a low-affinity uptake system for
compatible solutes. Arch Microbiol 168:143-151.
38. Peter H, Weil B, Burkovski A, Krämer R, Morbach S. 1998. *Corynebacterium glutamicum* is
equipped with four secondary carriers for compatible solutes: Identification, sequencing, and
characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine
betaine carrier, EctP. J Bacteriol 180:6005-6012.
39. Rucktooa P, Antoine R, Herrou J, Huvent I, Locht C, Jacob-Dubuisson F, Villeret V, Bompard
C. 2007. Crystal structures of two *Bordetella pertussis* periplasmic receptors contribute to
defining a novel pyroglutamic acid binding DctP subfamily. J Mol Biol 370:93-106.
40. Morbach S, Krämer R. 2005. Osmoregulation, p 417-435. *In* Eggeling L, Bott M (ed), Handbook
of Corynebacterium glutamicum. CRC Press, Taylor & Francis Group, Boca Raton.
41. Trotsenko YA, Khmelenina VN. 2002. The biology and osmoadaptation of haloalkaliphilic
methanotrophs Microbiology (translated from Mikrobiologiya, Vol 71, No 2, 2002, pp 149–159)
71:123-132.
42. Novichkov PS, Kazakov AE, Ravcheev DA, Leyn SA, Kovaleva GY, Sutormin RA, Kazanov MD,
Riehl W, Arkin AP, Dubchak I, Rodionov DA. 2013. RegPrecise 3.0 - a resource for genome-
scale exploration of transcriptional regulation in bacteria. BMC Genomics 14:745.
43. Pfeifer-Sancar K, Mentz A, Rückert C, Kalinowski J. 2013. Comprehensive analysis of the
Corynebacterium glutamicum transcriptome using an improved RNAseq technique. BMC
Genomics 14:888.
44. Beck HJ, Moll I. 2018. Leaderless mRNAs in the spotlight: ancient but not outdated! Microbiol
Spectr 6:RWR-0016-2017.
45. Silberbach M, Hüser A, Kalinowski J, Puhler A, Walter B, Krämer R, Burkovski A. 2005. DNA
microarray analysis of the nitrogen starvation response of *Corynebacterium glutamicum*. J
Biotechnol 119:357-367.
46. Silberbach M, Schäfer M, Hüser AT, Kalinowski J, Pühler A, Krämer R, Burkovski A. 2005.
Adaptation of *Corynebacterium glutamicum* to ammonium limitation: A global analysis using
transcriptome and proteome techniques. Appl Environm Microbiol 71:2391-2402.
47. Beckers G, Strösser J, Hildebrandt U, Kalinowski J, Farwick M, Krämer R, Burkovski A. 2005.
Regulation of AmtR-controlled gene expression in *Corynebacterium glutamicum*: mechanism
and characterization of the AmtR regulon. Mol Microbiol 58:580-595.
48. Burkovski A. 2007. Nitrogen control in *Corynebacterium glutamicum*: proteins, mechanisms,
signals. J Microbiol Biotechnol 17.

49. Toyoda K, Teramoto H, Inui M, Yukawa H. 2011. Genome-wide identification of *in vivo* binding
sites of GlxR, a cyclic AMP receptor protein-type regulator in *Corynebacterium glutamicum*. J
Bacteriol 193:4123-4133.
50. Jungwirth B, Sala C, Kohl TA, Uplekar S, Baumbach J, Cole ST, Pühler A, Tauch A. 2013. High-
resolution detection of DNA binding sites of the global transcriptional regulator GlxR in
Corynebacterium glutamicum. Microbiology 159:12-22.
51. Eggeling L, Bott M, Marienhagen J. 2015. Novel screening methods-biosensors. Curr Opin
Biotechnol 35:30-36.
52. Frommer WB, Davidson MW, Campbell RE. 2009. Genetically encoded biosensors based on
engineered fluorescent proteins. Chem Soc Rev 38:2833-2841.
53. Bertani G. 1951. Studies on lysogenesis. The mode of phage liberation by lysogenic *Escherichia*
coli. J Bacteriol 62:293-300.
54. Keilhauer C, Eggeling L, Sahm H. 1993. Isoleucine synthesis in *Corynebacterium glutamicum*:
molecular analysis of the *ilvB-ilvN-ilvC* operon. J Bacteriol 175:5595-5603.
55. Kensy F, Zang E, Faulhammer C, Tan RK, Büchs J. 2009. Validation of a high-throughput
fermentation system based on online monitoring of biomass and fluorescence in continuously
shaken microtiter plates. Microb Cell Fact 8:31.
56. Green MR, Sambrook J. 2012. Molecular Cloning. A Laboratory Manual, 3rd ed. Cold Spring
Harbor Laboratory Press, Cold Spring Harbor, New York.
57. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic
assembly of DNA molecules up to several hundred kilobases. Nat Meth 6:343-345.
58. Hanahan D. 1983. Studies on transformation of *Escherichia coli* with plasmids. J Mol Biol
166:557-580.
59. van der Rest ME, Lange C, Molenaar D. 1999. A heat shock following electroporation induces
highly efficient transformation of *Corynebacterium glutamicum* with xenogeneic plasmid DNA.
Appl Microbiol Biotechnol 52:541-545.
60. Niebisch A, Bott M. 2001. Molecular analysis of the cytochrome bc_1-aa_3 branch of the
Corynebacterium glutamicum respiratory chain containing an unusual diheme cytochrome c_1.
Arch Microbiol 175:282-294.
61. Studier FW, Moffatt BA. 1986. Use of bacteriophage T7 RNA polymerase to direct selective
high-level expression of cloned genes. J Mol Biol 189:113-130.
62. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small mobilizable
multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19:
selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. Gene
145:69-73.
63. Bakkes PJ, Ramp P, Bida A, Dohmen-Olma D, Bott M, Freudl R. 2020. Improved pEKEx2-
derived expression vectors for tightly controlled production of recombinant proteins in
C*orynebacterium glutamicum*. Plasmid 112:102540.
64. Ramp P, Pfleger C, Dittrich J, Mack C, Gohlke H, Bott M. 2022. Physiological, biochemical, and
structural bioinformatic analysis of the multiple inositol dehydrogenases from *Corynebacterium*
glutamicum. Microbiol Spectr doi:10.1128/spectrum.01950-22:e0195022.
65. Cremer J, Eggeling L, Sahm H. 1990. Cloning the *dapA dapB* cluster of the lysine-secreting
bacterium *Corynebacterium glutamicum*. Mol Gen Genet 220:478-480.
549

Table 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Description	Reference or source
***C. glutamicum* strains**		
ATCC 13032	Biotin-auxotrophic wild-type strain	DSMZ
Δ*pxpABC*	Wild type derivative with in-frame deletion of *pxpABC* (cg1141-cg1140-cg1139)	This work
Δ*pxpTABC*	Wild type derivative with in-frame deletion of *pxpTABC* (cg1142-cg1141-cg1140-cg1139)	This work
Δ*pxpT*	Wild type derivative with in-frame deletion of *pxpT* (cg1142)	This work
Δ*pxpR*	Wild type derivative with in-frame deletion of *pxpR* (cg1143)	This work
***E. coli* strains**		
DH5α	F- *supE44* Δ*lacU*169 (Φ80*lacZ*Δ*M15*) *hsdR17 recA1 endA1 gyrA96 thi-1 relA1*	(58)
BL21(DE3)	F- *ompT hsdSB(rB-mB-) gal dcm* (λc*Its857 ind1* Sam7 *nin5 lacUV5*-T7 Gen 1	(61)
Plasmids		
pK19mobsacB	Kan^R; suicide vector for allelic exchange in *C. glutamicum*; *oriV*~E. coli~ *oriT sacB*	(62)
pK19mobsacB-Δ*pxpABC*	Kan^R; pK19mobsacB derivative containing a PCR product covering the upstream region of *pxpA* and the downstream region of *pxpC*	This work
pK19mobsacB-Δ*pxpTABC*	Kan^R; pK19mobsacB derivative containing a PCR product covering the upstream region of *pxpT* and the downstream region of *pxpC*	This work
pK19mobsacB-Δ*pxpT*	Kan^R; pK19mobsacB derivative containing a PCR product covering the upstream and downstream regions of *pxpT*	This work
pK19mobsacB-Δ*pxpR*	Kan^R; pK19mobsacB derivative containing a PCR product covering the upstream and downstream region of *pxpR*	This work
pPREx2	Kan^R; pPBEx2 derivative (P*tacI*, *lacI*q, *oriC.g* from pBL1.; *oriE.c.* ColE1 from pUC18), with a consensus RBS (AAGGAG) for *C. glutamicum*	(63)

pPREx2-*pxpABC*	Kan^R; pPREx2 derivative carrying the *pxpABC* genes	This work
pPREx2-*pxpTABC*	Kan^R; pPREx2 derivative carrying the *pxpTABC* genes	This work
pPREx2-*pxpT*	Kan^R; pPREx2 derivative carrying the *pxpT* gene	This work
pPREx2-*pxpR*-Strep	Kan^R; pPREx2 derivative carrying the pxpR gene fused to a Strep tag-II encoding sequence at the 3'-end	This work
pPREx6	Kan^R; pPREx2 derivative (P$_{T7}$, *lacI^q*, *oriC.g* from pBL1.; *oriE.c.* ColE1 from pUC18), with a consensus RBS (AAGGAG) for *C. glutamicum*	(64)
pPREx6-*pxpR*-Strep	pPREx6 derivative carrying the *pxpR* gene fused to a Strep -tag-II encoding sequence at the 3'-end under the control of the T7 promoter	This work
pJC1	Kan^R; *E. coli-C. glutamicum* shuttle vector	(65)
pJC1-P$_{pxpT}$-Venus	Kan^R; pJC1 derivative containing the Venus reporter gene under the control of the *pxpT* promoter (transcriptional fusion)	This work

Table 2. Oligonucleotides used in this study

Name	Sequence
Construction of pK19mobsacB-Δ*pxpABC*	
D1_cg1139-1141	TGCAGGTCGACTCTAGAGTCATGCGCGTGGTGCTCTTC
D2_cg1139-1141	GACTTTATCGCCCGGCAATTCGCCGAGGTCGCTGTTG
D3_cg1139-1141	AACAGCGACCTCGGCGAATTGCCGGGCGATAAAGTCAG
D4_cg1139-1141	GTAAAACGACGGCCAGTGTCAGGTGAGGGACATCTAC
Construction of pK19mobsacB-Δ*pxpTABC*	
D1_cg1139-1142	TGCAGGTCGACTCTAGAGTAGGCCTCTCGCATTACCC
D2_cg1139-1142	GACTTTATCGCCCGGCAATGTCGCGCTCCTTTCGTG
D3_cg1139-1142	CACGAAAGGAGCGCGACATTGCCGGGCGATAAAGTCAG
D4_cg1139-1142	GTAAAACGACGGCCAGTGTCAGGTGAGGGACATCTAC
Construction of pK19mobsacB-Δ*pxpT*	
D1_cg1142	TGCAGGTCGACTCTAGAGTAGGCCTCTCGCATTACCC
D2_cg1139-1142	AACCAGGCCGATGACTCCTGTCGCGCTCCTTTCGTG
D3_cg1142	CACGAAAGGAGCGCGACAGGAGTCATCGGCCTGGTTATC
D4_cg1142	GTAAAACGACGGCCAGTGAGCCAGGGCCATGAGGGAG
Construction of pK19mobsacB-Δ*pxpR*	
D1_cg1143	TTGTTCCGCCTCTTCGAACATGTGACTGATCTTAAAGGAC
D2_cg1143	TGCAGGTCGACTCTAGAGTAACCGTGTTGCCGAGTTC
D3_cg1143	TTTAAGATCAGTCACATGTTCGAAGAGGCGGAACAATAC
D4_cg1143	GTAAAACGACGGCCAGTGAGCTGCATTCATCGCGAG
Confirmation of gene deletion by colony-PCR	
Dcg1139-1142 rv	TGGCAGAACCGACAGTTTG
Dcg1139-1142 fw	GTAAAACGACGGCCAGTGTCAGGTGAGGGACATCTAC
Dcg1143 fw	TCCAGTGCTGGCTGAGAC
Dcg1143 rv	TGAACGCTGAATCATGGC
Constrtuction of pPREx2-*pxpABC*	
pPREx2-cg1141 fw	TGCAGAAGGAGATATACATATGACCACCATCGATCTCAAC
pPREx2-cg1139 rv	AAACGACGGCCAGTGAATTCTTAAAGCAATTTAAATCTGACTTTATC
Construction of pPREx2-*pxpTABC*	
pPREx2-cg1142 fw	TGCAGAAGGAGATATACATATGGCCGACAACAAAAATG
pPREx2-cg1139 rv	AAACGACGGCCAGTGAATTCTTAAAGCAATTTAAATCTGACTTTATC
Construction of pPREx2-*pxpT*	
pPREx2-cg1142 fw	TGCAGAAGGAGATATACATATGGCCGACAACAAAAATG
pPREx2-cg1142 rv	AAACGACGGCCAGTGAATTCTTAACTGCTAAGAAGATCGAAG
Construction of pPREx2-*pxpR*-Strep	
cg1143-Strep fw	TGCAGAAGGAGATATACAtATGACGCAGGCAATAGCAG
cg1143-Strep rv	GAACTGTGGGTGGGACCAGCTAGCAAATTCCGGTAGGTGCGCAG
Construction of pJC1-P*pxpT*-venus	
Pcg1142 fw	TCAGCGACGCCGCAGGGGATGTGACTGATCTTAAAGGAC
Pcg1142 rv	CATGATATCCCTCCTCTAAGTGCTGGCTGAGACTAGCTG
RBS-Venus fw	TAGAGGAGGGATATCATGGTG
venus-rv	GTTGCCATTGCTGCAGGTCGAC

Fig. 1. Growth of *C. glutamicum* WT on 5-OP as sole carbon and nitrogen source. Cells were cultivated either in standard CGXII medium with 20 g/L glucose as the carbon source and 20 g/L ammonium sulfate plus 5 g/L urea as nitrogen sources or in modified CGXII medium lacking glucose, ammonium sulfate, and urea and containing 12.9 g/L 5-OP as the sole carbon and nitrogen source. The pre-cultures were grown either in glucose medium (unadapted) or in 5-OP medium (adapted). Cultivation was performed in a BioLector microcultivation system at 30°C and 1200 rpm.

Fig. 2. Map of the *C. glutamicum* genome region with the 5-oxoprolinase genes

pxpABC and for comparison the corresponding region of *B. subtilis*. The %-numbers

indicate the protein sequence identity to the homologous *C. glutamicum* proteins.

A

B

C

D

E

Fig. 3. Dependency of growth with 5-OP on the *pxpTABC* genes. The indicated

C. glutamicum strains were cultivated either in CGXII glucose medium as a control (A)

or in modified CGXII medium containing 5-OP as the sole carbon and nitrogen source

(B-E). All media contained 50 µg/mL kanamycin. "Adapted" indicates that the

corresponding cultures were pre-grown in 5-OP medium, whereas all others were pre-

grown in glucose medium. Cultivation was performed in a BioLector microcultivation

system at 30°C and 1200 rpm.

Fig. 4. Influence of the transcriptional regulator PxpR on growth with 5-OP. The

indicated *C. glutamicum* strains were cultivated either in CGXII glucose medium as

control (A) or in modified CGXII medium containing 5-OP as sole carbon and nitrogen

source (B), all containing 50 µg/ml kanamycin. Where indicated, the medium contained

10 µM IPTG for enhancing the expression of *pxpR*-Strep.

Fig. 5. Purification of PxpR and ligand binding studies by isothermal titration

calorimetry. (A) Size exclusion chromatography of PxpR-Strep purified by. (B) SDS-

PAGE of PxpR-Strep purified by Strep-tag HP affinity and size exclusion

chromatography. (B) Raw data for a representative ITC experiment with 40 µM PxpR-

Strep and titration with 150 µM 5-OP. (C) Binding isotherm calculated from the raw

data shown in Fig. 5C. DP, differential power.

Fig. 6. Overview on the formation and degradation of 5-OP in *C. glutamicum* and

transcriptional regulation by PxpR.