1

Cooperative Binding of Cytosolic Type III Secretion System Proteins to the Injectisome

Revealed by Live-Cell Single-Molecule Localization Microscopy

Olivia I.C. de Cuba[1], Gopika A. Lekshmi[2], Caroline F. Tyndall[2], Andreas Gahlmann[1,3,4,#]

[1] Department of Molecular Physiology & Biological Physics, University of Virginia School of

Medicine, Charlottesville, Virginia.

[2] Department of Biology, University of Virginia, Charlottesville, Virginia.

[3] Department of Chemistry, University of Virginia, Charlottesville, Virginia.

[4] Department of Biomolecular Engineering, University of Virginia, Charlottesville, Virginia.

[#] Address correspondence to agahlmann@virginia.edu

ABSTRACT

Type III secretion systems (T3SSs) are employed by many Gram-negative bacteria to translocate virulent effector proteins into host cells. Secretion occurs through the T3SS injectisome, a multi-membrane-spanning biomolecular machine. While the overall structure of the injectisome is becoming increasingly well understood, the structural dynamics that regulate secretion remain unresolved. Particularly important for the functional regulation of type 3 secretion are the cytosolic injectisome proteins which transiently associate with injectisomes and whose structural dynamics and function remain incompletely understood. Here, we use long-exposure single-molecule localization microscopy to quantify the bound times of the cytosolic components SctQ, SctL, and SctN, at the injectisome of *Yersinia enterocolitica* in chemically induced secretion ON and OFF states and in the presence and absence of effector proteins. In the absence of effector proteins, each component exhibits distinct short-lived binding behavior on a timescale of seconds. The presence of the effector protein YopE increases bound times and synchronizes binding behavior across components, even without active secretion. Upon activation of secretion, a subpopulation of long-lived binding events emerges for SctQ, SctL, and SctN even within the same injectisomes, indicating that binding behavior between these proteins is secretion state-dependent and cooperative. These findings establish that the dynamic interactions between cytosolic injectisome proteins are modulated by the presence of secretion substrates and by secretion activating signals. Understanding the protein structural dynamics that underpin these modulations thus likely holds the key to understanding functional regulation of type 3 secretion.

INTRODUCTION

Type III secretion systems (T3SSs) are conserved multi-membrane spanning protein complexes used by many Gram-negative pathogens to assemble either flagella or injectisomes that translocate effector proteins directly into eukaryotic cells[1]. The injectisome includes a membrane-embedded export apparatus connected through the bacterial envelope to an extracellular needle (Fig. 1a). Substrates are engaged and prioritized for secretion through a cytosolic protein network[2-5].

In *Yersinia enterocolitica*, the expression and secretion of effector proteins are tightly regulated by the *yop* regulon, which responds to environmental cues such as temperature and calcium concentration. Induction of the *yop* regulon, along with the low-calcium response, provides tunable experimental parameters for toggling secretion activity and has been used extensively to model secretion-ON and -OFF states of the system[6-8]. Under high-calcium conditions, secretion is repressed and effectors accumulate within the bacterial cytosol, whereas calcium depletion activates the injectisome and triggers effector translocation across the host membrane. This reversible calcium switch enables controlled interrogation of the system's functional state. The post-secretion state, i.e. a secretion OFF state after secretion has occurred at least once, offers mechanistic value because it reveals how the cytosolic components are distinct from their involvement in initial injectisome assembly, thereby probing the involvement of cytosolic components in ON/OFF transitions.

Cryo-electron tomography and in situ structural studies have converged on an

architectural model in which the export apparatus is capped on its cytosolic face by a

multiprotein assembly comprising SctK, SctQ, and SctL, arranged around the hexameric

ATPase SctN and its central stalk[2,3,9,10]. Initial in situ observations using *Shigella* and

Salmonella minicell-producing mutants revealed six distinct cytoplasmic "pods" that

tether SctN to the membrane-embedded SctD ring[2,11,12]. In contrast, cryo-focused ion

beam milling followed by tomography in *Yersinia* revealed more continuous, ring-like

densities beneath the injectisome, reminiscent of the flagellar C ring[13,14]. These densities

are composed of SctK, SctQ, and likely $SctQ_C$, the alternatively expressed C-terminal

fragment of SctQ[15,16], while SctL retains a six-fold symmetric spoke-like arrangement

cradling SctN. Protein-protein interaction studies using bacterial hybrid systems[17-19] and

photo-crosslinking[20] have mapped a sequential SctK–Q–L–N arrangement, with SctQ

playing a particularly enigmatic role. Its N-terminal domain has been implicated in

interactions with SctK[20], $SctQ_C$[21], SctN[18], SctD[22], and even chaperone–effector

complexes[18,19,23]. Together, these studies define the physical stage on which dynamic

interactions occur, but they do not resolve how the cytosolic architecture reconfigures

under secretion-ON versus secretion-OFF conditions in live cells. Despite evidence for

complex formation in vitro, the functional relevance of these interactions, and how they

respond to environmental cues that toggle secretion remains unclear.

Complementing these structural insights, biochemical and genetic studies have

illuminated a functional role for the cytosolic injectisome proteins in substrate hierarchy. A

high molecular weight complex comprising SctK, SctQ, and SctL was shown to coordinate

the sequential loading of translocases and effector proteins, guided by dedicated

chaperones that ensure proper secretion order[24]. Using native gel electrophoresis and

subcellular fractionation in *Salmonella*, this study demonstrated that the so-called "sorting

platform" remains intact even in the absence of the needle complex, suggesting it

assembles independently and may serve as a pre-assembled substrate queue. Disruption

of this sorting platform impairs secretion efficiency and abolishes substrate prioritization,

underscoring the regulatory importance of this cytosolic interface. Whether similar

organizational principles apply across species remains to be determined. Thus, although

the static molecular framework of the injectisome and cytosolic injectisome complexes is

increasingly well-studied, the kinetics and regulation of protein exchange at the

injectisome remain largely unknown.

Fluorescence-recovery-after-photobleaching (FRAP) experiments in living cells have

shown that the cytosolic interface of the injectisome is highly dynamic. In *Yersinia*, SctQ

exchanges between injectisome-bound and cytosolic pools with a recovery half-time of

roughly one minute, accelerating under secreting conditions and requiring ATPase

activity[25]. These data indicate that SctQ binding and unbinding is connected to secretion

activity. Single-molecule tracking by our lab and others, in *Y. enterocolitica* subsequently

revealed that SctQ, SctL, and SctN populate multiple diffusive states in addition to an

injectisome-bound state, consistent with transient hetero-oligomeric subcomplexes that

assemble and disassemble on the timescale of seconds[26,27]. More recent work showed that

chaperones facilitate a cytosolic handover of effector proteins to SctQ-containing

complexes[28]. These SctQ- and effector-containing complexes are proposed to shuttle

substrates to the injectisome, implying that substrate availability informs both the

composition of cytosolic complexes and their binding/bound time to the injectisome[23,28,29].

Together, these observations suggest a dynamic picture in which cytosolic complex

formation, substrate loading, and injectisome engagement are kinetically coupled.

Here, we use long-exposure single-molecule localization to extract bound-time

distributions (also referred to as residence times) of SctQ, SctL, and SctN at the

injectisome of *Y. enterocolitica* in effector-free and YopE-expressing backgrounds under

secretion-OFF and ON conditions. In parallel, 3D single-molecule tracking characterizes

the bound fraction of SctQ as a proxy for k_{on} under different secretion states and the

presence of YopE. This approach allows us to test whether secretion state and effector

presence differentially regulate the binding kinetics and the associated number of

resolvable bound states. By quantifying these kinetics across defined environmental and

substrate conditions, we provide a quantitative framework describing how the T3SS

cytosolic proteins dynamically adjust the bound time in response to secretion demand and

effector availability.

RESULTS

To quantify the bound times of cytosolic T3SS components at the injectisome, we used a biosafety leveadjusttible *Yersinia enterocolitica* strain lacking six major effector proteins referred to as T3SS$^{\Delta effector}$ (also known as IML421asd)[4]. We expressed Halo-tagged SctQ and SctL from their native loci via allelic exchange, while Halo-SctN was expressed from a plasmid due to unsuccessful allelic replacement attempts. All fusion proteins were confirmed to be stable (Fig. S1a) and functional (Fig S1b), i.e. capable of reversible induction by the calcium effect (Fig S1c). These strains thus allowed us to probe the binding behavior of SctQ, SctL, and SctN in the absence of effector interactions.

Halo-fused proteins were visualized with a low concentration of dye coupled with long-exposure imaging, so that fast-moving molecules blur into the background, and stationary, injectisome-bound proteins are detected as sharp point spread functions (Fig 1b). The pooled bound times were displayed as probability density function (pdf)-normalized histograms and fit with single- or bi-exponential decays. The resulting bound time fit (τ) of each protein at the injectisome follows an exponential decay and is the inverse of the dissociation rate k_{off}.

Figure 1: **(a)** Schematic of the Type III Secretion System (T3SS) architecture in *Yersinia enterocolitica*, highlighting the substructures and the cytosolic components analyzed in this study: SctQ, SctL, and SctN. **(b)** Schematic of the long-exposure imaging strategy used to extract bound times from stationary single-molecule trajectories. Fast-diffusing cytosolic molecules blur into the background, while stationary, injectisome-bound molecules appear as sharp point spread functions. This strategy enables direct measurement of bound times at the injectisome.

Effector-Free Conditions Reveal Distinct Binding Dynamics for Injectisome-Associated Proteins SctQ, SctL, and SctN

In the T3SS$^{\Delta effector/OFF}$ condition, the bound time histograms of SctQ, SctL and SctN are explained best by single exponential decays (Fig. 2a-c). Notably, all three proteins exhibit bound times on the order of a few seconds, but the fitted average bound times are all distinct. SctQ exhibits the shortest average bound time, τ = 2.5 s [2.4, 2.5], while the average bound times of SctL and SctN are slightly longer, τ = 3.4 s [3.3, 3.4] and τ = 3.0 s [2.9, 3.0], respectively (99% confidence intervals are indicated in square brackets).

Figure 2: Distinct bound times for SctQ, SctL, and SctN and emergence of long-lived binding events under secretion ON conditions. Bound time distributions for Halo-tagged SctQ, SctL, and SctN under secretion OFF (top row) and ON (bottom row) conditions in the effector-free background. **(a–c)** Under secretion OFF conditions, all three proteins exhibit mono-exponential decay kinetics with distinct bound times: SctQ (τ = 2.5 s), SctL (τ = 3.4 s), and SctN (τ = 3.0 s). **(d–f)** Under secretion ON conditions, a subpopulation of long-lived binding events emerges for all proteins, requiring bi-exponential fits. Long-lived events comprise 9% (SctQ), 8% (SctL), and 12% (SctN) of trajectories, indicating two binding states at the injectisome. a) n = 4063 counts, b) n = 1973 counts, c) n = 5545 counts, d) n = 1417 counts, e) n = 1463 counts, f) n = 5857 counts.

Under secretion ON conditions, we observed a distinct shift in the binding behavior of SctQ, SctL, and SctN at the injectisome. Rather than a uniform change in average bound time, the data revealed a subpopulation of binding events with substantially longer durations (>10 s), prompting bi-exponential fits to the bound time distributions. These long-lived events comprised approximately 9% of SctQ, 8% of SctL, and 12% of SctN trajectories (Fig. 2d-f), suggesting the emergence of two binding modes at the injectisome. The finding

of long-lived states across all three proteins points to a coordinated and potentially

cooperative injectisome-bound configuration under secretion ON conditions.

Notably, the majority of binding events remained short-lived, and the fast

component of the bi-exponential fit was shorter under ON than OFF conditions: for

example, SctQ's short-term bound time decreased from 2.5 s (OFF) to 2.1 s (ON). When

considering both short- and long-lived events, the weighted average bound times were 2.7

s for SctQ, 4.4 s for SctL, and 3.1 s for SctN under T3SS$^{\Delta effector/ON}$ conditions.

As mentioned before, SctQ binding and unbinding have previously been measured

using fluorescence recovery after photobleaching (FRAP), an ensemble technique that

reports on the average recovery behavior of a population. Published FRAP measurements

showed a recovery half-time of 68.2 ± 7.9 s under secreting conditions, compared to 134.3

± 16.1 s under non-secreting conditions[25], representing an overall faster recovery for

T3SS$^{\Delta effector/ON}$. Interestingly, despite the dominance of short-lived events in our single-

molecule data, the average bound times did not skew toward faster exchange as might be

expected from the FRAP results. This raised the question of whether individual

injectisomes engage in both short- and long-lived binding, or whether distinct classes of

injectisomes specialize in one mode over the other.

The Same Injectisome Can Engage in Short- and Long-Term Binding

Each single-molecule binding event can be precisely localized to a specific

subcellular location. Within the same bacterial cell, we observed both short- and long-lived

binding events for SctQ, SctL, and SctN (Fig. 3a), establishing that the emergence of long-

lived states is not due to cell-to-cell variation. In this context, repeated binding at a given

spatial location, i.e. a cluster, can be used as a proxy for the position of injectisomes

(specifically its binding sites). To assess whether long-lived binding events preferentially

occur at a distinct subset of injectisomes, we clustered binding events within each field of

view using DBSCAN (ε = 150 nm, minPts = 2) and classified events as long if their bound

time exceeded 30 frames (i.e. 15 seconds). Across all three proteins, we observed a range

of behaviors; injectisomes with multiple binding events can be classified as showing

exclusively short-lived events, exclusively long-lived events, and mixed binding profiles (Fig.

3b, table S2).

Figure 3: Short- and long-lived binding events occur at the same injectisome. (a) Representative cells
showing spatial co-localization of short- and long-lived binding events for SctQ, SctL, and SctN, confirming
that both binding modes occur within the same cell and at the same subcellular locations. **(b)** Classification
of injectisomes based on binding profiles: mixed (both short and long), short-only, and long-only. **(c)**
Statistical analysis using a hypergeometric test with FDR correction reveals no significant enrichment of long-
lived events at specific clusters, suggesting that binding mode heterogeneity is not spatially segregated but
reflects dynamic behavior at individual injectisomes. Parameters: ε = 100 nm, minPts = 2; long = dwell ≥ 30
frames.

It is important to note that these data were obtained by sparely labeling the proteins of

interest. The classification could therefore be skewed because we are not sampling all the

possible binding events at a given injectisome. We therefore compared the observed

fraction of long events within clusters to the global fraction of long events across the field

using a hypergeometric test with Benjamini–Hochberg correction ($\alpha = 0.05$). No clusters

showed significant enrichment of long events for any protein, indicating that long-lived

binding events were not spatially localized beyond random expectation. We repeated this

analysis for short binding events and again observed no statistical evidence for

enrichment. We therefore conclude that long- and short- lived bound states are not

spatially segregated and occur at the same injectisome.

YopE Modulates Injectisome Binding Dynamics, With SctQ Showing Distinct

Sensitivity Across Secretion States

To determine how the presence of effector proteins influences injectisome-

associated protein dynamics, we quantified the bound lifetimes of SctQ, SctL, and SctN

under secretion ON and OFF conditions in the presence of the effector protein YopE. YopE

was constitutively expressed from a plasmid and confirmed to be successfully secreted in

all strains (Fig S1b).

Figure 4: Effector presence increases bound times and synchronizes binding behavior. Bound time distributions for SctQ, SctL, and SctN under secretion OFF (top row) and ON (bottom row) conditions in the presence of the effector protein YopE. **(a–c)** Under secretion OFF conditions, all three proteins show increased bound times compared to the effector-free background, with mono-exponential fits yielding τ ≈ 4 s for all proteins. **(d–f)** Under secretion ON conditions, SctQ exhibits both an increased short-term bound time (τ = 4.5 s) and a distinct long-lived subpopulation (τ = 23 s, 6% of events). SctL and SctN show smaller changes, indicating component-specific sensitivity to effector presence. a) n = 1345 counts, b) n = 1446 counts, c) n = 2292 counts, d) n = 1003 counts, e) n = 1902 counts, f) n = 1240 counts.

All three proteins exhibited increased bound times compared to the effector-free condition (T3SS$^{\Delta effector/OFF}$), and their averaged bound times converged to statistically indistinguishable values (Fig. 4a-c). Specifically, mono-exponential fits yielded average bound times of τ = 4.1 s [4.0, 4.2] for SctQ, τ = 3.9 s [3.8, 4.0] for SctL, and τ = 3.9 s [3.8, 3.9] for SctN. The increase was most pronounced for SctQ, which nearly doubled its average bound time relative to the effector-free condition. These findings indicate that effector presence alone,

independent of active secretion, can increase the bound time of cytosolic components at

the injectisome. We next asked whether these effects persist during active secretion

(T3SS$^{YopE/ON}$).

Under secretion ON conditions, the presence of YopE modulates the binding

dynamics of injectisome-associated proteins in a component-specific manner (Fig. 4d-f).

Like in T3SS$^{\Delta effector/ON}$, this T3SS$^{YopE/ON}$ secretion condition revealed a subpopulation of

binding events with substantially longer durations (>10 s), prompting bi-exponential fits to

the bound time distributions. These long-lived events comprised approximately 6% of

SctQ, 3% of SctL, and 13% of SctN trajectories (Fig. 4d-f), again suggesting the emergence

of two binding modes at the injectisome. These two binding modes were evaluated using

similar methods as before and we again observed mixed (long and short) binding behavior

at the same injectisome (Table S3). In addition, there was no statistical evidence for

enrichment of one type of binding at specific spatial locations (Fig S3), ruling out the

possibility of two distinct classes of injectisomes. SctQ exhibits the longest short-term

bound time (τ = 4.5 s [3.9, 4.9]), while the short-bound times of SctL and SctN are shorter, τ

= 3.1 s [2.8, 3.5] and τ = 2.1 s [1.7, 2.3], respectively.

Secretion State Modulates Cytosolic Diffusion, Not Injectisome-Bound Fraction, of

SctQ Independently of Effector Presence

The observation that SctQ exhibits slightly longer bound times on average under

secreting conditions does not match previous FRAP measurements that indicate increased

protein exchange under secreting conditions which are proposed to facilitate effector

secretion. However, our bound time measurements do not rule out changes in the

association rate constant (k_{on}), thereby resulting in a distinct secretion on/off states. To

explore this possibility, we quantified the rate of appearance of fluorescence foci in the

bound-state data as a proxy for k_{on}. This provides an estimate of the on-rate per cell over

the imaging period, normalized to the total number of fluorescently active molecules in a

cell (Fig. S4). This quantification suggests that SctQ appearance rate decreases 4-fold in

the secretion ON state while the effect of YopE presence is minimal. However, because

this metric is normalized to total cellular fluorescence, some variation is expected, and

small shifts are difficult to interpret with high confidence. To verify the role of cytosolic

and/or injectisome remodeling of Halo-SctQ in light of effector protein changes *and*

secretion-state dependence, which may be connected to k_{on}, we employed 3D single-

molecule localization and tracking microscopy.

We have previously published eYFP-SctQ tracking results for *Y. enterocolitica* under

secretion-on conditions and found cytosolic states in addition to a prominent stationary

state corresponding to injectisome-bound SctQ proteins[26,27]. Similarly, for Halo-SctQ,

under all conditions, there exist both slow, membrane associated states, as well as

cytosolic diffusing states (Fig. 5a). We focused on the slowly diffusing population; the

fraction of slow-moving SctQ trajectories (mean displacement < 4 nm/ms) serves as a

proxy for injectisome association and is conceptually linked to the association rate

constant (k_{on}). An increase in k_{on} would increase the fraction of injectisome bound

molecules, especially if bound time remains constant or increases. Alternatively, under

saturating conditions, all binding sites may already be occupied regardless of secretion

state, in which case we expect bound fraction to remain constant across conditions as

well. We quantified this fraction to assess injectisome association (Fig. 5b). While this

metric showed statistically significant (Kruskal–Wallis, $p<0.05$) differences across

conditions, the observed variation was small prompting us to verify other indicators of

injectisome engagement.

To alternatively assess potential variations in injectisome association, we

implemented an approach based on spatial clustering of single-molecule localizations. By

identifying localization clusters (these were found at the membrane, i.e. consistent with

injectisome locations) (Fig S5a) and classifying trajectories as cluster-associated or not, we

obtained an alternative more direct measure of injectisome binding. This analysis also

revealed that the fraction of cluster-associated trajectories did not change drastically

across conditions (Fig. 5c), indicating that the injectisome-bound fraction of SctQ does not

vary substantially with secretion state or effector presence. We can thus interpret the

bound time changes (i.e. k_{off} variations), in the absence of strong evidence for counter k_{on}

changes.

Figure 5: Secretion state modulates cytosolic diffusion of SctQ independently of effector presence. (a)

Representative single-molecule trajectories of Halo-SctQ under four experimental conditions. **(b)**

Quantification of the fraction of slow-moving trajectories (mean displacement < 4 nm/ms) per cell, used as a

proxy for injectisome association. **(c)** Cluster-based analysis of single-molecule localizations confirms

consistent injectisome-bound fractions across conditions, indicating that secretion state alters cytosolic

diffusion but not the fraction of injectisome-engaged SctQ.

The shape differences in the distributions shown in Fig 5a further show different

cytosolic diffusion behavior of SctQ in secretion ON and OFF conditions, potentially

reflecting changes in cytosolic subcomplex formation. This occurs independently of the

YopE-driven changes in k_{off} which we may conclude occur specifically at or when binding to

the injectisome. Together, these results reveal that secretion state, but not effector

presence, exert component-specific effects on cytosolic injectisome protein dynamics,

both at the injectisome and in the cytosol.

DISCUSSION

During infection, *Yersinia enterocolitica* must rapidly adapt to changing host environments, including rapid transitions between secretion ON and OFF states thought to be essential for both immune evasion and efficient colonization[30-34]. Activation of secretion, as well as deactivation, occurs rapidly[34,35] and the ability of *Y. enterocolitica* to respond to these external stimuli suggests a strong external sensing mechanism that modulate how (cytosolic) components of the injectisome function.

Here, we measured the bound times of SctQ, SctL, and SctN under secretion OFF and ON conditions in the presence and absence of the effector protein YopE. In secretion OFF conditions, we observed mono-exponential decay kinetics for SctQ, SctL, and SctN, indicating a single, transient bound state at the injectisome for each cytosolic complex protein. In the absence of effectors (T3SS$^{\Delta effector}$), each protein exhibited a distinct fitted bound time, SctQ having the shortest, followed by SctN and SctL, suggesting that these components do not both bind *and* unbind as a coordinated unit.

In contrast, the fitted bound times of all three proteins converged to approximately 4 seconds in the presence of the effector protein YopE. This synchronization of bound times suggests that effector presence alone, independent of active secretion, can modulate the binding kinetics of cytosolic components. This observation also suggests that effectors may be preloaded onto injectisomes in the secretion OFF state priming the system for rapid activation. At the molecular level, the convergence of bound times may reflect a shift toward a more uniform or stabilized interaction mode, potentially due to effector-induced

conformational remodeling of cytosolic injectisome complexes prior to secretion.

Convergence of bound times also raises the likelihood that SctQ, SctL, and SctN bind and

unbind as a part of the same subcomplex. An alternative explanation to intact subcomplex

dissociation would be that all other proteins disassemble immediately after at least one of

them unbinds from the injectisome. Such a model is supported gene deletion studies that

have established that cytosolic injectisome proteins are mutually dependent on each other

injectisome binding[2,25,36,37], expression of each protein is required for injectisome

assembly[2,11,38] and type 3 secretion[17-19,39,40].

Under secretion ON conditions, SctQ, SctL, and SctN all displayed a subpopulation

of long-lived binding events, resulting in bi-exponential decay kinetics. Detection of this

long-lived binding state is uniquely enabled by single-molecule measurements, which

allow direct quantification of injectisome bound times of individual protein molecules.

Importantly, we observed both short- and long-lived binding events at the same subcellular

locations, suggesting that individual injectisomes can engage proteins in different binding

modes, one mode being more stable than the other. This raises the intriguing possibility of

functional asymmetry within a single injectisome, either through structural heterogeneity

across pods or through dynamic switching between distinct operational states in time (our

pooled single-molecule data were obtained over a timescale of ~20 minutes). In support of

the structural heterogeneity hypothesis, cryoEM studies have shown that the ATPase SctN

forms an asymmetric hexamer at the base of the injectisome[41], raising the possibility that

asymmetry extends to other cytosolic injectisome components and contributes to the

observed kinetic heterogeneity. Future structural and computational studies will be

essential to determine whether the short and long bound states correspond to distinct

conformational or functional states of the injectisome under defined secretion conditions.

Among the three cytosolic components analyzed, SctQ consistently exhibited the

most pronounced changes in bound time across conditions and was particularly sensitive

to the presence of the effector protein YopE. Under secretion OFF conditions, the addition

of YopE nearly doubled SctQ bound time, bringing it into close alignment with SctL and

SctN. Under secretion ON conditions, SctQ again showed heightened sensitivity, with both

an increased short-term bound time and a long-lived subpopulation. This component-

specific response supports the notion that SctQ may serve in a key regulatory role within

the assembled injectisome.

Single-molecule tracking revealed that the injectisome-bound fraction of SctQ does

not vary substantially with secretion state or YopE effector presence. When modest

changes in bound time occur, such as a shift in secretion state within the same strain

(T3SS$^{\Delta effector}$ or T3SS^{YopE}), we can interpret these while recognizing that there are no

indications for large swings in k_{on}. Interestingly, our single-molecule tracking data suggests

that SctQ does undergo secretion-dependent remodeling in the cytosol, while the

presence of YopE did not appear to influence this behavior. Indeed, we did not observe

direct evidence for distinct YopE-containing SctQ complexes in the cytosol under secreting

or post-secreting conditions. The absence of a YopE effect is intriguing, especially given

prior reports suggesting cytosolic effector–chaperone handover. One possibility to be

explored is that because SctQ participates in a range of cytosolic complexes[20,36,42], only

some of them may include YopE and these subcomplexes could be of similar size, making

it difficult to resolve effector-specific interactions. Alternatively, the changes in bound time

may not originate from altered cytosolic subcomplexes at all, but rather from effector-

induced alterations at the injectisome itself. For example, YopE may stabilize the cytosolic

injectisome complexes or modify the binding interface through allosteric effects that

prolongs SctQ binding. The fact that we observe strong changes in bound time without

corresponding shifts in cytosolic diffusion supports this possibility.

Comparing effector-free and YopE-expressing strains allowed us to separate the

effects of secretion state from those with effector involvement. The fact that cytosolic

injectisome protein binding kinetics are altered even in the absence of active secretion

supports the idea that cytosolic remodeling, potentially through effector or possible

effector–chaperone interactions, can precondition cytosolic injectisome proteins before

injectisome association. With a known ability to form multiple cytosolic

subcomplexes[15,17,27] the cytosolic injectisome proteins can act as a dynamic mediator that

integrates substrate availability with injectisome engagement.

In conclusion, the results reported here establish that cytosolic interactions and

local injectisome binding kinetics both correlate with secretion state and the presence of

YopE. The emergence of two distinct binding states under secretion ON conditions, at the

same injectisome, points to a dynamic secretion mechanism, with possible functional

asymmetry in space or time. The kinetic heterogeneity may arise either from structural

variation across injectisomes or through temporal switching between functional modes.

Understanding how these states arise and are regulated will be essential for refining

models of substrate selection and export in the T3SS.

METHODS

Bacterial culture

Yersinia enterocolitica strains were grown overnight in Brain Heart Infusion (BHI, Sigma-Aldrich, St. Louis, MO) media supplemented with nalidixic acid (NaI) (35 µg/mL) and 2,6-diaminopimelic acid (DAP) (80 µg/mL) at 28°C with shaking. Additional antibiotics were included when appropriate: kanamycin (50 µg/mL) and/or chloramphenicol (25 µg/mL). On the day of imaging, overnight culture was back diluted to a target OD_{600} of 0.1-0.3 and grown at 28°C with shaking for 1-1.5 h. For secreting conditions, glycerol (4 mg/mL), $MgCl_2$ (20 mM), and EGTA (5 mM) were then added to the culture, and the culture was transferred to a 37°C water bath or incubator with shaking, for 3h to induce expression of the *yop* regulon and ensure secretion activation. For non-secreting conditions (post-secretion), cells were pelleted by centrifugation at 5,000 × *g* for 3 min and resuspended with fresh BHI supplemented with NaI, DAP, $CaCl_2$ (5 mM) and grown for an additional hour.

Strains and Plasmids

Plasmids were constructed using Gibson Assembly. Briefly, DNA fragments with overlapping ends were generated by Q5 PCR (New England Biolabs (NEB), Ipswich, MA) and assembled in a single isothermal reaction. Assembled plasmids were transformed into chemically competent *dh5α E coli* (NEB) and plated on LB agar supplemented with the appropriate antibiotic for selection. Individual colonies were cultured overnight, and plasmid DNA was isolated using a commercial miniprep kit (Qiagen, Hilden, Germany).

423   Plasmid sequences were verified by long-read sequencing (Plasmidsaurus, Eugene, OR).

424   Plasmids were transformed into the appropriate *Yersinia Enterocolitica* strain by

425   electroporation.

426   List of oligonucleotides

Name	Sequence	Used for plasmid
OC116	cacaggaaacacattcaacaatgGCAGAAATCGGTACTGGCTTTCCATTCGACCC	*pAH12::halo-sctN*
OC117	ccgacccacccaattgGCCGGAAATCTCGAGCGTCGACAGCCAG	*pAH12::halo-sctN*
OC118	cattgttgaatgtgtttcctgtgtgggcgcgcc	*pAH12::halo-sctN*
OC119	caattgggtgggtcgggcggct	*pAH12::halo-sctN*
OC163	GTA CCA CGC GGC AGT AAA ATA TCA TCA TTT ATT TCT ACA TCA CTG CCC CTG CCG GC	*pACAC::YopE*
OC164	AGC CTC CAC CTA GCC TTC ACA TCA ATG ACA GTA ATT GCT GCA TCT GTT GCC CC	*pACAC::YopE*

428   List of plasmids

Plasmid	Genotype	Reference
1	*pAH12::halo-sctN*	this study
2	*pAH12::halo*	this study
3	*pAH12::mEYFP*	this study
4	*pACAC::YopE*	this study

430   List of Strains

Strain	Genotype	Reference
IML421asd (ΔHOPEMTasd)	MRS40 *yopOΔ12-427 yopE21 yopHΔ11-352 yopM23 yopP23 yopT135 Δasd*	[4]
AD4461	IML421*asd halo-sctQ*	[36]
ADTM4525	IML421*asd halo-sctL*	[36]
AD4165	IML421*asd ΔsctN*	[25]

Secretion Assay

Cultures were grown as above, to desired secretion state. A small volume of culture was

taken to measure the OD_{600}. Main culture was pelleted (20,000 × *g*, 10 min, 4°C) and syringe

filtered with a 0.22 µm PVDF filter (Millipore, Burlington, MA). Supernatant was mixed with

trichloroacetic acid (TCA) to a final concentration of 10% w/v and incubated overnight at

4°C to precipitate proteins within the supernatant. Proteins were collected (20,000 × g,

20 min, 4°C) and washed with cold acetone (20,000 × g, 20 min, 4°C). The pellet was dried

at room temperature and resuspended in reducing SDS-PAGE loading buffer (SDS [3%,

wt/vol], glycerol [7.5%, wt/vol], 1.5% beta-mercaptoethanol, 0.0125% w/v Coomassie blue,

buffered with Tris [37.5 mM]; pH 7.0). Loading volume was normalized across samples ran

on the same gel, and proteins were separated by SDS-PAGE on 4–20% polyacrylamide gels

(Bio-rad, Hercules, CA). For visualization, the gels were stained with Coomassie Blue and

imaged with Bio-rad Gel Doc EZ imager.

In-gel fluorescence Assay

Cultures were grown as above, to desired secretion state. HaloTag fusions were incubated

with JFX-554 (Promega, Madison, WI) for the last 30 min of culturing. A small volume of

culture was taken to measure the OD_{600}. Culture was pelleted (5,000 x g, 5 min) and lysed in

a buffer of 20 mM Tris HCl, 50 mM NaCl, pH 7.4, supplemented with 0.2 mg/mL of lysozyme

and cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche, Basel, Switzerland).

Samples were resuspended in SDS-PAGE loading buffer and heated for 5 minutes at 98°C

before loading. Loading volume was normalized across samples ran on the same gel, and

proteins were separated by SDS-PAGE on 4–20% polyacrylamide gels (Bio-rad). For

visualization of in-gel JFX-554 fluorescence, a Biorad ChemiDoc MP system was used.

Cell Plating for Microscopy

Cultures were grown as described above. In the case of HaloTag visualization, Janelia Fluor

HaloTag ligand dye, JFX-554 (Promega) was added to the culture 30 minutes prior to

imaging. The appropriate dye concentration was strain dependent: 0.5 nM for Halo-SctQ, 2

nM for Halo-SctL, 2nM for Halo-SctN. Cells were pelleted by centrifugation at 5,000 × g for

3 min and washed three times in fresh M9 minimal media. After the three washes, the

remaining pellet was resuspended in M9 supplemented with DAP, $MgCl_2$, glycerol, and

EGTA (secreting conditions) or $CaCl_2$. For super-resolution fluorescent imaging, fluorescent

fiducial markers (Invitrogen) were added to cell suspensions. Cells were finally plated on

1.5% agarose pads in M9 containing DAP and EGTA or $CaCl_2$.

Optical Setup

Image data were acquired on a custom-built inverted fluorescence microscope based on

the RM21 platform (Mad City Labs, Madison, WI), as previously described[26,43] . Immersion

oil was placed between the lens objective (UPLSAPO 100; numerical aperture [NA], 1.4)

and the glass coverslip (VWR, Radnor, PA; number 1.5; 22 mm by 22 mm). A 561 nm laser

(Coherent, Santa Clara, CA; Genesis MX561-4000 MTM) was used for excitation of JFX-554

(Promega Madison, WI) (~600 mW cm^{-2}). Zero-order quarter-wave plates (Thorlabs,

Newton, NJ, WPQ05M-561) were used to circularly polarize the excitation laser. A dichroic

mirror (Chroma zt440/488/561rpc) was used to separate excitation/activation light from

fluorescence emission of JFX-554. Fluorescence emission was passed through a filter set

including long-pass (514nm), short- pass (800 nm), and notch filter 561 nm/10bp

(Semrock, Rochester, NY; LP02-514RU-25, Chroma, Bellows Falls, VT; ET700SP-2P8, and

Chroma, Bellows Falls, VT; ZET561NF, respectively). In the case of 3D super-resolution

imaging, the emission path contains a wavelength-specific dielectric phase mask (Double

Helix, LLC, Boulder, CO) that is placed in the Fourier plane of the microscope to generate a

double-helix point spread function (DHPSF)[44,45]. The fluorescence signal is detected on a

scientific complementary metal oxide semiconductor (sCMOS) camera (Hamamatsu,

Bridgewater, NJ; ORCA-Flash 4.0 V2). A flip mirror in the emission pathway enables toggling

the microscope between fluorescence imaging and phase-contrast imaging modes without

having to change the lens objective of the microscope. Illumination for phase contrast

imaging was achieved with an LED at 625 nm (Thorlabs, Newton NJ; M625L3).

Imaging was performed on a custom-built, dual-color inverted fluorescence microscope.

The laser intensity was attenuated significantly. Up to 2500 frames were collected per field-

of-view with an exposure time of 500 ms.

Long Exposure Data processing

Broadly, image processing occurred in a two-step process. First, potential binding events

were identified using ThunderSTORM[46] generating regions of interest for further analysis.

Second, intensity traces over time were fitted with a Hidden Markov Model to distinguish

between a bound and background state.

Specifically, prior to PSF fitting, images from each field-of-view were background

subtracted and cropped as needed. PSFs were fit using ThunderSTORM in ImageJ using the

difference of Gaussians (DoG) filter option with a threshold of 1.5*std(Wave.F1)[46].

Candidate binding events were further narrowed down with a requirement for there to be a

localization within 50 nm at least 2 times per 1000 frames. The image stack and its

corresponding ThunderSTORM data was imported into MATLAB. Images were background

subtracted, normalized and converted to photoelectron units. A simple nearest-neighbor

linker associated detections across frames (distance threshold 300 nm; maximum frame

gap 15) to form candidate bound trajectories.

For every frame within a trajectory, signal was quantified as the sum in a 5×5 pixel ROI

centered on the trajectory position. Pooled photon values were modeled with a Gaussian

mixture model (GMM; 3 components) to estimate background and signal intensity

distributions. Per-track ROI time series were then segmented with a two-state hidden

Markov model (HMM; background ↔ signal): initial state means/variances were set from

the GMM, a conservative transition matrix was used and the most probable state sequence

was decoded by Viterbi. Quality gates rejected traces with (i) peak intensity ≤ background

upper bound, (ii) minimum intensity ≥ signal lower bound, (iii) dynamic range below the

GMM-implied separation, or (iv) substantial signal/background standard-deviation overlap.

Error codes were logged for each exclusion. In addition, any bound state event longer than

40 frames (20 seconds) was inspected and manually approved. Remaining signal segments

were taken as valid dwell events; durations (frames) were converted to time using 0.5

s/frame and compiled across tracks.

Determination of t$_{bound}$

To characterize the distribution of bound-state lifetimes, dwell time measurements were

extracted for each condition and protein variant and binned into normalized histograms. To

reduce the influence of short-duration events on the fit, the first histogram bin (0-0.5

seconds), a group that is disproportionally undercounted, was excluded from model fitting.

For each distribution, both single- and double-exponential decay models were fitted to the

data using nonlinear least-squares regression. The single-exponential model was defined

as $y(t) = A\,e^{-\lambda t}$ and the double-exponential model as $y(t) = A_1 e^{-\lambda_1 t} + A_2 e^{-\lambda_2 t}$ with

time constants $\tau_i = \frac{1}{\lambda_i}$.

Model selection was based on comparison of the Akaike Information Criterion (AIC), with

the bi-exponential model favored if AIC was reduced by more than 2 units. In addition, if a

state in the bi-exponential fit contributed less than 2% of the population, the mono-

exponential model was favored. Goodness-of-fit was assessed via the coefficient of

determination (R^2). For each fit, 99% confidence intervals were estimated by propagation

of uncertainty using the Jacobian and parameter covariance matrix. Fits and their

confidence intervals were overlaid on probability density plots to visualize lifetime

distributions for each experimental condition.

Cluster detection and enrichment analysis

Single-molecule trajectories of SctQ, SctL, and SctN recorded under secretion-ON

conditions were analyzed to test whether long-lived binding events were spatially enriched.

Approved trajectories were converted into individual binding events, each represented by

the mean localization position and event duration. Events lasting ≥ 30 frames were

classified as long.

Spatial event clusters were identified using DBSCAN (ε = 100 nm, minPts = 2). For each

field of view, the observed fraction of long events within each cluster was compared with

the global fraction of long events using a one-sided hypergeometric test. Resulting p-

values were corrected for multiple testing by the Benjamini–Hochberg procedure (α =

0.05), and significant clusters were optionally validated by 2,000 random label-shuffle

permutations. Clusters were categorized as mixed, pure-long, or pure-short based on the

composition of event durations.

Super-resolution fluorescence imaging & Data Processing

Up to 20,000 frames are collected per field of view with an exposure time of 25 ms. Phase

contrast images were taken before and after fluorescence imaging. Raw data were

processed in MATLAB using a modified version of easy-DHPSF software[26,47]. Fluorescent

fiducial markers were used for sample drift correction. Phase contrast images were

segmented using Omnipose[48]. Single-molecule localizations were registered to phase

contrast masks using a two-step affine transformation in MATLAB, as described

previously[26]. Localizations outside of cells were discarded from further analysis using axial

bounds and Omnipose-derived cell outlines.

MSD / cumulative displacement tracking

To derive single-molecule displacements, localizations in subsequent frames were linked

into trajectories. A maximum linking distance of 2.5 µm was used for linking analysis, and

multiple localizations that were present at the same time within a single cell were

discarded to prevent misassignment of molecules.

For each trajectory, the cumulative displacement was calculated as the running sum of all

stepwise displacements between consecutive localizations,

576
$$CD_n = \sum_{i=2}^{n} \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2 + (z_i - z_{i-1})^2}$$

where x_i, y_i, z_i are the 3D coordinates of the molecule at frame i. The cumulative

displacement versus time relationship was used to visualize overall molecular motion and

to compute the mean step size or slope of displacement accumulation over time. This

slope provided a direct, model-free metric of apparent mobility for each trajectory. The

plotted mean displacement normalized by time is thus an average of this displacement,

which is equivalent to the slope of the cumulative displacement curve over time.

REFERENCES

Galán, J. E. & Collmer, A. Type III Secretion Machines: Bacterial Devices for Protein Delivery into Host Cells. *Science* **284**, 1322-1328 (1999). https://doi.org/doi:10.1126/science.284.5418.1322

Hu, B., Lara-Tejero, M., Kong, Q., Galán, J. E. & Liu, J. In Situ Molecular Architecture of the Salmonella Type III Secretion Machine. *Cell* **168**, 1065-1074.e1010 (2017). https://doi.org/http://dx.doi.org/10.1016/j.cell.2017.02.022

Butan, C., Lara-Tejero, M., Li, W., Liu, J. & Galán, J. E. High-resolution view of the type III secretion export apparatus in situ reveals membrane remodeling and a secretion pathway. *Proc Natl Acad Sci U S A* **116**, 24786-24795 (2019). https://doi.org/10.1073/pnas.1916331116

Kudryashev, M. *et al.* In situ structural analysis of the Yersinia enterocolitica injectisome. *Elife* **2**, e00792 (2013). https://doi.org/10.7554/eLife.00792

Miletic, S. *et al.* Substrate-engaged type III secretion system structures reveal gating mechanism for unfolded protein translocation. *Nat Commun* **12**, 1546 (2021). https://doi.org/10.1038/s41467-021-21143-1

Cornelis, G. R. The Yersinia Ysc-Yop 'type III' weaponry. *Nat Rev Mol Cell Biol* **3**, 742-752 (2002). https://doi.org/10.1038/nrm932

Dewoody, R. S., Merritt, P. M. & Marketon, M. M. Regulation of the Yersinia type III secretion system: traffic control. *Front Cell Infect Microbiol* **3**, 4 (2013). https://doi.org/10.3389/fcimb.2013.00004

8   Mehigh, R. J., Sample, A. K. & Brubaker, R. R. Expression of the low calcium

response in Yersinia pestis. *Microb Pathog* **6**, 203-217 (1989).

https://doi.org/10.1016/0882-4010(89)90070-3

9   Diepold, A. & Wagner, S. Assembly of the bacterial type III secretion machinery.

FEMS Microbiol Rev **38**, 802-822 (2014). https://doi.org/10.1111/1574-6976.12061

10   Bergeron, J. R. C. & Marlovits, T. C. Cryo-EM of the injectisome and type III secretion

systems. *Curr Opin Struct Biol* **75**, 102403 (2022).

https://doi.org/10.1016/j.sbi.2022.102403

11   Hu, B. *et al*. Visualization of the type III secretion sorting platform of Shigella

flexneri. *Proc Natl Acad Sci U S A* **112**, 1047-1052 (2015).

https://doi.org/10.1073/pnas.1411610112

12   Makino, F. *et al*. The Architecture of the Cytoplasmic Region of Type III Secretion

Systems. *Scientific Reports* **6**, 33341 (2016). https://doi.org/10.1038/srep33341

http://www.nature.com/articles/srep33341#supplementary-information

13   Berger, C. *et al*. Structure of the Yersinia injectisome in intracellular host cell

phagosomes revealed by cryo FIB electron tomography. *Journal of Structural Biology*

213, 107701 (2021). https://doi.org/https://doi.org/10.1016/j.jsb.2021.107701

14   Diepold, A. & Armitage, J. P. Type III secretion systems: the bacterial flagellum and

the injectisome. *Philos Trans R Soc Lond B Biol Sci* **370** (2015).

https://doi.org/10.1098/rstb.2015.0020

15   Bzymek, K. P., Hamaoka, B. Y. & Ghosh, P. Two translation products of Yersinia yscQ

assemble to form a complex essential to type III secretion. *Biochemistry* **51**, 1669-

1677 (2012). https://doi.org/10.1021/bi201792p

16   Yu, X. J., Liu, M., Matthews, S. & Holden, D. W. Tandem translation generates a

chaperone for the Salmonella type III secretion system protein SsaQ. *J Biol Chem*

286, 36098-36107 (2011). https://doi.org/10.1074/jbc.M111.278663

17   Jackson, M. W. & Plano, G. V. Interactions between type III secretion apparatus

components from *Yersinia pestis* detected using the yeast two-hybrid system. *FEMS*

Microbiology Letters **186**, 85-90 (2000). https://doi.org/10.1111/j.1574-

6968.2000.tb09086.x

18   Morita-Ishihara, T. *et al*. Shigella Spa33 is an essential C-ring component of type III

secretion machinery. *J Biol Chem* **281**, 599-607 (2006).

https://doi.org/10.1074/jbc.M509644200

19   Spaeth, K. E., Chen, Y. S. & Valdivia, R. H. The Chlamydia type III secretion system C-

ring engages a chaperone-effector protein complex. *PLoS Pathog* **5**, e1000579

(2009). https://doi.org/10.1371/journal.ppat.1000579

20   Soto, J. E., Galan, J. E. & Lara-Tejero, M. Assembly and architecture of the type III

secretion sorting platform. *Proc Natl Acad Sci U S A* **119**, e2218010119 (2022).

https://doi.org/10.1073/pnas.2218010119

21   Bernal, I. *et al*. Molecular Organization of Soluble Type III Secretion System Sorting

Platform Complexes. *J Mol Biol* (2019). https://doi.org/10.1016/j.jmb.2019.07.004

22   Barison, N., Lambers, J., Hurwitz, R. & Kolbe, M. Interaction of MxiG with the

641   cytosolic complex of the type III secretion system controls Shigella virulence. *FASEB*

642   *J* **26**, 1717-1726 (2012). https://doi.org/10.1096/fj.11-197160

643   23   Wimmi, S. *et al.* Cytosolic sorting platform complexes shuttle type III secretion

644   system effectors to the injectisome in Yersinia enterocolitica. *Nat Microbiol* **9**, 185-

645   199 (2024). https://doi.org/10.1038/s41564-023-01545-1

646   24   Lara-Tejero, M., Kato, J., Wagner, S., Liu, X. & Galan, J. E. A Sorting Platform

647   Determines the Order of Protein Secretion in Bacterial Type III Systems. *Science*

648   **331**, 1188-1191 (2011). https://doi.org/10.1126/science.1201476

649   25   Diepold, A., Kudryashev, M., Delalez, N. J., Berry, R. M. & Armitage, J. P.

650   Composition, formation, and regulation of the cytosolic c-ring, a dynamic

651   component of the type III secretion injectisome. *PLoS Biol* **13**, e1002039 (2015).

652   https://doi.org/10.1371/journal.pbio.1002039

653   26   Rocha, J. M. *et al.* Single-molecule tracking in live Yersinia enterocolitica reveals

654   distinct cytosolic complexes of injectisome subunits. *Integr Biol (Camb)* **10**, 502-

655   515 (2018). https://doi.org/10.1039/c8ib00075a

656   27   Prindle, J. R., Wang, Y., Rocha, J. M., Diepold, A. & Gahlmann, A. Distinct Cytosolic

657   Complexes Containing the Type III Secretion System ATPase Resolved by Three-

658   Dimensional Single-Molecule Tracking in Live Yersinia enterocolitica. *Microbiol*

659   *Spectr* **10**, e0174422 (2022). https://doi.org/10.1128/spectrum.01744-22

28   Pintor, K. *et al*. In situ analysis of type III secretion chaperone proteins indicates a

cytosolic handover of virulence effectors. *FEMS Microbes* **6**, xtaf010 (2025).

https://doi.org/10.1093/femsmc/xtaf010

29   Gilzer, D., Schreiner, M. & Niemann, H. H. Direct interaction of a chaperone-bound

type III secretion substrate with the export gate. *Nat Commun* **13**, 2858 (2022).

https://doi.org/10.1038/s41467-022-30487-1

30   Grosdent, N., Maridonneau-Parini, I., Sory, M. P. & Cornelis, G. R. Role of Yops and

adhesins in resistance of Yersinia enterocolitica to phagocytosis. *Infect Immun* **70**,

4165-4176 (2002). https://doi.org/10.1128/iai.70.8.4165-4176.2002

31   Navarro, L., Alto, N. M. & Dixon, J. E. Functions of the Yersinia effector proteins in

inhibiting host immune responses. *Curr Opin Microbiol* **8**, 21-27 (2005).

https://doi.org/10.1016/j.mib.2004.12.014

32   Galán, J. E. Common themes in the design and function of bacterial effectors. *Cell*

Host Microbe **5**, 571-579 (2009). https://doi.org/10.1016/j.chom.2009.04.008

33   Zwack, E. E. *et al*. Inflammasome activation in response to the Yersinia type III

secretion system requires hyperinjection of translocon proteins YopB and YopD.

mBio **6**, e02095-02014 (2015). https://doi.org/10.1128/mBio.02095-14

34   Milne-Davies, B. *et al*. Life After Secretion-Yersinia enterocolitica Rapidly Toggles

Effector Secretion and Can Resume Cell Division in Response to Changing External

Conditions. *Front Microbiol* **10**, 2128 (2019).

https://doi.org/10.3389/fmicb.2019.02128

35   Enninga, J., Mounier, J., Sansonetti, P. & Tran Van Nhieu, G. Secretion of type III

effectors into host cells in real time. *Nat Methods* **2**, 959-965 (2005).

https://doi.org/10.1038/nmeth804

36   Diepold, A. *et al*. A dynamic and adaptive network of cytosolic interactions governs

protein export by the T3SS injectisome. *Nat Commun* **8**, 15940 (2017).

https://doi.org/10.1038/ncomms15940

37   Diepold, A. *et al*. Deciphering the assembly of the Yersinia type III secretion

injectisome. *EMBO J* **29**, 1928-1940 (2010). https://doi.org/10.1038/emboj.2010.84

38   Jouihri, N. *et al*. MxiK and MxiN interact with the Spa47 ATPase and are required for

transit of the needle components MxiH and MxiI, but not of Ipa proteins, through the

type III secretion apparatus of Shigella flexneri. *Molecular Microbiology* **49**, 755-767

(2004). https://doi.org/10.1046/j.1365-2958.2003.03590.x

39   Johnson, S. & Blocker, A. Characterization of soluble complexes of the Shigella

flexneri type III secretion system ATPase. *FEMS Microbiol Lett* **286**, 274-278 (2008).

https://doi.org/10.1111/j.1574-6968.2008.01284.x

40   Hueck, C. J. Type III Protein Secretion Systems in Bacterial Pathogens of Animals

and Plants. *Microbiology and Molecular Biology Reviews* **62**, 379-433 (1998).

41   Majewski, D. D. *et al*. Cryo-EM structure of the homohexameric T3SS ATPase-central

stalk complex reveals rotary ATPase-like asymmetry. *Nature Communications* **10**,

626 (2019). https://doi.org/10.1038/s41467-019-08477-7

42   Marcos-Vilchis, A. *et al*. On the role of the sorting platform in hierarchical type III

secretion regulation in enteropathogenic Escherichia coli. *Journal of Bacteriology*

207 (2025). https://doi.org/10.1128/jb.00446-24

43   Rocha, J., Corbitt, J., Yan, T., Richardson, C. & Gahlmann, A. Resolving Cytosolic

Diffusive States in Bacteria by Single-Molecule Tracking. *Biophys J* **116**, 1970-1983

(2019). https://doi.org/10.1016/j.bpj.2019.03.039

44   Pavani, S. R. P. *et al*. Three-dimensional, single-molecule fluorescence imaging

beyond the diffraction limit by using a double-helix point spread function.

Proceedings of the National Academy of Sciences of the United States of America

106, 2995-2999 (2009). https://doi.org/10.1073/pnas.0900245106

45   Thompson, M. A., Lew, M. D., Badieirostami, M. & Moerner, W. E. Localizing and

tracking single nanoscale emitters in three dimensions with high spatiotemporal

resolution using a double-helix point spread function. *Nano Letters* **10**, 211-218

(2010). https://doi.org/10.1021/nl903295p

46   Ovesny, M., Krizek, P., Borkovec, J., Svindrych, Z. & Hagen, G. M. ThunderSTORM: a

comprehensive ImageJ plug-in for PALM and STORM data analysis and super-

resolution imaging. *Bioinformatics* **30**, 2389-2390 (2014).

https://doi.org/10.1093/bioinformatics/btu202

47   Lew, M. D., von Diezmann, A. R. S. & Moerner, W. E. Easy-DHPSF open-source

software for three-dimensional localization of single molecules with precision

beyond the optical diffraction limit. *Protocol Exchange*,

doi:10.1038/protex.2013.1026 (2013). https://doi.org/doi:10.1038/protex.2013.026

48   Cutler, K. J. *et al.* Omnipose: a high-precision morphology-independent solution for

bacterial cell segmentation. *Nat Methods* **19**, 1438-1448 (2022).

https://doi.org/10.1038/s41592-022-01639-4