Title: Molecular evolution of octopamine receptors in *Drosophila*

Authors: Mengye Yang[1], Jolie A. Carlisle[1], Ben R. Hopkins[2,3], Mariana F. Wolfner[1*]

* Corresponding author: mfw5@cornell.edu

Affiliations:

1. Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853

2. Department of Evolution and Ecology, University of California, Davis, CA 95616

3. Department of Molecular Genetics and Microbiology, University of Florida, Gainesville, FL
32610

Running head: Octopamine receptor evolution in *Drosophila*

Keywords: *Drosophila*; octopamine receptor; molecular evolution; positive selection;
conservation; reproduction

13
14

Abstract

Octopamine (OA), the insect analog of noradrenaline, plays important roles in diverse behavioral and physiological processes, from modulating fight-or-flight behavior to regulating post-mating ovulation. In *Drosophila*, six OA receptors have been identified: Oamb, Octα2R, Octβ1R, Octβ2R, Octβ3R, and Oct-TyrR, and they have been linked to different behavioral and physiological processes. Here, we investigated the evolutionary characteristics of these receptors across *Drosophila* species. We found that OA receptors are generally found as single-copy genes. Notably, Octβ2R and Octβ3R exhibit positive selection within the *melanogaster* group, though in different structural regions from one another. The positively selected sites in Octβ2R are exclusively located in regions important for ligand binding, whereas those in Octβ3R are predominantly found in regions crucial for signal transduction. Interestingly, Octβ2R remains highly conserved outside the *melanogaster* group, so the detection of positive selection in its ligand-binding related domains within this clade raises the possibility that it has evolved an additional, *melanogaster*-specific ligand interaction(s), among other potential reasons. These findings highlight the evolutionary flexibility of aminergic signaling and suggest lineage-specific adaptations of OA receptor function in *Drosophila*, likely shaped by lineage-specific selective pressures.

Article summary

Octopamine is a chemical messenger in insects, analogous to noradrenaline in vertebrates, and influences many processes such as fight-or-flight responses, locomotion, and reproduction. In fruit flies and other insects, multiple octopamine receptors detect and respond to this signal. We explored how these receptors have changed over millions of years across various fly species. We found that while some receptors remain similar, others have evolved more rapidly, possibly due to changes in the insects' lifestyles, environments, or other biological demands. Understanding these differences can help reveal how insects fine-tune their behaviors and adapt to diverse challenges.

Introduction

Biogenic amines are small signaling molecules that play multiple roles in regulating physiological and behavioral functions across both mammals and insects. In mammals, noradrenaline and adrenaline mediate fight-or-flight responses and other key processes. In insects, however, these catecholamines are absent or present only at very low levels. Instead, insects rely on structurally and functionally analogous molecules: tyramine (TA) and octopamine (OA). The amino acid tyrosine is first converted to TA by tyrosine decarboxylase, and TA is subsequently converted to OA by tyramine β-hydroxylase. Although TA serves as the biosynthetic precursor to OA, both can act as independent neurotransmitters (Roeder, 2005). In *Drosophila melanogaster* (fruit fly), OA function has been particularly well-studied, revealing its pleiotropic roles in diverse processes including locomotion, aggression, learning, memory, sleep, energy intake and expenditure, as well as reproduction (Farooqui, 2012; Roeder, 2005, 2020; White et al., 2021).

OA exerts its diverse physiological and behavioral effects by binding to and activating its receptors, all of which belong to the G protein-coupled receptor (GPCR) family. In *Drosophila*, these receptors are classified into four groups based on their structural and signaling similarities to vertebrate adrenergic receptors: Oamb (α-adrenergic-like octopamine receptor), Octα2R (α$_2$-adrenergic-like octopamine receptor), three β-adrenergic-like receptors (Octβ1R, Octβ2R, and Octβ3R), and the octopamine/tyramine receptor Oct-TyrR (Evans & Maqueira, 2005; Qi et al.,

2017). Despite being functionally related proteins belonging to the same protein family, the
relationships among the OA receptors remains unresolved. Previously, two independent
evolutionary studies constructed phylogenetic trees of biogenic amine receptors across species.
As expected, β-adrenergic-like octopamine receptors clustered together; however, the
phylogenetic and evolutionary relationships with and among the other OA receptors remained
unclear due to weak branch support (Qi et al., 2017; Zhang et al., 2023). Upon activation, Oamb
promotes increases in intracellular calcium and cAMP levels and has been implicated in
processes such as follicle rupture, ovulation, sperm storage, sleep/wake regulation, appetitive
learning and memory, insulin-like peptide transcription, and male aggression (Avila et al., 2012;
Balfanz et al., 2005; Burke et al., 2012; Crocker et al., 2010; Deady & Sun, 2015; Han et al.,
1998; Huetteroth et al., 2015; Kim et al., 2013; Lee et al., 2009; Luo et al., 2014). Octα2R, by
contrast, reduces cAMP production and has been linked to locomotion, grooming behavior, and
starvation-induced hyperactivity (Nakagawa et al., 2022; Qi et al., 2017). The OctβRs
predominantly signal through cAMP, and Octβ1R is involved in olfactory learning, exercise
adaptation, and hunger-driven modulation of female receptivity (Sabandal et al., 2020; Sujkowski
et al., 2020; Sun et al., 2023); Octβ2R regulates ovulation, locomotor activity, anesthesia-
resistant memory, and sleep, as well as stimulates synaptic growth, an effect antagonized by
Octβ1R (Koon et al., 2011; Koon & Budnik, 2012; Li et al., 2015; Lim et al., 2014; Wu et al., 2013;
Zhao et al., 2021); while Octβ3R has been implicated in metamorphosis and appetitive motivation
(Ohhara et al., 2015; Zhang et al., 2013). Finally, Oct-TyrR is sensitive to both TA and OA, with
TA being slightly more potent in inhibiting adenylate cyclase activity, while OA more strongly
stimulates calcium signaling (Robb et al., 1994). Oct-TyrR has been shown to affect chemotaxis
behavior and startle responses by modulating downstream dopaminergic neuron activity (Ma et
al., 2016).

Genes associated with reproduction not infrequently evolve more rapidly than non-reproductive
genes, sometimes exhibiting elevated sequence divergence potentially driven by sexual
selection, sexual conflict, or relaxed selective constraints (Carlisle & Swanson, 2021; Clark et al.,
2006; Dapper & Wade, 2020; Swanson & Vacquier, 2002; Wilburn & Swanson, 2016). Male
seminal fluid proteins trigger the female post-mating response, a potential battleground of male x
female interaction. Indeed, 7-12% of male seminal fluid proteins show signatures of positive
selection potentially driven by sexual selection or sexual conflict (Haerty et al., 2007; Patlar et al.,
2021; Swanson, Clark, et al., 2001; Wong & Wolfner, 2012). Although fewer studies have
addressed the evolution of female reproductive proteins, growing evidence suggests that positive
selection also acts on the female side (Galindo et al., 2003; McGeary & Findlay, 2020; Moyle et
al., 2021; Swanson et al., 2001). Given that multiple OA receptors are involved in reproductive
processes (Avila et al., 2012; Deady & Sun, 2015; Lee et al., 2009; Li et al., 2015; Lim et al.,
2014; Sun et al., 2023), it is of particular interest to explore the evolutionary dynamics of this
receptor group.

Here, we identified OA receptor orthologs across 27 *Drosophila* species and detected two
lineage-specific tandem duplication events using synteny analysis. Nevertheless, OA receptors
are generally maintained as single-copy genes. Using a suite of molecular evolution analyses, we
found evidence of positive selection on Octβ2R and Octβ3R within the *melanogaster* group.
Interestingly, we did not detect evidence for positive selection of Octβ2R in the *virilis-repleta*
radiation nor in mosquitoes, suggesting that the positive selection observed in residues important
for ligand binding within *melanogaster* species may reflect lineage-specific selective pressures.
These selective pressures could include sexual selection on reproductive function, pathogen-
mediated selection, ecological adaptation involving receptor modulation, or perhaps co-option to
bind a ligand other than OA. Taken together, these results reveal distinct evolutionary trajectories
shaping OA receptor evolution.

Materials and Methods

Synteny analysis and tandem duplication detection

To identify potential tandem duplications of OA receptors, we first determined their syntenic regions by focusing on the three closest upstream and downstream genes flanking each OA receptor in the *Drosophila melanogaster* genome, as identified using the JBrowse map within FlyBase (flybase.org). Using three neighboring genes provides sufficient local context to detect tandem duplications without extending into unrelated genomic regions. We then identified orthologs of these seven genes (including the focal OA receptor) in 27 *Drosophila* species whose genomes had been annotated by the automated NCBI Gnomon prediction pipeline. The *D. melanogaster* sequence of each gene was used as the query in the tBLASTn search against the annotated transcriptomes for each species. The species we tested were: *melanogaster*, *simulans*, *mauritiana*, *sechellia*, *erecta*, *yakuba*, *santomea*, *teissieri*, *takahashii*, *suzukii*, *subpulchrella*, *biarmipes*, *elegans*, *rhopaloa*, *ficusphila*, *kikkawai*, *ananassae*, *persimilis*, *pseudoobscura*, *willistoni*, *busckii*, *grimshawi*, *mojavensis*, *hydei*, *virilis*, *innubila*, and *nasuta*. We also performed reciprocal tBLASTn searches against *D. melanogaster* transcriptome to strengthen the confidence in ortholog prediction. Their chromosomal locations were confirmed using the NCBI Genome Data Viewer, and OA receptor copy number was determined for each species. For species with more than one copy of an OA receptor, RNA-seq exon coverage tracks were inspected on the species' genome browsers to confirm transcription of each identified paralog. For OA receptors, at least one side of the flanking genes showed some level of conservation, supporting their placement within the expected genomic context. For flanking genes that differed from those in *D. melanogaster*, we used Gnomon predictions available on NCBI to confirm their identities. Synteny metrics are provided in File S1, and more details are available in File S6.

High-throughput methods such as those used in OrthoDB (Tegenfeldt et al., 2025) and DROSOMA (Thiébaut et al., 2024) provide large-scale resources for ortholog annotation, but they rely on automated pipelines that can misclassify relationships, particularly in cases of gene duplication or lineage-specific loss (Carlisle et al., 2024). Our approach involves nonautomated, detailed, and targeted characterization of receptor genes across species. While our results largely agree with OrthoDB and DROSOMA for most octopamine receptors existing as single-copy in species examined, these databases did not reliably find the duplicates that we detected. The duplicate in *D. busckii* of *Octβ3R* was not detected in either DROSOMA or OrthoDB and the duplicate in *D. sechellia* of *Octα2R* was not detected in DROSOMA but was detected in OrthoDB. This inconsistency between our results and ortholog predictors and the inconsistency between ortholog predictors themselves highlight the value of our detailed, targeted investigations, such as that presented here.

Molecular evolution analyses

We performed PAML analyses on each OA receptor across 15 species within the *melanogaster* group, as listed in Fig S4. Only isoforms detectable in more than 10 species by Gnomon prediction were included in the analysis. For Octα2R, *Dsec* orthologs were excluded due to the presence of two copies in this species. Protein sequence alignments were generated by Clustal Omega (Madeira et al., 2024), which were then converted into codon-based DNA alignments with PAL2NAL (Suyama et al., 2006). Protein alignment statistics can be found in File S7. Maximum likelihood trees were constructed with reference to two established phylogenies: a high-confidence phylogeny of 155 *Drosophila* species (Suvorov et al., 2022) and an additional tree including *D. santomea* (Hopkins et al., 2024) to determine its phylogenetic position. The codeml program from the PAML package (Yang, 2007) was used to calculate an overall ω estimate for the whole sequence under model M0 and to perform site tests by comparing model M8, which allows for a class of sites with $\omega > 1$, against the null models M7 and M8a using likelihood ratio tests (Swanson et al., 2003; Yang et al., 2000). The 'cleandata' option was enabled in codeml to remove alignment sites with gaps or ambiguous data. For genes where model M8 provided a

significantly better fit than models M7 and M8a, the BEB approach was applied to identify
positively selected sites at a 0.9 confidence level.

For Octβ2R, PAML analyses were also performed within the *Drosophila virilis-repleta* radiation
and across mosquito species, following the similar procedures described above. Ortholog
prediction was conducted in the same manner as for the 27 *Drosophila* species (see the Synteny
analysis and tandem duplication detection section), and PAML analyses were carried out as
described in the preceding paragraph. Within the *virilis-repleta* radiation, seven species (*virilis*,
novamexicana, *hydei*, *navojoa*, *mojavensis*, *arizonae*, and *montana*) were chosen based on the
availability of transcriptome data. In mosquitoes, PAML tests were performed separately within
the *Cellia*, *Anophelinae*, and *Culicidae* lineages. The species included in these analyses, along
with their phylogenetic relationships (Neafsey et al., 2015; Suvorov et al., 2022), are shown in Fig
4B.

MEME (Mixed Effects Model of Evolution) analyses were conducted using the Datamonkey
Adaptive Evolution Server (Weaver et al., 2018) with default parameters and 100 resamples. The
maximum likelihood trees and DNA alignments used in the PAML analyses were also used as
input for these analyses.

Codon usage bias and GC content analysis

CDS of Octβ2R and Octβ3R from 15 species in the *Drosophila melanogaster* group were
analyzed to assess codon usage bias and GC content. Sequences were first cleaned by
removing gaps (-). Codon usage metrics were calculated using the coRdon R package (v1.20.0),
including the effective number of codons (ENC) to quantify overall codon usage bias. Overall GC
content and GC content in the third codon position (GC3) were computed using the seqinr R
package (v 4.2.36). For GC3, the third nucleotide of each codon was extracted and the proportion
of G or C nucleotides was calculated per sequence.

ENC values range from 20 (extreme codon bias) to 61 (no bias), with intermediate values
indicating moderate codon usage bias. GC and GC3 metrics were used to evaluate potential
nucleotide composition shifts that might influence ω estimates. All analyses were performed in R
(v4.3.2) on gap-free, full-length CDS alignments.

Results and Discussion

OA receptors are found in single copy in most of the *Drosophila* genomes we analyzed

To examine the evolutionary history of OA receptors, we first identified the orthologs of OA
receptors across 27 *Drosophila* species with available transcriptome data, which, combined with
the NCBI Gnomon gene predictions, provides comprehensive and well-supported receptor
annotations. Searches against transcriptomes, rather than genomes, are more sensitive to detect
paralogs since expressed coding sequences do not have introns that complicate genome-based
searches and the database that is being searched against is much smaller. By examining
syntenic regions, we found that most OA receptor family members are retained as single-copy
genes across species (Fig 1-2, Fig S1-3, File S1), reflecting evolutionary conservation and
functional constraint. This is likely important for maintaining a tightly regulated neuromodulatory
system, where altered gene dosage or expression levels could lead to signaling imbalances and
physiological dysfunction.

Interestingly and in contrast, the *Drosophila sechellia* (*Dsec*) genome possesses two copies of
Octα2R (Fig 1) and *Drosophila busckii* (*Dbus*) genome has two copies of *Octβ3R* (Fig 2).
Although the evolutionary forces underlying this variation in OA receptor copy number are
unknown, it is intriguing that both *Dsec* and *Dbus* have adapted to thrive on toxic hosts: *Dsec*
feeds on *Morinda* fruit (Legal et al., 1992; Louis, 1986), while *Dbus* consumes rotting vegetables

such as potatoes (Atkinson, 1977; Buda et al., 2009). This raises the possibility that OA signaling
may play a role in adaptation to environmental conditions through lineage-specific diversification
or amplification. Since dopamine and octopamine signaling interact functionally in flies (Burke et
al., 2012; Ma et al., 2016; Sabandal et al., 2020; Schwaerzel et al., 2003), and the dopaminergic
system is crucial for *Dsec*'s reproductive success and specialization on its toxic host (Lavista-
Llanos et al., 2014), it is plausible that octopaminergic signaling may also have contributed to
such species-specific adaptations. Gene copy number changes have been linked to ecological
adaptation in other species. For example, human populations with high-starch diets exhibit
increased amylase gene copy number relative to those with lower-starch diets (Perry et al.,
2007).Together, these findings raise the possibility that OA receptor copy number variation may
contribute to ecological specialization and environmental adaptation. Alternatively, this variation
may reflect reduced selective restraint on these genes in these species, leading to toleration of
duplication.

Octβ2R and Octβ3R have undergone positive selection in the *Drosophila melanogaster*
group

We then used PAML to investigate whether any of the OA receptors contain positively selected
sites (Yang, 2007), focusing our analysis on species within the *melanogaster* group (Fig 1, S4),
which corresponds to Clade 4 in the phylogeny of Suvorov et al. (2022). Because the functions of
these genes have been characterized in *D. melanogaster*, limiting our selection analyses to this
clade allows for evolutionary insights that are more directly relevant to the experimentally studied
functions. Model M0 was used to estimate the overall d_N/d_S (ω) ratio across the entire protein-
coding sequence of each gene; and models M7 vs M8 and M8a vs M8 comparisons (where M7
and M8a are neutral models and M8 is a model that allows for positively selected sites) were
used to identify genes evolving under positive selection and identify specific positively selected
sites (Swanson et al., 2003; Yang et al., 2000). The M8a null model explicitly includes neutrally
evolving sites, and the M8a/M8 comparison is therefore a more sensitive alternative to M7/M8
and less likely to yield false positives. We consider a gene to be under positive selection only if
both comparisons are significant. For genes under positive selection, we use the posterior
probabilities from the Bayes Empirical Bayes (BEB) test included in the output for codeml's M8 for
identifying sites under positive selection with a posterior probability threshold of 0.9.

Overall, OA receptors exhibit relatively low M0 ω estimates (d_N/d_S across the entire gene), which
may reflect their conserved roles in neuromodulatory and physiological processes. Octβ2R and
Octβ3R have the two highest M0 ω estimates among them, and interestingly, we found significant
evidence of positive selection acting on several amino acid sites within each of these two
receptors (Fig 3A). These genes have multiple isoforms identified and some of these isoforms
vary in coding DNA sequence (CDS) which could lead to isoforms having differing results from
site selection tests. We performed our analysis on all isoforms for our genes of interest that are
sufficiently annotated across species. The results for different isoforms were largely consistent
with one another, however, there was a difference in results for Octβ3R isoforms.

Under our criteria for the detection of positive selection, we identified sites under positive
selection in the Octβ3R-PG isoform but not the PF isoform (Fig 3A). The only difference in CDS
between these isoforms is the presence of an extra exon in the PF isoform (Fig 3B). Notably, the
positively selected sites (PSSs) identified in the PG isoform using the BEB posterior probabilities
from codeml's M8 output are all located in regions shared by both isoforms (Fig 3A, 3B). To
confirm the reliability of PSS detection, we performed an additional analysis focusing solely on
this shared region (PF_PG_5' shared), which yielded results consistent with those of the PG
isoform, showing significant differences in both the M7/M8 and M8a/M8 comparisons, with most
PSSs overlapping with those of PG (Fig 3A). Analysis of the PF-unique exon also detected
positive selection under both M7/M8 and M8a/M8 comparisons and exhibited the highest M0 ω
estimate relative to the shared regions (Fig 3A, 3B). Interestingly, when this exon is included in
the full PF isoform, the signal of positive selection is lost. This apparent loss of significance may

result from the unstructured nature of intracellular loop 3 encoded by the PF-unique exon, which
is likely evolving under reduced constraint. Such variation could alter the underlying ω distribution
and reduce the sensitivity of PSS detection in the full isoform. Moreover, the p-values for Octβ2R
and Octβ3R-PG remain significant after Benjamini-Hochberg false discovery rate (FDR)
correction (q-value; Fig 3A), supporting the robustness of our results.

The PSSs in Octβ2R and Octβ3R are predominantly clustered within unstructured loops (Fig 3C,
3D). In Octβ2R, the PSSs are primarily located in the N terminus, with one additional site in
extracellular loop 2 (ECL2) (Fig 3C). Notably, in the human β2-adrenergic receptor, a predicted
ortholog of Octβ2R, the N terminus and ECL2 have been reported to be important for ligand
binding and accessibility (Isin et al., 2012; Shahane et al., 2014). This suggests that Octβ2R may
have evolved to optimize interactions with its ligand. In some *melanogaster* group species,
Octβ2R may have evolved to bind OA with altered affinity. Alternatively, since OA is identical
across species, this raises the intriguing possibility that Octβ2R could have adapted to bind an
additional ligand. In Octβ3R, the PSSs are located in the N terminus and predominantly in
intracellular loop 3 (ICL3) (Fig 3D). In class A GPCRs, ICL3 plays a crucial role in signal
transduction and receptor activation, with its conformation and length influencing G protein
accessibility and selectivity (Sadler et al., 2023). The adaptive evolution of ICL3 in Octβ3R may
partially explain why Octβ3R is unable to fully compensate for Octβ2R's function in ovulation,
unlike Octβ1R (Lim et al., 2014). Additionally, Octβ3R may have evolved to interact with different
G proteins and/or acquired novel or specialized signaling functions in certain species.

To validate the PAML-based results, we performed complementary analyses, including
assessment of mutational bias and additional model-based tests (MEME). To exclude mutational
bias as a driver of elevated ω values, we examined codon usage bias, measured by the effective
number of codons (ENC), and GC content, including GC3 (GC content in the third codon position),
in the *melanogaster* group. Both genes show moderate codon usage bias (ENC: Octβ2R ~42-50;
Octβ3R-PG ~41-49) and relatively stable GC content across species (Octβ2R: GC ~0.55-0.58,
GC3 ~0.72-0.81; Octβ3R-PG: GC ~0.58-0.61, GC3 ~0.72-0.79) (Files S2), indicating that
mutational bias is unlikely to explain the observed patterns. Finally, we used MEME analysis
(Murrell et al., 2012) to investigate whether an alternate approach also detected PSSs in Octβ2R
and Octβ3R. Unlike PAML, which tests for pervasive selection, MEME detects episodic selection
using a different statistical framework. MEME identified more PSSs than PAML, and although
only a few sites overlapped between the two analyses (Fig 3A), the PSSs detected by either
method were localized in similar regions of the proteins. Detailed discussion of these analyses
and supporting data can be found in the supplement (File S3-S5).

In addition, Octα2R and certain isoforms of Oamb also showed sites under positive selection in
the M7/M8 comparison but lost significance in the M8a/M8 test (Fig 3A). Since the M8a null
model is an adaptation of M7 that includes an additional category of sites where ω = 1 (neutrally
evolving), this result suggests that these sites are more likely evolving under neutral rather than
adaptive selection. The possibility of redundancy with some of the other OA receptors could also
contribute to relaxed selective constraint on Octβ2R and Octβ3R, allowing them to evolve more
rapidly.

From a broader perspective, the observation that only some OA receptors have sites that have
evolved under positive selection in distinct functional domains highlights the heterogeneous
selective pressures shaping the evolution of this receptor family.

Octβ2R does not contain positively selected sites in the *Drosophila virilis-repleta* radiation nor in mosquitoes

We were curious whether positive selection of Octβ2R was restricted to the *melanogaster* clade,
or part of a broader pattern of its evolution. To explore this, we examined whether Octβ2R is also
under positive selection in other clades. We first analyzed species from the *virilis-repleta* radiation,

which last shared a common ancestor with the *melanogaster* group approximately 40-60 million
years ago within the *Drosophila* genus (Russo et al., 1995; Tamura et al., 2004). Using PAML, we
found that both the M7/M8 and more sensitive M8a/M8 comparisons showed no significant
difference, suggesting that Octβ2R is under purifying selection in this clade (Fig 4A). To further
broaden our analysis, we extended our tests to mosquitoes, a more distantly related group of
Dipterans. To minimize the risk of spurious signals of positive selection driven by synonymous
site saturation due to excessive evolutionary distance, we adopted a stepwise approach to the
PAML analyses. We began with the closely related *Cellia* species, where the M7/M8 and M8a/M8
comparisons revealed no evidence of positive selection (Fig 4A, 4B). We then expanded our
analysis to *Anophelinae* and subsequently to the broader *Culicidae* species (Fig 4B), despite the
divergence between *Culicinae* and *Anophelinae* exceeding 100 million years (Krzywinski et al.,
2006). Across all comparisons, we consistently found no evidence of positive selection (Fig 4A).

Positively selected sites were detected in Octβ2R within the *Drosophila melanogaster* group, but
not in the *Drosophila virilis‐repleta* radiation or mosquito species, suggesting a lineage-specific
change in evolutionary selective pressure. One possibility could be that there is lineage-specific
functionality in a process such as reproduction. Previous studies have shown that Octβ2R is
essential for ovulation (Li et al., 2015; Lim et al., 2014) and is expressed in both the oviduct
epithelium and OA neurons that project into the reproductive tract (Deshpande et al., 2022; Koon
et al., 2011). *D. virilis* and *D. melanogaster* vary greatly in the coterie of seminal fluid proteins
identified in their genomes and ejaculate (Garlovsky & Ahmed-Braimah, 2023), several of these
proteins have been observed to undergo rapid sequence divergence, sometimes driven by
positive selection (Haerty et al., 2007; Patlar et al., 2021; Swanson, Clark, et al., 2001; Wong &
Wolfner, 2012). Among other hypotheses, it is intriguing to wonder whether rapid evolution of the
female reproductive tract-expressed Octβ2R in the *melanogaster* clade might be driven by
interaction with a rapidly-evolving ejaculate protein in this clade, such as ovulin, which stimulates
ovulation through modulating OA signaling (Aguade, 1998; Aguade et al., 1992; Heifetz et al.,
2000; Rubinstein & Wolfner, 2013; Tsaur et al., 1998; Tsaur & Wu, 1997; Wong et al., 2006) and
is absent from species like *D. virilis* and *D. mojavensis*. However, experimental validation would
be needed to confirm any potential interaction between *D. melanogaster* Octβ2R and ovulin, or
any other protein.

Conclusion

OA receptors are key modulators of physiology and behavior in insects, yet their evolutionary
features remain largely unexplored. While their essential roles imply strong functional constraints,
their involvement in reproduction-related processes may also subject them to shifting evolutionary
selective pressures. In this study, we examined the molecular evolution of OA receptors across
Drosophila species and uncovered signatures of both conservation and diversification. Most OA
receptors were found as single-copy genes, consistent with stringent functional constraints acting
on them. However, we identified copy number changes in two species, *Drosophila sechellia*
(Octα2R) and *Drosophila busckii* (Octβ3R), which may reflect lineage-specific adaptations.
Among the six OA receptors, Octβ2R and Octβ3R within the *Drosophila melanogaster* clade
evolve under positive selection and contain positively selected sites in functionally distinct regions
of the GPCR domain. This heterogeneity suggests that different OA receptors are subject to
distinct selective pressures across the genus *Drosophila* and between paralogs, likely reflecting
divergent roles, molecular partners, or regulatory mechanisms. Notably, the detection of positive
selection in the ligand-binding involved regions of Octβ2R within the *melanogaster* group, but not
in other lineages, raises the possibility of lineage-specific selective pressure on ligand interaction.
Together, our findings illustrate the diverse evolutionary trajectories of a closely related receptor
family and motivate future functional studies aimed at understanding the molecular and ecological
roles of OA receptors in *Drosophila*.

Data Availability

359 PAML analysis data, MEME output files and jsons, and PSE files (PyMOL session files) showing
360 the structures of Octβ2R and Octβ3R with annotated PSSs and functional domains (extracellular,
361 transmembrane, and intracellular regions) are provided in the supplementary files.

Acknowledgments

364 We thank Drs. Willie Swanson, Geoffrey Findlay, Yasir Ahmed-Braimah, and anonymous
365 reviewers for advice on the selection analyses and/or comments on the manuscript, and
366 members of the Wolfner lab for helpful discussions and suggestions.

Funding

369 This work was supported by NIH grant R37-HD038921 to MFW. JAC was supported by NIH
370 postdoctoral fellowship F32-HD111231.

Conflicts of interest

The authors declare no conflicts of interest.

Literature Cited

Aguade, M. (1998). Different forces drive the evolution of the Acp26Aa and Acp26Ab accessory
gland genes in the Drosophila melanogaster species complex. *Genetics*, *150*(3), 1079-
1089. https://doi.org/10.1093/genetics/150.3.1079
Aguade, M., Miyashita, N., & Langley, C. H. (1992). Polymorphism and divergence in the Mst26A
male accessory gland gene region in Drosophila. *Genetics*, *132*(3), 755-770.
https://doi.org/10.1093/genetics/132.3.755
Atkinson, W. (1977). *Ecological studies of the breeding sites and reproductive strategies of
domestic species of Drosophila* University of Leeds].
Avila, F. W., Bloch Qazi, M. C., Rubinstein, C. D., & Wolfner, M. F. (2012). A requirement for the
neuromodulators octopamine and tyramine in Drosophila melanogaster female sperm
storage. *Proc Natl Acad Sci U S A*, *109*(12), 4562-4567.
https://doi.org/10.1073/pnas.1117689109
Balfanz, S., Strunker, T., Frings, S., & Baumann, A. (2005). A family of octopamine [corrected]
receptors that specifically induce cyclic AMP production or Ca2+ release in Drosophila
melanogaster. *J Neurochem*, *93*(2), 440-451. https://doi.org/10.1111/j.1471-
4159.2005.03034.x
Buda, V., Radziute, S., & Lutovinovas, E. (2009). Attractant for vinegar fly, Drosophila busckii,
and cluster fly, Pollenia rudis (Diptera: Drosophilidae et Calliphoridae). *Z Naturforsch C J
Biosci*, *64*(3-4), 267-270. https://doi.org/10.1515/znc-2009-3-419
Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., Gohl, D., Silies, M.,
Certel, S., & Waddell, S. (2012). Layered reward signalling through octopamine and

dopamine in Drosophila. *Nature*, *492*(7429), 433-437.
https://doi.org/10.1038/nature11614
Carlisle, J. A., Gurbuz, D. H., & Swanson, W. J. (2024). Recurrent Independent Pseudogenization
Events of the Sperm Fertilization Gene ZP3r in Apes and Monkeys. *Journal of molecular*
evolution, *92*(6), 695–702. https://doi.org/10.1007/s00239-024-10192-x
Carlisle, J. A., & Swanson, W. J. (2021). Molecular mechanisms and evolution of fertilization
proteins. Journal of Experimental Zoology Part B: Molecular and Developmental
Evolution, 336(8), 652-665.
Clark, N. L., Aagaard, J. E., & Swanson, W. J. (2006). Evolution of reproductive proteins from
animals and plants. *Reproduction*, *131*(1), 11-22. https://doi.org/10.1530/rep.1.00357
Crocker, A., Shahidullah, M., Levitan, I. B., & Sehgal, A. (2010). Identification of a neural circuit
that underlies the effects of octopamine on sleep:wake behavior. *Neuron*, *65*(5), 670-681.
https://doi.org/10.1016/j.neuron.2010.01.032
Dapper, A. L., & Wade, M. J. (2020). Relaxed Selection and the Rapid Evolution of Reproductive
Genes. *Trends Genet*, *36*(9), 640-649. https://doi.org/10.1016/j.tig.2020.06.014
Deady, L. D., & Sun, J. (2015). A Follicle Rupture Assay Reveals an Essential Role for Follicular
Adrenergic Signaling in Drosophila Ovulation. *PLoS Genet*, *11*(10), e1005604.
https://doi.org/10.1371/journal.pgen.1005604
Deshpande, S. A., Rohrbach, E. W., Asuncion, J. D., Harrigan, J., Eamani, A., Schlingmann, E.
H., Suto, D. J., Lee, P. T., Schweizer, F. E., Bellen, H. J., & Krantz, D. E. (2022).
Regulation of Drosophila oviduct muscle contractility by octopamine. *iScience*, *25*(8),
104697. https://doi.org/10.1016/j.isci.2022.104697
Evans, P. D., & Maqueira, B. (2005). Insect octopamine receptors: a new classification scheme
based on studies of cloned Drosophila G-protein coupled receptors. *Invert Neurosci*, *5*(3-
4), 111-118. https://doi.org/10.1007/s10158-005-0001-z
Farooqui, T. (2012). Review of octopamine in insect nervous systems. *Open access insect*
physiology, 1-17.
Galindo, B. E., Vacquier, V. D., & Swanson, W. J. (2003). Positive selection in the egg receptor
for abalone sperm lysin. *Proc Natl Acad Sci U S A*, *100*(8), 4639-4643.
https://doi.org/10.1073/pnas.0830022100
Garlovsky, M. D., & Ahmed-Braimah, Y. H. (2023). Evolutionary quantitative proteomics of
reproductive protein divergence in Drosophila. *Mol Cell Proteomics*, 22(8), 100610.
https://doi.org/10.1016/j.mcpro.2023.100610
Haerty, W., Jagadeeshan, S., Kulathinal, R. J., Wong, A., Ravi Ram, K., Sirot, L. K., Levesque,
L., Artieri, C. G., Wolfner, M. F., Civetta, A., & Singh, R. S. (2007). Evolution in the fast
lane: rapidly evolving sex-related genes in Drosophila. *Genetics*, *177*(3), 1321-1335.
https://doi.org/10.1534/genetics.107.078865
Han, K. A., Millar, N. S., & Davis, R. L. (1998). A novel octopamine receptor with preferential
expression in Drosophila mushroom bodies. *J Neurosci*, *18*(10), 3650-3658.
https://doi.org/10.1523/JNEUROSCI.18-10-03650.1998
Heifetz, Y., Lung, O., Frongillo, E. A., Jr., & Wolfner, M. F. (2000). The Drosophila seminal fluid
protein Acp26Aa stimulates release of oocytes by the ovary. *Curr Biol*, *10*(2), 99-102.
https://doi.org/10.1016/s0960-9822(00)00288-8
Hopkins, B. R., Angus-Henry, A., Kim, B. Y., Carlisle, J. A., Thompson, A., & Kopp, A. (2024).
Decoupled evolution of the Sex Peptide gene family and Sex Peptide Receptor in
Drosophilidae. *Proc Natl Acad Sci U S A*, *121*(3), e2312380120.
https://doi.org/10.1073/pnas.2312380120
Huetteroth, W., Perisse, E., Lin, S., Klappenbach, M., Burke, C., & Waddell, S. (2015). Sweet
taste and nutrient value subdivide rewarding dopaminergic neurons in Drosophila. *Curr*
Biol, *25*(6), 751-758. https://doi.org/10.1016/j.cub.2015.01.036
Isin, B., Estiu, G., Wiest, O., & Oltvai, Z. N. (2012). Identifying ligand binding conformations of the
beta2-adrenergic receptor by using its agonists as computational probes. *PLoS One*,
7(12), e50186. https://doi.org/10.1371/journal.pone.0050186

Kim, Y. C., Lee, H. G., Lim, J., & Han, K. A. (2013). Appetitive learning requires the alpha1-like
octopamine receptor OAMB in the Drosophila mushroom body neurons. *J Neurosci*,
33(4), 1672-1677. https://doi.org/10.1523/JNEUROSCI.3042-12.2013
Koon, A. C., Ashley, J., Barria, R., DasGupta, S., Brain, R., Waddell, S., Alkema, M. J., & Budnik,
V. (2011). Autoregulatory and paracrine control of synaptic and behavioral plasticity by
octopaminergic signaling. *Nat Neurosci*, *14*(2), 190-199. https://doi.org/10.1038/nn.2716
Koon, A. C., & Budnik, V. (2012). Inhibitory control of synaptic and behavioral plasticity by
octopaminergic signaling. *J Neurosci*, *32*(18), 6312-6322.
https://doi.org/10.1523/JNEUROSCI.6517-11.2012
Krzywinski, J., Grushko, O. G., & Besansky, N. J. (2006). Analysis of the complete mitochondrial
DNA from Anopheles funestus: an improved dipteran mitochondrial genome annotation
and a temporal dimension of mosquito evolution. *Mol Phylogenet Evol*, *39*(2), 417-423.
https://doi.org/10.1016/j.ympev.2006.01.006
Lavista-Llanos, S., Svatos, A., Kai, M., Riemensperger, T., Birman, S., Stensmyr, M. C., &
Hansson, B. S. (2014). Dopamine drives Drosophila sechellia adaptation to its toxic host.
Elife, *3*. https://doi.org/10.7554/eLife.03785
Lee, H. G., Rohila, S., & Han, K. A. (2009). The octopamine receptor OAMB mediates ovulation
via Ca2+/calmodulin-dependent protein kinase II in the Drosophila oviduct epithelium.
PLoS One, *4*(3), e4716. https://doi.org/10.1371/journal.pone.0004716
Legal, L., David, J. R., & Jallon, J. M. (1992). Toxicity and attraction effects produced by Morinda
citrifolia fruits on the Drosophila melanogaster complex of species. *Chemoecology*, *3*,
125-129.
Li, Y., Fink, C., El-Kholy, S., & Roeder, T. (2015). The octopamine receptor octss2R is essential
for ovulation and fertilization in the fruit fly Drosophila melanogaster. *Arch Insect Biochem
Physiol*, *88*(3), 168-178. https://doi.org/10.1002/arch.21211
Lim, J., Sabandal, P. R., Fernandez, A., Sabandal, J. M., Lee, H. G., Evans, P., & Han, K. A.
(2014). The octopamine receptor Octbeta2R regulates ovulation in Drosophila
melanogaster. *PLoS One*, *9*(8), e104441. https://doi.org/10.1371/journal.pone.0104441
Louis, J. (1986). Ecological specialization in the Drosophila melanogaster species subgroup: a
case study of D. sechellia. *Ecol. Genet.*, *7*, 215-229.
Luo, J., Lushchak, O. V., Goergen, P., Williams, M. J., & Nassel, D. R. (2014). Drosophila insulin-
producing cells are differentially modulated by serotonin and octopamine receptors and
affect social behavior. *PLoS One*, *9*(6), e99732.
https://doi.org/10.1371/journal.pone.0099732
Ma, Z., Stork, T., Bergles, D. E., & Freeman, M. R. (2016). Neuromodulators signal through
astrocytes to alter neural circuit activity and behaviour. *Nature*, *539*(7629), 428-432.
https://doi.org/10.1038/nature20145
Madeira, F., Madhusoodanan, N., Lee, J., Eusebi, A., Niewielska, A., Tivey, A. R. N., Lopez, R., &
Butcher, S. (2024). The EMBL-EBI Job Dispatcher sequence analysis tools framework in
2024. *Nucleic Acids Res*, *52*(W1), W521-W525. https://doi.org/10.1093/nar/gkae241
McGeary, M. K., & Findlay, G. D. (2020). Molecular evolution of the sex peptide network in
Drosophila. *J Evol Biol*, *33*(5), 629-641. https://doi.org/10.1111/jeb.13597
Moyle, L. C., Wu, M., & Gibson, M. J. S. (2021). Reproductive Proteins Evolve Faster Than Non-
reproductive Proteins Among Solanum Species. *Front Plant Sci*, *12*, 635990.
https://doi.org/10.3389/fpls.2021.635990
Murrell, B., Wertheim, J. O., Moola, S., Weighill, T., Scheffler, K., & Kosakovsky Pond, S. L.
(2012). Detecting individual sites subject to episodic diversifying selection. *PLoS
genetics*, *8*(7), e1002764.
Nakagawa, H., Maehara, S., Kume, K., Ohta, H., & Tomita, J. (2022). Biological functions of
alpha2-adrenergic-like octopamine receptor in Drosophila melanogaster. *Genes Brain
Behav*, *21*(6), e12807. https://doi.org/10.1111/gbb.12807
Neafsey, D. E., Waterhouse, R. M., Abai, M. R., Aganezov, S. S., Alekseyev, M. A., Allen, J. E.,
Amon, J., Arcà, B., Arensburger, P., & Artemov, G. (2015). Highly evolvable malaria
vectors: the genomes of 16 Anopheles mosquitoes. *Science*, *347*(6217), 1258522.

Ohhara, Y., Shimada-Niwa, Y., Niwa, R., Kayashima, Y., Hayashi, Y., Akagi, K., Ueda, H., Yamakawa-Kobayashi, K., & Kobayashi, S. (2015). Autocrine regulation of ecdysone synthesis by beta3-octopamine receptor in the prothoracic gland is essential for Drosophila metamorphosis. *Proc Natl Acad Sci U S A*, *112*(5), 1452-1457. https://doi.org/10.1073/pnas.1414966112

Patlar, B., Jayaswal, V., Ranz, J. M., & Civetta, A. (2021). Nonadaptive molecular evolution of seminal fluid proteins in Drosophila. *Evolution*, *75*(8), 2102-2113. https://doi.org/10.1111/evo.14297

Perry, G. H., Dominy, N. J., Claw, K. G., Lee, A. S., Fiegler, H., Redon, R., Werner, J., Villanea, F. A., Mountain, J. L., Misra, R., Carter, N. P., Lee, C., & Stone, A. C. (2007). Diet and the evolution of human amylase gene copy number variation. *Nat Genet*, *39*(10), 1256-1260. https://doi.org/10.1038/ng2123

Qi, Y. X., Xu, G., Gu, G. X., Mao, F., Ye, G. Y., Liu, W., & Huang, J. (2017). A new Drosophila octopamine receptor responds to serotonin. *Insect Biochem Mol Biol*, *90*, 61-70. https://doi.org/10.1016/j.ibmb.2017.09.010

Robb, S., Cheek, T. R., Hannan, F. L., Hall, L. M., Midgley, J. M., & Evans, P. D. (1994). Agonist-specific coupling of a cloned Drosophila octopamine/tyramine receptor to multiple second messenger systems. *EMBO J*, *13*(6), 1325-1330. https://doi.org/10.1002/j.1460-2075.1994.tb06385.x

Roeder, T. (2005). Tyramine and octopamine: ruling behavior and metabolism. *Annu Rev Entomol*, *50*, 447-477. https://doi.org/10.1146/annurev.ento.50.071803.130404

Roeder, T. (2020). The control of metabolic traits by octopamine and tyramine in invertebrates. *J Exp Biol*, *223*(Pt 7). https://doi.org/10.1242/jeb.194282

Rubinstein, C. D., & Wolfner, M. F. (2013). Drosophila seminal protein ovulin mediates ovulation through female octopamine neuronal signaling. *Proc Natl Acad Sci U S A*, *110*(43), 17420-17425. https://doi.org/10.1073/pnas.1220018110

Russo, C. A., Takezaki, N., & Nei, M. (1995). Molecular phylogeny and divergence times of drosophilid species. *Mol Biol Evol*, *12*(3), 391-404. https://doi.org/10.1093/oxfordjournals.molbev.a040214

Sabandal, J. M., Sabandal, P. R., Kim, Y. C., & Han, K. A. (2020). Concerted Actions of Octopamine and Dopamine Receptors Drive Olfactory Learning. *J Neurosci*, *40*(21), 4240-4250. https://doi.org/10.1523/JNEUROSCI.1756-19.2020

Sadler, F., Ma, N., Ritt, M., Sharma, Y., Vaidehi, N., & Sivaramakrishnan, S. (2023). Autoregulation of GPCR signalling through the third intracellular loop. *Nature*, *615*(7953), 734-741. https://doi.org/10.1038/s41586-023-05789-z

Schwaerzel, M., Monastirioti, M., Scholz, H., Friggi-Grelin, F., Birman, S., & Heisenberg, M. (2003). Dopamine and octopamine differentiate between aversive and appetitive olfactory memories in Drosophila. *J Neurosci*, *23*(33), 10495-10502. https://doi.org/10.1523/JNEUROSCI.23-33-10495.2003

Shahane, G., Parsania, C., Sengupta, D., & Joshi, M. (2014). Molecular insights into the dynamics of pharmacogenetically important N-terminal variants of the human beta2-adrenergic receptor. *PLoS Comput Biol*, *10*(12), e1004006. https://doi.org/10.1371/journal.pcbi.1004006

Sujkowski, A., Gretzinger, A., Soave, N., Todi, S. V., & Wessells, R. (2020). Alpha- and beta-adrenergic octopamine receptors in muscle and heart are required for Drosophila exercise adaptations. *PLoS Genet*, *16*(6), e1008778. https://doi.org/10.1371/journal.pgen.1008778

Sun, M., Ma, M., Deng, B., Li, N., Peng, Q., & Pan, Y. (2023). A neural pathway underlying hunger modulation of sexual receptivity in Drosophila females. *Cell Rep*, *42*(10), 113243. https://doi.org/10.1016/j.celrep.2023.113243

Suvorov, A., Kim, B. Y., Wang, J., Armstrong, E. E., Peede, D., D'Agostino, E. R. R., Price, D. K., Waddell, P., Lang, M., Courtier-Orgogozo, V., David, J. R., Petrov, D., Matute, D. R., Schrider, D. R., & Comeault, A. A. (2022). Widespread introgression across a phylogeny

of 155 Drosophila genomes. *Curr Biol*, *32*(1), 111-123 e115.
https://doi.org/10.1016/j.cub.2021.10.052

Suyama, M., Torrents, D., & Bork, P. (2006). PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. *Nucleic Acids Res*, *34*(Web Server
issue), W609-612. https://doi.org/10.1093/nar/gkl315

Swanson, W. J., Clark, A. G., Waldrip-Dail, H. M., Wolfner, M. F., & Aquadro, C. F. (2001).
Evolutionary EST analysis identifies rapidly evolving male reproductive proteins in
Drosophila. *Proc Natl Acad Sci U S A*, *98*(13), 7375-7379.
https://doi.org/10.1073/pnas.131568198

Swanson, W. J., Nielsen, R., & Yang, Q. (2003). Pervasive adaptive evolution in mammalian
fertilization proteins. *Mol Biol Evol*, *20*(1), 18-20.
https://doi.org/10.1093/oxfordjournals.molbev.a004233

Swanson, W. J., & Vacquier, V. D. (2002). The rapid evolution of reproductive proteins. *Nat Rev*
Genet, *3*(2), 137-144. https://doi.org/10.1038/nrg733

Swanson, W. J., Yang, Z., Wolfner, M. F., & Aquadro, C. F. (2001). Positive Darwinian selection
drives the evolution of several female reproductive proteins in mammals. *Proc Natl Acad*
Sci U S A, *98*(5), 2509-2514. https://doi.org/10.1073/pnas.051605998

Tamura, K., Subramanian, S., & Kumar, S. (2004). Temporal patterns of fruit fly (Drosophila)
evolution revealed by mutation clocks. *Mol Biol Evol*, *21*(1), 36-44.
https://doi.org/10.1093/molbev/msg236

Tegenfeldt, F., Kuznetsov, D., Manni, M., Berkeley, M., Zdobnov, E. M., & Kriventseva, E. V.
(2025). OrthoDB and BUSCO update: annotation of orthologs with wider sampling of
genomes. *Nucleic acids research*, *53*(D1), D516–D522.
https://doi.org/10.1093/nar/gkae987

Thiébaut, A., Altenhoff, A. M., Campli, G., Glover, N., Dessimoz, C., & Waterhouse, R. M. (2024).
DrosOMA: the *Drosophila* Orthologous Matrix browser. *F1000Research*, *12*, 936.
https://doi.org/10.12688/f1000research.135250.2

Tsaur, S. C., Ting, C. T., & Wu, C. I. (1998). Positive selection driving the evolution of a gene of
male reproduction, Acp26Aa, of Drosophila: II. Divergence versus polymorphism. *Mol*
Biol Evol, *15*(8), 1040-1046. https://doi.org/10.1093/oxfordjournals.molbev.a026002

Tsaur, S. C., & Wu, C. I. (1997). Positive selection and the molecular evolution of a gene of male
reproduction, Acp26Aa of Drosophila. *Mol Biol Evol*, *14*(5), 544-549.
https://doi.org/10.1093/oxfordjournals.molbev.a025791

Weaver, S., Shank, S. D., Spielman, S. J., Li, M., Muse, S. V., & Kosakovsky Pond, S. L. (2018).
Datamonkey 2.0: A Modern Web Application for Characterizing Selective and Other
Evolutionary Processes. *Molecular biology and evolution*, *35*(3), 773–777.
https://doi.org/10.1093/molbev/msx335

White, M. A., Chen, D. S., & Wolfner, M. F. (2021). She's got nerve: roles of octopamine in insect
female reproduction. *J Neurogenet*, *35*(3), 132-153.
https://doi.org/10.1080/01677063.2020.1868457

Wilburn, D. B., & Swanson, W. J. (2016). From molecules to mating: Rapid evolution and
biochemical studies of reproductive proteins. *J Proteomics*, *135*, 12-25.
https://doi.org/10.1016/j.jprot.2015.06.007

Wong, A., Albright, S. N., & Wolfner, M. F. (2006). Evidence for structural constraint on ovulin, a
rapidly evolving Drosophila melanogaster seminal protein. *Proc Natl Acad Sci U S A*,
103(49), 18644-18649. https://doi.org/10.1073/pnas.0601849103

Wong, A., & Wolfner, M. F. (2012). 144Evolution of Drosophila seminal proteins and their
networks. In R. S. Singh, J. Xu, & R. J. Kulathinal (Eds.), *Rapidly Evolving Genes and*
Genetic Systems (pp. 0). Oxford University Press.
https://doi.org/10.1093/acprof:oso/9780199642274.003.0015

Wu, C. L., Shih, M. F., Lee, P. T., & Chiang, A. S. (2013). An octopamine-mushroom body circuit
modulates the formation of anesthesia-resistant memory in Drosophila. *Curr Biol*, *23*(23),
2346-2354. https://doi.org/10.1016/j.cub.2013.09.056

Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*, *24*(8),
612   1586-1591. https://doi.org/10.1093/molbev/msm088
613   Yang, Z., Nielsen, R., Goldman, N., & Pedersen, A. M. (2000). Codon-substitution models for
614   heterogeneous selection pressure at amino acid sites. *Genetics*, *155*(1), 431-449.
615   https://doi.org/10.1093/genetics/155.1.431
616   Zhang, T., Branch, A., & Shen, P. (2013). Octopamine-mediated circuit mechanism underlying
617   controlled appetite for palatable food in Drosophila. *Proc Natl Acad Sci U S A*, *110*(38),
618   15431-15436. https://doi.org/10.1073/pnas.1308816110
619   Zhang, Y., Yu, Y., Qian, M., Gui, W., Shah, A. Z., Xu, G., & Yang, G. (2023). Characterization and
620   functional analysis of an alpha-adrenergic-like octopamine receptor in the small brown
621   planthopper Laodelphax striatellus. *Pestic Biochem Physiol*, *194*, 105509.
622   https://doi.org/10.1016/j.pestbp.2023.105509
623   Zhao, Z., Zhao, X., He, T., Wu, X., Lv, P., Zhu, A. J., & Du, J. (2021). Epigenetic regulator
624   Stuxnet modulates octopamine effect on sleep through a Stuxnet-Polycomb-Octbeta2R
625   cascade. *EMBO Rep*, *22*(2), e47910. https://doi.org/10.15252/embr.201947910

Figures

Figure 1. Syntenic region of *Octα2R* across *Drosophila* species. The phylogeny is based on
Suvorov et al. (2022) and Hopkins et al. (2024). Surrounding gene names correspond to
orthologs in *D. melanogaster.* The species within the gray box represent the *melanogaster* group
as defined in the paper.

Figure 2. Syntenic region of *Octβ2R* and *Octβ3R* across *Drosophila* species. The phylogeny is
based on Suvorov et al. (2022) and Hopkins et al. (2024). Surrounding gene names correspond
to orthologs in *D. melanogaster.*

Figure 3. Octβ2R and Octβ3R are under recurrent positive selection in the *melanogaster* group.
A. PAML test results for all six OA receptors across the *melanogaster* group. Significant p-values
(raw p-values and Benjamini-Hochberg FDR adjusted p-values (q-values, rounded to four decimal
places)) are shown in bold, and coding sequences with significant evidence of positive selection
are highlighted in green. The italicized sites in the last column (mapped to *D. melanogaster* amino
acid sequence) are located in the intracellular domain, while the others are extracellular.
Asterisks in the last column indicate sites also detected as positively selected by MEME (For
Octβ3R, gray asterisks indicate sites also detected in the PF isoform MEME analysis, black
asterisks indicate sites detected in the PG isoform analysis, and sites marked with both gray and
black asterisks were found in both MEME analyses) (File S4, S5). BEB: Bayes empirical Bayes.
B. Schematic of the PF and PG isoforms of Octβ3R. Boxes represent exons, lines indicate
introns, and asterisks mark exons containing positively selected sites. The "PF_PG_5' shared"
and "PF_unique" regions tested in panel A, as well as the "PF_PG_3' shared" exons, are outlined
with dotted borders. M0 ω estimations for each region are indicated. C. Positively selected amino
acid sites in Octβ2R (BEB posterior probability > 0.90) are shown in magenta and mapped onto
the *D. melanogaster* protein sequence and AlphaFold-predicted structure. Nine sites are in the N
terminus (orange) and one site is in the extracellular loop 2 (deep olive); both regions are
extracellular. D. Positively selected amino acid sites in the PG isoform of Octβ3R (BEB posterior
probability > 0.90) are shown in magenta and mapped onto the *D. melanogaster* protein
sequence and AlphaFold-predicted structure. Three sites are in the N terminus (orange;
extracellular) and eight sites are in the intracellular loop 3 (cyan; intracellular).

Figure 4. Octβ2R is highly conserved in the *Drosophila virilis-repleta* radiation and in mosquitoes.
A. PAML test results for Octβ2R across the *virilis-repleta* radiation and mosquito (*Cellia*,
Anophelinae, and *Culicidae*) species. No significant differences were detected in any M7 vs. M8
or M8a vs. M8 comparisons, indicating that Octβ2R is under purifying selection in these groups.
B. Phylogeny of the *virilis-repleta* radiation and mosquito species used in the PAML analyses,
based on the species tree reported by Suvorov et al. (2022) and Neafsey et al. (2015). The
melanogaster group is also included to illustrate relationships among the three clades in which
Octβ2R was analyzed.

( ) Genes on opposite strand relative to *Dmel*

A

Gene	Isoforms	M0 ω estimate	M7 vs. M8			M8a vs. M8			% sites in ω > 1	Sites with BEB Pr > 0.90
			2*|ΔlnL|	p-value (raw)	q-value (FDR)	2*|ΔlnL|	p-value (raw)	q-value (FDR)		
Octβ2R	PA/PC/PD/PE/PG	0.13231	16.366	**.000279**	**.0031**	10.884	**.00097**	**.0100**	3.936% (ω=2.28245)	35D, 36G, 48I, 49V, 51N, 53I, 98I, 130A, 142N*, 308T*
Oamb	PB/PE	0.03577	4.210	.121846	.1489	0.060	.806496	1.0000	-	-
	PC/PD/PF/PG	0.03167	9.450	**.008871**	**.0163**	0.012	.912771	1.0000	-	-
Octβ1R	PA/PE	0.04122	2.169	.338071	.3568	0	1	1.0000	-	-
Octβ3R	PF	0.07295	8.228	**.016342**	**.0225**	0	1	1.0000	-	-
	PG	0.07347	14.338	**.00077**	**.0042**	8.758	**.003082**	**.0113**	1.597% (ω=2.29822)	22A, 23G , 28Q*, *476I, 486S, 611L, 737I* , 833G, 841S, 893Q , 956G*
	PF_PG_ 5' shared	0.07465	12.697	**.001749**	**.0064**	6.598	**.010209**	**.0281**	1.995% (ω=1.96236)	22A, 23G , *486S, 611L, 737I* , 833G, 839S* , 841S,891N*, 893Q*
	PF_unique	0.23771	11.935	**.002561**	**.0067**	9.720	**.001823**	**.0100**	5.696% (ω=3.79922)	*1101Q , 1112S*
Octα2R	PA/PB	0.05676	11.602	**.003025**	**.0067**	0.812	.36753	.8086	-	-
	PC	0.05742	8.609	**.013508**	**.0212**	0.006	.938258	1.0000	-	-
Oct-TyrR	PA/PB	0.04271	2.061	.356829	.3568	0	1	1.0000	-	-

B

PF_unique (M0 ω= 0.23771)

Octβ3R

PF

PG

PF_PG_3' shared (M0 ω= 0.00010)

PF_PG_5' shared (M0 ω= 0.07465)

C

Octβ2R

D

Octβ3R-PG

A

Gene	Species	M0 ω estimate	M7 vs. M8		M8a vs. M8	
			2*|ΔlnL|	p-value	2*|ΔlnL|	p-value
Octβ2R	*Drosophila virilis-repleta* species	0.11871	2×10^{-5}	~1	0	1
	Cellia species	0.06588	1×10^{-4}	~1	0	1
	Anophelinae species	0.07351	0.032	.984127	0.032	.858028
	Culicidae species	0.07054	0.017	.991536	0	1

B