NRT1.1s and NRT2.1 affect rhizosphere and apoplastic pH during plant nitrate uptake

Qiying Xiao[a], Junjie Wang[a], Yi Chen[b], Fuyu Li[a], Wuge Jiwu[c], Anthony J. Miller[b], Jeremy D. Murray*[, a, b]

[a]Department of Botany, CAS-JIC Centre of Excellence for Plant and Microbial Science (CEPAMS), Centre for Excellence in Molecular Plant Sciences (CEMPS), Shanghai Institute of Plant Physiology and Ecology (SIPPE), Chinese Academy of Sciences, 300 Feng Lin Road, Shanghai 200032, China; [b]Department Biochemistry & Metabolism, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK; [c]College of Life Sciences, Shanghai Normal University, Shanghai, China

*Corresponding author: Jeremy D. Murray

Email: jeremy.murray@jic.ac.uk

Author Contributions: QX, JW, YC and WJ performed experimental work and data analysis. JDM and AM supervised and co-directed the research. QX, AJM, FL and JDM conceived the experiments and wrote the article.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences, Plant Biology

Keywords: nitrate transport, apoplast, cell wall, pH,

This file includes:

Main Text

Figures 1 to 2

1 Abstract

Nitrate is the primary source of nitrogen for plants in most environments. Nitrate uptake from the
rhizosphere into root cells is mediated by Nitrate Transporter 2 (NRT2) and NRT1.1 proteins via
proton symport, with the apoplast serving as an intermediate compartment. Using a simple root
immersion system, we investigated the effects of NRT transporters on external media pH (pH_{med})
and apoplastic pH (pH_{apo}) in *M. truncatula* nitrate transporter mutants. Monitoring of pH_{med} showed
that loss of NRT2.1 largely prevented nitrate induced alkalization of the media, while loss of
NRT1.1A/B reduced media alkalization only during the initial stages of nitrate uptake, suggesting
they have fundamentally different physiological roles. Use of the pH sensitive dye HPTS showed
that the pH_{apo} of root cells initially decreased during nitrate uptake, and this required NRT1.1A/B.
Unexpectedly, the observed changes in pH_{med} during nitrate uptake in WT and the mutants were
not reflected in pH_{apo}. To explain these discrepancies, we propose a model in which the depletion
of protons in the media occurring during nitrate symport is offset by the buffering of protons in the
cell wall and the activity of the H^+-ATPase, allowing a lower pH_{apo} to be maintained. Such a
mechanism would enable efficient nitrate uptake across a range of external pH values.

17 Introduction

Nitrate uptake in plants depends on proton symport, which requires a supply of protons from the
apoplast. This requirement is reflected in the apoplastic pH (pH_{apo}), which typically ranges from 4.5-
5.5 (1). Nitrate symport results in alkalization of the rhizosphere which occurs through the
consumption of protons by nitrate proton symporters with a stoichiometry of at least 2 H^+ per nitrate
absorbed (2). In *Arabidopsis thaliana*, this occurs through the activities of the high affinity nitrate
uptake transporter AtNRT2.1 and the low or dual affinity transporter AtNRT1.1, also called Nitrate
Transporter 1/Peptide Transporter 6.3 (3,4). AtNRT1.1 also plays an important role in nitrate
sensing (5). Loss of AtNRT1.1 greatly diminishes nitrate-induced rhizosphere alkalization (6), but
NRT2.1's role in this phenomenon has not been studied. Despite it playing a critical role in nitrate
uptake, the dynamics of pH_{apo} in response to nitrate remains uninvestigated. Here we addressed
these questions in *Medicago truncatula* (Gaertn) using NRT2.1 and NRT1.1A/B mutants, the
respective orthologs of Arabidopsis NRT2.1 and NRT1.1 (7).

32 Results

34 To observe the effect of nitrate uptake on medium pH (pH_{med}), *M. truncatula* seedlings were grown
35 on agar plates containing nitrate-free Fåhraeus nutrient solution (FP) for one week. The seedlings

were then transplanted into non-aerated liquid FP medium supplemented with different concentrations of KNO_3 (initial pH_{med} of 6.0) with their roots fully submerged. In the absence of nitrate, the seedlings acidified the FP medium to at ~4.0-4.2 by 24h (Fig. 1*A*). Treatment with high concentrations of nitrate ($\geq$5mM) significantly lowered pH_{med} at 3 and 6h compared to the FP control by 0.2-0.4 pH units (Dunnett's test, p-value <0.05; inset Fig. 1*A*), but this trend reversed at 24h with the nitrate containing samples becoming alkalized, with a pH_{med} ~1 pH unit higher than the nitrate-free control, with even larger differences (~1.5-2.2 pH units) occurring at later time points. In comparison, treatment with low nitrate (0.5 mM KNO_3) had only a slight effect on pH_{med}. These results are consistent with media alkalization by the seedlings being due to nitrate/H^+ symport mediated depletion of protons.

To determine if the medium acidification in our system was caused by flood-induced hypoxia, we repeated the experiment with gentle aeration. This increased the O_2 concentration of the media from 7.2 ($\pm$0.04 SEM) to 9.6 ($\pm$0.17) ppm and largely attenuated the acidification (Fig. 1*B*). We then investigated the potential involvement of the PM H^+-ATPase using the specific inhibitor protonstatin-1 (PS-1). Treatment with PS-1 essentially blocked root proton extrusion, with 91 and 95% inhibition by 25 and 50mM PS-1, respectively (Fig. 1*C*). These results suggest that the observed media acidification is mediated by the PM H^+-ATPase and is induced by short-term hypoxia resulting from the immersion.

The root immersion system provides a means of evaluating the H^+ consumption by NRTs during nitrate uptake. First, we tested whether the acidification of the media was affected by the absence of NRTs. Five mutants were tested, *nrt2.1*, *nrt1.1a*, *nrt1.1b*, *nrt1.1c*, and the *nrt1.1a/b* double mutant. Every mutant acidified pH_{med} to ~3.8-4.0 within 24h, similar to WT (Fig. 1*D*). We then tested the effect of these mutations on pH_{med} in the presence of various concentrations of KNO_3. We found that the *nrt1.1a/b* double mutant and *nrt2.1* mutant affected pH_{med} in different ways (Fig. 1*E-G*). The *nrt1.1a/b* mutant showed a small but significant (~0.1 pH unit; Dunnett's test, p-value <0.05) decrease in pH_{med} at 3 and 6h after nitrate addition at all nitrate concentrations. This effect grew stronger at 24h (~0.2-0.4 pH units, depending on the nitrate concentration) but was greatly diminished or absent at the later time points. This sharply contrasted with *nrt2.1*, which showed no phenotype at 3 and 6h, but had a significantly diminished ability to alkalize the media at 24 and 48 under all nitrate concentrations, which grew more pronounced at 80h (~1.4-1.7 pH units) under 5 and 10 mM KNO_3. Under 0.5mM the pH_{med} phenotype of *nrt2.* at 24h and 48h was weaker, and no difference was seen at 80h.

We then investigated whether changes occur in pH_{apo} occur during nitrate uptake. We used HPTS, a widely used fluorescent membrane-impermeant pH sensitive dye, to stain immersed roots and made observations on root tip cells using confocal microscopy (see *S1 Appendix*). After transfer to 5 mM nitrate the pH_{apo} of WT seedlings significantly decreased, indicating a pH drop, at

4h (Fig. 2*A*). The *nrt1.1a/b* mutant failed to respond, indicating it has a reduced ability to acidify the

apoplast. Observations at the 24- and 48-hour marks indicate that pH_{apo} of WT recovered, while

that of *nrt1.1a/b* was higher at 48h.

Discussion

The seedling immersion approach provides a convenient means to observe the effect of root nitrate

uptake on external pH. Our results indicate that the initial drop in pH_{med} after seedling transfer was

caused by hypoxia. A similar drop in pH_{med} was reported in *A. thaliana* after root immersion, which

was shown to require H^+-ATPases 1 and 2 (8). Furthermore, it was shown that hypoxia induces

increased proton efflux in barley root tips, which was attributed to PM H^+-ATPase activity (9),

suggesting this is a general hypoxia-related phenomenon. Using this approach, we found that

nitrate initially induced a small drop in pH_{med} and pH_{apo}. It was shown in maize that following an

initial depolarization after exposure to nitrate, the PM becomes hyperpolarized through

overcompensation of the H^+-ATPase (10). Such hyperpolarisation could explain the observed drop

in pH_{med} and pH_{apo} at early time points. Consistent with this, reports in maize showed an increase

in root PM H^+-ATPase activity after nitrate supply (11,12).

The initial nitrate-induced drop in pH_{apo} was compromised in the *nrt1.1a/b* mutant. This

requires an explanation, since decreased nitrate/H^+ symport would be expected to lower, not raise,

the pH of this compartment. We suggest that this may occur through reduced activation of the PM

H^+-ATPase due to lower nitrate uptake in the mutants. While the trigger for nitrate induced

hyperpolarization is unknown, it is possible that the increased influx of protons into the cytoplasm

by nitrate/H^+ symport could trigger higher H^+-ATPase activity to maintain a stable cytoplasmic pH

(13). In this case, a decrease in nitrate/H^+ symport activity would lessen the stimulation of the PM

H^+-ATPase by alleviating symport-mediated cytosolic acidification, resulting in a higher pH_{apo} in the

mutants. Another, non-exclusive possibility, is the inhibitory effect of nitrate on vacuolar (V)-

ATPases (14). Such downregulation of the V-ATPase would serve a complementary role,

preventing competition with the PM-ATPase for cytosolic protons during nitrate uptake.

The strong alkalization observed in the medium at later time points did not occur in the

apoplast. Similarly, the much lower pH_{med} of *nrt2.1* compared to WT at 48h was not reflected in its

pH_{apo}, which was similar to WT. It is generally assumed that pH_{apo} and pH_{med} are the same since

these compartments are physically connected. These discrepancies can be explained by the role

the cell wall plays in maintaining a low pH_{apo}. The cell wall's apoplast has two zones: the Water

Free Space, which mirrors the ionic concentration of the external solution (pH_{med}), and the Donnan

Free Space, where electrostatic charges from the wall concentrate cations (including protons) and

depletes anions (15). We propose that the proton binding capacity of the cell wall results in pH_{apo}

being lower than pH_{med}, providing the PM ATPase is sufficently active. In line with this, the apoplast

contained within the cell wall was found to be as much as 2 units more acidic than the remaining

apoplastic liquid, which was attributed the cell wall's high cation exchange capacity (16). Moreover,

Martinière et al. (17) found that the apoplast is considerably more acidic than the apoplastic liquid

immediately adjacent to the PM. Hence, the cell wall may act as a proton reservoir, that charges

itself through the action of the PM ATPase and by acquiring protons from the surrounding medium,

thus helping to sustain apoplastic acidity. To explain how NRT1.1A/B enhances apoplast

acidification, we posit that nitrate symport enhances the activation of H^+-ATPase, in order to

maintain a stable cytosolic pH (Fig. 2*B*).

Materials and Methods

The detailed methods are described in SI Appendix. Briefly, *M. truncatula* seedlings were grown

on agar plates containing nitrate-free FP nutrient solution for one week. The seedlings were then

transplanted into non-aerated liquid FP medium supplemented with different concentrations of

KNO_3 with their roots fully submerged and pH was monitored using a pH meter. For HPTS staining,

the seedlings were prepared in the same manner and then transferred to FP medium containing 5

mM KNO_3 for the given time, and then stained with HPTS for 30 minutes, rinsed three times with

FP, then root epidermal cells near the root tip were imaged using a confocal microscope. The *M.*

truncatula nrt1.1 and *nrt2.1* mutants were previously described (7, 18).

Acknowledgments

We would like to acknowledge Ping Xu, Yisheng Wang and Chao-Feng Huang for their comments

on the manuscript and Wenxian Lan and Wenjuan Cai of the CEMPS core facility for technical

support. Seeds of the *nrt2.1* mutant were kindly provided by Dr. Fang Xie.

References

1. H.-H. Tsai, and W. Schmidt, The enigma of environmental pH sensing in plants. *Nat.*

Plants 7:106–115 (2021).

2. A.J. Miller, and S.J. Smith, Nitrate transport and compartmentation in cereal root cells. *J.*

Exp. Bot. 47: 843–54 (1996).

3. M. Cerezo, *et al.*, Major alterations of the regulation of root NO$_3$- uptake are associated
with the mutation of *Nrt2.1* and *Nrt2.2* genes in *Arabidopsis*. *Plant Phys.* 127:262-71
(2001).

4. Y.F. Tsay, J.I. Schroeder, K.A. Feldmann, N.M. Crawford, The herbicide sensitivity gene
CHL1 of *Arabidopsis* encodes a nitrate-inducible nitrate transporter. *Cell* 72:705-13
(1993).

5. A. Maghiaoui, A. Gojon, L. Bach, NRT1.1-centered nitrate signaling in plants. *J. Exp. Bot.*
71:6226-6237 (2020).

6. X.Z. Fang *et al.*, Alleviation of proton toxicity by nitrate uptake specifically depends on
nitrate transporter 1.1 in *Arabidopsis*. *New Phyt.* 211:149-58 (2016).

7. Q. Xiao, et al., MtNPF6.5 mediates chloride uptake and nitrate preference in *Medicago*
roots. *EMBO J.* 40(21):e106847 (2021).

8. M. Haruta *et al.*, Molecular characterization of mutant *Arabidopsis* plants with reduced
plasma membrane proton pump activity. *J. Biol. Chem.* 285:17918-29 (2010).

9. J.Y. Pang, I. Newman, N. Mendham, M. Zhou, S. Shabala, Microelectrode ion and O$_2$
fluxes measurements reveal differential sensitivity of barley root tissues to hypoxia. *Plant,*
Cell & Environ. 29:1107-21 (2006).

10. P.R. McClure, L.V. Kochian, R.M. Spanswick, J.E. Shaff, Evidence for cotransport of
nitrate and protons in maize roots: I. Effects of nitrate on the membrane potential. *Plant*
Phys. 93:281-9 (1990).

11. S. Santi, G. Locci, R. Pinton, S. Cesco, Z. Varanini, Plasma Membrane H$^+$-ATPase in
maize roots induced for NO$_3$$^-$ uptake. *Plant Phys.*109:1277-1283 (1995).

12. S. Santi, G. Locci, R. Monte, R. Pinton, Z. Varanini, Induction of nitrate uptake in maize
roots: expression of a putative high-affinity nitrate transporter and plasma membrane H$^+$-
ATPase isoforms. *J. Exp. Bot.* 54:1851-64 (2003).

13. L. Espen, F.F. Nocito, M. Cocucci, Effect of NO$_3$$^-$ transport and reduction on intracellular
pH: an *in vivo* NMR study in maize roots. *J. Exp. Bot.* 55:2053-61 (2004).

14. H. Sze, J.M. Ward, S. Lai, Vacuolar H$^+$-translocating ATPases from plants: structure,
function, and isoforms. *J Bioenerg. and Biomemb.* 24:371-81 (1992).

15. D. S. Bush, and J. G. McColl, Mass-action expressions of ion exchange applied to Ca^{2+},
H$^+$, K$^+$, and Mg^{2+} sorption on isolated cell walls of leaves from *Brassica oleracea*. *Plant*
Physiol. 85:247-60 (1987).

16. H. Sentenac, C. Grignon, Effect of H$^+$ excretion on the surface pH of corn root cells
evaluated by using weak acid influx as a pH probe. *Plant Phys.* 84:1367–1372 (1987).

17. A. Martinière *et al.*, Uncovering pH at both sides of the root plasma membrane interface
using noninvasive imaging. *Proc. Nat. Acad. Sci. U.S.A.* 115:6488-6493 (2018).

18. Z. Luo, *et al*., The small peptide CEP1 and the NIN-like protein NLP1 regulate NRT2.1 to
mediate root nodule formation across nitrate concentrations. *Plant Cell* 35:776-794
(2023).
183
184

Figure 1. NRT2.1 and NRT1.1s affect nitrate-induced media alkalization. WT *M. truncatula*
seedlings were transferred to either FP (control), or FP supplemented with varying concentrations
of KNO_3 and pH_{med} was monitored (*A*). *M. truncatula* seedlings were transferred to either FP or FP
containing 5 mM KNO_3 with and without aeration (aer.) and pH_{med} was monitored (*B*). Seedlings
grown on FP were transferred to FP containing the indicated concentrations of PS-1 for 6h and
then pH_{med} was determined (*C*). WT and mutant *M. truncatula* seedlings were transferred to either
FP (*D*), or FP with varying concentrations of KNO_3 (*E-G*) and pH_{med} was monitored. In (*A*), the inset
shows a detailed view of the 3 and 6 h timepoints. For all experiments *N*=3, with 5 plants per
biological replicate. Error bars indicate SEM.

Figure 2. Nitrate induces a short-term decrease in pH_{apo} which requires NRT1.1A/B. Roots of *M. truncatula* seedlings were stained with HPTS and 458/405 ratios of the root tip cells were determined by confocal microscopy after immersion in FP (0 h) or at different times after addition of FP containing 5mM KNO_3 (*A*). *P-value<0.05, Dunnett's test, mutants vs WT at each time point. *N*=6-32. A model for nitrate uptake in plants (*B*). (left) Without aeration and in the absence of nitrate the rhizosphere and apoplast of immersed plants are acidified through the activity of H^+-ATPases that serve to maintain cytosolic pH during hypoxia. (right) When nitrate is added to the media, the activation of NRT1.1 and NRT2.1 symporters triggers an overcompensation of the PM H^+-ATPase through an unknown mechanism, which more than offsets H^+ uptake via nitrate symport, leading to a small decrease in pH_{apo}. The ongoing uptake of nitrate and protons via the apoplast by the symporters eventually leads to the alkalization of the rhizosphere.

Extended methods

Seed sterilization

Wild type *M. truncatula* (ecotype R108) and *nrt1.1a*, *nrt1.1b*, *nrt1.1c, nrt2.1* and the *nrt1.1a/b*
double mutant seeds were scarified with sandpaper, then surface-sterilized with 70% ethanol for
1 minute, then 10% bleach for 3 minutes, and then rinsed five times with sterile distilled water.

Plant culture

Sterilized *M. truncatula* (ecotype R108) and *nrt1.1a*, *nrt1.1b*, *nrt1.1c*, *nrt2.1* and the *nrt1.1a/b*
double mutant seeds were placed on 0.8 % agar plates and stratified in darkness at 4°C for 3 days.
The plates containing the seedlings were then placed upside down in the dark at 22°C to allow the
roots to extend overnight. The seedlings were then transferred to fresh 0.8 % agar plates containing
Fahraeus nitrogen free nutrient medium (FP) for one week with the plates vertically positioned.

pH measurements

For the pH measurements five of the seedlings were transferred to graduated 15 ml plastic tubes
filled to the 15 ml mark with FP medium (~13.2 ml of FP) supplemented with different concentrations
of nitrate with the pH adjusted using H_2SO_4 or KOH. Plants were grown under controlled conditions
with a 16-hour light period at 22°C and an 8-hour dark period at 20°C. The pH of the liquid medium
(pH_{med}) was measured every few hours using a pH meter (Mettler Toledo FE28) with a reported
accuracy of ± 0.01.

HPTS staining and imaging

HPTS is a membrane impermeant pH sensitive fluorescent dye[1,2,3]. For HPTS staining, plants were
grown on FP medium for one week as described above and then transferred to FP medium
containing 5 mM KNO_3 for 0, 1.5, 4, 24, or 48h. Seedlings were then transferred to the same
medium supplemented with 1 mM 8-hydroxypyrene-1,3,6-trisulfonic acid (HPTS) (Abcam, USA) for
30 minutes. Roots were then rinsed three times with HPTS-free growth medium, placed on a
microscope slide, and covered with a coverslip. Imaging of epidermal cells near the root tip was
carried out using an inverted Leica SP8 confocal microscope with a 20X water immersion objective.

249   HPTS fluorescence was detected using dual excitation at 405 nm and 458 nm, with emission
250   collected at 514 nm. The overall 458/405 ratio was calculated for each image using Fiji software.
251
252   **Seed sterilization**
253
254   Wild type *M. truncatula* (ecotype R108) and *nrt1.1a*, *nrt1.1b*, *nrt1.1c, nrt2.1* and the *nrt1.1a/b*
255   double mutant seeds were scarified with sandpaper, then surface-sterilized with 70% ethanol for
256   1 minute, then 10% bleach for 3 minutes, and then rinsed five times with sterile distilled water.
257
258   **Aeration experiments**
259
260   To provide gentle aeration of the medium a perforated plastic tube (0.5 mm outer diameter) was
261   placed in each 15 ml tube and connected to an SB-748 silent air pump (Zhongshan SOBO
262   electric appliance co., China) on its lowest setting.
263
264   **Measurement of oxygen levels**
265
266   Plants were grown on FP medium for one week as described above and then transferred to
267   aerated or non-aerated FP or FP supplemented with 5 mM KNO_3 and left for approximately 30
268   hours. Oxygen concentrations were then measured using an optical oxygen microsensor
269   (PreSens; Regensburg, Germany).
270
271   **PS-1 treatment**
272
273   PS-1 (TargetMol Chemicals Inc., USA) was stored as a 25 mM stock solution in DMSO. One-
274   week-old seedlings were prepared as described above were transferred to 15 mL tubes
275   containing aerated FP solution supplemented with PS-1 or mock treatment (equivalent amount of
276   25 µM DMSO) for 6 hours. The pH of the liquid medium was then measured.
277
278
279   **Statistical analyses**
280
281   All means were compared using Student's *t*-test or, when more than two means were compared
282   to controls using Dunett's Test to correct for multiple comparisons.
283
284

1. Lin W, Zhou X, Tang W, Takahashi K, Pan X, Dai J, Ren H, Zhu X, Pan S, Zheng H, *et al.* 2021. TMK-based cell-surface auxin signalling activates cell-wall acidification. *Nature* 599: 278–282.

2. Wang J, Jin D, Deng Z, Zheng L, Guo P, Ji Y, Song Z, Zeng HY, Kinoshita T, Liao Z, *et al.* 2025. The apoplastic pH is a key determinant in the hypocotyl growth response to auxin dosage and light. *Nature Plants* 11:279–294.

3. Barbez E, Dünser K, Gaidora A, Lendl T, Busch W. 2017. Auxin steers root cell expansion via apoplastic pH regulation in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 114(24):E4884-E4893.