Microbial-inspired antidotes to repurpose toxic compounds as antibiotics

AUTHORS: Tahoura Samad[1,7], Chayanon Ngambenjawong[1,5], Henry Ko[1], Savan Patel[6], Cristiana DeAgazio[1], Heather E. Fleming[1,2,8], Sangeeta N. Bhatia[1,2,4,6,7,8]

AFFILIATIONS:

1. Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology, Cambridge, MA 02139

2. Marble Center for Cancer Nanomedicine, Massachusetts Institute of Technology, Cambridge, MA 02139

3. Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139

4. Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Cambridge, MA 02139

5. School of Biomolecular Science and Engineering, Vidyasirimedhi Institute of Science and Technology (VISTEC), Rayong, Thailand

6. Harvard-MIT Division of Health Sciences and Technology, Cambridge, MA 02139

7. Institute of Medical Engineering and Science, Massachusetts Institute of Technology, Cambridge, MA 02139

8. Howard Hughes Medical Institute, Massachusetts Institute of Technology, Cambridge, MA 02139

ABSTRACT:

Antibiotic-resistant (AMR) bacterial infections are a major global health threat. Despite the critical need for new antimicrobials, progress is constrained by protracted development timelines, as well as the requirement for chemical novelty to avoid cross-resistance. Repurposing therapeutics approved for indications other than infection offers a potential shortcut to rapidly develop new antibiotics that are outside of existing antibiotic classes and therefore less susceptible to resistance. Cancer therapeutics, which often disrupt essential cellular processes that are evolutionarily conserved within bacteria, are a promising source of novel antimicrobial candidates. However, the narrow therapeutic index of these compounds limits their application for infection. Here, we establish a two-component strategy to repurpose highly toxic therapeutics as antimicrobials by improving their safety profile, using the cytotoxin calicheamicin as a proof-of-concept. In the first arm, we engineer a conditionally-active drug conjugate that restricts calicheamicin activity to infected tissue, improving its therapeutic index. In the second arm, we administer a re-engineered self-resistance enzyme from *Micromonospora echinospora*, the natural producer of calicheamicin, as an "antidote" to neutralize calicheamicin released outside of infected tissue, further widening its therapeutic index. The calicheamicin conjugate exhibits activity against a panel of Gram-negative and Gram-positive pathogens in response to a protease present within the infected microenvironment. Delivered in combination with the antidote, we demonstrate that antibacterial efficacy is maintained, and off-target toxicity is reduced in a mouse model of bacterial pneumonia. We anticipate that our dual strategy, which enhances drug safety by combining conditional activation of drug activity with neutralization of off-target effects, provides a generalizable framework for repurposing other promising but toxic compounds as antimicrobials.

INTRODUCTION:

Antimicrobial-resistant (AMR) infections are a critical health threat. The World Health Organization (WHO) estimates that resistance to existing drugs directly caused 1.27 million deaths globally and contributed to an additional 4.95 million deaths in 2019 (*1*). Development of new agents to address AMR infections is slow, hindered by the lengthy and costly processes of drug discovery and regulatory approval (*2*), and further challenged by cross-resistance, in which newly developed compounds that are similar to antibiotics are subject to the rapid development of resistance(*3*). Additionally, the economics of antibiotic treatment, which can involve short courses of treatment and stewardship programs to limit use, disincentivize investment in antimicrobial development by industry (*2*). Together, these scientific, regulatory, and economic hurdles have resulted in a widening gap between the rise of AMR and the pace of antibiotic development. Given this landscape, there is a critical need for innovative approaches to create antibiotics rapidly, while introducing novel mechanisms of action that help insulate against cross-resistance.

Current methods to develop new antibiotics often involve high throughput screening of small molecule libraries against predicted antimicrobial targets or whole cells. While tools such as machine learning (*4–7*), large-scale chemical-genetics (*8*), computational modeling of compound-target interactions (*9*), and genome mining (*10*, *11*) have greatly enhanced this process, promising lead compounds still face a prolonged and expensive development pipeline. Screening for companion compounds that can amplify the effectiveness of existing antibiotics (*9*, *12*) or mitigate drug resistance (*13*) is another strategy to accelerate drug development and has yielded notable successes, such as Augmentin (*13*, *14*). However, the use of existing antibiotics in these combination approaches may still leave them vulnerable to common resistance mechanisms (*13*). Alternatively, leveraging drugs approved by regulators for conditions other than infection, known as drug repurposing, might offer a pool of antibiotic candidates that are distinct

from existing approved antimicrobials while also shortening the development timeline (*15*).

Cancer therapeutics, many of which disrupt fundamental, conserved biological processes such

as DNA replication and metabolism, are a rich source of compounds with potential antimicrobial

activity. However, the narrow therapeutic index of these agents—the difference between the

effective therapeutic dose and the dose that causes substantial toxicity (Fig. 1A)—limits their

direct translation for infectious disease applications. Toxicities routinely tolerated in oncology,

such as myelosuppression (*16*), neuropathy (*17*), or even secondary malignancies (*18*), would be

unacceptable for patients with acute infections, who are frequently otherwise healthy.

Repurposing chemotherapeutic agents for antimicrobial use will therefore require innovative

strategies to widen their therapeutic index.

In this study, we present a two-armed, generalizable strategy for repurposing toxic

compounds as antimicrobials by widening their therapeutic index. As a proof-of-concept, we

apply this strategy to calicheamicin, a potent DNA-damaging agent that is approved for cancer

treatment, but which causes significant hepatotoxicity that would preclude administration to

patients with infections (*19, 20*). In the first arm of our approach, we formulate calicheamicin as

a conditionally-active conjugate, adapting design principles that have been used to deliver

potent cytotoxic payloads in oncology (17,18). In this format, we tether calicheamicin to a

carrier protein through a protease-cleavable linker. While conjugated, the cytotoxic activity of

calicheamicin is masked until an infection-associated protease releases the drug, thereby

confining activity to infected tissue and reducing toxicity to healthy tissue, widening the

therapeutic index (Fig. 1B). Recognizing that even limited exposure of healthy tissue to a

compound as potent as calicheamicin can cause severe damage, we introduced a second arm

into our strategy to neutralize any drug released or diffusing beyond the infection site**.** To this

end, we adapted a natural microbial strategy, the self-resistance mechanism of

Micromonospora echinospora, the natural producer of calicheamicin. *M. echinospora* produces

several enzymes that neutralize calicheamicin to protect itself from autotoxicity (*21, 22*); we

hypothesized that we could engineer and administer these enzymes as "antidotes" to neutralize

off-target calicheamicin, further widening the therapeutic index (Fig. 1C). We first show that our

calicheamicin-containing drug conjugate exhibits activity dependent on an infection-associated

protease against a panel of Gram-negative and -positive priority pathogens, and that prior to

cleavage, this conjugated format masks calicheamicin toxicity to mammalian cells. We further

show that bacterial-derived calicheamicin self-resistance enzymes can be re-engineered to act

as long-circulating "antidotes" that neutralize off-target calicheamicin *in vitro* and *in vivo*,

reducing liver toxicity. Finally, we validated that the combination of these two strategies

achieves therapeutic efficacy in a mouse model of pneumonia and reduces systemic toxicity.

Our dual approach provides a modular framework for the rapid repurposing of other promising-

yet-toxic natural products for use as antibiotics.

RESULTS

Calicheamicin has broad spectrum activity against Gram-positive and Gram-negative

pathogens

We first evaluated the antimicrobial activity of calicheamicin against a panel of Gram-

negative and Gram-positive bacterial pathogens, including several designated as priority

pathogens by the World Health Organization (*23*). Measuring microbial susceptibility to

calicheamicin using broth microdilution assays, we determined the minimum inhibitory

concentration (MIC) of calicheamicin across representative strains (Table 1). We found that

calicheamicin had potent activity against the Gram-positive bacterium *Staphylococcus aureus*,

with an MIC of 0.0006 µg/mL, a potency that was comparable to or greater than that of a

clinically used antibiotic vancomycin (Table 1). Notably, this potency was maintained against a

methicillin-resistant *S. aureus* (MRSA) isolate, BA1556. Calicheamicin also demonstrated

activity against Gram-negative pathogens including *Pseudomonas aeruginosa*, *Escherichia coli*,

and *Klebsiella pneumoniae*, with MIC values ranging from 0.015 to 0.5 µg/mL (Table 1).

Together, these data demonstrate that calicheamicin exhibits broad-spectrum antimicrobial

activity, with efficacy extending to multidrug-resistant strains, and at concentrations that would

be realistic for clinical translation.

130 Conditionally-active calicheamicin conjugate masks cytotoxicity while retaining

131 antibacterial activity *in vitro*

While the potent antimicrobial activity of calicheamicin is a promising starting point, its

substantial toxicity toward mammalian cells, particularly liver cells, precludes its use as a free

compound. In oncology, this limitation has been addressed by formulating calicheamicin as an

antibody–drug conjugate (*24, 25*), where targeting and conditional activity improve the

therapeutic index. Building on this approach, we developed a conditionally-active calicheamicin

conjugate tailored for the infection context. In this design, calicheamicin was tethered to an

albumin-binding domain (ABD) via a protease-sensitive linker that masks its cytotoxic activity

until cleavage by infection-associated proteases (Fig. 2A). Use of albumin as the carrier protein

leverages endogenous trafficking pathways to enhance accumulation (*26, 27*) , retention (*28*),

and mucopenetration (*29*) at sites of infection. As a linker between albumin and the drug, we

used a peptide sequence cleavable by neutrophil elastase, which is enriched at sites of bacterial

infection due to immune cell recruitment.

To generate the ABD–calicheamicin conjugate (ABD-Cal), we prepared a calicheamicin–

peptide intermediate and coupled it to a recombinantly-expressed ABD using strain-promoted

azide–alkyne cycloaddition. We next tested whether conjugation masked the cytotoxic activity

of calicheamicin and allowed for protease-dependent activation by assessing antibacterial

activity and mammalian cell toxicity in the presence or absence of neutrophil elastase *in vitro*.

The conjugate was incubated with neutrophil elastase or vehicle control for 4 hours prior to

evaluation. We then measured antibacterial activity against a representative Gram-positive (*S.*

aureus) (Fig. 2B) and Gram-negative (*K. pneumoniae*) pathogen (Fig. 2C). In the absence of

protease, we observed no antimicrobial activity; in the presence of protease, we observed a

recovery in antimicrobial activity with a (~60-120-fold) change in the minimum inhibitory

concentration (MIC) between the intact and cleaved conjugates, depending on the pathogen

tested. We also assessed whether formulation of calicheamicin as a conditionally-active

conjugate protected mammalian cells from toxicity. We found that in the absence of protease,

the intact conjugate demonstrated minimal mammalian toxicity to HepG2 liver cells after 24

hours of exposure (Fig. 2D). Together, these data show that calicheamicin, when formulated as

a conditional conjugate, exhibits reduced toxicity toward mammalian cells while conjugated, but

regains potent antibacterial activity once released by protease cleavage.

Conditional calicheamicin conjugate demonstrates improved toxicity relative to free drug

in vivo

Having shown in vitro that conjugation masks calicheamicin's cytotoxicity and that

protease cleavage restores antibacterial activity, we next asked whether restricting activation to

the infection microenvironment reduces systemic, particularly hepatic, toxicity in vivo. To test

this, we compared the liver toxicity of the conditionally active calicheamicin conjugate with free

calicheamicin *in vivo*. Healthy mice were intravenously administered either free calicheamicin or

the conjugate (dose matched by drug concentration) (Fig. 2E). Liver injury was assessed 36 h

post-injection by measuring serum aspartate aminotransferase (AST) levels (Fig. 2F), alanine

aminotransferase (ALT) (Fig. 2G) and by histological analysis (Fig. S1). Mice treated with free

calicheamicin exhibited elevated AST (Fig. 2F) and ALT (Fig. 2G) levels. Mice receiving the

conjugate showed a significant reduction in toxicity relative to free drug, although serum liver

enzymes were significantly elevated above the baseline of mice treated with vehicle alone.

Histology revealed moderate to severe injury in mice treated with free drug, evidenced by

diffuse presence of hepatocytes with swollen cytoplasm or shrunken, hypereosinophilic cells

(Fig. S1) The livers of conjugate-treated mice showed more mild changes, including some

apoptotic cells with swollen cytoplasm, and some healthy cells, consistent with more limited

toxicity (Fig. S1). These findings indicate that formulation of calicheamicin as the conditionally-

active conjugate significantly reduces off-target hepatotoxicity *in vivo.*

Microbial-derived antidotes can neutralize calicheamicin toxicity *in vitro* and *in vivo*

Formulation of calicheamicin as a conditional conjugate significantly reduced

hepatotoxicity in healthy mice but did not eliminate liver injury, which we hypothesized arose

from a fraction of drug released outside the infection site, potentially via low levels of protease-

mediated activation outside the infection environment, or diffusion of released drug from

infected tissue back into the bloodstream. We therefore developed a second complementary

strategy to address these residual sources of toxicity and further improve the safety profile of

calicheamicin. To achieve a second layer of toxicity mitigation, we drew inspiration from the

natural self-resistance mechanism of *Micromonospora echinospora*, the native producer of

calicheamicin, which avoids self-toxicity by enzymatically inactivating the compound. We

envisioned adapting this self-resistance strategy for therapeutic use by engineering long-

circulating variants of these calicheamicin-neutralizing enzymes to act as "antidotes",

intercepting and neutralizing calicheamicin that escapes the infection microenvironment before it

reaches sensitive tissues (Fig. 3A).

We first evaluated which of the native self-resistance enzymes would be most suitable

for adaptation as a circulating antidote. The calicheamicin biosynthetic cluster encodes three

self-resistance enzymes, CalC, CalU16, and CalU19 (*21*, *22*). Reduction of calicheamicin's

trisulfide bond generates the reactive species responsible for its cytotoxicity. CalC and CalU16

act only on this reduced form, whereas CalU19 also neutralizes the unreduced molecule,

making it best suited for calicheamicin neutralization in circulation, which is not a reducing

environment. We recombinantly expressed CalU19 and sought to confirm its neutralizing activity

in vitro, by evaluating if administration of CalU19 could protect mammalian cells from

calicheamicin-induced toxicity. HepG2 cells were incubated with serial dilutions of calicheamicin

in the presence or absence of a fixed concentration of CalU19 for 24 hours, and cell viability

was assessed by MTS assay (Fig. 3B). CalU19 conferred protection, with the highest protection

observed at lower calicheamicin concentrations, consistent with the stoichiometric mechanism

of inactivation in which one molecule of CalU19 is consumed per molecule of calicheamicin

neutralized (Fig. 3B).

Following *in vitro* validation of CalU19 activity, we next assessed its ability to selectively

neutralize free calicheamicin without affecting the conjugated drug. During calicheamicin

neutralization, CalU19 undergoes a unique self-cleavage event, which can be used as an

indicator of neutralizing activity. SDS-PAGE of CalU19 incubated with calicheamicin, ABD-Cal

or vehicle, revealed CalU19 self-cleavage fragments only when CalU19 was incubated with free

calicheamicin (Fig. S2), confirming that neutralizing activity is restricted to free drug.

To realize our design goal of a long-circulating antidote capable of soaking up off-target

drug *in vivo*, we next engineered CalU19x for *in vivo* administration. We first removed two

unpaired cysteines, which could promote off-target interactions with other proteins in vivo,

substituting them with alanine to generate CalU19x, which retained neutralizing activity

comparable to the wild-type enzyme (Fig. 3B). We then assessed CalU19 *pharmacokinetics in*

vivo (Fig. 3C, D). Fluorophore-labeled CalU19 and CalU19x were both rapidly cleared from

circulation, with plasma half-lives of ~1 h. To extend the circulation half-life of CalU19x, we

generated fusions to single (C-CalU19x) and tandem (LC-CalU19x) albumin-binding domains,

as these extend half-life through association with serum albumin. LC-CalU19x exhibited the

longest half-life, remaining detectable for at least 48 h in plasma (Fig. 3D), and maintained

neutralizing activity *in vitro* comparable to wild-type CalU19 (Fig. 3B).

Having established a long-circulating antidote *in vitro*, we next evaluated whether LC-

CalU19x could protect against calicheamicin toxicity *in vivo*. Healthy animals were intravenously

administered LC-CalU19x or vehicle followed by dosing with calicheamicin (Fig. 3E). Serum and

tissue were collected 36-48 h later, and serum AST and ALT levels were quantified. Mice

administered calicheamicin and LC-CalU19x showed significantly reduced serum AST (Fig. 3F)

and ALT (Fig. 3G) compared with mice administered calicheamicin and vehicle, indicating

mitigation of calicheamicin-induced hepatotoxicity. Histology (Fig. S3) agreed with these

findings: livers from mice administered calicheamicin and LC-CalU19x-treated mice were largely

indistinguishable from healthy controls, whereas mice treated with calicheamicin alone

displayed hepatocellular degeneration and necrosis, indicated by swollen cells with eosinophilic

cytoplasm (Fig. S3). Together, these results demonstrate that LC-CalU19x acts as a circulating

antidote that mitigates off-target toxicity without interfering with the activity of the conditionally-

active calicheamicin conjugate, supporting their use as a combined therapy.

Calicheamicin conjugate and antidote combination achieves antimicrobial efficacy in

mouse models of pneumonia

We proceeded to test the preclinical potential of ABD-calicheamicin conjugate and LC-

CalU19x in combination, evaluating whether co-administration of the antidote with the

conditionally-active drug would reduce bacterial burden, while minimizing liver toxicity in a mouse

model of bacterial pneumonia. Mice were infected with bacterial pathogen *S. aureus*, followed by

intratracheal administration of vehicle, ABD-calicheamicin alone, or ABD-calicheamicin in

combination with intravenously administered LC-CalU19x (Fig. 4A-B). Treatment of *S. aureus*-

infected mice with the conditionally-active conjugate and antidote resulted in a 4-log reduction in

lung bacterial burden 6 h after administration, compared with the vehicle control, demonstrating

that on-target antimicrobial efficacy is preserved in the presence of the antidote (Fig. 4B).

Comparing liver toxicity between mice treated with ABD-Cal alone or ABD-Cal with CalU19x, we

observed a significant reduction in toxicity for mice treated with conjugate and antidote relative to

the conjugate-only and vehicle groups, as measured by serum AST (Fig. 4C) and ALT (Fig. 4D),

with levels in the antidote-treated mice comparable to the untreated healthy control. Together,

these findings demonstrate that pairing a conditionally-active calicheamicin conjugate with a long-

circulating enzyme antidote maintains antimicrobial efficacy while substantially reducing systemic

toxicity.

DISCUSSION

In this work, we demonstrate two complementary strategies to expand the therapeutic

window of a chemotherapeutic drug, calicheamicin, drawing inspiration from 1) engineered

oncology approaches for managing toxic therapeutics and 2) natural microbial mechanisms for

detoxification. In oncology, antibody–drug conjugates (ADCs) take advantage of

microenvironmental cues to restrict payload activity to the site of disease; we created a

calicheamicin conjugate that restricts drug activation to infected tissue, reducing systemic

toxicity. In parallel, we take advantage of the natural self-resistance mechanism from the

calicheamicin producing organism, optimizing and then administering a calicheamicin-

neutralizing "antidote" to inactivate off-target free drug. The combination of conjugate and

antidote successfully treated bacterial pneumonia in mice while reducing hepatotoxicity,

demonstrating that the two complementary strategies, conditional activation and antidote co-

administration, can widen the therapeutic window of a potent cytotoxin.

Conditional drug conjugates hold considerable potential for infectious disease, but their

development in that context is far less common than in oncology, where antibody–drug

conjugates (ADCs) have achieved notable clinical success (*26*, *27*, *30*, *31*). The distinct

biological context of bacterial infections requires revising design criteria that have guided ADC

development in oncology. While oncology relies on antibodies to direct cytotoxic payloads to

defined tumor antigens, infection-focused conjugates would benefit from carrier proteins that

target the site of infection in an antigen-independent manner and are thus compatible with

broad-spectrum activity against a variety of pathogens. Our design (Fig. 2) uses an albumin-

binding domain (ABD) as our drug carrier, with the intent to promote retention (*28*) and

mucopenetration (*29*) at the sites of infection when administered into the lung. For the linker,

bacterial infection most likely will require extracellular drug release (except in the case of

intracellular pathogens), which is a heterogeneous environment, where disease-associated

triggers act amid variable pH and redox states in the presence of immune cells and cellular

debris. In contrast, ADCs for cancer typically exploit tumor-associated proteases such as

cathepsins to release payloads intracellularly within lysosomes, a defined environment with

acidic pH, reducing conditions, and a well-characterized protease repertoire (*32*). To achieve

robust and selective activation in this setting, we incorporated a peptide linker cleavable by

neutrophil elastase, which is enriched extracellularly at sites of bacterial infection after release

by recruited immune cells. The criteria for payload selection are consistent between the two

indications: potency is indispensable for efficacy. Early preclinical ADCs used conventional

chemotherapeutics with modest potency, which limited their efficacy (33). Recognition that

greater potency was required led to the development of gemtuzumab ozogamicin (Mylotarg),

the first FDA-approved ADC, which used calicheamicin as its payload. Although it achieved

strong clinical responses, its severe dose-limiting toxicities underscored the risks associated

with ultrapotent payloads (34). The use of calicheamicin in our conjugate therefore represents a

stringent test of whether such highly potent molecules can be safely repurposed for

antimicrobial use.

Formulation of calicheamicin as a conjugate reduced liver toxicity but did not fully

eliminate it (Fig. 2), consistent with both preclinical and clinical data from oncology. While

conditional drug conjugates have demonstrated clear clinical success (*33*), some evidence

suggests that the maximum tolerated doses of some ADCs approach those of the unconjugated

payload (*20*, *34–36*). Thus, ADCs improve efficacy but do not always reduce toxicity, and the full

therapeutic potential of this modality has not been realized. Prior efforts have focused on

refining the antibody, linker, or payload to reduce toxicity, but off-target damage in the liver,

bone marrow, and gastrointestinal tract remains a major barrier. To move beyond these

limitations in the context of infection, we turned to microbes themselves, which not only produce

some of the cytotoxic natural products used as ADC payloads but also encode self-resistance

enzymes to protect against autotoxicity. Inspired by this microbial strategy, we introduced a

second layer of protection: delivery of an engineered self-resistance enzyme derived

from *Micromonospora echinospora*, the natural producer of calicheamicin, as an antidote to

neutralize circulating free drug outside the site of infection (Fig. 3). While antibiotic adjuncts that

enhance antimicrobial activity are under development (*37–40*), coadministration of agents

specifically designed to mitigate host toxicity is less explored in infectious disease. This model is

under active consideration within oncology contexts, for example in preclinical work examining

neutralization of toxicity for radiation (*41*), and clinically in the use of dasatinib (*42*) to mitigate

cytokine release syndrome after administration of CAR-T therapy.

Despite the promise of our drug conjugate and antidote platform, several challenges

may remain before translation to clinical settings. Microbial-derived proteins like the antidote we

developed may be immunogenic upon repeated administration, as is often required in the

treatment of bacterial infections. Existing strategies such as PEGylation (*43*), domain

humanization (*44*), or epitope masking (*45*) could reduce immunogenic potential, and there are

encouraging examples of regulator-approved and clinically-used products derived from

microbial enzymes (*46*, *47*). Second, the success of this dual system hinges on spatial, kinetic,

or temporal separation between on-target activity and off-target detoxification. In our current

model, we achieve this using compartmentalization, via local pulmonary delivery of the drug

conjugate and systemic administration of the antidote (Fig 4). This separation is also achieved

by virtue of the conjugate, which protects the drug from premature neutralization by the antidote,

as the antidote can only bind and inactivate the released, active form of the drug (Fig. S2).

However, extending the paired conjugate-antidote system to other sites of infection will require

context-specific tailoring. For example, we hypothesize that treatment of systemic bloodstream

infections would likely require prophylactic delivery of CalU19, either as a protein or mRNA

encapsulated in lipid nanoparticles, such that they can be taken up by liver sinusoidal

endothelial cells or hepatocytes to protect these tissues prior to systemic administration and

release of calicheamicin from the conjugate in response to cues from the bloodstream infection.

The modularity of our platform, with the ability to exchange the carrier protein and linker within

the conjugate and the possibility of fusing other targeting partners to CalU19, to control half-life

or localization within the body, makes such adaptation feasible. Finally, the kinetics of enzyme-

mediated neutralization of "off-target" drug relative to bacterial uptake and killing by "on target"

drug remain incompletely understood. Future studies may benefit from structure-guided

engineering or directed evolution of CalU19 to modulate its catalytic efficiency to further

enhance the therapeutic window of the conjugate and antidote pair.

Beyond calicheamicin, our dual strategy—1) leveraging selective activity within the

infected microenvironment and (2) engineering of microbial resistance proteins as adjuncts to

neutralize off-target toxicity--offers a generalizable framework to rapidly mobilize other

promising but previously unusable compounds for antimicrobial development. The modular

nature of our system, which allows for the easy swapping of targeting moieties, linker and drug,

enables facile adaptation to varied infection contexts. Potentially valuable classes of compounds

for this dual strategy include cytotoxic therapeutics with narrow therapeutic indices, antibiotic

candidates that failed in late-stage clinical trials due to toxicity, and preclinical compounds

developed in pharmaceutical pipelines but later abandoned, due to toxicities and/or poor

pharmacokinetics, with eventual disinvestment by industry (*48–51*). By mobilizing such agents

with conditional activation and antidote-mediated neutralization, these candidates could be re-

evaluated with the advantage of an accelerated development timelines and reduced costs from

leveraging existing chemistry, pharmacology, and, in some cases, clinical data. Looking ahead,

genome mining and artificial intelligence could accelerate the discovery and prioritization of

naturally produced drug–antidote pairs, leveraging microbial self-resistance strategies such as

enzymatic inactivation or compound sequestration. Beyond mining natural mechanisms, these

same tools may also enable *de novo* design of antidotes, creating entirely new ways to widen

the therapeutic index.

More broadly, the combined use of the drug conjugate and antidote represents a

conceptual shift in antibiotic treatment—towards a form of molecular stewardship—in which

each molecule of the antimicrobial is accounted for, with on-target directed precisely to the site

of infection while off-target drug is actively neutralized. Although mitigation of liver toxicity was a

primary goal of this work, the same principle could reduce collateral damage to other sensitive

tissues, such as the bone marrow. Antibiotic-induced dysbiosis of the gut microbiome is another

major concern, such that antibiotic treatment is frequently associated with shifts in microbial

composition, overgrowth of opportunistic pathogens such as *Clostridioides difficile*, and impaired

gut barrier function (*52, 53*). We hypothesize that co-administration of an antidote could

attenuate antibiotic-associated microbiome disruption by neutralizing drug before it reaches the

gastrointestinal tract. Beyond the host, excreted antibiotics that enter into the environment

through sewage and wastewater exert selective pressure on environmental bacteria, which can

drive the emergence of resistance through exposure to sublethal concentrations (*54*). The

excretion of highly potent cytotoxic compounds such as calicheamicin also raises concerns

about ecological toxicity (*55–57*). Incorporation of antidotes into therapeutic regimens may

therefore not only improve safety within the patient but also mitigate the broader ecological

consequences of antimicrobial use. Finally, outside of the context of infectious disease, we

envision antidotes as an enabling technology for conditionally-active conjugates more generally,

with the potential to mitigate toxicity in oncology and other therapeutic indications where potent,

conditionally-active drugs are administered.

ACKNOWLEDGEMENTS, FUNDING AND COMPETING INTERESTS

We are grateful to Dr. Laura Kiessling and Dr. Deborah Hung for their scientific insight and support over the course of the work. We are grateful to the Koch Institute Swanson Biotechnology Center, specifically the Histology core (Kathy Cormier, Magalie Boucher) and the Biopolymer and Proteomics core (Heather Amoroso). We are grateful to Kasturi Chakraborthy, Susanna Elledge and Vardhman Kumar for feedback on the manuscript. This work was supported by R01 AI132413 and U19 AI142780 grants from the National Institute of Allergy and Infectious Diseases. This study was supported in part by the Koch Institute Support Grant P30-CA14051 from the National Cancer Institute (Swanson Biotechnology Center) and Core Center Grant P30-ES002109 from the National Institute of Environmental Health Sciences. Additional support was received from the Koch Institute's Marble Center for Cancer Nanomedicine. This research was supported in part by a generous gift from Joel Gantcher. T.S. was supported by a postdoctoral fellowship from the Ludwig Center at MIT's Koch Institute for Integrative Cancer Research. S.N.B. is a Howard Hughes Institute Investigator. Competing interests: S.N.B. reports consulting roles and/or equity Amplifyer Bio, Catalio Capital, Danaher, Earli Inc., Impilo Therapeutics, Matrisome Bio, Ochre Bio, Port Therapeutics, Satellite Bio, Sunbird Bio, and Xilio Therapeutics, which were not involved in this study.

MATERIALS and METHODS

Bacterial strains and culture

Escherichia coli ATCC 29522, *Klebsiella pneumoniae* ATCC 43816, *Pseudomonas*

aeruginosa PAO1, *Staphylococcus aureus* 502A, and *Staphylococcus*

aureus USA300_FPR3757 were used in this study. Strains were obtained from the American

Type Culture Collection (ATCC) or as indicated. Bacteria were cultured in Mueller–Hinton broth

(MHB; BD Difco) at 37 °C with shaking at 250 r.p.m. Growth was monitored by optical density at

600 nm (OD600).

Molecular cloning

gBlock™ gene fragments encoding ABD with a C-terminal cysteine ((ABD)$_2$-(EEG)$_6$-Cys), CalC,

and CalU19 with flanking restriction sites NcoI and XhoI were ordered from Integrated DNA

Technologies, cloned into pET28(a) by restriction enzyme digestion and ligation, and transformed

into *E. coli* DH5α competent cells (New England Biolabs). Colonies encoding the correct inserts

were confirmed by Sanger sequencing (QuintaraBio), and plasmids were purified using a plasmid

miniprep kit (Thermo Fisher Scientific). For CalU19 variants, clonal genes were directly obtained

from Twist Bioscience already cloned into the pET28(a) expression vector. All expression

plasmids were subsequently transformed into *E. coli* BL21(DE3) cells for protein expression.

Protein expression

A secondary culture (500 mL LB broth (50 µg/mL kanamycin)) of *E coli.* BL21(DE3) encoding the

protein of interest was expanded from an overnight primary culture (3 mL) in an incubator shaker

at 37 °C until OD600 reached about 0.6 (~ 4hours). Protein expression was induced by addition

of an IPTG solution to a final concentration of 1 mM. The induced culture was incubated in an

incubator shaker for 2 h at 37 °C after which the culture was centrifuged and the bacteria pellet

was frozen and stored in a -80 °C freezer. For purification, the bacteria pellet was thawed in a 37

°C water bath and lysed with a B-PER™ complete bacterial protein extraction reagent

(ThermoFisher). The lysed solution was centrifuged and the supernatant was incubated with

Qiagen Ni-NTA agarose for 30minutes at 4 °C to capture the product. The agarose was

transferred to a fritted column, washed sequentially with a wash buffer A (50 mM Tris, 500 mM

NaCl, 2% Triton X-114) (10 column volumes (CVs)) and a wash buffer B (50 mM Tris, 500 mM

NaCl) (10 column volumes) before elution with an elution buffer (500 mM imidazole, 50 mM Tris,

500 mM NaCl, 10% glycerol) (3 CVs). The purified protein was confirmed via SDS-PAGE analysis

with Coomassie blue staining.

Synthesis of Conjugate Azide-AAPV-calicheamicin

An N-terminal azidoacetylated peptide substrate with a C-terminal penicillamine (Pen) (Azide-

AAPV-Pen) was synthesized by CPC Scientific via a standard Fmoc-based solid phase peptide

synthesis. A solution of calicheamicin (1 eq.) (MedChemExpress) in N,N-dimethylformamide

(DMF) was added to a solution of Azide-AAPV-Pen (3 eq.) in DMF followed by addition of

triethylamine (8 eq.) to the reaction solution. The reaction solution was purged with nitrogen for

5minutes and incubated for 24hours at 4 °C. The Azide-AAPV-calicheamicin product was purified

by the Koch Institute Biopolymers & Proteomics core facility using a semi-preparative HPLC

column and confirmed with MALDI-ToF MS.

(ABD)$_2$-(EEG)$_6$-Cys was incubated with TCEP (10 eq.) for 30minutes at RT and buffer-exchanged

into PBS (1 mM EDTA, pH 6.5) using an Amicon 10-kDa, ultra-0.5 centrifugal filter unit (14,000

rpm, 2minutes, 4 times). The reduced protein was incubated with DBCO-Maleimide (4 eq.) for

4hours at RT. The ABD-DBCO product was purified and buffer-exchanged into PBS using a PD-

10 desalting column (Cytiva).

Synthesis of ABD-calicheamicin conjugates

Azide-Sx-calicheamicin (2 eq.) was dissolved in DMF and added to a solution of ABD-DBCO (1

eq.) in PBS to a final concentration of 50% DMF. The reaction solution was incubated for 24hours

at RT and centrifuged to remove precipitate. The product in supernatant was immobilized on Ni-

NTA agarose, washed with a wash buffer C (50 mM Tris, 500 mM NaCl, 20% DMF) (10 CVs) and

a wash buffer B (10 CVs) before elution with the elution buffer. The product was buffer-exchanged

into PBS using a PD-10 desalting column and quantified via absorbance at 200 nm. The product

was confirmed and assessed for percentage purity using SDS-PAGE analysis.

***In vitro* activity and toxicity masking**

To evaluate conditional activation, ABD–calicheamicin conjugates were incubated with

recombinant human neutrophil elastase (Enzo Life Sciences) or vehicle control in PBS at 37 °C

for 4 h. Reaction mixtures were then diluted into bacterial or mammalian assay media as

described below. Minimum inhibitory concentrations (MICs) were determined in accordance with

CLSI guidelines. Briefly, bacteria were seeded at OD 0.005 in 96-well plates containing MHB

supplemented with conjugate, protease-treated conjugate, or controls. Plates were incubated at

37 °C for 24 h, and bacterial growth was assessed by measuring optical density at 600 nm

(SpectraMax i3, Molecular Devices). MICs were defined as the lowest concentration of

conjugate preventing visible bacterial growth. Human hepatocellular carcinoma cells (HepG2,

ATCC HB-8065) were maintained in Dulbecco's Modified Eagle Medium (DMEM; Thermo

Fisher Scientific) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin–

streptomycin. For cytotoxicity assays, HepG2 cells were seeded at 1×10^4 cells per well in

collagen-coated 96-well plates and allowed to adhere overnight. Cells were then exposed to

intact or protease-treated ABD–calicheamicin conjugates for 24 h. Viability was quantified using

the CellTiter AQueous MTS assay (Promega) according to the manufacturer's instructions, and

absorbance was read at 490 nm using a plate reader. In vitro activity and toxicity data were fit

using a four-parameter logistic (4PL) sigmoidal model, where X represents concentration, in

GraphPad Prism (GraphPad Software).

472   *In vitro* **CalU19 neutralization evaluation**

473   *SDS-PAGE* CalU19 was labeled with Cy7 through random lysine labeling and incubated with

474   vehicle, calicheamicin, or ABD-Cal for 4 hours at 37C. At 4h, reactions were run on a gel

475   (NuPAGE™ 4-12% Bis-Tris Protein Gels, 1.0 mm, 12-well, Thermofisher) which was then

476   imaged using an Odyssey CLx imager in the 800 nm channel, for detection of Cy7 fluorescence,

477   monitoring specifically for the presence of the intact protein at ~27 kDa, as well as the self-

478   cleavage fragments at 6kDa and 21 kDa.

479   *Cytotoxicity Assay* HepG2 cells were seeded at 1×10^4 cells per well in collagen-coated 96-

480   well plates and allowed to adhere overnight. Cells were then exposed to serial dilutions of

481   calicheamicin, in the presence of absence of 5 µM CalU19, CalU19x, C-CalU19X, LC-CalU19x,

482   or vehicle (for the control). Viability was quantified at 24 h using the CellTiter AQueous MTS

483   assay (Promega) according to the manufacturer's instructions, and absorbance was read at 490

484   nm using a plate reader. Data were fit using a four-parameter logistic (4PL) sigmoidal model,

485   where X represents concentration, in GraphPad Prism (GraphPad Software).

486   *In vivo* **pharmacokinetics, activity and toxicity evaluation**

487   All animal studies were approved by the Massachusetts Institute of Technology's Committee on

488   Animal Care (MIT CAC protocol 2203000310). For infected mice, bacterial inocula were

489   prepared from mid-logarithmic phase cultures grown in Mueller–Hinton broth, washed twice with

490   PBS, and quantified by optical density and serial dilution plating. Following administration, mice

491   were monitored continuously until recovery from anesthesia and then observed at regular

492   intervals for clinical signs of infection, including weight loss, respiratory distress, and reduced

493   activity. For lung infections, CD-1 mice (10-12 weeks old) were administered 1.25×10^8 CFU of

494   *S. aureus* (502A) (suspended in PBS) by intratracheal administration using a 22G blunt end

catheter and monitored after administration. At designated time points post-infection, mice were

euthanized, lungs were aseptically harvested, and/or blood was collected.

CFU quantification Lungs were weighed and homogenized in sterile PBS using gentleMACS M-

tubes (Miltenyi Biotec). Homogenates were serially diluted and plated on mannitol salt agar (BD

Difco) to selectively recover *S. aureus*, with colonies were enumerated after 24 h incubation at

37 °C.

Histology Lungs were fixed in 10% neutral buffered formalin for at least 48 h, processed, and

paraffin-embedded by the Koch Institute Histology Core Facility. Tissue sections (X µm) were

stained with hematoxylin and eosin (H&E) and imaged using brightfield microscopy. Histological

evaluation was performed by

Toxicity evaluation To evaluate liver toxicity, blood was collected from mice at designated time

points via terminal cardiac puncture. Whole blood was allowed to clot at room temperature in

BD Microtainer SST tubes with gold tops (BD Biosciences, Cat. No. 365967). The resulting

serum was aliquoted and stored at –80 °C until analysis. Serum alanine aminotransferase (ALT)

activity was quantified using a colorimetric assay kit (Thermo Fisher Scientific, ALT/GPT Activity

Assay Kit, Cat. No. EEA001, and serum aspartate aminotransferase (AST) activity was

measured using a colorimetric assay kit (MilliporeSigma, AST Activity Assay Kit, Cat. No.

MAK055), following the manufacturers' protocols. Absorbance was read using a microplate

reader (SpectraMax i3, Molecular Devices).

Pharmacokinetics CalU19 variants were conjugated with Cy7 through random lysine labeling.

Labeled proteins were administered intravenously to CD-1 mice. At the indicated time points,

blood was collected from the tail vein using heparinized capillary tubes, diluted in PBS

containing EDTA, and centrifuged to remove cells. Fluorescence in plasma supernatants was

quantified using a microplate reader to determine circulating protein concentrations. Percentage

of antidote protein remaining in circulation was calculated as the percentage of antidote at each

timepoint relative to the 1 min timepoint.

Statistical Analysis and Schematic Representation

Statistical analyses was performed with GraphPad Prism. Data are plotted as mean ± SD.

Comparisons among different treatment groups are based on one-way ANOVA with Tukey *post*

hoc tests. A *P* value < 0.05 was considered statistical significance. Parts of the schematics in

this publication were created with BioRender.

REFERENCES

1.   M. Naghavi *et al.*, *The Lancet*. **404**, 1199–1226 (2024).

2.   C. Årdal *et al.*, *Nat Rev Microbiol*. **18**, 267–274 (2020).

3.   F. Ruggieri *et al.*, *European Journal of Medicinal Chemistry*. **256**, 115413 (2023).

4.   G. Liu *et al.*, *Nat Chem Biol*. **19**, 1342–1350 (2023).

5.   F. Wan, F. Wong, J. J. Collins, C. de la Fuente-Nunez, *Nat Rev Bioeng*. **2**, 392–407
(2024).

6.   C. D. Santos-Júnior *et al.*, *Cell*. **187**, 3761-3778.e16 (2024).

7.   F. Wong, C. de la Fuente-Nunez, J. J. Collins, *Science*. **381**, 164–170 (2023).

8.   N. K. Khandelwal *et al.*, *Nat Commun*. **16**, 3139 (2025).

9.   M. W. Tse *et al.*, *Proc Natl Acad Sci U S A*. **122**, e2402017122 (2025).

10.   M. D. T. Torres *et al.*, *Cell*. **187**, 5453-5467.e15 (2024).

11.   W. Liang, J. Diana, *EMBO Mol Med*. **14**, e15409 (2022).

12.   M. Chawla, J. Verma, R. Gupta, B. Das, *Front. Microbiol*. **13** (2022),
doi:10.3389/fmicb.2022.887251.

13.   B. Bognár, R. Spohn, V. Lázár, *npj Antimicrob Resist*. **2**, 1–8 (2024).

14.   Z. Si, K. Pethe, M. B. Chan-Park, *JACS Au*. **3**, 276–292 (2023).

15. M. A. Farha, E. D. Brown, *Nat Microbiol*. **4**, 565–577 (2019).

16. E. Mitchell *et al.*, *Nat Genet*. **57**, 1684–1694 (2025).

17. G. Cavaletti, *Nat Rev Cancer*. **8**, 72–72 (2008).

18. M. E. McNerney, L. A. Godley, M. M. Le Beau, *Nat Rev Cancer*. **17**, 513–527 (2017).

19. F. J. Giles *et al.*, *Cancer*. **92**, 406–413 (2001).

20. M. Guffroy *et al.*, *Clin Cancer Res*. **23**, 1760–1770 (2017).

21. J. B. Biggins, K. C. Onwueme, J. S. Thorson, *Science*. **301**, 1537–1541 (2003).

22. S. I. Elshahawi *et al.*, *ACS Chem Biol*. **9**, 2347–2358 (2014).

23. WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health
importance to guide research, development and strategies to prevent and control
antimicrobial resistance, (available at
https://www.who.int/publications/i/item/9789240093461).

24. E. L. Sievers *et al.*, *Blood*. **93**, 3678–3684 (1999).

25. E. L. Sievers *et al.*, *J Clin Oncol*. **19**, 3244–3254 (2001).

26. C. Ngambenjawong, L. W. Chan, H. E. Fleming, S. N. Bhatia, *ACS Nano*. **16**, 15779–
15791 (2022).

27. C. Ngambenjawong *et al.*, *ACS Nano*. **19**, 9958–9970 (2025).

28. L. W. Chan *et al.*, *Nat. Nanotechnol.* **15**, 792–800 (2020).

29. K. Rakhra *et al.*, *Science Immunology*. **6**, eabd8003 (2021).

30. S. M. Lehar *et al.*, *Nature*. **527**, 323–328 (2015).

31. K. K. Kajihara *et al.*, *mBio*. **12**, e00202-21.

32. J. D. Bargh, A. Isidro-Llobet, J. S. Parker, D. R. Spring, *Chem. Soc. Rev.* **48**, 4361–4374
(2019).

33. K. Liu *et al.*, *Molecular Cancer*. **23**, 62 (2024).

34. R. Colombo, J. R. Rich, *Cancer Cell* (2022), doi:10.1016/j.ccell.2022.09.016.

35. H.-P. Gerber, S. Gangwar, A. Betts, *MAbs*. **15**, 2230618.

36. J. E. Fisher, *Antibodies (Basel)*. **10**, 15 (2021).

37. J. Joseph, S. Boby, S. Mooyottu, M. S. Muyyarikkandy, *npj Antimicrob Resist*. **3**, 53 (2025).

38. C. Reading, M. Cole, *Antimicrob Agents Chemother*. **11**, 852–857 (1977).

39.   J. M. Stokes *et al.*, *Nat Microbiol*. **2**, 17028 (2017).

40.   N. Blondiaux *et al.*, *Science*. **355**, 1206–1211 (2017).

41.   A. R. Kirtane *et al.*, *Nat. Biomed. Eng.* **9**, 1240–1253 (2025).

42.   K. Mestermann *et al.*, *Science Translational Medicine*. **11**, eaau5907 (2019).

43.   J. M. Harris, R. B. Chess, *Nat Rev Drug Discov*. **2**, 214–221 (2003).

44.   P. J. Carter, A. Rajpal, *Cell*. **185**, 2789–2805 (2022).

45.   L. V. Zinsli, N. Stierlin, M. J. Loessner, M. Schmelcher, *Comput Struct Biotechnol J*. **19**,
580   315–329 (2021).

581   46.   A. E. Place *et al.*, *Lancet Oncol*. **16**, 1677–1690 (2015).

582   47.   L. C. Hurst *et al.*, *New England Journal of Medicine*. **361**, 968–979 (2009).

583   48.   N. K. Prasad, I. B. Seiple, R. T. Cirz, O. S. Rosenberg, *Antimicrobial Agents and*
584   *Chemotherapy*. **66**, e00054-22 (2022).

585   49.   L. J. V. Piddock, R. Malpani, A. Hennessy, *ACS Infect. Dis.* **10**, 2445–2447 (2024).

586   50.   U. Theuretzbacher, K. Outterson, A. Engel, A. Karlén, *Nat Rev Microbiol*. **18**, 275–285
587   (2020).

588   51.   L. L. Silver, *Clin Microbiol Rev*. **24**, 71–109 (2011).

589   52.   J. de la Cuesta-Zuluaga, L. Boldt, L. Maier, *Cell Host & Microbe*. **32**, 786–793 (2024).

590   53.   B. P. Willing, S. L. Russell, B. B. Finlay, *Nat Rev Microbiol*. **9**, 233–243 (2011).

591   54.   D. G. J. Larsson, C.-F. Flach, *Nat Rev Microbiol*. **20**, 257–269 (2022).

592   55.   Fate and effects of cytostatic pharmaceuticals in the environment and the identification of
593   biomarkers for and improved risk assessment on environmental exposure | FP7. *CORDIS |*
594   *European Commission*, (available at https://cordis.europa.eu/project/id/265264/reporting).

595   56.   A. Castellano-Hinojosa, M. J. Gallardo-Altamirano, J. González-López, A. González-
596   Martínez, *J Hazard Mater*. **447**, 130818 (2023).

597   57.   S. Mukherjee, D. Mehta, K. Dhangar, M. Kumar, *Chemical Engineering Journal*. **407**,
598   127184 (2021).

FIGURES AND TABLES

Table 1

	Calicheamicin γ1
Pseudomonas aeruginosa PAO1	0.11 µg/mL (78.1 nM)
Escherichia coli KS1	0.013 µg/mL (9.23 nM)
Klebsiella pneumoniae ATCC43816	0.05 µg/mL (312.5 nM)
Staphylococcus aureus 502A*	0.00061 µg/mL (835 pM)
BA1566 (MRSA)	0.00061 µg/mL (835 pM)

*For comparison, MIC of vancomycin for SA 502A was measured

as 1 µg/mL

Table 1 Antibacterial activity of calicheamicin against representative Gram-negative and Gram-positive pathogens.
Minimum inhibitory concentrations (MICs) were determined by microdilution assay in Mueller–Hinton medium supplemented with 3% DMSO. Data represent biological triplicates.

Figure 1. Dual strategy to expand the therapeutic index of toxic compounds for antimicrobial use. (Top) A narrow therapeutic index, defined as the difference between the median therapeutic effect and the median toxic effect, constrains the utility of cytotoxins (blue star) as antimicrobials. (Middle) Formulation of cytotoxins as conditionally active conjugates can widen the therapeutic index by restricting drug activity to an infection site, where proteases (red pac-man) release drug from the conjugate. (Bottom) "Antidotes" derived from self-resistance enzymes further widen the therapeutic index by neutralization off-target drug.

Figure 2. ABD-calicheamicin conjugate demonstrates masked mammalian toxicity and protease-triggered antibacterial activity *in vitro* and improved mammalian toxicity *in vivo.* (A) Schematic of ABD-calicheamicin conjugate (ABD-Cal) activation within the infection microenvironment. Within the infection environment, 1) neutrophils are recruited and 2) secrete neutrophil elastase which 3) cleaves and releases calicheamicin from the conjugate 4) such that it can kill bacteria. **(B-C)** Antimicrobial activity of ABD-Cal evaluated via microdilution assays against representative **(B)** Gram-positive and **(C)** Gram-negative bacterial strains in the presence or absence of recombinant neutrophil elastase. Bacterial growth measured by absorbance at 600 nm and normalized to untreated controls. Fold change in antimicrobial activity calculated as the ratio of MIC values of intact vs. cleaved conjugates. (mean ± sd, n = 3 biological replicates) **(D)** Masked mammalian toxicity of ABD-Cal evaluated via MTS cell proliferation assay in HepG2 cells. Cell viability measured by absorbance at 490 nm and normalized to untreated controls. Fold change in cytotoxicity calculated as the ratio of IC_{50} values of intact vs. cleaved conjugates. (mean ± sd, n = 3 biological replicates) **(E)** Schematic of experimental timeline of mice administered free calicheamicin (0.025 mg/kg) compared with ABD-Cal (0.025 mg/kg drug eq.) **(F)** Serum AST and **(G)** ALT. Dashed lines indicate normal healthy range. (n = 5 biological replicates. Statistical significance was determined by one-way ANOVA followed by Tukey's post hoc test). *P < 0.001 (***)*; *P < 0.0001 (****)*

Figure 3. Microbial-inspired antidotes protect against calicheamicin-mediated liver toxicity *in vitro* and *in vivo.* Schematic illustrating neutralization of calicheamicin by engineered microbial self-resistance enzymes ("antidotes"). After neutrophil-elastase–mediated cleavage of the ABD–calicheamicin conjugate, either on-target (within the infection microenvironment) or off-target (within healthy tissue) released calicheamicin can 1) enter circulation, where it is 2) bound and inactivated by LC-CalU19x, such that exposure of healthy tissue to active drug is reduced. (B) Calicheamicin neutralization by engineered variants of CalU19 (evaluated via cell proliferation assay in HepG2 cells. Cell viability measured by absorbance at 490 nm and normalized to untreated controls. mean ± sd, n = 4 biological replicates. (C) Schematic of timeline during which Cy7-conjugated proteins were assayed for clearance from plasma. (D) Pharmacokinetics of antidote variants following systemic injection, with plasma concentrations of Cy7 quantified over time. (mean ± sd, n = 4–5 mice per group) (E) Schematic of *in vivo* neutralization of free calicheamicin by LC-CalU19x. (F) Serum AST and (G) ALT levels measured 36 h after calicheamicin administration with or without antidote treatment. Dashed lines indicate normal healthy range (n = 5 Statistical significance determined by one-way ANOVA followed by Tukey's post hoc test.) *P* < 0.0001 (****); ns: not significant.

Fig. 4 Therapeutic efficacy of calicheamicin conjugate and neutralizing antidote in a murine model of bacterial pneumonia. (A) Schematic of experimental timeline. (B) Bacterial burden in lungs 6 h post-treatment in mice infected intratracheally with *Staphylococcus aureus* 502A and treated intratracheally with vehicle or conjugated calicheamicin (ABD-Cal) and then intravenously with vehicle or antidote (LC-CalU19x). Each point represents an individual animal (n = 5 per group) Statistical significance was determined by unpaired two-tailed *t*-test. (C) Serum levels of AST and (D) ALT measured 36-48 h after infection and treatment with ABD-Cal, in the presence or absence of antidote. Each point represents an individual animal (n = 4–5 per group). Dashed lines indicate the normal healthy range. Statistical significance was determined by unpaired t-test. *P* < 0.0001 (****)

SUPPLEMENTARY FIGURES

Figure S1. H&E-stained liver sections from mice receiving calicheamicin vs. ABD–Cal

Figure S2. CalU19 neutralizes free calicheamicin but not ABD-Cal

Figure S3. H&E-stained liver sections from mice receiving calicheamicin with or without LC-CalU19x.