The Phosphate-Specific Transport System Gene *pstA1* Contributes to Rifampin

Tolerance in *Mycobacterium tuberculosis*

<u>Authors:</u>

Carina Danchik[1, †], G. V. Anoushka Chinmayi[1, †], Gi Yong Lee[2], Joel S. Bader[3,4],

Hyungjin Eoh[2,5], Petros C. Karakousis[1,6,7,*]

† These authors contributed equally to this work.

<u>Authors' affiliations:</u>

[1] Tuberculosis Research Advancement Center, Department of Medicine, Johns Hopkins

School of Medicine, Baltimore, MD, USA

[2] Department of Immunology and Immune Therapeutics, Keck School of Medicine,

University of Southern California, Los Angeles, CA, USA

[3] Institute for Computational Medicine, Johns Hopkins University, Baltimore, MD, USA

[4] Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD, USA

[5] Department of Ophthalmology, Keck School of Medicine, University of Southern

California, Los Angeles, CA, USA

[6] Department of International Health, Johns Hopkins Bloomberg School of Public Health,

Baltimore, MD, USA

[7] Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg

School of Public Health, Baltimore, MD, USA

* Corresponding Author: Petros C. Karakousis, Center for Tuberculosis Research,

Division of Infectious Diseases, School of Medicine, 1550 Orleans Street, Room 110,

Baltimore, MD 21287, (410) 502-8233 / FAX (410) 614-8173, Email: <u>petros@jhmi.edu</u>

Keywords: *Mycobacterium tuberculosis*, PstA1, rifampin, antibiotic tolerance,

phosphate-specific transport

Abstract

Tuberculosis (TB) caused an estimated 10.8 million new cases and 1.25 million deaths in 2023. Antibiotic tolerance, the ability of bacteria to survive exposure to bactericidal antibiotics without genetic resistance mutations, contributes to the need for prolonged treatment. Targeting antibiotic tolerance mechanisms could promote accelerated clearance of *Mycobacterium tuberculosis* (*Mtb*), thereby improving medical adherence. Our previous forward genetic screen for rifampin tolerance genes identified *pstA1*, which is involved in phosphate-specific import. The ABC-type transporter permease PstA1 has previously been implicated in *Mtb* virulence, as mutants lacking this gene exhibit defective survival during phosphate limitation *in vitro*, in infected macrophages, and in immunocompetent mice. The minimum inhibitory concentration (MIC) of rifampin was not altered in a *pstA1* deletion mutant (Δ*pstA1*), suggesting this difference in susceptibility is not due to antibiotic resistance. Consistent with a role in rifampin tolerance, time-kill assays revealed a shift in the mean duration of killing (2-log reduction) from 1.6 days in wild-type to 1.0 day in Δ*pstA1*, and complementation partially restored this phenotype. We found that *pstA1* is specifically required for *Mtb* survival in the absence of exogenous inorganic phosphate and is important for adaptation to growth in culture without detergent, and within macrophages in an interferon-γ-dependent manner. Differential expression analysis revealed that Δ*pstA1* exhibited substantial transcriptional reprogramming with 58 differentially expressed genes, including altered expression of metabolic, DNA damage repair, and secretory pathways. PstA1 represents a novel drug target, and inhibitors could serve as adjunctive therapies to shorten treatment times, reducing opportunities for drug resistance emergence.

Importance

Tuberculosis remains one of the world's deadliest infectious diseases, causing over a million deaths annually. Current treatment requires months of antibiotic therapy, and poor adherence to these lengthy regimens contributes to the emergence of drug-resistant strains that are increasingly difficult to treat. A major barrier to shorter treatment is antibiotic tolerance, which allows bacteria to survive drug exposure without genetic resistance mutations. This study identifies the phosphate transport component PstA1 as a critical factor enabling *Mycobacterium tuberculosis* to tolerate the frontline antibiotic rifampin. Bacteria lacking *pstA1* are eliminated more rapidly by rifampin, demonstrating that this transporter actively promotes bacterial survival during treatment. These findings suggest that drugs targeting PstA1 could be combined with standard antibiotics to accelerate bacterial clearance, potentially shortening treatment duration and improving patient adherence. Such adjunctive therapies targeting tolerance mechanisms represent a promising strategy to combat tuberculosis and reduce the global burden of drug-resistant disease.

Introduction

Tuberculosis (TB) remains the leading cause of death from a single infectious agent, causing an estimated 10.8 million new cases and 1.25 million deaths in 2023 alone [1]. While effective regimens exist and three new drugs have been approved to treat TB since 2010, additional advances are needed to shorten treatment duration and improve treatment completion [2]. Current first-line TB treatment requires the use of multiple antibiotics for a minimum of six months to prevent relapse. A key reason for this lengthy treatment time is the presence of residual populations of bacteria that exhibit antibiotic tolerance– a phenotypic and reversible state in which bacteria become transiently insusceptible to antibiotics, requiring longer exposures to achieve bacterial eradication [3]. Bacteria can use a variety of adaptive mechanisms, collectively termed phenotypic tolerance, to adapt to and survive under stress conditions experienced within the host as well as antibiotic pressure [4]. The molecular mechanisms underlying antibiotic tolerance in *Mtb* are complex, involving transcriptional stress responses, metabolic adaptations, activation of drug efflux pumps, and changes in cell wall permeability [5]. Direct disruption

of *Mtb* antibiotic tolerance mechanisms is expected to result in more rapid bacillary killing,

leading to shorter treatment times that may improve medical adherence and decrease

opportunities for the development of antibiotic resistance [6].

We previously used transposon mutant sequencing (Tn-seq) to identify *Mtb* genes

required for survival during exposure to bactericidal concentrations of the first-line drug

rifampin. Among the top hits from this unbiased genome-wide approach were the

phosphate-specific transport (Pst) genes *rv0820/phoT*, *rv0929/pstC2*, and *rv0930/pstA1*

[7]. Notably, their homologs were also identified in our similar forward genetic screen in

which *M. avium* was exposed to rifabutin, providing additional confidence in their

significance and suggesting that Pst genes contribute to rifamycin tolerance across

mycobacterial species [6]. Pst systems are comprised of four essential components: PstA

and PstC are permeases that form a transmembrane channel complex; PstS is an

extracellular protein that senses and captures phosphate from the environment; and PstB

is an intracellular ATPase that provides energy for active transport of phosphate against

its concentration gradient [8-10]. PstA1, which is co-expressed as part of the putative *Mtb*

operon *pstS3-pstC2-pstA1* [11], is an important virulence factor required for *Mtb* survival

during acidic stress [12] and phosphate depletion [13, 14], as well as for *Mtb* survival

within macrophages [15] and for resistance to IFN-γ dependent immunity in mice [14].

PstA1 also plays important regulatory roles, as several substrates of the ESX-5 secretion

system, as well as membrane vesicles, are hypersecreted in Δ*pstA1* mutants [16, 17].

Each of these phenotypes is mediated at least in part through the constitutive activation

of RegX3, a response regulator of the PhoPR two-component system, in the absence of

PstA1 [18, 19]. This regulatory connection links phosphate sensing to broader stress

responses and virulence factor expression. Previous studies have shown that a *pstA1*

deletion mutant was more susceptible to several drug combinations *in vitro* and to

rifampin in a mouse model [20], providing initial evidence for a role in antibiotic tolerance.

However, the molecular mechanisms underlying this phenotype have not been fully

elucidated.

In this study, we have comprehensively characterized the role of *pstA1* in rifampin

tolerance using phenotypic assays, transcriptomics, and metabolomics to explore the

underlying molecular mechanisms of its increased susceptibility. We have also studied the role of PstA1 in *Mtb* growth and survival in phosphate-rich and phosphate-depleted models of nutrient starvation and within macrophages, providing insights into how phosphate transport contributes to bacterial adaptation under diverse stress conditions. Our findings reveal new mechanistic insights into antibiotic tolerance and highlight the potential for targeting phosphate transport pathways in TB treatment.

Results

Phosphate transport gene deletions confer rifampin hypersusceptibility

Our Tn-seq screen to identify *Mtb* genes required for tolerance to the sterilizing drug rifampin identified three Pst genes, *rv0820/phoT*, *rv0929/pstC2*, and *rv0930/pstA1*, among the top hits [7]. Their homologs were also found in a similar Tn-seq study of *M. avium* exposed to rifabutin [6]. Notably, *pstC2* and *pstA1* are downstream components of an *Mtb* operon including *pstS3* [22, 37]. To confirm that these three genes are co-transcribed in the H37Rv strain under our experimental conditions, we performed RT-PCR analysis using primers spanning the intergenic and intragenic regions of this operon. Our results confirmed that *pstS3*, *pstC2*, and *pstA1* are co-expressed as a single polycistronic mRNA transcript (Fig 1), validating the operonic structure for subsequent studies.

Next, we generated individual targeted deletion strains for each of these three genes using the ORBIT cloning system [21]. Cloning was validated via PCR of the 5' and 3' ends of the insertion site, as well as with whole genome sequencing for Δ*pstA1* (SeqCenter, Pittsburgh, PA) (Supplementary Figure S1). To validate our Tn-seq findings, we assessed the activity of rifampin against each deletion strain using standard time-kill assays following addition of 2.0 µg/ml rifampin. All three deletion strains showed increased susceptibility to rifampin treatment compared to wildtype controls harboring empty vectors. At day 6, Δ*phoT* showed a non-significant 0.9 $\log_{10}$ reduction in CFU compared to control (adjusted P = 0.358; Fig. 2A). Δ*pstC2* demonstrated a more pronounced phenotype with an additional 1.3 $\log_{10}$ reduction (adjusted P = 0.000189, Fig. 2B). Δ*pstA1*

exhibited the strongest susceptibility to rifampin, showing an additional 1.9 $\log_{10}$ reduction
in viable bacteria (adjusted P = 0.00742) relative to the isogenic wild-type control (Fig.
2C). The minimum duration of killing (MDK), defined as the time required to achieve a 2-
$\log_{10}$ reduction from the starting bacterial density, decreased by 0.4 days for Δ*phoT*, 1.1
days for Δ*pstC2*, and 0.6 days for Δ*pstA1* relative to controls. Consistent with a role for
each of these genes in *Mtb* tolerance, rather than resistance, to rifampin, each deletion
mutant showed minimal alterations in rifampin MIC values. The MIC values of rifampin
against Δ*phoT* and Δ*pstC2* were 0.03 µg/ml and that of Δ*pstA1* was 0.016 µg/ml (wild
type, 0.03-0.06 µg/ml).

Complementation of *pstA1* partially restores wild-type rifampin tolerance

Given the robust phenotype observed with Δ*pstA1*, and the possibility that deletion of
pstC2 could have polar effects on *pstA1*, our subsequent studies focused on Δ*pstA1*. To
validate the *pstA1* deletion phenotype, we constructed a genetic complementation strain
(Δ*pstA1::pstA1*). Since *pstS3*, *pstC2*, and *pstA1* are co-transcribed as an operon, we
designed our complementation construct to include appropriate regulatory elements. The
pstA1 coding sequence fused to the putative operonic promoter region (422 bp upstream
of *pstS3* plus the first 75 bp of *pstS3*) was cloned into the episomal mycobacterial vector
pMH94 and introduced into the Δ*pstA1* strain. Complementation was confirmed by PCR
of insert junctions and whole genome sequencing (Supplementary Figure S1). Time-kill
assays revealed that the minimum duration of killing (MDK) of rifampin for the
complement strain was 1.3 days, as compared to 1.6 days for the wild type and 1.0 day
for ΔpstA1 (Fig. 2D). MDK values differed significantly between all three strains (WT vs
ΔpstA1: p = 0.00004; WT vs pstA1 comp: p = 0.004; ΔpstA1 vs pstA1 comp: p = 0.0011,
one-way ANOVA with Tukey's multiple comparisons test). Thus, genetic complementation
of Δ*pstA1* partially restored wild-type rifampin tolerance.

Δ*pstA1* exhibits phosphate-specific survival defects while maintaining rifampin
susceptibility under starvation conditions

To characterize the broader physiological role of *pstA1*, we examined bacterial survival
under various stress conditions relevant to mycobacterial pathogenesis and persistence.
As multiple forms of nutrient starvation are known to induce a tolerance phenotype [13,
22, 23], we tested the ability of Δ*pstA1* to survive in both the PBS nutrient starvation model
used in our prior screen [7] and during inorganic phosphate limitation [13].

During nutrient starvation in phosphate-buffered saline containing the detergent tween-
80, both wild-type and Δ*pstA1* strains showed equivalent survival over a 14-day
adaptation period, with no significant differences observed between strains (Fig. 3A).
ΔpstA1 retained enhanced rifampin susceptibility under nutrient-starved conditions.
When rifampin (2.0 μg/ml) was added after the 14-day starvation period, the deletion
mutant showed significantly enhanced killing compared to wild-type, with additional
reductions in viability becoming apparent by day 2 (1.4 $\log_{10}$ reduction; adjusted P =
0.000184) and persisting through day 6 (1.6 $\log_{10}$ reduction; adjusted P = 0.000184) (Fig.
3B).

Since phosphate depletion is known to induce the stringent response, a key antibiotic
tolerance pathway in *Mtb* [24, 25], and the *pstS3-pstC2-pstA1* operon and *phoT* are
upregulated under low phosphate conditions [13], we tested the ability of Δ*pstA1* to
survive in the absence of exogenous inorganic phosphate. Phosphate-specific depletion
revealed a clear survival disadvantage for Δ*pstA1*. While both wild-type and Δ*pstA1*
strains exhibited similar growth and survival characteristics in standard phosphate-rich
7H9 broth, Δ*pstA1* showed significantly reduced viability under phosphate-depleted
conditions. In modified 7H9 broth lacking inorganic phosphate, the deletion strain
demonstrated an additional survival defect with a 0.7 $\log_{10}$/ml decrease by day 7 (adjusted
P = 0.0115) and a 0.6 $\log_{10}$/ml decrease by day 14 (adjusted P = 0.00181) relative to wild
type (Fig. 3C). These results confirm the role of *pstA1* in phosphate acquisition under
limiting conditions and demonstrate its requirement for growth in low-phosphate
environments.

Δ*pstA1* exhibits survival defects in IFN-γ-activated macrophages

To assess the relevance of our findings to mycobacterial pathogenesis, we next examined the role of *pstA1* in macrophage infection and intracellular survival as a model of the natural host environment. Macrophages represent the primary host cell niche for *Mtb* and create a hostile intracellular environment characterized by nutrient limitation, acidic pH, and antimicrobial factor exposure [15]. We infected PMA-differentiated THP-1 macrophages with wild-type and Δ*pstA1* strains at a multiplicity of infection (MOI) of 1:1 and monitored intracellular bacterial survival over 4 days. In resting macrophages, there was no significant difference in bacterial burden between wild-type and Δ*pstA1* at any time point measured (days 0, 2, and 4 post-infection; Fig. 4A), suggesting that *pstA1* is not required for initial invasion or survival in resting macrophages. However, when macrophages were pre-activated with IFN-γ, a key cytokine that enhances macrophage antimicrobial functions, Δ*pstA1* showed a significant survival defect that became pronounced by day 4 post-infection. While wild-type bacteria maintained relatively high intracellular numbers ($4.2×10^6$ CFU/ml), the Δ*pstA1* strain showed reduced survival ($2.43×10^6$ CFU/ml), representing a statistically significant difference (adjusted P = 0.0098; Fig. 4B). This result confirms previous reports that *pstA1* is specifically required for resistance to IFN-γ-mediated host immunity and suggests a role in counteracting activated macrophage antimicrobial mechanisms [15].

The contribution of PstA1 to *Mtb* rifampin tolerance is not mediated through the accumulation of inorganic polyphosphate

Having established that Δ*pstA1* exhibits enhanced rifampin susceptibility, we next sought to investigate the underlying mechanisms responsible for this phenotype. Induction of the *Mtb* stringent response and accumulation of the multifunctional stress signaling molecule, inorganic polyphosphate (polyP), has been linked to antibiotic tolerance in *Mtb* [26-30]. Since RegX3, a two-component response regulator that controls the *Mtb* phosphate starvation and stringent responses [13, 31], is constitutively activated upon *pstA1* deletion [14], we measured intracellular polyP levels as a proxy for stringent response activation. No significant differences in polyP accumulation were observed between wild-type and

227 Δ*pstA1* strains across different growth phases (Fig. 5A). Wild-type polyP content ranged

228 from 3.072 to 5.536 ng/µg total protein, while Δ*pstA1* levels ranged from 3.673 to 5.915

229 ng/µg total protein.

Δ*pstA1* exhibits increased rifampin uptake

We hypothesized that *pstA1* deficiency might be associated with increased intrabacterial

accumulation of rifampin. Using a previously validated device that enables timed start-

stop measurements of antibiotics remaining in spent media [32], we quantified rifampin

content in the wild-type and mutant strains The kinetics of rifampin uptake revealed that

Δ*pstA1* displayed significantly faster and greater rifampin accumulation compared to wild-

type at 6 hours post-exposure, suggesting increased uptake and/or decreased efflux of

rifampin (Fig. 5B-C).

Cell morphology analysis reveals minimal structural differences

To determine whether the observed increase in rifampin uptake was attributable to gross

alterations in cell wall structure of the mutant, we examined bacterial morphology using

electron microscopy. Under standard growth conditions including Tween-80 to prevent

bacterial clumping, we observed no significant difference in cell length between the two

strains (average bacterial length= 2.34 µm for wild type and 2.26 µm for Δ*pstA1;* Fig. 5D).

Likewise, the strains showed similar average cell wall thicknesses of 21.18 nm and 19.61

nm, respectively (Fig. 5E). Substantial variations in cell wall thickness were observed for

both wild-type and Δ*pstA1* which appear similar to previously reported lipid inclusions

[33]. There were no significant differences in minimum, maximum, or average cell wall

thickness between the two strains.

Since use of detergent can alter cell wall permeability [34], the experiment was repeated

using cultures grown without Tween-80 (Fig. 5F-G). Cell length decreased for both strains

in detergent-free broth, from an average of 2.34 µm to 1.64 µm for wild-type and from

2.26 µm to 1.95 µm for Δ*pstA1* (difference= 0.31 µm; P = 0.0329; Fig. 5F). Cell wall

thickness did not vary notably between strains with averages of 15.37 nm for wild type

and 16.06 nm for Δ*pstA1*. The large variations in cell wall thickness seen in cultures grown

with Tween-80 were not present under detergent-free conditions (Fig. 5H). These results
demonstrate that Δ*pstA1* exhibits increased rifampin accumulation without major changes
in cellular morphology.

Transcriptomic analysis reveals extensive metabolic and regulatory
reprogramming in Δ*pstA1*

To further explore the molecular mechanisms underlying the increased rifampin
susceptibility of Δ*pstA1* and identify potential cellular pathways contributing to enhanced
drug uptake, we performed transcriptomic analysis using RNA sequencing. Wild-type and
Δ*pstA1* strains were exposed to a sub-inhibitory concentration of rifampin (7.5 ng/ml,
representing 1/4 MIC) for 30 or 60 minutes. This concentration and timing were selected
to capture early transcriptional responses while minimizing the global transcriptional
inhibition that occurs at bactericidal concentrations of rifampin. During axenic growth in
nutrient-rich broth lacking rifampin, Δ*pstA1* exhibited substantial transcriptional
reprogramming compared to wild type, with 58 differentially expressed genes (DEGs)
identified: 49 upregulated and 9 downregulated (adjusted $P < 0.05$, |$\log_2$ fold change| >
1; Fig. 6A). Several genes known to be part of the phosphate regulon showed significant
upregulation, including *pstS3* ($\log_2$ fold change = 3.22), *pstC2* ($\log_2$ fold change = 2.50),
and the two-component regulatory system genes *senX3* ($\log_2$ fold change = 1.59) and
regX3 ($\log_2$ fold change = 1.77), indicating constitutive activation of phosphate starvation
signaling.

KEGG pathway enrichment analysis revealed that genes involved in biosynthesis and
metabolic pathways were prominently represented among the DEGs (Fig. 6B). The
differentially expressed genes included fourteen PE/PPE family genes (9 upregulated, 5
downregulated), with upregulation of members of the ESX-5 locus and downregulation of
pe31 ($\log_2$ fold change = -6.02) and *ppe60* ($\log_2$ fold change = -3.37). Genes encoding
four ESAT-6-like proteins were upregulated, namely *esxI* ($\log_2$ fold change = 3.65), *esxJ*
($\log_2$ fold change = 2.59), esxV ($\log_2$ fold change = 3.14), and *esxW* ($\log_2$ fold change =
2.35). The redox-responsive transcriptional regulator *whiB3* ($\log_2$ fold change = 2.60), a
key regulator of thiol metabolism and oxidative stress responses, was also significantly

upregulated [12]. Additionally, eleven genes involved in DNA metabolism were upregulated in the mutant at baseline, including resolvases, transposases, and DNA helicases. Rifampin treatment led to a progressive reduction in the number of DEGs between strains. After 30 minutes of rifampin exposure, 41 genes remained differentially expressed between strains. After 60 minutes, 41 genes were differentially expressed, with 36 genes showing consistent differential expression across all three conditions (baseline, 30 min, and 60 min). A comprehensive Venn diagram analysis of DEG overlap across all treatment conditions is shown in Supplementary Figure S2. The DNA metabolism genes were no longer significantly differentially expressed following rifampin treatment, while the core set of 36 genes included the phosphate regulon genes, all four ESAT-6-like protein genes, twelve PE/PPE family genes, and *whiB3*.

RT-qPCR analysis of 11 selected genes confirmed the expression patterns observed in the RNA-seq dataset, with correlation coefficients exceeding 0.7 for all comparisons (Fig. 6D; Supplementary Figure S3). Validated genes included members of the phosphate starvation response (*pstC2*, $\log_2$ fold change = 2.50; *regX3*, $\log_2$ fold change = 1.77; *senX3*, $\log_2$ fold change = 1.59), metabolic enzymes (*udgA*, $\log_2$ fold change = 3.08; *mgtA*, $\log_2$ fold change = 3.04), the stress response regulator *whiB3* ($\log_2$ fold change = 2.60), and PE/PPE family protein genes (*ppe60*, $\log_2$ fold change = -3.37; *ppe65*, $\log_2$ fold change = 1.89). A complete list of all DEGs across all experimental conditions is provided in Supplementary Table S2.

Metabolomic analysis reveals altered antioxidant metabolites in Δ*pstA1*

To understand the biochemical consequences of increased rifampin accumulation in the deletion mutant, we performed targeted metabolomics analysis of rifampin-treated bacteria and analysis was focused on antioxidant pathways. We analyzed the intracellular abundance of 16 metabolites involved in antioxidant biosynthesis and redox homeostasis in the Δ*pstA1* mutant, wild-type, and complement strains at 6 hours post-rifampin exposure. The heatmap analysis of all measured antioxidant metabolites revealed distinct metabolic profiles between strains (Fig. 7A). While most metabolites, including L-glutamate, glycine, NADPH, glutathionylspermidine, and various glutathione derivatives,

showed trends toward increased abundance in Δ*pstA1*, these changes did not reach
statistical significance. The complement strain generally displayed metabolite levels
which were intermediate between wild-type and the deletion mutant, consistent with
partial restoration of the metabolic phenotype.

Following rifampin treatment, two antioxidant pathway metabolites showed trends toward
altered abundance in Δ*pstA1* compared to wild-type: ergothioneine and 5-oxoproline ($p <$
0.1, ANOVA; Fig. 7B). Ergothioneine, a sulfur-containing antioxidant, showed a trend
toward elevation in Δ*pstA1* compared to both wild-type and complement strains. This
trend is notable given that *whiB3*, which was upregulated 2.6-fold in the transcriptomic
analysis and confirmed by RT-qPCR, is a metabolic regulator involved in the production
of ergothioneine and mycothiol, which play an important role in detoxifying the oxidative
stress [35]. Similarly, 5-oxoproline, a glutamate derived metabolite, showed a trend
toward increased abundance in the deletion mutant [36]. The observed trends suggest
that Δ*pstA1* may exhibit hypersensitivity to rifampin largely because of alterations in
antioxidant metabolism following increased rifampin exposure.

Discussion

Components of the Pst system have been implicated in virulence, stress tolerance, and
antibiotic susceptibility in several bacterial pathogens, suggesting that phosphate-specific
transport pathways may represent promising targets for therapeutic intervention [14, 37].
This study establishes PstA1 as a critical determinant of rifampin tolerance in *Mtb* through
mechanisms distinct from drug resistance. Deletion of *pstA1* resulted in enhanced
rifampin killing without substantial change in MIC, a phenotype that remained consistent
across nutrient starvation conditions and correlated with reduced survival in IFN-γ-
activated macrophages. Integrative multi-omics analysis revealed that *pstA1* loss triggers
constitutive phosphate starvation signaling, leading to PE/PPE porin upregulation. These
findings raise the possibility that PE/PPE porin upregulation enhances membrane
permeability, thereby promoting rifampin uptake. Metabolic profiling revealed
nonsignificant upregulation of antioxidant pathways including ergothioneine and

glutathione biosynthesis, suggesting elevated oxidative stress resulting from increased
intracellular rifampin accumulation due to hyperpermeability [35].

In contrast to prior studies showing that *pstA1* deficiency reduces the MIC of rifampin by
8-fold, we found a more modest 2-fold decrease in the rifampin MIC against Δ*pstA1*. This
discrepancy may be due to differences in the parent strain used (Erdman vs. H37Rv) [20].
Notwithstanding these findings, we observed significantly increased rifampin activity
against Δ*pstA1* in time-kill assays, the standard approach for assessing antibiotic
tolerance [5].

Δ*pstA1* retained its rifampin hypersusceptibility under nutrient starvation and showed
reduced survival under phosphate depletion, consistent with previous reports [14, 15] and
validating its core metabolic function in phosphate acquisition. This metabolic vulnerability
extended to host-pathogen interactions. While a previous Tn-seq study found that *pstS3*,
pstC2, *pstA1*, and *phoT* were each important for macrophage survival under all activation
conditions tested [15], Δ*pstA1* showed a more nuanced phenotype in the current study,
with no survival defect in resting macrophages but significant reduction in intracellular
survival in IFN-γ-activated macrophages by day 4. This finding aligns well with that from
a separate study reporting that Δ*pstA1* retained full virulence in IFN-γ$^{-/-}$ mice but showed
reduced chronic-phase survival in wild-type mice, establishing that PstA1 is specifically
required for surviving IFN-γ-mediated immunity [14].

To understand the mechanistic basis for enhanced rifampin susceptibility, we examined
drug uptake dynamics and cellular structure. We observed accelerated rifampin uptake
with significantly elevated drug accumulation by 6 hours, implicating increased drug
permeability and/or reduced efflux. Electron microscopy revealed no major structural
changes in cellular morphology. Our transcriptomic analysis provided insight into the
molecular basis for this altered permeability state. In the absence of rifampin exposure,
Δ*pstA1* showed extensive cellular reprogramming with 58 genes differentially expressed.
Most striking was the constitutive activation of the phosphate starvation regulon, including
pstS3, *pstC2*, *senX3*, and *regX3,* indicating that cells lacking functional PstA1 exist in a
state of perceived phosphate limitation regardless of environmental phosphate

availability. The marked upregulation of *pstS3* and *pstC2* likely represents compensatory

transcriptional upregulation consistent with constitutively activated RegX3, as multiple

studies have demonstrated RegX3-dependent regulation of the *pstS3-pstC2-pstA1*

operon [13, 14, 18]. Without the functional PstA1 permease, however, this transcriptional

response cannot restore phosphate transport, resulting in persistent phosphate starvation

signaling.

This baseline stress state likely compromises bacterial ability to respond effectively to

additional challenges like antibiotic exposure. There is substantial concordance between

our transcriptomic data and those of another study evaluating a *pstA1* deletion mutant in

the Erdman background, as 22 common DEGs were also identified in both studies,

including *senX3*, *regX3*, the *pstS3-pstC2-pstA1* operon, *whiB3*, and 9 PE/PPE genes,

validating the core transcriptional signature of *pstA1* deficiency across strain backgrounds

[14].

Among the 36 genes differentially expressed in *ΔpstA1* regardless of rifampin treatment,

12 encoded PE/PPE family genes and 4 encoded ESAT-6-like proteins. Four of these

PE/PPE genes (PE18, PPE19, PPE25, PPE27) are part of the *esx-5* locus, consistent

with established regulation of ESX-5 secretion by the Pst/SenX3-RegX3 axis and the

finding that *pstA1* deletion causes ESX-5 substrate hypersecretion. This upregulation

provides a direct mechanistic link to the increased rifampin uptake we observed [16].

Recent work demonstrated that *ΔpstA1* mutants overcome acid growth arrest by

upregulating PE19-PPE51 porin expression to enhance glycerol uptake [12], while

another study established that PE/PPE proteins function as substrate-specific outer

membrane channels [38]. The constitutive overexpression of multiple PE/PPE proteins in

ΔpstA1 facilitates not only glycerol uptake but also rifampin entry through increased

membrane permeability, potentially explaining the accelerated drug accumulation and

enhanced killing [12]. Transcriptomic analysis revealed no downregulation of known

rifampin efflux pump genes, further supporting the hypothesis that enhanced drug

accumulation results from increased membrane permeability rather than impaired efflux.

The absence of gross structural defects on electron microscopy supports the

interpretation that increased permeability results from specific changes in channel
composition rather than general membrane disruption.

Interestingly, DNA-interacting genes were only upregulated in Δ*pstA1* in the absence of
rifampin, suggesting that drug's inhibitory effects on RNA polymerase may suppress
some baseline transcriptional differences. We identified only a few genes differentially
regulated exclusively upon rifampin exposure, likely because of the relatively low drug
concentration and brief timepoints, which were selected to minimize global transcriptional
inhibition.

Our integrated metabolomic and transcriptomic analysis revealed activation of antioxidant
pathways in rifampin-treated Δ*pstA1*, consistent with elevated oxidative stress from
increased intracellular drug accumulation. We observed an upward trend of metabolites
involved in antioxidant defense, including 5-oxoproline, glutamate,
glutathionylspermidine, and notably ergothioneine. The ergothioneine response is
particularly informative: this mycobacterial antioxidant provides protection against
rifampin-induced damage, as a study demonstrated that Δ*egtD* mutants lacking
ergothioneine show enhanced susceptibility to rifampin and other first-line drugs [39]. The
apparent paradox in Δ*pstA1*, where ergothioneine is elevated yet susceptibility increases,
could reflect compensatory antioxidant upregulation that is insufficient to overcome the
primary defect of PE/PPE-mediated drug hyperpermeability.

The concurrent 2.6-fold upregulation of *whiB3*, a redox-responsive regulator of thiol
metabolism, further supports activation of oxidative stress defenses. While one study
showed that WhiB3 negatively regulates ergothioneine biosynthesis under standard
conditions, the simultaneous increase in both *whiB3* expression and ergothioneine levels
in rifampin-treated Δ*pstA1* suggests that rifampin-induced oxidative stress may trigger
compensatory antioxidant production [35]. The elevation of 5-oxoproline, a glutamate-
derived metabolite, indicates broad activation of thiol-based defenses beyond
ergothioneine alone [40]. Critically, despite these activated protective pathways, Δ*pstA1*
exhibits marked hypersusceptibility to rifampin, suggesting that the primary driver of
enhanced killing is increased drug accumulation rather than deficient antioxidant capacity.

This is further supported by recent findings showing that WhiB3 expression is induced by
both acidic pH and phosphate limitation through RegX3-dependent mechanisms, and that
elevated WhiB3 levels in Δ*pstA1* correlate with reduced acid-induced ROS and preserved
GAPDH activity [12].

While increased rifampin uptake provides mechanistic connection to enhanced killing, the
molecular basis for altered drug accumulation requires further investigation. The link
between PE/PPE upregulation, ESX-5 hypersecretion, and rifampin permeability needs
direct demonstration. The transcriptomic analysis captures a snapshot of cellular
responses and may miss dynamic changes occurring at other timepoints. The targeted
metabolomics, while revealing specific antioxidant pathway alterations, can be extended
to look at relevant perturbations in central metabolism or lipid composition that could
contribute to the phenotype.

The identification of *pstA1* as a tolerance determinant opens potential therapeutic
avenues, and targeting this pathway requires careful consideration. Our findings show
that loss of PstA1 function creates multiple vulnerabilities, including enhanced drug
uptake, metabolic dysregulation, and impaired stress responses, suggesting that
constitutive activation of phosphate-sensing pathways during antibiotic treatment
becomes a liability rather than an adaptive advantage. The specific mechanisms identified
are intriguing targets: the WhiB3-ergothioneine stress response axis and PE/PPE-
mediated membrane changes represent downstream effectors that could be modulated
without disrupting fundamental phosphate homeostasis. Understanding how phosphate
transport influences antibiotic tolerance could inform adjunctive strategies to eliminate
persister populations, potentially shortening the lengthy treatment regimens currently
required for TB cure.

Materials and Methods

Bacterial Strains and Plasmids

Mycobacterium tuberculosis H37Rv served as the wild-type strain for all experiments. Gene deletion strains were constructed using ORBIT (Oligonucleotide-Recombineering followed by Bxb1 Integrase Targeting) cloning as previously described [94]. Wild-type H37Rv was first transformed with pKM444, a non-integrating plasmid encoding RecT recombinase and Bxb1 integrase (gift from Kenan Murphy, University of Massachusetts). In a second electroporation, cells received pKM464 (also from Kenan Murphy) along with a gene-specific oligonucleotide designed to delete pstA1. This oligonucleotide contained the first and last 11 codons of pstA1 flanked by 37-bp homology arms, with the Bxb1 attP integration site positioned centrally. Transformants were selected on antibiotic-selective plates, and deletion junctions were verified by PCR amplification followed by Sanger sequencing.

For complementation, the pstA1 coding sequence plus its upstream regulatory region (422 bp upstream of pstS3 plus the first 75 bp of pstS3) was PCR-amplified from wild-type H37Rv genomic DNA. This fragment was cloned into pMH94, an episomal mycobacterial vector modified with apramycin resistance (gift from Dr. Gyanu Lamichhane, Johns Hopkins University), using In-Fusion Snap Assembly (Takara Bio). The vector was first linearized by restriction digestion with EcoRI and XbaI, gel-purified using QIAquick Gel Extraction Kit (Qiagen), then assembled with the PCR product. Cloned constructs were transformed into Stellar Competent Cells (Takara Bio), selected on LB plates with appropriate antibiotics, and verified by PCR before electroporation into the ΔpstA1 strain.

Culture Conditions and Media

M. tuberculosis strains were routinely cultured in Middlebrook 7H9 broth supplemented with 10% OADC enrichment, 0.2% glycerol, and 0.05% Tween-80, or on Middlebrook 7H11 agar plates incubated at 37°C. Phosphate-depleted medium was prepared as described previously [24] by omitting inorganic phosphate salts from the 7H9 formulation.

483   THP-1 monocytic cells were maintained in RPMI-1640 medium (Gibco) containing 2 mM

484   L-glutamine and 10% heat-inactivated fetal bovine serum at 37°C with 5% CO_2.

Operon Structure Validation

To confirm co-expression of *pstS3*, *pstC2*, and *pstA1* as a single operon, RNA was

extracted from logarithmically growing *Mtb* H37Rv cultures using TRIzol reagent.

Following DNase-I treatment to eliminate genomic DNA contamination, RNA was reverse

transcribed using SuperScript VILO cDNA Synthesis Kit (Invitrogen). PCR primers were

designed to amplify both intragenic regions (within individual genes) and intergenic

regions (spanning gene boundaries).

Whole Genome Sequencing Validation

High-molecular-weight genomic DNA was extracted from 10 mL log-phase cultures using

an extended protocol optimized for mycobacteria. Cultures were lysed by sequential

treatments with lysozyme (20 hours at 37°C), SDS/proteinase K (50°C), and CTAB,

followed by chloroform/isoamyl alcohol extraction and isopropanol precipitation. To verify

deletion and complementation, genomic DNA from wild type, Δ*pstA1*, and complement

strains were submitted to SeqCenter (Pittsburgh, PA) for whole genome sequencing.

Libraries were prepared using the Illumina DNA Prep kit with tagmentation-based

fragmentation and sequenced on an Illumina NovaSeq X Plus platform (2×151 bp paired-

end reads). Raw sequencing reads were aligned to the *M. tuberculosis* H37Rv reference

genome, and coverage analysis was performed using SAMtools v1.18. Deletion

boundaries were confirmed by examining read coverage across the *pstA1* locus.

Drug Susceptibility Testing

Minimum inhibitory concentrations (MICs) were determined by broth microdilution in 96-

well plates. Bacterial suspensions were prepared by washing log-phase cultures and

resuspending in Tween-80-free 7H9 medium to OD_{600} 0.01. Equal volumes (100 µL) of

bacterial suspension and 2× serial drug dilutions were combined and incubated at 37°C

for 6 days. Metabolic activity was assessed by adding 50 µL alamarBlue reagent (1:1

mixture of 10× alamarBlue and 10% Tween-80) and measuring fluorescence after 24

hours (544/590 nm excitation/emission) on a Fluostar OPTIMA plate reader (BMG

Labtech).

Transmission Electron Microscopy

Transmission electron microscopy was performed previously [27]. Briefly, mid-log phase

bacteria were collected and resuspended in 3% formaldehyde, 2.0% glutaraldehyde, 80

mM Sorenson's phosphate, and 4 mM MgCl2, pH 7.2 with duplicate samples for negative

staining and sectioning. Samples were fixed overnight at 4°C before processing and

imaging. Cell wall thickness was measured at two locations per bacterium: one in a thin

region and one in a thick region when present, to account for regional variations.

Time-Kill Assays

For nutrient-rich conditions, bacterial cultures were grown to mid-logarithmic phase in

complete 7H9 medium, then diluted to OD_{600} 0.05 one day prior to drug exposure. On

day 0, rifampin was added to a final concentration of 2 µg/mL unless otherwise noted

(stock: 2 mg/mL in DMSO), with vehicle controls receiving equivalent DMSO volumes.

Each condition was performed in triplicate using 5 mL cultures in 50 mL conical tubes,

incubated at 37°C with shaking. Bacterial viability was quantified by serial dilution plating

on 7H11 agar at days 2, 4, and 6 post-treatment. Plates were incubated at 37°C for 14-

21 days before colony counting.

For nutrient starvation conditions, log-phase cultures were washed and resuspended in

PBS containing 0.05% Tween-80 to OD_{600} 0.5, then incubated without shaking at 37°C

for 14 days to establish nutrient-deprivation. Rifampin (2 µg/mL) or vehicle control was

then added, and bacterial survival was monitored by CFU plating as described above.

For phosphate depletion experiments, cultures were washed twice and resuspended in

phosphate-free 7H9 medium to OD_{600} 0.05. Growth and survival were monitored over

time by CFU enumeration, with samples taken at indicated intervals during incubation at

37°C with shaking.

Macrophage Infection Assays

THP-1 cells were differentiated by treatment with 50 ng/mL phorbol 12-myristate 13-acetate (PMA) for 48 hours at 37°C with 5% CO_2. Differentiated macrophages were optionally activated with 20 ng/mL recombinant IFN-γ for 24 hours before infection. Bacterial suspensions were prepared from log-phase cultures and used to infect macrophages at a multiplicity of infection (MOI) of 1:1. After 4 hours, extracellular bacteria were removed by washing three times with PBS, and fresh medium was added. At specified time points, macrophages were lysed with 0.05% SDS and lysates were plated on 7H11 agar for CFU determination.

Polyphosphate Quantification

Polyphosphate levels were measured as described previously [41]. Bacterial pellets were lysed in guanidine thiocyanate buffer (5 M guanidine thiocyanate, 50 mM Tris-HCl pH 7.0) using bead beating (3 cycles, 20 seconds at 6,800 rpm). After centrifugation, supernatants were stored at -20°C, and protein concentrations were determined using DC Protein Assay (Bio-Rad) for normalization. Polyphosphate was isolated using the GeneClean III Kit (MP Biomedicals) following manufacturer protocols, including RNase/DNase treatment to remove nucleic acids. Polyphosphate concentrations were determined fluorometrically using DAPI staining (100 μg/mL final concentration) with excitation/emission at 415/525 nm.

Transcriptomic Analysis

Wild-type and Δ*pstA1* cultures were grown to mid-log phase (OD_{600} ~0.5) in 45 mL 7H9 medium. To capture early transcriptional responses, cultures were treated with subinhibitory rifampin (7.5 ng/mL, 1/4 MIC) or vehicle. RNA samples were collected at baseline, 30- or 60-minutes post-treatment with three independent biological replicates per condition.

Bacterial pellets (15 mL culture) were resuspended in TRIzol with 0.1 mm zirconia beads and lysed using a Precellys Evolution homogenizer (3 cycles, 30 seconds at 7,400 rpm).

RNA was extracted with chloroform, precipitated with ethanol, and purified on RNeasy
columns (Qiagen) with on-column DNase I digestion. Following polyA selection, strand-
specific libraries were prepared using NEBNext Ultra II Directional RNA Library Prep Kit
and sequenced on Illumina NovaSeq X Plus (2×150 bp paired-end) at the Johns Hopkins
Genetic Resources Core Facility.

Raw reads were quality-assessed (FastQC v0.12.1) [30] and aligned to the *M.*
tuberculosis H37Rv reference genome (Bowtie2 v2.5.1) [31]. Gene counts were
quantified with featureCounts (Subread v2.0.0) [32]. Differential expression analysis used
DESeq2 v1.44.0 with significance thresholds of adjusted p-value < 0.05 (Benjamini-
Hochberg) and $|log_2FC| > 1$. Pathway analysis was performed using KEGGREST to query
KEGG pathways [34] for significantly differentially expressed genes.

RT-qPCR Validation

Twelve genes were analyzed by RT-qPCR, including eleven target genes and sigA as a
housekeeping control (18-22 nt, 100-147 bp amplicons) designed with IDT PrimerQuest
(Supplementary Table S1). Total RNA (500 ng) was DNase-treated, purified on RNeasy
columns, and reverse transcribed with iScript RT Master Mix (Bio-Rad) alongside no-RT
controls. qPCR used 10 µL reactions with SYBR Green Supermix on a QuantStudio 3
system (95°C/10 min; 40 cycles: 95°C/15s, 60°C/1 min). Relative expression was
calculated using the ΔΔCt method with sigA normalization.

Metabolite extraction and LC-MS analysis

Mtb-laden filters were incubated on m7H10 plates at 37°C for 5 days to mid-log phase
[35]. Cells were quenched with prechilled acetonitrile:methanol:H_2O (40:40:20, v:v:v) at
−40°C. Metabolites were extracted through mechanical lysis using 0.1-mm zirconia beads
in a Precellys tissue homogenizer for 2 min at 6,000 rpm, repeated three times under
continuous cooling at ≤2°C. Cell lysates were clarified by filtration through 0.22-µm Spin-
X columns. Residual protein content was measured with a BCA protein assay kit (Thermo
Scientific) to normalize metabolite ion counts to cell biomass.

Extracted metabolites were separated using a Cogent Diamond Hydride Type C column
with mobile phases of solvent A (ddH$_2$O with 0.2% formic acid) and solvent B (acetonitrile
with 0.2% formic acid). An Agilent 6230 TOF mass spectrometer was coupled to an
Agilent 1290 LC system. Dynamic mass axis calibration was maintained via continuous
infusion of reference mass solution through an isocratic pump with a 100:1 splitter. This
setup achieved mass errors of ~5 ppm and mass resolution of 10,000-25,000 over m/z
range 62–966, with a dynamic range of 5 log$_{10}$. Ions were identified based on unique
accurate mass and retention time. Data analysis used Agilent Profinder B.07.00 with
mass tolerance <0.005 Da. All metabolomics data represent the average of at least two
independent triplicates.

Antibiotic permeability (rifampin uptake)

Rifampin uptake was measured using a previously validated device that enables timed
start-stop measurements of antibiotics in spent media [35, 36]. The device comprises a
plastic inset containing m7H9 with known antibiotic concentration. *Mtb*-laden filters were
generated using wild-type, ΔpstA1, and complement strains and placed onto media
containing 1 µg/mL rifampin at 37°C. After incubating for 2, 6, and 24 hours, bacterial
lysates and spent m7H9 were analyzed for rifampin content by LC-MS. Rifampin uptake
was calculated as: [RIF]drug only – [RIF]filtrate, with three biological replicates.

Statistical Analysis

Data are presented as mean ± standard deviation unless otherwise noted. Statistical
comparisons were performed using appropriate tests as specified in figure legends
(unpaired t-tests, one-way or two-way ANOVA with post-hoc corrections). Multiple
comparisons were corrected using the Holm-Sidak method or Tukey's Multiple
Comparisons where applicable. Statistical significance was defined as $p < 0.05$. Sample
sizes are indicated in individual figure legends.

Code and Data Availability

Acknowledgements:

We thank Dr. Kenan Murphy (University of Massachusetts) for generously providing the pKM444 and pKM464 plasmids used for ORBIT cloning, and Dr. Gyanu Lamichhane (Johns Hopkins University) for providing the pMH94 episomal vector. We are grateful to Megan Gelement (Johns Hopkins University) for contributions to methodology and data analysis discussions. We are grateful to SeqCenter (Pittsburgh, PA) for whole genome sequencing services. We thank the Johns Hopkins Genetic Resources Core Facility (GRCF) for RNA sequencing services.

Funding:

These studies were supported by NIAID/NIH grants R21 AI167027 to PCK ; R01AI168088 to HE. The funding source had no role in the study design, data collection, data analysis, data interpretation or writing of the report. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions:

Conceptualization: CD, PCK; Methodology: CD, GVAC, JSB, GYL, HE, PCK; Data acquisition: CD, GVAC, GYL; Data analysis: CD, GVAC, GYL; Supervision: JSB, HE, PCK; Funding acquisition: PCK, HE; Writing of original draft: CD, GVAC, PCK; Writing— review & editing: CD, GVAC, JSB, GYL, HE, PCK

Competing Interests:

All authors declare that they have no competing interests.

References:

1. *Global tuberculosis report*. 2023, World Health Organization: Geneva.

2. Nuermberger, E.L. and R.E. Chaisson, *Restocking the tuberculosis drug arsenal.* Nature medicine, 2024. **30**(3): p. 642–643.

3. Balaban, N.Q., et al., *Definitions and guidelines for research on antibiotic persistence.* Nature Reviews Microbiology, 2019. **17**(7): p. 441–448.

4. Joshi, H., D. Kandari, and R. Bhatnagar, *Insights into the molecular determinants involved in Mycobacterium tuberculosis persistence and their therapeutic implications.* Virulence, 2021. **12**(1): p. 2721–2749.

5. Brauner, A., et al., *Distinguishing between resistance, tolerance and persistence to antibiotic treatment.* Nature Reviews Microbiology, 2016. **14**(5): p. 320–330.

6. Matern, W.M., et al., *Genetic Determinants of Intrinsic Antibiotic Tolerance in Mycobacterium avium.* Microbiology Spectrum, 2021. **9**(2).

7. Matern, W.M., et al., *Functional Whole Genome Screen of Nutrient-Starved Mycobacterium tuberculosis Identifies Genes Involved in Rifampin Tolerance.* Microorganisms, 2023. **11**(9): p. 2269–2269.

8. Braibant, M., et al., *Identification of a second Mycobacterium tuberculosis gene cluster encoding proteins of an ABC phosphate transporter.* FEBS Letters, 1996. **394**(2): p. 206–212.

9. Pelly, S., et al., *REMap: Operon map of M. tuberculosis based on RNA sequence data.* Tuberculosis, 2016. **99**: p. 70–80.

10. Tischler, A.D., et al., *Mycobacterium tuberculosis Phosphate Uptake System Component PstA2 Is Not Required for Gene Regulation or Virulence.* PLOS ONE, 2016. **11**(8): p. e0161467–e0161467.

11. Vanzembergh, F., et al., *Effect of PstS sub-units or PknD deficiency on the survival of Mycobacterium tuberculosis.* Tuberculosis (Edinburgh, Scotland), 2010. **90**(6): p. 338–45.

12.   Healy, C., S. Ehrt, and A. Gouzy, *An exacerbated phosphate starvation response*
triggers Mycobacterium tuberculosis glycerol utilization at acidic pH. mBio, 2025.
16(1).

13.   Rifat, D., William R. Bishai, and Petros C. Karakousis, *Phosphate Depletion: A*
Novel Trigger for Mycobacterium tuberculosis Persistence. The Journal of
Infectious Diseases, 2009. **200**(7): p. 1126–1135.

14.   Tischler, A.D., et al., *Mycobacterium tuberculosis Requires Phosphate-Responsive*
Gene Regulation To Resist Host Immunity. Infection and Immunity, 2013. **81**(1): p.
317–328.

15.   Rengarajan, J., B.R. Bloom, and E.J. Rubin, *Genome-wide requirements for*
Mycobacterium tuberculosis adaptation and survival in macrophages. Proceedings
of the National Academy of Sciences, 2005. **102**(23): p. 8327–8332.

16.   Elliott, S.R. and A.D. Tischler, *Phosphate responsive regulation provides insights*
for ESX-5 function in Mycobacterium tuberculosis. Current Genetics, 2016. **62**(4):
p. 759–763.

17.   White, D.W., et al., *Mycobacterium tuberculosis Pst/SenX3-RegX3 Regulates*
Membrane Vesicle Production Independently of ESX-5 Activity. mBio, 2018. **9**(3).

18.   Rifat, D., D.A. Belchis, and P.C. Karakousis, *senX3-independent contribution of*
regX3 to Mycobacterium tuberculosis virulence. BMC microbiology, 2014. **14**: p.
265–265.

19.   Rifat, D. and P.C. Karakousis, *Differential regulation of the two-component*
regulatory system senX3-regX3 in Mycobacterium tuberculosis. Microbiology
(Reading, England), 2014. **160**(Pt 6): p. 1125–1133.

20.   Namugenyi, S.B., et al., *Mycobacterium tuberculosis PhoY Proteins Promote*
Persister Formation by Mediating Pst/SenX3-RegX3 Phosphate Sensing. mBio,
2017. **8**(4).

21.   Murphy, K.C., et al., *ORBIT: a New Paradigm for Genetic Engineering of*
Mycobacterial Chromosomes. mBio, 2018. **9**(6).

22.   Betts, J.C., et al., *Evaluation of a nutrient starvation model of Mycobacterium*
tuberculosis persistence by gene and protein expression profiling. Molecular
Microbiology, 2002. **43**(3): p. 717–731.

23.   Xie, Z., N. Siddiqi, and E.J. Rubin, *Differential Antibiotic Susceptibilities of Starved*
Mycobacterium tuberculosis Isolates. Antimicrobial Agents and Chemotherapy,
2005. **49**(11): p. 4778–4780.

24.   Danchik, C., S. Wang, and P.C. Karakousis, *Targeting the Mycobacterium*
tuberculosis Stringent Response as a Strategy for Shortening Tuberculosis
Treatment. Frontiers in microbiology, 2021. **12**: p. 744167–744167.

25.   Dutta, N.K., et al., *Inhibiting the stringent response blocks Mycobacterium*
tuberculosis entry into quiescence and reduces persistence. Science Advances,
2019. **5**(3).

26.   Thayil, S.M., et al., *The role of the novel exopolyphosphatase MT0516 in*
Mycobacterium tuberculosis drug tolerance and persistence. PloS one, 2011.
6(11): p. e28076–e28076.

27.   Chuang, Y.-M., et al., *Deficiency of the Novel Exopolyphosphatase Rv1026/PPX2*
Leads to Metabolic Downshift and Altered Cell Wall Permeability in Mycobacterium
tuberculosis. mBio, 2015. **6**(2).

28.   Chuang, Y.-M., et al., *Stringent Response Factors PPX1 and PPK2 Play an*
Important Role in Mycobacterium tuberculosis Metabolism, Biofilm Formation, and
Sensitivity to Isoniazid In Vivo. Antimicrobial agents and chemotherapy, 2016.
60(11): p. 6460–6470.

29.   Chugh, S., et al., *Polyphosphate kinase-1 regulates bacterial and host metabolic*
pathways involved in pathogenesis of Mycobacterium tuberculosis. Proceedings
of the National Academy of Sciences of the United States of America, 2024. **121**(2):
p. e2309664121–e2309664121.

30.   Singh, R., et al., *Polyphosphate deficiency in Mycobacterium tuberculosis is*
associated with enhanced drug susceptibility and impaired growth in guinea pigs.
Journal of bacteriology, 2013. **195**(12): p. 2839–51.

31.   Sureka, K., et al., *Polyphosphate kinase is involved in stress-induced mprAB-sigE-*
rel signalling in mycobacteria. Molecular microbiology, 2007. **65**(2): p. 261–76.

32.   Eoh, H. and K.Y. Rhee, *Multifunctional essentiality of succinate metabolism in*
adaptation to hypoxia in Mycobacterium tuberculosis. Proceedings of the National
Academy of Sciences, 2013. **110**(16): p. 6554–6559.

33.   Vijay, S., et al., *Ultrastructural Analysis of Cell Envelope and Accumulation of Lipid Inclusions in Clinical Mycobacterium tuberculosis Isolates from Sputum, Oxidative Stress, and Iron Deficiency.* Frontiers in Microbiology, 2018. **8**.

34.   Leisching, G., et al., *Virulence, biochemistry, morphology and host-interacting properties of detergent-free cultured mycobacteria: An update.* Tuberculosis, 2016. **100**: p. 53–60.

35.   Saini, V., et al., *Ergothioneine Maintains Redox and Bioenergetic Homeostasis Essential for Drug Susceptibility and Virulence of Mycobacterium tuberculosis.* Cell Reports, 2016. **14**(3): p. 572–585.

36.   Rieck, B., et al., *PknG senses amino acid availability to control metabolism and virulence of Mycobacterium tuberculosis.* PLoS pathogens, 2017. **13**(5): p. e1006399–e1006399.

37.   Lamarche, M.G., et al., *The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis.* FEMS microbiology reviews, 2008. **32**(3): p. 461–73.

38.   Wang, Q., et al., *PE/PPE proteins mediate nutrient transport across the outer membrane of Mycobacterium tuberculosis.* Science (New York, N.Y.), 2020. **367**(6482): p. 1147–1151.

39.   Richard-Greenblatt, M., et al., *Regulation of Ergothioneine Biosynthesis and Its Effect on Mycobacterium tuberculosis Growth and Infectivity.* The Journal of biological chemistry, 2015. **290**(38): p. 23064–76.

40.   Trivedi, A., et al., *Thiol reductive stress induces cellulose-anchored biofilm formation in Mycobacterium tuberculosis.* Nature Communications, 2016. **7**(1): p. 11392–11392.

41.   Chuang, Y.-M., D.A. Belchis, and P.C. Karakousis, *The Polyphosphate Kinase Gene ppk2 Is Required for Mycobacterium tuberculosis Inorganic Polyphosphate Regulation and Virulence.* mBio, 2013. **4**(3).

764 **Figures**

Lane	Region Amplified	Band Size
1	pstS3	1028 bp
2	pstS3-pstC2 junction	414 bp
3	pstC2	755 bp
4	pstC2-pstA1 junction	483 bp
5	pstA1	810 bp

Figure 1. The *pstS3-pstC2-pstA1* genes are co-transcribed as an operon. RT-PCR
analysis of H37Rv total RNA using primers spanning intragenic regions within *pstS3* (lane
1), *pstC2* (lane 3), and *pstA1* (lane 5), or intergenic junctions between *pstS3-pstC2* (lane
2) and *pstC2-pstA1* (lane 4). Amplification of all regions from cDNA confirms continuous
transcription across the three genes. No-RT controls (right panel) show the absence of
genomic DNA contamination. Expected amplicon sizes are indicated in the table.

Figure 2. Deletion of phosphate-specific transport genes enhances rifampin susceptibility. Time-kill assays were performed in 7H9 medium with 2 µg/mL rifampin for (A) ΔphoT, (B) ΔpstC2, and (C) ΔpstA1 in parallel with an empty vector control. (D) Time-kill assays performed with 7H9 medium with 0.5 µg/mL rifampin on wild-type, ΔpstA1 and pstA1 complement strains. Dotted lines indicate 2-$\log_{10}$ reduction from starting inoculum. Tables show minimum duration of killing (MDK) to achieve 2-$\log_{10}$ reduction and rifampin MIC values for each strain. Statistical significance for CFU comparisons at individual timepoints (A-C) was determined by unpaired t-test with Holm-Šídák correction for multiple comparisons. Statistical significance for MDK values (D) was determined by one-way ANOVA with Tukey's multiple comparisons test (*$P \le 0.05$; **$P \le 0.01$; ***$P \le 0.001$).

Figure 3. Δ*pstA1* survival under nutrient starvation and phosphate depletion. (**A**) Wild-type H37Rv and Δ*pstA1* survival after exposure to PBS-Tween for 14 days. No significant differences were observed by one-way ANOVA followed by Tukey's multiple comparisons test. (**B**) Δ*pstA1* maintained increased rifampin sensitivity after 2 weeks of PBS pre-treatment. Bacteria were treated with 2 µg/mL rifampin or DMSO vehicle control. Statistical differences indicated are for WT (RIF) vs Δ*pstA1* (RIF). Representative of two independent experiments.

(**C**) Δ*pstA1* shows no growth defect in phosphate-rich 7H9 medium but exhibits accelerated death under phosphate depletion (no Pi). Statistical differences shown are between WT no Pi and Δ*pstA1* no Pi conditions. Representative of three independent experiments.

Figure 4. ΔpstA1 exhibits an IFN-γ-dependent growth defect within macrophages.

Bacterial burden within PMA-differentiated THP1 cells after 4 hours of infection (D0) or 2

or 4 days without (A) or with (B) 24 hr IFN-γ polarization before infection. One way ANOVA

was performed followed by Sidak's multiple comparisons test to determine significance.

Adjusted p- value ≤0.05 = *; ≤0.01 = **; ≤0.001 = ***.

Figure 5. Rifampin uptake kinetics and cell morphology analysis of Δ*pstA1*. (A) Intracellular polyphosphate levels across different growth phases for wild-type and Δ*pstA1* normalized to total protein concentration. No significant differences were observed by one-way ANOVA followed by Tukey's multiple comparisons test. **(B-C)** Rifampin uptake kinetics in wild-type, Δ*pstA1*, and *pstA1* complement strains measured in a filter culture system. Bacteria were treated with 1.0 µg/mL rifampin and uptake was calculated as (initial rifampin concentration in media at 0 h - rifampin concentration at each timepoint) normalized to protein concentration. **(D-E)** Cell length and average cell

wall thickness of wild-type and Δ*pstA1* cultures grown in 7H9 medium with 0.05% Tween-80 during log phase. (**F-G**) Cell length and average cell wall thickness of cultures grown without Tween-80. Cell length measurements: n=11 bacteria per strain (with Tween-80), n=29 bacteria per strain (without Tween-80). Cell wall thickness was determined by averaging two measurements per bacterium: n=33 bacteria per strain (with Tween-80), n=29 bacteria per strain (without Tween-80). (**H**) Representative transmission electron microscopy images of whole bacteria or thin sections from log-phase cultures grown with or without 0.05% Tween-80. Data represent mean ± SD. Statistical significance was determined by one-way ANOVA followed by Tukey's multiple comparisons test (**B-C**) or unpaired t-test (**D-G**). **$P \le 0.01$; *$P \le 0.05$; ns, not significant.

Figure 6. Transcriptional profiling of ΔpstA1 with and without rifampin treatment.
(**A**) Heatmap of differentially expressed genes (DEGs) in wild-type and ΔpstA1 strains at
baseline and after rifampin treatment (30 and 60 min). Genes were filtered for absolute
log₂ fold change ≥ 1 and adjusted P < 0.05. Each row represents a gene, and each column
represents a biological replicate. Color scale indicates log₂ fold change relative to wild-
type control. (**B**) Functional enrichment analysis of DEGs for key comparisons: ΔpstA1
vs wild-type control (top left), wild-type control vs wild-type with 60 min rifampin treatment
(top right), ΔpstA1 control vs ΔpstA1 with rifampin treatment at 30 and 60 min (bottom
left), and wild-type control vs ΔpstA1 with 60 min rifampin treatment (bottom right). Bar
plots show the number of DEGs in each functional category. Correlation between RNA-
seq and RT-qPCR validation for selected genes across three comparisons. Each point
represents a gene, with correlation coefficients (R²) indicated for each comparison.

A

B

Figure 7. Metabolomic profiling reveals elevated antioxidant metabolites in rifampin-treated Δ*pstA1*. (**A**) Heatmap showing relative abundances of intracellular metabolites involved in antioxidant and redox pathways in wild type, Δ*pstA1*, and *pstA1* complement strains after 6 hours of rifampin exposure. Color scale indicates relative metabolite abundance (normalized to internal standards and protein concentration). (**B**) Quantification of selected antioxidant pathway metabolites. Bars represent mean ± SEM from three biological replicates. Statistical significance was determined by one-way ANOVA followed by Tukey's multiple comparisons test ($P < 0.1$).

Supplementary Figures

Figure S1. Whole genome sequencing of pstA1 deletion and complementation
strains.

Read depth coverage across the *pstA1* locus in wild-type H37Rv (blue), ΔpstA1 deletion
mutant (red), and Δ*pstA1::pstA1* complement strain (orange). Gray shaded region
indicates pstA1 gene boundaries (coordinates 1,036,999-1,037,925 bp). Vertical dashed
lines mark gene start and end positions. Wild-type exhibits uniform coverage (~150-200×
read depth) throughout the *pstA1* coding sequence. The Δ*pstA1* strain shows near-zero
coverage within the deleted region with maintained coverage in flanking sequences,
confirming successful targeted deletion. The complement strain displays reduced
coverage (~50-100× read depth) within *pstA1* relative to wild-type, consistent with
episomal expression from the pMH94 plasmid. Individual points represent sequencing
depth at each nucleotide position; smoothed trend lines (LOESS, span = 0.1) are shown
for wild-type and *pstA1* complement strains. Coverage was calculated from reads aligned
to the *M. tuberculosis* H37Rv reference genome using samtools depth and visualized in
R.

Figure S2. Venn diagram analysis of differentially expressed genes between ΔpstA1 and wild-type strains. Overlap of differentially expressed genes (DEGs) identified by RNA-seq (log2 fold change > 1, adjusted p-value < 0.05). Top panels show upregulated (left) and downregulated (right) genes in ΔpstA1 compared to WT across control, 30 min, and 60 min rifampin treatment conditions. Bottom panels show upregulated (left) and downregulated (right) genes comparing WT and ΔpstA1 responses to 60 min rifampin treatment relative to their respective untreated controls. Numbers indicate the count of unique or shared DEGs in each category.

Figure S3. Validation of RNA-seq results by quantitative PCR. Comparison of transcript levels measured by RNA-seq (light blue) and qPCR (dark blue) for selected differentially expressed genes in wild-type *M. tuberculosis* under control conditions and following 30- or 60-min rifampin treatment. Genes were selected to represent diverse functional categories including transcriptional regulators (whiB3, regX), metabolic enzymes (udgA, mgtA), PE/PPE family proteins (pe_pgrs10, ppe60, ppe65, rv3479), phosphate transport (pstC2), and ESX secretion (esxJ). Relative transcript levels are normalized to control conditions. Strong correlation between RNA-seq and qPCR measurements validates the RNA-seq dataset.

896 **Supplementary Table S1. Oligonucleotide primers used for RT-qPCR analysis of**

897 *Mycobacterium tuberculosis* **genes**

Target Gene	Direction	Sequence (5′→3′)	Length (bp)	GC Content (%)	Melting Temperature (°C)	Amplicon Size (bp)
sigA	Forward	CGACGAAGACCACGAAGAC	19	57.9	62	131
sigA	Reverse	CGTCTTCATCCCAGACGAAAT	21	47.6	62	
pstA1	Forward	CGTTGGTCCTGCTGATGTT	19	52.6	62	111
pstA1	Reverse	TCCATTTCGGAACGCCTAAC	20	50.0	62	
whiB3	Forward	ACTCATCGATGTTCTTCCATCC	22	45.5	62	147
whiB3	Reverse	AAACGCCATAGGGCTCAC	18	55.6	62	
udgA	Forward	GTCCACGTGTACGATCCAAAG	21	52.4	63	123
udgA	Reverse	CCATTCGGTAAGCACCAACA	20	50.0	63	
regX	Forward	CTACGTGACCAAGCCCTATTC	21	52.4	62	132
regX	Reverse	TCCACATCCATGCGAACC	18	55.6	62	
mgtA	Forward	CACCTTCCGAATGCTGAGA	19	52.6	62	145
mgtA	Reverse	CGGAACATCGGTTTGGTAGA	20	50.0	62	
rv0785	Forward	GGCACGGTATCCGGATTATT	20	50.0	62	132
rv0785	Reverse	CATGTCCAGACTTGGTGATGTA	22	45.5	62	
pe_pgrs10	Forward	CAACACGACGGGATTGTTGA	20	50.0	63	131
pe_pgrs10	Reverse	GCACGAACTGGGCGTAAA	18	55.6	63	
ppe60	Forward	TCAGCAACGTCAGTTCGATAG	21	47.6	62	100
ppe60	Reverse	GGAGCTAAGCCCTTCAACAT	20	50.0	62	
pstC2	Forward	GGCGAATCACGCCAATTTC	19	52.6	62	117
pstC2	Reverse	AGAGCCGTTATCGAACTCAAC	21	47.6	62	
esxJ	Forward	GACACCATGACCCAGATGAAT	21	47.6	62	108
esxJ	Reverse	CTGCTCTTGCTGTTCGTAGT	20	50.0	62	
rv3479	Forward	TCCGATCTCAGTGTGCTAGT	20	50.0	62	112
rv3479	Reverse	CCGTGGATTGTTGGAGATAGTC	22	50.0	62	
ppe65	Forward	GCAGTTGTCGCCATTCAATC	20	50.0	62	144
ppe65	Reverse	CTAAGCCCAAACAACGCTTTAG	22	45.5	62	

Abbreviations: bp, base pairs; GC, guanine-cytosine; RT-qPCR, reverse transcription

quantitative polymerase chain reaction.

All primers were designed to amplify *Mycobacterium tuberculosis* target genes. Primer
pairs generate amplicons ranging from 100 to 147 bp in length.

Supplementary Table S2. All genes meeting the significance threshold (|log2 fold
change| > 1, adjusted p-value < 0.05) across seven pairwise comparisons are listed.
Comparisons include temporal responses to rifampin treatment in wild-type and Δ*pstA1*
strains (30 and 60 min versus untreated controls), and strain-specific differences between
Δ*pstA1* and wild-type under control and rifampin-treated conditions. For each gene, the
following information is provided: comparison name, *M. tuberculosis* H37Rv gene
identifier (Rv number), gene name, log2 fold change, adjusted p-value (Benjamini-
Hochberg correction), and KEGG pathway annotations where available. A total of 58
genes were differentially expressed between Δ*pstA1* and wild-type under control
conditions (49 upregulated, 9 downregulated), 41 at 30 min rifampin (33 upregulated, 8
downregulated), and 41 at 60 min rifampin (31 upregulated, 10 downregulated). Only 6
genes responded to rifampin in wild-type (4 upregulated, 2 downregulated at 60 min), and
3 genes in Δ*pstA1* (all upregulated at 60 min), highlighting the substantial transcriptional
differences between the strains.