Mechanistic dissection of a dopamine-gated cation channel from

Daphnia reveals key determinants of ligand selectivity and

sensitivity

Tina M. McRunnel[1], Thomas E. Reynoldson[1], Iris Hardege[1*]

1. Department of Zoology, University of Cambridge, UK
*   Corresponding author

Abstract

The pentameric ligand-gated ion channels (pLGICs) are a deeply conserved superfamily that

includes nicotinic acetylcholine (nAChR), $GABA_A$Rs, glycine, and serotonin receptors and

mediates fast neurotransmission. While vertebrate pLGICs have been extensively

characterized, recent discoveries in invertebrates have revealed atypical members with

novel ligand specificities, including dopamine-gated ion channels (dop-LGICs) that enable

direct ionotropic dopaminergic signaling. Here, we investigate the molecular determinants of

catecholamine sensitivity and selectivity in a cation-permeable dopamine-gated receptor

from *Daphnia magna* (Dm-DopC1), which shares high structural homology with vertebrate

nAChRs. We find that mutations in loop B (D188N) near-abolish catecholamine responses,

while substitutions in loops A (S123P) and C (S227G, E230P) significantly alter sensitivity

and ligand selectivity, highlighting the cooperative role of these loops in shaping receptor

pharmacology. Complementary face mutations in loops D and E (N92T, A158G) further

refined dopamine versus norepinephrine sensitivity. Finally, we also show that this dopamine

receptor can be antagonized by both nicotinic and dopaminergic compounds, underscoring

its hybrid pharmacological profile. Together, these results reveal how subtle changes in

conserved binding motifs can shift receptor specificity from cholinergic to dopaminergic

signaling, providing new insights into the evolutionary flexibility of pLGICs and the molecular

basis of ionotropic catecholamine transmission in invertebrates.

1. Introduction

The processes governing neural communication are dynamic and highly evolutionarily conserved in animals across phyla (Moroz et al., 2021). Relying on the highly orchestrated molecular mechanisms that mediate the release and detection of neurotransmitters, converting extracellular chemical cues into either rapid electrical shifts in cell membrane potential or slower intracellular signaling cascades. Neurotransmission occurs via two distinct pathways: ionotropic and metabotropic. Ionotropic receptors, mediate fast synaptic transmission while in contrast, metabotropic receptors act through slow and second messenger pathways and produce longer lasting, modulatory effects.

The most widespread ionotropic receptors in animals fall into the pentameric or Cys-loop ligand gated ion channel superfamily (pLGIC), which includes nicotinic acetylcholine receptors (nAChR), $GABA_A$ receptors, glycine receptors (GlyR) and serotonin receptors ($5HT_3R$), each typically associated with distinct neurotransmitters and ionic conductances. The structural architecture of pLGICs is highly conserved and consists of five subunits arranged symmetrically around a central ion pore (Figure 1A). Each subunit contributes a β-sandwich extracellular domain (ECD) and four transmembrane α-helices. Neurotransmitter binding occurs at the interfaces between subunits, where on subunit serves as the 'principal face' contributing three extracellular loops A-C and the other contributes loops D-F to form a composite ligand binding site (Figure 1B) (Lynagh and Pless, 2014). While some pLGICs function as homomers most receptors are formed of heteromeric combinations of up to 5 different subunits (Boulin et al., 2008) with the ligand binding interface only forming between specific subunit combinations. Despite this conserved scaffold, structural studies have revealed how small differences in loop composition and conformation underlie substantial diversity in ligand recognition and receptor pharmacology (Corringer et al., 2010; Dalal et al., 2025; Dani, 2015; Pan et al., 2012; Sauguet et al., 2014; Unwin, 2003). This class of receptors have been extensively studied in vertebrate systems, and these receptors have become major pharmacological targets, forming the basis of anesthetics, antiparasitic and anxiolytics. This vertebrate-centric knowledge base has shaped much of our understanding of synaptic transmission. However, studies in invertebrate models have uncovered atypical pLGICs with unexpected ligand specificity, gating properties and synaptic roles (Cully et al., 1996; Ringstad et al., 2009). Characterizing these divergent receptors is essential for understanding the evolutionary diversity and functional plasticity possible in neurotransmission. For example, it was once assumed that each neurotransmitter exerts either an excitatory or an inhibitory effect depending on the receptor class that it activates. However, the discovery of both excitatory and inhibitory glutamate-gated chloride channels (GluCls) in *Caenorhabditis elegans* and *Drosophila melanogaster* challenged this view (Cully et al., 1996, 1994). These invertebrate-

specific channels are the targets for the antiparasitic treatment ivermectin, which locks the
channel in an open state, causing paralysis in parasitic worms.

One of the most intriguing discoveries in invertebrate systems is the identification of ligand-
gated ion channels gated by dopamine (dop-LGICs), which enable direct, ionotropic
dopaminergic signalling (Courtney et al., 2025; Morud et al., 2021; Ringstad et al., 2009). This
mode of neurotransmission contrasts with canonical dopaminergic signalling in mammals,
which is mediated by GPCRs. The most recently identified cation permeable dop-LGIC which
is gated by the catecholamines dopamine, norepinephrine and epinephrine has been shown
to be conserved in multiple invertebrate species, including *Daphnia magna, Apis mellifera*,
Aplysia california and *Octopus sinensis* (Courtney et al., 2025). In contrast, the anion dop-
LGICs studied previously appear to be nematode specific and gated by dopamine and to a
lesser extent tyramine (Morud et al., 2021; Ringstad et al., 2009). Intriguingly the invertebrate
cation dop-LGIC shares close sequence homology with the vertebrate α9α10 nicotinic
acetylcholine receptor (nAChR), including within the classic ligand binding domains. A great
deal of previous research, including more recent structural studies have revealed residues
that contribute to the binding of cholinergic ligands for nAChRs (Dani, 2015; Morales-Perez et
al., 2016; Su et al., 2025); however, we found that many of these are conserved in cationic
dop-LGICs (Figure 1C). In this study we sought to unravel the contribution of putative ligand
binding domains and individual residues in the homomeric *Daphnia magna* dop-LGIC, Dm-
DopC1, in the sensitivity of cationic dop-LGICs to catecholamines rather than cholinergic
ligands like its close relatives the nAChRs.

Figure 1 A. Full length AlphaFold2 model of Dm-DopC1 with the extracellular (ECD),

transmembrane (TM) and intracellular (ICD) domains highlighted. **B.** Close up of the ECD and

ligand binding domain made up of the interface between two adjacent subunits (light and dark

grey). Extracellular loops associated with ligand binding are labeled as follows; loop A

(orange), loop B (pink), loop C (green), loop D (turquoise), loop E (blue), loop F (yellow). **C.**

Trimmed alignment of the ECD of dop-LGICs, nAChRs and other pLGICs showing the ligand

binding loops. Alignment generated with MAFFT and residues colored with CLUSTAL

coloring.

2. Materials and methods

2.1 Structural Modelling and Sequence Alignments

Amino acid sequences of a range of pLGICs (full sequences are provided in Table S 1) were

aligned using the MAFTT algorithm and visualized with SnapGene (Dotmatics). Structural

models were predicted using AlphaFold ColabFold. For *D. magna* Dm-DopC1, five predicted

models were generated and overlaid in PyMOL. Residue positions were evaluated for

consistency and model confidence (pLDDT > 70). Sites exhibiting conserved structural

placement and high-confidence predictions were considered for site-directed mutagenesis.

2.2 Molecular Biology and Mutagenesis

Mutations were introduced using the Q5® Site-Directed Mutagenesis Kit (New England Biolabs). Constructs were linearized using NotI. cRNA was synthesized using the T3 mMessage mMachine Kit (ThermoFisher) and purified with the GeneJET RNA Cleanup Kit (ThermoFisher Scientific).

2.3 *Xenopus laevis* Oocyte Preparation and RNA Injection

Stage V-VI defolliculated *Xenopus laevis* oocytes were obtained from EcoCyte Bioscience and incubated in ND96 solution (in mM): 96 NaCl, 2 KCl, 1.8 CaCl2, 1 MgCl2, 5 HEPES, pH 7.5. Oocytes were stored at 18°C and injected within two days of delivery. For RNA injection, oocytes were placed in 96-well plates and injected using the Roboinject system (Multi Channel Systems, Germany). Each oocyte was injected with 50 nL of cRNA (250ng/µL). Oocytes were incubated at 18°C for 1 day prior to electrophysiological recordings.

2.4 Two-Electrode Voltage Clamp (TEVC) Recordings

TEVC recordings were performed using the Robocyte2 automated platform (Multi Channel Systems, Germany). Microelectrodes (0.5-1.5 MΩ resistance) were pulled using a P-97 micropipette puller (Sutter Instrument) and filled with 1 M KCl and 1.5 M acetic acid. Oocytes were voltage-clamped at -60 mV and recordings were sampled at 500 Hz throughout. Agonist solutions (dopamine, norepinephrine, epinephrine, tyramine, acetylcholine and serotonin) were prepared in ND96 on the day of recording and applied at concentrations ranging from 0.1 µM to 10 mM. Each ligand was perfused for 10 s followed by a 60 s wash. Wild-type oocytes were included in every recording batch. For antagonist experiment, compound solutions (50 nM to 500 µM, unless otherwise stated) were prepared in ND96 on the day of recording with 50 µM dopamine (Dm-DopC1). Before and after compound application, response to dopamine-only was recorded to obtain an Imax value per oocyte and to screen for potential irreversible binding of the antagonist. All compounds were perfused for 7 s followed by a 60 s wash. Perfusion speed was set to 3 ml/min throughout. Compounds used as antagonists: Acetylcholine chloride (Sigma-Aldrich (A6625)), Apomorphine hydrochloride (Abcam (ab269887)), Atropine (Abcam (ab145582)), Isoprenaline hydrochloride (Sigma-Aldrich (I5627)), Levamisole hydrochloride (MedChemExpress (HY-13666)), (-)-Nicotine detartrate (Abcam (ab120562)), Propranolol hydrochloride (Santa-Cruz Biotechnology (sc-3580)), (+)-Tubocurarine chloride pentahydrate (Sigma-Aldrich (93750)).

2.5 Data Analysis

All current amplitudes were exported using the Roboocyte2+ software (Multi Channel Systems) and normalized per oocyte to the maximum response (I/Imax) to account for

variation in expression levels using a custom python script adapted from (Morud et al., 2021).
Outliers were identified for each construct and removed within GraphPad Prism 10 using the
ROUT method (Q=1%) for every individual agonist/antagonist concentration. Agonist and
antagonist dose response curves were fitted in GraphPad Prism 10 using a three-parameter
logistic equation with a fixed Hill Slope of 1, unless otherwise stated. EC_{50} and IC_{50} values
were calculated from the fitted curves and reported as mean ± standard error of mean (SEM).
Statistical significance between wild-type and mutant dose-response curves was assessed
using a sum-of-squares F test. Peak current values were extracted using an existing Python
script (Morud et al., 2021). Normalized peak currents from the panel of ligands (1mM each:
tyramine, acetylcholine, norepinephrine, serotonin and epinephrine) were analyzed using a
one-way ANOVA followed by a Fisher's Least Significant Difference (LSD) post hoc test. All
TEVC recordings were baseline-subtracted unless otherwise stated. The same pooled wild-
type data was used for all comparisons.

2.6 Software and Code Availability

TEVC data were analyzed using custom Python scripts (https://github.com/hiris25/TEVC-
analysis-scripts). GraphPad Prism 10 was used for data plotting and fitting. MAFFT (Katoh
and Standley, 2013) was used for sequence alignment, alignments were visualized in
snapgene and ChimeraX 1.9 was used for 3D structural visualization. All constructs and raw
data are available upon request.

3. Results

3.1 Despite strong conservation with nAChRs, residues in loop C contribute to
catecholamine sensitivity and selectivity

The principle or α-type subunit contributes three structurally distinct loops to the ligand
binding domain, with loop C forming the front outer side of the binding pocket that closes
around the ligand during receptor activation and loops A and B forming the back of the
binding pocket (Dani, 2015; Morales-Perez et al., 2016). Loop C contains two conserved
structural features in nAChRs. The first is a pair of tyrosines implicated in aromatic cage
formation (Braun et al., 2016; Celie et al., 2004; Tomaselli et al., 1991) and the second is a
double cystine motif (Tomizawa et al., 2008) together forming a 'YXCC' motif. Both features
are present in dop-LGICs and in Dm-DopC1 are represented by Y226 and Y233, and the
C228 - C229, respectively (Figure 2E-F). Notably though both human and chicken nAChR
α9 and α10 subunits contain a glycine at the 'X' position, while the dop-LGICs contain serine
at this position. When we substituted S227 for glycine at this position in Dm-DopC1 we

observed a significant reduction in sensitivity to all catecholamines (Figure 2A-C). The
largest shift was seen with dopamine, shifting 7-fold from a WT EC_{50} of 84µM to 589µM,
norepinephrine EC_{50} also shifted 4-fold and epinephrine 1.8-fold (Figure 2A-C). This
suggests that the native serine may stabilise local interactions required for maintaining the C
loop conformation or supporting neighbouring aromatic residues for catecholamine
recognition.

We were also interested in the residue immediately following the 'YXCC' motif, which is
divergent across species and receptor types. However, both Dm-DopC2 and Am-DopC1
(from *Apis mellifera*), which have a higher relative selectivity for dopamine vs. the other
catecholamines, encode a proline at this position. A substitution to proline in this position in
Dm-DopC1 (E230P) produced a selective decrease in norepinephrine sensitivity only, with a
4-fold shift of EC_{50} from 98µM to 401µM, while maintaining wild type sensitivity for dopamine
and epinephrine (Figure 2A-C). However, despite altering the relative sensitivities, the
E230P mutation did not alter the ligand binding profile of the receptor, with norepinephrine
still eliciting similar relative peak currents to dopamine and epinephrine at 1mM (Figure 2D).
In contrast S227G which altered sensitivity to all catecholamines, abelite with more
pronounced shift in dopamine sensitivity, did alter the ligand binding profile such that
dopamine induced currents at 1mM fell significantly to approximately 50% of the maximum
currents achieved with norepinephrine and epinephrine (Figure 2D). Together these data
suggest that despite key nAChR features of loop C being conserved between cholinergic
and catecholamine binding receptors, the surrounding residues also contribute to ligand
specificity and binding capability.

Figure 2 Mutations in Loop C of a *D. magna* dop-LGIC reduce catecholamine sensitivity and alter agonist selectivity. A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type, S227G and E230P Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001, * = P<0.05, ns = not significant calculated by sum of squares F test. WT: *n* = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); S227G: *n* = 6 (dopamine), 3 (norepinephrine), 5 (epinephrine); E230P: *n* = 10 (dopamine), 4 (norepinephrine), 9 (epinephrine). **D.** Relative maximum I/Imax currents of *Xenopus* oocytes expressing wild-type, S227G and E230P Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: *n* = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); S227G: *n* = 11 per agonist; E230P: *n* = 6 per agonist. ** = P<0.01, ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the Loop C region. Residues S227 and E230 indicated by arrows, loop C is highlighted by the green bar. **F.** AlphaFold2 model of the loop C (green) region of Dm-DopC1 with S227 and E230 relative to Loop B (pink) at the ligand-binding site.

3.2 A conserved aspartate in loop B of principle face is critical to catecholamine sensitivity

Loop B forms part of the back wall of the ligand binding domains on the principle face and contains one of the conserved tryptophans forming the aromatic cage which is part of a

conserved region encoded the consensus motif 'SWTY' (Figure 3E-F). A semi conserved glycine, 4 amino acids downstream of the tryptophan has been implicated in defining high and low potency nAChRs (Puskar et al., 2012). In the dop-LGICs this post-tryptophan glycine is broadly conserved and directly preceded by an aspartate which is conserved across the dop-LGICs but not found in vertebrate α9/10 receptors. Substitution of this position (D188) in Dm-DopC1 with asparagine, more commonly found in nAChRs, led to a striking reduction in catecholamine sensitivity, leading to EC_{50} values for all three ligands of over 9mM, although true EC_{50} values could not be calculated due to lack of saturation even at 10mM (Figure 3A-C). The D188N mutation also led to significant changes in ligand binding profiles at 1mM with norepinephrine currents significantly reduced to 8% of the maximal currents, dopamine reduced to 37% and serotonin induced currents also falling to 19% (Figure 3D). However, it is worth noting that there was no appreciable difference in acetylcholine currents, which along with the residual currents observed from the catecholamines suggests that while loop B is critical to catecholamine gating, this single residue switch is insufficient in defining catecholamine vs cholinergic activation.

We also noted an additional residue of interest downstream of loop B, which differed only in Dm-DopC1 compared to other dop-LGICs and vertebrate α9/10. Given the differences in selectivity between Dm-DopC1 and Dm-DopC2 we hypothesised that the conserved aspartate at this position may also be contributing to ligand selectivity. Mutation at this location in Dm-DopC1 (N193D), led to a significant increase in EC_{50} for all catecholamines, most notably for norepinephrine with a 11-fold shift resulting in an EC_{50} of 1087μM, with smaller yet statistically significant shifts for dopamine and epinephrine of approximately 2-fold (Figure 3A-C). There was no change in overall ligand binding profile with all 3 catecholamines still eliciting the largest currents (Figure 3D). This suggests that the conserved aspartate at this position may contribute to selectivity between catecholamines, with receptors bearing this residue being relatively more selective for dopamine.

Figure 3 Mutations in Loop B of Dm-DopC1 disrupt catecholamine sensitivity and alter ligand selectivity. A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type, D188N and N193D Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001, *** = P<0.001, * = P<0.05 calculated by sum of squares F test. WT: *n* = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); D188N: *n* = 10 (dopamine), 5 (norepinephrine), 9 (epinephrine); N193D: *n* = 15 (dopamine), 5 (norepinephrine), 7 (epinephrine). **D.** Relative maximum I/Imax currents of *Xenopus* oocytes expressing wild-type, D188N and N193D Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: *n* = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); D188N: *n* = 11 per agonist; N193D: *n* = 11 per agonist. **** = P<0.0001, ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the Loop B region. Residues D188 and N193 indicated by arrows, loop B is highlighted by the pink bar. **F.** AlphaFold2 model of the loop B (pink) region of Dm-DopC1 with D188N and N193D relative to Loop C (green) at the ligand-binding site.

3.3 Residues on the complementary face of the binding pocket contribute to dopamine sensitivity and selectivity

In addition to the principal subunit, two loops on the complementary face of the binding pocket have also been shown to be involved in ligand binding in nAChRs. Loop D is composed of a single β strand and forms the base of the binding site, while loop E is

composed of two β strands connected by a short loop. We observed that loop D was broadly

conserved between both nAChRs and dop-LGICs, however we again noted that the two

dop-LGICs with higher dopamine selectivity, Dm-DopC2 and Am-DopC1, encoded a

threonine in the centre of the loop (Figure 4E-F). Creating this change in Dm-DopC1, N92T,

led to a striking and significant reduction of norepinephrine sensitivity, with an EC_{50} increase

from 98µM to 439µM, with no effect on dopamine or epinephrine (Figure 4A-C) and no

changes to overall ligand binding profile at 1mM (Figure 4D).This suggests that the threonine

at this position is important for discerning between norepinephrine and the other

catecholamines.

Despite overall high similarity within loop E, sequence alignments revealed a strongly

conserved aspartate at the proximal end of loop E in α9/10 that is missing from the dop-

LGICs, which frequently encode a hydrophobic leucine or phenylalanine at this position

(Figure 4E-F). When we substituted leucine with aspartate at this location in Dm-DopC1 we

could not generate functional receptors that responded to any of the ligands tested, although

due to leak current in oocytes we hypothesise that the protein was expressed (Figure S 1G).

This suggests that leucine at this position is critical to either channel formation or ligand

binding. Additionally, we also noted that the two more-dopamine selective dop-LGICs, Dm-

DopC2 and Am-DopC1, had a substitution of an alanine conserved across both receptor

types with a glycine (Figure S 1E-F). Despite being a minor alteration in sequence we found

that this substitution in Dm-DopC1, A158G, led to small but significant increases in EC_{50} for

all catecholamines to similar extents (EC_{50}s: dopamine: 235µM, norepinephrine: 440µM,

epinephrine: 438µM) (Figure S 1A-D). Again, supporting the notion that residues

surrounding key ligand binding loops contribute to receptor function, likely through the

stabilisation of these regions.

Figure 4 The N92T mutation in Loop D of a _D. magna_ dop-LGIC reduces norepinephrine sensitivity but preserves agonist selectivity. A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type and N92T Dm-DopC1 expressing _Xenopus_ oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001, ns = not significant calculated by sum of squares F test. WT: _n_ = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); N92T: _n_ = 10 (dopamine), 5 (norepinephrine), 8 (epinephrine). **D.** Relative maximum I/Imax currents of _Xenopus_ oocytes expressing wild-type and N92T Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: _n_ = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); N92T: _n_ = 11 per agonist. ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the loop D region. Residue N92 is indicated by an arrow, loop D is highlighted by the turquoise bar. **F.** AlphaFold2 model of the loop D (turquoise) region of Dm-DopC1 with N92 relative to loop B (pink), loop C (green), loop F (yellow), loop A (orange) and loop E (blue) at the ligand-binding site.

3.4 The conserved proline of loop A enhances catecholamine sensitivity

In line with previous findings, our sequence alignments identified a proline residue conserved across all pLGICs that sits within the conserved 'WXPD' motif (Figure 5E-F) (Braun et al., 2016). However, in Dm-DopC1 this location encodes a serine, which AlphaFold2 structural modelling placed near the inter-subunit ligand-binding interface, in proximity to Loops B and C (Figure 5F). Strikingly, restoring the proline at this position in Dm-DopC1 (S123P)

significantly increased sensitivity to all catecholamines (Figure 5A-C). Dopamine EC$_{50}$

decreased from $84.82 \pm 9.1\ \mu M$ to $39.84 \pm 6.06\ \mu M$, norepinephrine from $98.78 \pm 14.09\ \mu M$ to

$36.52 \pm 7.86\ \mu M$, and epinephrine from $169.17 \pm 29.5\ \mu M$ to $47.87 \pm 14.05\ \mu M$ (Figure 5A-C).

Interestingly, unlike the other mutations examined in this study, S123P also led to an increase

in Hill slope for norepinephrine (2.16 ± 0.19) compared to WT (1.31 ± 0.30), suggesting

enhanced co-operativity (Figure S 2). Overall, the relative ligand binding profile was unaltered,

although there was a small but non-significant increase in relative acetylcholine current (Figure

5D).

Figure 5 The conserved proline of loop A is required for ligand sensitivity in dop-LGICs.

A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-

type and S123P Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by I/Imax. EC$_{50}$

values ($\pm$ SEM) shown below each plot. **** = $P<0.0001$ calculated by sum of squares F test.

WT: n = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); S123P: n = 12, (dopamine), 4

(norepinephrine) and 8 (epinephrine). **D.** Relative maximum I/Imax currents of *Xenopus*

oocytes expressing wild-type and A158G Dm-DopC1, perfused for 10 s with 1 mM each:

dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and

acetylcholine (ACh). WT: n = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine,

serotonin); S123P: n = 12 per agonist. ns = not significant by one-way ANOVA and a Fisher's

LSD. **E.** Sequence alignment of the loop E region. Residue S123 is indicated by an arrow,

loop A is highlighted by the orange bar. **F.** AlphaFold2 model of the loop A (orange) region of

Dm-DopC1 with S123 relative to loop B (pink), loop C (green) and loop E (blue) at the ligand-

binding site.

3.5 Cholinergic, dopaminergic and adrenergic compounds antagonise dop-LGICs

To further dissect the pharmacology of the cationic dop-LGICs we tested a range of

cholinergic, dopaminergic and adrenergic compounds for their ability to activate or

antagonise Dm-DopC1. We found that both Dm-DopC1 dopamine induced currents could be

blocked by the nicotinic antagonist tubocurarine, with an IC_{50} of 0.9µM (Figure 6A). In

contrast to other nAChRs, the vertebrate a9/10 nAChRs are antagonised by nicotine

(Elgoyhen et al., 2001, 1994). In line with this we found that the dopamine currents of Dm-

DopC1 could be blocked at high concentrations of nicotine with and IC_{50} of 109µM (Figure

6A). We also tested the muscarinic blocker, atropine, which has been shown to act on

nAChRs (Parker et al., 2003; Zwart and Vijverberg, 1997), for its ability to block dopamine

currents and found that it behaved similarly to nicotine with relatively poor blocking (IC_{50}:

101µM (Figure 6A). This family of dop-LGICs have also been shown to be minimally

activated by acetylcholine (Courtney et al., 2025), and due to their close homology with

nAChRs and antagonism by nicotine we sought to determine if acetylcholine is able to

interact with the dopamine induced current, however we saw no evidence of this (Figure 6A).

In addition, we also tested apomorphine, a non-selective vertebrate dopamine receptor

agonist and the β-adrenoceptor agonist and antagonist, isoprenaline and propranolol

respectively. Apomorphine was able to block the dopamine induced current of Dm-DopC1,

with an IC_{50} of 6µM (Figure 6B). Meanwhile propranolol but not isoprenaline was able to

block dopamine currents (IC_{50}: 100µM) (Figure 6B). We also tested the ability of these

compounds to activate the receptors and observed no activation at 100µM (Figure S 3).

Figure 6 Dm-DopC1is sensitive to antagonism by cholinergic and dopaminergic

compounds. **A-B.** Inhibitory dose response curves for a panel of cholinergic (A) and

dopaminergic and adrenergic compounds (B). Error bars represent SEM. *n* = 4

(apomorphine, isoprenaline), 6 (atropine, propranolol), 9 (tubocurarine, acetylcholine), 12

(nicotine). IC50s are shown below each plot and calculated with a 4-parameter curve with

constrained bottom = 0, nd = not determined.

4. Discussion

Despite high levels of conservation of key ligand binding motifs with vertebrate and

invertebrate nAChRs this new class of dopamine-gated cation channels displays striking

differences in ligand sensitivity and antagonism by a range of compounds. In this study we

highlight some of the key residues that likely contribute to this functional difference (Table 1),

with many surrounding previously defined key motifs for cholinergic receptors.

Table 1 **Summary of Ec50 values for all tested mutations.** All values are given in µM and

calculated using a 3-parameter curve fit with a Hill slope of 1. P values were calculated using

the sum of squares F test.

		Dopamine			Norepinephrine			Epinephrine		
		E_C50	SEM	P value	E_C50	SEM	P value	E_C50	SEM	P value
Loop	**Construct**									
	WT	84.8	8.9	N/A	98.8	14.1	N/A	169.2	29.5	N/A
A	S123P	39.8	6.1	<0.0001	36.5	7.9	<0.0001	47.9	14.05	<0.0001
B	D188N	9466.9	3110	<0.0001	Unstable	Unstable	<0.0001	13153.8	2861.8	<0.0001
	N193D	172.6	31.1	0.0003	1087.9	190.6	<0.0001	384.8	109.9	0.0163
C	S227G	589.1	92.2	<0.0001	394.4	186.6	<0.0001	307.8	33.5	0.0262

	E230P	119.2	22.6	0.0812	401.1	73.6	<0.0001	166.2	34.8	0.9494
D	N92T	90.5	18.1	0.7445	439.8	114.1	<0.0001	170.2	57	0.9868
E	L155D	-	-	-	-	-	-	-	-	-
	A158G	234.6	30.7	<0.0001	439.8	57.5	<0.0001	438.2	57.97	<0.0001

The most striking effect was the almost complete loss of catecholamine gating upon
replacement of an aspartate in loop B (D188N) which follows the conserved 'SWTY' motif and
which is conserved in dop-LGICs. To our knowledge, this site, which in vertebrate α9/10
encodes an asparagine or glycine, has not been previously directly implicated in ligand binding
for nAChRs. However, it is unlikely that this residue alone contributes to dopamine selectivity
over cholinergic ligands given that aspartate is also found at this position in classic nAChRs
such as human α4 and the lack of significant acetylcholine gating over the wild type for this
mutation in Dm-DopC1. Interestingly, though immediately following this aspartate is a semi
conserved glycine residue which has been shown to distinguish high affinity vertebrate
nAChRs (eg. α4 which encodes lysine at this position) from low affinity receptors (eg. α7 which
encodes glycine). The increased affinity is thought to be due to stabilization of the open and
desensitized states by virtue of a hydrogen bond between this position in loop B and a proline
in loop C that forms in absence of glycine (Grutter et al., 2003; Kouvatsos et al., 2016; Puskar
et al., 2012). Interestingly glycine is present at this position in all but one dop-LGIC (which
instead encodes an alanine) further supporting the hypothesis that this hydrogen bond is key
to high affinity acetylcholine receptors. Decoupling of loops B and C may be important for
gating by aminergic ligands given that monoamine gated receptors from other subfamilies
such as the vertebrate $5HT_3R$ and invertebrate chloride permeable dopamine and histamine
receptors lack the hydrogen forming proline within loop C. In addition, loop B also contains
one of the conserved tryptophans associated with aromatic cage formation. Preceding this
tryptophan by 2 amino acids is another glycine residue conserved in nAChRs α subunits but
encoding an arginine in β subunits, which causes movements in the tryptophan preventing it
from forming a functional binding site (Morales-Perez et al., 2016), in dop-LGICs this site
encodes a alanine or serine. Mutations in loop B have also been shown to be more broadly
required for monoamine binding in $5HT_3Rs$ for example (Thompson et al., 2008). It would be
interesting to see if the D188N reverse mutation in vertebrate receptors alters acetylcholine
gating, although given the divergence of sequences at this position across nAChRs in both

vertebrates and invertebrates this residue may not be contributing in a significant way to
acetylcholine recognition.

We also confirmed the importance, yet surprising flexibility, of a highly conserved proline
residue within the 'WXPD' motif of loop A, which has been hypothesized to be critical for
maintaining structural integrity and receptor assembly in pLGIC receptors from bacteria
through to humans (Braun et al., 2016). We observed that Dm-DopC1, which exhibits a higher
EC_{50} for dopamine than other dop-LGICs (Courtney et al., 2025), diverged from this motif to
encode serine at this position (WKSD), which when substituted for proline, led to significant
increases in catecholamine sensitivity. The mirrors previous work in glycine receptors where
the same proline was replaced by an alanine leading to a decrease in ligand sensitivity
(Hughes et al., 2020; Lummis and Dougherty, 2022). Not only does our finding further highlight
the importance of a proline at this position, but it also shows that there is flexibility in the motif,
as the natural variant still displays micromolar ligand binding. As this serine is not conserved
within the rest of the dop-LGIC group, it is unlikely to be directly participating in catecholamine
recognition. Rather this suggests, as has been previously hypothesized, that the conserved
proline of the 'WXPD' motif contributes indirectly by stabilizing loop A conformation or
facilitating ligand alignment within the binding pocket.

An intriguing feature of the invertebrate cationic dop-LGICs is the presence of key motifs
usually attributed to acetylcholine and nicotine binding in nAChRs. This is highlighted by loop
C, which in dop-LGICs also encodes the conserved 'YXCC' motif (Karlin, 2002; Tomizawa et
al., 2008) as well as a pair of tyrosines implicated in aromatic cage formation (Braun et al.,
2016). However, the at 'X' position of the 'YXCC' motif the dop-LGICs encode a conserved
serine, which is not present in nAChRs. By exchanging this serine for glycine (present in
α9/10) in Dm-DopC1, we found that this serine is indeed important for catecholamine gating
but did not confer acetylcholine gating despite restoring the motif often found in nAChRs.
Separately, we also found that the residue immediately downstream of the 'YXCC' motif was
important for determining differences in ligand sensitivity between *Daphnia* dop-LGICs. The
functional impact of mutations at these positions supports the established role of loop C in
maintaining the geometry of the ligand binding pocket, particularly in orienting conserved
aromatic residues that contribute to catecholamine recognition. It also echoes our findings
within loop B, which suggests that a series of changes in the binding regions are required to
completely alter receptor selectivity from cholinergic to dopaminergic.

We also explored the role of loops D, E and F that form the complimentary face of the ligand
binding pocket and have been shown to contribute to stabilizing the extracellular domain. A
key feature of Loop D is the 'WXD' motif that is conserved across nAChRs, dop-LGICs,

5HT$_3$Rs and even some chloride channels (Braun et al., 2016; Corringer et al., 1995; Spier

and Lummis, 2000). Loop D is remarkably well conserved between dop-LGICs and nAChRs,

suggesting that its main function is to stabilize the ligand binding regions. However, we did

note that both Dm-DopC2 and Am-DopC1, which are more selective for dopamine than the

other dop-LGICs encoded a threonine 2 amino acids before the 'WXD' motif, rather than an

asparagine. When we substituted this asparagine for threonine in Dm-DopC1 we noted a

specific alteration in ligand sensitivity with only the norepinephrine EC$_{50}$ being adversely

affected while dopamine and epinephrine responses remained like wild type, making the

receptor overall more selective for dopamine. Interestingly, vertebrate nAChRs typically

encode a lysine or arginine at this position which is also conserved across other pLGICs

including glycine and GABA receptors, where disruption of this residue leads to almost

complete loss of ligand activation (Goldschen-Ohm et al., 2011; Grudzinska et al., 2005). The

cationic dop-LGICs, in line with hypotheses drawn up by Lynagh and Pless in 2014, also lack

positively charged residues in this position instead favoring polar residues also found in other

amine gated receptors including HisCl and the anionic dop-LGICs from nematodes, which

may assist in binding the hydroxyl termini or these ligands (Lynagh and Pless, 2014).

Collectively these mutations have identified residues critical for catecholamine recognition

across various conserved ligand binding domains, revealing that like in other pLGICs, both

faces of the ligand binding domain are critical not only for ligand gating but also relative ligand

selectivity. However, this study also demonstrates that ligand switching from acetylcholine to

(or from) dopamine requires a series of alterations across evolutionary time.

Finally, we also examined the pharmacology of dop-LGICs by testing a range of cholinergic,

dopaminergic and adrenergic compounds for their ability to interact with the dopamine induced

currents from both Dm-DopC1. The competitive nAChR antagonist, tubocurarine (Rahman et

al., 2022), efficiently blocked currents in the low µM range, broadly in line with classic nAChRs

(Zwart and Vijverberg, 1998). In contrast, other cholinergic compounds such as nicotine and

atropine (a muscarinic antagonist) had effects only at high concentrations. This is in line with

evidence that tubocurarine can act on a broader range of pLGICs including vertebrate 5HT$_3$Rs

(Peters et al., 1990; Yan et al., 1998) and invertebrate chloride gated cholinergic receptors

(Hardege et al., 2023). In addition, dop-LGICs could be blocked by apomorphine (non-

selective vertebrate dopamine agonist) and propranolol (β-adrenoceptor antagonist) in the µM

range suggesting that dop-LGICs have a distinct pharmacology from their closely related

nAChRs, with the potential of novel chemistries to be used to target this subfamily.

5. Conclusion

In this study we detail the mechanistic map of ligand selectivity of the invertebrate dopamine-
gated cation channel (dop-LGIC) family, extending the known functional repertoire of these
receptors. Our primary aim was to identify how variations in conserved ligand binding domains
enable dop-LGICs to respond selectively to catecholamines rather than cholinergic ligands,
despite their close sequence homology with vertebrate nAChRs. In line with this we found that
while core structural features of the pLGIC superfamily are conserved, functional divergence
is driven by key residue variations surrounding these canonical motifs in the extracellular
loops. We identified residues critical for catecholamine recognition across loops typically
associated with both the principal (A-C) and complementary faces (D-F) of the ligand binding
domain. Pharmacological analysis of dop-LGICs also confirmed a distinct profile from their
nAChR relatives, with dopamine-induced currents efficiently blocked in the micromolar range
by cholinergic, dopaminergic and adrenergic compounds.

Our results provide insight into the functional plasticity within invertebrate neurotransmission
systems and reveal how subtle differences in loop composition enable the switch from
cholinergic to aminergic ligand recognition. The fact that cationic dop-LGICs are conserved
across multiple invertebrate species, confirms the widespread nature of this signaling
mechanism. Furthermore, the demonstrated requirement for a series of evolutionary changes
to achieve ligand switching raises fundamental questions about the origin of ionotropic
neurotransmission, specifically whether dopamine or acetylcholine signaling emerged first.
Future research should focus on comparative structural analysis of dop-LGICs across basal
phyla, such as Platyhelminths or Nematodes (where unique anionic dop-LGICs are already
found), to trace the evolutionary trajectory and structural requirements that drove the transition
between cholinergic and dopaminergic ionotropic signaling.

Funding

I.H. and this work was kindly supported by a Royal Society University Research Fellowship
(URF\R1\231648).

Acknowledgements

The authors gratefully acknowledge Howard Baylis and all present and past members of the
Hardege lab for helpful discussions.

Author Contributions

T. M. M: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Visualization. **T. E. R:** Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Review & Editing. **I. H:** Conceptualization, Formal analysis, Resources, Data Curation, Writing - Original Draft, Visualization, Supervision, Project administration, Funding acquisition.

Declaration of interest

The authors declare no competing interests.

References

Boulin, T., Gielen, M., Richmond, J.E., Williams, D.C., Paoletti, P., Bessereau, J.-L., 2008. Eight genes are required for functional reconstitution of the Caenorhabditis elegans levamisole-sensitive acetylcholine receptor. Proceedings of the National Academy of Sciences 105, 18590–18595. https://doi.org/10.1073/pnas.0806933105

Braun, N., Lynagh, T., Yu, R., Biggin, P.C., Pless, S.A., 2016. Role of an Absolutely Conserved Tryptophan Pair in the Extracellular Domain of Cys-Loop Receptors. ACS Chem. Neurosci. 7, 339–348. https://doi.org/10.1021/acschemneuro.5b00298

Celie, P.H.N., van Rossum-Fikkert, S.E., van Dijk, W.J., Brejc, K., Smit, A.B., Sixma, T.K., 2004. Nicotine and carbamylcholine binding to nicotinic acetylcholine receptors as studied in AChBP crystal structures. Neuron 41, 907–914. https://doi.org/10.1016/s0896-6273(04)00115-1

Corringer, P.-J., Baaden, M., Bocquet, N., Delarue, M., Dufresne, V., Nury, H., Prevost, M., Van Renterghem, C., 2010. Atomic structure and dynamics of pentameric ligand-gated ion channels: new insight from bacterial homologues. J Physiol 588, 565–572. https://doi.org/10.1113/jphysiol.2009.183160

Corringer, P.J., Galzi, J.L., Eiselé, J.L., Bertrand, S., Changeux, J.P., Bertrand, D., 1995. Identification of a new component of the agonist binding site of the nicotinic alpha 7 homooligomeric receptor. J Biol Chem 270, 11749–11752. https://doi.org/10.1074/jbc.270.20.11749

Courtney, A., Styfhals, R., Dijck, M.V., Boulanger, J., Geenen, L., Lanoizelet, M., Hardege, I., Lassnig, M., Jansson, A.M., Gehler-Rahman, J., Almansa, E., Obenhaus, H.A., Seuntjens, E., Schafer, W.R., 2025. Novel dopaminergic neurotransmission in the Octopus visual system. https://doi.org/10.1101/2025.04.02.646353

Cully, D.F., Paress, P.S., Liu, K.K., Schaeffer, J.M., Arena, J.P., 1996. Identification of a
Drosophila melanogaster glutamate-gated chloride channel sensitive to the
antiparasitic agent avermectin. J Biol Chem 271, 20187–20191.
https://doi.org/10.1074/jbc.271.33.20187
Cully, D.F., Vassilatis, D.K., Liu, K.K., Paress, P.S., Van der Ploeg, L.H., Schaeffer, J.M.,
Arena, J.P., 1994. Cloning of an avermectin-sensitive glutamate-gated chloride
channel from Caenorhabditis elegans. Nature 371, 707–711.
https://doi.org/10.1038/371707a0
Dalal, V., Tan, B.K., Xu, H., Cheng, W.W., 2025. Cryo-EM structures of a pentameric ligand-
gated ion channel in liposomes. eLife 14, RP106728.
https://doi.org/10.7554/eLife.106728
Dani, J.A., 2015. Neuronal Nicotinic Acetylcholine Receptor Structure and Function and
Response to Nicotine. Int Rev Neurobiol 124, 3–19.
https://doi.org/10.1016/bs.irn.2015.07.001
Elgoyhen, A.B., Johnson, D.S., Boulter, J., Vetter, D.E., Heinemann, S., 1994. α9: An
acetylcholine receptor with novel pharmacological properties expressed in rat
cochlear hair cells. Cell 79, 705–715. https://doi.org/10.1016/0092-8674(94)90555-X
Elgoyhen, A.B., Vetter, D.E., Katz, E., Rothlin, C.V., Heinemann, S.F., Boulter, J., 2001. α10:
A determinant of nicotinic cholinergic receptor function in mammalian vestibular and
cochlear mechanosensory hair cells. Proceedings of the National Academy of
Sciences 98, 3501–3506. https://doi.org/10.1073/pnas.051622798
Goldschen-Ohm, M.P., Wagner, D.A., Jones, M.V., 2011. Three Arginines in the GABAA
Receptor Binding Pocket Have Distinct Roles in the Formation and Stability of
Agonist- versus Antagonist-Bound Complexes. Molecular Pharmacology 80, 647–
656. https://doi.org/10.1124/mol.111.072033
Grudzinska, J., Schemm, R., Haeger, S., Nicke, A., Schmalzing, G., Betz, H., Laube, B.,
2005. The β Subunit Determines the Ligand Binding Properties of Synaptic Glycine
Receptors. Neuron 45, 727–739. https://doi.org/10.1016/j.neuron.2005.01.028
Grutter, T., Prado de Carvalho, L., Le Novère, N., Corringer, P.J., Edelstein, S., Changeux,
J., 2003. An H-bond between two residues from different loops of the acetylcholine
binding site contributes to the activation mechanism of nicotinic receptors. The
EMBO Journal 22, 1990–2003. https://doi.org/10.1093/emboj/cdg197
Hardege, I., Morud, J., Courtney, A., Schafer, W.R., 2023. A Novel and Functionally Diverse
Class of Acetylcholine-Gated Ion Channels. J Neurosci 43, 1111–1124.
https://doi.org/10.1523/JNEUROSCI.1516-22.2022

Hughes, M.E., Mesoy, S.M., Capes, E., Lummis, S.C.R., 2020. Many Proline Residues in the
Extracellular Domain Contribute to Glycine Receptor Function. ACS Chem. Neurosci.
11, 2658–2665. https://doi.org/10.1021/acschemneuro.0c00320
Karlin, A., 2002. Emerging structure of the Nicotinic Acetylcholine receptors. Nat Rev
Neurosci 3, 102–114. https://doi.org/10.1038/nrn731
Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol 30, 772–780.
https://doi.org/10.1093/molbev/mst010
Kouvatsos, N., Giastas, P., Chroni-Tzartou, D., Poulopoulou, C., Tzartos, S.J., 2016. Crystal
structure of a human neuronal nAChR extracellular domain in pentameric assembly:
Ligand-bound α2 homopentamer. Proceedings of the National Academy of Sciences
113, 9635–9640. https://doi.org/10.1073/pnas.1602619113
Lummis, S.C.R., Dougherty, D.A., 2022. Expression of Mutant Glycine Receptors in
Xenopus Oocytes Using Canonical and Non-Canonical Amino Acids Reveals Distinct
Roles of Conserved Proline Residues. Membranes 12, 1012.
https://doi.org/10.3390/membranes12101012
Lynagh, T., Pless, S.A., 2014. Principles of agonist recognition in Cys-loop receptors. Front.
Physiol. 5. https://doi.org/10.3389/fphys.2014.00160
Morales-Perez, C.L., Noviello, C.M., Hibbs, R.E., 2016. X-ray structure of the human α4β2
nicotinic receptor. Nature 538, 411–415. https://doi.org/10.1038/nature19785
Moroz, L.L., Romanova, D.Y., Kohn, A.B., 2021. Neural versus alternative integrative
systems: molecular insights into origins of neurotransmitters. Philosophical
Transactions of the Royal Society B: Biological Sciences 376, 20190762.
https://doi.org/10.1098/rstb.2019.0762
Morud, J., Hardege, I., Liu, H., Wu, T., Choi, M.-K., Basu, S., Zhang, Y., Schafer, W.R.,
2021. Deorphanization of novel biogenic amine-gated ion channels identifies a new
serotonin receptor for learning. Curr Biol 31, 4282-4292.e6.
https://doi.org/10.1016/j.cub.2021.07.036
Pan, J., Chen, Q., Willenbring, D., Yoshida, K., Tillman, T., Kashlan, O.B., Cohen, A., Kong,
X.-P., Xu, Y., Tang, P., 2012. Structure of the pentameric ligand-gated ion channel
ELIC cocrystallized with its competitive antagonist acetylcholine. Nat Commun 3,
714. https://doi.org/10.1038/ncomms1703
Parker, J.C., Sarkar, D., Quick, M.W., Lester, R.A.J., 2003. Interactions of atropine with
heterologously expressed and native α3 subunit-containing nicotinic acetylcholine
receptors. Br J Pharmacol 138, 801–810. https://doi.org/10.1038/sj.bjp.0705124

Peters, J.A., Malone, H.M., Lambert, J.J., 1990. Antagonism of 5-HT3 receptor mediated
currents in murine N1E-115 neuroblastoma cells by (+)-tubocurarine. Neuroscience
Letters 110, 107–112. https://doi.org/10.1016/0304-3940(90)90796-C

Puskar, N.L., Lester, H.A., Dougherty, D.A., 2012. Probing the Effects of Residues Located
Outside the Agonist Binding Site on Drug-Receptor Selectivity in the Nicotinic
Receptor. ACS Chem Biol 7, 841–846. https://doi.org/10.1021/cb200448j

Rahman, M.M., Basta, T., Teng, J., Lee, M., Worrell, B.T., Stowell, M.H.B., Hibbs, R.E.,
2022. Structural mechanism of muscle nicotinic receptor desensitization and block by
curare. Nat Struct Mol Biol 29, 386–394. https://doi.org/10.1038/s41594-022-00737-3

Ringstad, N., Abe, N., Horvitz, H.R., 2009. Ligand-gated chloride channels are receptors for
biogenic amines in C. elegans. Science 325, 96–100.
https://doi.org/10.1126/science.1169243

Sauguet, L., Shahsavar, A., Poitevin, F., Huon, C., Menny, A., Nemecz, À., Haouz, A.,
Changeux, J.-P., Corringer, P.-J., Delarue, M., 2014. Crystal structures of a
pentameric ligand-gated ion channel provide a mechanism for activation.
Proceedings of the National Academy of Sciences 111, 966–971.
https://doi.org/10.1073/pnas.1314997111

Spier, A.D., Lummis, S.C.R., 2000. The Role of Tryptophan Residues in the 5-
Hydroxytryptamine3 Receptor Ligand Binding Domain*. Journal of Biological
Chemistry 275, 5620–5625. https://doi.org/10.1074/jbc.275.8.5620

Su, J., Yu, Z., Yin, Z., Zhang, Z., Zhao, J., Meng, Y., Li, R., Gao, Y., Zhang, H., Yu, R., Zhao,
Y., 2025. Molecular insights into the α6β4 nicotinic acetylcholine receptor function
and ligand recognition. Nat Commun 16, 3153. https://doi.org/10.1038/s41467-025-
58333-0

Thompson, A.J., Lochner, M., Lummis, S.C.R., 2008. Loop B Is a Major Structural
Component of the 5-HT3 Receptor. Biophysical Journal 95, 5728–5736.
https://doi.org/10.1529/biophysj.108.135624

Tomaselli, G.F., McLaughlin, J.T., Jurman, M.E., Hawrot, E., Yellen, G., 1991. Mutations
affecting agonist sensitivity of the nicotinic acetylcholine receptor. Biophysical Journal
60, 721–727. https://doi.org/10.1016/S0006-3495(91)82102-6

Tomizawa, M., Maltby, D., Talley, T.T., Durkin, K.A., Medzihradszky, K.F., Burlingame, A.L.,
Taylor, P., Casida, J.E., 2008. Atypical nicotinic agonist bound conformations
conferring subtype selectivity. Proc Natl Acad Sci U S A 105, 1728–1732.
https://doi.org/10.1073/pnas.0711724105

Unwin, N., 2003. Structure and action of the nicotinic acetylcholine receptor explored by
electron microscopy. FEBS Lett 555, 91–95. https://doi.org/10.1016/s0014-
5793(03)01084-6

Yan, D., Pedersen, S.E., White, M.M., 1998. Interaction of D-tubocurarine analogs with the

5HT3 receptor. Neuropharmacology 37, 251–257. https://doi.org/10.1016/s0028-

3908(98)00010-0

Zwart, R., Vijverberg, H.P.M., 1998. Four Pharmacologically Distinct Subtypes of α4β2

Nicotinic Acetylcholine Receptor Expressed in Xenopus laevisOocytes. Molecular

Pharmacology 54, 1124–1131. https://doi.org/10.1124/mol.54.6.1124

Zwart, R., Vijverberg, H.P.M., 1997. Potentiation and Inhibition of Neuronal Nicotinic

Receptors by Atropine: Competitive and Noncompetitive Effects. Molecular

Pharmacology 52, 886–895. https://doi.org/10.1124/mol.52.5.886

Supplementary data

Table S 1 Protein sequences used for alignments.

Name	Protein Sequence
O_sinensis_DopC1	MNYRLYIAMTLMKDIRILMSAFHLLMGASIALCDENEYRLVRDLMKNYNKQV RPSSNFSEALNVTFGVALAQIIDVDEKDQIITTNCWINQNWIDYKLQWDPIKY GNVSVIRLPFSDIWRPDILLYNNADVKAFESSISTNVIVTSEGNVTWLSMVIFK SSCSIDVKFFPFDEQNCSMEFASWTYDAYQVNILTNGEDNGDMSNYIENSE WSLIGFQQKRHVVRFSCCEEPYIIIHYHILIKRRPLFYLFNMVMPCILITLVALLG FYMPSDSGEKISMGITTLLSMTVFLMIVAEKMPPTSDVLPLIGLYYGITIAIVSF ATGMTVLTLNIHHKGTRGYEVPMILKKICFGVVAKILFIKLEVPLGSKESMPAA PVADYYRRFDTDKLPENGRLSPRFIRKLNPSASATASETFERQFMRVLQKVY QTIERNEMRLAEQDRRDLIKQEWQQLALIIDRVLLLIFLGLTITVTLTVFFQGP-- --- ----------------------------------
O_sinensis_DopC2	MEFWASITLTIFSLCLPSSSADDVELSLVNRLLKNYDKRVRPSLNSSEAINVTF GVALAQIIDVDERNQILTTNCWLNQVVWLDVNLRWDPDKYNGIKVVRIPFTDV WKPDILLYNNADVTSYFSSLSSNVIVTSEGNVTWLSMVIFKSSCRIDVRYFPF DEQNCSMQFASWTYDGFQLQLRMNSEEGDISNYIRNSEWVLIKLHVEWNLV YYSCCIEPYPDITYYIQIRRRPLFYVFNMILPCILITLVALLGFYIPSDSGEKVTM GITTLLSMTVFMMLVAENMPPTSDVLPLIGIYYGITISIVSFATAMTVFTLNIHH KGARGHEVPIIIKKICFGILAKIFCMRLEKPNNKSSNVAHLSDMSSSSEIPEMF VPCSSQSNSSYIRFKLSSQEESDQNAKPLLEMNPTLSQQQSMIRETRFHGD LETSNGGILSPRFSRLRGLQSRLTSPTHSSDDFERQFKRVLTKVYQTIERNEI

	RLAEQDRRDVIKLEWQQVAQIVDRLFLTIFVCVTILITCIVMFKAPHSTESLIS--- --
A_californica _DopC1	MADGINLILSLLVLASCAVMWTRGDPNEYRLVRDLMKGYDKRIRPSLNHTDS LNVTFGLALVQIIDVDEKNQILTTNCWLNQIWTDYNLRWDPAKYGGIKVLRLP YDEVWKPDILLHNNAEVSTFVSSTSSNLIVTNDGNVTWFSMVIFKSSCSINVR YFPFDAQNCSMQFASWSYDGYQVNLIMNTEKGDTSNYIHNSEWELIYLQVE RTVRIFSCCEEPYPDIHFHILIKRRPLFYIFNMILPCILITLVALLGFYIPSDSGEK VTMGITTLLSMTVFMMLVTENMPPTSDVLPLIGIYYGMTIFIVSFATAMTVFTL NIHHKGSRGKAVPMVIKKICFDFLAKMFCMKVDTWDEKHFTPDCNGTPGGA SNYIRFKVANQQDTDQLNMELKQTHEMSPLNRVELPLDSESFQSNGAVPRA FTSFRSSNGNGGSSSEQQNTELTGFEAYFSRVLQRVYQTIERNDIRLAEQDK REGIKLEWQQVAQITDRLLLTCFVLVTLTITGVVLLASPASVKV-------------------- --
D_magna_D opC1	MLRSHKMTQRIWIAFVTVLAIPGNRGDDGEYTRPIVSHEYRLIQHLMANYDR SVRPSRNAAEPLNITFGLALTQIIDVDERNQILTTNCWMNQMWTDYSLVWNV SEFGGIRVVRIPADKVWKSDIILYNNADSEYNSALLSTNVIVTHDGNLTWLSS AIFKSSCGINVEYFPFDEQKCSMKFSSWTYDGLQVNLLNQAEEGDLSNYVP NGEWDLIDLVVERNVVFYSCCEEPYPDITFHIVLRRRPLFYVFNLILPCVLITGI ALMSFYMPSDSGEKVTLGITTLLSMTVFLMVIGESMPPTSEKLPLIGLYYGVTI SLVSFATGLSVVTLNIHHRGMRGRGVPPMVKKIVFGVLAKLLFIQLDIAEGCS CSTKAKEFQQNNQFEMNERFNVDIGDSLKNESNGSLPHRLVVVFVAAAWFH VRQSSFGANSGLINNATSNIGPTDSFERQFLRVLNKVYQHIERSEARLADQD RKDAIKLEWQQVALVIDRFLLWIFIIATVAATFGILYMSPHSRLFSM--------------- --
D_magna_D opC2	MESKAMSPQSVSVRTAGRLLLSLGLFVLLPEAVHSDLNEYRLLRHLMENYDS NARPVANSTKALQVIFGMSLHLLIDVDERNQILTTNCWLTQKWNDSHLTWNA SDYGGIHVVRIPYHRVWRPDIILYNNADAQYTSSVINTNVIVSDNGEVVWLSH GIFRSSCDINVEYFPFDLQSCKMKWASWTYDGFQIDLKKQTDEGDVSNYQT NGEFDLVDFEAIHHEEYYSCCPEPYPDITYVIKMRRRPLFYVFNLILPCLLINGI ALLVFYVPSESGEKVTLGISSLLSMTVFLMTIRESLPPTEKTPLISLYYGVSICL VSFASGLSVVTLNLHHRGLRGTEVPYIMRRIILGGLARLVFLRFDVDKRHSPA STPTSAPAGEPSSSHSGDPIPASRLASRFEESPTKTCASGDPRCTCGKSRNI NQRGRHQPVGVSATRSSGWYSTPVNGGPSCSRNSQERLRLRLQELEPRAT DVYLDPEFSDKFQSAELDFLGIGPPSPKFYSRRVPSTSKAPHFQQQQQQPLI NPESASGSSSAEIGGGFGLLSQGSIYEQRLTELMARVSTAVERNENRLEEQ DRREAIALEWKQLALVCDRILLLTFVIITTVATCVVLTSSPYGP

A_mellifera_DopC1	MIPTSYKQPVCLTFVSALFETPARIFLGIEVRWSRGSAGFRVEIARWLKSIGY QSRTGGLAFKWSINSMSPLVLFFHYGVLAIIFGNGFGDEHEYRLTKYLLDGY DAGVRPAENSSQPLAVVFGLSLHHIIDVDEKNQILTTNCWVTQIWTDHHLKW NASEFAGIRVIRVPYNRVWRPDTILYNNADPQYSSAVINTNVIVSHTGEVVWL SHGIFRSSCDIDVEFFPFDEQRCVLKWASWTYDGYQLELEKQSEQGDVSNY QANGEFDLVNFSARRNVEYYSCCPEPYPDITYEIRLRRRPMFYVFNLILPCILI NSVALLVFYVPSESGEKVTLGISALLSMTVFLMTIRESLPPTEKTPLISLYYGV SICLVTFASALAVVTLNLHHRGVRGTRVPGIVRSLVLDKLARIVLLNFQEERRS EPVEPPRRKNNCPLHCKPELGQSQSSPKFVARREESNSSSPSLDLGKEGGL EAQWSRVLGRVHATIERNEKRLAEQDRRERMEFDWKQVALVSDRALLCVF FLTTIVSTTVILCGSPPSANIAKEG--
H_sapiens_ACHA9	MNWSHSCISFCWIYFAASRLRAAETADGKYAQKLFNDLFEDYSNALRPVED TDKVLNVTLQITLSQIKDMDERNQILTAYLWIRQIWHDAYLTWDRDQYDGLD SIRIPSDLVWRPDIVLYNKADDESSEPVNTNVVLRYDGLITWDAPAITKSSCV VDVTYFPFDNQQCNLTFGSWTYNGNQVDIFNALDSGDLSDFIEDVEWEVHG MPAVKNVISYGCCSEPYPDVTFTLLLKRRSSFYIVNLLIPCVLISFLAPLSFYLP AASGEKVSLGVTILLAMTVFQLMVAEIMPASENVPLIGKYYIATMALITASTALT IMVMNIHFCGAEARPVPHWARVVILKYMSRVLFVYDVGESCLSPHHSRERD HLTKVYSKLPESNLKAARNKDLSRKKDMNKRLKNDLGCQGKNPQEAESYCA QYKVLTRNIEYIAKCLKDHKATNSKGSEWKKVAKVIDRFFMWIFFIMVFVMTIL IIARAD
G_gallus_ACHA9	MKRNNLSSFYVSLWLLFTATMLQAVESAKGKYAQMLFNELFEDYSNALRPV EDTDKVLNVTLQITLSQIKDMDERNQILTAYLWIRQSWYDAYLKWDKDKYDG LDSIRIPSNLVWRPDIVLYNKADDDFSEPVNTNVVLRYDGKITWDAPAITKSS CVVDVSYFPFDSQQCNLTFGSWTYNGNQVDIINSLDSGDLSDFVEDVEWEI HGMPAVKNVITYGCCSEPYPDVTFTLILKRKSSFYIFNLLLPCILISFLAPLGFY LPADSGEKVSLGVTVLLALTVFQLMVAEIMPPSENVPLIGKYYIATMTMITAST ALTIIIMNVHHCGSEAKPVPQWARVVILDYMSKIFFVYDVGENCTSPRREKEQ EHRLEGGDMCRGGDGKSHLSSRNDDSDLKENLGNWNKSFGVHGENVRE NVNCCSCYKMLIKNIEYIANCVRDHKANRAKGIEWKKVAKVMDRFFMWIFFI MVFFMSVLIIGKAA
G_gallus_ACH10	MGSAPLPACLLALSLAGTVLAPGCGAAQGRLAHKLLHDLFANYSSALRPAED TERALNVTLQVTLSQIIDMDERNQVLTSYLWVRQAWLDAHLAWDKDAYGGI DSIRIPSSYVWRPDIVLYNNADERFGGSMETNVVLRSDGHIMWDSPAITKSS CKVDVSYFPFDGQQCRLTFGSWTYNGNQIDLRNLLDTGDLTDFVENVEWEV

	LGMPATRNVVTYGCCSEPYPDVTYTLLLRRRASFYIFNLLLPCVMISFLAPLG FYLPADSGEKVSLGVTVLLALTVFQLLVAESMPPSESVPLIGKYYIATMTMITA STALTIFIMNIHHCGPGARPVPPWARRLILHHLARALCVCEVGESCGRPQRE GTGGMGPRDPPGEGVEPGLCPRSRCLCHHHAVLSSVGYIAGVFRRHRTAQ RRAAEWKKVAKVMDRFFMWVFFLMVFLMSVLVIGKAA
H_sapiens_ ACH10	MGLRSHHLSLGLLLLFLLPAECLGAEGRLALKLFRDLFANYTSALRPVADTDQ TLNVTLEVTLSQIIDMDERNQVLTLYLWIRQEWTDAYLRWDPNAYGGLDAIRI PSSLVWRPDIVLYNKADAQPPGSASTNVVLRHDGAVRWDAPAITRSSCRVD VVAAFPFDAQHCGLTFGSWTHGGHQLDVRPRGAAASLADFVENVEWRVLG MPARRRVLTYGCCSEPYPDVTFTLLLRRRAAAYVCNLLLPCVLISLLAPLAFH LPADSGEKVSLGVTVLLALTVFQLLLAESMPPAESVPLIGKYYMATMTMVTFS TALTILIMNLHYCGPSVRPVPAWARALLLGHLARGLCVRERGEPCGQSRPPE LSPSPQSPEGGAGPPAGPCHEPRCLCRQEALLHHVATIANTFRSHRAAQRC HEDWKRLARVMDRFFLAIFFSMALVMSLLVLVQAL
H_sapiens_ ACHA7	MRCSPGGVWLALAASLLHVSLQGEFQRKLYKELVKNYNPLERPVANDSQPL TVYFSLSLLQIMDVDEKNQVLTTNIWLQMSWTDHYLQWNVSEYPGVKTVRF PDGQIWKPDILLYNSADERFDATFHTNVLVNSSGHCQYLPPGIFKSSCYIDVR WFPFDVQHCKLKFGSWSYGGWSLDLQMQEADISGYIPNGEWDLVGIPGKR SERFYECCKEPYPDVTFTVTMRRRTLYYGLNLLIPCVLISALALLVFLLPADSG EKISLGITVLLSLTVFMLLVAEIMPATSDSVPLIAQYFASTMIIVGLSVVVTVIVL QYHHHDPDGGKMPKWTRVILLNWCAWFLRMKRPGEDKVRPACQHKQRRC SLASVEMSAVAPPPASNGNLLYIGFRGLDGVHCVPTPDSGVVCGRMACSPT HDEHLLHGGQPPEGDPDLAKILEEVRYIANRFRCQDESEAVCSEWKFAACV VDRLCLMAFSVFTIICTIGILMSAPNFVEAVSKDFA
H_sapiens_ ACHA3	MGSGPLSLPLALSPPRLLLLLLLSLLPVARASEAEHRLFERLFEDYNEIIRPVA NVSDPVIIHFEVSMSQLVKVDEVNQIMETNLWLKQIWNDYKLKWNPSDYGG AEFMRVPAQKIWKPDIVLYNNAVGDFQVDDKTKALLKYTGEVTWIPPAIFKSS CKIDVTYFPFDYQNCTMKFGSWSYDKAKIDLVLIGSSMNLKDYWESGEWAII KAPGYKHDIKYNCCEEIYPDITYSLYIRRLPLFYTINLIIPCLLISFLTVLVFYLPS DCGEKVTLCISVLLSLTVFLLVITETIPSTSLVIPLIGEYLLFTMIFVTLSIVITVFV LNVHYRTPTTHTMPSWVKTVFLNLLPRVMFMTRPTSNEGNAQKPRPLYGAE LSNLNCFSRAESKGCKEGYPCQDGMCGYCHHRRIKISNFSANLTRSSSSES VDAVLSLSALSPEIKEAIQSVKYIAENMKAQNEAKEIQDDWKYVAMVIDRIFL WVFTLVCILGTAGLFLQPLMAREDA
H_sapiens_ ACHA4	MELGGPGAPRLLPPLLLLLGTGLLRASSHVETRAHAEERLLKKLFSGYNKWS RPVANISDVVLVRFGLSIAQLIDVDEKNQMMTTNVWVKQEWHDYKLRWDPA

	DYENVTSIRIPSELIWRPDIVLYNNADGDFAVTHLTKAHLFHDGRVQWTPPAI YKSSCSIDVTFFPFDQQNCTMKFGSWTYDKAKIDLVNMHSRVDQLDFWESG EWVIVDAVGTYNTRKYECCAEIYPDITYAFVIRRLPLFYTINLIIPCLLISCLTVLV FYLPSECGEKITLCISVLLSLTVFLLLITEIIPSTSLVIPLIGEYLLFTMIFVTLSIVIT VFVLNVHHRSPRTHTMPTWVRRVFLDIVPRLLLMKRPSVVKDNCRRLIESMH KMASAPRFWPEPEGEPPATSGTQSLHPPSPSFCVPLDVPAEPGPSCKSPS DQLPPQQPLEAEKASPHPSPGPCRPPHGTQAPGLAKARSLSVQHMSSPGE AVEGGVRCRSRSIQYCVPRDDAAPEADGQAAGALASRNTHSAELPPPDQP SPCKCTCKKEPSSVSPSATVKTRSTKAPPPHLPLSPALTRAVEGVQYIADHL KAEDTDFSVKEDWKYVAMVIDRIFLWMFIIVCLLGTVGLFLPPWLAGMI
C_elegans_ UNC-29	MRTNRLSWILVLSVVIFLVIINTINASDDEERLMVDVFRGYNSLIQPVRNSSEL PLIVKMALQLVLLINVDEKDQVMHTNVWLTLQWHDFQMKWNPVNYGEIKQIR VSPDKVWLPDIVLFNNADGNYEVSFMCNVVINHKGDMLWVPPAIYKSSCIID VEFFPFDEQVCTLVFGSWTYNENEIKLEFVQAELVDVSEYSASSIWDVIDVPA SLVNKRSRIEFQVRIRRKTLFYTVVLIIPTVLMAFLSMAVFFLPTDSGEKITLTIS VLLSIVVFLLLVSKILPPTSSTIPLMAKYLLLTFVLNVITILVTVIIINVYFRGPRTH RMPQWVRVVFLQFLPKLVCMKRPKSASERSAVRSGMAQLPGVGQFTLSPS AHHPLCPSADDRTTTIRNTASNETSAYYPLSTDALRAIDAIEYITEHLKRDEQH KSFRDDWKYVAMIIDRLLLYVFFGITVGGTCGILFSAPHVFQRIDQQEMLDRL KEKYDTASNIP
D_chrysanth emi_ELIC	APADNAADARPVDVSVSIFINKIYGVNTLEQTYKVDGYIVAQWTGKPRKTPG DKPLIVENTQIERWINNGLWVPALEFINVVGSPDTGNKRLMLFPDGRVIYNAR FLGSFSNDMDFRLFPFDRQQFVLELEPFSYNNQQLRFSDIQVYTENIDNEEID EWWIRKASTHISDIRYDHLSSVQPNQNEFSRITVRIDAVRNPSYYLWSFILPL GLIIAASWSVFWLESFSERLQTSFTLMLTVVAYAFYTSNILPRLPYTTVIDQMII AGYGSIFAAILLIIFAHHRQAMGVEDDLLIQRCRLAFPLGFLAIGCVLVIRGITL
H_sapiens_ 5HT3A	MLLWVQQALLALLLPTLLAQGEARRSRNTTRPALLRLSDYLLTNYRKGVRPV RDWRKPTTVSIDVIVYAILNVDEKNQVLTTYIWYRQYWTDEFLQWNPEDFDN ITKLSIPTDSIWVPDILINEFVDVGKSPNIPYVYIRHQGEVQNYKPLQVVTACSL DIYNFPFDVQNCSLTFTSWLHTIQDINISLWRLPEKVKSDRSVFMNQGEWEL LGVLPYFREFSMESSNYYAEMKFYVVIRRRPLFYVVSLLLPSIFLMVMDIVGF YLPPNSGERVSFKITLLLGYSVFLIIVSDTLPATAIGTPLIGVYFVVCMALLVISL AETIFIVRLVHKQDLQQPVPAWLRHLVLERIAWLLCLREQSTSQRPPATSQAT KTDDCSAMGNHCSHMGGPQDFEKSPRDRCSPPPPPREASLAVCGLLQELS SIRQFLEKRDEIREVARDWLRVGSVLDKLLFHIYLLAVLAYSITLVMLWSIWQY A

H_spaiens_ACHB2	MARRCGPVALLLGFGLLRLCSGVWGTDTEERLVEHLLDPSRYNKLIRPATN GSELVTVQLMVSLAQLISVHEREQIMTTNVWLTQEWEDYRLTWKPEEFDNM KKVRLPSKHIWLPDVVLYNNADGMYEVSFYSNAVVSYDGSIFWLPPAIYKSA CKIEVKHFPFDQQNCTMKFRSWTYDRTEIDLVLKSEVASLDDFTPSGEWDIV ALPGRRNENPDDSTYVDITYDFIIRRKPLFYTINLIIPCVLITSLAILVFYLPSDCG EKMTLCISVLLALTVFLLLISKIVPPTSLDVPLVGKYLMFTMVLVTFSIVTSVCVL NVHHRSPTTHTMAPWVKVVFLEKLPALLFMQQPRHHCARQRLRLRRRQRE REGAGALFFREAPGADSCTCFVNRASVQGLAGAFGAEPAPVAGPGRSGEP CGCGLREAVDGVRFIADHMRSEDDDQSVSEDWKYVAMVIDRLFLWIFVFVC VFGTIGMFLQPLFQNYTTTTFLHSDHSAPSSK
H_sapines_ACHB4	MRRAPSLVLFFLVALCGRGNCRVANAEEKLMDDLLNKTRYNNLIRPATSSSQ LISIKLQLSLAQLISVNEREQIMTTNVWLKQEWTDYRLTWNSSRYEGVNILRIP AKRIWLPDIVLYNNADGTYEVSVYTNLIVRSNGSVLWLPPAIYKSACKIEVKYF PFDQQNCTLKFRSWTYDHTEIDMVLMTPTASMDDFTPSGEWDIVALPGRRT VNPQDPSYVDVTYDFIIKRKPLFYTINLIIPCVLTTLLAILVFYLPSDCGEKMTLC ISVLLALTFFLLLISKIVPPTSLDVPLIGKYLMFTMVLVTFSIVTSVCVLNVHHRS PSTHTMAPWVKRCFLHKLPTFLFMKRPGPDSSPARAFPPSKSCVTKPEATA TSTSPSNFYGNSMYFVNPASAASKSPAGSTPVAIPRDFWLRSSGRFRQDVQ EALEGVSFIAQHMKNDDEDQSVVEDWKYVAMVVDRLFLWVFMFVCVLGTV GLFLPPLFQTHAASEGPYAAQRD
C_elegans_LGC-53	MIISYLARIVLLLPELSSGDTDRLAEIAANQEHSIQLDQMPPDRTFHYPNIPSS QHDRRREGGKRKKFRKERLDEPCTAINETLRKNLLQELFDDSYDKNNLPSA NSTEVIVELTVQSITEISEFSSSFKADVWFSQIWRDPRLDFSDRNYCLKNISLA SHKLPTMWSPNVCFVNSKKVEIHASPSQNILLLIFPNGTVWLNFRVSLTGPC KLDLTYFPMDRQSCNLVFESYSYNTAEVRIVWRDWDAVTIPDPDSKNLPDFE LVNIEHMNATLVYTAGLWDQLEVKFTFRRLYGYYVLQAYMPTYLSVFISWIAF WIDTKALPARITLGVSSLMALTFQFGNIVKNLPRVSYVKALDIWMFGCVGFIFL SLVELAVVGFADKLEAKRRRHNRCKEQLMMRSDSEQQWLSRLSGQRPQVS ETNSDATHTIQINDGNGNVRRRKSEEKRQKLLLQMESPMFVNGEKIDEISAK LFPLLFTAFNIFYWFYYIGMSGGFF
C_elegans_LGC-52	MIYSIQALFIETILMSARIHDVTYQHIEDFTCDDIGLSSLSGAKATDLARILMTNY SRNALPEPAPVKVEVEITIQDISDISAITGTFTIDYWISAIWNDKRLAFSHLDPC RKNLSLDHDMEPKLWSPNVCIVNSKSTKVHDSPKPNILLMIFPNGTIWLNYRI RSEAPCQMNLRNFPLDSIRCSLVFESYSYNAADVELQWLEWSPVSTVRNDY NLPDFRMTNITYGSVTEAYTAGMWHRLSVSIHFERLYGFYILQMYLPTYISVFI SWIAFWMDTKALPARITLSVSSLMALTFQFGNIVKSLPKASYVKAIDIWMFSC

	VGFIFFSLIELAIVAYNDKMHDQRLREVRCSVGNLNSNGIGQYNSTTRRSFVE LSARRAKGSELGASIDRVASMLFPTMFALFNLVYWAYYLTTSMNPVYT
D_melanoga ster_HisCl1	MQSPTSKLVEFRCLIALAIYLHALEQSIQHCHCVHGYRNNTESAELVSHYESS LSLPDILPIPSKTYDKNRAPKLLGQPTVVYFHVTVLSLDSINEESMTYVTDIFLA QSWRDPRLRLPENMSEQYRILDVDWLHSIWRPDCFFKNAKKVTFHEMSIPN HYLWLYHDKTLLYMSKLTLVLSCAMKFESYPHDTQICSMMIESLSHTVEDLV FIWNMTDPLVVNTEIELPQLDISNNYTTDCTIEYSTGNFTCLAIVFNLRRRLGY HLFHTYIPSALIVVMSWISFWIKPEAIPARVTLGVTSLLTLATQNTQSQQSLPP VSYVKAIDVWMSSCSVFVFLSLMEFAVVNNFMGPVATKAMKGYSDENISDL DDLKSALQHHRESIIEPQYDTFCHGHATAIYIDKFSRFFFPFSFFILNIVYWTTF L
H_sapiens_ GBRA1	MRKSPGLSDCLWAWILLLSTLTGRSYGQPSLQDELKDNTTVFTRILDRLLDG YDNRLRPGLGERVTEVKTDIFVTSFGPVSDHDMEYTIDVFFRQSWKDERLKF KGPMTVLRLNNLMASKIWTPDTFFHNGKKSVAHNMTMPNKLLRITEDGTLLY TMRLTVRAECPMHLEDFPMDAHACPLKFGSYAYTRAEVVYEWTREPARSV VVAEDGSRLNQYDLLGQTVDSGIVQSSTGEYVVMTTHFHLKRKIGYFVIQTY LPCIMTVILSQVSFWLNRESVPARTVFGVTTVLTMTTLSISARNSLPKVAYAT AMDWFIAVCYAFVFSALIEFATVNYFTKRGYAWDGKSVVPEKPKKVKDPLIK KNNTYAPTATSYTPNLARGDPGLATIAKSATIEPKEVKPETKPPEPKKTFNSV SKIDRLSRIAFPLLFGIFNLVYWATYLNREPQLKAPTPHQ

Figure S 1 The A158G mutation in Loop E of a _D. magna_ dop-LGIC decreases catecholamine sensitivity.

A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type and A158G Dm-DopC1 expressing _Xenopus_ oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001, ns = not significant calculated by sum of squares F test. WT: n = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); A158G: n = 10 (dopamine), 11 (norepinephrine), 10 (epinephrine). **D.** Relative maximum I/Imax currents of _Xenopus_ oocytes expressing wild-type and A158G Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: n = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); A158G: n = 6 per agonist. ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the loop E region. Residues A158 and L155 are indicated by arrows, loop E is highlighted by the blue bar. **F.** AlphaFold2 model of the loop E (blue) region of Dm-DopC1 with A158 and L155 relative to loop B (pink), loop C (green), loop F (yellow), loop A (orange) and loop D (turquoise) at the ligand-binding site. **G.**

 Baseline currents in uninjected oocytes or oocytes expressing WT or L155D Dm-DopC1.

669 Baseline currents measured 5s before agonist application. Error bars represent SEM, *n* = 3

670 (WT), 6 (uninjected), 7 (L155D).

Figure S 2 S123P exhibits a steeper Hill slope than wild type for norepinephrine.

A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-
type and S123P Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by I/Imax. Hill
slope values are shown underneath each plot with SEM, calculated with a 4-parameter curve.
WT: *n* = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); S123P: n = 12, (dopamine), 4
(norepinephrine) and 8 (epinephrine).

Figure S 3 Continuous traces of Dm-DopC1 oocytes exposed to a panel of compounds.
Red bar indicates application of 100μM compound.

683