Mechanistic dissection of a dopamine-gated cation channel from

***Daphnia* reveals key determinants of ligand selectivity and**

sensitivity

Tina M. McRunnel[a], Thomas E. Reynoldson[a], Yu Zhu[b], Taufiq Rahman[b], Iris Hardege[a*]

[a] Department of Zoology, University of Cambridge, CB2 3EJ, UK

[b] Department of Pharmacology, University of Cambridge, CB2 1QR, UK

*   Corresponding author

Abstract

The pentameric ligand-gated ion channels (pLGICs) are a deeply conserved superfamily that

includes nicotinic acetylcholine (nAChR), $GABA_A Rs$, glycine, and serotonin receptors and

mediates fast neurotransmission. While vertebrate pLGICs have been extensively

characterized, recent discoveries in invertebrates have revealed atypical members with

novel ligand specificities, including dopamine-gated ion channels (dop-LGICs) that enable

direct ionotropic dopaminergic signaling. Here, we investigate the molecular determinants of

catecholamine sensitivity and selectivity in a cation-permeable dopamine-gated receptor

from *Daphnia magna* (Dm-DopC1), which shares high structural homology with vertebrate

nAChRs. We find that mutations in loop B (D188N) near-abolish catecholamine responses,

while substitutions in loops A (S123P) and C (S227G, E230P) significantly alter sensitivity

and ligand selectivity, highlighting the cooperative role of these loops in shaping receptor

pharmacology. Complementary face mutations in loops D and E (N92T, A158G) further

refined dopamine versus norepinephrine sensitivity. Finally, we also show that this dopamine

receptor can be antagonized by both nicotinic and dopaminergic compounds, underscoring

its hybrid pharmacological profile. Together, these results reveal how subtle changes in

conserved binding motifs can shift receptor specificity from cholinergic to dopaminergic

signaling, providing new insights into the evolutionary flexibility of pLGICs and the molecular

basis of ionotropic catecholamine transmission in invertebrates.

1. Introduction

The processes governing neural communication are dynamic and highly evolutionarily conserved in animals across phyla (Moroz et al., 2021). Relying on the highly orchestrated molecular mechanisms that mediate the release and detection of neurotransmitters, converting extracellular chemical cues into either rapid electrical shifts in cell membrane potential or slower intracellular signaling cascades. Neurotransmission occurs via two distinct pathways: ionotropic and metabotropic. Ionotropic receptors, mediate fast synaptic transmission while in contrast, metabotropic receptors act through slow and second messenger pathways and produce longer lasting, modulatory effects.

The most widespread ionotropic receptors in animals fall into the pentameric or Cys-loop ligand gated ion channel superfamily (pLGIC), which includes nicotinic acetylcholine receptors (nAChR), $GABA_A$ receptors, glycine receptors (GlyR) and serotonin receptors ($5HT_3R$), each typically associated with distinct neurotransmitters and ionic conductances. The structural architecture of pLGICs is highly conserved and consists of five subunits arranged symmetrically around a central ion pore (Figure 1A). Each subunit contributes a β-sandwich extracellular domain (ECD) and four transmembrane α-helices. Neurotransmitter binding occurs at the interfaces between subunits, where one subunit serves as the 'principal face' contributing three extracellular loops A-C and the other contributes loops D-F to form a composite ligand binding site (Figure 1B) (Lynagh and Pless, 2014). While some pLGICs function as homomers most receptors are formed of heteromeric combinations of up to 5 different subunits (Boulin et al., 2008) with the ligand binding interface only forming between specific subunit combinations. Despite this conserved scaffold, structural studies have revealed how small differences in loop composition and conformation underlie substantial diversity in ligand recognition and receptor pharmacology (Corringer et al., 2010; Dalal et al., 2025; Dani, 2015; Pan et al., 2012; Sauguet et al., 2014; Unwin, 2003). This class of receptors have been extensively studied in vertebrate systems, and these receptors have become major pharmacological targets, forming the basis of anesthetics, antiparasitic and anxiolytics. This vertebrate-centric knowledge base has shaped much of our understanding of synaptic transmission. However, studies in invertebrate models have uncovered atypical pLGICs with unexpected ligand specificity, gating properties and synaptic roles (Cully et al., 1996; Ringstad et al., 2009). Characterizing these divergent receptors is essential for understanding the evolutionary diversity and functional plasticity possible in neurotransmission. For example, it was once assumed that each neurotransmitter exerts either an excitatory or an inhibitory effect depending on the receptor class that it activates. However, the discovery of both excitatory and inhibitory glutamate-gated chloride channels (GluCls) in *Caenorhabditis elegans* and *Drosophila melanogaster* challenged this view (Cully

et al., 1996, 1994). These invertebrate-specific channels are the targets for the antiparasitic
treatment ivermectin, which locks the channel in an open state, causing paralysis in parasitic
worms.

One of the most intriguing discoveries in invertebrate systems is the identification of ligand-
gated ion channels gated by dopamine (dop-LGICs), which enable direct, ionotropic
dopaminergic signalling (Courtney et al., 2025; Mitchell et al., 2025; Morud et al., 2021;
Ringstad et al., 2009). This mode of neurotransmission contrasts with canonical
dopaminergic signalling in mammals, which is mediated by GPCRs. The most recently
identified cation permeable dop-LGIC which is gated by the catecholamines dopamine,
norepinephrine and epinephrine has been shown to be conserved in multiple invertebrate
species, including *Daphnia magna, Apis mellifera*, *Aplysia california* and *Octopus sinensis*
(Courtney et al., 2025; Mitchell et al., 2025). In contrast, the anion dop-LGICs studied
previously appear to be nematode specific and gated by dopamine and to a lesser extent
tyramine (Morud et al., 2021; Ringstad et al., 2009). Intriguingly the invertebrate cation dop-
LGIC shares close sequence homology with the vertebrate α9α10 nicotinic acetylcholine
receptor (nAChR), including within the classic ligand binding domains. A great deal of
previous research, including more recent structural studies have revealed residues that
contribute to the binding of cholinergic ligands for nAChRs (Dani, 2015; Morales-Perez et al.,
2016; Su et al., 2025); however, we found that many of these are conserved in cationic dop-
LGICs (Figure 1C). In this study we sought to unravel the contribution of putative ligand
binding domains and individual residues in the homomeric *Daphnia magna* dop-LGIC, Dm-
DopC1, in the sensitivity of cationic dop-LGICs to catecholamines rather than cholinergic
ligands like its close relatives the nAChRs.

Figure 1 A. Full length AlphaFold2 model of Dm-DopC1 with the extracellular (ECD),

transmembrane (TM) and intracellular (ICD) domains highlighted. **B.** Close up of the ECD

and ligand binding domain made up of the interface between two adjacent subunits (light

and dark grey). Extracellular loops associated with ligand binding are labeled as follows;

loop A (orange), loop B (pink), loop C (green), loop D (turquoise), loop E (blue), loop F

(yellow). **C.** Trimmed alignment of the ECD of dop-LGICs, nAChRs and other pLGICs

showing the ligand binding loops. Alignment generated with MAFFT and residues colored

with CLUSTAL coloring.

2. Materials and methods

2.1 Structural Modelling and Sequence Alignments

Amino acid sequences of a range of pLGICs (full sequences are provided in Table S 1) were

aligned using the MAFTT algorithm and visualized with SnapGene (Dotmatics). Structural

models were predicted using AlphaFold ColabFold. For *D. magna* Dm-DopC1, five predicted

models were generated and overlaid in PyMOL. Residue positions were evaluated for

consistency and model confidence (pLDDT > 70). Sites exhibiting conserved structural

placement and high-confidence predictions were considered for site-directed mutagenesis.

2.2 Molecular Biology and Mutagenesis

Mutations were introduced using the Q5® Site-Directed Mutagenesis Kit (New England
Biolabs). Constructs were linearized using NotI. cRNA was synthesized using the T3
mMessage mMachine Kit (ThermoFisher) and purified with the GeneJET RNA Cleanup Kit
(ThermoFisher Scientific).

2.3 *Xenopus laevis* Oocyte Preparation and RNA Injection

Stage V-VI defolliculated *Xenopus laevis* oocytes were obtained from EcoCyte Bioscience
and incubated in ND96 solution (in mM): 96 NaCl, 2 KCl, 1.8 CaCl2, 1 MgCl2, 5 HEPES, pH
7.5. Oocytes were stored at 18°C and injected within two days of delivery. For RNA injection,
oocytes were placed in 96-well plates and injected using the Roboinject system (Multi
Channel Systems, Germany). Each oocyte was injected with 50 nL of cRNA (250ng/µL).
Oocytes were incubated at 18°C for 1 day prior to electrophysiological recordings.

2.4 Two-Electrode Voltage Clamp (TEVC) Recordings

TEVC recordings were performed using the Robocyte2 automated platform (Multi Channel
Systems, Germany). Microelectrodes (0.5-1.5 MΩ resistance) were pulled using a P-97
micropipette puller (Sutter Instrument) and filled with 1 M KCl and 1.5 M acetic acid. Oocytes
were voltage-clamped at -60 mV and recordings were sampled at 500 Hz throughout.
Agonist solutions (dopamine, norepinephrine, epinephrine, tyramine, acetylcholine and
serotonin) were prepared in ND96 on the day of recording and applied at concentrations
ranging from 0.1 µM to 10 mM. Each ligand was perfused for 10 s followed by a 60 s wash.
Wild-type oocytes were included in every recording batch. For antagonist experiment,
compound solutions (50 nM to 500 µM, unless otherwise stated) were prepared in ND96 on
the day of recording with 50 µM dopamine (Dm-DopC1). Before and after compound
application, response to dopamine-only was recorded to obtain an Imax value per oocyte
and to screen for potential irreversible binding of the antagonist. All compounds were
perfused for 7 s followed by a 60 s wash. Perfusion speed was set to 3 ml/min throughout.
Compounds used as antagonists: Acetylcholine chloride (Sigma-Aldrich (A6625)),
Apomorphine hydrochloride (Abcam (ab269887)), Atropine (Abcam (ab145582)),
Isoprenaline hydrochloride (Sigma-Aldrich (I5627)), Levamisole hydrochloride
(MedChemExpress (HY-13666)), (-)-Nicotine detartrate (Abcam (ab120562)), Propranolol
hydrochloride (Santa-Cruz Biotechnology (sc-3580)), (+)-Tubocurarine chloride pentahydrate
(Sigma-Aldrich (93750)).

2.5 Data Analysis

All current amplitudes were exported using the Roboocyte2+ software (Multi Channel Systems) and normalized per oocyte to the maximum response (I/Imax) to account for variation in expression levels using a custom python script adapted from (Morud et al., 2021). Outliers were identified for each construct and removed within GraphPad Prism 10 using the ROUT method (Q=1%) for every individual agonist/antagonist concentration. Agonist and antagonist dose response curves were fitted in GraphPad Prism 10 using a three-parameter logistic equation with a fixed Hill Slope of 1, unless otherwise stated. EC_{50} and IC_{50} values were calculated from the fitted curves and reported as mean $\pm$ standard error of mean (SEM). Statistical significance between wild-type and mutant dose-response curves was assessed using a sum-of-squares F test. Peak current values were extracted using an existing Python script (Morud et al., 2021). Normalized peak currents from the panel of ligands (1mM each: tyramine, acetylcholine, norepinephrine, serotonin and epinephrine) were analyzed using a one-way ANOVA followed by a Fisher's Least Significant Difference (LSD) post hoc test. All TEVC recordings were baseline-subtracted unless otherwise stated. The same pooled wild-type data was used for all comparisons.

2.6 *In silico* prediction of Dm-DopC1 protein in complex with catecholamines and acetylcholine

To predict the possible binding mode of different ligands onto Dm-DopC1 protein (GenBank accession no. XP_032790382.1), we used AlphaFold3-based co-folding approach (Abramson et al., 2024). The N-terminal signal peptide sequence (residues 1-26) was excluded prior to modelling, and the remaining sequence was used to model as homodimer whilst each of the small-molecule ligands (dopamine, norepinephrine, epinephrine and acetylcholine) was incorporated directly into the modelling process using their corresponding Chemical Component Dictionary (CCD) codes. For each ligand, the protein dimer and a single ligand molecule were provided simultaneously as inputs, enabling joint prediction of protein conformation and ligand binding pose within a single inference step. Five independent structures were generated for each protein–ligand combination using an identical protocol to ensure methodological consistency and to account for stochastic variation in model sampling. The best model was considered for each ligand and the corresponding 2D ligand interaction diagram was generated using PoseView™ (Stierand and Rarey, 2010) accessed through the Protein Plus server (https://proteins.plus/).

2.7 Software and Code Availability

TEVC data were analyzed using custom Python scripts (https://github.com/hiris25/TEVC-
analysis-scripts). GraphPad Prism 10 was used for data plotting and fitting. MAFFT (Katoh
and Standley, 2013) was used for sequence alignment, alignments were visualized in
snapgene and ChimeraX 1.9 and Pymol were used for 3D structural visualization. All
constructs and raw data are available upon request.

3. Results

3.1 Despite strong conservation with nAChRs, residues in loop C contribute to
catecholamine sensitivity and selectivity

The principle or α-type subunit contributes three structurally distinct loops to the ligand
binding domain, with loop C forming the front outer side of the binding pocket that closes
around the ligand during receptor activation and loops A and B forming the back of the
binding pocket (Dani, 2015; Morales-Perez et al., 2016). Loop C contains two conserved
structural features in nAChRs. The first is a pair of tyrosines implicated in aromatic cage
formation (Braun et al., 2016; Celie et al., 2004; Tomaselli et al., 1991) and the second is a
double cystine motif (Tomizawa et al., 2008) together forming a 'YXCC' motif. Both features
are present in dop-LGICs and in Dm-DopC1 are represented by Y226 and Y233, and the
C228 - C229, respectively (Figure 2E-F). Notably though both human and chicken nAChR
α9 and α10 subunits contain a glycine at the 'X' position, while the dop-LGICs contain serine
at this position. When we substituted S227 for glycine at this position in Dm-DopC1 we
observed a significant reduction in sensitivity to all catecholamines (Figure 2A-C). The
largest shift was seen with dopamine, shifting 7-fold from a WT EC_{50} of 84μM to 589μM,
norepinephrine EC_{50} also shifted 4-fold and epinephrine 1.8-fold (Figure 2A-C). This
suggests that the native serine may stabilise local interactions required for maintaining the C
loop conformation or supporting neighbouring aromatic residues for catecholamine
recognition.

We were also interested in the residue immediately following the 'YXCC' motif, which is
divergent across species and receptor types. However, both Dm-DopC2 and Am-DopC1
(from *Apis mellifera*), which have a higher relative selectivity for dopamine vs. the other
catecholamines, encode a proline at this position. A substitution to proline in this position in
Dm-DopC1 (E230P) produced a selective decrease in norepinephrine sensitivity only, with a
4-fold shift of EC_{50} from 98μM to 401μM, while maintaining wild type sensitivity for dopamine
and epinephrine (Figure 2A-C). However, despite altering the relative sensitivities, the
E230P mutation did not alter the ligand binding profile of the receptor, with norepinephrine

still eliciting similar relative peak currents to dopamine and epinephrine at 1mM (Figure 2D).

In contrast S227G which altered sensitivity to all catecholamines, abelite with more

pronounced shift in dopamine sensitivity, did alter the ligand binding profile such that

dopamine induced currents at 1mM fell significantly to approximately 50% of the maximum

currents achieved with norepinephrine and epinephrine (Figure 2D). Together these data

suggest that despite key nAChR features of loop C being conserved between cholinergic

and catecholamine binding receptors, the surrounding residues also contribute to ligand

specificity and binding capability.

Figure 2 Mutations in Loop C of a *D. magna* dop-LGIC reduce catecholamine

sensitivity and alter agonist selectivity. A-C. Dose response curves for dopamine (A),

norepinephrine (B) and epinephrine (C) in wild-type, S227G and E230P Dm-DopC1

expressing *Xenopus* oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown

below each plot. **** = $P<0.0001$, * = $P<0.05$, ns = not significant calculated by sum of

squares F test. WT: $n = 24$ (dopamine), 12 (norepinephrine), 11 (epinephrine); S227G: $n = 6$

(dopamine), 3 (norepinephrine), 5 (epinephrine); E230P: $n = 10$ (dopamine), 4

(norepinephrine), 9 (epinephrine). **D.** Relative maximum I/Imax currents of *Xenopus* oocytes

expressing wild-type, S227G and E230P Dm-DopC1, perfused for 10 s with 1 mM each:

dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr),

and acetylcholine (ACh). WT: $n = 18$ (dopamine, norepinephrine, ACh, tyramine), 17

(epinephrine, serotonin); S227G: $n = 11$ per agonist; E230P: $n = 6$ per agonist. ** = $P<0.01$,

ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the

Loop C region. Residues S227 and E230 indicated by arrows, loop C is highlighted by the

green bar. **F.** AlphaFold2 model of the loop C (green) region of Dm-DopC1 with S227 and

E230 relative to Loop B (pink) at the ligand-binding site.

3.2 A conserved aspartate in loop B of principle face is critical to catecholamine
sensitivity

Loop B forms part of the back wall of the ligand binding domains on the principle face and

contains one of the conserved tryptophans forming the aromatic cage which is part of a

conserved region encoded the consensus motif 'SWTY' (Figure 3E-F). A semi conserved

glycine, 4 amino acids downstream of the tryptophan has been implicated in defining high

and low potency nAChRs (Puskar et al., 2012). In the dop-LGICs this post-tryptophan

glycine is broadly conserved and directly preceded by an aspartate which is conserved

across the dop-LGICs but not found in vertebrate α9/10 receptors. Substitution of this

position (D188) in Dm-DopC1 with asparagine, more commonly found in nAChRs, led to a

striking reduction in catecholamine sensitivity, leading to EC_{50} values for all three ligands of

over 9mM, although true EC_{50} values could not be calculated due to lack of saturation even

at 10mM (Figure 3A-C). The D188N mutation also led to significant changes in ligand

binding profiles at 1mM with norepinephrine currents significantly reduced to 8% of the

maximal currents, dopamine reduced to 37% and serotonin induced currents also falling to

19% (Figure 3D). However, it is worth noting that there was no appreciable difference in

acetylcholine currents, which along with the residual currents observed from the

catecholamines suggests that while loop B is critical to catecholamine gating, this single

residue switch is insufficient in defining catecholamine vs cholinergic activation.

We also noted an additional residue of interest downstream of loop B, which differed only in

Dm-DopC1 compared to other dop-LGICs and vertebrate α9/10. Given the differences in

selectivity between Dm-DopC1 and Dm-DopC2 we hypothesised that the conserved

aspartate at this position may also be contributing to ligand selectivity. Mutation at this

location in Dm-DopC1 (N193D), led to a significant increase in EC_{50} for all catecholamines,

most notably for norepinephrine with a 11-fold shift resulting in an EC_{50} of 1087µM, with

smaller yet statistically significant shifts for dopamine and epinephrine of approximately 2-

fold (Figure 3A-C). There was no change in overall ligand binding profile with all 3

catecholamines still eliciting the largest currents (Figure 3D). This suggests that the

conserved aspartate at this position may contribute to selectivity between catecholamines,

with receptors bearing this residue being relatively more selective for dopamine.

Figure 3 Mutations in Loop B of Dm-DopC1 disrupt catecholamine sensitivity and alter ligand selectivity. A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type, D188N and N193D Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001, *** = P<0.001, * = P<0.05 calculated by sum of squares F test. WT: *n* = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); D188N: *n* = 10 (dopamine), 5 (norepinephrine), 9 (epinephrine); N193D: *n* = 15 (dopamine), 5 (norepinephrine), 7 (epinephrine). **D.** Relative maximum I/Imax currents of *Xenopus* oocytes expressing wild-type, D188N and N193D Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: *n* = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); D188N: *n* = 11 per agonist; N193D: *n* = 11 per agonist. **** = P<0.0001, ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the Loop B region. Residues D188 and N193 indicated by arrows, loop B is highlighted by the pink bar. **F.** AlphaFold2 model of the loop B (pink) region of Dm-DopC1 with D188N and N193D relative to Loop C (green) at the ligand-binding site.

3.3 Residues on the complementary face of the binding pocket contribute to dopamine sensitivity and selectivity

In addition to the principal subunit, two loops on the complementary face of the binding pocket have also been shown to be involved in ligand binding in nAChRs. Loop D is composed of a single β strand and forms the base of the binding site, while loop E is

composed of two β strands connected by a short loop. We observed that loop D was broadly

conserved between both nAChRs and dop-LGICs, however we again noted that the two

dop-LGICs with higher dopamine selectivity, Dm-DopC2 and Am-DopC1, encoded a

threonine in the centre of the loop (Figure 4E-F). Creating this change in Dm-DopC1, N92T,

led to a striking and significant reduction of norepinephrine sensitivity, with an EC_{50} increase

from 98μM to 439μM, with no effect on dopamine or epinephrine (Figure 4A-C) and no

changes to overall ligand binding profile at 1mM (Figure 4D).This suggests that the threonine

at this position is important for discerning between norepinephrine and the other

catecholamines.

Despite overall high similarity within loop E, sequence alignments revealed a strongly

conserved aspartate at the proximal end of loop E in α9/10 that is missing from the dop-

LGICs, which frequently encode a hydrophobic leucine or phenylalanine at this position

(Figure 4E-F). When we substituted leucine with aspartate (L155D) at this location in Dm-

DopC1 we could not generate functional receptors that responded consistently to any of the

ligands tested, although due to leak current in oocytes we hypothesise that the protein was

expressed (Figure S 1G-H). This suggests that leucine at this position is critical to either

channel formation or ligand binding. Additionally, we also noted that the two more-dopamine

selective dop-LGICs, Dm-DopC2 and Am-DopC1, had a substitution of an alanine conserved

across both receptor types with a glycine (Figure S 1E-F). Despite being a minor alteration in

sequence we found that this substitution in Dm-DopC1, A158G, led to small but significant

increases in EC_{50} for all catecholamines to similar extents (EC_{50}s: dopamine: 235μM,

norepinephrine: 440μM, epinephrine: 438μM) (Figure S 1A-D). Again, supporting the notion

that residues surrounding key ligand binding loops contribute to receptor function, likely

through the stabilisation of these regions.

Figure 4 The N92T mutation in Loop D of a _D. magna_ dop-LGIC reduces norepinephrine sensitivity but preserves agonist selectivity. A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type and N92T Dm-DopC1 expressing _Xenopus_ oocytes. Current normalized by I/Imax. EC$_{50}$ values (± SEM) shown below each plot. **** = P<0.0001, ns = not significant calculated by sum of squares F test. WT: n = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); N92T: n = 10 (dopamine), 5 (norepinephrine), 8 (epinephrine). **D.** Relative maximum I/Imax currents of _Xenopus_ oocytes expressing wild-type and N92T Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: n = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); N92T: n = 11 per agonist. ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the loop D region. Residue N92 is indicated by an arrow, loop D is highlighted by the turquoise bar. **F.** AlphaFold2 model of the loop D (turquoise) region of Dm-DopC1 with N92 relative to loop B (pink), loop C (green), loop F (yellow), loop A (orange) and loop E (blue) at the ligand-binding site.

3.4 The conserved proline of loop A enhances catecholamine sensitivity

In line with previous findings, our sequence alignments identified a proline residue conserved across all pLGICs that sits within the conserved 'WXPD' motif (Figure 5E-F) (Braun et al., 2016). However, in Dm-DopC1 this location encodes a serine, which AlphaFold2 structural modelling placed near the inter-subunit ligand-binding interface, in proximity to Loops B and C (Figure 5F). Strikingly, restoring the proline at this position in

Dm-DopC1 (S123P) significantly increased sensitivity to all catecholamines (Figure 5A-C).
Dopamine EC_{50} decreased from 84.82 ± 9.1 µM to 39.84 ± 6.06 µM, norepinephrine
from 98.78 ± 14.09 µM to 36.52 ± 7.86 µM, and epinephrine from
169.17 ± 29.5 µM to 47.87 ± 14.05 µM (Figure 5A-C). Interestingly, unlike the other
mutations examined in this study, S123P also led to an increase in Hill slope for
norepinephrine (2.16 ± 0.19) compared to WT (1.31 ± 0.30), suggesting enhanced co-
operativity (Figure S 2). Overall, the relative ligand binding profile was unaltered, although
there was a small but non-significant increase in relative acetylcholine current (Figure 5D).

Figure 5 The conserved proline of loop A is required for ligand sensitivity in dop-
LGICs. A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine
(C) in wild-type and S123P Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by
I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001 calculated by sum of
squares F test. WT: *n* = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); S123P: n =
12, (dopamine), 4 (norepinephrine) and 8 (epinephrine). **D.** Relative maximum I/Imax
currents of *Xenopus* oocytes expressing wild-type and A158G Dm-DopC1, perfused for 10 s
with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT),
tyramine (Tyr), and acetylcholine (ACh). WT: *n* = 18 (dopamine, norepinephrine, ACh,
tyramine), 17 (epinephrine, serotonin); S123P: n = 12 per agonist. ns = not significant by
one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the loop A region. Residue
S123 is indicated by an arrow, loop A is highlighted by the orange bar. **F.** AlphaFold2 model

of the loop A (orange) region of Dm-DopC1 with S123 relative to loop B (pink), loop C

(green) and loop E (blue) at the ligand-binding site.

3.5 Cholinergic, dopaminergic and adrenergic compounds antagonise dop-LGICs

To further dissect the pharmacology of the cationic dop-LGICs we tested a range of

cholinergic, dopaminergic and adrenergic compounds for their ability to activate or

antagonise Dm-DopC1. We found that both Dm-DopC1 dopamine induced currents could be

blocked by the nicotinic antagonist tubocurarine, with an IC_{50} of 0.9µM (Figure 6A). In

contrast to other nAChRs, the vertebrate a9/10 nAChRs are antagonised by nicotine

(Elgoyhen et al., 2001, 1994). In line with this we found that the dopamine currents of Dm-

DopC1 could be blocked at high concentrations of nicotine with and IC_{50} of 109µM (Figure

6A). We also tested the muscarinic blocker, atropine, which has been shown to act on

nAChRs (Parker et al., 2003; Zwart and Vijverberg, 1997), for its ability to block dopamine

currents and found that it behaved similarly to nicotine with relatively poor blocking (IC_{50}:

101µM (Figure 6A). This family of dop-LGICs have also been shown to be minimally

activated by acetylcholine (Courtney et al., 2025), and due to their close homology with

nAChRs and antagonism by nicotine we sought to determine if acetylcholine is able to

interact with the dopamine induced current, however we saw no evidence of this (Figure 6A).

In addition, we also tested apomorphine, a non-selective vertebrate dopamine receptor

agonist and the β-adrenoceptor agonist and antagonist, isoprenaline and propranolol

respectively. Apomorphine was able to block the dopamine induced current of Dm-DopC1,

with an IC_{50} of 6µM (Figure 6B). Meanwhile propranolol but not isoprenaline was able to

block dopamine currents (IC_{50}: 100µM) (Figure 6B). We also tested the ability of these

compounds to activate the receptors and observed no activation at 100µM (Figure S 3).

Figure 6 Dm-DopC1 is sensitive to antagonism by cholinergic and dopaminergic

compounds. **A-B.** Inhibitory dose response curves for a panel of cholinergic (A) and

dopaminergic and adrenergic compounds (B). Error bars represent SEM. $n = 4$

(apomorphine, isoprenaline), 6 (atropine, propranolol), 9 (tubocurarine, acetylcholine), 12

(nicotine). IC50s are shown below each plot and calculated with a 4-parameter curve with

constrained bottom = 0, nd = not determined.

3.6 *In silico* ligand binding predicts involvement of conserved aromatic cage

residues and loop E in ligand recognition

To further predict the binding modality of catecholamines in this novel group of LGICs we

employed structural modelling using an AlphaFold3-based co-folding approach (Abramson

et al., 2024) in which Dm-DopC1 dimers were co-folded with catecholamines and

acetylcholine, 2D interaction predictions were then generated using PoseView (Stierand and

Rarey, 2010) (Figure 7). As anticipated in all cases the ligand was predicted to bind within

the conserved ligand binding domain composed of the interface between the two adjacent

subunits. We also observed that catecholamine binding was predicted to involve many of the

highly conserved aromatic cage residues, usually associated with nAChR recognition of

acetylcholine. In line with our pharmacological data, dopamine was predicted to have the

most direct interactions with amino acids in the binding pocket, including at least one residue

from each of the conserved loops (Figure 7A-B). In addition to the aromatic residue

interactions dopamine was also predicted to form hydrophobic interactions with L155 of loop

E and hydrogen bond with D202 of loop F (Figure 7B). Norepinephrine and epinephrine

appeared to have fewer direct contacts, again in line with pharmacological observations in

which these ligands have lower receptor sensitivity, however epinephrine maintained an

interaction with L155 of loop E (Figure 7C-F). Meanwhile acetylcholine was also predicted to

bind in a similar position within the binding pocket however no direct contacts could be

predicted (Figure S4).

**Figure 7 *In silico* binding of catecholamines with Dm-DopC1 predicts involvement of
conserved aromatic residues. A, C & E.** Predicted AlphaFold co-folded models of Dm-

DopC1 in complex with dopamine (A), norepinephrine (C) and epinepheline (E). Ligands are

shown in beige with polar contacts between the ligand and adjacent residues shown by

dashed yellow lines. Colors represent conserved ligand binding loops: A: orange, B: pink, C:

green, D: teal, E: blue, F: yellow. Amino acid side chains are shown for residues which

appear in 2D poses, as well as S227 which forms a hydrogen bond with dopamine and

epinephrine. **B, D & F.** 2D ligand poses for dopamine (B), norepinephrine (D) and

epinepheline (F) showing predicted interactions between the ligand and adjacent residues.

4. Discussion

Despite high levels of conservation of key ligand binding motifs with vertebrate and
invertebrate nAChRs this new class of dopamine-gated cation channels displays striking
differences in ligand sensitivity and antagonism by a range of compounds. In this study we
highlight some of the key residues that likely contribute to this functional difference (Table 1),
with many surrounding previously defined key motifs for cholinergic receptors.

Table 1 **Summary of Ec50 values for all tested mutations.** All values are given in µM and
calculated using a 3-parameter curve fit with a Hill slope of 1. P values were calculated using
the sum of squares F test.

		Dopamine			Norepinephrine			Epinephrine		
		E_C50	SEM	P value	E_C50	SEM	P value	E_C50	SEM	P value
Loop	Construct									
	WT	84.8	8.9	N/A	98.8	14.1	N/A	169.2	29.5	N/A
A	S123P	39.8	6.1	<0.0001	36.5	7.9	<0.0001	47.9	14.05	<0.0001
B	D188N	9466.9	3110	<0.0001	Unstable	Unstable	<0.0001	13153.8	2861.8	<0.0001
	N193D	172.6	31.1	0.0003	1087.9	190.6	<0.0001	384.8	109.9	0.0163
C	S227G	589.1	92.2	<0.0001	394.4	186.6	<0.0001	307.8	33.5	0.0262

	E230P	119.2	22.6	0.0812	401.1	73.6	<0.0001	166.2	34.8	0.9494
D	N92T	90.5	18.1	0.7445	439.8	114.1	<0.0001	170.2	57	0.9868
E	L155D	-	-	-	-	-	-	-	-	-
	A158G	234.6	30.7	<0.0001	439.8	57.5	<0.0001	438.2	57.97	<0.0001

The most striking effect was the almost complete loss of catecholamine gating upon
replacement of an aspartate in loop B (D188N) which follows the conserved 'SWTY' motif
and which is conserved in dop-LGICs. To our knowledge, this site, which in vertebrate α9/10
encodes an asparagine or glycine, has not been previously directly implicated in ligand
binding for nAChRs. However, it is unlikely that this residue alone contributes to dopamine
selectivity over cholinergic ligands given that aspartate is also found at this position in classic
nAChRs such as human α4 and the lack of significant acetylcholine gating over the wild type
for this mutation in Dm-DopC1. Our *in silico* ligand binding models also place D188 outside
of the core ligand binding residues (Figure S 4), suggesting that it may contributing
structurally to the placement of loop B. Interestingly, though immediately following this
aspartate is a semi conserved glycine residue which has been shown to distinguish high
affinity vertebrate nAChRs (eg. α4 which encodes lysine at this position) from low affinity
receptors (eg. α7 which encodes glycine). The increased affinity is thought to be due to
stabilization of the open and desensitized states by virtue of a hydrogen bond between this
position in loop B and a proline in loop C that forms in absence of glycine (Grutter et al.,
2003; Kouvatsos et al., 2016; Puskar et al., 2012). Interestingly glycine is present at this
position in all but one dop-LGIC (which instead encodes an alanine) further supporting the
hypothesis that this hydrogen bond is key to high affinity acetylcholine receptors. Decoupling
of loops B and C may be important for gating by aminergic ligands given that monoamine
gated receptors from other subfamilies such as the vertebrate $5HT_3R$ and invertebrate
chloride permeable dopamine and histamine receptors lack the hydrogen forming proline
within loop C. In addition, loop B also contains one of the conserved tryptophans associated
with aromatic cage formation. Preceding this tryptophan by 2 amino acids is another glycine
residue conserved in nAChRs α subunits but encoding an arginine in β subunits, which
causes movements in the tryptophan preventing it from forming a functional binding site
(Morales-Perez et al., 2016), in dop-LGICs this site encodes a alanine or serine. Mutations in
loop B have also been shown to be more broadly required for monoamine binding in $5HT_3Rs$

for example (Thompson et al., 2008). It would be interesting to see if the D188N reverse

mutation in vertebrate receptors alters acetylcholine gating, although given the divergence of

sequences at this position across nAChRs in both vertebrates and invertebrates this residue

may not be contributing in a significant way to acetylcholine recognition.

We also confirmed the importance, yet surprising flexibility, of a highly conserved proline

residue within the 'WXPD' motif of loop A, which has been hypothesized to be critical for

maintaining structural integrity and receptor assembly in pLGIC receptors from bacteria

through to humans (Braun et al., 2016). We observed that Dm-DopC1, which exhibits a

higher EC_{50} for dopamine than other dop-LGICs (Courtney et al., 2025), diverged from this

motif to encode serine at this position (WKSD), which when substituted for proline, led to

significant increases in catecholamine sensitivity. The mirrors previous work in glycine

receptors where the same proline was replaced by an alanine leading to a decrease in

ligand sensitivity (Hughes et al., 2020; Lummis and Dougherty, 2022). Not only does our

finding further highlight the importance of a proline at this position, but it also shows that

there is flexibility in the motif, as the natural variant still displays micromolar ligand binding.

As this serine is not conserved within the rest of the dop-LGIC group, it is unlikely to be

directly participating in catecholamine recognition. Rather this suggests, as has been

previously hypothesized, that the conserved proline of the 'WXPD' motif contributes

indirectly by stabilizing loop A conformation or facilitating ligand alignment within the binding

pocket.

An intriguing feature of the invertebrate cationic dop-LGICs is the presence of key motifs

usually attributed to acetylcholine and nicotine binding in nAChRs. This is highlighted by our

in silico binding experiments which suggest that many of the key residues involved in

aromatic cage formation, usually associated with acetylcholine binding participate in the

binding of dopamine and the other catecholamines. In particular this can be seen in loop C,

which in dop-LGICs also encodes the conserved 'YXCC' motif (Karlin, 2002; Tomizawa et

al., 2008) as well as a pair of tyrosines implicated in aromatic cage formation (Braun et al.,

2016). However, the at 'X' position of the 'YXCC' motif the dop-LGICs encode a conserved

serine, which is not present in nAChRs. By exchanging this serine for glycine (present in

α9/10) in Dm-DopC1, we found that this serine is indeed important for catecholamine gating

but did not confer acetylcholine gating despite restoring the motif often found in nAChRs.

Interestingly our binding pose predictions did not identify direct interactions of the ligands

with this serine, however for both dopamine and epinephrine a hydrogen bond between

serine and the ligand was predicted. Separately, we also found that the residue immediately

downstream of the 'YXCC' motif was important for determining differences in ligand

sensitivity between *Daphnia* dop-LGICs. The functional impact of mutations at these
positions supports the established role of loop C in maintaining the geometry of the ligand
binding pocket, particularly in orienting conserved aromatic residues that contribute to
catecholamine recognition. It also echoes our findings within loop B, which suggests that a
series of changes in the binding regions are required to completely alter receptor selectivity
from cholinergic to dopaminergic.

We also explored the role of loops D, E and F that form the complimentary face of the ligand
binding pocket and have been shown to contribute to stabilizing the extracellular domain. A
key feature of Loop D is the 'WXD' motif that is conserved across nAChRs, dop-LGICs,
$5HT_3Rs$ and even some chloride channels (Braun et al., 2016; Corringer et al., 1995; Spier
and Lummis, 2000). Loop D is remarkably well conserved between dop-LGICs and nAChRs,
suggesting that its main function is to stabilize the ligand binding regions. However, we did
note that both Dm-DopC2 and Am-DopC1, which are more selective for dopamine than the
other dop-LGICs encoded a threonine 2 amino acids before the 'WXD' motif, rather than an
asparagine. When we substituted this asparagine for threonine in Dm-DopC1 we noted a
specific alteration in ligand sensitivity with only the norepinephrine EC_{50} being adversely
affected while dopamine and epinephrine responses remained like wild type, making the
receptor overall more selective for dopamine. Interestingly, vertebrate nAChRs typically
encode a lysine or arginine at this position which is also conserved across other pLGICs
including glycine and GABA receptors, where disruption of this residue leads to almost
complete loss of ligand activation (Goldschen-Ohm et al., 2011; Grudzinska et al., 2005).
The cationic dop-LGICs, in line with hypotheses drawn up by Lynagh and Pless in 2014, also
lack positively charged residues in this position instead favoring polar residues also found in
other amine gated receptors including HisCl and the anionic dop-LGICs from nematodes,
which may assist in binding the hydroxyl termini or these ligands (Lynagh and Pless, 2014).
In addition, our *in silico* model also predicted direct hydrophobic interactions between L155
of loop E and dopamine and epinephrine. Hydrophobic residues at this position are
conserved across dop-LGICs in contrast to vertebrate α9/10 nAChRs which encode a
charged aspartate at this position. Interestingly when we exchanged leucine for aspartate at
this position, we could not generate significant dose dependent ligand-induced currents for
any of the catecholamines nor acetylcholine. This is mirrored by a previous study of Am-
DopC1 which identified the adjacent serine residue, which is conserved across dop-LGICs,
as critical for dopamine binding (Mitchell et al., 2025). Taken together these data suggest
this region of loop B is important for ligand recognition in dop-LGICs. Collectively these
mutations have identified residues critical for catecholamine recognition across various
conserved ligand binding domains, revealing that like in other pLGICs, both faces of the

ligand binding domain are critical not only for ligand gating but also relative ligand selectivity.
However, this study also demonstrates that ligand switching from acetylcholine to (or from)
dopamine requires a series of alterations across evolutionary time and across multiple
regions of the binding domain.

Finally, we also examined the pharmacology of dop-LGICs by testing a range of cholinergic,
dopaminergic and adrenergic compounds for their ability to interact with the dopamine
induced currents from Dm-DopC1. The competitive nAChR antagonist, tubocurarine
(Rahman et al., 2022), efficiently blocked currents in the low µM range, broadly in line with
classic nAChRs (Zwart and Vijverberg, 1998). In contrast, other cholinergic compounds such
as nicotine and atropine (a muscarinic antagonist) had effects only at high concentrations.
This is in line with evidence that tubocurarine can act on a broader range of pLGICs
including vertebrate 5HT$_3$Rs (Peters et al., 1990; Yan et al., 1998) and invertebrate chloride
gated cholinergic receptors (Hardege et al., 2023). In addition, dop-LGICs could be blocked
by apomorphine (non-selective vertebrate dopamine agonist) and propranolol (β-
adrenoceptor antagonist) in the µM range suggesting that dop-LGICs have a distinct
pharmacology from their closely related nAChRs, with the potential of novel chemistries to
be used to target this subfamily.

5. Conclusion

In this study we detail the mechanistic map of ligand selectivity of the invertebrate dopamine-
gated cation channel (dop-LGIC) family, extending the known functional repertoire of these
receptors. Our primary aim was to identify how variations in conserved ligand binding
domains enable dop-LGICs to respond selectively to catecholamines rather than cholinergic
ligands, despite their close sequence homology with vertebrate nAChRs. In line with this we
found that while core structural features of the pLGIC superfamily are conserved, functional
divergence is driven by key residue variations surrounding these canonical motifs in the
extracellular loops. We identified residues critical for catecholamine recognition across loops
typically associated with both the principal (A-C) and complementary faces (D-F) of the
ligand binding domain. Pharmacological analysis of dop-LGICs also confirmed a distinct
profile from their nAChR relatives, with dopamine-induced currents efficiently blocked in the
micromolar range by cholinergic, dopaminergic and adrenergic compounds.

Our results provide insight into the functional plasticity within invertebrate neurotransmission
systems and reveal how subtle differences in loop composition enable the switch from
cholinergic to aminergic ligand recognition. The fact that cationic dop-LGICs are conserved
across multiple invertebrate species, confirms the widespread nature of this signaling

mechanism. Furthermore, the demonstrated requirement for a series of evolutionary
changes to achieve ligand switching raises fundamental questions about the origin of
ionotropic neurotransmission, specifically whether dopamine or acetylcholine signaling
emerged first. Future research should focus on comparative structural analysis of dop-LGICs
across basal phyla, such as Platyhelminths or Nematodes (where unique anionic dop-LGICs
are already found), to trace the evolutionary trajectory and structural requirements that drove
the transition between cholinergic and dopaminergic ionotropic signaling.

Funding

I.H, T.M.M, T.E.R. and this work was kindly supported by a Royal Society University
Research Fellowship (URF\R1\231648). Y.Z. was supported kindly by the Percy Lander
Studentship from Downing College, Cambridge.

Acknowledgements

The authors kindly thank Amy Courtney & William Schafer for providing the plasmid
containing wild type Dm-DopC1. The authors also gratefully acknowledge Howard Baylis
and all present and past members of the Hardege lab for helpful discussions.

Author Contributions

T. M. M: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data
Curation, Writing - Original Draft, Visualization.

T. E. R: Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing -
Review & Editing.

Y. Z: Formal analysis, Investigation, Data Curation.

T. R: Supervision, Project administration.

I. H: Conceptualization, Formal analysis, Resources, Data Curation, Writing - Original Draft,
Visualization, Supervision, Project administration, Funding acquisition.

Declaration of interest

The authors declare no competing interests.

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,
Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.-C.,
O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E.,
Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers,
A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R., Khan, Y.A., Low,
C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram,
A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Žídek, A., Bapst, V., Kohli, P.,
Jaderberg, M., Hassabis, D., Jumper, J.M., 2024. Accurate structure prediction of
biomolecular interactions with AlphaFold 3. Nature 630, 493–500.
https://doi.org/10.1038/s41586-024-07487-w
Boulin, T., Gielen, M., Richmond, J.E., Williams, D.C., Paoletti, P., Bessereau, J.-L., 2008.
Eight genes are required for functional reconstitution of the Caenorhabditis elegans
levamisole-sensitive acetylcholine receptor. Proceedings of the National Academy of
Sciences 105, 18590–18595. https://doi.org/10.1073/pnas.0806933105
Braun, N., Lynagh, T., Yu, R., Biggin, P.C., Pless, S.A., 2016. Role of an Absolutely
Conserved Tryptophan Pair in the Extracellular Domain of Cys-Loop Receptors. ACS
Chem. Neurosci. 7, 339–348. https://doi.org/10.1021/acschemneuro.5b00298
Celie, P.H.N., van Rossum-Fikkert, S.E., van Dijk, W.J., Brejc, K., Smit, A.B., Sixma, T.K.,
2004. Nicotine and carbamylcholine binding to nicotinic acetylcholine receptors as
studied in AChBP crystal structures. Neuron 41, 907–914.
https://doi.org/10.1016/s0896-6273(04)00115-1
Corringer, P.-J., Baaden, M., Bocquet, N., Delarue, M., Dufresne, V., Nury, H., Prevost, M.,
Van Renterghem, C., 2010. Atomic structure and dynamics of pentameric ligand-
gated ion channels: new insight from bacterial homologues. J Physiol 588, 565–572.
https://doi.org/10.1113/jphysiol.2009.183160
Corringer, P.J., Galzi, J.L., Eiselé, J.L., Bertrand, S., Changeux, J.P., Bertrand, D., 1995.
Identification of a new component of the agonist binding site of the nicotinic alpha 7
homooligomeric receptor. J Biol Chem 270, 11749–11752.
https://doi.org/10.1074/jbc.270.20.11749
Courtney, A., Styfhals, R., Dijck, M.V., Boulanger, J., Geenen, L., Lanoizelet, M., Hardege,
I., Lassnig, M., Jansson, A.M., Gehler-Rahman, J., Almansa, E., Obenhaus, H.A.,
Seuntjens, E., Schafer, W.R., 2025. Novel dopaminergic neurotransmission in the
Octopus visual system. https://doi.org/10.1101/2025.04.02.646353
Cully, D.F., Paress, P.S., Liu, K.K., Schaeffer, J.M., Arena, J.P., 1996. Identification of a
Drosophila melanogaster glutamate-gated chloride channel sensitive to the
antiparasitic agent avermectin. J Biol Chem 271, 20187–20191.
https://doi.org/10.1074/jbc.271.33.20187
Cully, D.F., Vassilatis, D.K., Liu, K.K., Paress, P.S., Van der Ploeg, L.H., Schaeffer, J.M.,
Arena, J.P., 1994. Cloning of an avermectin-sensitive glutamate-gated chloride
channel from Caenorhabditis elegans. Nature 371, 707–711.
https://doi.org/10.1038/371707a0
Dalal, V., Tan, B.K., Xu, H., Cheng, W.W., 2025. Cryo-EM structures of a pentameric ligand-
gated ion channel in liposomes. eLife 14, RP106728.
https://doi.org/10.7554/eLife.106728
Dani, J.A., 2015. Neuronal Nicotinic Acetylcholine Receptor Structure and Function and
Response to Nicotine. Int Rev Neurobiol 124, 3–19.
https://doi.org/10.1016/bs.irn.2015.07.001

Elgoyhen, A.B., Johnson, D.S., Boulter, J., Vetter, D.E., Heinemann, S., 1994. α9: An
acetylcholine receptor with novel pharmacological properties expressed in rat
cochlear hair cells. Cell 79, 705–715. https://doi.org/10.1016/0092-8674(94)90555-X
Elgoyhen, A.B., Vetter, D.E., Katz, E., Rothlin, C.V., Heinemann, S.F., Boulter, J., 2001. α10:
A determinant of nicotinic cholinergic receptor function in mammalian vestibular and
cochlear mechanosensory hair cells. Proceedings of the National Academy of
Sciences 98, 3501–3506. https://doi.org/10.1073/pnas.051622798
Goldschen-Ohm, M.P., Wagner, D.A., Jones, M.V., 2011. Three Arginines in the GABAA
Receptor Binding Pocket Have Distinct Roles in the Formation and Stability of
Agonist- versus Antagonist-Bound Complexes. Molecular Pharmacology 80, 647–
656. https://doi.org/10.1124/mol.111.072033
Grudzinska, J., Schemm, R., Haeger, S., Nicke, A., Schmalzing, G., Betz, H., Laube, B.,
2005. The β Subunit Determines the Ligand Binding Properties of Synaptic Glycine
Receptors. Neuron 45, 727–739. https://doi.org/10.1016/j.neuron.2005.01.028
Grutter, T., Prado de Carvalho, L., Le Novère, N., Corringer, P.J., Edelstein, S., Changeux,
J., 2003. An H-bond between two residues from different loops of the acetylcholine
binding site contributes to the activation mechanism of nicotinic receptors. The
EMBO Journal 22, 1990–2003. https://doi.org/10.1093/emboj/cdg197
Hardege, I., Morud, J., Courtney, A., Schafer, W.R., 2023. A Novel and Functionally Diverse
Class of Acetylcholine-Gated Ion Channels. J Neurosci 43, 1111–1124.
https://doi.org/10.1523/JNEUROSCI.1516-22.2022
Hughes, M.E., Mesoy, S.M., Capes, E., Lummis, S.C.R., 2020. Many Proline Residues in the
Extracellular Domain Contribute to Glycine Receptor Function. ACS Chem. Neurosci.
11, 2658–2665. https://doi.org/10.1021/acschemneuro.0c00320
Karlin, A., 2002. Emerging structure of the Nicotinic Acetylcholine receptors. Nat Rev
Neurosci 3, 102–114. https://doi.org/10.1038/nrn731
Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol 30, 772–780.
https://doi.org/10.1093/molbev/mst010
Kouvatsos, N., Giastas, P., Chroni-Tzartou, D., Poulopoulou, C., Tzartos, S.J., 2016. Crystal
structure of a human neuronal nAChR extracellular domain in pentameric assembly:
Ligand-bound α2 homopentamer. Proceedings of the National Academy of Sciences
113, 9635–9640. https://doi.org/10.1073/pnas.1602619113
Lummis, S.C.R., Dougherty, D.A., 2022. Expression of Mutant Glycine Receptors in
Xenopus Oocytes Using Canonical and Non-Canonical Amino Acids Reveals Distinct
Roles of Conserved Proline Residues. Membranes 12, 1012.
https://doi.org/10.3390/membranes12101012
Lynagh, T., Pless, S.A., 2014. Principles of agonist recognition in Cys-loop receptors. Front.
Physiol. 5. https://doi.org/10.3389/fphys.2014.00160
Mitchell, E.L., Armstrong, E.B., Viscarra, F., Bermudez, I., Biggin, P.C., Goodchild, J.A.,
Jones, A.K., 2025. Characterisation of an unusual nicotinic acetylcholine receptor
subtype preferentially sensitive to biogenic amines. Commun Biol 8, 1764.
https://doi.org/10.1038/s42003-025-09143-z
Morales-Perez, C.L., Noviello, C.M., Hibbs, R.E., 2016. X-ray structure of the human α4β2
nicotinic receptor. Nature 538, 411–415. https://doi.org/10.1038/nature19785
Moroz, L.L., Romanova, D.Y., Kohn, A.B., 2021. Neural versus alternative integrative
systems: molecular insights into origins of neurotransmitters. Philosophical
Transactions of the Royal Society B: Biological Sciences 376, 20190762.
https://doi.org/10.1098/rstb.2019.0762
Morud, J., Hardege, I., Liu, H., Wu, T., Choi, M.-K., Basu, S., Zhang, Y., Schafer, W.R.,
2021. Deorphanization of novel biogenic amine-gated ion channels identifies a new
serotonin receptor for learning. Curr Biol 31, 4282-4292.e6.
https://doi.org/10.1016/j.cub.2021.07.036
Pan, J., Chen, Q., Willenbring, D., Yoshida, K., Tillman, T., Kashlan, O.B., Cohen, A., Kong,
X.-P., Xu, Y., Tang, P., 2012. Structure of the pentameric ligand-gated ion channel

ELIC cocrystallized with its competitive antagonist acetylcholine. Nat Commun 3,
714. https://doi.org/10.1038/ncomms1703
Parker, J.C., Sarkar, D., Quick, M.W., Lester, R.A.J., 2003. Interactions of atropine with
heterologously expressed and native α3 subunit-containing nicotinic acetylcholine
receptors. Br J Pharmacol 138, 801–810. https://doi.org/10.1038/sj.bjp.0705124
Peters, J.A., Malone, H.M., Lambert, J.J., 1990. Antagonism of 5-HT3 receptor mediated
currents in murine N1E-115 neuroblastoma cells by (+)-tubocurarine. Neuroscience
Letters 110, 107–112. https://doi.org/10.1016/0304-3940(90)90796-C
Puskar, N.L., Lester, H.A., Dougherty, D.A., 2012. Probing the Effects of Residues Located
Outside the Agonist Binding Site on Drug-Receptor Selectivity in the Nicotinic
Receptor. ACS Chem Biol 7, 841–846. https://doi.org/10.1021/cb200448j
Rahman, M.M., Basta, T., Teng, J., Lee, M., Worrell, B.T., Stowell, M.H.B., Hibbs, R.E.,
2022. Structural mechanism of muscle nicotinic receptor desensitization and block by
curare. Nat Struct Mol Biol 29, 386–394. https://doi.org/10.1038/s41594-022-00737-3
Ringstad, N., Abe, N., Horvitz, H.R., 2009. Ligand-gated chloride channels are receptors for
biogenic amines in C. elegans. Science 325, 96–100.
https://doi.org/10.1126/science.1169243
Sauguet, L., Shahsavar, A., Poitevin, F., Huon, C., Menny, A., Nemecz, À., Haouz, A.,
Changeux, J.-P., Corringer, P.-J., Delarue, M., 2014. Crystal structures of a
pentameric ligand-gated ion channel provide a mechanism for activation.
Proceedings of the National Academy of Sciences 111, 966–971.
https://doi.org/10.1073/pnas.1314997111
Spier, A.D., Lummis, S.C.R., 2000. The Role of Tryptophan Residues in the 5-
Hydroxytryptamine3 Receptor Ligand Binding Domain*. Journal of Biological
Chemistry 275, 5620–5625. https://doi.org/10.1074/jbc.275.8.5620
Stierand, K., Rarey, M., 2010. Drawing the PDB: Protein–Ligand Complexes in Two
Dimensions. ACS Med. Chem. Lett. 1, 540–545. https://doi.org/10.1021/ml100164p
Su, J., Yu, Z., Yin, Z., Zhang, Z., Zhao, J., Meng, Y., Li, R., Gao, Y., Zhang, H., Yu, R., Zhao,
Y., 2025. Molecular insights into the α6β4 nicotinic acetylcholine receptor function
and ligand recognition. Nat Commun 16, 3153. https://doi.org/10.1038/s41467-025-
58333-0
Thompson, A.J., Lochner, M., Lummis, S.C.R., 2008. Loop B Is a Major Structural
Component of the 5-HT3 Receptor. Biophysical Journal 95, 5728–5736.
https://doi.org/10.1529/biophysj.108.135624
Tomaselli, G.F., McLaughlin, J.T., Jurman, M.E., Hawrot, E., Yellen, G., 1991. Mutations
affecting agonist sensitivity of the nicotinic acetylcholine receptor. Biophysical Journal
60, 721–727. https://doi.org/10.1016/S0006-3495(91)82102-6
Tomizawa, M., Maltby, D., Talley, T.T., Durkin, K.A., Medzihradszky, K.F., Burlingame, A.L.,
Taylor, P., Casida, J.E., 2008. Atypical nicotinic agonist bound conformations
conferring subtype selectivity. Proc Natl Acad Sci U S A 105, 1728–1732.
https://doi.org/10.1073/pnas.0711724105
Unwin, N., 2003. Structure and action of the nicotinic acetylcholine receptor explored by
electron microscopy. FEBS Lett 555, 91–95. https://doi.org/10.1016/s0014-
5793(03)01084-6
Yan, D., Pedersen, S.E., White, M.M., 1998. Interaction of D-tubocurarine analogs with the
5HT3 receptor. Neuropharmacology 37, 251–257. https://doi.org/10.1016/s0028-
3908(98)00010-0
Zwart, R., Vijverberg, H.P.M., 1998. Four Pharmacologically Distinct Subtypes of α4β2
Nicotinic Acetylcholine Receptor Expressed in Xenopus laevisOocytes. Molecular
Pharmacology 54, 1124–1131. https://doi.org/10.1124/mol.54.6.1124
Zwart, R., Vijverberg, H.P.M., 1997. Potentiation and Inhibition of Neuronal Nicotinic
Receptors by Atropine: Competitive and Noncompetitive Effects. Molecular
Pharmacology 52, 886–895. https://doi.org/10.1124/mol.52.5.886
738
739

740 **Supplementary data**

741 Table S 1 Protein sequences used for alignments.

Name	Protein Sequence
O_sinensis_ DopC1	MNYRLYIAMTLMKDIRILMSAFHLLMGASIALCDENEYRLVRDLMKNYNKQV RPSSNFSEALNVTFGVALAQIIDVDEKDQIITTNCWINQNWIDYKLQWDPIKY GNVSVIRLPFSDIWRPDILLYNNADVKAFESSISTNVIVTSEGNVTWLSMVIFK SSCSIDVKFFPFDEQNCSMEFASWTYDAYQVNILTNGEDNGDMSNYIENSE WSLIGFQQKRHVVRFSCCEEPYIIIHYHILIKRRPLFYLFNMVMPCILITLVALLG FYMPSDSGEKISMGITTLLSMTVFLMIVAEKMPPTSDVLPLIGLYYGITIAIVSF ATGMTVLTLNIHHKGTRGYEVPMILKKICFGVVAKILFIKLEVPLGSKESMPAA PVADYYRRFDTDKLPENGRLSPRFIRKLNPSASATASETFERQFMRVLQKVY QTIERNEMRLAEQDRRDLIKQEWQQLALIIDRVLLLIFLGLTITVTLTVFFQGP
O_sinensis_ DopC2	MEFWASITLTIFSLCLPSSSADDVELSLVNRLLKNYDKRVRPSLNSSEAINVTF GVALAQIIDVDERNQILTTNCWLNQVWLDVNLRWDPDKYNGIKVVRIPFTDV WKPDILLYNNADVTSYFSSLSSNVIVTSEGNVTWLSMVIFKSSCRIDVRYFPF DEQNCSMQFASWTYDGFQLQLRMNSEEGDISNYIRNSEWVLIKLHVEWNLV YYSCCIEPYPDITYYIQIRRRPLFYVFNMILPCILITLVALLGFYIPSDSGEKVTM GITTLLSMTVFMMLVAENMPPTSDVLPLIGIYYGITISIVSFATAMTVFTLNIHH KGARGHEVPIIIKKICFGILAKIFCMRLEKPNNKSSNVAHLSDMSSSSEIPEMF VPCSSQSNSSYIRFKLSSQEESDQNAKPLLEMNPTLSQQQSMIRETRFHGD LETSNGGILSPRFSRLRGLQSRLTSPTHSSDDFERQFKRVLTKVYQTIERNEI RLAEQDRRDVIKLEWQQVAQIVDRLFLTIFVCVTILITCIVMFKAPHSTESLIS
A_californica _DopC1	MADGINLILSLLVLASCAVMWTRGDPNEYRLVRDLMKGYDKRIRPSLNHTDS LNVTFGLALVQIIDVDEKNQILTTNCWLNQIWTDYNLRWDPAKYGGIKVLRLP YDEVWKPDILLHNNAEVSTFVSSTSSNLIVTNDGNVTWFSMVIFKSSCSINVR YFPFDAQNCSMQFASWSYDGYQVNLIMNTEKGDTSNYIHNSEWELIYLQVE RTVRIFSCCEEPYPDIHFHILIKRRPLFYIFNMILPCILITLVALLGFYIPSDSGEK VTMGITTLLSMTVFMMLVTENMPPTSDVLPLIGIYYGMTIFIVSFATAMTVFTL NIHHKGSRGKAVPMVIKKICFDFLAKMFCMKVDTWDEKHFTPDCNGTPGGA SNYIRFKVANQQDTDQLNMELKQTHEMSPLNRVELPLDSESFQSNGAVPRA FTSFRSSNGNGGSSSEQQNTELTGFEAYFSRVLQRVYQTIERNDIRLAEQDK REGIKLEWQQVAQITDRLLLTCFVLVTLTITGVVLLASPASVKV
D_magna_D opC1	MLRSHKMTQRIWIAFVTVLAIPGNRGDDGEYTRPIVSHEYRLIQHLMANYDR SVRPSRNAAEPLNITFGLALTQIIDVDERNQILTTNCWMNQMWTDYSLVWNV SEFGGIRVVRIPADKVWKSDIILYNNADSEYNSALLSTNVIVTHDGNLTWLSS

	AIFKSSCGINVEYFPFDEQKCSMKFSSWTYDGLQVNLLNQAEEGDLSNYVP NGEWDLIDLVVERNVVFYSCCEEPYPDITFHIVLRRRPLFYVFNLILPCVLITGI ALMSFYMPSDSGEKVTLGITTLLSMTVFLMVIGESMPPTSEKLPLIGLYYGVTI SLVSFATGLSVVTLNIHHRGMRGRGVPPMVKKIVFGVLAKLLFIQLDIAEGCS CSTKAKEFQQNNQFEMNERFNVDIGDSLKNESNGSLPHRLVVVFVAAAWFH VRQSSFGANSGLINNATSNIGPTDSFERQFLRVLNKVYQHIERSEARLADQD RKDAIKLEWQQVALVIDRFLLWIFIIATVAATFGILYMSPHSRLFSM
D_magna_D opC2	MESKAMSPQSVSVRTAGRLLLSLGLFVLLPEAVHSDLNEYRLLRHLMENYDS NARPVANSTKALQVIFGMSLHLLIDVDERNQILTTNCWLTQKWNDSHLTWNA SDYGGIHVVRIPYHRVWRPDIILYNNADAQYTSSVINTNVIVSDNGEVVWLSH GIFRSSCDINVEYFPFDLQSCKMKWASWTYDGFQIDLKKQTDEGDVSNYQT NGEFDLVDFEAIHHEEYYSCCPEPYPDITYVIKMRRRPLFYVFNLILPCLLINGI ALLVFYVPSESGEKVTLGISSLLSMTVFLMTIRESLPPTEKTPLISLYYGVSICL VSFASGLSVVTLNLHHRGLRGTEVPYIMRRIILGGLARLVFLRFDVDKRHSPA STPTSAPAGEPSSSHSGDPIPASRLASRFEESPTKTCASGDPRCTCGKSRNI NQRGRHQPVGVSATRSSGWYSTPVNGGPSCSRNSQERLRLRLQELEPRAT DVYLDPEFSDKFQSAELDFLGIGPPSPKFYSRRVPSTSKAPHFQQQQQQPLI NPESASGSSSAEIGGGFGLLSQGSIYEQRLTELMARVSTAVERNENRLEEQ DRREAIALEWKQLALVCDRILLLTFVIITTVATCVVLTSSPYGP
A_mellifera_ DopC1	MIPTSYKQPVCLTFVSALFETPARIFLGIEVRWSRGSAGFRVEIARWLKSIGY QSRTGGLAFKWSINSMSPLVLFFHYGVLAIIFGNGFGDEHEYRLTKYLLDGY DAGVRPAENSSQPLAVVFGLSLHHIIDVDEKNQILTTNCWVTQIWTDHHLKW NASEFAGIRVIRVPYNRVWRPDTILYNNADPQYSSAVINTNVIVSHTGEVVWL SHGIFRSSCDIDVEFFPFDEQRCVLKWASWTYDGYQLELEKQSEQGDVSNY QANGEFDLVNFSARRNVEYYSCCPEPYPDITYEIRLRRRPMFYVFNLILPCILI NSVALLVFYVPSESGEKVTLGISALLSMTVFLMTIRESLPPTEKTPLISLYYGV SICLVTFASALAVVTLNLHHRGVRGTRVPGIVRSLVLDKLARIVLLNFQEERRS EPVEPPRRKNNCPLHCKPELGQSQSSPKFVARREESNSSSPSLDLGKEGGL EAQWSRVLGRVHATIERNEKRLAEQDRRERMEFDWKQVALVSDRALLCVF FLTTIVSTTVILCGSPPSANIAKEG
H_sapiens_ ACHA9	MNWSHSCISFCWIYFAASRLRAAETADGKYAQKLFNDLFEDYSNALRPVED TDKVLNVTLQITLSQIKDMDERNQILTAYLWIRQIWHDAYLTWDRDQYDGLD SIRIPSDLVWRPDIVLYNKADDESSEPVNTNVVLRYDGLITWDAPAITKSSCV VDVTYFPFDNQQCNLTFGSWTYNGNQVDIFNALDSGDLSDFIEDVEWEVHG MPAVKNVISYGCCSEPYPDVTFTLLLKRRSSFYIVNLLIPCVLISFLAPLSFYLP AASGEKVSLGVTILLAMTVFQLMVAEIMPASENVPLIGKYYIATMALITASTALT

	IMVMNIHFCGAEARPVPHWARVVILKYMSRVLFVYDVGESCLSPHHSRERD HLTKVYSKLPESNLKAARNKDLSRKKDMNKRLKNDLGCQGKNPQEAESYCA QYKVLTRNIEYIAKCLKDHKATNSKGSEWKKVAKVIDRFFMWIFFIMVFVMTIL IIARAD
G_gallus_A CHA9	MKRNNLSSFYVSLWLLFTATMLQAVESAKGKYAQMLFNELFEDYSNALRPV EDTDKVLNVTLQITLSQIKDMDERNQILTAYLWIRQSWYDAYLKWDKDKYDG LDSIRIPSNLVWRPDIVLYNKADDDFSEPVNTNVVLRYDGKITWDAPAITKSS CVVDVSYFPFDSQQCNLTFGSWTYNGNQVDIINSLDSGDLSDFVEDVEWEI HGMPAVKNVITYGCCSEPYPDVTFTLILKRKSSFYIFNLLLPCILISFLAPLGFY LPADSGEKVSLGVTVLLALTVFQLMVAEIMPPSENVPLIGKYYIATMTMITAST ALTIIIMNVHHCGSEAKPVPQWARVVILDYMSKIFFVYDVGENCTSPRREKEQ EHRLEGGDMCRGGDGKSHLSSRNDDSDLKENLNGNWNKSFGVHGENVRE NVNCCSCYKMLIKNIEYIANCVRDHKANRAKGIEWKKVAKVMDRFFMWIFFI MVFFMSVLIIGKAA
G_gallus_A CH10	MGSAPLPACLLALSLAGTVLAPGCGAAQGRLAHKLLHDLFANYSSALRPAED TERALNVTLQVTLSQIIDMDERNQVLTSYLWVRQAWLDAHLAWDKDAYGGI DSIRIPSSYVWRPDIVLYNNADERFGGSMETNVVLRSDGHIMWDSPAITKSS CKVDVSYFPFDGQQCRLTFGSWTYNGNQIDLRNLLDTGDLTDFVENVEWEV LGMPATRNVVTYGCCSEPYPDVTYTLLLRRRASFYIFNLLLPCVMISFLAPLG FYLPADSGEKVSLGVTVLLALTVFQLLVAESMPPSESVPLIGKYYIATMTMITA STALTIFIMNIHHCGPGARPVPPWARRLILHHLARALCVCEVGESCGRPQRE GTGGMGPRDPPGEGVEPGLCPRSRCLCHHHAVLSSVGYIAGVFRRHRTAQ RRAAEWKKVAKVMDRFFMWVFFLMVFLMSVLVIGKAA
H_sapiens_ ACH10	MGLRSHHLSLGLLLLFLLPAECLGAEGRLALKLFRDLFANYTSALRPVADTDQ TLNVTLEVTLSQIIDMDERNQVLTLYLWIRQEWTDAYLRWDPNAYGGLDAIRI PSSLVWRPDIVLYNKADAQPPGSASTNVVLRHDGAVRWDAPAITRSSCRVD VAAFPFDAQHCGLTFGSWTHGGHQLDVRPRGAAASLADFVENVEWRVLG MPARRRVLTYGCCSEPYPDVTFTLLLRRRAAAYVCNLLLPCVLISLLAPLAFH LPADSGEKVSLGVTVLLALTVFQLLLAESMPPAESVPLIGKYYMATMTMVTFS TALTILIMNLHYCGPSVRPVPAWARALLLGHLARGLCVRERGEPCGQSRPPE LSPSPQSPEGGAGPPAGPCHEPRCLCRQEALLHHVATIANTFRSHRAAQRC HEDWKRLARVMDRFFLAIFFSMALVMSLLVLVQAL
H_sapiens_ ACHA7	MRCSPGGVWLALAASLLHVSLQGEFQRKLYKELVKNYNPLERPVANDSQPL TVYFSLSLLQIMDVDEKNQVLTTNIWLQMSWTDHYLQWNVSEYPGVKTVRF PDGQIWKPDILLYNSADERFDATFHTNVLVNSSGHCQYLPPGIFKSSCYIDVR WFPFDVQHCKLKFGSWSYGGWSLDLQMQEADISGYIPNGEWDLVGIPGKR

	SERFYECCKEPYPDVTFTVTMRRRTLYYGLNLLIPCVLISALALLVFLLPADSG EKISLGITVLLSLTVFMLLVAEIMPATSDSVPLIAQYFASTMIIVGLSVVVTVIVL QYHHHDPDGGKMPKWTRVILLNWCAWFLRMKRPGEDKVRPACQHKQRRC SLASVEMSAVAPPPASNGNLLYIGFRGLDGVHCVPTPDSGVVCGRMACSPT HDEHLLHGGQPPEGDPDLAKILEEVRYIANRFRCQDESEAVCSEWKFAACV VDRLCLMAFSVFTIICTIGILMSAPNFVEAVSKDFA
H_sapiens_ ACHA3	MGSGPLSLPLALSPPRLLLLLLLLSLLPVARASEAEHRLFERLFEDYNEIIRPVA NVSDPVIIHFEVSMSQLVKVDEVNQIMETNLWLKQIWNDYKLKWNPSDYGG AEFMRVPAQKIWKPDIVLYNNAVGDFQVDDKTKALLKYTGEVTWIPPAIFKSS CKIDVTYFPFDYQNCTMKFGSWSYDKAKIDLVLIGSSMNLKDYWESGEWAII KAPGYKHDIKYNCCEEIYPDITYSLYIRRLPLFYTINLIIPCLLISFLTVLVFYLPS DCGEKVTLCISVLLSLTVFLLVITETIPSTSLVIPLIGEYLLFTMIFVTLSIVITVFV LNVHYRTPTTHTMPSWVKTVFLNLLPRVMFMTRPTSNEGNAQKPRPLYGAE LSNLNCFSRAESKGCKEGYPCQDGMCGYCHHRRIKISNFSANLTRSSSSES VDAVLSLSALSPEIKEAIQSVKYIAENMKAQNEAKEIQDDWKYVAMVIDRIFL WVFTLVCILGTAGLFLQPLMAREDA
H_sapiens_ ACHA4	MELGGPGAPRLLPPLLLLLGTGLLRASSHVETRAHAEERLLKKLFSGYNKWS RPVANISDVVLVRFGLSIAQLIDVDEKNQMMTTNVWVKQEWHDYKLRWDPA DYENVTSIRIPSELIWRPDIVLYNNADGDFAVTHLTKAHLFHDGRVQWTPPAI YKSSCSIDVTFFPFDQQNCTMKFGSWTYDKAKIDLVNMHSRVDQLDFWESG EWVIVDAVGTYNTRKYECCAEIYPDITYAFVIRRLPLFYTINLIIPCLLISCLTVLV FYLPSECGEKITLCISVLLSLTVFLLLITEIIPSTSLVIPLIGEYLLFTMIFVTLSIVIT VFVLNVHHRSPRTHTMPTWVRRVFLDIVPRLLLMKRPSVVKDNCRRLIESMH KMASAPRFWPEPEGEPPATSGTQSLHPPSPSFCVPLDVPAEPGPSCKSPS DQLPPQQPLEAEKASPHPSPGPCRPPHGTQAPGLAKARSLSVQHMSSPGE AVEGGVRCRSRSIQYCVPRDDAAPEADGQAAGALASRNTHSAELPPPDQP SPCKCTCKKEPSSVSPSATVKTRSTKAPPPHLPLSPALTRAVEGVQYIADHL KAEDTDFSVKEDWKYVAMVIDRIFLWMFIIVCLLGTVGLFLPPWLAGMI
C_elegans_ UNC-29	MRTNRLSWILVLSVVIFLVIINTINASDDEERLMVDVFRGYNSLIQPVRNSSEL PLIVKMALQLVLLINVDEKDQVMHTNVWLTLQWHDFQMKWNPVNYGEIKQIR VSPDKVWLPDIVLFNNADGNYEVSFMCNVVINHKGDMLWVPPAIYKSSCIID VEFFPFDEQVCTLVFGSWTYNENEIKLEFVQAELVDVSEYSASSIWDVIDVPA SLVNKRSRIEFQVRIRRKTLFYTVVLIIPTVLMAFLSMAVFFLPTDSGEKITLTIS VLLSIVVFLLLVSKILPPTSSTIPLMAKYLLLTFVLNVITILVTVIIINVYFRGPRTH RMPQWVRVVFLQFLPKLVCMKRPKSASERSAVRSGMAQLPGVGQFTLSPS AHHPLCPSADDRTTTIRNTASNETSAYYPLSTDALRAIDAIEYITEHLKRDEQH

	KSFRDDWKYVAMIIDRLLLYVFFGITVGGTCGILFSAPHVFQRIDQQEMLDRL KEKYDTASNIP
D_chrysanth emi_ELIC	APADNAADARPVDVSVSIFINKIYGVNTLEQTYKVDGYIVAQWTGKPRKTPG DKPLIVENTQIERWINNGLWVPALEFINVVGSPDTGNKRLMLFPDGRVIYNAR FLGSFSNDMDFRLFPFDRQQFVLELEPFSYNNQQLRFSDIQVYTENIDNEEID EWWIRKASTHISDIRYDHLSSVQPNQNEFSRITVRIDAVRNPSYYLWSFILPL GLIIAASWSVFWLESFSERLQTSFTLMLTVVAYAFYTSNILPRLPYTTVIDQMII AGYGSIFAAILLIIFAHHRQAMGVEDDLLIQRCRLAFPLGFLAIGCVLVIRGITL
H_sapiens_ 5HT3A	MLLWVQQALLALLLPTLLAQGEARRSRNTTRPALLRLSDYLLTNYRKGVRPV RDWRKPTTVSIDVIVYAILNVDEKNQVLTTYIWYRQYWTDEFLQWNPEDFDN ITKLSIPTDSIWVPDILINEFVDVGKSPNIPYVYIRHQGEVQNYKPLQVVTACSL DIYNFPFDVQNCSLTFTSWLHTIQDINISLWRLPEKVKSDRSVFMNQGEWEL LGVLPYFREFSMESSNYYAEMKFYVVIRRRPLFYVVSLLLPSIFLMVMDIVGF YLPPNSGERVSFKITLLLGYSVFLIIVSDTLPATAIGTPLIGVYFVVCMALLVISL AETIFIVRLVHKQDLQQPVPAWLRHLVLERIAWLLCLREQSTSQRPPATSQAT KTDDCSAMGNHCSHMGGPQDFEKSPRDRCSPPPPPREASLAVCGLLQELS SIRQFLEKRDEIREVARDWLRVGSVLDKLLFHIYLLAVLAYSITLVMLWSIWQY A
H_spaiens_ ACHB2	MARRCGPVALLLGFGLLRLCSGVWGTDTEERLVEHLLDPSRYNKLIRPATN GSELVTVQLMVSLAQLISVHEREQIMTTNVWLTQEWEDYRLTWKPEEFDNM KKVRLPSKHIWLPDVVLYNNADGMYEVSFYSNAVVSYDGSIFWLPPAIYKSA CKIEVKHFPFDQQNCTMKFRSWTYDRTEIDLVLKSEVASLDDFTPSGEWDIV ALPGRRNENPDDSTYVDITYDFIIRRKPLFYTINLIIPCVLITSLAILVFYLPSDCG EKMTLCISVLLALTVFLLLISKIVPPTSLDVPLVGKYLMFTMVLVTFSIVTSVCVL NVHHRSPTTHTMAPWVKVVFLEKLPALLFMQQPRHHCARQRLRLRRRQRE REGAGALFFREAPGADSCTCFVNRASVQGLAGAFGAEPAPVAGPGRSGEP CGCGLREAVDGVRFIADHMRSEDDDQSVSEDWKYVAMVIDRLFLWIFVFVC VFGTIGMFLQPLFQNYTTTTFLHSDHSAPSSK
H_sapines_ ACHB4	MRRAPSLVLFFLVALCGRGNCRVANAEEKLMDDLLNKTRYNNLIRPATSSSQ LISIKLQLSLAQLISVNEREQIMTTNVWLKQEWTDYRLTWNSSRYEGVNILRIP AKRIWLPDIVLYNNADGTYEVSVYTNLIVRSNGSVLWLPPAIYKSACKIEVKYF PFDQQNCTLKFRSWTYDHTEIDMVLMTPTASMDDFTPSGEWDIVALPGRRT VNPQDPSYVDVTYDFIIKRKPLFYTINLIIPCVLTTLLAILVFYLPSDCGEKMTLC ISVLLALTFFLLLISKIVPPTSLDVPLIGKYLMFTMVLVTFSIVTSVCVLNVHHRS PSTHTMAPWVKRCFLHKLPTFLFMKRPGPDSSPARAFPPSKSCVTKPEATA TSTSPSNFYGNSMYFVNPASAASKSPAGSTPVAIPRDFWLRSSGRFRQDVQ

	EALEGVSFIAQHMKNDDEDQSVVEDWKYVAMVVDRLFLWVFMFVCVLGTV GLFLPPLFQTHAASEGPYAAQRD
C_elegans_ LGC-53	MIISYLARIVLLLPELSSGDTDRLAEIAANQEHSIQLDQMPPDRTFHYPNIPSS QHDRRREGGKRKKFRKERLDEPCTAINETLRKNLLQELFDDSYDKNNLPSA NSTEVIVELTVQSITEISEFSSSFKADVWFSQIWRDPRLDFSDRNYCLKNISLA SHKLPTMWSPNVCFVNSKKVEIHASPSQNILLLIFPNGTVWLNFRVSLTGPC KLDLTYFPMDRQSCNLVFESYSYNTAEVRIVWRDWDAVTIPDPDSKNLPDFE LVNIEHMNATLVYTAGLWDQLEVKFTFRRLYGYYVLQAYMPTYLSVFISWIAF WIDTKALPARITLGVSSLMALTFQFGNIVKNLPRVSYVKALDIWMFGCVGFIFL SLVELAVVGFADKLEAKRRRHNRCKEQLMMRSDSEQQWLSRLSGQRPQVS ETNSDATHTIQINDGNGNVRRRKSEEKRQKLLLQMESPMFVNGEKIDEISAK LFPLLFTAFNIFYWFYYIGMSGGFF
C_elegans_ LGC-52	MIYSIQALFIETILMSARIHDVTYQHIEDFTCDDIGLSSLSGAKATDLARILMTNY SRNALPEPAPVKVEVEITIQDISDISAITGTFTIDYWISAIWNDKRLAFSHLDPC RKNLSLDHDMEPKLWSPNVCIVNSKSTKVHDSPKPNILLMIFPNGTIWLNYRI RSEAPCQMNLRNFPLDSIRCSLVFESYSYNAADVELQWLEWSPVSTVRNDY NLPDFRMTNITYGSVTEAYTAGMWHRLSVSIHFERLYGFYILQMYLPTYISVFI SWIAFWMDTKALPARITLSVSSLMALTFQFGNIVKSLPKASYVKAIDIWMFSC VGFIFFSLIELAIVAYNDKMHDQRLREVRCSVGNLNSNGIGQYNSTTRRSFVE LSARRAKGSELGASIDRVASMLFPTMFALFNLVYWAYYLTTSMNPVYT
D_melanoga ster_HisCl1	MQSPTSKLVEFRCLIALAIYLHALEQSIQHCHCVHGYRNNTESAELVSHYESS LSLPDILPIPSKTYDKNRAPKLLGQPTVVYFHVTVLSLDSINEESMTYVTDIFLA QSWRDPRLRLPENMSEQYRILDVDWLHSIWRPDCFFKNAKKVTFHEMSIPN HYLWLYHDKTLLYMSKLTLVLSCAMKFESYPHDTQICSMMIESLSHTVEDLV FIWNMTDPLVVNTEIELPQLDISNNYTTDCTIEYSTGNFTCLAIVFNLRRRLGY HLFHTYIPSALIVVMSWISFWIKPEAIPARVTLGVTSLLTLATQNTQSQQSLPP VSYVKAIDVWMSSCSVFVFLSLMEFAVVNNFMGPVATKAMKGYSDENISDL DDLKSALQHHRESIIEPQYDTFCHGHATAIYIDKFSRFFFPFSFFILNIVYWTTF L
H_sapiens_ GBRA1	MRKSPGLSDCLWAWILLLSTLTGRSYGQPSLQDELKDNTTVFTRILDRLLDG YDNRLRPGLGERVTEVKTDIFVTSFGPVSDHDMEYTIDVFFRQSWKDERLKF KGPMTVLRLNNLMASKIWTPDTFFHNGKKSVAHNMTMPNKLLRITEDGTLLY TMRLTVRAECPMHLEDFPMDAHACPLKFGSYAYTRAEVVYEWTREPARSV VVAEDGSRLNQYDLLGQTVDSGIVQSSTGEYVVMTTHFHLKRKIGYFVIQTY LPCIMTVILSQVSFWLNRESVPARTVFGVTTVLTMTTLSISARNSLPKVAYAT AMDWFIAVCYAFVFSALIEFATVNYFTKRGYAWDGKSVVPEKPKKVKDPLIK

KNNTYAPTATSYTPNLARGDPGLATIAKSATIEPKEVKPETKPPEPKKTFNSV SKIDRLSRIAFPLLFGIFNLVYWATYLNREPQLKAPTPHQ

Figure S 1 The A158G mutation in Loop E of a _D. magna_ dop-LGIC decreases catecholamine sensitivity.

A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in wild-type and A158G Dm-DopC1 expressing _Xenopus_ oocytes. Current normalized by I/Imax. EC_{50} values (± SEM) shown below each plot. **** = P<0.0001, ns = not significant calculated by sum of squares F test. WT: n = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); A158G: n = 10 (dopamine), 11 (norepinephrine), 10 (epinephrine). **D.** Relative maximum I/Imax currents of _Xenopus_ oocytes expressing wild-type and A158G Dm-DopC1, perfused for 10 s with 1 mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi), serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). WT: n = 18 (dopamine, norepinephrine, ACh, tyramine), 17 (epinephrine, serotonin); A158G: n = 6 per agonist. ns = not significant by one-way ANOVA and a Fisher's LSD. **E.** Sequence alignment of the loop E

region. Residues A158 and L155 are indicated by arrows, loop E is highlighted by the blue
bar. **F.** AlphaFold2 model of the loop E (blue) region of Dm-DopC1 with A158 and L155
relative to loop B (pink), loop C (green), loop F (yellow), loop A (orange) and loop D
(turquoise) at the ligand-binding site. **G.** Baseline currents in uninjected oocytes or oocytes
expressing WT or L155D Dm-DopC1. Baseline currents measured 5s before agonist
application. Error bars represent SEM, n = 3 (WT), 6 (uninjected), 7 (L155D). **H.** Peak
currents in response to 10s agonist application of *Xenopus* oocytes expressing wild-type and
L155D Dm-DopC1 with 1mM each: dopamine (DA), norepinephrine (Nor), epinephrine (Epi),
serotonin (5HT), tyramine (Tyr), and acetylcholine (ACh). Error bars represent SEM. WT: n =
5 (dopamine), 4 (norepinephrine, ACh, tyramine), 3 (epinephrine, serotonin); L155D: n = 9
(dopamine, tyramine, ACh), 6 (norepinephrine, serotonin), 5 (epinephrine). Note scale bar on
$\log_{10}$ scale.

Figure S 2 S123P exhibits a steeper Hill slope than wild type for norepinephrine.

A-C. Dose response curves for dopamine (A), norepinephrine (B) and epinephrine (C) in
wild-type and S123P Dm-DopC1 expressing *Xenopus* oocytes. Current normalized by
I/Imax. Hill slope values are shown underneath each plot with SEM, calculated with a 4-
parameter curve. WT: n = 24 (dopamine), 12 (norepinephrine), 11 (epinephrine); S123P: n =
12, (dopamine), 4 (norepinephrine) and 8 (epinephrine).

Figure S 3 Continuous traces of Dm-DopC1 oocytes exposed to a panel of compounds. Red bar indicates application of 100µM compound.

Figure S 4 *In silico* **binding models of Dm-DopC1 with acetylcholine (A) and dopamine (B). A & B.** Predicted AlphaFold3 co-fold models of Dm-DopC1 in complex with acetylcholine (A) and dopamine (B). Ligands are shown in beige with polar contacts between the ligand and adjacent residues shown by dashed yellow lines. Colors represent conserved ligand binding loops: A: orange, B: pink, C: green, D: teal, E: blue, F: yellow. Amino acid side chains are shown for residues which appear in 2D poses for dopamine, as well as D188 and S227 which forms a hydrogen bond with dopamine.