Poultry Litter Microbiome Shifts in Commercial Broiler Houses Following a Biotic-Based Litter Treatment

Elena G. Olson[1], Brett Hale[2], Steven C. Ricke[1]

[1]Meat Science and Animal Biologics Discovery Program, Department of Animal and Dairy Sciences, University of Wisconsin, Madison, WI, 53706, USA

[2]AgriGro, Inc, Doniphan, MO, 63935, USA

Abstract

Litter management plays a critical role in broiler production, affecting bird health, performance, and environmental impact. This preliminary study evaluated the effects of IndigoLT™, a biological amendment, in combination with a 40% sodium bisulfate ($NaHSO_4$) regimen on litter microbiome composition, compared to a standard $NaHSO_4$-only treatment. Litter samples ($n = 18$) were collected from three commercial broiler houses following flock removal: one control ($NaHSO_4$-only), one treated with IndigoLT for a single flock, and one treated for two consecutive flocks. The V4 region of the 16S rRNA gene was sequenced to assess prokaryotic community composition. Alpha diversity metrics (Faith's phylogenetic diversity, observed ASVs, Pielou's evenness, Shannon diversity) did not differ significantly across treatments (ANOVA, $p > 0.05$). In contrast, PERMANOVA analyses of Bray-Curtis, Jaccard, Weighted and Unweighted UniFrac distances revealed significant shifts in β diversity between IndigoLT-amended and control groups ($q < 0.05$), with no significant differences between the one- and two-flock IndigoLT treatments. Core microbiome and differential abundance analyses suggested that IndigoLT, when paired with reduced $NaHSO_4$ input, may accelerate organic matter decomposition, promote nitrogen retention, and suppress potentially pathogenic taxa. Although limited by the absence of baseline sampling and biological replication, these findings suggest that IndigoLT influences litter microbial succession. Future work should aim to optimize inclusion rates most complementary between $NaHSO_4$ and IndigoLT to enhance litter quality, reduce NH_3 volatilization, and support bird health.

Keywords: Broiler litter, Microbiome, IndigoLT, 16S rRNA sequencing, Litter amendment

Introduction

Poultry litter management is critical to broiler production systems, directly impacting bird health and welfare, environmental sustainability, and food safety. Litter serves as a biologically complex substrate composed of fecal matter, feed residues, bedding material, and diverse microbial populations that interact dynamically with environmental and management factors. These microbial communities are shaped by variables such as stocking density, litter age, and the application of chemical or biological amendments (Bolan et al., 2010). As the poultry industry shifts toward antimicrobial stewardship and microbiome-informed production strategies, characterizing the microbial ecology of broiler litter has gained prominence (Pedroso et al., 2013).

Broiler producers commonly apply chemical amendments to improve litter conditions, reduce moisture, suppress pathogen proliferation, and limit ammonia (NH_3) volatilization. Acidifying agents such as sodium bisulfate ($NaHSO_4$) are widely used to lower litter pH, thereby reducing the conversion of nonvolatile ammonium (NH_4^+) into volatilized NH_3 (Ritz et al., 2004; Rothrock et al., 2010). Although the effectiveness of $NaHSO_4$-based amendments in reducing NH_3 emissions is well established (Johnson et al., 2011), their impacts on litter microbiome structure and composition remain less understood (Toppel et al., 2019; de Toledo et al., 2020; Johnson et al., 2021). While pH reduction is associated with a general decrease in total bacterial load (Johnson et al., 2021 and references therein), some evidence suggests that it may inadvertently promote the persistence of potentially pathogenic bacterial taxa, including *Salmonella* (Williams et al., 2012) and *Escherichia* (Joerger et al., 2020; Johnson et al., 2021). This phenomenon may be attributed to physiological characteristics such as acid tolerance and nitrogen metabolism, allowing select organisms to survive under lowered pH conditions and reduced competitive pressures resulting from microbial load decline (Bucher et al., 2020; Johnson et al., 2021; Swelum et al., 2021). The potential for dysbiosis following acidification highlights the need to investigate litter microbiome dynamics more holistically (e.g., via metabarcoding approaches) and suggests that complementary interventions may be warranted to preserve microbiome stability while optimizing NH_3 reduction to support bird health and welfare.

Biological amendments, including prebiotics, probiotics, and postbiotics, are increasingly employed in poultry production systems to reduce reliance on antibiotic growth promoters (Ricke 2018; Callaway and Ricke, 2023). These technologies are formulated primarily as feed or water additives (Pedroso et al., 2013; Swanson et al. 2025) and have been shown to elevate mutualistic microorganisms that competitively exclude pathogens in the gastrointestinal tract, directly suppress pathogen proliferation through specialized metabolites or antimicrobial compounds, prime the host immune system, and improve production outcomes by enhancing nutrient uptake, digestibility, and intestinal integrity (Reuben et al., 2021). However, the application of biological amendments to poultry litter remains comparatively underexplored relative to their use as ingestible interventions (Pedroso et al., 2013), and much of the existing literature reflects controlled microcosm conditions that may not fully capture microbial dynamics in commercial production environments (Johnson et al., 2021). Expanding biological intervention strategies to the litter environment is particularly relevant given that poultry litter harbors zoonotic pathogens,

including *Salmonella*, *Campylobacter*, and *Staphylococcus* species, which are prominently
associated with the acquisition and dissemination of antimicrobial resistance (Gilchrist et al.,
2007). Thus, biological amendments designed for litter application may offer a novel opportunity
to enhance both food safety and bird performance.

IndigoLT™ (LA-I, AgriGro, Inc.) is a fermentation-based biotic litter amendment
formulated to support microbial stability in poultry litter. The biotic-based technology contains a
prebiotic/postbiotic fraction composed of organic acids, hygroscopic sugar alcohols, nitrogen-
containing compounds, and antimicrobial compounds/specialized metabolites derived from a
proprietary microbial consortium. In addition to the biological fraction, LA-I includes liquid $CaCl_2$
formulated with additional organic acid chelation. Although its commercial application is
expanding, the effects of LA-I on litter microbial community structure and ecological function
remain largely unexplored.

This preliminary study aimed to evaluate the effects of LA-I litter amendment on the
prokaryotic litter microbiome in a commercial broiler operation. Litter samples were collected
from three poultry houses: one treated only with $NaHSO_4$ (control), one treated with LA-I for a
single flock, and one treated across two consecutive flocks. Both LA-I regimens included reduced
rates of $NaHSO_4$. Microbial communities were characterized using 16S rRNA gene sequencing
targeting the V4 region, and shifts in diversity, composition, and taxon abundance were assessed.
Findings from this study provide initial insight into how different litter amendments may modulate
microbial ecology and support the development of microbiome-targeted strategies for sustainable
poultry production.

Materials and Methods
<u>Study design</u>

The test site was identified at Gillham, Arkansas, USA (34.16214°N, 94.35262°W) and
encompassed three commercial broiler houses with 500 X 43 ft dimensions (21,500 ft²; ≈ 1,997
m²). Bedding material consisted of locally sourced rice (*Oryza sativa*) hulls and had been used for
18 consecutive broiler flocks prior to study initiation. Two houses (denoted hereafter as "1" and
"2") received the prebiotic amendment IndigoLT ("LA-I"; AgriGro Inc, Doniphan, MO, USA) at
a rate of 48 oz/1,000 ft² (≈ 1.47 L/93 m²) 72 hrs before chick placement. Treatment 2 also received
IndigoLT during the preceding flock (**Table 1**). Prebiotic applications were made with a spot
sprayer between litter windrowing and leveling.

Approximately 16,333 Ross 308 chicks (Aviagen, Huntsville, AL, USA) were placed per
house, corresponding to a stocking density of 1.32 birds/ft² (8.18 birds/m²). Less than 24 hours
prior to chick placement, an $NaHSO_4$-based litter amendment (Jones-Hamilton Co., Maumee, OH,
USA) was applied to houses 1 and 2 at a rate of 40 lbs/1,000 ft² (≈ 195 g/m²) and to house 3
("Control") at 100 lbs/1,000 ft² (≈ 488 g/m²) (**Table 1**). All other production parameters remained
constant across treatments.

Table 1 Litter amendment regimen for each treatment		
Treatment	**LA-I**	**NaHSO₄**
1	1.47 L/93 m² (1st exposure)	195 g/m²
2	1.47 L/93 m² (2nd exposure)	195 g/m²
Control	——	488 g/m²

Sample collection and DNA isolation

On the day following bird removal, six composite litter samples ($n = 18$) were collected from each house to capture spatial heterogeneity. Each composite consisted of five subsamples collected to a depth of 10 cm using a sanitized trowel, with a new trowel used for each house to prevent cross-contamination. Subsamples were pooled into sterile ziplock bags and immediately flash-frozen on dry ice. Samples were then shipped on dry ice to the University of Missouri (MU) Genomics Technology Core, where they were stored at −80 °C until DNA isolation.

DNA was isolated using the QIAamp PowerFecal Pro DNA Kit (Cat. #51804; Qiagen, Hilden, Germany) according to the manufacturer's instructions, except that samples were homogenized using a TissueLyser II (Qiagen) at 30 Hz for 10 minutes in place of the vortex adapter specified in the protocol. DNA concentrations were determined fluorometrically using a Qubit 2.0 fluorometer (Thermo Fisher Scientific, Waltham, MA, USA) with the Quant-iT dsDNA Broad-Range Assay Kit (Cat. #Q33130) and subsequently normalized to a consistent concentration and volume.

16S rRNA library preparation and sequencing

Library preparation and sequencing were conducted at the MU Genomics Technology Core. Prokaryotic 16S rRNA amplicons were generated by targeting the V4 region of the 16S rRNA gene using universal primers U515F/806R, flanked by Illumina standard adapter sequences (Caporaso et al., 2011; Walters and Caporaso et al., 2011). All reactions employed dual-indexed forward and reverse primers. PCR was performed in 50 μL reactions containing 100 ng DNA, 0.2 μM of each primer, 200 μM of each dNTP, 1 U of Phusion High-Fidelity DNA Polymerase (Cat #F530S; Thermo Fisher Scientific), and the appropriate volume of 5X Phusion HF Buffer (Cat. #F518L; Thermo Fisher Scientific). Amplification conditions included initial denaturation at 98 °C for 3 min, followed by 25 cycles of 98 °C for 15 s, 50 °C for 30 s, and 72 °C for 30 s, with a final extension at 72 °C for 7 min.

Following amplification, 5 μL of product from each reaction was pooled, mixed thoroughly, and purified using Axygen Axyprep MagPCR Clean-up Beads (Cat. #14-223-227; Thermo Fisher Scientific) at a 1:1 bead-to-amplicon volume ratio. The mixture was incubated for 15 min at room temperature, washed twice with 80% ethanol, and air-dried. The purified pellet

was resuspended in 32.5 µL of EB buffer (Cat. #19086; Qiagen), incubated at room temperature
for 2 min, and placed on a magnetic stand for 5 min to recover the supernatant.
The final pooled amplicon library was evaluated using the Advanced Analytical Fragment
Analyzer (Agilent Technologies, Santa Clara, CA, USA), quantified with the Quant-iT High
Sensitivity dsDNA Assay Kit (Cat. #Q33120; Thermo Fisher Scientific), and diluted according to
Illumina's protocol for sequencing on the MiSeq platform as 2×250 bp paired-end reads.
152
Bioinformatic analysis
DNA sequences were assembled and annotated at the MU Bioinformatics and Analytics
Core. Primers were designed to match the 5′ ends of the forward and reverse reads. Adapter and
primer removal was conducted using Cutadapt (v2.6) (Martin, 2011). The forward primer was
trimmed from the 5′ end of the forward read, and, if present, the reverse complement of the reverse
primer was removed from the 3′ end of the same read along with all downstream bases, resulting
in bidirectional trimming if the insert was shorter than the amplicon. The reverse read was
processed analogously, with primer roles reversed. Read pairs were discarded if either read failed
to match a 5′ primer, with an allowed error rate of 0.1 and a minimum overlap of 3 bp at the 3′ end
of the primer for removal. Each read underwent two passes to ensure complete primer removal.
Amplicon sequence variant (ASV) inference was performed using the DADA2 plugin
(v1.30.0) (Callahan et al., 2016) within QIIME2 (v2024.5.1) (Bolyen et al., 2019). Denoising
parameters included truncating both forward and reverse reads to 150 bases, discarding reads with
more than 2.0 expected errors, and removing chimeras using the 'consensus' method. The QIIME2
environment incorporated Python (v3.9.19) and BIOM (v2.1.15) for downstream processing.
Taxonomic classification was performed using the 'classify-sklearn' method with the SILVA
v138.1 reference database (Quast et al., 2012).
170
Statistical analysis
The QIIME2 artifacts were imported into RStudio (v4.4.0) (R Core Team, 2020) with the
qiime2R package (v0.99.6) (Bisanz et al., 2018) and were used to create a "phyloseq" S4 object
(v1.48.0) (McMurdie and Holmes, 2013). One sample with <1,000 reads was removed prior to
analysis. Per-sample coverage/sequencing depth was then estimated using Good's coverage,
calculated with the 'phyloseq_coverage' function from the metagMisc package (v0.5.0)
(Mikryukov, 2020) with the formula *1 - (F1/N)*, where *F1* is the number of singleton ASVs and *N*
is the read sum for all ASVs (Good, 1953; Good and Toulmin, 1956; Chao and Jost, 2012).
Observed ASVs (richness) were evaluated as a function of sequencing depth using the 'rarecurve'
function from the vegan package (v2.6.1) (Dixon, 2003). All visualizations, including family-level
relative abundance plots, were generated with ggplot2 (v3.4.2) (Wickham and Sievert, 2009). The
rooted phylogenetic tree was converted to a "phylo" object using the ape package (v5.8) (Paradis
et al., 2004) and visualized as a cladogram with ggtree (v3.12.0) (Yu et al., 2017).
Mean estimates for within-sample (α) diversity were calculated by subsampling 1,000
sequences per sample across 1,000 iterations. Richness and Shannon diversity (Hill, 1973) were

determined with the 'estimate_richness' phyloseq function, Pielou's evenness (Pielou, 1966) with
the 'evenness' function from the microbiome package (v1.26.0) (Lahti and Shetty, 2018), and
Faith's phylogenetic diversity (Faith, 1992) with the 'pd' function in picante (v1.8.2) (Kembel et
al., 2010). For each metric, Levene's test (Levene, 1960) was used to assess homogeneity of
variance across treatments via the 'leveneTest' function in the car package (v3.1-2) (Fox et al.,
2012), and the Shapiro-Wilk test (Shapiro and Wilk, 1965) was used to evaluate normality within
each treatment group with 'shapiro.test' from the stats package (v4.4.0). All assumptions of
homogeneity and normality were met ($p > 0.05$), and treatment effects were evaluated using
analysis of variance (ANOVA) models fitted with the 'aov' function in stats. The α diversity
estimates were visualized with ggplot2.
Prior to β diversity analysis, reads were normalized using cumulative sum scaling (CSS)
with the 'cumNormStat' and 'cumNorm' functions from the metagenomeSeq package (v1.46.0)
(Paulson et al., 2013). Global dissimilarity matrices were then constructed using Bray-Curtis (Bray
and Curtis, 1957), Jaccard (1908), Weighted UniFrac, and Unweighted UniFrac (Lozupone and
Knight, 2005) distances via the 'distance' function in phyloseq. Ordination was performed using
Principal Coordinates Analysis (PCoA) with the 'ordinate' function in phyloseq, and results were
visualized with ggplot2. Compositional dissimilarity among treatment levels was assessed
statistically using permutational multivariate analysis of variance (PERMANOVA) with the
'adonis2' function from vegan (v2.6-6.1) with 9,999 permutations. As PERMANOVA identified
significant treatment effects ($p < 0.05$) for all distance metrics, multilevel pairwise comparisons
were performed using 'pairwise.adonis' from the pairwiseAdonis package (v0.4.1) (Martinez
Arbizu, 2020), applying false discovery rate (FDR) correction to control for multiple testing.
Homogeneity of multivariate dispersions was evaluated with the 'betadisper' function from vegan,
and group centroid distances were compared across treatment levels using a permutational
ANOVA model implemented with the 'anova' function from the stats package (9,999
permutations).
Global and treatment-specific core microbiome features were defined as ASVs present in
all but one sample per condition, following the "relaxed core" concept proposed by Mejia et al.
(2025). Per-sample relative abundances of core ASVs were calculated from CSS-normalized
counts using the 'transform_sample_counts' function in phyloseq, and visualized with the
ComplexHeatmap package (v2.20.0) (Gu et al., 2016). The top 10 most represented genera,
families, and orders by ASV count within each core microbiome were identified and visualized
using ggplot2. Shared and exclusive ASVs among core groups were further visualized using the
ggvenn package (v0.1.10) (Yan, 2021). Core microbiomes were clustered using Jaccard distance,
calculated with the 'vegdist' function from the vegan package and visualized in ggplot2.
Features differentially abundant among treatment levels were identified using the
MaAsLin2 package (Microbiome Multivariable Associations with Linear Models) (v1.18.0)
(Mallick et al., 2021). Log-transformed linear models were constructed using total sum scaling
(TSS)-normalized count data following MaAsLin2 default parameters. Pairwise treatment
comparisons were performed independently. To reduce the influence of spurious or

environmentally stochastic features in the absence of a shared baseline, analyses were restricted to
features present in the core microbiomes of at least two treatment groups. Genus-, family-, and
order-level counts were additionally generated using the 'tax_glom' function from the phyloseq
package and reanalyzed using the same MaAsLin2 parameters. Features with a Benjamini-
Hochberg-adjusted *p*-value ($q < 0.25$) (Benjamini and Hochberg, 1995) were considered
differentially abundant as recommended by MaAsLin2 documentation (Mallick et al., 2021). The
log-normalized FDR [$-sign(coefficient) \times log(q\text{-value})$] (Boolchandani et al., 2022), which is
equivalent to log_2(fold change) (Reasoner et al., 2024), was calculated for significant features and
visualized using the ComplexHeatmap package. Shared and unique features across pairwise
comparisons were visualized with the ggvenn package.
236
Results
<u>Sequencing data overview</u>
The objective of the present study was to determine the impacts of LA-I, a commercial
amendment, on poultry litter microbiome dynamics when integrated into a conventional litter
management system. Comparisons were made between a house receiving LA-I plus 40% of the
recommended $NaHSO_4$ application for the first time (treatment 1), a house treated with the same
combination for a second consecutive flock (treatment 2), and a control house receiving only the
recommended $NaHSO_4$ application. Six composite litter samples were collected per house, and the
V4 region of the 16S rRNA gene was amplified and sequenced to assess prokaryotic microbiome
shifts across treatments.
Illumina V4 sequencing yielded $333,596 \pm 48,470$ reads per sample (mean $\pm$ 1 SD), with
Good's coverage estimates ($0.999994 \pm 6e{-}6$) and rarefaction curves supporting adequate
sampling depth (**Figure 1a**–**c**). Denoising rendered 2,938 ASVs, with individual samples
encompassing 531.94 ± 157.88. The dominant phyla were Firmicutes (59.87% of reads, 69.8% of
ASVs), Actinobacteriota (35.95% of reads, 11.37% of ASVs), and Proteobacteria (3.31% of reads,
7.26% of ASVs). The most abundant classes were Bacilli (54.58% of reads, 25.61% of ASVs),
Actinobacteria (35.94% of reads, 10.51% of ASVs), and Clostridia (5.27% of reads, 43.48% of
ASVs). Dominant orders included Staphylococcales (30.48% of reads, 4.16% of ASVs),
Micrococcales (19.80% of reads, 5.92% of ASVs), and Lactobacillales (17.74% of reads, 7.84%
of ASVs). At the family level, Staphylococcaceae (30.48% of reads, 4.16% of ASVs),
Dermabacteraceae (10.01% of reads, 1.10% of ASVs), and Lactobacillaceae (9.40% of reads,
3.30% of ASVs) were most prevalent (**Figure 1d**). Dominant families were distributed in
taxonomically consistent clades within a rooted phylogenetic tree, reflecting anticipated sequence
homology (**Figure 1e**).

Figure 1 Overview of the poultry litter microbiome. (**a**) Sequencing depth distribution. (**b**) Distribution of Good's coverage determined with the formula: *1 - (F1/N)*, where *F1* is the number of singleton OTUs and *N* is the read sum for all OTUs. (**c**) Rarefaction curve showing observed ASVs (richness) as a function of sequencing depth (reads per sample). (**d**) Stacked bar plot of family-level per-sample composition. Families contributing less than 5% of total read abundance across all samples were grouped into "<5% abund." after agglomerating counts to the Family level. (**e**) Rooted phylogenetic tree. Tip color corresponds to family utilizing the same key as the stacked bar plot.

<u>LA-I administration influenced prokaryotic microbiome composition</u>

Treatment effects on α diversity were evaluated using mean estimates derived from 1,000 rarefied iterations for Faith's phylogenetic diversity, observed ASVs, Shannon diversity, and Pielou's evenness. No statistically significant differences were detected among treatments for any metric (ANOVA, $p > 0.05$) (**Figure 2**). Following α diversity analysis, count matrices were normalized via cumulative sum scaling (CSS), and β diversity was assessed to infer treatment-driven compositional shifts.

Despite the absence of significant α diversity differences, treatment exerted a significant effect on microbial community composition (PERMANOVA, $p < 0.05$) across all tested dissimilarity measures, including Bray–Curtis, Jaccard, Weighted UniFrac, and Unweighted UniFrac distances (**Figure 3**). These global results warranted pairwise comparisons to identify specific treatment-level differences. Compositional differences were detected between treatment 1 and control across all distance metrics except Weighted UniFrac ($q = 0.29$, $R^2 = 0.15$) (**Figure**

3c), and between treatment 2 and control across all metrics (**Figure 3**). No significant differences
were observed between treatments 1 and 2 ($q > 0.05$), suggesting that the LA-I regimen, regardless
of duration of use, may have consistently altered the litter microbiome composition relative to the
control. These compositional shifts were not attributable to differences in within-group variation,
as global estimates of β dispersion were statistically insignificant (ANOVA, $p > 0.05$) for all
distance metrics (**Figure 3**) (Anderson et al., 2006).

Figure 2 Estimation of α diversity with Faith's phylogenetic diversity, observed ASVs (richness), Pielou's evenness, and Shannon diversity. Mean values were discerned following 1,000 iterations of subsampling to a depth of 1,000 reads. Shapiro-Wilks tests were performed to determine if values for each treatment-metric combination were normally distributed, and Levene's test was used to discern if variance was homogenous between treatments for each metric. Mean separation was performed with ANOVA.

<u>Treatment altered core microbiome membership without impacting global diversity</u>

Given the perceived role of the core microbiome in promoting ecological stability (Neu et
al., 2021; Jiao et al., 2022), core taxa were identified as ASVs present in at least all but one sample
across the collective dataset (Mejia et al., 2025). A total of 170 ASVs were detected in the global
core microbiome (**Figure 4a**, **d**). The most represented genus was *Christensenellaceae_R-7_group*
(6 ASVs), followed by eight genera each represented by four ASVs (**Figure 4c**). The most
prevalent families included Lachnospiraceae (19 ASVs), Bacillaceae (16 ASVs), and
Oscillospiraceae (14 ASVs), while dominant orders included Oscillospirales (31 ASVs),
Lactobacillales (25 ASVs), and Micrococcales (21 ASVs) (**Figure 4c**).

Treatment-specific core microbiomes were subsequently characterized, revealing
distinction in ASV presence with LA-I regimens. Across all treatments, 334 core ASVs were
identified, representing the 170 ASVs shared globally and 164 that were treatment specific (**Figure**

4a–b, d). Treatment 1 shared 215 of 334 (64.37%) core ASVs with the control, while treatment 2
shared 184 of 334 (55.09%) (**Figure 4d**). Treatments 1 and 2 shared 189 of 334 (56.59%) core
ASVs (**Figure 4d**). Forty-eight ASVs were unique to the control core microbiome, including ASVs
mapped to *Faecalibacterium* and *Nocardiopsis salina* (**Figure 4b**). Twenty-three ASVs were
unique to treatment 1 and fifteen to treatment 2 (**Figure 4b**). Nineteen ASVs were shared between
treatments 1 and 2 but absent from the control, such as *Lactobacillus aviarius* and *Bifidobacterium*
pullorum (**Figure 4b**). Clustering of ASVs based on Jaccard distances supported these findings,
with greater dissimilarity observed between the control and treatment 2 relative to the control and
treatment 1 (**Figure 4e**).

Figure 3 β diversity assessment leveraging (**a**) Bray-Curtis, (**b**) Jaccard, (**c**) Weighted UniFrac, and (**d**) Unweighted UniFrac distance matrices. Due to statistical significance with global PERMANOVA ($p < 0.05$), pairwise PERMANOVA was used to identify compositional dissimilarity between treatments. Pairwise PERMANOVA and global ẟ dispersion results are in the top right corner of each plot.

Figure 4 Global and treatment-specific core microbiome analysis. (**a**) Relative abundance (log₁₀) heatmap of ASVs represented in one or more core microbiomes. Column annotations reflect

representation in control (red), 1 (yellow), and 2 (black) treatments. (**b**) Relative abundance heatmap of ASVs represented exclusively in control, 1, 2, or 1 and 2 (yellow and black row annotations). (**c**) Top ten genera, families, and orders represented in global and treatment-specific core microbiomes. (**d**) Venn diagram depicting shared and exclusive ASV counts across the treatment-specific core microbiomes. (**e**) Clustering of treatment-specific core microbiomes with Jaccard distances.

At the genus level, shifts in core microbiome composition were evident across treatments. In the control, the most dominant genera included *Clostridia_vadinBB60_group* and *Christensenellaceae_R-7_group* (8 ASVs each), followed by *Faecalibacterium*, *UCG-005*, and *Bacteroides* (7, 6, and 6 ASVs, respectively) (**Figure 4c**). Additional genera such as *RF39*, *Nocardioides*, *Limosilactobacillus*, and *Corynebacterium* were also present, reflecting a diverse community enriched in Firmicutes and Actinobacteria (**Figure 4c**). In treatment 1, *Garciella* and *Christensenellaceae_R-7_group* (6 ASVs each) were the most abundant, followed by eight genera represented by five ASVs each (**Figure 4c**). Treatment 2 was composed primarily of *Christensenellaceae_R-7_group* and *RF39* (7 ASVs each), along with *Garicola*, *Limosilactobacillus*, and *Salinicoccus*, each represented by five ASVs (**Figure 4c**). Treatment-specific core microbiomes displayed similar compositions at the family and order levels (**Figure 4c**).

<u>Differential abundance analysis revealed consistent taxonomic shifts following LA-I treatment</u>

Differential abundance testing was performed at ASV level using log-transformed linear models, with the control treatment specified as a reference. A total of 115 ASVs were differentially abundant ($q < 0.25$) between the LA-I regimens and the control, with 82 identified between treatment 1 and the control (39 enriched, 43 depleted) and 81 between treatment 2 and the control (45 enriched, 36 depleted). Among these, 48 of 115 (41.74%) exhibited consistent directional changes in response to the LA-I application, with 23 enriched and 25 depleted.

To assess the biological relevance of these shifts, abundance matrices were agglomerated at higher taxonomic ranks and reanalyzed using MaAsLin2. Because incomplete taxonomic assignment can bias interpretation at broader ranks, the proportion of ASVs and reads mapped to each taxonomic level was calculated. Taxa were considered assigned if annotations were non-missing and excluded ambiguous terms such as "unknown," "uncultured," "uncultured_bacterium," or "uncultured organism," consistent with the logic applied to heatmap and Venn diagram visualizations. Based on these criteria, 100% of ASVs were assigned at the kingdom level, followed by 99% at phylum and class, 97% at order, 93% at family, 73% at genus, and 15% at species. Read-level assignments followed a similar trend, with 100% of reads mapped through the family level, 91% at genus, and 22% at species. Therefore, order-, family-, and genus-level agglomerations were selected for secondary differential abundance testing.

At the genus level, 20 features were differentially abundant in one or both LA-I treatments relative to the control (**Figure 5a**). Of these, 10 were detected in treatment 1 vs. control (6 enriched,

4 depleted) and 19 in treatment 2 vs. control (16 enriched, 3 depleted), with nine of 20 (45%) shared and showing consistent directional changes (6 enriched, 3 depleted). Genera enriched in both treatments included *Bifidobacterium* (1 vs. ctrl: coef = 3.56, *q* = 0.04; 2 vs. ctrl: coef = 2.26, *q* = 0.01) and *Brevibacterium* (1 vs. ctrl: coef = 1.10, *q* = 0.04; 2 vs. ctrl: coef = 1.09, *q* = 0.14), while *Staphylococcus* was consistently depleted (1 vs. ctrl: coef = –2.68, *q* = 0.16; 2 vs. ctrl: coef = –2.44, *q* = 0.11). The genus-level trend unique to treatment 1 was the depletion of *Campylobacter* (coef = –5.33, *q* = 9e-5), while those exclusive to treatment 2 included enrichment of *Atopostipes* (coef = 0.89, *q* = 0.23), *Halomonas* (coef = 2.38, *q* = 0.11), *Nocardiopsis* (coef = 1.69, *q* = 0.14), *Oceanobacillus* (coef = 0.85, *q* = 0.14), and nine additional genera (**Figure 5a**).

Family-level analysis identified 15 differentially abundant features, with 10 in treatment 1 vs. control (6 enriched, 4 depleted) and 14 in treatment 2 vs. control (11 enriched, 3 depleted). Nine of 15 (60%) were shared and showed consistent directional changes (6 enriched, 3 depleted), largely mirroring genus-level trends (**Figure 5a**). Consistent shifts included the enrichment of Bifidobacteriaceae (1 vs. ctrl: coef = 3.55, *q* = 0.02; 2 vs. ctrl: coef = 2.25, *q* = 0.005), Brevibacteriaceae (1 vs. ctrl: coef = 1.16, *q* = 0.02; 2 vs. ctrl: coef = 1.15, *q* = 0.1), and four additional families, and the depletion of Marinococcaceae (1 vs. ctrl: coef = -4.05, *q* = 0.005; 2 vs. ctrl: coef = -3.94, *q* = 0.01), Micrococcaceae (1 vs. ctrl: coef = -0.35, *q* = 0.02; 2 vs. ctrl: coef = -0.51, *q* = 0.04), and Sutterellaceae (1 vs. ctrl: coef = -3.21, *q* = 0.01; 2 vs. ctrl: coef = -2.71, *q* = 0.1). Furthermore, the treatment 1 vs. control comparison saw the depletion of Campylobacteraceae (coef = –5.26, *q* = 5.3e-5), while treatment 2-specific enrichments included Actinomycetaceae (coef = 1.79, *q* = 0.1), Bogoriellaceae (coef = 1.57, *q* = 0.21), Carnobacteriaceae (coef = 1.0, *q* = 0.11), Nocardiopsaceae (coef = 1.76, *q* = 0.1), and Paenibacillaceae (coef = 1.37, *q* = 0.12) (**Figure 5a**).

At the order level, 14 features were differentially abundant across comparisons. Treatment 1 vs. control accounted for 11 orders (7 enriched, 4 depleted), while 9 were detected in treatment 2 vs. control (7 enriched, 2 depleted). Six of 14 (42.86%) were shared and showed consistent directional change (5 enriched, 1 depleted), with most patterns reflecting shifts observed at lower ranks (**Figure 5a**).

Consistent with β diversity trends, treatments 1 and 2 exhibited broadly similar differential abundance profiles relative to the control. Specifically, 24 of 48 (50%) genus-to-order features (excluding unannotated genera) and 72 of 164 (43.9%) total differentially abundant features overlapped between comparisons (**Figure 5b,c**), with all shared features (72/72) showing consistent directional change (**Figure 5b,c**). Notably, 41 of 164 (25%) and 51 of 164 (31.1%) features were unique to treatment 1 vs. control and treatment 2 vs. control, respectively, indicating potential distinctions between early and sustained exposure to the LA-I regimen (**Figure 5b,c**).

Figure 5 Differential abundance testing of LA-I vs. control treatments. (**a**) Heatmap of differentially abundant features ($q \leq 0.25$) identified at genus, family, and order levels using a log-transformed linear model. (**b**) Venn diagram of all shared and distinct differentially abundant features among treatment comparisons. (**c**) Venn diagram excluding ASVs and genera without taxonomic information (only features in the heatmap).

Discussion

<u>LA-I application restructured the broiler litter microbiome</u>

This work demonstrates how LA-I application, in conjunction with reduced $NaHSO_4$, may alter the structure and composition of the broiler litter microbiome in a commercial setting. Despite no statistically significant differences in α diversity metrics (i.e., richness, evenness, phylogenetic diversity), β diversity analyses revealed that LA-I treatments yielded comparable community profiles that were distinct from the control across all dissimilarity measures. These results align with Do et al. (2024), who reported minimal effects of an organic litter amendment on α diversity despite improved flock performance. In contrast, Johnson et al. (2021) observed increased richness shortly after $NaHSO_4$ application, but found no major changes to bacterial community structure unless topdressing was also applied. The absence of significant differences between the two LA-I

regimens in the present study suggests that a single flock exposure may be sufficient to restructure
the microbial community in a way that persists across subsequent production cycles.

The core microbiome is thought to encompass the most functionally important microbiota
which interact synergistically/antagonistically with transient taxa (Neu et al., 2021; Jiao et al.,
2022). In the present study, the core microbiome remained largely stable across treatments,
although genus-level shifts became more pronounced with prolonged application of LA-I. The
most represented genus in the global core was *Christensenellaceae_R-7_group*, a cellulolytic
bacterium commonly detected in the gastrointestinal tract of livestock and previously linked to
host metabolic traits in beef cattle (Zhang et al., 2025), pigs (Shang et al., 2022), and goats (Liu et
al., 2019). *Christensenellaceae_R-7_group* was also identified as the second most abundant genus
in broiler feces (Shi et al., 2023). However, findings on its role in poultry remain inconsistent.
Higher abundance of *Christensenellaceae_R-7_group* has been associated with improved feed
conversion and residual feed intake (He et al., 2023), yet negative associations have also been
reported with body weight and average daily gain (Xu et al., 2025), alongside reductions in
abundance following dietary supplementation with bio-fermented distillers grain (Xu et al., 2025),
N-carbamylglutamate (Liu et al., 2025), and fungal-derived bioactive compounds (Shi et al.,
2023). These findings suggest that the ecological function of *Christensenellaceae_R-7_group* may
be context-dependent or shaped by interactions with other microorganisms. For instance, in studies
where negative associations with growth were observed, similar patterns were also reported for
ASVs assigned to *Clostridia_vadinBB60_group* and *Oscillospiraceae*, indicating possible co-
occurrence dynamics (Xu et al., 2025). Further investigation using larger sample sizes and
microbial co-occurrence network analysis (Röttjers and Faust, 2018) will be necessary to resolve
the ecological role of *Christensenellaceae_R-7_group* in poultry litter microbial communities.

Various ASVs of potential biological relevance demonstrated exclusivity across treatment-
specific core microbiomes. For example, those exclusively present in LA-I–amended groups
included an ASV encoding the carbohydrate-degrading bacterium *Bifidobacterium pullorum*
(Baseer et al., 2023), an ASV mapped to the family *Paenibacillaceae* (members of which have
demonstrated probiotic potential in poultry [Wu et al., 2019; Ekim et al., 2020; Lee et al., 2025]),
a *Bacillus* ASV (Grant et al., 2018), and an ASV mapped to *Butyricimonas*, a genus positively
associated with broiler growth performance (Johnson et al., 2018; Kairmi et al., 2024). The genera
Salinicoccus and *Atopostipes* were also more represented in the LA-I–amended core microbiomes,
with the former being characteristic of reused poultry litter (Wang et al., 2016; Kubasova et al.,
2022), and both being indicator taxa in active poultry compost (Wang et al., 2022) and represented
in the broiler litter core microbiome of Johnson et al. (2021).

In contrast, the control core microbiome exhibited exclusivity of three *Clostridia UCG-
014* ASVs, a group reported to improve yellow broiler performance and influence tryptophan
metabolism (Dai et al., 2023), while also shown to be significantly enriched following
Campylobacter jejuni challenge (Valečková et al., 2023). The latter pattern is consistent with that
reported by Dai et al. (2023), where *Clostridia UCG-014* co-occurred with *Oscillospiraceae* and
Faecalibacterium, both of which were more abundant in the control core microbiome in the present

study. Similarly, *Faecalibacterium* spp. were enriched in response to NaHSO₄ in the work of Johnson et al. (2021), further supporting trends observed here. The exclusion of *Clostridia UCG-014* and reduced abundance of *Clostridiales vadinBB60 group* in LA-I–amended cores aligns with prior findings showing that both taxa respond to *Campylobacter jejuni* infection (Yan et al., 2021; Ty et al., 2022; Valečková et al., 2023). Notably, *Campylobacter* genus-level depletion was also observed in the present study in treatment 1 relative to the control. Additional trends that warrant further investigation included the exclusivity of an ASV of the broiler probiotic *Lactobacillus reuteri* (Chai et al., 2023) in the treatment 1 core, an ASV mapped to *Helicobacter pullorum* (Abd El-Ghany, 2020) in the treatment 2 core, and the increased representation of *Nocardiopsis* in LA-I–amended cores, a genus known for the synthesis of bioactive compounds (Shi et al., 2022) and that is negatively associated with *Salmonella* abundance in poultry litter (Bucher et al., 2020). These findings collectively suggest that employed LA-I regimens modulate core microbiome composition, enriching taxa with potential probiotic or metabolic relevance while reducing groups associated with pathogen challenge. Future efforts incorporating baseline measurements are needed to validate core microbiome trends.

<u>Differentially abundant taxa reflected functional reorganization of the litter microbiome following LA-I amendment</u>

Differential abundance testing was performed at the ASV, genus, family, and order levels. The decision to agglomerate and analyze at higher taxonomic ranks, rather than infer patterns solely from ASV-level assignments (as done in core microbiome analysis), was guided by the proportion of ASVs and reads that could be confidently mapped at each rank and by recent benchmarking of 16S differential abundance workflows (Wallen et al., 2021; Nearing et al., 2022). To accentuate taxa most likely governed by deterministic selection rather than stochastic or transient signals (Hayashi et al., 2024), features were limited to those comprising the treatment-specific core microbiomes of at least two groups. This filtering ensured that analyses would focus on reproducible, treatment-responsive community members. The following section highlights genus-level patterns associated with LA-I regimens.

The LA-I–amended treatments exhibited consistent increases in *Jeotgalibaca*, a genus positively associated with body weight, feed intake, and peripheral immune function in broilers (Song et al., 2022). *Brevibacterium*, a common inhabitant of poultry litter microbiota (Dumas et al., 2011) and a marker of nutrient cycling, organic matter decomposition, and litter maturation (Wang et al., 2016; Peng et al., 2021; Kubasova et al., 2022), was also enriched. Similarly, *Gottschalkia* was consistently elevated following LA-I amendment. This purinolytic genus degrades avian uric acid to NH_3 under anaerobic conditions, thereby contributing to nitrogen mineralization and early-stage litter decomposition (Poehlein et al., 2017). While uric acid degradation is a significant contributor to NH_3 emissions in poultry litter, *Gottschalkia* is urease-negative and therefore does not contribute to urea hydrolysis, the enzymatic step directly responsible for the production of NH_3 from urea (Rothrock et al., 2010). Its persistence later in the production cycle may reflect the diminished efficacy of NaHSO₄ over time (Hunolt et al., 2015),

as litter pH rebounds and conditions become more favorable for uricolytic activity. Moreover,
enrichment of *Gottschalkia* may be linked to reduced $NaHSO_4$ input in LA-I treatments (relaxing
the acid-mediated suppression of uricase-active taxa) or indirectly tied to the early post-application
NH_3 spike reported with LA-I. Consistent elevation was also observed for *Pseudactinotalea*, a
genus previously linked to poultry litter composting (Chen et al., 2025), and *Bifidobacterium*, a
saccharolytic genus negatively associated with *Campylobacter* in poultry (Valeris-Chacin et al.,
2021) and widely used in commercial probiotics (Halder et al., 2024).

Prolonged LA-I exposure led to the enrichment of several ecologically relevant taxa. These
included the alkaliphilic genera *Alkalibacterium* and *Oceanobacillus*, both identified as broiler
litter hub taxa negatively associated with *Campylobacter* (Valeris-Chacin et al., 2022), with
Oceanobacillus also known to dominate microbial communities during poultry litter composting
(Awasthi et al., 2021; Chen et al., 2024). *Facklamia* was similarly enriched; this genus is among
the most abundant in poultry litter (Lu et al., 2003) and has been positively associated with
peripheral and intestinal mucosal immune indicators in the broiler ileum upon exposure to litter
(Song et al., 2022). Additionally, *Facklamia* abundance was elevated in normal ileal feces
compared to samples exhibiting feed passage, suggesting a potential link to gut microbial stability
(Alvarenga et al., 2023). *Flaviflexus* also showed consistent elevation and has been reported at
higher abundance in *Salmonella* Pullorum-negative hens than in diseased counterparts (Niu et al.,
2025). *Halomonas* was enriched with prolonged LA-I exposure, consistent with its known increase
in biochar-treated poultry litter composts, where it is associated with reduced nitrification and
enhanced NH_4^+ retention (Zainudin et al., 2020). It has also been identified as a core member of
the broiler litter microbiome (Johnson et al., 2021). Given that *Gottschalkia* was also elevated, co-
enrichment with *Halomonas* may reflect cooperation that enhances nitrogen conservation and
suppresses nitrifier activity. Such shifts could improve the fertilizer value of reused litter while
limiting NH_3 volatilization, even under reduced $NaHSO_4$ input. Additional enrichment patterns,
including *Atopostipes* and *Nocardiopsis*, mirrored their presence in the treatment-specific core
microbiomes.

Conversely, LA-I treatment was associated with consistent depletion of the halophilic,
mesophilic genera *Natribacillus* and *Gracilibacillus*, a pattern potentially linked to reduced
$NaHSO_4$ input and a presumable decrease in litter salinity. *Gracilibacillus* has been identified as a
key contributor to humic acid synthesis during the mature phase of poultry litter composting (Yang
et al., 2024; Bucher et al., 2020), and its decline may indicate altered conditions for late-stage
organic matter decomposition. This trend coincided with the enrichment and core microbiome
representation of *Nocardiopsis*, which is also associated with humic acid synthesis during the
composting maturity phase (Yang et al., 2024). The reduction in *Gracilibacillus* likely increased
substrate availability and reduced competitive pressure, facilitating the growth of *Nocardiopsis*
under LA-I conditions. This transition may help sustain humic acid production and enhance the
fertilizer value of reused litter even under modified salinity and acidification dynamics (Sánchez
et al., 2017).

Lastly, *Staphylococcus* was depleted in both LA-I–treated houses compared to the control.
This genus, commonly present in poultry litter and on bird skin and mucosa, includes opportunistic
pathogens such as *S. aureus* with zoonotic potential (Lowder et al., 2009). Its depletion under LA-
I may reflect unfavorable conditions for persistence or competitive exclusion by enriched taxa
such as *Actinomycetaceae. In vitro* assessments of minimum inhibitory concentration (MIC) will
be necessary to determine whether LA-I directly suppresses *Staphylococcus* or exerts indirect
effects via microbial community restructuring.

<u>Implications for litter management and future research</u>
Findings herein demonstrate the potential of microbiome-targeted litter amendments to
restructure poultry litter microbial communities in ways that reduce pathogen load and modulate
ecological function. Taxonomic shifts were largely consistent across LA-I regimens, suggesting
that LA-I establishes a microbial signature early in the flock cycle that persists through to flock
removal.
Given that measurements were taken weeks after application, observed compositional
differences likely reflect early alterations in microbial succession that influenced downstream
community assembly. First, LA-I may have accelerated the thermophilic phase of litter
decomposition, advancing the transition to oxidative and maturation stages. This supposition is
supported by enrichment of *Halomonas* and *Brevibacterium*, genera associated with nitrogen
retention and compost stabilization, as well as *Salinicoccus* and *Atopostipes*, two poultry litter–
compost indicators that presumably played a more prominent role in the core microbiomes of LA-
I–amended groups. *Bifidobacterium*, including *B. pullorum*, was also elevated in LA-I treatments;
while mesophilic, members of this saccharolytic genus produce β-xylanase and preferentially
metabolize xylooligosaccharides, components of rice hull hemicellulose, while contributing to
volatile fatty acid production (Parajó et al., 2004; Baseer et al., 2023). These shifts suggest that
LA-I may promote nitrogen conservation and targeted hemicellulose degradation over prolonged
thermophilic breakdown of complex structural biomass.
This accelerated succession may also contribute to the depletion of *Staphylococcus* and
Campylobacter by flock removal through a combination of elevated thermophilic conditions and
sustained microbial dynamics unfavorable to opportunists (Tuomela et al., 2000). Future work
incorporating temporal sampling, physicochemical profiles (e.g., pH, moisture, elemental
composition), and fungal community dynamics (Manogaran et al., 2022) will be critical for
resolving how early interventions influence microbiome compositional changes linked to litter
decomposition and poultry health under commercial conditions.

<u>Study limitations and future directions</u>
Several limitations should be considered when interpreting the findings of the current
study. First, using composite samples from each house introduced pseudoreplication, limiting the
ability to infer within-treatment variability and reducing statistical power. While distinct microbial
signatures were observed, overlapping compositional profiles between treatments, particularly

between treated houses, suggest that some observed effects may be subtle or subject to external
influences. Additionally, changes attributed to LA-I administration may partly reflect the gradual
removal or reduced top-dressing of $NaHSO_4$ over time, as indicated in previous studies
demonstrating microbial shifts following the withdrawal of conventional litter acidifiers (Johnson
et al., 2021). This complicates the attribution of microbiome changes solely to treatment.

Another important consideration is the absence of performance and physicochemical data
(e.g., ammonia concentration, moisture content, bird productivity), which are necessary to
contextualize microbiome shifts regarding functional or production-related outcomes. Ammonia-
binding products are well-documented for their importance in maintaining poultry welfare and
respiratory health (Johnson et al., 2011), and without these benchmarks, it is difficult to evaluate
the practical impact of LA-I fully.

Moreover, microbiome composition is susceptible to environmental factors such as litter
age, sampling time, and seasonality. These variables were not systematically controlled or
analyzed in this study and could influence the generalizability of the findings. Future studies
should incorporate replication at the flock or pen level, longitudinal sampling across production
cycles, and concurrent measurement of litter chemistry and bird performance metrics. *In vitro* litter
models could also help isolate the effects of specific amendment components under controlled
conditions. Collectively, these approaches will enable a more mechanistic understanding of how
LA-I can shape poultry litter microbiomes and whether those changes confer meaningful
improvements to poultry health and environmental outcomes.

Conclusion

The current study demonstrated that integrating a prebiotic litter amendment (LA-I) with
reduced $NaHSO_4$ application significantly altered the microbial composition of poultry litter,
promoting shifts toward taxa potentially associated with nutrient cycling, organic matter
degradation, and reduced pathogen presence. While α diversity remained stable, consistent
changes in community structure and differential abundance patterns were observed between
treated and control houses. These findings highlight the potential for targeted litter amendments to
modulate microbial communities in ways that enhance environmental sustainability and poultry
health. Complementing $NaHSO_4$-based acidifiers with IndigoLT™ may offer a synergistic
strategy to promote microbiome stability, accelerate litter decomposition, and reduce NH_3
volatilization, thereby improving litter quality and overall flock performance. Future work should
include optimizing inclusion rates most complementary between $NaHSO_4$ and IndigoLT™ to
maximize efficacy across production systems.

Data availability

The FASTQ files associated with this work have been deposited in the NCBI Sequence
Read Archive database under submission SUB15391559.

Conflict of Interest

Authors declare there are no competing interests. BH is employed by AgriGro (Doniphan, MO, United States) but his affiliation did not influence the results of the current work.

References

1. Bolan, N. S., Szogi, A. A., Chuasavathi, T., Seshadri, B., Rothrock Jr, M. J., & Panneerselvam, P. (2010). Uses and management of poultry litter. *World's Poultry Science Journal*, *66*(4), 673-698. https://doi.org/10.1017/S0043933910000656

2. Pedroso, A. A., Hurley-Bacon, A. L., Zedek, A. S., Kwan, T. W., Jordan, A. P., Avellaneda, G., ... & Lee, M. D. (2013). Can probiotics improve the environmental microbiome and resistome of commercial poultry production?. International journal of environmental research and public health, 10(10), 4534-4559. https://doi.org/10.3390/ijerph10104534

3. Ritz, C. W., Fairchild, B. D., & Lacy, M. P. (2004). Implications of ammonia production and emissions from commercial poultry facilities: A review. *Journal of applied poultry research*, *13*(4), 684-692. https://doi.org/10.1093/japr/13.4.684

4. Rothrock, M. J., Cook, K. L., Warren, J. G., Eiteman, M. A., & Sistani, K. (2010). Microbial mineralization of organic nitrogen forms in poultry litters. Journal of environmental quality, 39(5), 1848-1857. https://doi.org/10.2134/jeq2010.0024

5. Johnson, T. M., Fairchild, B., & Ritz, C. W. (2011). The use of sodium bisulfate as a best management practice for reducing ammonia emissions from poultry manures.

6. Toppel, K., Kaufmann, F., Schön, H., Gauly, M., & Andersson, R. (2019). Effect of pH-lowering litter amendment on animal-based welfare indicators and litter quality in a European commercial broiler husbandry. *Poultry Science*, *98*(3), 1181-1189. https://doi.org/10.3382/ps/pey489

7. de Toledo, T. D. S., Roll, A. A. P., Rutz, F., Dallmann, H. M., Dai Pra, M. A., Leite, F. P. L., & Roll, V. F. B. (2020). An assessment of the impacts of litter treatments on the litter quality and broiler performance: A systematic review and meta-analysis. *PLoS One*, *15*(5), e0232853. https://doi.org/10.1371/journal.pone.0232853

8. Johnson, J., Zwirzitz, B., Oladeinde, A., Milfort, M., Looft, T., Chai, L., ... & Aggrey, S. E. (2021). *Succession patterns of the bacterial community in poultry litter after bird removal and sodium bisulfate application* (Vol. 50, No. 4, pp. 923-933). https://doi.org/10.1002/jeq2.20248

9. Williams, Z. T., Blake, J. P., & Macklin, K. S. (2012). The effect of sodium bisulfate on Salmonella viability in broiler litter. *Poultry Science*, *91*(9), 2083-2088. https://doi.org/10.3382/ps.2011-01976

10. Joerger, R. D., Ganguly, A., de Los Santos, M., & Li, H. (2020). Effect of sodium bisulfate amendments on bacterial populations in broiler litter. *Poultry Science*, *99*(11), 5560-5571. https://doi.org/10.1016/j.psj.2020.08.013

11. Bucher, M. G., Zwirzitz, B., Oladeinde, A., Cook, K., Plymel, C., Zock, G., ... & Sistani, K. R. (2020). *Reused poultry litter microbiome with competitive exclusion potential against Salmonella Heidelberg* (Vol. 49, No. 4, pp. 869-881). https://doi.org/10.1002/jeq2.20081

12. Swelum, A. A., El-Saadony, M. T., Abd El-Hack, M. E., Ghanima, M. M. A., Shukry, M., Alhotan, R. A., ... & El-Tarabily, K. A. (2021). Ammonia emissions in poultry houses and microbial nitrification as a promising reduction strategy. *Science of The Total Environment*, *781*, 146978. https://doi.org/10.1016/j.scitotenv.2021.146978

13. Callaway, T.R. and S.C. Ricke (Eds.), 2023. *Direct Fed Microbials/Prebiotics for Animals: Science and Mechanisms of Action*, 2^nd Edition. Springer Science, New York, NY, 348 pp.

14. Ricke, S.C. 2018. Impact of prebiotics on poultry production and food safety. Yale Journal of Biology and Medicine. 91: 151-159.

15. Swanson, K.S., K. Allenspach, G. Amos, T.A. Auchtung, S.A. Bassett, C.R. Bjørnvad, N. Everaert, S.M. Martín-Orúe, S.C. Ricke, E.P. Ryan, and G.C. Fahey, Jr. 2025. Use of biotics in animals: Impact on nutrition, health, and food production. J. Animal Sci. 103: ska061. doi.org/10.1093/jas/skaf061.

16. Reuben, R. C., Sarkar, S. L., Roy, P. C., Anwar, A., Hossain, M. A., & Jahid, I. K. (2021). Prebiotics, probiotics and postbiotics for sustainable poultry production. *World's Poultry Science Journal*, *77*(4), 825-882. https://doi.org/10.1080/00439339.2021.1960234

17. Gilchrist, M. J., Greko, C., Wallinga, D. B., Beran, G. W., Riley, D. G., & Thorne, P. S. (2007). The potential role of concentrated animal feeding operations in infectious disease epidemics and antibiotic resistance. *Environmental health perspectives*, *115*(2), 313-316. https://doi.org/10.1289/ehp.8837

18. Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., ... & Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the national academy of sciences*, *108*(supplement_1), 4516-4522. https://doi.org/10.1073/pnas.1000080107

19. Walters, W. A., Caporaso, J. G., Lauber, C. L., Berg-Lyons, D., Fierer, N., & Knight, R. (2011). PrimerProspector: de novo design and taxonomic analysis of barcoded polymerase chain reaction primers. Bioinformatics, 27(8), 1159-1161. https://doi.org/10.1093/bioinformatics/btr087

20. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet. journal*, *17*(1), 10-12. https://doi.org/10.14806/ej.17.1.200

21. Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods*, *13*(7), 581-583. https://doi.org/10.1038/nmeth.3869

22. Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ... & Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome

data science using QIIME 2. *Nature biotechnology*, *37*(8), 852-857. https://doi.org/10.1038/s41587-019-0209-9

23. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic acids research*, *41*(D1), D590-D596. https://doi.org/10.1093/nar/gks1219

24. Team, R. C. (2020). RA language and environment for statistical computing, R Foundation for Statistical. *Computing*.

25. Bisanz, J. E. (2018). qiime2R: Importing QIIME2 artifacts and associated data into R sessions. *Version 0.99*, *13*.

26. McMurdie, P. J., & Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS one*, *8*(4), e61217. https://doi.org/10.1371/journal.pone.0061217

27. Mikryukov, V. (2020). metagMisc: Miscellaneous functions for metagenomic analysis. R package version 0.0. 4.

28. Good, I. J. (1953). The population frequencies of species and the estimation of population parameters. *Biometrika*, *40*(3-4), 237-264. https://doi.org/10.1093/biomet/40.3-4.237

29. Good, I. J., & Toulmin, G. H. (1956). The number of new species, and the increase in population coverage, when a sample is increased. *Biometrika*, *43*(1-2), 45-63. https://doi.org/10.1093/biomet/43.1-2.45

30. Chao, A., & Jost, L. (2012). Coverage-based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology*, *93*(12), 2533-2547. https://doi.org/10.1890/11-1952.1

31. Dixon, P. (2003). VEGAN, a package of R functions for community ecology. *Journal of vegetation science*, *14*(6), 927-930. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

32. Wickham, H., & Sievert, C. (2009). *ggplot2: elegant graphics for data analysis* (Vol. 10, pp. 978-0). New York: springer. http://dx.doi.org/10.18637/jss.v077.b02

33. Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics*, *20*(2), 289-290. https://doi.org/10.1093/bioinformatics/btg412

34. Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T. Y. (2017). ggtree: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution*, *8*(1), 28-36. https://doi.org/10.1111/2041-210X.12628

35. Hill, M. O. (1973). Diversity and evenness: a unifying notation and its consequences. *Ecology*, *54*(2), 427-432. https://doi.org/10.2307/1934352

36. Pielou, E. C. (1966). The measurement of diversity in different types of biological
collections. *Journal of theoretical biology*, *13*, 131-144. https://doi.org/10.1016/0022-
5193(66)90013-0

37. Lahti, L., & Shetty, S. (2018). Introduction to the microbiome R package. *Preprint at
https://microbiome. github. io/tutorials*.

38. Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biological
conservation*, *61*(1), 1-10. https://doi.org/10.1016/0006-3207(92)91201-3

39. Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D.
D., ... & Webb, C. O. (2010). Picante: R tools for integrating phylogenies and ecology.
Bioinformatics, *26*(11), 1463-1464. https://doi.org/10.1093/bioinformatics/btq166

40. Levene, H. (1960). Robust tests for equality of variances. *Contributions to probability and
statistics*, 278-292.

41. Fox, J., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., ... & Heiberger, R.
(2012). Package 'car'. *Vienna: R Foundation for Statistical Computing*, *16*(332), 333.

42. Shapiro, S. S., & Wilk, M. B. (1965). An analysis of variance test for normality (complete
samples). *Biometrika*, *52*(3-4), 591-611. https://doi.org/10.1093/biomet/52.3-4.591

43. Paulson, J. N., Stine, O. C., Bravo, H. C., & Pop, M. (2013). Differential abundance
analysis for microbial marker-gene surveys. *Nature methods*, *10*(12), 1200-1202.
https://doi.org/10.1038/nmeth.2658

44. Bray, J. R., & Curtis, J. T. (1957). An ordination of the upland forest communities of
southern   Wisconsin.   *Ecological   monographs*,   *27*(4),   326-349.
https://doi.org/10.2307/1942268

45. Jaccard, P. (1908). Nouvelles recherches sur la distribution florale. *Bull. Soc. Vaud. Sci.
Nat.*, *44*, 223-270.

46. Lozupone, C., & Knight, R. (2005). UniFrac: a new phylogenetic method for comparing
microbial communities. *Applied and environmental microbiology*, *71*(12), 8228-8235.
https://doi.org/10.1128/AEM.71.12.8228-8235.2005

47. Martinez Arbizu, P. (2020). pairwiseAdonis: Pairwise multilevel comparison using adonis.
R package version 0.4, *1*.

48. Mejia, G., Jara-Servin, A., Hernández-Álvarez, C., Romero-Chora, L., Peimbert, M., Cruz-
Ortega, R., & Alcaraz, L. D. (2025). Rhizosphere microbiome influence on tomato growth
under   low-nutrient   settings.   *FEMS   Microbiology   Ecology*,   *101*(3),   fiaf019.
https://doi.org/10.1093/femsec/fiaf019

49. Gu, Z., Eils, R., & Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. *Bioinformatics*, *32*(18), 2847-2849.
https://doi.org/10.1093/bioinformatics/btw313

50. Yan, L., & Yan, M. L. (2021, June). *Package "ggvenn."*.

51. Mallick, H., Rahnavard, A., McIver, L. J., Ma, S., Zhang, Y., Nguyen, L. H., ... &
Huttenhower, C. (2021). Multivariable association discovery in population-scale meta-
omics studies. *PLoS computational biology*, *17*(11), e1009442.
https://doi.org/10.1371/journal.pcbi.1009442

52. Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and
powerful approach to multiple testing. *Journal of the Royal statistical society: series B
(Methodological)*, *57*(1), 289-300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

53. Boolchandani, M., Blake, K. S., Tilley, D. H., Cabada, M. M., Schwartz, D. J., Patel, S., ...
& Dantas, G. (2022). Impact of international travel and diarrhea on gut microbiome and
resistome dynamics. *Nature communications*, *13*(1), 7485. https://doi.org/10.1038/s41467-
022-34862-w

54. Reasoner, S. A., Bernard, R., Waalkes, A., Penewit, K., Lewis, J., Sokolow, A. G., ... &
Nicholson, M. R. (2024). Longitudinal profiling of the intestinal microbiome in children
with cystic fibrosis treated with elexacaftor-tezacaftor-ivacaftor. *Mbio*, *15*(2), e01935-23.
https://doi.org/10.1128/mbio.01935-23

55. Neu, A. T., Allen, E. E., & Roy, K. (2021). Defining and quantifying the core microbiome:
challenges and prospects. *Proceedings of the National Academy of Sciences*, *118*(51),
e2104429118. https://doi.org/10.1073/pnas.2104429118

56. Jiao, S., Chen, W., & Wei, G. (2022). Core microbiota drive functional stability of soil
microbiome in reforestation ecosystems. *Global Change Biology*, *28*(3), 1038-1047.
https://doi.org/10.1111/gcb.16024

57. Do, A. D. T., Lozano, A., Van Laar, T. A., Mero, R., Lopez, C., Hisasaga, C., ... & Tarrant,
K. J. (2024). Evaluating microbiome patterns, microbial species, and leg health associated
with reused litter in a commercial broiler barn. *Journal of Applied Poultry Research*, *33*(4),
100490. https://doi.org/10.1016/j.japr.2024.100490

58. Zhang, R., Mei, S., He, G., Wei, M., Chen, L., Chen, Z., ... & Chen, C. (2025). Multi-omics
analyses reveal fecal microbial community and metabolic alterations in finishing cattle fed
probiotics-fermented distiller's grains diets. *Microbiology Spectrum*, *13*(5), e00721-24.
https://doi.org/10.1128/spectrum.00721-24

59. Shang, P., Wei, M., Duan, M., Yan, F., & Chamba, Y. (2022). Healthy gut microbiome
composition enhances disease resistance and fat deposition in Tibetan pigs. *Frontiers in
Microbiology*, *13*, 965292. https://doi.org/10.3389/fmicb.2022.965292

60. Liu, J., Xue, C., Sun, D., Zhu, W., & Mao, S. (2019). Impact of high-grain diet feeding on
mucosa-associated bacterial community and gene expression of tight junction proteins in
the small intestine of goats. *Microbiologyopen*, *8*(6), e00745.
https://doi.org/10.1002/mbo3.745

61. Shi, C., Guo, L., Yang, H., Leng, X., Deng, P., Bi, J., & Wang, Y. (2023). Effect of dietary supplementation of Ganoderma lucidum residue on growth performance, immune organ index and faecal microbial community diversity in broiler chickens. https://doi.org/10.21203/rs.3.rs-2976135/v1

62. He, Z., Liu, R., Wang, M., Wang, Q., Zheng, J., Ding, J., ... & Zhao, G. (2023). Combined effect of microbially derived cecal SCFA and host genetics on feed efficiency in broiler chickens. *Microbiome*, *11*(1), 198. https://doi.org/10.1186/s40168-023-01627-6

63. Xu, P., Liu, C., Ding, H., Chen, P., Fan, X., Wang, X., ... & Xiao, Y. (2025). Effect of Bio-Fermented Distillers Grain on Growth, Intestines, and Caecal Microbial Community in Broilers. *Fermentation*, *11*(3), 118. https://doi.org/10.3390/fermentation11030118

64. Liu, N., Zhang, Z., Zhang, J., Ma, W., & Wang, C. (2025). Effects of N-Carbamylglutamate supplementation on cecal morphology, microbiota composition, and short-chain fatty acids contents of broiler breeder roosters. *Scientific Reports*, *15*(1), 7489. https://doi.org/10.1038/s41598-025-91577-w

65. Röttjers, L., & Faust, K. (2018). From hairballs to hypotheses–biological insights from microbial networks. *FEMS microbiology reviews*, *42*(6), 761-780. https://doi.org/10.1093/femsre/fuy030

66. Baseer, A. Q., Mushfiq, S., Monib, A. W., Hassand, M. H., & Niazi, P. (2023). Computational Structural Analysis and Homology Modelling of Beta-Xylanase from Bifidobacterium pullorum: A Comprehensive In-Silico Investigation. *Computational Structural Analysis and Homology Modelling of Beta-Xylanase from Bifidobacterium pullorum: A Comprehensive In-Silico Investigation.*, (6). https://doi.org/10.55544/jrasb.2.6.9

67. Wu, Y., Wang, B., Zeng, Z., Liu, R., Tang, L., Gong, L., & Li, W. (2019). Effects of probiotics Lactobacillus plantarum 16 and Paenibacillus polymyxa 10 on intestinal barrier function, antioxidative capacity, apoptosis, immune response, and biochemical parameters in broilers. *Poultry Science*, *98*(10), 5028-5039. https://doi.org/10.3382/ps/pez226

68. Ekim, B., Calik, A., Ceylan, A., & Saçaklı, P. (2020). Effects of Paenibacillus xylanexedens on growth performance, intestinal histomorphology, intestinal microflora, and immune response in broiler chickens challenged with Escherichia coli K88. *Poultry science*, *99*(1), 214-223. https://doi.org/10.3382/ps/pez460

69. Lee, A. R., Kim, S. H., & Kim, S. K. (2025). Effects of supplementation with a novel strain of Paenibacillus konkukensis on laying performance, tibial characteristics, and caecal microbiota in laying hens. *Italian Journal of Animal Science*, *24*(1), 827-841. https://doi.org/10.1080/1828051X.2025.2477029

70. Grant, A. Q., Gay, C. G., & Lillehoj, H. S. (2018). Bacillus spp. as direct-fed microbial antibiotic alternatives to enhance growth, immunity, and gut health in poultry. *Avian Pathology*, *47*(4), 339-351. https://doi.org/10.1080/03079457.2018.1464117

71. Johnson, T. J., Youmans, B. P., Noll, S., Cardona, C., Evans, N. P., Karnezos, T. P., ... & Lee, C. W. (2018). A consistent and predictable commercial broiler chicken bacterial microbiota in antibiotic-free production displays strong correlations with performance. *Applied and environmental microbiology*, *84*(12), e00362-18. https://doi.org/10.1128/AEM.00362-18

72. Kairmi, S. H., Abdelaziz, K., Spahany, H., Astill, J., Trott, D., Wang, B., ... & Sharif, S. (2024). Intestinal microbiome profiles in broiler chickens raised without antibiotics exhibit altered microbiome dynamics relative to conventionally raised chickens. *Plos one*, *19*(4), e0301110. https://doi.org/10.1371/journal.pone.0301110

73. Wang, L., Lilburn, M., & Yu, Z. (2016). Intestinal microbiota of broiler chickens as affected by litter management regimens. *Frontiers in microbiology*, *7*, 593. https://doi.org/10.3389/fmicb.2016.00593

74. Kubasova, T., Faldynova, M., Crhanova, M., Karasova, D., Zeman, M., Babak, V., & Rychlik, I. (2022). Succession, replacement, and modification of chicken litter microbiota. *Applied and Environmental Microbiology*, *88*(24), e01809-22. https://doi.org/10.1128/aem.01809-22

75. Wang, H., Shankar, V., & Jiang, X. (2022). Compositional and functional changes in microbial communities of composts due to the composting-related factors and the presence of Listeria monocytogenes. *Microbiology Spectrum*, *10*(4), e01845-21. https://doi.org/10.1128/spectrum.01845-21

76. Dai, Z., Wang, X., Liu, Y., Liu, J., Xiao, S., Yang, C., & Zhong, Y. (2023). Effects of dietary microcapsule sustained-release sodium butyrate on the growth performance, immunity, and gut microbiota of yellow broilers. *Animals*, *13*(23), 3598. https://doi.org/10.3390/ani13233598

77. Valečková, E., Sun, L., Wang, H., Dube, F., Ivarsson, E., Kasmaei, K. M., ... & Wall, H. (2023). Intestinal colonization with Campylobacter jejuni affects broiler gut microbiota composition but is not inhibited by daily intake of Lactiplantibacillus plantarum. *Frontiers in Microbiology*, *14*, 1205797. https://doi.org/10.3389/fmicb.2023.1205797

78. Yan, W., Zhou, Q., Yuan, Z., Fu, L., Wen, C., Yang, N., & Sun, C. (2021). Impact of the gut microecology on Campylobacter presence revealed by comparisons of the gut microbiota from chickens raised on litter or in individual cages. *BMC microbiology*, *21*, 1-15. https://doi.org/10.1186/s12866-021-02353-5

79. Ty, M., Taha-Abdelaziz, K., Demey, V., Castex, M., Sharif, S., & Parkinson, J. (2022). Performance of distinct microbial based solutions in a Campylobacter infection challenge model in poultry. *Animal microbiome*, *4*, 1-19. https://doi.org/10.1186/s42523-021-00157-6

80. Chai, C., Guo, Y., Mohamed, T., Bumbie, G. Z., Wang, Y., Zeng, X., ... & Sun, W. (2023). Dietary Lactobacillus reuteri Sl001 improves growth performance, health-related parameters, intestinal morphology and microbiota of broiler chickens. *Animals*, *13*(10), 1690. https://doi.org/10.3390/ani13101690

81. Abd El-Ghany, W. A. (2020). Helicobacter pullorum: A potential hurdle emerging pathogen for public health. *The Journal of Infection in Developing Countries*, *14*(11), 1225-1230. https://doi.org/10.3855/jidc.12843

82. Shi, T., Wang, Y. F., Wang, H., & Wang, B. (2022). Genus Nocardiopsis: a prolific producer of natural products. *Marine Drugs*, *20*(6), 374. https://doi.org/10.3390/md20060374

83. Wallen, Z. D. (2021). Comparison study of differential abundance testing methods using two large Parkinson disease gut microbiome datasets derived from 16S amplicon sequencing. *BMC bioinformatics*, *22*(1), 265. https://doi.org/10.1186/s12859-021-04193-6

84. Nearing, J. T., Douglas, G. M., Hayes, M. G., MacDonald, J., Desai, D. K., Allward, N., ... & Langille, M. G. (2022). Microbiome differential abundance methods produce different results across 38 datasets. *Nature communications*, *13*(1), 342. https://doi.org/10.1038/s41467-022-28034-z

85. Hayashi, I., Fujita, H., & Toju, H. (2024). Deterministic and stochastic processes generating alternative states of microbiomes. *ISME communications*, *4*(1), ycae007. https://doi.org/10.1093/ismeco/ycae007

86. Song, B., Li, P., Xu, H., Wang, Z., Yuan, J., Zhang, B., ... & Guo, Y. (2022). Effects of rearing system and antibiotic treatment on immune function, gut microbiota and metabolites of broiler chickens. *Journal of Animal Science and Biotechnology*, *13*(1), 144. https://doi.org/10.1186/s40104-022-00788-y

87. Dumas, M. D., Polson, S. W., Ritter, D., Ravel, J., Gelb Jr, J., Morgan, R., & Wommack, K. E. (2011). Impacts of poultry house environment on poultry litter bacterial community composition. *PLoS One*, *6*(9), e24785. https://doi.org/10.1371/journal.pone.0024785

88. Peng, M., Tabashsum, Z., Millner, P., Parveen, S., & Biswas, D. (2021). Influence of manure application on the soil bacterial microbiome in integrated crop-livestock farms in Maryland. *Microorganisms*, *9*(12), 2586. https://doi.org/10.3390/microorganisms9122586

89. Poehlein, A., Yutin, N., Daniel, R., & Galperin, M. Y. (2017). Proposal for the reclassification of obligately purine-fermenting bacteria Clostridium acidurici (Barker 1938) and Clostridium purinilyticum (Dürre et al. 1981) as Gottschalkia acidurici gen. nov. comb. nov. and Gottschalkia purinilytica comb. nov. and of Eubacterium angustum (Beuscher and Andreesen 1985) as Andreesenia angusta gen. nov. comb. nov. in the family Gottschalkiaceae fam. nov. *International journal of systematic and evolutionary microbiology*, *67*(8), 2711-2719. https://doi.org/10.1099/ijsem.0.002008

90. Hunolt, A. E., Maguire, R. O., Ogejo, J. A., Badgley, B. D., Frame, W. H., & Reiter, M. S. (2015). Multiple applications of sodium bisulfate to broiler litter affect ammonia release and litter properties. *Journal of environmental quality*, *44*(6), 1903-1910. https://doi.org/10.2134/jeq2015.05.0214

91. Chen, X., Cai, D., Shen, X., Liao, Y., Ren, D., & Ding, T. (2025). Exploring the changes in chicken manure bacterial community structure and function after antibiotic and composting: Implications for food safety. *Food and Humanity*, *4*, 100571. https://doi.org/10.1016/j.foohum.2025.100571

92. Valeris-Chacin, R., Pieters, M., Hwang, H., Johnson, T. J., & Singer, R. S. (2021). Association of broiler litter microbiome composition and Campylobacter isolation. *Frontiers in Veterinary Science*, *8*, 654927. https://doi.org/10.3389/fvets.2021.654927

93. Halder, N., Sunder, J., De, A. K., Bhattacharya, D., & Joardar, S. N. (2024). Probiotics in poultry: a comprehensive review. *The Journal of Basic and Applied Zoology*, *85*(1), 23. https://doi.org/10.1186/s41936-024-00379-5

94. Valeris-Chacin, R., Weber, B., Johnson, T. J., Pieters, M., & Singer, R. S. (2022). Longitudinal changes in Campylobacter and the litter microbiome throughout the broiler production cycle. *Applied and environmental microbiology*, *88*(17), e00667-22. https://doi.org/10.1128/aem.00667-22

95. Awasthi, S. K., Duan, Y., Liu, T., Zhang, Z., Pandey, A., Varjani, S., ... & Taherzadeh, M. J. (2021). Can biochar regulate the fate of heavy metals (Cu and Zn) resistant bacteria community during the poultry manure composting?. *Journal of Hazardous Materials*, *406*, 124593. https://doi.org/10.1016/j.jhazmat.2020.124593

96. Chen, L., Han, M., Xu, J., Cao, Z., Chen, W., Jing, B., ... & Wen, X. (2024). Firmicutes primarily drive odor emission profiles in poultry manure treatments. *Poultry Science*, *103*(11), 104250. https://doi.org/10.1016/j.psj.2024.104250

97. Lu, J., Sanchez, S., Hofacre, C., Maurer, J. J., Harmon, B. G., & Lee, M. D. (2003). Evaluation of broiler litter with reference to the microbial composition as assessed by using 16S rRNA and functional gene markers. *Applied and environmental microbiology*, *69*(2), 901-908. https://doi.org/10.1128/AEM.69.2.901-908.2003

98. Song, B., Yan, S., Li, P., Li, G., Gao, M., Yan, L., ... & Guo, Y. (2022). Comparison and correlation analysis of immune function and gut microbiota of broiler chickens raised in double-layer cages and litter floor pens. *Microbiology Spectrum*, *10*(4), e00045-22. https://doi.org/10.1128/spectrum.00045-22

99. Alvarenga, B. O., Paiva, J. B., Souza, A. I., Rodrigues, D. R., Tizioto, P. C., & Ferreira, A. J. P. (2023). Metagenomics analysis of the morphological aspects and bacterial composition of broiler feces. *Poultry Science*, *102*(2), 102401. https://doi.org/10.1016/j.psj.2022.102401

100. Niu, Q., Yang, K., Zhou, Z., Huang, Q., & Wang, J. (2025). Intergenerational Transmission of Gut Microbiome from Infected and Non-Infected Salmonella pullorum Hens. *Microorganisms*, *13*(3), 640. https://doi.org/10.3390/microorganisms13030640

101. Zainudin, M. H., Mustapha, N. A., Maeda, T., Ramli, N., Sakai, K., & Hassan, M. (2020). Biochar enhanced the nitrifying and denitrifying bacterial communities during the

composting of poultry manure and rice straw. *Waste Management*, *106*, 240-249. https://doi.org/10.1016/j.wasman.2020.03.029

102.   Yang, J., Du, Z., Huang, C., Li, W., Xi, B., Zhu, L., & Wu, X. (2024). Dynamics of microbial functional guilds involved in the humification process during aerobic composting of chicken manure on an industrial scale. *Environmental Science and Pollution Research*, *31*(14), 21044-21056. https://doi.org/10.1007/s11356-024-32390-2

103.   Sánchez, Ó. J., Ospina, D. A., & Montoya, S. (2017). Compost supplementation with nutrients and microorganisms in composting process. *Waste management*, *69*, 136-153. https://doi.org/10.1016/j.wasman.2017.08.012

104.   Lowder, B. V., Guinane, C. M., Ben Zakour, N. L., Weinert, L. A., Conway-Morris, A., Cartwright, R. A., ... & Fitzgerald, J. R. (2009). Recent human-to-poultry host jump, adaptation, and pandemic spread of Staphylococcus aureus. *Proceedings of the National Academy of Sciences*, *106*(46), 19545-19550. https://doi.org/10.1073/pnas.0909285106

105.   Parajó, J. C., Garrote, G., Cruz, J. M., & Dominguez, H. (2004). Production of xylooligosaccharides by autohydrolysis of lignocellulosic materials. *Trends in Food Science & Technology*, *15*(3-4), 115-120. https://doi.org/10.1016/j.tifs.2003.09.009

106.   Tuomela, M., Vikman, M., Hatakka, A., & Itävaara, M. (2000). Biodegradation of lignin in a compost environment: a review. *Bioresource technology*, *72*(2), 169-183. https://doi.org/10.1016/S0960-8524(99)00104-2

107.   Manogaran, M. D., Shamsuddin, R., Yusoff, M. H. M., Lay, M., & Siyal, A. A. (2022). A review on treatment processes of chicken manure. *Cleaner and Circular Bioeconomy*, *2*, 100013. https://doi.org/10.1016/j.clcb.2022.100013