Whole-genome sequencing of CRFK and PG-4 cells to infer the phenotype of the original

donor cats

Ganma Tanaka[1]*, Rikuto Goto[1]*, Tetsushi Komoto[1,2], Akiko Kubota[1], Reo Hayashi[1], Takeshi

Igawa[1,2], Naoaki Sakamoto[1,3], Akinori Awazu[1,3]

[1]Graduate School of Integrated Sciences for Life, Hiroshima University, Higashihiroshima,

Hiroshima, Japan

[2]Amphibian Research Center, Hiroshima University, Higashihiroshima, Hiroshima, Japan

[3]Research Center for the Mathematics on Chromatin Live Dynamics, Hiroshima University,

Higashihiroshima, Hiroshima, Japan

* Ganma Tanaka and Rikuto Goto contributed equally to this work and share first authorship.

Corresponding author:

Akinori Awazu

Graduate School of Integrated Sciences for Life, Hiroshima University, 1-3-1, Kagamiyama,

Higashi-Hiroshima 739-8526, JAPAN,

email address: awa@hiroshima-u.ac.jp

Abstract

Background Crandell-Rees Feline Kidney (CRFK; kidney-derived) cells and PG-4 cells

(astrocyte-derived) have been in use and have been passaged for decades in laboratories

worldwide; however, no detailed information on the genetic background of the donor

individuals is available, particularly regarding phenotype characteristics such as coat and iris

color.

Results We performed whole-genome sequencing of CRFK and PG-4 cells. We analyzed the

resulting data to infer the phenotype of the individual from which the cells were derived,

specifically for the coat color, coat length, coat pattern, and iris color. Our data suggested that

CRFK cells originated from a cat with long black fur lacking stripes and with non-blue irises;

PG-4 cells originated from a cat with long bicolored white and black fur without stripes, and

with non-blue irises. Analysis of publicly available RNA-seq data confirmed that genes

associated with coat phenotype and iris color are expressed in the skin and eyes, as well as in

various other organs.

Conclusions Variants of the genes affecting coat phenotype and iris color may influence

physiological functions throughout the body. These results shed light on the previously

unknown genetic background of commonly used feline cultured cells and the phenotype of the

donor individuals. These findings may facilitate more accurate interpretation of data obtained

from cultured feline cells and provide guidelines for developing cell lines with domestic cat

genotypes exhibiting diverse phenotypes through genome editing. This will help to provide

clarity regarding deafness in cats with white fur and blue irises and elucidate the effects of

melanocyte destruction caused by KIT gene mutations on the nervous system and the influence

of other coat-related factors in organs other than the skin or functions independent of coat

formation.

Keywords: genomics; coat pattern; iris color; RNA sequencing;

Background

Domestic cats (*Felis catus*) are widely kept as companion animals around the world, and the phenotypic diversity of their coat color, coat pattern, coat length, and iris color has been a topic of interest to breeders and scientists alike. Recent advances in genomics have markedly improved our understanding of the genetic basis controlling these diverse phenotypes, and the respective genotype-phenotype relationships are being increasingly elucidated.[1] Examples include the *TYR* gene, which is the color locus associated with albinism and a color point phenotype[2–4]; the *TYRP1* gene, conveying brown coat color[2,5]; the *MC1R* genes, associated with in red coat color[6]; the *KIT* gene, determining white coat and white spotting[7–11]; the *ARHGAP36* gene, involved in orange coat color[12,13]; the *MLPH* gene, causing dilute pigmentation[14]; the *ASIP* gene, controlling the expression of the agouti pattern[15]; the *LVRN* gene, determining the type of tabby pattern[16]; the *DKK4* gene, associated with the ticked tabby pattern[17]; the *FGF5* gene, involved in long-fur variants[18,19]; the *KRT71* and *LPAR6* genes, involved in hairless and rexing (curly hair) phenotypes[20–22]; and the *PAX3* gene, which is associated with blue irises[23,24]. Furthermore, technological advancements such as long-read sequencing facilitated the construction of a more continuous and accurate draft genome of the domestic cat[25–27], and a plethora of genetic polymorphisms associated with disease have been identified[17,25,28–32].

Immortalized cell lines are an indispensable tool in current biomedical and veterinary science. Crandell-Rees Feline Kidney (CRFK) cells, derived from the kidneys of a domestic cat[33–34], have been used for numerous virological studies, including feline calicivirus,[35–37] feline immunodeficiency virus[38–41], feline coronavirus[42–45], and various feline endogenous retroviruses such as RD114[46-48], as well as for vaccine development[49–51]. PG-4 cells, derived from feline fetal astrocytes, have also played an important role in retrovirus research[52–53].

Additionally, Crandell et al.[33] reported that CRFK cells exhibited aneuploidy (2n = 37) compared with the normal feline autosome count of 2n = 38, and demonstrated a characteristic karyotype with three marker chromosomes. Lasfargues et al.[54] also observed that the chromosome count fluctuated between 36 and 38 in each cell. By contrast, no such findings have been reported for PG-4 cells.

These cell lines have been passaged for decades in laboratories worldwide and have facilitated numerous scientific breakthroughs. However, despite their widespread use, no detailed information on the genetic background of the donor individuals, particularly with regard to their phenotype, such as coat and iris color, is available. Genomic information on such cell lines is

crucial for a deeper understanding of their biological properties, and elucidating the phenotypes

of their donors may provide new insights into the influence of specific genotypes on cell

properties.

In this study, we performed whole-genome sequencing of CRFK and PG-4 cell lines. The

sequencing data were used for in-depth analysis of the nucleotide sequences of genes known to

be associated with coat and iris phenotypes (i.e., *TYR*, *TYRP1*, *MC1R*, *KIT*, *ARHGAP36*,

MLPH, *ASIP*, *LVRN*, *DKK4*, *FGF5*, *KRT71*, *LPAR6*, and *PAX3*; Table 1). Based on the

genotyping results for each gene, we inferred the previously unknown phenotypes of the donor

individuals. Furthermore, analysis of publicly available RNA-seq data confirmed that these

genes were expressed in skin and eyes as well as in various other tissues throughout the body.

The findings and genome sequence data are expected to provide fundamental insights into the

genetic background of extensively used feline cell lines. This information will support the

interpretation of future research outcomes involving these cell lines and contribute to the

development of guidelines for establishing cell lines with various genotypes through genome

editing.

Methods

Cell lines and culture conditions

CRFK cells (JCRB9035, RRID: CVCL_2426) and a feline sarcoma-positive leukemia-negative

(S+L−) astrocyte cells termed PG-4 (S+L-) (JCRB9125, RRID: CVCL_3322) were obtained

from the JCRB Cell Bank (National Institutes of Biomedical Innovation, Health and Nutrition,

Osaka, Japan). CRFK cells were cultured in Dulbecco's Modified Eagle Medium (Gibco,

Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 10% fetal bovine serum

(FBS) (Gibco), penicillin (100 IU/mL), streptomycin (100 ng/mL) (Sigma-Aldrich, St. Louis,

MO, USA), and 1% non-essential amino acids (FUJIFILM Wako Pure Chemical Corporation,

Osaka, Japan) at 37 °C and 5% CO_2. PG-4 cells were cultured in McCoy's 5A Modified

Medium (Gibco) supplemented with 10% heat-inactivated FBS (Gibco), penicillin (100 IU/mL),

and streptomycin (100 ng/mL) (Sigma-Aldrich) at 37 °C and 5% CO_2. For cell passaging, cells

were washed using BASIC DPBS (no calcium or magnesium; Gibco) and then treated with

TrypLE™ Express Enzyme (1X; Gibco) for 5 min at 37 °C and 5% CO_2 conditions.

Paired-end short-read whole-genome sequencing

Genomic DNA was extracted from logarithmically growing cells (approximately 1×10^6 cells)

using the DNeasy Blood & Tissue Kit (QIAGEN, Hilden, Germany) according to the

manufacturer's protocol. The concentration and purity (A260/A280 ratio) of the extracted DNA

were measured using a NanoDrop 2000c Spectrophotometer (Thermo Fisher Scientific).

Genomic DNA extracted from CRFK cells and PG-4 cells was used for library preparation and

paired-end sequencing (150 bp read length) at Novogene (Beijing, China) using a Rapid Plus

DNA Lib Prep Kit for Illumina V2 (Illumina) and an Illumina NovaSeq X Plus platform

(Illumina). Library quality control was performed by Novogene using qPCR and fragment size

analysis according to Novogene's standard protocols. The raw sequence data were obtained

as .fastq files.

Mapping of paired-end reads and quantification

Adapter sequences were trimmed, and low-quality reads were removed from the original .fastq

files using fastp software (ver. 0.21.0) with default parameters. Reads in processed .fastq files

were mapped to the reference genome sequence of *Felis catus*, F.catus_Fca126_mat1.0 (RefSeq

assembly GCF_018350175.1) or AnAms1.0 (GenBank assembly GCA_013340865.2)[27] using

bwa mem software (ver. 0.7.19) with the setting "-R

"@RG¥tID:sample1¥tSM:sample1¥tLB:lib1¥tPL:ILLUMINA"" and default parameters

otherwise. Mapping results were obtained as .sam files. Text data in .sam files were transformed

to binary data in sorted .bam files using Samtools (ver. 1.7) with default parameters. Index files

were created for the obtained .bam files in the form of .bam.bai files.

Small variations in coat morphology-related genes

The variations of coat morphology-related phenotypes of the donor cats from which CRFK and

PG-4 cell lines originated were inferred based on the comparisons between the wild-type

sequences and mapped sequences on variant positions of each coat morphology-related gene

(Table 1). Coat phenotype-related polymorphisms in *TYR*, *TYRP1*, *MC1R*, *KIT*, *MLPH*, *ASIP*,

LVRN, *DKK4*, *FGF5*, *KRT71*, and *LPAR6* genes, which relate to phenotypes such as the color,

brown, red, white glove, dilution, agouti, tabby, ticked, long-fur, hairless, and rexing (Table 1),

were examined using alignments of paired-end sequence reads with wild-type sequences.

Large deletions in intron 1 of ARHGAP36 and the intergenic region between *KIT* and

***KDR* genes**

The deletion of a part of intron 1 of the *ARHGAP36* gene (X: 109,186,183_109,191,258 in

AnAms1.0; Table 1) was reported to produce orange coat color[12,13], and that of the intergenic

region between the *KIT* and *KDR* genes (B1: 161142880_161237957 in F.catus_Fca126_mat1.0,

Table 1) produces salmiak coat color[11]. The occurrence of these deletions in CRFK and PG-4

was estimated by comparing the mapped read number distributions between these regions and

their neighboring regions with the same region length. It should be noted that

F.catus_Fca126_mat1.0 has a deletion of the genomic region corresponding to X:

109186183_109191258 in AnAms1.0 (Table 1). Thus, AnAms1.0 (or felCat9[25]) was used as the

reference genome for analysis of the orange locus.

Insertions of FERV1-LTR and RD114-LTR into *PAX3*

The insertions of homologous sequences of the long terminal repeat (LTR) of the feline

endogenous retrovirus 1 (FERV1), termed FERV1-LTR, into a specific region in intron 4 of

PAX3 (C1: g.205833101_205833102 in F.catus_Fca126_mat1.0, Table 1) or the LTR of RD114

retrovirus, termed RD114-LTR, into another specific region in the same intron (C1:

g.205834854_205834855 in F.catus_Fca126_mat1.0, Table 1) are associated with blue iris color

in cats.[23,24] The occurrence of such insertions was estimated by extracting the genome reads that

were mapped in both FERV1-LTR and intron 4 of *PAX3* or both RD114-LTR sequences and

intron 4 of *PAX3*. The genome sequence of RD114-LTR was obtained from the NCBI

database[55].

Long-read whole-genome sequencing

Genomic DNA of CRFK cells and PG-4 cells was extracted for long-read genome sequencing

using the NucleoBond HMW DNA kit (TaKaRa, Tokyo, Japan), and libraries for the Oxford

Nanopore Technologies (ONT, UK) sequencer were constructed using a ligation library

preparation kit (SQK-LSK114, ONT). All procedures were performed according to the

manufacturer's instructions, and the quality and molecular weight of the genomic DNA were

measured using the Qubit fluorometric quantification system (Thermo Fisher Scientific).

Sequencing was conducted using a PromethION sequencer (PromethION 2 Solo, ONT) and

R10.4.1 flow cells. POD5 files were basecalled to .fastq files using MinKNOW software

v25.05.14.

Insertions of FERV1 homolog into *KIT* examined by long-read sequencing

The insertion of FERV1 into a specific region of intron 1 of the *KIT* gene (B1:

161388687_161388688 in F.catus_Fca126_mat1.0, Table 1) and that of FERV1-LTR make the

coat color of cats white-spotted and white throughout, respectively [8]. The occurrence of such

insertions in CRFK and PG-4 cells was estimated by mapping the reads of long-read sequencing

to the genome sequence named KIT-FERV1 sequence using minimap2 (ver. 2.30), where the

KIT-FERV1 sequence is a part of the *KIT* intron 1 sequence with the insertion of the FERV1

sequence, as shown in Supplementary Figure 1 of the study by David et al[8].

Transcriptome data analysis of coat phenotype- and iris color-related genes in various

tissues of domestic cats

Mapped RNA-seq data on the F.catus_Fca126_mat1.0 were obtained in .bam and .bam.bai

formats for various tissues from the Ensembl public database[56], specifically, the file transfer

protocol (ftp) site[57]. The annotation file of the F.catus_Fca126_mat1.0 in gene transfer format

(GTF) was obtained from the NCBI database (Bethesda, MD, USA). In the file, the

chromosome numbers were converted from NC_058368.1, NC_058369.1, etc. to A1, A2, etc.

Using FeatureCounts with the options --M-O--fraction,[58] read counts data for each gene

described in the GTF file were obtained from the .bam and .bam.bai formatted files. This count

data could also be obtained from a file

"F.catus_Fca126_mat1.0.ENA_gene_exp_mat_by_FeatureCounts.csv" deposited in F.catlas[59].

From these counts, the transcript level of each gene was calculated by dividing the read count

by the total exon length. The total transcript level was defined as the sum of all the transcript

levels of the genes. Based on this result, the transcripts per million (TPM) of each gene were

calculated as 1,000,000*(transcript level)/(total transcript level).

Results

Short- and long-read whole-genome sequencing

For CRFK and PG-4 cells, the total output of raw sequence data of 150 bp paired-end

sequencing was 60.6 Gb and 60.1 Gb, and the [average] $\pm$ [standard deviation] of mapping

depth of the reads for F.catus_Fca126_mat1.0 were 23.59 ± 9.79 and 23.45 ± 8.17,

respectively; the total output of raw sequence data of long-read sequencing was 9.6 Gb and 11.8

Gb, N50 values were 23.5 kb and 38.2 kb, and the [average] $\pm$ [standard deviation] of mapping

depth of reads for F.catus_Fca126_mat1.0 were 3.91 ± 3.11 and 4.79 ± 3.34, respectively.

The average and standard deviations specified above were estimated through the entire genome

region without the regions with singular sequences where the read coverage exhibited >100.

Predicted phenotype of the CRFK cell donor

The mapping of paired-end reads from the whole-genome sequences of the CRFK cell revealed that most coat morphology-related genes were conserved (Table 1). However, the following variations were identified in mapped genome reads from CRFK cells (Table 1): a homozygous mutation of c.123_124del of the *ASIP* gene (*nonagouti* allele), a homozygous mutation of c.475A>C of the *FGF5* gene (*long coat* allele; Fig. 1), a homozygous mutation of c.2522G>A of the *LVRN* gene (*blotched-tabby* allele), and a heterozygous mutation of c.83delT of the *MLPH* gene (*dilute* allele). No 95 kbp deletion was detected in the intergenic region between the *KIT* and *KDR* genes, nor a 5 kbp deletion was observed in intron 1 of the *ARHGAP36* gene (Fig. 2). Additionally, no genome reads were identified that could be simultaneously mapped to both the *PAX3* intron 4 and FERV1-LTR or to both the *PAX3* intron 4 gene and RD114-LTR sequences. Furthermore, four reads from the long-read sequencing mapped to the KIT-FERV1 sequence; however, these reads contained no insertions of FERV1 and FERV1-LTR (Fig. 3).

The abovementioned homozygous mutation of the *FGF5* gene was known to exhibit long fur[18]. However, the mutation of the *LVRN* did not influence coat morphology in this case because of the mutation of the *ASIP* gene that suppresses the stripe pattern on the cat's skin coat, and the mutant genotype of the *MLPH* gene was recessive.[1] Therefore, the cat from which the CRFK cells originated is expected to have had long, black, unstriped fur and non-blue eyes (Fig. 4).

Predicted phenotype of the PG-4 cell donor

Mapped paired-end reads from the whole-genome sequences of the PG-4 cell indicated that most coat morphology-related genes were conserved but revealed the following variations (Table 1): a homozygous mutation of c.123_124del of the *ASIP* gene, a compound heterozygous mutation of c.474delT and c.475A>C of the *FGF5* gene (Fig. 1) where one allele exhibits c.474delT and the other allele exhibits c.475A>C as mentioned below, the heterozygous mutations of c.416C>A and c.2522G>A of *LVRN* gene, and a heterozygous mutation of c.904G>A of the *TYR* gene (*Siamese-color* allele). No 95 kbp deletion was identified in the intergenic region between the *KIT* and *KDR* genes, nor a 5 kbp deletion was found in intron 1 of the *ARHGAP36* gene (Fig. 2), as was similarly observed with CRFK. Additionally, we did not identify any genome reads that mapped concurrently to both the *PAX3* intron 4 and FERV1-LTR or to both the *PAX3* intron 4 gene and RD114-LTR sequences.

Notably, the genome sequence reads obtained from PG-4 cells containing bases 474 and 475 of

the CDS of the FGF5 gene revealed that seven reads exhibited only a deletion at base 474, while

the remaining 13 reads showed only a substitution at base 475 (Fig. 1). Accordingly, we

expected a compound heterozygous mutation in the FGF5 gene of PG-4 cells. Similarly, PG-4

cells exhibited heterozygous substitutions at bases 416 and 2522 of the CDS of the LVRN gene.

However, owing to the considerable distance between the genomic locations of these two bases,

none of our sequencing data, including the long-read sequencing data, produced reads that

contained both bases together. Therefore, we were unable to conclude whether the LVRN gene

in PG-4 cells harbors a compound heterozygous mutation.

Six long-read sequences mapped to the KIT-FERV1 sequence, with four of them containing the

insertions of the FERV1 homolog sequence exceeding 7 kbp (Fig. 3). This indicates that the PG-

4 genome contains the heterozygous insertion of the entire FERV1 homolog sequence.

The mutant genotype of the *TYR* gene was recessive, and in this case, the two heterozygous

mutations of the *LVRN* did not influence coat morphology because of the *ASIP* gene mutation,

the same as in CRFK[1]. However, the effect of *FERV1* is known to be dominant[8]. Additionally,

the compound heterozygous mutation of the *FGF5* gene produces long fur[18]. Therefore, the PG-

4 cells likely originated from a cat with white and black (white spotting), unstriped, long fur

(Fig. 4).

Notably, certain cats with white spotting may have blue eyes or atypical eye coloration;

however, this occurrence is less common than that in cats exhibiting the Dominant White allele

in the KIT gene[8]. Additionally, no insertions of FERV1-LTR and RD114-LTR in intron 4 of the

PAX3 gene were observed. Therefore, the PG-4 cells likely originated from a cat with non-blue

irises (Fig. 4).

Expression features of coat phenotype- and iris color-related genes in various tissues

The analysis of publicly available RNA-seq data from various organs of domestic cats in the

Ensembl database

(https://ftp.ensembl.org/pub/data_files/felis_catus/F.catus_Fca126_mat1.0/rnaseq/) revealed that

coat phenotype-related genes are expressed in various organs at levels (TPM) comparable to or

greater than those in the skin (Fig. 5, Table S2). For example, *TYR*, *TYRP1*, and *MLPH* were

highly expressed in the ear tip, retina, and optic nerve, and *MC1R*, *KIT*, *DKK4*, and *FGF5* were

highly expressed in the brain. *KIT*, *DKK4*, and *LPAR6* tend to be expressed throughout the body,

with prominent expression in the reproductive organs. Additionally, *ARHGAP36* was highly

expressed in the spinal cord, and *ASIP*, along with *MLPH*, was highly expressed in the lungs.

The iris color-related gene *PAX3* was also highly expressed in the cerebellum, in addition to

visual organs, and, along with *DKK4*, in the kidney and embryonal tissue.

Discussion

We performed whole-genome sequencing of CRFK, a cultured feline kidney cell line, and PG-4,

a feline fetal astrocyte-derived cell line, and analyzed the genome data to determine the coat

color, hair length, coat pattern, and iris color of the donor individuals. The results suggested that

the CRFK cell line originated from a cat with long black hair without stripes and non-blue

irises, and the PG-4 cell line came from a cat with long fur, black and white bicolor coat without

stripes, and non-blue irises.

Coat morphology and iris color are phenotypes that can be readily identified and used to predict

genotypes. However, the expressions of these genes are not limited to the skin and eyes; rather,

they are expressed in various tissues throughout the body (Fig. 5). Therefore, differences in

these genotypes may contribute to disease susceptibility in specific organs. The effects of these

genetic variations are likely weaker than those of the rearing environment and may not be

significant, especially in young, healthy individuals. However, as domestic cats have become

longer-lived in recent years, individual-specific risks may become more apparent, and the

underlying genotypes may exert increasing effects as the cats age. Therefore, elucidating the

relationships between genotypes underlying readily recognizable phenotypes and disease risk is

crucial from the perspectives of both medical care and welfare, such as prevention and symptom

suppression/alleviation.

Results from research on cultured cells to elucidate disease mechanisms and develop therapeutic

drugs must be interpreted with caution, i.e., not only on the experimental results obtained using

established, immortalized cells but also with regard to their genotype and expression profiles of

the genes harboring polymorphisms in various tissues and organs. However, so far, such

information has not been available, even for certain frequently used cell lines such as CRFK.

The present study and genome sequence data provide such information and facilitate a more

comprehensive interpretation of research on feline cultured cell lines.

Furthermore, with the availability of detailed genomic information, genome editing can be

applied to these cell lines to establish various derivative lines that differ only in specific aspects,

e.g., regarding coat morphology or other traits, and compare their properties. This is expected to

advance research on the regulatory mechanisms of coat color, for example, the validation of the

role of the novel *KIT* isoform caused by FERV1-LTR insertion in the formation of white coat[60],

as well as the relationships between the visually identifiable phenotypes examined in the present

study and disease risk and treatment methods. Simultaneously, this comprehensive genomic

information can help to develop guidelines for establishing cell lines with various disease-

associated genotypes using genome editing. The development of these derivative cell lines may

provide valuable insights into the mechanism underlying deafness in cats with white fur and

blue irises, as well as elucidate the influence of other coat-related factors in organs beyond the

skin or functions unrelated to coat formation.

The limitations and future directions of this study are outlined below.

First, notably, the findings of this study are specific to the cell types used. The cells used in this

study were CRFK cells (JCRB9035, RRID: CVCL_2426), which were prepared from CRFK

(ATCC: CCL 94) and PG-4 (S+L-) cells (JCRB9125, RRID: CVCL_3322) obtained from PG-4

(S+L-) cells (ATCC: CRL-2032). Consequently, these cells were expected to have similar

characteristics to those of the CRFK and PG-4 cells used in numerous previous studies. For

example, however, the number of chromosomes has been reported to vary from 36 to 38 in

CRFK[33,54]. Additionally, the karyotype of PG-4 cells, which has not yet been reported, likely

differs from the normal domestic cat genome. Therefore, using the same cell line may still

produce results that differ slightly from those used by other researchers. Investigating additional

variation among cells in these lines remains an important future consideration.

As mentioned previously, the karyotype of CRFK differs from that of the normal domestic cat

genome[33,54]. Additionally, the specific chromosomes that cause aneuploidy remain unidentified.

However, identifying large structural variations using only whole-genome analysis of the

present sequence read data, approximately 60 Gb paired-end sequencing data and 10 Gb long-

read data, was challenging. Similarly, accurately identifying sequences and locations of various

long repeat sequences, endogenous retroviral sequences, and their homologs was challenging

owing to limited read coverage, particularly for long-read coverage. However, these sequences

may also be associated with various diseases, as demonstrated by the involvement of FERV1-

LTR in deafness in cats[8,23,24] and the positive and negative roles of enFeLV-LTR and enFeLV-

env in resistance to exogenous feline leukemia virus[61-63]. The increased volume of ongoing

long-read sequencing, along with the draft genome assembly using these data, is expected to

330 address these challenges in the future.

331 This study focused on coat and iris phenotypes in domestic cats. However, major

332 histocompatibility complex (MHC) polymorphisms are strongly correlated with breed in

333 domestic dogs[64-68]. Although research on MHC polymorphisms in domestic cats is ongoing[69-71],

334 the relationship with breed remains unclear. In this study, we showed the results of the primary

335 analysis of the small indels and SNPs in MHC class I and II regions of CRFK and PG-4 cells

336 described by Variant Call format (VCF)[72] (Table S3). However, because the MHC gene cluster

337 comprises a complex and large genomic region, it is difficult to identify the donor breeds of

338 CRFK and PG-4 using only the sequence data and analytical methods presented in this study.

339 However, as mentioned above, further long-read sequencing and its data analysis may enable

340 breed estimation of these cell lines.

341 In addition, the present genome sequence data also included mitochondrial DNA data. In this

342 study, we presented the results of the primary analysis of the small indels and SNPs in the

343 mitochondrial DNA of CRFK and PG-4 cells described by VCF[72] (Table S4[72]). Although the

344 correlations are not substantially strong, an association may exist between the polymorphisms of

345 mitochondrial DNA and breeds of domestic cats[73-76]. Therefore, a comprehensive analysis of

346 this data may also contribute to breed estimation.

347 These analyses are also critical future challenges.

349 **Conclusions**

350 We performed whole-genome sequencing of CRFK and PG-4 cells and inferred the phenotype

351 of the donors of these cells. We suggested that CRFK cells originated from a cat with long,

352 black fur lacking stripes and non-blue irises; PG-4 cells originated from a cat with long,

353 bicolored white and black fur without stripes, and with non-blue irises. Additionally, analysis of

354 publicly available RNA-seq data confirmed that genes associated with coat phenotype and iris

355 color are expressed in the skin and eyes, as well as in various other organs, indicating that

356 variants of these genes, which affect coat phenotype and iris color, may influence physiological

357 functions throughout the body. These insights may facilitate a more accurate interpretation of

358 data derived from feline cultured cells and inform guidelines for developing cell lines with

359 domestic cat genotypes that exhibit various phenotypes using genome editing techniques. This

360 will help to elucidate the role of various coat phenotype-related and iris color-related factors in

organs other than the skin and functions independent of coat formation, including the effects of
melanocyte destruction caused by KIT gene mutations on the nervous system.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

These raw read data of the short and long read sequencings were deposited in the DDBJ
Sequence Read Archive (DRA) with BioProject accession PRJDB37530 (DRR748391-
DRR748394, DRR786948, and DRR786949). All other data generated or analyzed during this
study were included in this published article (and Supplementary Information files).

Conflicting interests

The authors have no conflicting interests to declare.

Funding

This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI
grants (award numbers 24K01783 (T. I.) and 24K09248 (A. A.)).

Author contributions

Conceptualization: A.A., Data curation: T.K., A.K., R.H., and T.I., Formal analysis: G.T. and
R.G., Funding acquisition: A.A., Investigation: G.T., R.G., T.K., R.H., and A.A., Methodology:

G.T., R.G., T.K., T.I., and A.A., Project administration: N.S. and A.A., Supervision: N.S. and

A.A., Validation: R.G. and A.A., Visualization: T.G., N.S., and A.A., Writing - original draft:

A.A., Writing - review & editing: T.G., T.K., T.I., and N.S.

Acknowledgements

Not applicable.

References

1.   Lyons LA. DNA mutations of the cat: The good, the bad and the ugly: The good, the bad

and the ugly. *J Feline Med Surg*. 2015; 17:203-219. doi:10.1177/1098612X15571878

2.   Schmidt-Kuntzel A, Eizirik E, O'Brien SJ, et al. Tyrosinase and tyrosinase related protein 1

alleles specify domestic cat coat color phenotypes of the Albino and Brown loci. *J Hered*

2005; 96: 289–301

3.   Lyons LA, Imes DL, Rah HC, et al. Tyrosinase mutations associated with Siamese and

Burmese patterns in the domestic cat (*Felis catus*). *Anim Genet* 2005; 36: 119–126.

4.   Imes DL, Geary LA, Grahn RA, et al. Albinism in the domestic cat (*Felis catus*) is

associated with a tyrosinase (TYR) mutation. *Anim Genet* 2006; 37: 175–178

5.   Lyons LA, Foe IT, Rah HC, et al. Chocolate coated cats: TYRP1 mutations for brown color

in domestic cats. *Mamm Genome* 2005; 16: 356–366

6.   Peterschmitt M, Grain F, Arnaud B, et al. Mutation in the melanocortin 1 receptor is

associated with amber colour in the Norwegian forest cat. *Anim Genet* 2009; 40: 547–552

7.   Montague MJ, Li G, Gandolfi B, et al. Comparative analysis of the domestic cat genome

reveals genetic signatures underlying feline biology and domestication. *Proc Natl Acad Sci*

USA 2014; 111: 17230–17235

8.   David VA, Menotti-Raymond M, Wallace AC, et al. Endogenous retrovirus insertion in the

KIT oncogene determines white and white spotting in domestic cats. G3 (Bethesda) 2014;

4: 1881–1891

9.   Frischknecht M, Jagannathan V, Leeb T. Whole genome sequencing confirms KIT insertions

in a white cat. *Anim. Genet* 2015; 46: 98

10. Górska A, Drobik-Czwarno W, Bryś, J. Genetic determination of the amount of white

spotting: a case study in Siberian cats. *Genes* (Basel) 2022; 13(6)

11. Anderson H, Salonen M, Toivola S, Blades M, Lyons L A, Forman OP, Hytönen MK, Lohi
H. A new Finnish flavor of feline coat coloration, "salmiak," is associated with a 95-kb
deletion downstream of the KIT gene. *Anim. Genet* 2024; 55: 676-680

12. Kaelin CB, McGowan KA, Trotman JC, Koroma DC, David VA, Menotti-Raymond M,
Graff EC, Schmidt-Küntzel A, Oancea E, Barsh GS. Molecular and genetic characterization
of sex-linked orange coat color in the domestic cat. *Curr Biol* 2025; 35:2826-2836.e5.
Pubmed reference: 40378841. DOI: 10.1016/j.cub.2025.04.055

13. Toh H, Au Yeung WK, Unoki M, Matsumoto Y, Miki Y, Matsumura Y, Baba Y, Sado T,
Nakamura Y, Matsuda M, Sasaki H. A deletion at the X-linked ARHGAP36 gene locus is
associated with the orange coloration of tortoiseshell and calico cats. *Curr Biol* 2025;
35:2816-2825.e3, 2025. Pubmed reference: 40378840. DOI: 10.1016/j.cub.2025.03.075

14. Ishida Y, David VA, Eizirik E, et al. A homozygous single-base deletion in MLPH causes
the dilute coat color phenotype in the domestic cat. *Genomics* 2006; 88: 698–705

15. Eizirik E, Yuhki N, Johnson WE, et al. Molecular genetics and evolution of melanism in the
cat family. *Curr Biol* 2003; 13: 448–453

16. Kaelin CB, Xu X, Hong LZ, et al. Specifying and sustaining pigmentation patterns in
domestic and wild cats. *Science* 2012; 337: 1536–1541

17. Lyons LA, Buckley RM, Harvey RJ, 99 Lives Cat Genome Consortium. Mining the 99
Lives Cat Genome Sequencing Consortium database implicates genes and variants for the
Ticked locus in domestic cats (*Felis catus*). *Anim Genet* 2021; 52: 321–332
https://doi.org/10.1111/age.13059

18. Kehler JS, David VA, Schaffer AA, et al. Four independent mutations in the feline fibroblast
growth factor 5 gene determine the long-haired phenotype in domestic cats. *J Hered* 2007;
98: 555–566

19. Drogemuller C, Rufenacht S, Wichert B, et al. Mutations within the FGF5 gene are
associated with hair length in cats. *Anim Genet* 2007; 38: 218–221

20. Gandolfi B, Outerbridge C, Beresford L, et al. The naked truth: Sphynx and Devon Rex cat
breed mutations in KRT71. *Mamm Genome* 2010; 21; 509–515

21. Gandolfi B, Alhaddad H, Joslin SE, et al. A splice variant in KRT71 is associated with curly
coat phenotype of Selkirk Rex cats. *Sci Rep* 2013; 3: 2000

22. Gandolfi B, Alhaddad H, Affolter VK, et al. To the root of the curl: a signature of a recent
selective sweep identifies a mutation that defines the Cornish Rex cat breed. *PloS One*
2013; 8: e67105

23. Abitbol M, Couronné A, Dufaure de Citres C, Gache V. A PAX3 insertion in the Celestial

breed and certain feline breeding lines with dominant blue eyes. *Anim Genet* 2024; 55: 670-
675

24. Abitbol M, Dufaure de Citres C, Rudd Garces G, Lühken G, Lyons LA, Gache V. Different
founding effects underlie dominant blue eyes (DBE) in the domestic cat. *Animals (Basel)*
2024; 14:1845. Pubmed reference: 38997957. DOI: 10.3390/ani14131845

25. Buckley RM, Davis BW, Brashear WA, Farias FHG, Kuroki K, Graves T, et al. A new
domestic cat genome assembly based on long sequence reads empowers feline genomic
medicine and identifies a novel gene for dwarfism. *PLoS Genet* 2020; 16(10): e1008926.
https://doi.org/10.1371/journal.pgen.1008926

26. Bredemeyer KR, Harris AJ, Li G, Zhao L, Foley NM, Roelke-Parker M, O'Brien SJ, Lyons
LA, Warren WC, Murphy WJ. Ultracontinuous single haplotype genome assemblies for the
domestic cat (*Felis catus*) and Asian leopard cat (*Prionailurus bengalensis*). *J Hered* 2021
112(2), 165–173. https://doi.org/10.1093/jhered/esaa057

27. Matsumoto Y, Chung CYL, Isobe S, Sakamoto M, Lin X, et al. Chromosome-scale
assembly with improved annotation provides insights into breed-wide genomic structure
and diversity in domestic cats. *J Adv Res* 2025; 75, 863-874

28. Lyons LA, Creighton EK, Alhaddad H. et al. Whole genome sequencing in cats, identifies
new models for blindness in AIPL1 and somite segmentation in HES7. *BMC Genomics*
2016; 17: 265. https://doi.org/10.1186/s12864-016-2595-4

29. Rodney AR, Buckley RM, Fulton RS et al. A domestic cat whole exome sequencing
resource for trait discovery. *Sci Rep* 2021; 11: 7159. https://doi.org/10.1038/s41598-021-
86200-7

30. Anderson H, Davison S, Lytle KM, Honkanen L, Freyer J, Mathlin J, et al. Genetic
epidemiology of blood type, disease and trait variants, and genome-wide genetic diversity in
over 11,000 domestic cats. *PLoS Genet* 2022; 18(6): e1009804.
https://doi.org/10.1371/journal.pgen.1009804

31. Hernandez I, Hayward JJ, Brockman JA, White ME et al. Complex feline disease mapping
using a dense genotyping array. *Front Vet Sci* 2022; 9, 862414

32. Raffle J, Novo Matos J, Wallace M, et al. Identification of novel genetic variants associated
with feline cardiomyopathy using targeted next-generation sequencing. *Sci Rep* 2025; 15:
3871

33. Crandell RA, Fabricant CG, Nelson-Rees WA. Development, characterization, and viral
susceptibility of a feline (*Felis catus*) renal cell line (CRFK). *In Vitro* 1973; 9: 176-185

34. Lawson JS, Syme HM, Wheeler-Jones CPD, Elliott J. Characterisation of Crandell-Rees

486  Feline Kidney (CRFK) cells as mesenchymal in phenotype. *Res Vet Sci* 2019; 127: 99-102

487 35. Sosnovtsev SV, Prikhod'ko EA, Belliot G, Cohen JI, Green KY. Feline calicivirus

488  replication induces apoptosis in cultured cells. *Virus Res* 2003; 94: 1-10

489 36. Bidawid S, Malik N, Adegbunrin O, Sattar SA, Farber JM. A feline kidney cell line-based

490  plaque assay for feline calicivirus, a surrogate for Norwalk virus. *J Virol Methods* 2003;

491  107: 163-167

492 37. Tian J, Zhang X, Wu H, Liu C, Liu J, Hu X, Qu L. Assessment of the IFN-β response to

493  four feline caliciviruses: Infection in CRFK cells. *Infect Genet Evol* 2015; 34: 352-360.

494 38. Siebelink KH, Karlas JA, Rimmelzwaan GF, Osterhaus AD, Bosch ML. A determinant of

495  feline immunodeficiency virus involved in Crandell feline kidney cell tropism. *Vet Immunol*

496  *Immunopathol* 1995; 46: 61-69

497 39. Verschoor EJ, Boven LA, Blaak H, Van Vliet AL, Horzinek MC, De Ronde A. A single

498  mutation within the V3 envelope neutralization domain of feline immunodeficiency virus

499  determines its tropism for CRFK cells. *J Virol* 1995; 69: 4752-4757

500 40. Baldinotti F, Matteucci D, Mazzetti P, Giannelli C, Bandecchi P, Tozzini F, Bendinelli M.

501  Serum neutralization of feline immunodeficiency virus is markedly dependent on passage

502  history of the virus and host system. *J Virol* 1994; 68: 4572-4579

503 41. Mizuno T, Goto Y, Baba K, Masuda K, Ohno K, Tsujimoto H. TNF-α-induced cell death in

504  feline immunodeficiency virus-infected cells is mediated by the caspase cascade. *Virology*

505  2001; 287: 446-455

506 42. Drechsler Y, Vasconcelos EJ, Griggs LM, Diniz PP. CRFK and primary macrophages

507  transcriptomes in response to feline coronavirus infection differ significantly. *Front Genet*

508  2020; 11: 584744

509 43. Drechsler Y, Vasconcelos EJ, Diniz PP. Host transcriptome studies in response to feline

510  coronavirus reveal differences in macrophages vs CRFK. *J Immunol*, 2020;

511  204(1_Supplement): 92-27

512 44. Drechsler Y, Vasconcelos EJ, Griggs LM, Diniz PP. Host responses to feline coronavirus are

513  significantly different in primary macrophages compared to CRFK cells. *J Immunol* 2021;

514  206(1_Supplement): 19-13

515 45. Camero M, Lanave G, Catella C, Lucente MS, et al. ERDRP-0519 inhibits feline

516  coronavirus in vitro. *BMC Vet Res* 2022; 18: 55

517 46. Baumann JG, Günzburg WH, Salmons B. CrFK feline kidney cells produce an RD114-like

518  endogenous virus that can package murine leukemia virus-based vectors. *J Virol* 1998; 72:

519  https://doi.org/10.1128/jvi.72.9.7685-7687.1998

47. Shimode S, Nakagawa S, Miyazawa T. Multiple invasions of an infectious retrovirus in cat
genomes. *Sci Rep* 2015; 5, 8164

48. Shimode S, Sakuma T, Yamamoto T, Miyazawa T. Establishment of CRFK cells for vaccine
production by inactivating endogenous retrovirus with TALEN technology. *Sci Rep* 2022;
12: 6641

49. Lappin MR, Jensen WA, Jensen TD, Basaraba RJ, Brown CA, Radecki SV, Hawley JR.
Investigation of the induction of antibodies against Crandell-Rees feline kidney cell lysates
and feline renal cell lysates after parenteral administration of vaccines against feline viral
rhinotracheitis, calicivirus, and panleukopenia in cats. *Am J Vet Res* 2005; 66: 506-511.

50. Whittemore JC, Hawley JR, Jensen WA, Lappin MR. Antibodies against Crandell Rees
Feline Kidney (CRFK) cell line antigens, α-enolase, and annexin A2 in vaccinated and
CRFK hyperinoculated cats. *J Vet Intern Med* 2010; 24: 306-313

51. Yoshikawa R, Sato E, Igarashi T, Miyazawa T. Characterization of RD-114 virus isolated
from a commercial canine vaccine manufactured using CRFK cells. *J Clin Microbiol* 2010;
48: 3366-3369

52. Haapala DK, Robey WG, Oroszlan SD, Tsai WP, Isolation from cats of an endogenous type
C virus with a novel envelope glycoprotein. *J Virol* 1985; 53: 827-33

53. Bassin RH, Ruscetti S, Ali I, Haapala DK, Rein A, Normal DBA/2 mouse cells synthesize a
glycoprotein which interferes with MCF virus infection. *Virol* 1982; 123: 139-51

54. Lasfargues EY, Lasfargues JC, Dion AS, Greene AE, Moore DH. Experimental infection of
a cat kidney cell line with the mouse mammary tumor virus. Cancer Res. 1976; 36: 67-72

55. RD114 retrovirus, complete genome:
https://www.ncbi.nlm.nih.gov/nuccore/NC_009889.1/. Accessed 4 August 2025

56. Cunningham F, Allen JE, Allen J, Amode MR, Armean IM, Azov AG, Barnes I, Bennett R,
Berry A, Bhai J, Bignell A, Billis K, Boddu S, et al. Ensembl 2022. *Nucl Acids Res* 2022;
50(D1), D988–D995

57. Ensembl Complete datasets and databases:
https://ftp.ensembl.org/pub/data_files/felis_catus/F.catus_Fca126_mat1.0/rnaseq/. Accessed
26 January 2023

58. Liao Y, Smyth GK, Shi W. FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* 2014; 30: 923–930

59. F. catlas: https://sites.google.com/view/fcatlas/. Accessed 1 November 2025

60. Awazu A, Takemoto D, Watanabe K, Sakamoto N. Possibilities of skin coat color-dependent
risks and risk factors of squamous cell carcinoma and deafness of domestic cats inferred via

RNA-seq data. *Genes Cells* 2023; 28: 893-905

61. Chiu ES, VandeWoude S. Presence of endogenous viral elements negatively correlates with

feline leukemia virus susceptibility in puma and domestic cat Cells. *J Virol* 2020; 94:

10.1128/jvi.01274-20

62. Erbeck K, Gagne RB, Kraberger S,Chiu ES, Roelke-Parker M, VandeWoude S. Feline

leukemia virus (FeLV) endogenous and exogenous recombination events result in multiple

FeLV-B subtypes during natural infection. *J Virol* 2021; 95: 10.1128/jvi.00353-21

63. Pramono D, Takeuchi D, Katsuki M, AbuEed L, Abdillah D, Kimura T, Kawasaki J, Miyake

A, Nishigaki K. FeLIX is a restriction factor for mammalian retrovirus infection. *J Virol*

2024; 98:e01771-23

64. Kennedy LJ, Barnes A, Happ GM, Quinnell RJ, Bennett D, Angles JM, Day MJ,

Carmichael N, Innes JF, Isherwood D, Carter SD, Thomson W, Ollier WE. Extensive

interbreed, but minimal intrabreed, variation of DLA class II alleles and haplotypes in dogs.

Tissue Antigens. 2002; 59: 194-204

65. Angles JM, Kennedy LJ, Pedersen NC. Frequency and distribution of alleles of canine

MHC-II DLA-DQB1, DLA-DQA1 and DLA-DRB1 in 25 representative American Kennel

Club breeds. Tissue Antigens. 2005; 66:173-84

66. Miyamae J, Okano M, Katakura F, Kulski JK, Moritomo T, Shiina T. Large-Scale

Polymorphism Analysis of Dog Leukocyte Antigen Class I and Class II Genes (*DLA-*

88, *DLA-12/88L* and *DLA-DRB1*) and Comparison of the Haplotype Diversity between

Breeds in Japan. Cells. 2023;12: 809

67. Graumann MB, DeRose SA, Ostrander EA, Storb R. Polymorphism analysis of four canine

MHC class I genes. Tissue Antigens. 1998; 51(4 Pt 1): 374-81

68. Wagner JL, Burnett RC, Storb R. Organization of the canine major histocompatibility

complex: current perspectives. J Hered. 1999; 90 :35-8

69. Morris KM, Kirby K, Beatty JA, Barrs VR, Cattley S, David V, O'Brien SJ, Menotti-

Raymond M, Belov K. Development of MHC-Linked Microsatellite Markers in the

Domestic Cat and Their Use to Evaluate MHC Diversity in Domestic Cats, Cheetahs, and

Gir Lions. J Hered. 2014; 105: 493-505

70. Okano M, Miyamae J, Suzuki S, Nishiya K, Katakura F, Kulski JK, Moritomo T, Shiina T.

Identification of Novel Alleles and Structural Haplotypes of Major Histocompatibility

Complex Class I and DRB Genes in Domestic Cat (*Felis catus*) by a Newly Developed

NGS-Based Genotyping Method. Front Genet. 2020; 11: 750

71. Holmes JC, Holmer SG, Ross P, Buntzman AS, Frelinger JA, Hess PR. Polymorphisms and

tissue expression of the feline leukocyte antigen class I loci FLAI-E, FLAI-H, and FLAI-K.

Immunogenetics. 2013; 65: 675-89

72. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,

Lunter G, Marth GT, Sherry ST, McVean G, Durbin R; 1000 Genomes Project Analysis

Group. The variant call format and VCFtools. Bioinformatics. 2011; 27: 2156-2158

73. Grahn RA, Kurushima JD, Billings NC, Grahn JC, Halverson JL, Hammer E, Ho CK, Kun

TJ, Levy JK, Lipinski MJ, Mwenda JM, Ozpinar H, Schuster RK, Shoorijeh SJ, Tarditi CR,

Waly NE, Wictum EJ, Lyons LA. Feline non-repetitive mitochondrial DNA control region

database for forensic evidence. Forensic Sci Int Genet. 2011;5: 33-42

74. Wesselink M, Bergwerff L, Hoogmoed D, Kloosterman AD, Kuiper I. Forensic utility of the

feline mitochondrial control region - A Dutch perspective. Forensic Sci Int Genet. 2015; 17:

25-32

75. Bilgen N, Güvendi M, Sezgin E, Can H, Köseoğlu AE, Erkunt Alak S, Döşkaya M, Ün C.

Mitotyping of random bred cats and pure breed cats (Turkish Angora and Turkish Van)

using non-repetitive mitochondrial DNA control region. Gene. 2024; 892: 147849

76. Jaito W, Singchat W, Patta C, Thatukan C, Kumnan N, Chalermwong P, Budi T, Panthum T,

Wongloet W, Wattanadilokchatkun P, Thong T, Muangmai N, Han K, Duengkae P,

Phatcharakullawarawat R, Srikulnath K. Shared alleles and genetic structures in different

Thai domestic cat breeds: the possible influence of common racial origins. Genomics

Inform. 2024; 22: 12

Table 1: Phenotypic traits determined by DNA variants, and genotypes of CRFK and PG-4
cells.

Locus Gene	Phenotype (Alleles)	Mutation in CDS (or related noncoding region): F.catus_Fca126_mat1.0 Genome: F.catus_Fca126_mat1.0 Genome: AnAms1.0	Depth of alleles, and Genotype in CRFK	Depth of alleles, and Genotype in PG-4
Color TYR[2–4]	Burmese (c^b)	c.679G>T	+ : 18	+ : 16
		D1: g.44022819C>A	c^b : 0	c^b : 0
		D1: g.46081417C>A	+ / +	+ / +
	Siamese (c^s)	c.904G>A	+ : 25	+ : 17
		D1: g.44013031C>T	c^s : 0	c^s : 20
		D1: g.46071661C>T	+ / +	+ / c^s
	Albino (c^a)	c.939del	+ : 25	+ : 34
		D1: g.44012996del	c^a : 0	c^a : 0
		D1: g.46071626del	+ / +	+ / +
	Mocha (c^m)	c.820_936delinsAATCTC	+ : 40	+ : 54
		D1: g.44012836_44012998delinsGAGATT	c^m : 0	c^m : 0
		D1: g.46071466_46071628delinsGAGATT	+ / +	+ / +
Brown TYRP1[2,5]	Chocolate (b)	c.1261+5G>A	+ : 22	+ : 26
		D4: g.38142265G>A	b : 0	b : 0
		D4: g.39391308G>A	+ / +	+ / +
	Cinnamon (b^l)	c.298C>T	+ : 20	+ : 19
		D4: g.38130163C>T	b^l : 0	b^l : 0
		D4: g.39379209C>T	+ / +	+ / +
Amber MC1R[6]	Red (e)	c.250G>A	+ : 35	+ : 34
		E2: g.61570294G>A	e : 0	e : 0
		E2: g.62623146G>A	+ / +	+ / +

White KIT[7–11]	White (w^w)	(Intron 1: FERV1-LTR insertion) B1: g.161388687_161388688insFERV1-LTR B1: g.163130991_163130992insFERV1-LTR	Fig. 3 + / +	Fig. 3 + / +
	White spotted (w^s)	(Intron 1: FERV1 insertion) B1: g.161388687_161388688insFERV1 B1: g.163130991_163130992insFERV1	Fig. 3 + / +	Fig. 3 + / w^s
	White glove (w^g)	c.1035_1036delinsCA B1: g.161337385_161337386delinsTG B1: g.163079561_163079562delinsTG	+ : 27 w^g : 0 + / +	+ : 28 w^g : 0 + / +
	Salmiak (w^a)	(KIT-KDR Intergenic region: 95 kb deletion) B1: g.161142880_161237957del B1: g.162885414_162980052del	Fig. 2a + / +	Fig. 2a + / +
Orange ARHGAP36 [12,13]	Orange (o)	(Intron 1: 5 kb deletion) (Lacking) X: g.109186183_109191258del	Fig. 2b + / +	Fig. 2b + / +
Dilution MLPH[14]	Dilute (d)	c.83del C1: g.218197448del C1: g.219535339del	+ : 15 d : 4 + / d	+ : 38 d : 0 + / +
Agouti ASIP[15]	Nonagouti (a)	c.123_124del A3: g.24831083-24831084del A3: g.25283750_25283751del	+: 0 a: 48 a / a	+: 0 a: 25 a / a
Tabby LVRN[16]	Blotched tabby (t^b)	c.176C>A A1: g.94477275C>A A1: g.97133509C>A	+ : 16 t^b : 0 + / +	+ : 17 t^b : 0 + / +
	Blotched tabby (t^b)	c.416C>A A1: g.94477515C>A A1: g.97133749C>A	+ : 34 t^b : 0 + / +	+ : 10 t^b : 12 + / t^b
	Blotched tabby (t^b)	c.682G>A A1: g.94477781G>A A1: g.97134015G>A	+ : 33 t^b : 0 + / +	+ : 22 t^b : 0 + / +

	Blotched	c.2522G>A	+ : 0	+ : 5
	tabby	A1: g.94536796G>A	t^b : 10	t^b : 3
	(t^b)	A1: g.97192983G>A	t^b / t^b	+ / t^b
Ticked DKK4[17]	Ticked (ti)	c.188G>A B1: g.40429492G>A B1: g.41805975G>A	+ : 36 ti : 0 + / +	+ : 20 ti : 0 + / +
	Ticked (ti)	c.53C>T B1: g.40428846C>T B1: g.41805329C>T	+ : 33 ti : 0 + / +	+ : 30 ti : 0 + / +
Longhair FGF5[18,19]	Long fur (l)	c.356_357insT B1: g.139634675_139634676insA B1: g.141368532_141368533insA	+ : 24 1 : 0 + / +	+ : 22 1 : 0 + / +
	Long fur (l)	c.406C > T B1: g.139645946G>A B1: g.141379803G>A	+ : 24 1 : 0 + / +	+ : 25 1 : 0 + / +
	Long fur (l)	c.474del B1: g.139634261del B1: g.141368118del	+ : 26 1 : 0 + / +	+ : 0 1 : 7 + / 1
	Long fur (l)	c.475A > C B1: g.139634260T>G B1: g.141368117T>G	+ : 0 1 : 26 1 / 1	+ : 0 1 : 13 1 / +
	Long fur (l)	c.577G>A B1: g.139634158C>T B1: g.141368015C>T	+ : 37 1 : 0 + / +	+ : 27 1 : 0 + / +
Hairless *Rexing* KRT71 [20,21]	Hairless (hr)	c.816+1G>A B4: g.78941699C>T B4: g.80471148C>T	+ : 40 hr : 0 + / +	+ : 25 hr : 0 + / +
	Rexing (re)	c.1108-4_1184del, c.1184_1185insAGTTGGAG, and c.1196_1197insT B4: g.78939390_78939470del,	+ : 51 re : 0 + / +	+ : 32 re : 0 + / +

		B4: g.78939389_78939390insCTCCAACT,		
		and B4: g.78939377_78939378insA		
		B4: g.80468919_80468919del,		
		B4: g.80468918_80468919insCTCCAACT,		
		and B4: g.80468826_80468827insA		
	Rexing	c.445-1G>C	+ : 37	+ : 29
	(re)	B4: g.78943283C>G	re : 0	re : 0
		B4: g.80472733C>G	+ / +	+ / +
Rexing LPAR6[22]	Rexing (re)	c.250_253del A1: g.22865220_22865223del A1: g.23372953_23372956del	+ : 16 re : 0 + / +	+ : 19 re : 0 + / +
Blue iris PAX3[23,24]	Blue iris (bi)	(Inrton4: FERV1-LTR homolog insertion) C1: g.205833101_205833102insN[395] C1: g.207180017_207180018insN[395]	+ : bi : 0 + / +	+ : bi : 0 + / +
	Blue iris (bi)	(Intron4: RD114-LTR homolog insertion) C1: g.205834854_205834855insN[433] C1: g.207181773_207181774insN[433]	+ : bi : 0 + / +	+ : bi : 0 + / +

611 List of genes and phenotypes we focused on in this study, mutation position on CDS and

612 genomic coordinates provided by Fca126_mat1.0 and AnAms1.0, coverage depth of alleles, and

613 a summary of genotypes of CRFK and PG-4 cells as determined in the present study.

    Figure captions

Figure 1: Alignment results of whole genome sequencing reads of CRFK and PG-4 cells

with WT and mutant sequences of FGF5 genes. Alignment results of genome sequencing

reads around FGF5: c.474 and c.475; CRFK (a) and PG-4 (b) cells. Left string indicates the

name of sequenced read, and red colored symbols indicate coat phenotype-related mutations

Figure 2: Read coverage depth distributions of CRFK and PG-4 cells around the salmiak

and Orange loci. Read coverage depth distributions from 95 kbp upper stream to 95 kbp lower

stream of the salmiak locus (a), and those from 5.1 kbp upper stream to 5.1 kbp lower stream of

the Orange locus (b) of CRFK and PG-4 cells. Bar graph colored orange indicates coverage

depth of reads mapped to the salmiak (a) and Orange loci (b), respectively. In all distributions,

no notable declines in read coverage depth were observed in both salmiak and Orange loci,

indicating that both cats from which the CRFK and PG-4 cells express no salmiak and orange

coat.

Figure 3: Parts of upper and lower streams of alignment results of long-read sequencing

reads of CRFK and PG-4 cells with WT and KIT-FERV1 sequences of KIT intron 1.

Alignment results of long-read genome sequencing reads from CRFK and PG-4 cells to the

KIT-FERV1 that contains sequence around the region B1: g.161388687_161388688

(F.catus_Fca126_mat1.0). Left string indicates [the name of cell (CRFK or PG-4): name of

sequenced read], and red colored sequences indicate FERV1 and aligned sequences with it. Four

long reads from CRFK could align, but no reads contained the FERV1 homologue, while 6

reads from PG-4 could align, and four of the six reads contained the FERV1 homologue. This

figure shows only a part of the results at the upper and lower streams of the KIT-FERV1

sequence. See Table S1 for the alignment result with the full-length KIT-FERV1 sequence

Figure 4: Example illustrations of inferred original cats of CRFK and PG-4 cells. An

example illustration of the inferred phenotypes of coat morphologies and iris colors of the

original cats of the CRFK cell (a) and the PG-4 cell (b). CRFK cells are expected to originate

from a cat with long, black fur, lacking stripes, and with non-blue irises. PG-4 cells were

expected to be derived from a cat with long, white and black bicolored fur without stripes and

non-blue irises

Figure 5: Expression levels of coat phenotype-related and iris color-related genes in

various tissues. The expression level of each gene in each tissue is quantified as $\log_2$(transcripts

per million+1). The length of the red bar for gene A for each tissue is proportional to the

[expression levels of gene A in each tissue] / [the maximum expression level of gene A among

all focused tissues]. Most coat phenotype-related genes were found to exhibit similar or higher

expressions in various tissues other than skin, and the iris color-related gene (PAX3) was also

found to exhibit similar or higher expressions in a variety of tissues other than the opt-functional

organs

(a) CRFK

FGF5: c. 474, c. 475
↓

```
WT                                                 tccccccaccaggccaaatt ta ccgatgactgcaagttcagggagcgattccaagaaaac
FGF5:c.474del                                      tccccccaccaggccaaatt -a ccgatgactgcaagttcagggagcgattccaagaaaac
FGF5:c.475A>C                                      tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
```
Read name and sequence
```
_R_LH00504:186:22YKYJLT3:1:1136:23098:8018/2       -------------------- tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:1192:25992:16846/1      ---------------aatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:1187:1481:21444/1       ---------caggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:1220:37134:16734/1      ----cccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:2219:7629:10021/2          --cccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:2212:14989:24120/1      tccccccaccaggccaaatt tc ccgatgactacaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:2139:51206:15580/1         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1247:37291:5630/1          tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:2246:43264:25113/2      tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:2246:43255:25129/2      tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:2185:31669:12744/2         tccccccaccaggccaaatt t- ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1132:13233:8867/2          tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1106:21286:25834/1         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1172:1749:11559/2          tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1172:1740:11575/2          tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1181:35396:19906/1         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:2274:13473:12616/2         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:2274:18586:13209/2         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1267:31882:27341/2         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
_R_LH00504:186:22YKYJLT3:1:1204:11014:26171/1      tccccccaccgggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1192:25992:16846/2         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1161:30662:6960/2          tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:1161:30671:6976/2          tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaac
LH00504:186:22YKYJLT3:1:2212:14989:24120/2         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccacgaaaac
LH00504:186:22YKYJLT3:1:2157:26510:25049/2         tccccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaaa
LH00504:186:22YKYJLT3:1:2144:20306:10389/2         tccccccacccggccaaatt tc ccgatgac------------------------------
```

(b) PG-4

FGF5: c. 474, c. 475
↓

```
WT                                                 cccccaccaggccaaatt ta ccgatgactgcaagttcagggagcgattccaagaaaaca-
FGF5:c.474del                                      cccccaccaggccaaatt -a ccgatgactgcaagttcagggagcgattccaagaaaaca-
FGF5:c.475A>C                                      cccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaaca-
```
Read name and sequence
```
_R_LH00504:188:22Y5KLLT3:4:2128:16034:3147/2       -------------aaatt -a ccgatgactgcaagttcagggagcgattccaagaaaacag
_R_LH00504:186:22YKYJLT3:1:1209:31290:3628/1       --------aggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2102:7361:11959/1          -------caggccaaatt -a ccgatgactgcaagttcagggagcgattccaagaaaacag
_R_LH00504:186:22YKYJLT3:1:1153:39436:4957/2       --cccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2154:16238:26411/2         cccccaccaggccaaatt -a ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2217:22645:11783/2         ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
_R_LH00504:186:22YKYJLT3:1:2203:41359:11014/2      ccccaccaggccaaact -a ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:188:22Y5KLLT3:4:1147:27379:6303/2          ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2249:9710:19490/2          ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:1153:39436:4957/1          ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2149:37162:21492/2         ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:1270:2452:27453/1          ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2276:50356:6095/2          ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2154:27490:12200/2         ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2136:34998:5022/1          ccccaccaggccaaatt -a ccgatgactgcaagttcagggagcgattccaagaaaacag
_R_LH00504:186:22YKYJLT3:1:1233:2174:8963/2        ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
_R_LH00504:186:22YKYJLT3:1:1233:2165:8979/2        ccccaccaggccaaatt tc ccgatgactgcaagttcagggagcgattccaagaaaacag
LH00504:186:22YKYJLT3:1:2287:47203:29279/2         ccccaccaggccaaatt -a ccgatgactgcaagttcagggagcgattccaagaaaaca-
_R_LH00504:186:22YKYJLT3:1:1202:22719:14218/1      ccccaccaggccaaatt tc ccgatgactgcaagttc---------------------
LH00504:186:22YKYJLT3:1:1222:6446:7233/1           ccccaccaggccaaatt -a ccgatgactgc---------------------------
```

Figure 1

Figure 2

Figure 3

(a) Original cat of CRFK cell

Coat: Long

Color: black

Stripes: No

Irises: Not blue

(b) Original cat of PG-4 cell

Coat: Long

Color: White and black

Stripes: No

Irises: Not blue

Figure 4

	TYR	TYRP1	MC1R	KIT	ARHGAP36	MLPH	ASIP	LVRN	Dkk4	FGF5	KRT71	LPAR6	PAX3
skin	0.97949123	4.979143826	0.907122448	3.586949825	0.595540759	1.890367316	0.617256057	0.790593088	2.788113859	1.322693065	10.420824	4.264054274	1.420870477
skin (orange color)	1.287074139	4.681136779	1.514178733	4.728150746	0.92351867	1.992335312	0.560564858	0.730570842	2.385149963	2.506351052	11.57185462	4.780163285	1.519920078
skin (white color)	0.106508109	0	0.153761119	4.416760098	0.281844346	1.38368236	0.375739542	0.623645798	1.860833766	1.628767281	10.27795109	4.513211603	0.498482548
fibroblast adult	0.201311442	0.021819551	0.730664277	0.704854625	0.003685788	0.186873412	0.100232329	0.124979228	1.500135676	0	0.019069387	1.488602888	0.019355019
muscle	0.056483647	0	0.092118876	1.414603576	0.067172023	0.052994709	0.077034761	0.278525184	2.672768585	0.040640712	0	2.118117862	0.303226425
tooth	0.266573907	2.524633257	0.217491846	4.150102427	0.297112095	1.773640024	0.567123047	0.264163973	2.596387987	0.317583819	1.358416774	6.087545257	0.549355469
brain	0.074138775	0.244408995	6.154690407	6.125305599	0.488360957	0.106920683	0.445307287	0.482900784	2.997739706	1.214526236	0.082337234	2.495234234	0.920727113
hippocampus brain	0.104049158	0.986809378	7.141695431	6.633483047	1.263545739	0.381895847	0.462627145	0.267753817	3.196347533	2.064892	0.362603007	3.018231011	0
occipital brain	0.071192089	0.039376982	7.448888471	3.963105509	0.312113692	0.193440716	0.170021122	0.101684789	4.119749695	1.540684343	0.011456374	1.605354354	0
parietal lobe brain	0	0	7.186145561	4.875402302	0.660400161	0.64629313	0.441065494	0.042402384	3.887808995	1.329709621	0.132720662	2.664134262	0
temperal lobe brain	0	0	7.599838074	5.105537251	0.42981211	0.928785547	0.501174959	0.149430158	4.035299228	0.314304896	0.031937969	2.246188765	0
cerebellum brain	0.022500547	0.331065581	5.309532593	7.45398614	0.251584181	0.41274569	0.149029272	0.025240316	4.380271309	2.432348486	0.042684587	1.665399115	3.984523055
spinal cord	0	0.374623015	6.555970102	3.816470814	4.054688656	0	0.529844749	0.020240874	3.532849165	0.36837159	0	2.899959482	1.204872472
ear cartilage	0.103888854	0	1.723159734	5.477792674	0.473531193	1.938874602	0.644717239	2.07648309	2.615927848	0.123064897	0.133560503	3.788115166	0.40246664
ear tip	2.697859441	5.195579694	2.58708282	4.675513343	1.711850594	3.075465823	0.814532665	0.396620186	1.503211535	1.193018933	11.47542623	4.398175564	1.791957956
optic nerve head glial	4.130895559	8.012365116	1.665723362	3.485028822	0.400480608	1.551928804	0.247193444	0.058079205	4.050267469	0.184663749	0.013562176	4.736100557	2.687600024
retina	4.87612993	8.045708036	4.178402823	4.258174233	1.512735409	4.36109265	0.314914546	0.039583889	3.898728626	0.436863555	0.047022265	2.713403406	4.612167496
iridiocorneal angle	0	0	0	3.398316494	1.340508557	0	0.832811799	0.294549849	3.890824087	0.021837031	0.167713357	4.251573057	0
thymus	0	0	0	2.877720336	0	0.053402831	0.193039961	0.157927938	2.951650685	0	0	5.682280737	0.275646391
lung	0.46317526	1.08688907	1.00873704	4.4408657	0	3.890849264	2.986203115	0.073203084	2.716721885	0	0.036791184	3.877821731	0.02499963
blood	0	0	0	0.006915628	0.004805397	0.001327058	0	0.001050701	0	0	0	0.507191811	0
pbmc adult	0.074872749	0.063952512	0.502852639	1.648542894	0.074860346	0.082377222	0.074963936	0.028480146	1.806553716	0.036201871	0.260477102	2.751137087	0.043494327
pbmc kitten	0.028385287	0	0.12101391	0.275256019	0.026084545	0.034435013	0.006286765	0.002297281	1.584521711	0.020367796	0	2.368045487	0.018412154
peripheral blood	0	0	0.515689299	0.170624486	0.000698519	0.006921187	0.009584428	0.061271917	2.006770673	0	0	1.438811972	0.003684287
heart	0.029440623	0.063653346	0.446935034	3.011956496	0.021684551	0.60214092	0.712725987	0.406792622	2.349859914	0	0.109389212	3.178257348	0.056555256
liver	0.002312635	0	0.020241341	1.302396352	0.000849826	1.797793796	0.01719273	0.102107549	0.818617394	0.001655185	0	3.134084928	0.00448134
pancreas	0	0.013125298	0	0.496673638	0	1.264438441	0.021037088	0.006750463	0.281244484	0	0.017165925	1.504350567	0
spleen	0.012814297	0.095348535	0.25874521	2.535923479	0.028089502	0.571690571	0.093113907	0.028619065	1.858826078	0.089280815	0.024382448	2.679015895	0.016545063
kidney	0.096950442	0.362604563	3.114634737	0.733737527	0.043426715	0.056746401	0.215789184	0.42457845	2.804342234	0.117904886	0.028226921	1.230818933	2.817516663
small instestine	0.02306041	0.022430093	0.919040713	3.996618633	0.024456245	2.577793702	0.093414428	0.128730281	1.772402464	0.199878752	0.009835045	4.093872667	0.005000054
uterus	0	0	0.205531703	4.085978971	0	1.742410888	0.315615181	4.607945819	2.968203859	0.035454435	0	4.719142051	0.032066243
ovarian cortex	0.04283244	0.276187361	0.582670387	3.53684483	0.843720594	0.982349469	0.448130552	1.012993997	2.856184178	0.172768503	0.096062791	3.784122401	2.362116768
ovarian follicles	0.00495286	0.230281623	0	1.531380368	0.345167523	1.761071281	0.266114162	0.171579633	0.76826987	1.00405354	0.336193392	2.984532996	3.591937717
ovarian granulosa cell	0.008991358	0	1.641464675	1.34165827	0.003308901	0.007291401	0.008893111	1.374887713	1.83266614	0.06930108	0	1.809366183	0.017385505
sorted germ cells	0.642845699	0.171782823	1.686028601	4.851533678	0.507402211	1.575586521	1.613303246	0.984520286	2.500689812	0.319754165	0.724100131	3.873686288	0.73733179
testes	0.299475317	0	0.465669398	3.948613829	0.451191728	2.258861052	0.93877363	0.454226974	2.136381202	0.185946121	0.413609009	1.801861257	0.968554674
testis	0.24053022	0.125749477	1.0236267	3.966700341	0.83726628	1.840459266	0.733582575	0.640244481	2.440294652	0.169048244	0.723056047	2.635496549	0.961670128
seminiferous tubules	0.384927802	0.035315508	1.033753649	5.559293771	0.280038998	1.628919203	0.768256782	0.664462007	2.059442182	0.107003835	0.753137427	2.890890396	0.720638172
whole testes	0.242898475	0.107476514	0.504191868	3.382755818	0.445615224	2.131378771	0.599954091	0.501315507	2.128284656	0.053722971	0.727927623	3.216319425	0.293530387
embryo fetus	0.191399234	0.084838436	1.583090233	3.563052317	0.6927721	0.201677545	0.789584671	0.1446442	2.490336787	0.429363083	0.141768181	4.068385141	2.890874164
head embryo	0.277447948	0.107457441	1.063088406	4.490549175	1.944556238	0.246320218	0.978889696	0.151267039	3.035272045	1.711925434	0	4.256039654	4.241113543
body embryo	0.29294407	0.127453247	3.118263135	4.178248004	2.802053829	0.227815419	1.051822179	0.217242241	3.044565518	1.572135718	0.009663073	4.408907278	3.509347983

Figure 5

Table S1: Alignment results of long-read sequencing reads of CRFK and PG-4 cells

(File: Table_S1)

Alignment results of long-read sequencing reads of CRFK and PG-4 cells with WT and KIT-
FERV1 sequences of KIT intron 1.

Table S2: Gene expression levels in various tissues

(File: Table_S2)

Gene expression levels (transcripts per million) in various tissues estimated from domestic cats
(Felis catus) in the Ensembl database
(https://ftp.ensembl.org/pub/data_files/felis_catus/F.catus_Fca126_mat1.0/rnaseq/)

Table S3: Lists of polymorphisms in MHC class I and II regions of CRFK and PG-4 cells.

(File: Table_S3.zip)

Variant Call Format (VCF)[72] files, text files providing lists of SNPs and small indels, for MHC
class I region in CRFK (a) and PG-4 (b) cells, and those for MHC class II regions in CRFK (c)
and PG-4 (d) cells. These VCF files were obtained as follows: 1) Similar to "Mapping of paired-
end reads and quantification" in the Methods section, .bam and .bam.bai files of the mapping
results of the original paired-end short read data of the whole genome sequencing to the
reference genomes were obtained. Here, genome sequencing data were mapped to the reference
genomes "Felis catus FLA distal class I region genomic sequence" (GenBank: EU153402.1) and
"Felis catus FLA extended class II, class II, class III, proximal and central class I region
genomic sequence" (GenBank: EU153401.1) for MHC class I and II, respectively. 2) Using the
"MarkDuplicates" program of Picard (ver. 2.27.4) to obtain .bam files with the identification of
duplicate reads. 3) By applying the HaplotypeCaller program of GATK (ver. 4.3.0.0) to
previously obtained .bam file and .fasta file of reference genomes with default parameter values,
VCF files of SNPs and small indels lists were obtained.

Table S4: Lists of polymorphisms in mitochondrial DNA of CRFK and PG-4 cells.

(File: Table_S4.zip)

VCF[72] files for the mitochondrial DNA in CRFK (a) and PG-4 (b) cells. VCF files were

obtained similarly to those presented in Table S3 using "Felis catus mitochondrion, complete

genome" (GenBank: NC_001700.1) as the reference genome.