A catastrophic marine mortality event caused by a complex algal bloom including the novel brevetoxin producer, *Karenia cristata* **(Dinophyceae)**

Shauna A. Murray[1], Christopher J S Bolch[2], Steve Brett[3], Cheong Xin Chan[4], Mark Doubell[5], Hazel Farrell[6], Greta Gaiani[1], Hannah Greenhough[7], Gustaaf Hallegraeff[2], Tamsyne Smith Harding[3], D. Tim Harwood[7], Robert G. Hatfield[8], Neil MacDonald[9], Ian Moody[4], Hugo Bastos de Oliveira[10,12], Lesley Rhodes[7], Andrew Selwood[7], Justin Seymour[11], Nachshon Siboni[11], Anastasiia Snigirova[12,] Nikola Streiber[9], Anne Rolton[7], Clinton Wilkinson[10,13], Kirsty Smith[7]

[1]University of Technology Sydney, School of Life Sciences, PO Box 123 Broadway, Sydney, NSW 2007, Australia. https://orcid.org/0000-0001-7096-1307

[2]Institute for Marine and Antarctic Studies (IMAS), University of Tasmania, Private Bag 129, Hobart,7001, Tasmania Australia. https://orcid.org/0000-0001-8464-7343 (GMH)

[3]Microalgal Services, 308 Tucker Rd, Ormond VIC, 3204, Australia

[4]Australian Centre for Ecogenomics, School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD 4072, Australia. https://orcid.org/0000-0002-3729-8176

[5] South Australian Research and Development Institute (SARDI), PO Box 120, Henley Beach, SA 5024, Australia

[6]NSW Department of Primary Industries and Regional Development (DPIRD), Biosecurity and Food Safety, Taree, NSW 2430, Australia

[7]Cawthron Institute, 98 Halifax Street East, Nelson 7010, New Zealand

[8]Centre for Environment, Fisheries and Aquaculture Science, Dorset, United Kingdom

[9]Seaford Meadows SA, 5169 Australia

[10]Department of Primary Industries and Regions (PIRSA), SA, Australia

[11]University of Technology Sydney, Climate Change Cluster, Ultimo NSW 2007, Australia

[12]Flinders University, Sturt Road, Bedford Park 5042, SA, Australia

[13]South Australian Shellfish Quality Assurance Program (SASQAP), Port Lincoln, SA, Australia

Corresponding Author email address: Shauna.Murray@uts.edu.au

Abstract

Harmful algal blooms of *Karenia brevis* (Dinophyceae) are a global anomaly, occurring in one location worldwide, causing severe marine and acute human impacts via brevetoxins (BTX). During 2025 an unprecedented, currently ongoing, mass marine mortality occurred in South Australia, across an area of ~20,000 km^2, persisting for >6 months, resulting in the deaths of ~10^6 marine animals of >550 taxa, with human health impacts. Using custom metabarcoding, long-read sequencing and targeted quantitative PCR, we characterized the microalgal assemblage. *Karenia cristata* dominated over the sampling area, in an assemblage with four other *Karenia* species with varied abundances spatially and temporally. High abundances of *K. cristata* appeared in the austral autumn, and hydrodynamic processes appear to have entrained cells coastward in the semi-enclosed seas. We isolated the species and characterized it using light and electron microscopy, liquid chromatography mass spectrometry and a toxicity assay. We show for the first time that the rare and little known *K. cristata* produces significant BTX with a profile (BTX 2, 3, B5), differing from *K. brevis*, with toxicological effects. These findings reveal a novel, significant BTX-producing *Karenia*, which considering its substantial detrimental marine ecosystem impacts, is an emerging international threat with unknown consequences in changing ocean conditions.

Fish-killing marine harmful algal blooms (HABs), particularly of dinoflagellates (Dinophyceae), are relatively common worldwide, and the vast majority are of a short duration (weeks), with a localized impact[1]. Since March 2025, a marine harmful algal bloom (HAB) off coastal South Australia has spread to encompass an area of ~20,000 km^2, resulted in the deaths of millions of individuals of at least 550 marine species[2]. At the time of writing, October 2025, marine mortalities continue[2]. Acute, usually self-limiting human respiratory symptoms have been commonly reported over 6 months in South Australia including the city of Adelaide[3]. Two government inquiries have been established to investigate the causes and consequences of this HAB. Among 849 marine mass mortality incidents recorded in an international database, HAEDAT [4] (Supplementary Table 1), this situation ranks among the most impactful of ichthytoxic HABs in marine taxa killed.

In March, a species of *Karenia* (Dinophyceae), *K. mikimotoi,* was identified as present[5], and later, brevetoxins were detected in shellfish[3]. Since the mid 20th century[6,7] it has been recognized that *Karenia brevis* (Dinophyceae) produces potent toxins causing mortalities in marine biota[8] across trophic levels: zooplankton, invertebrates, fish, birds, marine mammals, in the south-eastern United States[8-10]. In the US, these toxins can become aerosolized and cause respiratory symptoms, as well as accumulate in seafood causing neurotoxic shellfish poisoning in humans [11,12]. *K. brevis* toxins were characterised as brevetoxins (BTX/PbTx), lipid soluble polyether compounds that act on voltage-gated sodium channels in vertebrates[13]. *K. brevis* HABs have been a long-standing international anomaly due to: 1) their presence and nearannual recurrence at only one site worldwide, 2) potentially very long durations of HABs (up to 20 months[14]) and 3) severe environmental and acute human health effects[9,12,14-16]. While at least 10 other *Karenia* species, including *K. mikimotoi* can cause large-scale HABs [15,17-21], BTX-producing *Karenia* are rare. Those that have been reported*,* in New Zealand in 1993 [22,23], and in the Mediterranean [24] have not re-occurred. *Karenia* species other than *K. brevis* producing BTX at a significant concentration[25] elsewhere have not been identified.

Complicating the investigation into causes of this HAB is that most *Karenia* are cosmopolitan and co-occur [15]. *Karenia* HABs often consist of 4-5 taxa that differ in toxins produced, ecology and physiology, but are morphologically highly similar or overlapping[17-19,26,27]. Besides BTX, *Karenia* species produce: gymnocin [28], gymnodimine [29], brevisulcenals, brevisulcatic acids [30], brevisin [31], brevisamide [32], tamulamides [33], and other toxins, as well as produce toxic effects due to polyunsaturated fatty acids, superoxide, and unidentified allelochemicals [1,34-36]. High concentrations of *Karenia* can produce mucilage and anoxia that contribute to marine mortalities[1]. Spatially and temporally distinct *Karenia* assemblages can cause differing impacts, such that the ichthyotoxic effects of a long-running HAB may vary over time and across areas [37].

A complex *Karenia* assemblage complicates our understanding of ecological factors contributing to HAB development. Physiology and ecology can differ markedly from one *Karenia* species to another

[15]. *K. brevis*-dominated HABs have been hypothesized to initiate in oligotrophic waters off-shore, followed by coastward entrainment via oceanographic processes [38]. The extent to which 'bottom up' factors, such as increasing temperatures and locally high nutrient coastal inputs are potentially accelerating *Karenia* growth, in comparison to autecological factors and physical concentration effects, is uncertain[38]. In the context of rapidly warming ocean waters[39] and increasing coastal phytoplankton blooms[40], the role of changing marine environments in enhancing and intensifying *Karenia* HABs is crucial to establish. Here, we applied a systematic approach to identify and characterize the causative species of the mass marine mortality event, its toxic effects, and factors contributing to this HAB.

Main

Environmental and single cell amplification and sequencing

South Australia has a coastline comprising the large semi-enclosed seas of Spencer Gulf (~320 km long, 116 km maximum width, mean depth 24 m) and Gulf St Vincent~170 km long, 50 km maximum width, mean depth 21 m), on the south-eastern side of which is the city of Adelaide (Fig. 1a). From March to September 2025, we collected water samples at 39 nearshore and ocean locations (Fig. 1a-c, Supplementary Table 2). Site locations 18-20 in the vicinity of the city of Adelaide (Fig. 1a) were sampled weekly in triplicate from 27[th] July 2025 – 13[th] September 2025, with weekly sampling continuing at the time of writing. Using a custom, dinoflagellate-specific, long-read MinION (Oxford Nanopore Technologies) sequencing[41] method focused on regions of the ribosomal RNA gene array on 18 samples, we recovered reads of ~3020 bp in length from six species belonging to the Kareniaceae family, all with potential fish-killing impacts (Fig. 1d). The Kareniaceae taxa identified were *K. cristata, K. longicanalis, K. mikimotoi, K. papilionacea,* as well as *Karlodinium veneficum* and *Takayama tasmanica* (Fig. 1d). In a phylogenetic analysis of recovered consensus sequences in relation to reference sequences from the NCBI database, each environmental sequence clustered with full support (Fig. 1d) into clades with voucher specimens or multiple identified cultured strains of that taxon (Fig. 1d, Supplementary Table 3).

We used a custom-dinoflagellate Illumina metabarcoding sequencing[42] method to select amplicon sequence variants (ASVs) identified as Kareniaceae species from 13 marine samples (Fig. 1a, b, Supplementary Tables 2 and 4). We recovered ASVs with high affinity to the species *K. cristata, K. longicanalis*, *K. mikimotoi, Karl. veneficum* and two unidentified *Karlodinium* spp (Fig. 1a, b, Supplementary Table 4). The proportions of the taxa present varied by sample, with the most abundant kareniacean taxa being *K. cristata, K. mikimotoi,* and an unidentified *Karlodinium sp* (Fig. 1a, b). *K. cristata* dominated at sites 2-4, 6, 26, 28 and 31. *K.mikimotoi* was abundant and dominant in three samples (1, 7, and 30, Fig. 1a, b), in line with the identification of that taxon based on light microscopical observation in samples from Victor Harbor (sites 1-4)[5]. At sites 5, 27, no *Karenia* were found (Fig. 1a, b). Single cell picking, followed by direct PCR and sequencing of an rRNA barcoding region (partial 28S rRNA) was conducted on cells from samples from sites 24, 35 and 36, and the sequences were found to be *Karenia cristata* (Fig 1e, samples labelled SC).

Species-specific *Karenia* quantification across 39 locations from March to September 2025

Following the identification of *Karenia* using high-throughput sequencing, we proceeded to characterize assemblages via quantitative PCR (qPCR), using published assays[43] for *K. mikimotoi, K. longicanalis,* and *K. papilionaceae* (Fig. 2, Supplementary Table 5). As critical *Karenia* diversity may not have been present in initial sequenced samples, we additionally tested for *K. brevisulcata*[43]*,* and *K.*

brevis[44]. We designed assays for *K. cristata* and *K. hui* (Fig. 2, Supplementary Fig. 1), tested them against a set of multiple strains of seven *Karenia* species (Supplementary Tables 3, 5) and found no cross-reactivity. Assays were highly efficient (Supplementary Fig 1). Across sites and months, we detected five species of *Karenia: K. cristata, K. mikimotoi, K. brevisulcata, K. longicanalis* and *K. papilionacea*, at concentrations of $1 - 10^6$ cells L^{-1} (Fig. 2a). No *Karenia brevis* or *K. hui* were detected. This showed us that *K, brevis*, the only known high BTX producing species, was not present and not a toxin source. *Karenia hui* is morphologically and genetically highly similar to *K. cristata*[18] (Fig 1d, e), and establishing its absence in this HAB was important. Of *Karenia* species present, approximately 90% of samples contained a majority of *K. cristata* compared to other *Karenia* spp, at abundances of $10^3 - 10^6$ cells L^{-1}, with lower levels of *K. mikimotoi* $\sim 10^3$ cells L^{-1} present (Fig. 2). *K. brevisulcata,* which had not been identified in the initial samples using metabarcoding or long read sequencing, was found (Fig. 2). Regional differences in *Karenia* assemblages and abundances were apparent. Samples from Spencer Gulf in June and July (sites 8, 15, Fig. 1a) and the western St Vincent Gulf (sites 12, 16, 17) were occasionally dominated by *K. papilionaceae* (Fig. 2). These samples also showed that *Karenia* could aggregate or accumulate in exceptionally dense layers in the water column. A surface slick showing a maximum concentration of 15×10^6 cells L^{-1} of *K. cristata* was found in one sample from July (site 17). Weekly samples from the sites around metropolitan Adelaide showed a consistent, high abundance of *K. cristata* from July-September, of $\sim 10^3 - 10^6$ cells L^{-1}, with lower levels of *K. mikimotoi* $\sim 10^3$ cells L^{-1} present, and trace *K. brevisulcata*, with little sign of increasing or decreasing trends over the period (Fig. 2).

Karenia cristata isolation into culture and morphological characterization

In early May 2025, shellfish farms at the western Gulf St Vincent and Kangaroo Island (Fig 1a) were closed due to the detection of BTX[3]. These were the first ever BTX detections in Australia. Hence, to characterize *Karenia* BTX producers, we isolated single cells into a seawater media in laboratory culture. In particular, we focused on *K. cristata* due to its ubiquity across locations (Fig. 1a-e). From sample 6 collected on 7th May (Fig. 1a), three non-clonal, non-axenic cultures (SM1, SM2, and SM3) were isolated. DNA was extracted and PCR was used to amplify a dinoflagellate barcoding gene (partial 28S rRNA). In the phylogenetic analysis of the resulting sequences (Fig. 1e), the cultured strains (SM1, SM2, and SM3) grouped with high support to a clade comprising the only two verified sequences of *Karenia cristata* isolated, respectively, from South Africa[45] and a French island south of Newfoundland, Canada [46].

Morphological investigations were undertaken on strain SM2 as well as *Karenia* cells from samples dominated by *K. cristata* (Fig. 3) using light and electron microscopy. *K. cristata* cells were fragile, fast swimming, dorso-ventrally compressed (Fig. 3), with numerous yellow-green irregularly rod-shaped chloroplasts to spherical (Fig, 3a-c, f), a short epicone with hemispherical outline, a longer

asymmetrical bi-lobed hypocone with the right lobe slightly longer than the left (Fig. 3c, g, h). Cingular displacement was ~28% of cell length (Fig. 3b). Cell length ranged from 21.0 to 32.2µm (mean = 24.8 µm, SD=2.32, n=40), 16.1 to 28.6 µm wide (mean=21.1 µm, SD=2.67; n=40), and 9.1-13.6 µm depth (mean=11.2 µm, SD=1.35, n=20). The length to width ratio of cells ranged from 1.06 to 1.22 (mean=1.14, SD=0.04). Living cells exhibited a straight apical that was elevated on the ventral side to form a small apical crest. (cr; Figs. 3b, c). The crest became less evident when cells were stressed or immobilised under a coverslip. Cells examined using scanning electron microscopy (SEM) showed the crest was formed by the right side of the apical groove being slightly raised (Figs. 3g-k). After Lugol's iodine preservation, some cells exhibited a more pointed epicone and a more pronounced apical crest, with the nucleus positioned lower in the hypocone (Fig. 3d). A straight apical groove extended on the ventral surface of the epicone, with a narrow depression (fe; Fig. 3h) extending dorsally to the hook-shaped ventral flange (vf; Figs. 3a, g). On the dorsal side, the apical groove extended to ~1/3 of epicone length (ag; Fig. 3i). Using DIC light microscopy, the sulcal intrusion (si) into the epicone appeared closed or slightly open (si, Figs. 3a, b), but SEM elucidated the sulcal intrusion to be either wide-open or end in a finger-like projection (Figs. 3g, h). Comparisons of morphological features to other similar *Karenia* species is given in Supplementary information 1.

Toxicology and toxicity

To examine the production of BTX by strains of *K. cristata*, an aliquot of dense culture of SM2 was extracted and examined using liquid chromatography tandem mass spectrometry (LC-MS/MS) against verified BTX reference standards for 8 analogs (Supplementary Table 7). Results showed a high concentration of the BTX analogues BTX 2, 3 and smaller amounts of B5 (Fig. 4a, b) in strain SM2 and in two samples, with an absence of other analogs at detection levels.

To examine *Karenia* related fish-killing effects, samples from locations 18 and 24, O'Sullivan Beach and West Lakes collected on 6[th] September (Fig. 1a, Supplementary Table 2), which were dominated by *K. cristata* (Fig. 2), were examined for their ichthyotoxicity using a Rainbow Trout (*Oncorhynchus mykiss*) gill cell line method [47,48]. Results showed that fish gill cells % viability after exposure to the samples declined with cell abundance, and the EC50 was 1468 cells mL^{-1} (Fig. 4c). Due to the lability of most ichthyotoxins[1] and research showing BTXs can stick to plastic[35], these findings of ichthytoxic effects may be underestimated.

Chlorophyll a anomaly and temperatures over the period

We used the remotely sensed chlorophyll-a concentration anomaly, as compared to a 21 year climatology (2002-2023) (Fig. 5a) to determine departures from this long-term daily climatology based on the previous 10 day median filter to show the potential distribution of *Karenia* species throughout the region. This method shows that chlorophyll a rose sharply during the study period in relation to

previous years, reaching concentrations well above the typical autumn baseline (Fig. 5a). Phytoplankton biomass expanded northwestward in autumn through the Investigator Strait and into western Gulf St Vincent (Figure 5b), reaching the Adelaide metropolitan coast in winter (Figure5c). Elevated biomass has persisted for several months, with highly abundant *Karenia* species observed even in deeper areas of the Gulf (~30m, Fig. 5 b,c). Throughout the period, temperatures in the area of the HAB in the austral autumn varied from around 14 – 21 ℃ (Fig 5d). During May – September, when *Karenia* cells were most abundant and widespread, temperatures were around 14 – 18 ℃ (Fig 5d).

Discussion

Here, we describe the catastrophic effects of a high brevetoxin producing marine dinoflagellate, *Karenia cristata*. Our analysis reveals that the *K. cristata* is the second identified copious Kareniaceae brevetoxin producer after *K. brevis*, a species which has a restricted distribution in the south-eastern United States[15]. *K.cristata* co-occurred in a multi-species kareniacean assemblage with four other species including *K. mikimotoi,* leading to a HAB that has, at the time of writing, lasted over >6 months, spread across an area of ~20,000 km^2 (Figs 1, 2), and resulted in devastating impacts on marine animals, as well as acute human health impacts. While marine ichthyotoxic HAB events can be common worldwide, including in Australia, of 849 reported events from 1985-2025[4] (http://haedat.iode.org), this is amongst the most destructive and widespread marine ichthyotoxic HAB, in terms of temporal and spatial extent and particularly in the diversity of marine taxa killed. Such wide ecosystem and human health effects of ichthyotoxic HABs are most pronounced for species producing known toxins [1], such as some of the Karenianceae (Supplementary Table 1), rather than raphidophytes (ie *Heterosigma*) and dictyochophytes (ie *Pseudochattonella*) that require surface contact between the microalga and marine animals for harmful effects [1].

We show that a strain of *Karenia cristata* that we isolated and cultured from the impacted region produced 3 analogues of BTXs in higher (Fig. 4, ~30- 40 pg cell^{-1}) quantities than in *K. brevis* cultures (~10-20 pg cell^{-1} [23] [49]). *K. brevis* produces brevetoxins with type A and type B polyether backbones, with BTX-1 (BTX-1) and BTX-2 (BTX-2) [13,50] the respective parent structures, with other known brevetoxins as structural derivatives of these core compounds[51]. While strains of *K. brevis* typically produce a majority of BTX 1, here, we found a notable absence of type A polyther backbones, marking a distinctive toxin profile for *K. cristata* (Fig. 4). Reports of brevetoxins not associated with *K. brevis* are very rare. A strain of *K. papilionacea* was reported to produce brevetoxins, albeit at fg levels [25]. Corsica in the Mediterranean Sea reported BTX-2 and BTX-3 in mussels once [24]. While no *K. cristata* was found at the time[24], given the rarity of BTX production in association with *Karenia* outside of the south eastern United States, it may be possible that *K. cristata* was present and unrecognised.

K. cristata has only ever been previously reported from two locations internationally: False Bay and Walker Bay, South Africa, in the mid-1990s, and a remote island off the coast of Newfoundland (French Territory) [46]. In South Africa, *K. cristata* caused an ichthyotoxic HAB (Supplementary Table 1), causing large mortalities of marine life in March 1989 [52] and in the same region in 1995-1996 [45,53]. Human health symptoms included eye, nose, throat and skin irritations, the exact cause of which was not further investigated. In HABs caused by *K. brevis*, brevetoxins commonly cause skin irritations, and can become aerosolised as the fragile cells break and cause respiratory distress in humans, particularly during onshore winds [54] [11,12]. The US Center for Disease Control (CDC) states that inhaling brevetoxins can lead to serious health effects including: shortness of breath, asthma

exacerbation, bronchoconstriction, bronchitis pneumonia (https://www.cdc.gov/harmful-algal-blooms/hcp/clinical-signs/symptoms-saltwater-harmful-algal-blooms.html), some of which have been very commonly reported in South Australia since March. Hence it seems likely that the reported human respiratory symptoms both in this event and the South African *K. cristata* HAB are caused by brevetoxins.

Understanding the sources of the ichthyotoxicity and marine animal mortalities in this event is complex, due to multiple Kareniacaea co-occurring and varying in abundance regionally (Figs 1-2, 4). Nonetheless, determining toxicological source is critical to decision-making regarding actions for mitigation, prevention or restoration of marine taxa, as methods may be more or less suitable depending on what exactly caused the marine mortalities. Some toxins such as brevetoxins persist environmentally for some time in the absence of *Karenia* cells, while other toxic effects are more volatile or disappear rapidly at cell death. Brevetoxins or other characterised toxins, on the other hand, may not be of high importance in causing ichythotoxicity, as pure brevetoxins may be only weakly toxic in fish gill bioassays [35, 55]. Besides *K. cristata,* the species *K. mikimotoi, K. brevisulcata, K. longicanalis* and *K. papilionacea* were present and occasionally dominated (Fig 2), and have documented ichthyotoxic effects, as reported internationally[17,18,27]. Similarly *Karlodinium veneficum* and *Takayama tasmanica* (Fig. 1, 2) have caused ichthyotoxic HABs [48]. In this sense, our results can are the starting point rather than the end in investigating the degree to which well-characterised toxins such as brevetoxins, as opposed to 1) toxins: gymnocin [28], gymnodimine [29], brevisulcanals, brevisulcatic acids [30], brevisin [31], brevisamide [32], tamulamides [33], and 2) compounds other than chemically characterised toxins, including polyunsaturated fatty acids, superoxide, and multiple unidentified allelochemicals [1,34-36] all contributed regionally or over time to the observed marine mortalities. *Karenia* species were also extremely abundant in the water column (10^6 cells L^{-1}) at some locations, in which case marine mortalities may have occurred simply due to anoxia [1].

Investigations of Australian *Karenia* diversity are important to determine whether *K. cristata* has long been present, or whether populations have established relatively recently, through processes such as ballast water exchange. *Karenia* blooms have occurred in Australia[26,56,57] but it has also been known for the past two decades that undescribed species exist in Australian waters [26,56]. Two previous HABs identified as *Karenia mikimotoi* affected Coffin Bay in March 1995 and February-March 2014, and resulted in deaths of fish and invertebrates [58]. A HAB of *K. longicanalis*[56] caused extensive fish kills in Tasmania. Australia has previously recorded 6 species of *Karenia,*7 species of *Karlodinium* and 3 species of *Takayama,* most of which are capable of ichthyotoxicity (Supplementary Fig. 2). In re-investigating the source of fish deaths in the 2014 South Australian (Coffin Bay) HAB using our environmental DNA samples[57] (Supplementary Fig 2), we confirmed that *K. mikimotoi* was the only *Karenia* species present at the time. Our inability to determine whether *K. cristata* has long been

present or not in Australian waters highlights an urgent need to investigate marine microbial diversity. Such studies need to use methods that allow for an understanding of the 'hidden flora' or rare species at low abundances, to establish baselines and to prepare for the changes we are experiencing in marine ecosystems.

The environmental drivers behind this unprecedented HAB are unresolved, but may be multifarious and linked to anomalous oceanographic conditions. The onset of the bloom closely coincided with a temporally extensive and periodically severe marine heatwave (MHW) in this region[59], with water temperature anomalies in the range of 2-3º C (Fig. 5). Globally, increased frequency of HABs has been linked to rising seawater temperatures[60]. An unprecedented HAB caused by another *Karenia* species, *K. selliformis*, was preceded by a significant MHW in Japan[61]. However, different *Karenia* species exhibit distinct temperature and nutrient preferences[15] and the temperature for optimum growth of *K. cristata* has not been defined. The only other previously reported HAB of *K. cristata* took place in South Africa during the austral autumn in relatively cool water temperatures of 14-17 ºC [45,52]. This pattern is consistent with the expansion of the HAB in South Australian waters after May 2025 (Fig. 5d), as the highest *Karenia* abundances coincided with lower temperature conditions. Therefore, it is possible that the timing of bloom onset was unrelated to the concurrent MHW, or was triggered by other physico-chemical conditions linked to the MHW (e.g. heightened water column stability, increased bacterial nutrient cycling). The relatively rapid and widespread geographic expansion of this event bears some of the hallmarks of interannual blooms of other *Karenia* species, which often build slowly offshore in oligotrophic waters and spread into coastal waters in high concentrations via hydrodynamic processes [38]. Temporal changes in remotely sensed chlorophyll-a anomalies during the HAB expansion across St Vincent and Spencer Gulfs (Fig, 5) align with the net seasonal circulation patterns in the region, which is governed by both wind and density-driven processes, and can entrain cells coastward in these highly saline (inverse) estuaries characterized by high evaporation and low river inputs and precipitation [62, 62, 63, 64].

Beyond the physical concentration effects of oceanographic processes, there remains the question of why this bloom has been characterised by such significant spatial expansion and temporal persistence. There is the potential that autecological processes, acting in synergy with oceanographic triggers, could be at play. For instance, several *Karenia* species show allelopathic capacity[65, 34,66,67], allowing them to kill or inhibit co-occurring phytoplankton or predators, leading to their domination of the micro-algal community. Laboratory experiments of *K. cristata* physiological ecology would be necessary to establish these factors. The sudden emergence of *K. cristata* as a potentially new cause of significant coastal HABs, specifically, only the second known *Karenia* HAB producing unprecedented concentrations of highly damaging brevetoxins, has occurred in the context of evidence that climate change has expanded conditions favourable for HABs worldwide [60, 40,68, ·] Given that *K. cristata* has a

distribution in the North and South Atlantic Ocean as well as the Great Australian Bight, and could potentially impact many countries with suitable conditions, this highlights an urgent new need to develop a solid mechanistic understanding of the triggers of the development and persistence of *Karenia cristata* HABs within multispecies Kareniaceae assemblages with varying toxicologies. Such an effort is critical to establish early warning and mitigation strategies to protect environments and human health from brevetoxin exposure.

Methods

Water sample collection

Water samples were collected between 18 March and 13 September 2025 from 39 sites off the South Australian coast (Supplementary Table 2, Fig. 1a-c), from nearshore locations (18[th] March-13[th] September 2025, Fig. 1a) and on two occasions samples at sea, onboard SARDI's RV Ngerin (22[nd]-23[rd] May 2025, Fig. 1b) and FPV Southern Ranger (16 July 2025, Fig. 1c). Locations were presented (Fig. 1a-c) using QGIS [69], and an Australian national basemap from Geoscience Australia. Where indicated (Supplementary Table 2), surface samples were a direct sample within the top 0.5 m of the water column either collected by boat or from the jetty or shore. Three samples were collected via a 25cm integrated tube sampler (4m). Sub-surface samples were collected via a 5 L Niskin bottle from 1m or at the deep chlorophyll maximum (DCM), as determined by CTD cast using a SBE 19+ CTD profiler (Sea-Bird Scientific).

Unpreserved (live) samples were stored cool, in the dark and transported as soon as possible to the laboratory at ambient temperature. Sub samples (0.1-1 L) of live samples were filtered on 5 μm Durapore (Merck, Millipore) filters using a vacuum pump and held at -20 ºC until further processing. Samples indicated by ^ in Supplementary Table 2 were fixed with acidified Lugol's solution (~5ml/L) and stored in the dark before filtration. At locations indicated by * in Supplementary Table 2, water samples were collected on filters and preserved in Longmire's buffer using an onsite gravity filtration method described by Ruvindy et al (2024).

Remote sensing

Samples in deeper water (>10m) were targeted for periodic blooms using chlorophyll-A IMOS Surface Remote Sensing products using the OCI Algorithm (Hu, Lee, & Franz, 2012) and quality control. We merged both Visible Infrared Imaging Radiometer Suite (VIIRS) abord the NOAA20 and SNPP satellites, and use the Moderate Resolution Imaging Spectroradiometer (MODIS) to compute a 21 year climatology (2002-2023), via LombScargle Periodogram (Scargle; 1982, VanderPlas 2017). Anomalies (Fig. X) are departures from this long-term daily climatology. Intermittent blooms were characterised by a previous 10 day median filter, due to cloud cover. Although estimates in very shallow water (<15m) are in generally highly biased and noisy, the signal increases detected in shallow waters were much higher than any previous estimates (not shown).

Microscope observation

Light microscopic examinations were undertaken on live and Lugol's preserved cells using Zeiss (Axiolab A1, Axioscope A1 and Axioplan 2 Imaging) upright compound microscopes equipped for bright field, phase-contrast, DIC and epifluorescence microscopy. Samples were examined directly as

wet mounts on slides, or after gentle concentration using gravity-assisted membrane filtration. Cell counts were conducted in Sedgewick-Rafter Counting cells. The abundance of cells in culture was established by counting subsamples preserved in Lugol's Solution using Utermöhl chambers and an inverted microscope (Olympus CK41 microscope). Cell abundance was used to calculate the number of cells in the DNA extractions to generate cell based standard curves for quantitative PCR analyses.

For scanning electron microscopy, live cells in seawater were fixed (2h) with freshly opened 1-2% osmium tetroxide, washed 3 times with distilled water, followed by a dehydration series of ethanol (30, 50, 70, 80, 95, 100%, 10min each), then chemically dried first with a 1:1 mixture of hexamethyldisilazane (HMDS):ethanol , followed by 100% HMDS (10 min each). Similar fixation protocols starting with Lugol or 1-4% glutaraldehyde preserved cells consistently failed to produce satisfactory results. Cells were collected on nucleopore filters, mounted on aluminium stubs, coated with platinum (5nm layer thickness), and examined with a Hitachi SU70 field emission scanning electron microscope (FESEM).

Culture methods

Live cells of *Karenia* were selected from samples enriched by addition of K medium [70] using flame-drawn Pasteur pipettes, using either a Nikon Inverted light microscope or a Leica MZ9.5 brightfield/darkfield stereomicroscope. To establish cultures, groups of 10-50 *Karenia* cells were drawn up from K/2 enrichment samples using a sterile 200 ul pipette/tip, transferred to a 12-well plate, and incubated at 17 C and 12:12 light (80 umoles PAR m-2s-2) for 1-3 days. After incubation, over successive weeks, any non-target species were carefully removed by micro-pipette, and plates returned to the same incubation conditions. From samples collected in May 2025 at site 6 (Fig. 1a), three non-clonal, non-axenic cultures (SM1, SM2, and SM3) were isolated and maintained.

DNA extraction

DNA was extracted from water samples and cultures using the DNeasy 96 PowerSoil Pro QIAcube HT Kit (Qiagen) following the manufacturer's instructions and stored at –20 ▢ until further analysis. Nineteen *Karenia* and *Karlodinium* cultures from the Cawthron Institute Culture Collection, New Zealand, were used as standards (Supplementary Table 3). Set volumes of culture were filtered onto 0.45 µm Durapore® PVDF Membrane filter (Sigma-Aldrich) and DNA was extracted from filters as described above.

qPCR analysis

All qPCR assays were prepared using the epMotion 5075I Automated Liquid Handling System and run on a Bio-Rad CFX384 Touch Real-Time PCR Detection System with CFX Maestro software v2.3. Each assay included three technical replicates, a standard curve, and negative controls. Standard curves

were generated using serial dilutions of DNA extracted from known cell numbers of *Karenia* and *Karlodinium* species maintained in culture (Supplementary Table 3). The reaction mixture for each assay consisted of 2.5 µL of iTaq Universal Probes Supermix (Bio-Rad), variable concentrations of forward and reverse primers and probe (Supplementary Table 5), 1 µL of template DNA, and sterile water to a final volume of 5 µL. Thermal cycling conditions were as follows: initial denaturation at 95 °C for 3 minutes, followed by up to 45 cycles of 95 °C for 15 seconds and 60 °C for 1 minute. For quality control, the coefficient of variation (%CV) was calculated for the Cq values of technical triplicates. If the %CV exceeded 2%, one replicate was excluded. Biological samples with a %CV greater than 5% were omitted from further analysis. Final data were normalized to cell copies per liter of seawater.

Single-cell qPCR

For the identification of individual cells of *Karenia* species present in samples, swimming live cells were picked individually from samples using a fine glass Pasteur Pipette, using a Nikon Inverted light microscope. Cells were individually photographed under 40 x magnification, then placed in PCR tubes, and frozen at -80 degrees. PCR reagents were added directly to tubes.

Illumina metabarcoding

The 28S ribosomal RNA (rRNA) gene was amplified from select environmental samples using primers targeting the D1-D2 region of the 28S rRNA and included the tails for NEXTERA sequencing: D1R-F (5´-ACC CGC TGA ATT TAA GCA TA-3´)[71] and 305-R (5´-TTT AAY TCT CTT TYC AAA GTC C-3´)[42]. PCR reactions were undertaken with 450 nM of each primer, 12.5 µL of 2 × MyFi™ Mix (Bioline), approximately 5 ng of DNA and sterile water for a total reaction volume of 25 µL[42]. Cycling conditions were: 95 °C for 5 min, followed by 35 cycles of 94 °C for 30 s, 54 °C for 30 s, 72 °C for 45 s, and a final extension of 72 °C for 7 min. PCR products were pooled and visualised on 1.5 % agarose gel with Red Safe™ DNA Loading Dye (Herogen Biotech) and ultraviolet illumination. PCR negatives were run to assess for contamination during the PCR steps. The PCR products were purified, cleaned of primer dimers, and normalised using SequalPrep™ Normalization Plate (ThermoFisher), and submitted to Sequench Ltd. (Nelson, Aotearoa-New Zealand) for library preparation.

For the bioinformatic analysis, raw reads were processed, after primers were removed with cutadapt[72], using the DADA2 package[73] within R. Reads were truncated to 200 base pairs (bp) and filtered with a maxEE (maximum number of 'expected errors') of 2 for both forward and reverse reads (reads not reaching this threshold were discarded). DADA2 constructs a parametric error matrix (based on the first 108 bp in the dataset), dereplicates the samples, and sequence variants for the forward and reverse reads are inferred based on the derived error profiles from the samples. Singletons observed in the

inference step are discarded. Subsequently, paired-end reads were merged with a maximum mismatch of 1 bp and a required minimum overlap of 10 bp. Forward and reverse reads that did not merge were not included in further analysis. Chimeras were removed using the function removeBimeraDenovo. The resulting chimera-checked, merged amplicon sequence variants (ASVs) were used for taxonomic classification using a BLAST against both the National Center for Biotechnology Information (NCBI) database and a custom BLAST database [42]. The results were parsed into a table using the phyloseq package [74]. Negative controls were assessed and the sum of reads from contaminating ASVs was subtracted from the samples. The number of reads identified as Kareniaceae species from each sample were calculated.

MinION amplicon sequencing

The primers 18ScomF1 [75] and D2C [71] were used to amplify ~3020 bp from select environmental samples as described in [41]. This region comprising the almost complete sequence of the 18S through to the D2 region of the 28S rRNA gene. All PCR reactions were undertaken in 50 μL volumes composed of: 25 μL of MyFi™ Mix (Bioline), 2.5 μL of each primer (10 μM) and 5 μL of DNA extract. Cycling conditions were: 98 °C for 60 s, followed by 30 cycles of 98 °C for 10 s, 63 °C for 20 s, 72 °C for 90 s and a final extension of 72 °C for 10 min. Resulting amplicons were cleaned using Agencourt AMPure XP, and quantified by Qubit 3.0, using the 1X dsDNA high sensitivity kit (Thermo Fisher Scientific). The Oxford Nanopore Techologies (ONT) protocol for "Ligation sequencing amplicons - NativeBarcoding Kit 24 V14 (SQK-NBD114.24)" was performed following manufacturers' instructions. The library was combined with loading beads and sequencing buffer before being transferred to a MinION FLO-MIN114 (R10.4.1) flow cell. The MinION was run for 36 h using the MinIT device. Negative control samples were prepared by using average sample volumes for each step to be representative of sample preparation. Local base calling of Fast5 files was performed using the MinIT (ONT-minit-release 19.05.2) device with the "flipflop" algorithm. Demultiplexing was performed using Porechop (v0.2.3). Analysis of the resulting sequences was performed on the custom alignment tool on the EPI2ME platform via the ONT website provided by Metrichor (Cambridge, United Kingdom).

FASTQ data was aligned to a custom reference list using Minimap2 (v2.28-r1209). The reference list comprised approximately 46 Kareniaceae sequences, each covering ~700 bp of the D1 and D2 regions of the LSU rDNA. Alignments were filtered based on alignment score (AS: >400) and divergence (de: <0.05) using Bamtools, and a coverage report was generated with Samtools. Reads aligning to each species, as identified in the coverage report, were extracted and compiled into individual FASTQ files. The fastq files were submitted to NGSpeciesID (v0.2.1) to generate consensus sequences, with polishing performed using Medaka (v2.1.0). Consensus sequences were reviewed for orientation and aligned against the reference list using BioEdit (v7.2.5), followed by trimming.

Phylogenetic Analysis

To verify identity of the large subunit RNA sequences we recovered from MinION amplicon sequencing and single-cell qPCR-amplified sequences, we inferred phylogenetic relationships of these sequences integrating high-quality LSU rDNA sequences from Kareniaceae taxa available in NCBI GenBank: 50 from diverse *Karenia* spp., and 12 from *Takayama* and *Karlodinium* taxa (6 per genus, as outgroup) (Supplementary Table 6). As a comparison to a prior HAB occurring in the same region [57], we re-analysed LSU rDNA sequences acquired during the 2014 event and inferred a phylogeny incorporating more taxa of *Takayama* and *Karlodinium* as outgroup (Supplementary Table 6).

For each sequence set, multiple sequence alignment was performed using MAFFT v7.471 [76] (*linsi* mode) at default setting. Spurious and non-phylogenetically informative sites in the alignment were removed using trimAl v1.5.0 [77] with the *-automated1* option. The trimmed alignment was used for phylogenetic inference using both maximum likelihood and Bayesian methods. Maximum likelihood tree was inferred using IQ-TREE v3.0.1 [78] with ultrafast bootstrap of 10,000 replicates (*-B 10000*), and incorporating ModelFinder to select the best-fit substitution model by default. Bayesian tree was inferred using MrBayes v3.2.7 [79] using the 4by4 nucleotide substitution model with gamma distribution (in four discrete categories), ran in four Markov chains over 2,500,000 generations (sample frequency 500 at default), and burn-in at 2000 samples.

Brevetoxin analysis of water and culture samples using LC-MS/MS

Seawater samples (40 mL) were acidified by the addition of 10 µL of phosphoric acid (H_3PO_4). Subsequently, 5 mL of methyl tert-butyl ether (MTBE) was added, and the mixture was vigorously shaken to ensure thorough phase interaction. The samples were then centrifuged at $4000 \times g$ for 5 minutes. Following centrifugation, the upper organic (MTBE) layer was carefully transferred to a 20 mL scintillation vial, avoiding carryover of the aqueous phase. An additional 1 mL of MTBE was gently added to the remaining sample and then removed and combined with the initial 5 mL extract. The combined MTBE extract was evaporated under a stream of nitrogen gas at 40 °C. The resulting residue was redissolved in 200 µL of methanol (MeOH) and transferred to an autosampler vial equipped with a 300 µL glass insert for subsequent analysis. he same protocol was followed for the sample of culture SM2, albeit with reduced volumes. Here 1.75 mL of culture was extracted with 1 mL MTBE, with the final extract resuspended in 200 µL MeOH and transferred into a glass insert for analysis.

LC-MS/MS analysis was performed using a Waters Xevo-TQS UPLC-MS instrument. For chromatographic separation a Waters Acquity BEH C8, 1.7 µm, 50×2.1 mm UPLC column held at 50 °C was used. The following gradient was used with a constant 0.3 mL min^{-1} flow rate (A: 0.1% ammonium hydroxide in water, B: MeOH). Initial conditions were 90% A (10% B) held for 0.5 min,

with a linear change to 55% A at 1.5 min, followed a linear gradient to 10% A at 5 min. The solvent composition was then changed to 5% A at 5.25 min and held at this composition until 6.25 min. The column was then re-equilibrated using a linear gradient back to 90% A at 6.5 min, and then held until 8 min. The injection volume was 1 μL with samples held at 4 °C prior to injection. Positive electrospray ionization was used with the following optimised source parameters; capillary 3 kV, cone 50 V, source offset 50 V, desolvation temp 500 °C, desolvation gas 1200 L/hr, cone gas 150 L/hr, nebuliser gas 7 Bar, source temperature 150 °C. Quantitative multiple reaction monitoring (MRM) transitions, *m/z* precursor ion>product ion (collision energy, eV), for each toxin are shown in bold, with additional confirmation transitions also shown for each analyte monitored. Transitions are included for brevetoxins reported from microalgae and some B-type backbone shellfish metabolites: **BTX-1, 867.5>849.5 (12)**, 867.5>221.3 (20), 867.5>385.3 (20); **BTX-2, 895.5>301.2 (25)**, 895.5>319.2 (21), 895.5>859.4 (20), 895.5>877.4 (15); **BTX-3, 897.5>725.3 (18)**, 897.5>769.2 (15), 897.5>807.3 (10); **BTX-B5, 911.5>875.5 (19)**, 911.5>319.2 (32), 911.5>455.3 (26); **BTX-B1, 1018.5>248.1 (40)**, 1018.5>204.1 (52), 1018.5>222.1 (36), 1018.5>276.2 (41); **dBTX-B2, 1018.6>204.1 (44)**, 1018.6>248.2 (39), 1018.6>753.5 (30), 1018.6>929.5 (30); **BTX-B2, 1034.6>929.5 (31)**, 1034.6>911.3 (32), 1034.6>947.3 (33); **BTX-B4 (C16 fatty acid), 1272.7>326.3 (43)**, 1272.7>239.3 (46), 1272.7>929.7 (37), 1272.7>1016.5 (38). For instrument calibration, working standards containing various brevetoxin analogues were prepared as a 5-point calibration series from 2 ng/mL to 50 ng/mL, which was determined to be linear over this range. Samples with brevetoxin levels above the calibration range were diluted and rerun, or a higher concentration standard prepared if the calibration is still linear. For those analogues that did not have calibration material, they were calibrated of the nearest structurally related analogue (Supplementary Table 7).

RTgill-W1 viability assay

Water samples were collected from O'Sullivan Beach (Fig. 1a, site 18) and West Lakes (Fig, 1a, site 24) on 6 September 2025. Three 1ml sub-samples were counted for each sample. Samples were prepared following the same protocol as the seawater extraction described above. For each sample, 90 mL were extracted with 15 mL MBTE by mixing, then centrifuging at 3200 g for 5 min. MTBE removed on N2 blowdown at 40 °C, then residue resuspended in 0.9 mL MeOH to give a 100x conc extract. These methanolic extracts from each sample were serially diluted in 100% MeOH to a maximum of 64-fold. All samples were then diluted 100X in L15/ex, resulting in RTgill-W1 exposure concentrations ranging from the original bloom concentration down to 1/64 of that level, in 1% MeOH. The *Oncorhynchus mykiss* rainbow trout cell line (RTgill-W1, CRL-2523, ATCC, USA), was cultured following ATTC recommendation and the method of [47] and [48]. Bioassays were performed according to [47] and [48]. Statistical and graphical analyses for the bioassays were conducted in R version 4.4.0[80]. To examine the relationship between viability and microalgal cell concentration, nonlinear

dose–response models were fitted using the drm() function from the drc[18] package. Several commonly applied sigmoidal functions were evaluated as candidate models. Model selection was guided by the Akaike Information Criterion (AIC), with the model showing the lowest AIC considered the best fit. The selected model was subsequently compared with a null model using a Fisher test and validated when the p-value was below 0.05.

Acknowledgements

We thank Sam Gaylard (SA EPA), Phil Dunne, Anke-Maria Hoeffer, Carolyn Ricci, Tracey Spokes, Serhii Snihirov and Nicola Lieff for sample collection. Jacob Thomson-Laing, Karthiga Kumanan, John Pearman (Cawthron Institute) and Rosie Buselli (UTS) helped with molecular analyses. Sandrin Feig and Olivier Bibari from the University of Tasmania and Maria Byrne from the University of Sydney provided help in sample fixation and Scanning Electron Microscopy. Lucy Thompson (Cawthron Institute) assisted with culturing and microscopy. Grant Pitcher (South Africa) and Nicolas Chomerat (France) provided helpful information. We thank the captain and crew of the RV Ngerin and FPV Southern Ranger. Satellite data was sourced from Australia's Integrated Marine Observing System (IMOS), enabled by the National Collaborative Research Infrastructure Strategy (NCRIS). It is operated by a consortium of institutions as an unincorporated joint venture, with the University of Tasmania as Lead Agent.

Funding

We thank the Australian Fisheries Research and Development Corporation (FRDC) project 2025-019 for partially funding isolation into culture. We thank the University of Technology Sydney and our host organisations for infrastructure and in-kind support. NZ Funding: New Zealand Ministry of Business, Innovation and Employment (MBIE): SSIF "Safe New Zealand Seafood Safety Research Platform" (contract number CAWX1801) and the Endeavour research programme "From Reactive to Resilient: Effectively managing our changing microalgal communities" (contract number CAW2438). The Department of Primary Industries and Regions (PIRSA) and South Australian Research and Development Institute (SARDI) for in-kind support of sample collection.

Figure Legends

Figure 1. a-c Map of South Australian coastline, showing 39 sites at which water was collected (March to September 2025), including two offshore surveys in the St Vincent Gulf. **b.** 22-23[rd] May and **c.** 16[th] July 2025**.** The relevant abundance of ASVs from metabarcoding analyses are shown as donut plots, highlighting the mixed presence of several *Karenia* species and *Karlodinium*. **d, e**. Maximum likelihood tree of Kareniaceae based on LSU rDNA sequences, incorporating **d**. MinION amplicon sequences and **e.** single-cell qPCR amplified sequences generated from this study. Each tree is rooted with the clade of *Takayama* and *Karlodinium* as outgroup to *Karenia.* Sequences generated from this study are highlighted in red. Ultrafast bootstrap support (>70%) is shown on each internal node, and a thick branch length indicates Bayesian posterior probability = 1.0 in the additionally inferred Bayesian tree (see Methods). Unit of branch length is number of substitutions per site.

Figure 2. Cell abundance of distinct *Karenia* species detected in South Australian waters throughout the period of March through September 2025, based on qPCR detection. **a**. Cells L^{-1} (in log10 scale) is shown for samples collected at 35 distinct sites in March, May, June, July, August, and September 2025, for each of the detected species of *K. brevisculcata*, *K. cristata*, *K. longicanalis*, *K. mikimotoi*, and *K. papilionacea*. *K. brevis* and *K. hui* were not detected in any samples. The sites were colour coded and grouped by the seven regions as shown in the legend key. Error bar indicating standard deviation is shown for site with two or more samples. **b**. Cells L^{-1} (in log10 scale) is shown for weekly samples collected at the O'Sullivan Beach Boat Ramp (site 18), Port Noarlunga Jetty (site 19) and Seaford Beach (site 20) in the vicinity of Port Noarlunga, between 27 July and 13 September 2025. Data are shown for each detected *Karenia* species across the 8 sampling occasions during the period. Mean and error bar indicating standard deviation are shown for triplicate samples where available.

Figure 3. Morphology of *Karenia cristata* from South Australian blooms and laboratory cultures. **a-f.** LM (DIC) of *K. cristata* from blooms. **a-b**. Live cells in ventral view showing key features including the apical groove (ag), apical crest (ac), sulcal intrusion (si) and ventral flange (vf). **c**. Live cell dorsal view (n=nucleus). **d**. Lugol's fixed bloom cell (n=nucleus) with a more prominent apical crest (cr). **e**. Composite DIC/FM image of a Lugol's fixed cell from *K. cristata* culture SM2; lugol's fixed cell image overlaid by FM image of cell after DAPI staining showing the circular to ovoid outline of the nucleus (n). **f**. Ventral view live cell with elongated irregular rod-like chloroplasts (ec). **g-k**. SEM of *K. cristata* SM2 cultured cells. **g**. Ventral view with key surface features including the apical crest (cr) formed by a raised flange (rf) on right edge of apical groove (ag). **h**. Ventral-lateral view showing sulcal intrusion (si) into the epicone and a finger-like depression (fe) extending from apical groove. **i**. Dorsal view showing apical groove extending approx. 1/3 down dorsal side of the epicone. **j**.

Antapico-lateral view showing dorso-ventral compression of cells. **k**. Apico-lateral view showing straight apical groove with raised flange (rf), extending over apex of cell.

Figure 4. Toxins present and toxicological impact on a fish gill cell line**. a.** LC-MS chromatogram showing BTX standards, and BTXs detected in a seawater sample collected during the 2025 South Australia bloom, and from a *Karenia* culture derived from the same bloom event; **b**. Quantification of brevetoxin analogues per cell, illustrating similarity in toxin profiles and cellular production rates. **c**. Rainbow trout fish gill cell viability assay showing the percentage mortality in relation to differing concentrations of a water sample from West Lakes on 6th September 2025.

Figure 5. Monthly maps (pseudo-colour) of Chlorophyll-A anomalies off the South Australian coastline based on a 21 year dataset, and temperature across the period. **a**. March mean anomalies (stations 1-4), **b.** May (with stations 27-31), **c**. July (stations 32-39). The anomalies are computed with relation to a 21 year climatology (2002-2023) from MODIS and VIIRS sensors (see Methods) using the OCI algorithm. The magenta circles represent the total abundance of *Karenia* spp in the selected samples, while the light black line represents the 10m and 30m isobaths. **d.**Temperatures in the area from March to September 2025, during the HAB,

References

Hallegraeff, G. M. et al. Fish-Killing Marine Algal Blooms: Causative Organisms, Ichthyotoxic Mechanisms, Impacts and Mitigation. (2023). UNESCO. IOC Manuals and Guides Vol. 93

SA Marine Mortality Events 2025 project. https://www.inaturalist.org/projects/sa-marine-mortality-events-2025/journal/117548 (accessed 16/10/25).

South Australian Government. https://pir.sa.gov.au/sardi/aquatic_sciences/ marine_ecosystems/algal_bloom?shorturl_algal-bloom (accessed 16/10/25).

Hallegraeff, G. M. et al. Perceived global increase in algal blooms is attributable to intensified monitoring and emerging bloom impacts. Communications Earth & Environment 2, 117 (2021).

Murray, S. A. & Gaiani, G. Mystery solved: our tests reveal the tiny algae killing fish and harming surfers on SA beaches. The Conversation (2025).

Davis, C. C. *Gymnodinium brevis* sp. nov., a cause of discolored water and animal mortality in the Gulf of Mexico. Botanical Gazette 109, 358-360 (1948).

Gunter, G., Williams, R. H., Davis, C. C. & Smith, F. W. Catastrophic mass mortality of marine animals and coincident phytoplankton bloom on the west coast of Florida, November 1946 to August 1947. Ecological Monographs, 310-324 (1948).

Flewelling, L. J. et al. Red tides and marine mammal mortalities. Nature 435, 755-756 (2005).

Landsberg, J. H., Flewelling, L. J. & Naar, J. *Karenia brevis* red tides, brevetoxins in the food web, and impacts on natural resources: Decadal advancements. Harmful Algae 8, 598-607 (2009).

Naar, J. P. et al. Brevetoxins, like ciguatoxins, are potent ichthyotoxic neurotoxins that accumulate in fish. Toxicon 50, 707-723 (2007).

Fleming, L. E., Backer, L. C. & Baden, D. G. Overview of aerosolized Florida red tide toxins: exposures and effects. Environmental Health Perspectives 113, 618-620 (2005.).

Fleming, L. E. et al. Review of Florida red tide and human health effects. Harmful Algae 10, 224-233 (2011).

Baden, D. G. Brevetoxins: unique polyether dinoflagellate toxins. The FASEB Journal 3, 1807-1817 (1989).

Stumpf, R. P. et al. Quantifying *Karenia brevis* bloom severity and respiratory irritation impact along the shoreline of Southwest Florida. PLoS One 17, p.e0260755 (2022).

Brand, L. E., Campbell, L. & Bresnan, E. *Karenia*: The biology and ecology of a toxic genus. Harmful Algae 14, 156-178 (2012).

Magaña, H. A., Contreras, C. & Villareal, T. A. A historical assessment of *Karenia brevis* in the western Gulf of Mexico. Harmful Algae 2, 163-171 (2003).

Al-Kandari, M. et al. Morphological and molecular characterization of Kareniaceae (Dinophyceae, Gymnodiniales) in Kuwait's waters. Botanica Marina 68, 155-173 (2025).

Cen, J. et al. Five *Karenia* species along the Chinese coast: With the description of a new species, *Karenia hui* sp. nov. (Kareniaceae, Dinophyta). Harmful Algae 137, 102645 (2024).

Iwataki, M. et al. Morphological variation and phylogeny of *Karenia selliformis* (Gymnodiniales, Dinophyceae) in an intensive cold-water algal bloom in eastern Hokkaido, Japan. Harmful Algae 114, 102204 (2022).

Steidinger, K. A., Wolny, J. L. & Haywood, A. J. Identification of Kareniaceae (Dinophyceae) in the Gulf of Mexico. Nova Hedwigia Beihefte 133, 269 (2008).

Xu, Y. et al. The first recorded fish-killing bloom in the Beibu Gulf, China: caused by dinoflagellate *Karenia selliformis*. Frontiers in Marine Science 12, 1582234 (2025).

Holland, P. T. et al. Novel toxins produced by the dinoflagellate *Karenia brevisulcata*. Harmful Algae 13, 47-57 (2012).

McNabb, P., Rhodes, L., Adamson, J. & Holland, P. Brevetoxin—an elusive toxin in New Zealand waters. African Journal of Marine Science 28, 375-377 (2006).

Amzil, Z. et al. Monitoring the emergence of algal toxins in shellfish: First report on detection of brevetoxins in French mediterranean mussels. Marine Drugs 19, 393 (2021).

Fowler, N., Tomas, C., Baden, D., Campbell, L. & Bourdelais, A. Chemical analysis of *Karenia papilionacea*. Toxicon 101, 85-91 (2015).

de Salas, M. F., Bolch, C. J. & Hallegraeff, G. M. *Karenia asterichroma* sp. nov.(Gymnodiniales, Dinophyceae), a new dinoflagellate species associated with finfish aquaculture mortalities in Tasmania, Australia. Phycologia 43, 624-631 (2004).

Orlova, T. Y. et al. A massive bloom of *Karenia* species (Dinophyceae) off the Kamchatka coast, Russia, in the fall of 2020. Harmful Algae 120, 102337 (2022).

Satake, M. et al. Gymnocin-A, a cytotoxic polyether from the notorious red tide dinoflagellate, *Gymnodinium mikimotoi*. Tetrahedron Letters 43, 5829-5832 (2002).

Seki, T., Satake, M., Mackenzie, L., Kaspar, H. F. & Yasumoto, T. *Gymnodimine*, a new marine toxin of unprecedented structure isolated from New Zealand oysters and the dinoflagellate, Gymnodinium sp. Tetrahedron Letters 36, 7093-7096 (1995).

Harwood, D. T., Shi, F., Satake, M. & Holland, P. T. A sensitive LC-MS/MS assay for brevisulcenal and brevisulcatic acid toxins produced by the dinoflagellate *Karenia brevisulcata*. Toxicon 84, 19-27 (2014).

Satake, M. et al. Brevisin: an aberrant polycyclic ether structure from the dinoflagellate *Karenia brevis* and its implications for polyether assembly. The Journal of Organic Chemistry 74, 989-994 (2009).

Satake, M., Bourdelais, A. J., Van Wagoner, R. M., Baden, D. G. & Wright, J. L. Brevisamide: an unprecedented monocyclic ether alkaloid from the dinoflagellate *Karenia brevis* that provides a potential model for ladder-frame initiation. Organic Letters 10, 3465-3468 (2008).

Truxal, L. T., Bourdelais, A. J., Jacocks, H., Abraham, W. M. & Baden, D. G. Characterization of tamulamides A and B, polyethers isolated from the marine dinoflagellate *Karenia brevis*. Journal of Natural Products 73, 536-540 (2010).

Kuhlisch, C., Shemi, A., Barak-Gavish, N., Schatz, D. & Vardi, A. Algal blooms in the ocean: hot spots for chemically mediated microbial interactions. Nature Reviews Microbiology 22, 138-154 (2024).

Dorantes-Aranda, J. J., Seger, A., Mardones, J. I., Nichols, P. D. & Hallegraeff, G. M. Progress in understanding algal bloom-mediated fish kills: the role of superoxide radicals, phycotoxins and fatty acids. PLoS One 10, e0133549 (2015).

Mooney, B. D., Hallegraeff, G. M. & Place, A. R. Ichthyotoxicity of four species of gymnodinioid dinoflagellates (Kareniaceae, Dinophyta) and purified karlotoxins to larval sheepshead minnow. Harmful Algae 9, 557-562 (2010).

Zhang, Q. et al. Genetic diversity and distribution of *Karenia* in the eastern coastal seas of China and implications for the trends in *Karenia* blooms under global environmental changes. Journal of Environmental Management 373, 123465 (2025).

Vargo, G. A. A brief summary of the physiology and ecology of *Karenia* brevis Davis red tides on the West Florida Shelf and of hypotheses posed for their initiation, growth, maintenance, and termination. Harmful Algae 8, 573-584 (2009).

Seymour, J. R. & McLellan, S. L. Climate change will amplify the impacts of harmful microorganisms in aquatic ecosystems. Nature Microbiology, 1-12 (2025).

Dai, Y. et al. Coastal phytoplankton blooms expand and intensify in the 21st century. Nature 615, 280-284 (2023).

Hatfield, R. G. et al. The application of Nanopore sequencing technology to the study of dinoflagellates: a proof of concept study for rapid sequence-based discrimination of potentially harmful algae. Frontiers in Microbiology 11, 844 (2020).

Smith, K. F., Kohli, G. S., Murray, S. A. & Rhodes, L. L. Assessment of the metabarcoding approach for community analysis of benthic-epiphytic dinoflagellates using mock communities. New Zealand Journal of Marine and Freshwater Research 51, 555-576 (2017).

Smith, K. F., De Salas, M., Adamson, J. & Rhodes, L. L. Rapid and accurate identification by real-time PCR of biotoxin-producing dinoflagellates from the family Gymnodiniaceae. Marine Drugs 12, 1361-1376 (2014).

Gray, M., Wawrik, B., Paul, J. & Casper, E. Molecular detection and quantitation of the red tide dinoflagellate *Karenia brevis* in the marine environment. Applied and Environmental Microbiology 69, 5726-5730 (2003).

Botes, L., Sym, S. D. & Pitcher, G. C. *Karenia cristata* sp. nov. and *Karenia bicuneiformis* sp. nov.(Gymnodiniales, Dinophyceae): two new *Karenia* species from the South African coast. Phycologia 42, 563-571 (2003).

Nézan, E. et al. Genetic diversity of the harmful family Kareniaceae (Gymnodiniales, Dinophyceae) in France, with the description of *Karlodinium gentienii* sp. nov.: A new potentially toxic dinoflagellate. Harmful Algae 40, 75-91 (2014).

Dorantes-Aranda, J. J. et al. Novel application of a fish gill cell line assay to assess ichthyotoxicity of harmful marine microalgae. Harmful Algae 10, 366-373 (2011).

Rolton, A. et al. Comparative toxicity of *Karlodinium veneficum* from France and New Zealand on fish gill cells and oyster gametes. Harmful Algae 102991 (2025).

Hardison, D. R., Sunda, W. G., Tester, P. A., Shea, D. & Litaker, W. R. Increased cellular brevetoxins in the red tide dinoflagellate *Karenia brevis* under CO_2 limitation of growth rate: Evolutionary implications and potential effects on bloom toxicity. Limnology and Oceanography 59, 560-577 (2014).

Baden, D. G., Bourdelais, A. J., Jacocks, H., Michelliza, S. & Naar, J. Natural and derivative brevetoxins: historical background, multiplicity, and effects. Environmental Health Perspectives, 113, 621-625 (2005).

Wang, Z. et al. LC/MS analysis of brevetoxin metabolites in the Eastern oyster (*Crassostrea virginica*). Toxicon 43, 455-465 (2004).

Horstman, D. et al. Red tides in False Bay, 1959-1989, with particular reference to recent blooms of *Gymnodinium* sp. Transactions of the Royal Society of South Africa 47, 611-628 (1991).

Pitcher, G. & Matthews, S. Noxious *Gymnodinium* species in South African waters. Harmful Algae News 15, 1-3 (1996).

FAO. Assessment and management of biotoxin risks in bivalve molluscs. FAO Fisheries and Aquaculture Technical Paper, 551 (2011).

Mardones, J. I. et al. Unraveling the *Karenia selliformis* complex with the description of a non-gymnodimine producing Patagonian phylotype. Harmful Algae 98, 101892 (2020).

de Salas, M. F., Bolch, C. J. & Hallegraeff, G. M. *Karenia umbella* sp. nov.(Gymnodiniales, Dinophyceae), a new potentially ichthyotoxic dinoflagellate species from Tasmania, Australia. Phycologia 43, 166-175 (2004).

Verma, A. et al. Characterization and inter-strain variability in ichthyotoxicity of *Heterocapsa ovata* (Peridiniales, Dinophyceae) from temperate waters of South Australia. Frontiers in Protistology 2, 1422481 (2024).

PIRSA. Fish Kill Investigation: Coffin Bay harmful algal (*Karenia mikimotoi*) bloom February 2014. PIRSA Fisheries and Aquaculture Division Report. Adelaide: Primary Industries and Regions South Australia 31 (2014).

Schlegel, R. W. Marine Heatwave Tracker. http://www.marineheatwaves.org/tracker. (2020).

Gobler, C. J. et al. Ocean warming since 1982 has expanded the niche of toxic algal blooms in the North Atlantic and North Pacific oceans. Proceedings of the National Academy of Sciences 114, 4975-4980 (2017).

Takagi, S. et al. Controlling factors of large-scale harmful algal blooms with *Karenia selliformis* after record-breaking marine heatwaves. Frontiers in Marine Science 9, 939393. (2022).

Lennon, G. W. et al. Gravity currents and the release of salt from an inverse estuary. Nature 327, 695-697 (1987).

de Silva Samarasinghe, J. R., B., L. & Mason, L. B. Modelled response of Gulf St Vincent (South Australia) to evaporation, heating and winds. Continental Shelf Research 23, 1285–1313. (2003).

Kämpf, J., Brokensha, C. & Bolton, T. Hindcasts of the fate of desalination brine in large inverse estuaries: Spencer Gulf and Gulf St. Vincent, South Australia. Desalination and Water Treatment 2, 335-344 (2009).

Li, X., Yan, T., Yu, R. & Zhou, M. A review of *Karenia mikimotoi:* Bloom events, physiology, toxicity and toxic mechanism. Harmful Algae 90, 101702 (2019).

Poulin, R. X. et al. *Karenia brevis* allelopathy compromises the lipidome, membrane integrity, and photosynthesis of competitors. Scientific Reports 8, 9572 (2018).

Liu, Z. et al. Cell density-dependent mutual growth inhibition in harmful dinoflagellate *Karenia selliformis*. Marine Environmental Research 211, 107374 (2025).

Silva, E. et al. Warming and freshening coastal waters impact harmful algal bloom frequency in high latitudes. Communications Earth & Environment 6, 445 (2025).

Team, Q. D. QGIS Geographic Information System. Open Source Geospatial Foundation Project. http://qgis.osgeo.org

Keller, M. D., Selvin, R. C., Claus, W. & Guillard, R. R. Media for the culture of oceanic ultraphytoplankton Journal of Phycology 23(4), 633-638. (1987).

Scholin, C. A., Herzog, M., Sogin, M. & Anderson, D. M. Identification of group and strain specific genetic markers for globally distributed *Alexandrium* (Dinophyceae). Journal of Phycology 30, 999-1011 (1994).

Martin, M. Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads. EMBnet Journal 17, 10-12 (2011).

Callahan, B. et al. DADA2: High resolution sample inference from Illumina amplicon data. Nature Methods 13, 581-583 (2016).

McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PloS One 8, e61217 (2013).

Zhang, H., Bhattacharya, D. & Lin, S. Phylogeny of dinoflagellates based on mitochondrial cytochrome b and nuclear small subunit rDNA sequence comparisons. Journal of Phycology 41, 411-420 (2005).

Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular Biology and Evolution 30, 772-780 (2013).

Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25, 1972-1973 (2009).

Minh, B. Q. et al. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. Molecular Biology and Evolution, 37, 1530-1534. (2020).

Ronquist, F. et al. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61, 539-542 (2012).

R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. (2024).

Figure 3. Morphology of *Karenia cristata* from South Australian blooms and laboratory cultures. a-f. LM (DIC) of *K. cristata* from blooms. a-b. Live cells in ventral view showing key features including the apical groove (ag), apical crest (ac), sulcal intrusion (si) and ventral flange (vf). c. Live cell dorsal view (n=nucleus). d. Lugol's fixed bloom cell (n=nucleus) with a more prominent apical crest (cr). e. Composite DIC/FM image of a Lugol's fixed cell from *K. cristata* culture SM2; lugol's fixed cell image overlaid by FM image of cell after DAPI staining showing the circular to ovoid outline of the nucleus (n). f. Ventral view live cell with elongated irregular rod-like chloroplasts (ec). g-k. SEM of *K. cristata* SM2 cultured cells. g. Ventral view with key surface features including the apical crest (cr) formed by a raised flange (rf) on right edge of apical groove (ag). h. Ventral-lateral view showing sulcal intrusion (si) into the epicone and a finger-like depression (fe) extending from apical groove. i. Dorsal view showing apical groove extending approx. 1/3 down dorsal side of the epicone. j. Antapico-lateral view showing dorso-ventral compression of cells. k. Apico-lateral view showing straight apical groove with raised flange (rf), extending over apex of cell.