Dynamical memory underlies prolonged plasmid persistence after transient antibiotic

treatment

Zhengqing Zhou[1,2,*], Andrea Weiss[1,2,*], Zhixiang Yao[1,2,6], Xiaoli Chen[1,2], Kristen Lok[1,2], Hye-in

Son[1,2], Lingchong You[1,2,3,4,5,+]

1 Department of Biomedical Engineering, Duke University, Durham, North Carolina, USA

2 Center for Quantitative Biodesign, Duke University, Durham, North Carolina, USA

3 Center for Biomolecular and Tissue Engineering, Duke University, Durham, North Carolina,

USA

4 Center for Genomic and Computational Biology, Duke University, Durham, North Carolina,

USA

5 Department of Molecular Genetics and Microbiology, Duke University School of Medicine,

Durham, North Carolina, USA

6 Present address: Innovative Genomics Institute, University of California, Berkeley, California,

USA

*These authors contribute equally to the work.

+Correspondence and requests for materials should be addressed to Lingchong You. E-mail:

lingchong.you@duke.edu; Tel: 919-660-8408; Fax: 919-668-0795.

Abstract

Plasmids play critical roles in spreading and maintaining antimicrobial resistance (AMR). They often exhibit prolonged persistence upon antibiotic treatment, even when they impose substantial burden on their hosts. This persistence has been primarily attributed to rapid horizontal transfer or low plasmid cost. However, these mechanisms cannot account for the slow decay of burdensome plasmids with poor mobility. Here, we show that the decoupling of time scales between slow segregation loss and fast growth competition leads to a slow-down in plasmid abundance decay at high initial plasmid abundance, reminiscent of the *ghost effect* from nonlinear dynamical systems. Integrating theory, simulations, and quantitative experiments across clonal populations and multi-species bacterial communities, we demonstrate that a transient antibiotic pulse can eliminate plasmid-free cells and create a *ghost* state that extends plasmid persistence from days to months. Our research reveals a generalizable mechanism for the prolonged ecological memory of antibiotic exposure and underscores the need for proactive strategies to curb the spread of AMR.

Introduction

Plasmids are self-replicating genetic elements that play a critical role in bacterial evolution by enabling the acquisition, maintenance, and dissemination of adaptive traits such as antimicrobial resistance (AMR)[1-3], giving rise to many multidrug-resistant high-risk clones[4,5] and accelerating the global AMR crisis[3]. Although plasmid carriage is often assumed to be burdensome in the absence of selection, leading to rapid plasmid loss[6,7], empirical studies suggest otherwise. Resistance frequency often remains high for months after short-term antibiotic treatments (days to weeks), despite the absence of continued selective pressure thereafter[8-12]. Given that many resistance determinants reside on plasmids[11,13-19], the slow resistance reversal suggests burdensome plasmid can still experience long transients after antibiotic exposure.

Different mechanisms may account for this prolonged persistence after antibiotic treatment. For instance, positive selection can increase the frequency of compensatory mutations that reduce the cost of plasmid carriage and slow down the decay of plasmids[20]. Long half-life of certain antibiotics can exert sustained selection for resistance after the treatment ceased[9]. More broadly, the persistence or slow reversal of resistance or plasmid carriage can be associated with low or zero fitness cost[6,21-23], high conjugation rate[24-26], or both[24,27]. Co-selection by antibiotics[6,28] or environmental contaminants[29] on the same mobilizable gene element has also been proposed. However, these mechanisms cannot account for the slow decay of plasmids whose baseline abundance is low, i.e., burdensome, low-mobility plasmids get lost fast.

Past theories suggest another under-appreciated possibility: the slow segregation loss of plasmids[30,31]. While largely ignored due to its relatively minor contribution to plasmid steady state persistence[24,25,27], slow segregation loss can sustain plasmid carriage in clones for a long time [32,33], before plasmid-encoded high-burden cargoes drive the plasmid to get completely lost. This provides a theoretical basis for the prolonged plasmid carriage following antibiotic exposure:

antibiotics can eliminate the plasmid-free competitor cells, so the plasmid needs to go through

the slow segregation loss process before rapidly decrease in their abundance due to the growth

competition from plasmid-free segregant.

Here, we combine theory, numerical simulations, and quantitative experiments to

demonstrate this principle in clonal populations, synthetic *E. coli* communities, and multi-species

communities of hospital sink isolates: the decoupling of time scales between segregation loss and

growth competition can lead to the observed prolonged ecological memory of plasmids. This

principle underscores the need to develop active intervention strategies beyond standard

antibiotic stewardship.

Results

Population ghost effect slows down plasmid abundance decay

We start from a simple ordinary differential equation (ODE) model depicting the dynamics

of a non-mobilizable plasmid within a single strain population (**Methods**). The dynamics of

plasmid loss in a population are mainly dictated by two processes: *(1) segregation loss*, where a

daughter cell fails to inherit the plasmid during cell division (**Fig. 1a**, pink), and *(2) growth

competition, where plasmid-carrying cells are outcompeted by plasmid-free population due to

plasmid-imposed metabolic burden on their host (**Fig. 1a**, purple). Segregation loss is generally

a slow process, where different plasmids can deploy active partitioning, post-segregational killing,

and high copy-number to minimize its loss during cell division[34,35]. Meanwhile, growth competition

operates on a faster timescale.

We used the ODE model to simulate plasmid abundance (fraction of plasmid carrying cells

in a population) over time (**Fig. 1b**), across different initial plasmid abundances of $P_0\% = 100\%$,

99%, 90%, 50%, and 20%. When starting from a pure plasmid-carrying population ($P_0\% = 100\%$),

plasmid abundance decays in two phases, corresponding to the two processes: an initial slow

decay driven by segregation loss, followed by a faster decay driven by growth competition once

plasmid-free cells have emerged. This two-phase behavior is consistent with previous

experimental observations[32,36], where the presence of post-segregational killing mechanism

results in a long plateau in which plasmid abundance remains ~100%. In the absence of active

partitioning or post-segregational killing mechanisms, this plateau was significantly shortened or

even eliminated. As the initial plasmid abundance decreases, growth competition becomes more

prominent, driving the plasmid abundance decay exponentially. Based on a previous work[30], we

can analytically derive the half-life of plasmid abundance decay (**Methods**) which nonlinearly

increases with initial plasmid abundance (**Fig. 1c**). When $P_0\% << 100\%$, the abundance decays

exponentially with half-life $\tau_{1/2} = \frac{1}{\delta}\log 2$. When $P_0\% = 100\%$, plasmid half-life becomes

$\tau_{1/2} \rightarrow \frac{1}{\delta}\log(\frac{\delta}{\kappa})$, where δ is the rate of growth competition between the plasmid-free and plasmid-

carrying populations, and κ is the segregation loss rate. In most cases, $\kappa \ll \delta$ [24], thus half-life

$\tau_{1/2}$ becomes large, and the high abundance state appear as the *ghost* of an equilibrium[37]. This

logarithmic scaling between κ and $\tau_{1/2}$ resembles the *ghost effect* coined by Strogatz[37], originally

referring to the loss of stability in a dynamical system during saddle-node bifurcation. The

prolonged plasmid carriage in our case instead arises from the remnant of a saddle point (when

κ increases from 0 to a small number).

To test our prediction of diverging half-lives near pure population, we monitored the

abundance of three non-mobilizable plasmids with different origin of replication, pSC101

(kanamycin resistant, Kan^R), colE1 (Kan^R), and pUC (spectinomycin, Spec^R), all encoding sfGFP,

in *E. coli* strain MG1655 over time (**Fig. 1d & Fig. S1**). To facilitate high-throughput measurements,

we used the GFP/OD ratio to quantify plasmid abundance (calibration details in **Methods**).

Plasmid-carrying MG1655 cells were mixed with plasmid-free MG1655 cells at different volume

ratios to generate populations where plasmid carrying cells make up of 100%, 99%, 90%, 50% or

20% of the total population. Populations were cultured in LB and diluted 1:500 every 24 hours

(**Methods**). As predicted by the mathematical model, a plateau in plasmid abundance was

observed at high initial abundances before exponential decay (**Fig. S1b**), significantly prolonging

the half-life of plasmid abundance. Thus, the initial purity of the community, in terms of the

plasmid-carrying cell population, dictates the plasmid half-life (**Fig. 1 e - g**). Interestingly, at P_0%

= 100%, pUC persisted in the community. As a high copy plasmid[38], the stochastic chance of pUC

to have segregation error becomes extremely small, preventing the birth of plasmid-free cells and

the subsequent growth competition. Moreover, for the populations with plasmid abundance decay

(pUC with P_0% = 99% and 90%, and colE1 with P_0% = 99%), the plasmid abundances sometimes

rebounded, suggesting potential compensatory mutations.

We further quantified the dynamics of two conjugative plasmids, pCU1 and R6K in *E. coli*

DA28102, with daily 100-fold dilution in LB media supplemented with 25 µg/mL chloramphenicol

(Cm) to prevent contamination. Under these conditions, pCU1 and R6K cannot persist at steady

state and were lost rapidly for P_0% < 90%. With P_0% = 90%, pCU1 had a half-life of 4.7 ± 0.4 days

(mean ± SD, n = 3), and R6K had a half-life of 6.8 ± 1.0 days (mean ± SD, n = 3). However, both

exhibited drastically prolonged persistence when P_0% is near 100% (**Fig. 1h & i, Fig. S2**).

Transient antibiotic selection extends plasmid half-life through purifying selection

Antibiotics can eliminate plasmid-free cells in a population and purify the population to

enter the ghost effect region (**Fig. 2a**). We thus hypothesize selection with antibiotics could

prolong plasmid persistence. We assembled populations of *E. coli* MG1655 with an equal mixture

of plasmid-free and plasmid-carrying cells (P_0%=50%). We used the non-mobilizable, sfGFP-

encoding plasmids pSC101, colE1, and pUC for high-throughput measurement. We applied a

pulse of antibiotic selection to the populations between day 2 and 3. After growing the population

in LB supplemented with a gradient of antibiotics (Kan for pSC101 and colE1, and Spec for pUC)

for 24 hours, we removed the selection and tracked the GFP/OD readout over time (**Fig. 2b – d,**

Fig. S3b). The half-lives of the three plasmids all substantially increased with the strength of the

pulsed antibiotic selection (**Fig. 2e - g**), and plateaued above critical concentrations. The critical

concentrations are 3.125 µg/mL Kan for pSC101, 6.25 µg/mL Kan for colE1, and 12.5 µg/mL Spec

for pUC.

Antibiotic pulse induces prolonged plasmid carriage in synthetic *E. coli* communities

To determine whether the ghost effect occurs in more complex microbial systems, we

constructed five synthetic communities of *E. coli* Keio strains[39] (Table S1 & S2). Each strain

contains a uniquely barcoded, non-mobilizable plasmid, enabling the quantification of community

structure using next generation sequencing (NGS). The first community, Comm87, consisted of

86 background strains plus one donor strain carrying conjugative plasmid R388 (trimethoprim

resistant, Trim^R). Four other communities (Comm57) shared the same background community of

56 barcoded Keio strains, in addition to a unique donor strain carrying one additional conjugative

plasmid, either R6K (streptomycin resistant, Strp^R), pCU1 (carbenicillin resistant, Carb^R), R388,

or RP4 (tetracycline resistant, Tet^R). The overnight cultures of the background strains were

equally combined, then mixed with the donor strain at a 3:1 ratio to assemble each of the five

communities. The communities were passaged daily with a 100-fold dilution in LB media

supplemented with 100 µg/mL Carb for Comm87 or 25 µg/mL Cm for Comm57 to maintain the

barcode plasmids. We tracked the abundance of the target plasmids over time through selective

plating.

Without any additional antibiotic intervention, the conjugative plasmids exponentially

decayed (**Fig. 3a - c**, purple), with half-lives of 2.2 ± 0.4 days (mean ± SD, n = 3) for R388 in

Comm87, 4.4 ± 0.1 days (mean ± SD, n = 3) for R6K in Comm57, and 1.0 ± 0.1 days (mean ±

SD, n = 3) for pCU1 in Comm57 (**Fig. 3d - f**). R388 and RP4 were able to persist in Comm57 for

the entire course of the 20-day experiment (**Fig. 3g & h**). As predicted by the model, antibiotic

selection for the plasmids, applied between day 2 and 3, increased plasmid abundances to

approach 100%. Driving the communities into the *ghost* state drastically extended plasmid half-

lives to 27.3 ± 1.7 days (mean ± SD, n = 3) for R388 in Comm87 and 9.9 ± 0.9 days (mean ± SD,

n = 3) for pCU1 in Comm57 (**Fig. 3 d-f,** $p<0.01$ or $p<0.001$ with Welch's t-test). Plasmids R6K,

R388, and RP4 all persisted in the communities at ~100% for more than 17 days after antibiotic

selection (**Fig. 3b, g, h**, orange). The four Comm57 communities, when passaged in a different

dilution rate (1:500), showed similar results (**Fig. S4**).

We used NGS to quantify the community composition over time (**Fig. 3i & Fig. S5**). Upon

antibiotic selection, a significant shift in each community structure was observed, and the

community diversity dropped significantly (**Fig. S6**). Upon selection, the relative abundances of

the plasmid donors were significantly amplified (**Fig. 3j**), while the donor abundances decreased

in most communities without antibiotic pulse. The decline in diversity, and the increase in donor

abundance, suggests that the plasmids have not yet spread to every member in the community.

One exception was Comm57 harboring RP4, where the donor abundance was significantly

decreased upon antibiotic selection (**Fig. 3j**). While in other communities the antibiotic pulse

resulted in a crash in diversity, the RP4 community instead increased its diversity (**Fig. S6**). This

could be due to the combination of two factors: the low fitness of the donor strain, and the fast

conjugation capacity of RP4[24,40]. By the time positive selection took place, most strains in the

community would have already received the plasmid and could outcompete the donor strain in

the presence of Tet.

These findings demonstrate that the ghost effect operates not only in clonal populations

but also in synthetic microbial communities. A brief antibiotic pulse can reorganize community

composition by selectively eliminating plasmid-free competitors, thereby stabilizing plasmid
persistence at the community level without requiring genetic adaptation.

Ghost effect in multi-specific communities with niche partition

In nature, microbes can co-exist in natural communities through niche partition, where
different members can take up different resources (**Fig. 4a**) or habitats (**Fig. 4b**) to avoid
competitive exclusion. In the above experiments, we showed the ghost effect taking place in
clonal populations and *E. coli* communities, where strong competitions exist between individual
strains, and plasmid-free populations are sensitive to the antibiotics. In these cases, ghost effect
takes place when the plasmid abundance reaches ~100%. However, numerical simulations
(**Methods**) suggest antibiotic-induced ghost effect can happen without community purity (**Fig. 4**
c & d) in the presence of niche partition. For instance, consider a mixed population of plasmid-
free strain 1, plasmid-carrying strain 1, and non-host strain 2 (**Methods**). In the simple case where
(1) strain 1 and 2 do not interact and (2) strain 2 has moderate resistance to the antibiotic, the
non-host strain 2 can survive the antibiotic selection, limiting the plasmid abundance to < 100%.
However, ghost effect can still happen within the purified strain 1 population, resulting in the
prolonged carriage of the plasmid in the two-member community (**Fig. 4c & d**). In essence, as
long as the plasmid-carrying population is approaching purity in a niche, ghost effect should take
place. Since natural microbiomes often have niche partition, with different species exhibiting
natural or acquired resistance, we expect general applicability of ghost effect following antibiotic
treatments, even when the plasmid or resistance abundance does not reach 100%.

To test this idea, we constructed three communities consisting of 8 isolates from Duke
Hospital sink p-traps[41] (**Methods**), plasmid-free and plasmid-carrying *E. coli* MG1655 with one of
the three non-mobilizable plasmids (pSC101, colE1 and pUC) expressing sfGFP as a surrogate
for their abundance. The sink isolates consist of one *Pseudomonas aeruginosa* strain, three

Bacillus trains, one *Citrobacter* strain, and three *Enterobacter* strains (**Fig. 4e**). To facilitate niche

partitioning, we used cellulose sponges immersed in LB media to create porous, physically

partitioned habitats (**Fig. 4e**), which were previously found to promote the diversity of microbial

communities[42].

In the absence of antibiotic selection, the GFP/OD level (**Methods**) dropped rapidly to

below 10^2. After transient Kan or Spec selection between day 1 and 2, GFP levels were increased.

pSC101 showed slowed decay in GFP/OD level, while colE1 and pUC persisted until day 10 (**Fig.**

4 f – h). This is consistent with our observations of the three plasmids in clonal populations (**Fig.**

1, Fig. S1). Importantly, for plasmids colE1 and pUC, the GFP/OD levels were consistently below

the baseline in pure plasmid-carrying MG1655 populations.

Through 16S sequencing (**Methods**), we quantified community dynamics in the presence

and absence of the transient antibiotic treatment (**Fig. 4 i – k**). On day 2, in the absence of

antibiotics, *P. aeruginosa* and *Enterobacter* became dominant in the communities, with the other

three genera decreasing in their fractions or even become undetectable in some of the replicates.

Antibiotic treatment increased the abundance of the two resistant strains, *E. coli* and *P.*

aeruginosa, and eliminated the rest of the community. The survival of *P. aeruginosa* confirms that

the communities do not purely contain plasmid-carrying cells, but a ghost effect within the *E. coli*

population still takes place. Consistent with the GFP/OD reading of pSC101, *P. aeruginosa* was

detectable in one of the three replicate communities harboring pSC101, potentially due to the

stochastic effect of high dilution rate (2×10^7) during daily passaging.

In LB cultures without cellulose sponges, the GFP/OD levels of the communities after

antibiotic treatment became comparable to the pure plasmid-carrying baselines (Fig. S7) and

exhibited slowed-down decay in comparison to the antibiotic-free control group. Since the

plasmids are non-mobilizable, the post-treatment communities likely consisted primarily of *E. coli*.

In contrast, the survival of *P. aeruginosa* should be attributed to the presence of sponges, which

possibly introduces spatial structures to mediate the competition[42] and promote biofilm formation[43].

Discussion

Resistance limits the clinical efficacy of antibiotic treatments and poses a major threat to

human health. Many antibiotic resistance genes are maintained and propagated by mobile genetic

elements including plasmids[1-3]. Understanding and predicting the persistence of plasmids is

critical for designing intervention strategies to contain and reverse the spread of AMR. In this work,

we systematically quantified the decay dynamics of multiple plasmids across different community

contexts, and established their dependence on community structure. We showed that the plasmid

half-life drastically increases in nearly pure plasmid-carrying populations, either from assembly or

induced by transient antibiotic pulse. Theoretically, this ghost effect arises from the decrease in

the strength of growth competition when plasmid-free cells are scarce in the population.

Our results reveal a common mechanism underlying slow reversal of plasmid-mediated

resistance after antibiotic treatments, a phenomenon that spans across microbial ecosystems

ranging from the human oral, urinary, and gut microbiomes, to those of livestock. Macrolide

antibiotics have been reported to increase the abundance of resistant streptococci (primarily

carrying *mef* gene or macrolide-lincosamide-streptogramin-B resistant cassette *erm*(B)) in oral

microbiomes to more than 80%, which required over 6 months to revert to baseline[9]. The authors

attributed this prolonged resistance carriage to the long half-life of azithromycin, and the low

burden of the *erm*(B) gene. Similarly, veterinary β-lactams strongly amplified the abundance of

resistant bacteria in the pig intestinal microbiome, persisting for more than 3 weeks beyond the

drug withdrawal. The extended-spectrum β-lactamase CTX-M-1 was carried by both an

exogenous conjugative plasmid and indigenous gut microbiota. Consistent with our results (**Fig.**

3, Fig. S5), the authors observed the significant amplification of resistance abundance and donor

abundance upon antibiotic selection, followed by rapid decline of the donor strain abundance after

antibiotic withdrawal. Importantly, both the *erm*(B) cassette [13-17] and the CTX-M-1 gene[18,19] are

commonly found on conjugative plasmids and other mobilizable gene elements. Yelin et al.

reported that resistance in urinary tract infections shows autocorrelation lasting more than six

months, with strong and persistent correlation between patients' antibiotic purchase history and

subsequent resistance[12]. Anthony et al. observed increases in the abundance of β-lactam,

macrolide, and tetracycline resistance following treatment with cephalosporin (β-lactam),

azithromycin (macrolide), or both[10]. During the development of the infant gut microbiome,

resistance genes on mobile genetic elements were found to persist longer than those encoded

on chromosomes[11]. The diversity of antibiotics, resistance genes, bacterial host, and ecosystems

all point to a general population-dynamical mechanism. As the ghost effect depends only on the

slow time scale of segregation loss, it offers a potential explanation for these observations.

While our work focuses on the population dynamics of the plasmids, the ghost effect also

has strong evolutionary consequences. Prolonged plasmid carriage enables co-evolution of the

plasmid and host, creating opportunities for compensatory mutations. Long-term co-evolution

could ameliorate the burden of the plasmid[20], and may even turn the deleterious cost of plasmid

carriage to a beneficial one[22]. Thus, despite predictions that certain plasmids should not persist

based on ecological models[24,27], their long-term persistence may occur before eventual extinction.

In our experiments, we observed potential evolutionary rescues for colE1 and pUC starting at high

initial plasmid abundances (**Fig. S1**), which was not observed in populations outside of the ghost

effect region. Therefore, plasmid persistence can have bistable outcomes when evolution forces

are taken into consideration.

The antibiotic-induced persistence also provides an environment-driven explanation for

the survival of burdensome plasmids, addressing the so-called plasmid paradox[7]. Previous

studies have attributed plasmid survival to biotic factors, such as fast conjugation[24,25,44], variable

fitness effects across different hosts[23], spatial partitioning[45], and compensatory mutations that

ameliorates the plasmid burden[20,22]. These works typically assume constant environments,

without positive selections on the plasmids. However, plasmids in natural and clinical settings

experience fluctuating selection forces[46-50]. In such cases, the burdensome plasmids can persist

when the frequency of positive selection is on the same order of magnitude as the plasmid half-

life within the ghost effect region. This mechanism further raises questions about the efficacy of

simply reducing antibiotic use to contain AMR[6] – similar persistence can be achieved with far less

amount of antibiotics with sporadic selection. Conversely, pulsed antibiotic treatments has been

proposed as a strategy to constrain the development of resistance[51-55]. Such strategies should be

designed to avoid the ghost effect region to prevent slow population responses that

disproportionally favor resistant bacteria.

The clinical, ecological, and evolutionary implications of ghost effect underscore the need

for strategies that counteract against unintended antibiotic exposure. Previous studies have used

chemicals to reduce plasmid abundance and reverse its persistence[24,56]. As both plasmid

abundance and half-life are governed by conjugation, growth, and segregation events[27], we

hypothesize that these chemicals could also accelerate plasmid decay. We found that curing

chemicals and their combinations were able to reduce the half-life of pSC101 almost by half (**Fig.

S8**). A single day of treatment using chemical combinations achieved comparable effects as full

course, suggesting these chemicals may accelerate the plasmid decay by increasing segregation

error.

Overall, our work established the timescale decoupling between segregation loss and

growth competition for burdensome plasmids and demonstrated the disproportionate impact of

transient antibiotic pulses on the plasmid persistence. Our findings provide a generalizable

mechanism for the prolonged ecological memory of plasmid carriage following antibiotic exposure,

and have implications for clinical management of antibiotic use and the ecology and evolution of

plasmids.

Acknowledgments

We thank Emma Chory, Teng Wang, Rohan Maddemsetti, Emrah Şimşek, Grayson Hamrick,

Yifan Dai, and Aaron Yip for helpful discussions. This work was supported by National Institute

of Health (L.Y.: R01AI125604, R01GM098642, R01EB031869) and the National Science

Foundation (L. Y.: Cooperative Agreement No. EEC-2133504)

Author contributions

Z.Z., A.W., and L.Y. conceived the research. Z.Z., A.W., Z.Y., X.C., and K.L. performed the

experiments. Z.Z. performed the modelling and statistical analysis. Z.Z., A.W., and L.Y. wrote

the manuscript. All authors read and contributed to the manuscript.

Declaration of interests

The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT to improve the language of the

manuscript. After using this tool, the authors reviewed and edited the content as needed and

take full responsibility for the content of the published article.

Resource availability

Materials availability

Genetic constructs generated for this study are available upon request.

Lead contact

354 Further information and requests for resources should be directed to and will be fulfilled by the

355 lead contact, Lingchong You (lingchong.you@duke.edu).

357 **Data and code availability**

358 All original code and data have been deposited at https://github.com/youlab/Plasmid-Ghost-

359 Effect.

References

1 Bethke, J. H. *et al.* Environmental and genetic determinants of plasmid mobility in

pathogenic <i>Escherichia coli</i>. *Science Advances* **6**, eaax3173 (2020).

https://doi.org:10.1126/sciadv.aax3173

2 Lipworth, S. *et al.* The plasmidome associated with Gram-negative bloodstream

infections: A large-scale observational study using complete plasmid assemblies. *Nature*

Communications **15** (2024). https://doi.org:10.1038/s41467-024-45761-7

3 Castañeda-Barba, S., Top, E. M. & Stalder, T. Plasmids, a molecular cornerstone of

antimicrobial resistance in the One Health era. *Nature Reviews Microbiology* **22**, 18-32

(2024). https://doi.org:10.1038/s41579-023-00926-x

4 Hawkey, P. M. & Jones, A. M. The changing epidemiology of resistance. *J Antimicrob*

Chemother **64 Suppl 1**, i3-10 (2009). https://doi.org:10.1093/jac/dkp256

5 Mathers, A. J., Peirano, G. & Pitout, J. D. The role of epidemic resistance plasmids and

international high-risk clones in the spread of multidrug-resistant Enterobacteriaceae.

Clin Microbiol Rev **28**, 565-591 (2015). https://doi.org:10.1128/cmr.00116-14

6 Andersson, D. I. & Hughes, D. Antibiotic resistance and its cost: is it possible to reverse

resistance? *Nature Reviews Microbiology* **8**, 260-271 (2010).

https://doi.org:10.1038/nrmicro2319

7 Brockhurst, M. A. & Harrison, E. Ecological and evolutionary solutions to the plasmid

paradox. *Trends in Microbiology* **30**, 534-543 (2022).

https://doi.org:10.1016/j.tim.2021.11.001

8 Cavaco, L. M., Abatih, E., Aarestrup, F. M. & Guardabassi, L. Selection and persistence

of CTX-M-producing Escherichia coli in the intestinal flora of pigs treated with amoxicillin,

ceftiofur, or cefquinome. *Antimicrob Agents Chemother* **52**, 3612-3616 (2008).

https://doi.org:10.1128/aac.00354-08

9 Malhotra-Kumar, S., Lammens, C., Coenen, S., Van Herck, K. & Goossens, H. Effect of

azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant

streptococci in healthy volunteers: a randomised, double-blind, placebo-controlled study.

The Lancet **369**, 482-490 (2007). https://doi.org:10.1016/S0140-6736(07)60235-9

10 Anthony, W. E. *et al.* Acute and persistent effects of commonly used antibiotics on the

gut microbiome and resistome in healthy adults. *Cell Reports* **39**, 110649 (2022).

https://doi.org:10.1016/j.celrep.2022.110649

11 Yassour, M. *et al.* Natural history of the infant gut microbiome and impact of antibiotic

treatment on bacterial strain diversity and stability. *Sci Transl Med* **8**, 343ra381 (2016).

https://doi.org:10.1126/scitranslmed.aad0917

12 Yelin, I. *et al.* Personal clinical history predicts antibiotic resistance of urinary tract

infections. *Nature Medicine* **25**, 1143-1152 (2019). https://doi.org:10.1038/s41591-019-

0503-6

13 Berryman, D. I. & Rood, J. I. The closely related ermB-ermAM genes from Clostridium

perfringens, Enterococcus faecalis (pAM beta 1), and Streptococcus agalactiae (pIP501)

are flanked by variants of a directly repeated sequence. *Antimicrob Agents Chemother*

39, 1830-1834 (1995). https://doi.org:10.1128/aac.39.8.1830

14 Clewell, D. B., Flannagan, S. E. & Jaworski, D. D. Unconstrained bacterial promiscuity:

the Tn916-Tn1545 family of conjugative transposons. *Trends Microbiol* **3**, 229-236

(1995). https://doi.org:10.1016/s0966-842x(00)88930-1

15 Horodniceanu, T., Bouanchaud, D. H., Bieth, G. & Chabbert, Y. A. R plasmids in

Streptococcus agalactiae (group B). *Antimicrob Agents Chemother* **10**, 795-801 (1976).

https://doi.org:10.1128/aac.10.5.795

16 Leblanc, D. J. & Lee, L. N. Physical and genetic analyses of streptococcal plasmid pAM

beta 1 and cloning of its replication region. *Journal of Bacteriology* **157**, 445-453 (1984).

https://doi.org:10.1128/jb.157.2.445-453.1984

17   Shaw, J. H. & Clewell, D. B. Complete nucleotide sequence of macrolide-lincosamide-

streptogramin B-resistance transposon Tn917 in Streptococcus faecalis. *J Bacteriol* **164**,

782-796 (1985). https://doi.org:10.1128/jb.164.2.782-796.1985

18   Cormier, A. C. *et al.* Diversity of blaCTX-M-1-carrying plasmids recovered from

Escherichia coli isolated from Canadian domestic animals. *PLOS ONE* **17**, e0264439

(2022). https://doi.org:10.1371/journal.pone.0264439

19   Dolejska, M., Villa, L., Hasman, H., Hansen, L. & Carattoli, A. Characterization of IncN

plasmids carrying bla CTX-M-1 and qnr genes in Escherichia coli and Salmonella from

animals, the environment and humans. *J Antimicrob Chemother* **68**, 333-339 (2013).

https://doi.org:10.1093/jac/dks387

20   San Millan, A. *et al.* Positive selection and compensatory adaptation interact to stabilize

non-transmissible plasmids. *Nature Communications* **5**, 5208 (2014).

https://doi.org:10.1038/ncomms6208

21   Enne, V. I. *et al.* Assessment of the fitness impacts on Escherichia coli of acquisition of

antibiotic resistance genes encoded by different types of genetic element. *Journal of*

Antimicrobial Chemotherapy **56**, 544-551 (2005). https://doi.org:10.1093/jac/dki255

22   Loftie-Eaton, W. *et al.* Compensatory mutations improve general permissiveness to

antibiotic resistance plasmids. *Nature Ecology & Evolution* **1**, 1354-1363 (2017).

https://doi.org:10.1038/s41559-017-0243-2

23   Alonso-del Valle, A. *et al.* Variability of plasmid fitness effects contributes to plasmid

persistence in bacterial communities. *Nature Communications* **12**, 2653 (2021).

https://doi.org:10.1038/s41467-021-22849-y

24   Lopatkin, A. J. *et al.* Persistence and reversal of plasmid-mediated antibiotic resistance.

Nature Communications **8** (2017). https://doi.org:10.1038/s41467-017-01532-1

25   Hall, J. P. J., Wood, A. J., Harrison, E. & Brockhurst, M. A. Source–sink plasmid transfer

dynamics maintain gene mobility in soil bacterial communities. *Proceedings of the*

National Academy of Sciences **113**, 8260-8265 (2016).

https://doi.org:10.1073/pnas.1600974113

26   Alonso-Del Valle, A. *et al.* Antimicrobial resistance level and conjugation permissiveness

shape plasmid distribution in clinical enterobacteria. *Proceedings of the National*

Academy of Sciences **120** (2023). https://doi.org:10.1073/pnas.2314135120

27   Wang, T. & You, L. The persistence potential of transferable plasmids. *Nature*

Communications **11** (2020). https://doi.org:10.1038/s41467-020-19368-7

28   Wang, X. *et al.* Inter-plasmid transfer of antibiotic resistance genes accelerates antibiotic

resistance in bacterial pathogens. *Isme j* **18** (2024).

https://doi.org:10.1093/ismejo/wrad032

29   Murray, L. M. *et al.* Co-selection for antibiotic resistance by environmental contaminants.

npj Antimicrobials and Resistance **2**, 9 (2024). https://doi.org:10.1038/s44259-024-

00026-7

30   Cooper, N. S., Brown, M. E. & Caulcott, C. A. A Mathematical Method for Analysing

Plasmid Stability in Micro-organisms. *Microbiology* **133**, 1871-1880 (1987).

https://doi.org:10.1099/00221287-133-7-1871

31   Summers, D. K. The kinetics of plasmid loss. *Trends in Biotechnology* **9**, 273-278

(1991). https://doi.org:https://doi.org/10.1016/0167-7799(91)90089-Z

32   Fedorec, A. J. H. *et al.* Two New Plasmid Post-segregational Killing Mechanisms for the

Implementation of Synthetic Gene Networks in Escherichia coli. *iScience* **14**, 323-334

(2019). https://doi.org:10.1016/j.isci.2019.03.019

33   Danino, T. *et al.* Programmable probiotics for detection of cancer in urine. *Science*

Translational Medicine **7**, 289ra284-289ra284 (2015).

https://doi.org:10.1126/scitranslmed.aaa3519

34   Nordström, K. & Austin, S. J. MECHANISMS THAT CONTRIBUTE TO THE STABLE

SEGREGATION OF PLASMIDS. *Annual Review of Genetics* **23**, 37-69 (1989).

https://doi.org:https://doi.org/10.1146/annurev.ge.23.120189.000345

35   Baxter, J. C. & Funnell, B. E. Plasmid Partition Mechanisms. *Microbiol Spectr* **2** (2014).

https://doi.org:10.1128/microbiolspec.PLAS-0023-2014

36   Cooper, T. F. & Heinemann, J. A. Postsegregational killing does not increase plasmid

stability but acts to mediate the exclusion of competing plasmids. *Proceedings of the*

National Academy of Sciences **97**, 12643-12648 (2000).

https://doi.org:10.1073/pnas.220077897

37   Strogatz, S. a. *Nonlinear dynamics and chaos : with applications to physics, biology,*

chemistry, and engineering. (Second edition. Boulder, CO : Westview Press, a member

of the Perseus Books Group, [2015], 2015).

38   Shao, B. *et al.* Single-cell measurement of plasmid copy number and promoter activity.

Nature Communications **12**, 1475 (2021). https://doi.org:10.1038/s41467-021-21734-y

39   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout

mutants: the Keio collection. *Mol Syst Biol* **2**, 2006.0008 (2006).

https://doi.org:10.1038/msb4100050

40   Wang, T. *et al.* Horizontal gene transfer enables programmable gene stability in

synthetic microbiota. *Nature Chemical Biology* (2022). https://doi.org:10.1038/s41589-

022-01114-3

41   Şimşek, E. *et al.* Keystone engineering enables collective range expansion in microbial

communities. *bioRxiv*, 2025.2001.2011.632568 (2025).

https://doi.org:10.1101/2025.01.11.632568

42   Wu, F. *et al.* Modulation of microbial community dynamics by spatial partitioning. *Nature*

Chemical Biology **18**, 394-402 (2022). https://doi.org:10.1038/s41589-021-00961-w

43 Agustín, M. D., Stengel, P., Kellermeier, M., Tücking, K.-S. & Müller, M. Monitoring

Growth and Removal of Pseudomonas Biofilms on Cellulose-Based Fabrics.

Microorganisms **11** (2023).

44 Stevenson, C., Hall, J. P. J., Harrison, E., Wood, A. J. & Brockhurst, M. A. Gene mobility

promotes the spread of resistance in bacterial populations. *The ISME Journal* **11**, 1930-

1932 (2017). https://doi.org:10.1038/ismej.2017.42

45 Weiss, A., Wang, T. & You, L. Promotion of plasmid maintenance by heterogeneous

partitioning of microbial communities. *Cell Systems* **14**, 895-905.e895 (2023).

https://doi.org:10.1016/j.cels.2023.09.002

46 Stevenson, C., Hall, J. P. J., Brockhurst, M. A. & Harrison, E. Plasmid stability is

enhanced by higher-frequency pulses of positive selection. *Proceedings of the Royal*

Society B: Biological Sciences **285**, 20172497 (2018).

https://doi.org:10.1098/rspb.2017.2497

47 Cai, M. *et al.* Occurrence and temporal variation of antibiotics and antibiotic resistance

genes in hospital inpatient department wastewater: Impacts of daily schedule of

inpatients and wastewater treatment process. *Chemosphere* **292**, 133405 (2022).

https://doi.org:https://doi.org/10.1016/j.chemosphere.2021.133405

48 Cao, S., Zhang, P., Halsall, C., Hou, Z. & Ge, L. Occurrence and seasonal variations of

antibiotic micro-pollutants in the Wei River, China. *Environmental Research* **252**, 118863

(2024). https://doi.org:https://doi.org/10.1016/j.envres.2024.118863

49 Schuster, D. *et al.* Antibiotic concentrations in raw hospital wastewater surpass minimal

selective and minimum inhibitory concentrations of resistant Acinetobacter baylyi strains.

Environ Microbiol **24**, 5721-5733 (2022). https://doi.org:10.1111/1462-2920.16206

50 Coutu, S., Rossi, L., Barry, D. A., Rudaz, S. & Vernaz, N. Temporal Variability of

Antibiotics Fluxes in Wastewater and Contribution from Hospitals. *PLOS ONE* **8**, e53592

(2013). https://doi.org:10.1371/journal.pone.0053592

51   Baker, C. M., Ferrari, M. J. & Shea, K. Beyond dose: Pulsed antibiotic treatment

schedules can maintain individual benefit while reducing resistance. *Scientific Reports* **8**,

5866 (2018). https://doi.org:10.1038/s41598-018-24006-w

52   Letten, A. D., Hall, A. R. & Levine, J. M. Using ecological coexistence theory to

understand antibiotic resistance and microbial competition. *Nature Ecology & Evolution*

5, 431-441 (2021). https://doi.org:10.1038/s41559-020-01385-w

53   Bauer, M., Graf, I. R., Ngampruetikorn, V., Stephens, G. J. & Frey, E. Exploiting ecology

in drug pulse sequences in favour of population reduction. *PLOS Computational Biology*

13, e1005747 (2017). https://doi.org:10.1371/journal.pcbi.1005747

54   Lin, W.-H. & Kussell, E. Complex Interplay of Physiology and Selection in the

Emergence of Antibiotic Resistance. *Current Biology* **26**, 1486-1493 (2016).

https://doi.org:10.1016/j.cub.2016.04.015

55   Nev, O. A., Jepson, A., Beardmore, R. E. & Gudelj, I. Predicting community dynamics of

antibiotic-sensitive and -resistant species in fluctuating environments. *Journal of The*

Royal Society Interface **17**, 20190776 (2020). https://doi.org:10.1098/rsif.2019.0776

56   Buckner, M. M. C., Ciusa, M. L. & Piddock, L. J. V. Strategies to combat antimicrobial

resistance: anti-plasmid and plasmid curing. *FEMS Microbiology Reviews* **42**, 781-804

(2018). https://doi.org:10.1093/femsre/fuy031

Figure 1. High plasmid abundance nonlinearly increases plasmid half-lives.

(a). Plasmid abundance decreases through two processes operating on different time scales, slow segregation loss of the plasmid (pink) and rapid out-competition of plasmid-carrying cells by plasmid-free cells (purple).

(b). Simulated time series of plasmid abundance decay. When starting from high initial abundance P_0% (teal: 100%, orange: 99%), plasmid abundance exhibits biphasic decay, with the slow phase driven by segregation loss (pink arrow) followed a faster phase driven by growth competition (purple arrow) later. When starting from lower P_0% values (dusty blue: 90%, pink: 50%, and lime green: 20%), the plasmid abundance follows exponential decay.

(c). Dependence of plasmid half-life $\tau_{1/2}$ on the initial abundance $P_0\%$. The scatter

points represent simulation results; the curve shows the best fit to Eq. (1) in **Methods**.

(d). Time series of average pSC101 plasmid abundance decay (mean ± SE, n = 3) in *E.*

coli MG1655 across different $P_0\%$ (teal: 100%, orange: 99%, dusty blue: 90%, pink:

50%, and lime green: 20%).

(e - g). Dependence of plasmid half-life $\tau_{1/2}$ (mean ± SE, delta method) on the initial

abundance $P_0\%$ for non-mobilizable plasmids **(e)** pSC101, **(f)** colE1, and **(g)** pUC.

Curves connect the median half-lives at each given initial abundance $P_0\%$. Right-

censored datapoints ($\tau_{1/2}$ > 12 days) are shown without error bar.

(h & i). Dependence of plasmid half-life $\tau_{1/2}$ (mean ± SE, delta method) on the initial

abundance $P_0\%$ for conjugative plasmids **(h)** pCU1 and **(i)** R6K. Curves connect the

median half-lives at each initial abundance $P_0\%$. Right-censored datapoints ($\tau_{1/2}$ > 25

days) are shown without error bar.

Figure 2. A transient antibiotic pulse prolongs plasmid persistence by inducing ghost effect.

(a). In a mixed population containing both plasmid-carrying and plasmid-free cells (purple), an antibiotic purifies the population by eliminating the plasmid-free cells,

moving the population into the *ghost* state (orange), and extending the half-life of

plasmid decay (right).

(b – d). Time series of average plasmid abundance (mean ± SE, n = 3) for non-

mobilizable plasmids **(b)** pSC101, **(c)** colE1, and **(d)** pUC under different selection

strength. Between day 2 and 3 (grey), antibiotics were applied at different

concentrations to induce varying levels of ghost effect. Three representative conditions

are shown here, corresponding to no antibiotic treatment (circle), 1/32 × 50 µg/mL

(triangle), and 50 µg/mL (square). pSC101 and colE1 populations were treated with

Kan, and pUC populations were treated with Spec.

(e – g). Dose-dependence of plasmid half-lives $\tau_{1/2}$ (mean ± SE, delta method) on the

pulsed antibiotic concentration for **(e)** pSC101, **(f)** colE1, and **(g)** pUC. Right-censored

datapoints ($\tau_{1/2}$ > 17 days) are shown without error bar.

Figure 3. Antibiotic pulse induces prolonged plasmid carriage in synthetic *E. coli*

communities.

(a – c). Plasmid abundance (mean ± SE based on technical triplicates of selective

plating) over time for **(a)** R388 in Comm87, **(b)** R6K in Comm60, **(c)** pCU1 in Comm 60.

Purple curves: communities were passaged daily at 1:100 dilution ratio in LB

supplemented with 100 µg/mL Carbenicillin (Comm87, **a**) or 25µg/mL Chloramphenicol

(Comm57, **b & c**). Orange curves: communities were treated with an additional

antibiotic pulse between day 2 and 3 (grey bar) to select for the plasmid-carrying

populations (R388: 10 µg/mL Trimethoprim, R6K: 100 µg/mL Streptomycin, pCU1: 100

µg/mL Carbenicillin).

(d - f). Antibiotic pulses extended plasmid half-lives (mean ± SE, delta method) for **(d)**

R388 in Comm87, **(e)** R6K in Comm57, **(f)** pCU1 in Comm57. Right-censored

datapoints ($\tau_{1/2}$ > 17 days) are shown without error bar. For hypothesis testing, half-

lives for R6K in the pulsed group (+Strp) were taken as $\tau_{1/2}$ = 17 days. **(d)** Average half-

life of R388 after Trim treatment was significantly higher than in LB group: 27.3 ± 1.7

(SD, n = 3) vs 2.2 ± 0.4 (SD, n = 3) days. Welch's t-test: t(2.21) = 24.91, p<0.001, mean

difference = 25.1 [95% CI: 21.1, 29.1], Cohen's d = 20.34. **(e)** Average half-life of R6K

after Trim treatment was significantly higher than in LB group: 17.0 ± 0.0 (SD, n = 3) vs

4.4 ± 0.1 (SD, n = 3) days. Welch's t-test: t(2.00) = 190.56, p<0.001, mean difference =

12.6 [95% CI: 12.4, 12.9], Cohen's d = 155.59. **(f)** Average half-life of pCU1 after Trim

treatment was significantly higher than in LB group: 9.9 ± 0.9 (SD, n = 3) vs 1.0 ± 0.1

(SD, n = 3) days. Welch's t-test: t(2.01) = 16.50, p=0.004, mean difference = 8.8 [95%

CI: 6.5, 11.1], Cohen's d = 13.47. In the figures, **: p < 0.01, ***: p < 0.001.

(g & h). Plasmid abundance (mean ± SE based on technical triplicates of selective

plating) over time for **(g)** R388 and **(h)** RP4 in Comm60. Antibiotic pulsing induced

alternative steady state for plasmid persistence. Purple curves: communities passaged

daily at 1:100 dilution ratio in LB supplemented with 25µg/mL Chloramphenicol. Orange:

communities were treated with additional antibiotic pulse between Day 2 and 3 (grey

bar) to select for the plasmid-carrying populations (R388: 10 µg/mL Trimethoprim, RP4:

10 µg/mL Tetracycline).

(i). Strain-level community dynamics (mean ± SE, n = 3) of Comm87 without (left) or

with (right) 1-day of Trimethoprim treatment. The five most abundant strains (based on

cumulative relative abundance across samples) are color-coded. Strain 1 corresponds

to the donor strain carrying plasmid R388.

(j). Change in the relative abundance of the donor strain (Δ donor) before (day 2) and

after (day 3) the antibiotic pulse. Bars represent mean ± SE (n = 3). Welch's t-test, **: p

< 0.01, ***: p < 0.001.

Figure 4. Ghost effect in niche partitioned communities.

(a & b). Microbes co-exist through partitioning of resource or habitat.

(c). Change in the community composition over time for a mixed population of two

strains (green and purple) with complete niche partitioning (no interaction). Orange

shading indicates the plasmid-carrying fraction of strain 1 (green). The community

experienced selection for the plasmid-carrying cells between day 2 and 3 (gray bar).

(d). Plasmid abundance over time with (orange) and without (purple) 1-day antibiotic

selection (gray bar) for the plasmid. The ghost effect occurs without plasmid abundance

reaching P% = 100%.

(e). Microbial communities containing plasmid-free and plasmid-carrying *E. coli* MG1655

and eight sink isolates were cultured in LB media with sponge-imposed structure.

Genus-level taxonomies of the eight sink isolates are shown. The isolates include one

Pseudomonas strain (*P. aeruginosa*), three *Bacillus* strains, one *Citrobacter* strain, and

three *Enterobacter* strains (including one strain identified as *E. cloacae*). Strains are

color-coded by genus correspondingly as in (**i – k**). Three plasmids (pSC101, colE1,

and pUC) were separately cultured in three independent communities.

(f – h). Time series of GFP/OD for GFP-encoding plasmids pSC101, colE1, and pUC.

Purple: communities were passaged daily at $1:2\times10^7$ dilution ratio in sponges immersed

in LB. Orange: communities were treated with additional antibiotic pulse between day 1

and 2 (grey bar) to select for the plasmid-carrying populations (pSC101 and colE1: 50

µg/mL Kanamycin, pUC: 50 µg/mL Spectinomycin). Green bar: mean ± SE GFP/OD for

E. coli MG1655 carrying the corresponding plasmid (n = 3). Error bars indicate

propagated measurement uncertainties (per-well SD).

(i – k). 16S-resolved community dynamics between day 0 (D0) and day 2 (D2) (n = 1 for

day 0, mean ± SE, n = 3 for other samples). After the antibiotic pulse, the communities

were dominated by *E. coli* and *P. aeruginosa*. Due to limited taxonomic resolution of

16S sequencing, *E. coli* was assigned as *Escherichia-Shigella*, here we denote it as

Escherichia. Taxa with maximum relative abundance < 1% across all samples are not

shown.