1 **Dynamical memory underlies prolonged plasmid persistence after transient antibiotic**

2 **treatment**

3 Zhengqing Zhou[1,2,*], Andrea Weiss[1,2,*], Zhixiang Yao[1,2,6], Xiaoli Chen[1,2], Kristen Lok[1,2], Hye-in

4 Son[1,2], Lingchong You[1,2,3,4,5,+]

6 1 Department of Biomedical Engineering, Duke University, Durham, North Carolina, USA
7
8 2 Center for Quantitative Biodesign, Duke University, Durham, North Carolina, USA
9
10 3 Center for Biomolecular and Tissue Engineering, Duke University, Durham, North Carolina,
11 USA
12
13 4 Center for Genomic and Computational Biology, Duke University, Durham, North Carolina,
14 USA
15
16 5 Department of Molecular Genetics and Microbiology, Duke University School of Medicine,
17 Durham, North Carolina, USA
18
19 6 Present address: Innovative Genomics Institute, University of California, Berkeley, California,
20 USA
21
22

23 *These authors contribute equally to the work.

24 +Correspondence and requests for materials should be addressed to Lingchong You. E-mail:

25 lingchong.you@duke.edu; Tel: 919-660-8408; Fax: 919-668-0795.

Abstract

Plasmids play critical roles in spreading and maintaining antimicrobial resistance (AMR). They often exhibit prolonged persistence upon antibiotic treatment, even when they impose substantial burden on their hosts. This persistence has been primarily attributed to rapid horizontal transfer or low plasmid cost. However, these mechanisms cannot account for the slow decay of non-mobilizable, burdensome plasmids. Here, we show that the decoupling of time scales between slow segregation loss and fast growth competition leads to a slow-down in plasmid abundance decay at high plasmid abundance, reminiscent of the "ghost effect" from nonlinear dynamical systems. Integrating theory, simulations, and quantitative experiments across clonal populations and multi-species bacterial communities, we demonstrate that a transient antibiotic pulse can eliminate plasmid-free cells and create the "ghost" state that extends plasmid persistence from days to months. Our research reveals a generalizable mechanism for the prolonged ecological memory of antibiotic exposure, and underscores the need for proactive strategies to curb the spread of AMR.

Introduction

Plasmids are self-replicating genetic elements that play a critical role in bacterial evolution by enabling the acquisition, maintenance, and dissemination of adaptive traits such as antimicrobial resistance (AMR)[1-3], giving rise to many multidrug-resistant high-risk clones[4,5] and accelerating the global AMR crisis[3]. Although plasmid carriage is often assumed to be burdensome in the absence of selection, leading to rapid plasmid loss[6,7], empirical studies suggest otherwise. Resistance frequency often remains high for months after short-term antibiotic treatments (days to weeks), despite the absence of continued selective pressure thereafter[8-12]. Given that many resistance determinants reside on plasmids[11,13-19], the slow resistance reversal suggests burdensome plasmid can still experience long transients after antibiotic exposure.

Different mechanisms may account for this prolonged persistence after antibiotic treatment. For instance, positive selection can increase the frequency of compensatory mutations that reduce the cost of plasmid carriage and slow down the decay of plasmids[20]. Long half-life of certain antibiotics can exert sustained selection for resistance after the treatment ceased[9]. More broadly, the persistence or slow reversal of resistance or plasmid carriage can be associated with low or zero fitness cost[6,21-23], high conjugation rate[24-26], or both[24,27]. Co-selection by antibiotics[6,28] or environmental contaminants[29] on the same mobilizable gene elements has also been proposed. However, these mechanisms cannot account for the slow decay of plasmids whose baseline abundance is low, i.e., burdensome, slow-transferring plasmids get lost fast.

Past theories suggest another under-appreciated possibility: the slow segregation loss of plasmids[30,31]. While largely ignored due to its relatively minor contribution to plasmid steady state persistence[24,25,27], slow segregation loss can sustain plasmid carriage in clones for a long time [32,33], before plasmid-encoded high-burden cargoes drive the plasmid to get completely lost. This provides a theoretical basis for the prolonged plasmid carriage following antibiotic exposure: antibiotics can eliminate the plasmid-free competitor cells, so the plasmid needs to go through the slow segregation loss process before rapidly decrease in their abundance due to the growth competition from plasmid-free segregant.

Here, we combine theory, numerical simulations, and quantitative experiments to demonstrate this principle in clonal populations, synthetic *E. coli* communities, and multi-species communities of hospital sink isolates: the decoupling of time scales between segregation loss and growth competition can lead to the observed prolonged ecological memory of plasmids. This

principle underscores the need to develop active intervention strategies beyond standard
antibiotic stewardship.

Results

Population ghost effect slows down plasmid abundance decay

We start from a simple ordinary differential equation (ODE) model depicting the dynamics
of a non-mobilizable plasmid within a single strain population (**Methods**). The dynamics of
plasmid loss in a population are mainly dictated by two processes: *(1) segregation loss*, where a
daughter cell fails to inherit the plasmid during cell division (**Fig. 1a**, pink)*, and *(2) growth
competition*, where plasmid-carrying cells are outcompeted by plasmid-free population due to
plasmid-imposed metabolic burden on their host (**Fig. 1a**, purple). Segregation loss is generally
a slow process, where different plasmids can deploy active partitioning, post-segregational killing,
and high copy-number to minimize its loss during cell division[34,35]. Meanwhile, growth competition
operates on a faster timescale.

We used the ODE model to simulate plasmid abundance (fraction of plasmid carrying cells
in a population) over time (**Fig. 1b**), across different initial plasmid abundances of $P_0\% = 100\%$,
99%, 90%, 50%, and 20%. When starting from a pure plasmid-carrying population ($P_0\% = 100\%$),
plasmid abundance decays in two phases, corresponding to the two processes: an initial slow
decay driven by segregation loss, followed by a faster decay driven by growth competition once
plasmid-free cells have emerged. This two-phase behavior is consistent with previous
experimental observations[32,36], where the presence of post-segregational killing mechanism
results in a long plateau in which plasmid abundance remains ~100%. In the absence of active
partitioning or post-segregational killing mechanisms, this plateau was significantly shortened or
even eliminated. As the initial plasmid abundance decreases, growth competition becomes more
prominent, driving the plasmid abundance decay exponentially. Based on a previous work[30], we
can analytically derive the half-life of plasmid abundance decay (**Methods**) which nonlinearly
increases with initial plasmid abundance (**Fig. 1c**). When $P_0\% << 100\%$, the abundance decays
exponentially with half-life $\tau_{1/2} = \frac{1}{\delta}\log 2$. When $P_0\% = 100\%$, plasmid half-life becomes
$\tau_{1/2} \rightarrow \frac{1}{\delta}\log(\frac{\delta}{\kappa})$, where δ is the rate of growth competition between the plasmid-free and plasmid-
carrying populations, and κ is the segregation loss rate. In most cases, $\kappa \ll \delta$ [24], thus half-life
$\tau_{1/2}$ becomes large, and the high abundance state appear as the "ghost" of an equilibrium[37]. This
logarithmic scaling between κ and $\tau_{1/2}$ resembles the "ghost effect" coined by Strogatz[37],

originally referring to the loss of stability in a dynamical system during saddle-node bifurcation.

The prolonged plasmid carriage in our case instead arises from the remnant of a saddle point

(when κ changes from 0 to a small number).

To verify our prediction of diverging half-lives near pure population, we monitored the

abundance of three non-mobilizable plasmids with different origin of replication, pSC101

(kanamycin resistant, Kan^R), colE1 (Kan^R), and pUC (spectinomycin, Spect^R), all encoding sfGFP,

in *E. coli* strain MG1655 over time (**Fig. 1d & Fig. S1**). To facilitate high-throughput measurements,

we used the GFP/OD ratio to quantify plasmid abundance (calibration details in **Methods**).

Plasmid-carrying MG1655 cells were mixed with plasmid-free MG1655 cells at different volume

ratios to generate populations where plasmid carrying cells make up of 100%, 99%, 90%, 50% or

20% of the total population. Populations were cultured in LB and diluted 1:500 every 24 hours

(**Methods**). As predicted by the mathematical model, a plateau in plasmid abundance was

observed at high initial abundances before exponential decay (**Fig. S1b**), significantly prolonging

the half-life of plasmid abundance. Thus, the initial purity of the community, in terms of the

plasmid-carrying cell population, dictates the plasmid half-life (**Fig. 1 e - g**). Interestingly, at P_0%

= 100%, pUC persisted in the community. As a high copy plasmid[38], the stochastic chance of pUC

to have segregation error becomes extremely small, preventing the birth of plasmid-free cells and

the subsequent growth competition. Moreover, for the populations with plasmid abundance decay

(pUC with P_0% = 99% and 90%, and colE1 with P_0% = 99%), the plasmid abundances sometimes

rebounded, suggesting potential compensatory mutations.

We further quantified the dynamics of two conjugative plasmids, pCU1 and R6K in *E. coli*

DA28102, with daily 100-fold dilution in LB media supplemented with 25 µg/mL chloramphenicol

(Cm) to prevent contamination. Both pCU1 and R6K exhibited slowed-down decays and diverging

half-lives when P_0% is near 100% (**Fig. 1h & i, Fig. S2**).

Transient antibiotic selection extends plasmid half-life through purifying selection

Antibiotics can eliminate plasmid-free cells in a population and purify the population to

enter the ghost effect region (**Fig. 2a**). We thus hypothesize selection with antibiotics could

prolong the carriage of a plasmid. We assembled populations of *E. coli* MG1655 with an equal

mixture of plasmid-free and plasmid-carrying cells (P_0%=50%). We used the non-mobilizable,

sfGFP-encoding plasmids pSC101, colE1, and pUC for high-throughput measurement. We

applied a pulse of antibiotic selection to the populations between day 2 and 3. After growing the

population in LB supplemented with a gradient of antibiotics (Kan for pSC101 and colE1, and
Spect for pUC) for 24 hours, we removed the selection and tracked the GFP/OD readout over
time (**Fig. 2b – d, Fig. S3b**). We found the half-lives of the three plasmids all positively correlate
with the dosage of antibiotic selection (**Fig. 2e - g**), and plateaus above critical concentrations.
The critical concentrations are 3.125 µg/mL Kan for pSC101, 6.25 µg/mL Kan for colE1, and 12.5
µg/mL Spect for pUC.
147
Antibiotic pulse induces prolonged plasmid carriage in synthetic *E. coli* communities
To determine whether the ghost effect occurs in more complex microbial systems, we
constructed five synthetic communities of *E. coli* Keio strains[39]. Each strain contains a uniquely
barcoded, non-mobilizable plasmid, enabling the quantification of community structure using next
generation sequencing (NGS). The first community, Comm87, consisted of 86 background strains
plus one donor strain carrying conjugative plasmid R388 (trimethoprim resistant, Trim^R). Four
other communities (Comm57) shared the same background community of 56 barcoded Keio
strains, in addition to a unique donor strain carrying one additional conjugative plasmid, either
R6K (streptomycin resistant, Strp^R), pCU1 (carbenicillin resistant, Carb^R), R388, and RP4
(tetracycline resistant, Tet^R). The overnight cultures of the background strains were equally
combined, then mixed with the donor strain at a 3:1 ratio to assemble each of the five communities.
The communities were passaged daily with a 100-fold dilution in LB media supplemented with
100 µg/mL Carb for Comm 87 or 25 µg/mL Cm for Comm 57 to maintain the barcode plasmids.
We tracked the abundance of the target plasmids over time through selective plating.
162
Without any additional antibiotic intervention, the conjugative plasmids exponentially
decayed (**Fig. 3a - c**, purple), with half-lives of 2.2 (±SD = 0.4, n=3) days for R388 in Comm87,
4.4 (±0.1) days for R6K in Comm57, and 1.0 (±0.1) days for pCU1 in Comm57 (**Fig. 3d - f**). R388
and RP4 were able to persist in Comm57 for the entire course of the 20-day xperiment (**Fig. 3g**
& h). As predicted by the model, antibiotic selection for the plasmids, applied between day 2 and
3, increased plasmid abundances to ~ 100%. Driving the communities into the ghost effect region
significantly extended plasmid half-lives to 27.3 (±1.7) days for R388 in Comm87 and 9.9 (±0.9)
days for pCU1 in Comm57 (**Fig. 3 d-f,** p<0.01 or p<0.001 with Welch's t-test). Plasmids R6K,
R388, and RP4 all persisted in the communities at ~100% for more than 17 days after antibiotic
selection (**Fig. 3b, g, h**, orange). The four Comm57 communities, when passaged in a different
dilution rate (1:500), showed similar results (**Fig. S4**).
174

We used NGS to quantify the community composition over time (**Fig. 3i & Fig. S5**). Upon
antibiotic selection, a significant shift in each community structure was observed, and the
community diversity dropped significantly (**Fig. S6**). Upon selection, the relative abundances of
the plasmid donors were significantly amplified (**Fig. 3j**), while the donor abundances decreased
in most communities without antibiotic pulse. The decline in diversity, and the increase in donor
abundance, suggests that the plasmids have not yet spread to every member in the community.
One exception was Comm57 harboring RP4, where the donor abundance was significantly
decreased upon antibiotic selection (**Fig. 3j**). While in other communities the antibiotic pulse
resulted in a crash in diversity, the RP4 community instead increased its diversity (**Fig. S6**). This
could be due to the combination of two factors: the low fitness of the donor strain, and the fast
conjugation capacity of RP4[24,40]. By the time positive selection takes place, most strains in the
community would have already received the plasmid, and could outcompete the donor strain in
the presence of Tet.
188
189   These findings demonstrate that the ghost effect operates not only in clonal populations
190   but also in synthetic microbial communities. A brief antibiotic pulse can reorganize community
191   composition by selectively eliminating plasmid-free competitors, thereby stabilizing plasmid
192   persistence at the community level without requiring genetic adaptation.
193

Ghost effect in multi-specific communities with niche partition

In natural ecosystems, microbes can co-exist in natural communities through niche
partition, where different species / strains can take up different resources (**Fig. 4a**) or habitats
(**Fig. 4b**) to avoid competitive exclusion. In the above experiments, we showed the ghost effect
taking place in clonal populations and *E. coli* communities, where strong competitions exist
between individual strains, and thus plasmid-free and plasmid-carrying populations. In these
cases, ghost effect takes place when the plasmid abundance reaches ~100%. However,
numerical simulations (**Methods**) suggest antibiotic-induced ghost effect can happen without
community purity (**Fig. 4 c & d**) with niche partition. For instance, consider a mixed population of
plasmid-free strain 1, plasmid-carrying strain 1, and non-host strain 2 (**Methods**). In the simple
case where (1) strain 1 and 2 do not interact and (2) strain 2 has moderate resistance to the
antibiotic, the non-host strain 2 can survive the antibiotic selection, limiting the plasmid abundance
to < 100%. However, ghost effect can still happen within the purified strain 1 population, resulting
in the prolonged carriage of the plasmid in the two-member community (**Fig. 4c & d**). In essence,
as long as the plasmid-carrying population is pure within a given niche, ghost effect should take

place. Since natural microbiomes often have niche partition, with different species exhibiting

natural or acquired resistance, we expect general applicability of ghost effect following antibiotic

treatments, even when the plasmid or resistance abundance does not reach 100%.

To demonstrate this idea, we constructed three communities consisting of 8 isolates from

Duke Hospital sink p-traps[41] (**Methods**), plasmid-free and plasmid-carrying *E. coli* MG1655 with

one of the three non-mobilizable plasmids (pSC101, colE1 and pUC) expressing sfGFP as a

surrogate for their abundance. The sink isolates consist of one *Pseudomonas Aeruginosa* strain,

three *Bacillus* trains, one *Citrobacter* strain, and three *Enterobacter* strains (**Fig. 4e**). To facilitate

niche partitioning, we used cellulose sponges immersed in LB media to create porous, physically

partitioned habitats (**Fig. 4e**), which were previously found to promote the diversity of microbial

communities[42].

In the absence of antibiotic selection, the GFP/OD level (**Methods**) dropped rapidly to

below 10^2. After transient Kan or Spect selection between day 1 and 2, GFP expression was

increased. pSC101 showed slowed decay in GFP/OD level, while colE1 and pUC persisted until

day 10 (**Fig. 4 f – h**). This is consistent with our observations of the three plasmids in clonal

populations (**Fig. 1, Fig. S1**). Importantly, for plasmids colE1 and pUC, the GFP/OD levels were

consistently below the baseline in pure plasmid-carrying MG1655 populations.

Through 16S sequencing (**Methods**), we quantified community dynamics in the presence

and absence of the transient antibiotic treatment (**Fig. 4 i – k**). On day 2, in the absence of

antibiotics, *P. aeruginosa* and *Enterobacter* became dominant in the communities, with the other

three genera decreasing in their fractions or even become undetectable in some of the replicates.

Antibiotic treatment increased the abundance of the two resistant strains, *E. coli* and *P.*

aeruginosa, and eliminated the rest of the community. The survival of *P. aeruginosa* confirms that

the communities do not purely contain plasmid-carrying cells, but a ghost effect within the *E. coli*

population still takes place. Consistent with the GFP/OD reading of pSC101, *P. aeruginosa* was

detectable in one of the three replicate communities harboring pSC101, potentially due to the

stochastic effect of high dilution rate (2×10^7) during daily passaging.

In LB cultures without cellulose sponges, the GFP/OD levels of the communities after

antibiotic treatment became comparable to the pure plasmid-carrying baselines (Fig. S7) and

exhibited slowed-down decay in comparison to the antibiotic-free control group. Since the

plasmids are non-mobilizable, the post-treatment communities likely consisted primarily of *E. coli*.

In contrast, the survival of *P. aeruginosa* should be attributed to the presence of sponges, which

possibly introduces spatial structures to mediate the competition[42] and promote biofilm formation[43].

Discussion

Resistance limits the clinical efficacy of antibiotic treatments and poses a major threat to human health. Many antibiotic resistance genes are maintained and propagated by mobile genetic elements including plasmids[1-3]. Understanding and predicting the persistence of plasmids is critical for designing intervention strategies to contain and reverse the spread of AMR. In this work, we systematically quantified the decay dynamics of multiple plasmids across different community contexts, and established their dependence on community structure. We showed that the plasmid half-life diverges in nearly pure plasmid-carrying populations, either from assembly or induced by transient antibiotic pulse, extending the plasmid's survival for months. Theoretically, this ghost effect arises from the decrease in the strength of growth competition when plasmid-free cells are scarce in the population.

Our results reveal a common mechanism underlying slow reversal of plasmid-mediated resistance after antibiotic treatments, a phenomenon that spans across microbial ecosystems ranging from the human oral, urinary, and gut microbiomes, to those of livestock. Macrolide antibiotics have been reported to increase the abundance of resistant streptococci (primarily carrying *mef* gene or macrolide-lincosamide-streptogramin-B resistant cassette *erm*(B)) in oral microbiomes to more than 80%, which requiring over 6 months to revert to baseline[9]. The authors attributed this prolonged resistance carriage to the long half-life of azithromycin, and the low burden of the *erm*(B) gene. Similarly, veterinary β-lactams strongly amplified the abundance of resistant bacteria in the pig intestinal microbiome, persisting for more than 3 weeks beyond the drug withdrawal. The extended-spectrum β-lactamase CTX-M-1 was carried by both an exogenous conjugative plasmid and indigenous gut microbiota. Consistent with our results (**Fig. 3, Fig. S5**), the authors observed the significant amplification of resistance abundance and donor abundance upon antibiotic selection, followed by rapid decline of the donor strain abundance after antibiotic withdrawal. Importantly, both the *erm*(B) cassette [13-17] and the CTX-M-1 gene[18,19] are commonly found on conjugative plasmids and other mobilizable gene elements. Yelin et al. reported that resistance in urinary tract infections shows autocorrelation lasting more than six months, with strong and persistent correlation between patients' antibiotic purchase history and subsequent resistance[12]. Anthony et al. observed increases in the abundance of β-lactam, macrolide, and tetracycline resistance following treatment with cephalosporin (β-lactam), azithromycin (macrolide), or both[10]. During the development of the infant gut microbiome, resistance genes on mobile genetic elements were found to persist longer than those encoded on chromosomes[11]. The diversity of antibiotics, resistance genes, bacterial host, and ecosystems

all point to a general population-dynamical mechanism. As the ghost effect depends only on the

slow time scale of segregation loss, it provides a generalizable theoretical framework for these

observed phenomena. Consequently, persistent resistance has important implications for clinical

practice and food production, as it provides a reservoir for the community spread of AMR.

While our work focuses on the population dynamics of the plasmids, the ghost effect also

has strong evolutionary consequences. Prolonged plasmid carriage enables co-evolution of the

plasmid and host, creating opportunities for compensatory mutations. Long-term co-evolution

could ameliorate the burden of the plasmid[20], and may even turn the deleterious cost of plasmid

carriage to a beneficial one[22]. Thus, despite predictions that certain plasmids should not persist

based on ecological models[24,27], their long-term persistence may occur before eventual extinction.

In our experiments, we observed potential evolutionary rescues for colE1 and pUC starting at high

initial plasmid abundances (**Fig. S1**), which was not observed in populations outside of the ghost

effect region. Therefore, plasmid persistence can have bistable outcomes when evolution forces

are taken into consideration.

The antibiotic-induced persistence also provides an environment-driven explanation for

the survival of burdensome plasmids, addressing the so-called plasmid paradox[7]. Previous

studies have attributed plasmid survival to biotic factors, such as fast conjugation[24,25,44], variable

fitness effects across different hosts[23], spatial partitioning[45], and compensatory mutations that

ameliorates the plasmid burden[20,22]. These works typically assume constant environments,

without positive selections on the plasmids. However, plasmids in natural and clinical settings

experience fluctuating selection forces[46-50]. In such cases, the burdensome plasmids can persist

when the frequency of positive selection is on the same order of magnitude as the plasmid half-

life within the ghost effect region. This mechanism further raises questions about the efficacy of

simply reducing antibiotic use to contain AMR[6] – similar persistence can be achieved with far less

amount of antibiotics with sporadic selection. Conversely, pulsed antibiotic treatments has been

proposed as a strategy to constrain the development of resistance[51-55]. Such strategies should

avoid the ghost effect region in design to prevent slow population responses that disproportionally

favor resistant bacteria.

The clinical, ecological, and evolutionary implications of ghost effect underscore the need

for strategies that counteract against unintended antibiotic exposure. Previous studies have used

chemicals to reduce plasmid abundance and reverse its persistence[24,56]. As both plasmid

abundance and half-life are governed by conjugation, growth, and segregation events[27], we
hypothesize that these chemicals could also accelerate plasmid decay. We found that curing
chemicals and their combinations were able to reduce the half-life of pSC101 almost by half (**Fig.**
S8). A single day of treatment using chemical combinations achieved comparable effects as full
course, suggesting these chemicals may accelerate the plasmid decay by increasing segregation
error.
321
Overall, our work established the timescale decoupling between segregation loss and
growth competition for burdensome plasmids, and demonstrated the disproportionate impact of
transient antibiotic pulses on the plasmid persistence. Our findings provide a generalizable
mechanism for the prolonged ecological memory of plasmid carriage following antibiotic exposure,
and have implications for clinical management of antibiotic use and the ecology and evolution of
plasmids.
328

Acknowledgments

We thank Emma Chory, Teng Wang, Rohan Maddemsetti, Emrah Şimşek, Grayson Hamrick, Yifan Dai, and Aaron Yip for helpful discussions. This work was supported by National Institute of Health (L.Y.: R01AI125604, R01GM098642, R01EB031869) and the National Science Foundation (L. Y.: Cooperative Agreement No. EEC-2133504)

Author contributions

Z.Z., A.W., and L.Y. conceived the research. Z.Z., A.W., Z.Y., X.C., and K.L. performed the experiments. Z.Z. performed the modelling and statistical analysis. Z.Z., A.W., and L.Y. wrote the manuscript. All authors read and contributed to the manuscript.

Declaration of interests

The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT to improve the language of the manuscript. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Resource availability

Materials availability

Genetic constructs generated for this study are available upon request.

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Lingchong You (you@duke.edu).

Data and code availability

All original code and data have been deposited at https://github.com/youlab/Plasmid-Ghost-Effect.

References

Bethke, J. H. *et al.* Environmental and genetic determinants of plasmid mobility in pathogenic <i>Escherichia coli</i>. *Science Advances* **6**, eaax3173 (2020). https://doi.org:10.1126/sciadv.aax3173

Lipworth, S. *et al.* The plasmidome associated with Gram-negative bloodstream infections: A large-scale observational study using complete plasmid assemblies. *Nature Communications* **15** (2024). https://doi.org:10.1038/s41467-024-45761-7

Castañeda-Barba, S., Top, E. M. & Stalder, T. Plasmids, a molecular cornerstone of antimicrobial resistance in the One Health era. *Nature Reviews Microbiology* **22**, 18-32 (2024). https://doi.org:10.1038/s41579-023-00926-x

Hawkey, P. M. & Jones, A. M. The changing epidemiology of resistance. *J Antimicrob Chemother* **64 Suppl 1**, i3-10 (2009). https://doi.org:10.1093/jac/dkp256

Mathers, A. J., Peirano, G. & Pitout, J. D. The role of epidemic resistance plasmids and international high-risk clones in the spread of multidrug-resistant Enterobacteriaceae. *Clin Microbiol Rev* **28**, 565-591 (2015). https://doi.org:10.1128/cmr.00116-14

Andersson, D. I. & Hughes, D. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nature Reviews Microbiology* **8**, 260-271 (2010). https://doi.org:10.1038/nrmicro2319

Brockhurst, M. A. & Harrison, E. Ecological and evolutionary solutions to the plasmid paradox. *Trends in Microbiology* **30**, 534-543 (2022). https://doi.org:10.1016/j.tim.2021.11.001

Cavaco, L. M., Abatih, E., Aarestrup, F. M. & Guardabassi, L. Selection and persistence of CTX-M-producing Escherichia coli in the intestinal flora of pigs treated with amoxicillin, ceftiofur, or cefquinome. *Antimicrob Agents Chemother* **52**, 3612-3616 (2008). https://doi.org:10.1128/aac.00354-08

Malhotra-Kumar, S., Lammens, C., Coenen, S., Van Herck, K. & Goossens, H. Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a randomised, double-blind, placebo-controlled study. *The Lancet* **369**, 482-490 (2007). https://doi.org:10.1016/S0140-6736(07)60235-9

Anthony, W. E. *et al.* Acute and persistent effects of commonly used antibiotics on the gut microbiome and resistome in healthy adults. *Cell Reports* **39**, 110649 (2022). https://doi.org:10.1016/j.celrep.2022.110649

Yassour, M. *et al.* Natural history of the infant gut microbiome and impact of antibiotic treatment on bacterial strain diversity and stability. *Sci Transl Med* **8**, 343ra381 (2016). https://doi.org:10.1126/scitranslmed.aad0917

Yelin, I. *et al.* Personal clinical history predicts antibiotic resistance of urinary tract infections. *Nature Medicine* **25**, 1143-1152 (2019). https://doi.org:10.1038/s41591-019-0503-6

Berryman, D. I. & Rood, J. I. The closely related ermB-ermAM genes from Clostridium perfringens, Enterococcus faecalis (pAM beta 1), and Streptococcus agalactiae (pIP501) are flanked by variants of a directly repeated sequence. *Antimicrob Agents Chemother* **39**, 1830-1834 (1995). https://doi.org:10.1128/aac.39.8.1830

Clewell, D. B., Flannagan, S. E. & Jaworski, D. D. Unconstrained bacterial promiscuity: the Tn916-Tn1545 family of conjugative transposons. *Trends Microbiol* **3**, 229-236 (1995). https://doi.org:10.1016/s0966-842x(00)88930-1

Horodniceanu, T., Bouanchaud, D. H., Bieth, G. & Chabbert, Y. A. R plasmids in Streptococcus agalactiae (group B). *Antimicrob Agents Chemother* **10**, 795-801 (1976). https://doi.org:10.1128/aac.10.5.795

16 Leblanc, D. J. & Lee, L. N. Physical and genetic analyses of streptococcal plasmid pAM beta 1 and
cloning of its replication region. *Journal of Bacteriology* **157**, 445-453 (1984).
https://doi.org:10.1128/jb.157.2.445-453.1984

17 Shaw, J. H. & Clewell, D. B. Complete nucleotide sequence of macrolide-lincosamide-
streptogramin B-resistance transposon Tn917 in Streptococcus faecalis. *J Bacteriol* **164**, 782-796
(1985). https://doi.org:10.1128/jb.164.2.782-796.1985

18 Cormier, A. C. *et al.* Diversity of blaCTX-M-1-carrying plasmids recovered from Escherichia coli
isolated from Canadian domestic animals. *PLOS ONE* **17**, e0264439 (2022).
https://doi.org:10.1371/journal.pone.0264439

19 Dolejska, M., Villa, L., Hasman, H., Hansen, L. & Carattoli, A. Characterization of IncN plasmids
carrying bla CTX-M-1 and qnr genes in Escherichia coli and Salmonella from animals, the
environment and humans. *J Antimicrob Chemother* **68**, 333-339 (2013).
https://doi.org:10.1093/jac/dks387

20 San Millan, A. *et al.* Positive selection and compensatory adaptation interact to stabilize non-
transmissible plasmids. *Nature Communications* **5**, 5208 (2014).
https://doi.org:10.1038/ncomms6208

21 Enne, V. I. *et al.* Assessment of the fitness impacts on Escherichia coli of acquisition of antibiotic
resistance genes encoded by different types of genetic element. *Journal of Antimicrobial*
Chemotherapy **56**, 544-551 (2005). https://doi.org:10.1093/jac/dki255

22 Loftie-Eaton, W. *et al.* Compensatory mutations improve general permissiveness to antibiotic
resistance plasmids. *Nature Ecology & Evolution* **1**, 1354-1363 (2017).
https://doi.org:10.1038/s41559-017-0243-2

23 Alonso-del Valle, A. *et al.* Variability of plasmid fitness effects contributes to plasmid persistence
in bacterial communities. *Nature Communications* **12**, 2653 (2021).
https://doi.org:10.1038/s41467-021-22849-y

24 Lopatkin, A. J. *et al.* Persistence and reversal of plasmid-mediated antibiotic resistance. *Nature*
Communications **8** (2017). https://doi.org:10.1038/s41467-017-01532-1

25 Hall, J. P. J., Wood, A. J., Harrison, E. & Brockhurst, M. A. Source–sink plasmid transfer dynamics
maintain gene mobility in soil bacterial communities. *Proceedings of the National Academy of*
Sciences **113**, 8260-8265 (2016). https://doi.org:10.1073/pnas.1600974113

26 Alonso-Del Valle, A. *et al.* Antimicrobial resistance level and conjugation permissiveness shape
plasmid distribution in clinical enterobacteria. *Proceedings of the National Academy of Sciences*
120 (2023). https://doi.org:10.1073/pnas.2314135120

27 Wang, T. & You, L. The persistence potential of transferable plasmids. *Nature Communications*
11 (2020). https://doi.org:10.1038/s41467-020-19368-7

28 Wang, X. *et al.* Inter-plasmid transfer of antibiotic resistance genes accelerates antibiotic
resistance in bacterial pathogens. *Isme j* **18** (2024). https://doi.org:10.1093/ismejo/wrad032

29 Murray, L. M. *et al.* Co-selection for antibiotic resistance by environmental contaminants. *npj*
Antimicrobials and Resistance **2**, 9 (2024). https://doi.org:10.1038/s44259-024-00026-7

30 Cooper, N. S., Brown, M. E. & Caulcott, C. A. A Mathematical Method for Analysing Plasmid
Stability in Micro-organisms. *Microbiology* **133**, 1871-1880 (1987).
https://doi.org:10.1099/00221287-133-7-1871

31 Summers, D. K. The kinetics of plasmid loss. *Trends in Biotechnology* **9**, 273-278 (1991).
https://doi.org:https://doi.org/10.1016/0167-7799(91)90089-Z

32 Fedorec, A. J. H. *et al.* Two New Plasmid Post-segregational Killing Mechanisms for the
Implementation of Synthetic Gene Networks in Escherichia coli. *iScience* **14**, 323-334 (2019).
https://doi.org:10.1016/j.isci.2019.03.019

33   Danino, T. *et al.* Programmable probiotics for detection of cancer in urine. *Science Translational*
Medicine **7**, 289ra284-289ra284 (2015). https://doi.org:10.1126/scitranslmed.aaa3519
34   Nordström, K. & Austin, S. J. MECHANISMS THAT CONTRIBUTE TO THE STABLE SEGREGATION OF
PLASMIDS. *Annual Review of Genetics* **23**, 37-69 (1989).
https://doi.org:https://doi.org/10.1146/annurev.ge.23.120189.000345
35   Baxter, J. C. & Funnell, B. E. Plasmid Partition Mechanisms. *Microbiol Spectr* **2** (2014).
https://doi.org:10.1128/microbiolspec.PLAS-0023-2014
36   Cooper, T. F. & Heinemann, J. A. Postsegregational killing does not increase plasmid stability but
acts to mediate the exclusion of competing plasmids. *Proceedings of the National Academy of*
Sciences **97**, 12643-12648 (2000). https://doi.org:10.1073/pnas.220077897
37   Strogatz, S. a. *Nonlinear dynamics and chaos : with applications to physics, biology, chemistry,*
and engineering. (Second edition. Boulder, CO : Westview Press, a member of the Perseus
Books Group, [2015], 2015).
38   Shao, B. *et al.* Single-cell measurement of plasmid copy number and promoter activity. *Nature*
Communications **12**, 1475 (2021). https://doi.org:10.1038/s41467-021-21734-y
39   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the
Keio collection. *Mol Syst Biol* **2**, 2006.0008 (2006). https://doi.org:10.1038/msb4100050
40   Wang, T. *et al.* Horizontal gene transfer enables programmable gene stability in synthetic
microbiota. *Nature Chemical Biology* (2022). https://doi.org:10.1038/s41589-022-01114-3
41   Şimşek, E. *et al.* Keystone engineering enables collective range expansion in microbial
communities. *bioRxiv*, 2025.2001.2011.632568 (2025).
https://doi.org:10.1101/2025.01.11.632568
42   Wu, F. *et al.* Modulation of microbial community dynamics by spatial partitioning. *Nature*
Chemical Biology **18**, 394-402 (2022). https://doi.org:10.1038/s41589-021-00961-w
43   Agustín, M. D., Stengel, P., Kellermeier, M., Tücking, K.-S. & Müller, M. Monitoring Growth and
Removal of Pseudomonas Biofilms on Cellulose-Based Fabrics. *Microorganisms* **11** (2023).
44   Stevenson, C., Hall, J. P. J., Harrison, E., Wood, A. J. & Brockhurst, M. A. Gene mobility promotes
the spread of resistance in bacterial populations. *The ISME Journal* **11**, 1930-1932 (2017).
https://doi.org:10.1038/ismej.2017.42
45   Weiss, A., Wang, T. & You, L. Promotion of plasmid maintenance by heterogeneous partitioning
of microbial communities. *Cell Systems* **14**, 895-905.e895 (2023).
https://doi.org:10.1016/j.cels.2023.09.002
46   Stevenson, C., Hall, J. P. J., Brockhurst, M. A. & Harrison, E. Plasmid stability is enhanced by
higher-frequency pulses of positive selection. *Proceedings of the Royal Society B: Biological*
Sciences **285**, 20172497 (2018). https://doi.org:10.1098/rspb.2017.2497
47   Cai, M. *et al.* Occurrence and temporal variation of antibiotics and antibiotic resistance genes in
hospital inpatient department wastewater: Impacts of daily schedule of inpatients and
wastewater treatment process. *Chemosphere* **292**, 133405 (2022).
https://doi.org:https://doi.org/10.1016/j.chemosphere.2021.133405
48   Cao, S., Zhang, P., Halsall, C., Hou, Z. & Ge, L. Occurrence and seasonal variations of antibiotic
micro-pollutants in the Wei River, China. *Environmental Research* **252**, 118863 (2024).
https://doi.org:https://doi.org/10.1016/j.envres.2024.118863
49   Schuster, D. *et al.* Antibiotic concentrations in raw hospital wastewater surpass minimal
selective and minimum inhibitory concentrations of resistant Acinetobacter baylyi strains.
Environ Microbiol **24**, 5721-5733 (2022). https://doi.org:10.1111/1462-2920.16206
50   Coutu, S., Rossi, L., Barry, D. A., Rudaz, S. & Vernaz, N. Temporal Variability of Antibiotics Fluxes
in Wastewater and Contribution from Hospitals. *PLOS ONE* **8**, e53592 (2013).
https://doi.org:10.1371/journal.pone.0053592

51 Baker, C. M., Ferrari, M. J. & Shea, K. Beyond dose: Pulsed antibiotic treatment schedules can
maintain individual benefit while reducing resistance. *Scientific Reports* **8**, 5866 (2018).
https://doi.org:10.1038/s41598-018-24006-w
52 Letten, A. D., Hall, A. R. & Levine, J. M. Using ecological coexistence theory to understand
antibiotic resistance and microbial competition. *Nature Ecology & Evolution* **5**, 431-441 (2021).
https://doi.org:10.1038/s41559-020-01385-w
53 Bauer, M., Graf, I. R., Ngampruetikorn, V., Stephens, G. J. & Frey, E. Exploiting ecology in drug
pulse sequences in favour of population reduction. *PLOS Computational Biology* **13**, e1005747
(2017). https://doi.org:10.1371/journal.pcbi.1005747
54 Lin, W.-H. & Kussell, E. Complex Interplay of Physiology and Selection in the Emergence of
Antibiotic Resistance. *Current Biology* **26**, 1486-1493 (2016).
https://doi.org:10.1016/j.cub.2016.04.015
55 Nev, O. A., Jepson, A., Beardmore, R. E. & Gudelj, I. Predicting community dynamics of
antibiotic-sensitive and -resistant species in fluctuating environments. *Journal of The Royal*
Society Interface **17**, 20190776 (2020). https://doi.org:10.1098/rsif.2019.0776
56 Buckner, M. M. C., Ciusa, M. L. & Piddock, L. J. V. Strategies to combat antimicrobial resistance:
anti-plasmid and plasmid curing. *FEMS Microbiology Reviews* **42**, 781-804 (2018).
https://doi.org:10.1093/femsre/fuy031
520
521

Figure 1. High plasmid abundance nonlinearly increases plasmid half-lives.

(a). Plasmid abundance decreases through two processes operating on different time scales, slow segregation loss of the plasmid (pink) and rapid out-competition of plasmid-carrying cells by plasmid-free cells (purple).

(b). Simulated time series of plasmid abundance decay. When starting from high initial abundance $P_0\%$ (teal: 100%, orange: 99%), plasmid abundance exhibits biphasic decay, with the slow phase driven by segregation loss (pink arrow) followed a faster phase driven by growth competition (purple arrow) later. When starting from lower $P_0\%$ values (dusty blue: 90%, pink: 50%, and lime green: 20%), the plasmid abundance follows exponential decay.

(c). Dependence of plasmid half-life $\tau_{1/2}$ on the initial abundance $P_0\%$. The scatter points represent simulation results; the curve shows the best fit to Eq. (1) in **Methods**.

(d). Time series of pSC101 plasmid abundance decay (mean ± SE, n = 3) in *E. coli*

MG1655 across different P_0% (teal: 100%, orange: 99%, dusty blue: 90%, pink: 50%,

and lime green: 20%).

(e - g). Dependence of plasmid half-life $\tau_{1/2}$ on the initial abundance P_0% for non-

mobilizable plasmids **(e)** pSC101, **(f)** colE1, and **(g)** pUC. Curves connect the median

half-lives at each given initial abundance P_0%. Error bars indicate propagated

measurement uncertainties (per-well SD, delta-method in log space). Right-censored

datapoints ($\tau_{1/2}$ > 12 days) are shown without error bar.

(h & i). Dependence of plasmid half-life $\tau_{1/2}$ (mean ± SE based on technical triplicates

of selective plating) on the initial abundance P_0% for conjugative plasmids **(h)** pCU1 and

(i) R6K. Curves connect the median half-lives at each initial abundance P_0%. Right-

censored datapoints ($\tau_{1/2}$ > 25 days) are shown without error bar.

Figure 2. A transient antibiotic pulse prolongs plasmid persistence by inducing ghost effect.

(a). In a mixed population containing both plasmid-carrying and plasmid-free cells (purple), an antibiotic purifies the population by eliminating the plasmid-free cells, moving the population into the ghost effect region (orange), and extending the half-life of plasmid decay (right).

(b – d). Time series of plasmid abundance (mean ± SE, n = 3) for non-mobilizable plasmids **(b)** pSC101, **(c)** colE1, and **(d)** pUC. Between day 2 and 3 (grey), antibiotics were applied at different concentrations to induce varying levels of ghost effect. Three

representative conditions are shown here, corresponding to no antibiotic treatment

(circle), 1/32 × 50 μg/mL (triangle), and 50 μg/mL (square). pSC101 and colE1

populations were treated with Kan, and pUC populations were treated with Spect.

(e – g). Dose-dependence of plasmid half-lives $\tau_{1/2}$ on the pulsed antibiotic

concentration for **(e)** pSC101, **(f)** colE1, and **(g)** pUC. Error bars indicate propagated

measurement uncertainties (per-well SD, delta-method in log space). Right-censored

datapoints ($\tau_{1/2}$ > 17 days) are shown without error bar.

Figure 3. Antibiotic pulse induces prolonged plasmid carriage in synthetic *E. coli*

communities.

(a – c). Plasmid abundance (mean ± SE based on technical triplicates of selective

plating) over time for **(a)** R388 in Comm87, **(b)** R6K in Comm60, **(c)** pCU1 in Comm 60.

Purple curves: communities were passaged daily at 1:100 dilution ratio in LB

supplemented with 100 µg/mL Carbenicillin (Comm87, **a**) or 25µg/mL Chloramphenicol

(Comm60, **b & c**). Orange curves: communities were treated with an additional

antibiotic pulse between day 2 and 3 (grey bar) to select for the plasmid-carrying

populations (R388: 10 µg/mL Trimethoprim, R6K: 100 µg/mL Streptomycin, pCU1: 100

µg/mL Carbenicillin,).

(d - f). Antibiotic pulses extended plasmid half-lives (mean ± SE) for **(d)** R388 in

Comm87, **(e)** R6K in Comm60, **(f)** pCU1 in Comm 60. Right-censored datapoints ($\tau_{1/2}$ >

17 days) are shown without error bar. For hypothesis testing, half-lives for R6K in the

pulsed group (+Strp) were taken as $\tau_{1/2}$ = 17 days. **(d)** R388 half-life after Trim

treatment was significantly higher than in LB group: 27.3 ± 1.7 (SD) vs 2.2 ± 0.4 (SD)

days. Welch's t-test: t(2.21) = 24.91, p<0.001, mean difference = 25.1 [95% CI: 21.1,

29.1], Cohen's d = 20.34. **(e)** R6K half-life after Trim treatment was significantly higher

than in LB group: 17.0 ± 0.0 (SD) vs 4.4 ± 0.1 (SD) days. Welch's t-test: t(2.00) =

190.56, p<0.001, mean difference = 12.6 [95% CI: 12.4, 12.9], Cohen's d = 155.59. **(f)**

pCU1 half-life after Trim treatment was significantly higher than in LB group: 9.9 ± 0.9

(SD) vs 1.0 ± 0.1 (SD) days. Welch's t-test: t(2.01) = 16.50, p=0.004, mean difference =

8.8 [95% CI: 6.5, 11.1], Cohen's d = 13.47. In the figures, **: p < 0.01, ***: p < 0.001.

(g & h). Plasmid abundance (mean ± SE) over time for **(g)** R388 and **(h)** RP4 in

Comm60. Antibiotic pulsing induced alternative steady state for plasmid persistence.

Purple curves: communities passaged daily at 1:100 dilution ratio in LB supplemented

with 25µg/mL Chloramphenicol. Orange: communities were treated with additional

antibiotic pulse between Day 2 and 3 (grey bar) to select for the plasmid-carrying

populations (R388: 10 µg/mL Trimethoprim, RP4: 10 µg/mL Tetracycline).

(i). Strain-level community dynamics (mean ± SE) of Comm87 without (left) or with

(right) 1-day of Trimethoprim treatment. The five most abundant strains (based on

cumulative relative abundance across samples) are color-coded. Strain 1 corresponds

to the donor strain carrying plasmid R388.

(j). Change in the relative abundance of the donor strain (Δ donor) before (day 2) and

after (day 3) the antibiotic pulse. Bars represent mean ± SE (n = 3). Welch's t-test, **: p

< 0.01, ***: p < 0.001.

Figure 4. Ghost effect in niche partitioned communities.

(a & b). Microbes co-exist through partitioning of resource or habitat.

(c). Change in the community composition over time for a mixed population of two strains (green and purple) with complete niche partitioning (no interaction). Orange shading indicates the plasmid-carrying fraction of strain 1 (green). The community experienced selection for the plasmid-carrying cells between day 2 and 3 (gray bar).

(d). Plasmid abundance over time with (orange) and without (purple) 1-day antibiotic selection (gray bar) for the plasmid. The ghost effect occurs without plasmid abundance reaching P% = 100%.

(e). Microbial communities containing plasmid-free and plasmid-carrying *E. coli* MG1655 and eight sink isolates were cultured in LB media with sponge-imposed structure. Genus-level taxonomies of the eight sink isolates are shown. The isolates include one *Pseudomonas* strain (*P. Aeruginosa*), three *Bacillus* strains, one *Citrobacter* strain, and three *Enterobacter* strains (including one strain identified as *E. cloacae*). Strains are

color-coded by genus correspondingly as in (**i – k**). Three plasmids (pSC101, colE1,

and pUC) were separately cultured in three independent communities.

(f – h). Time series of GFP/OD for GFP-encoding plasmids pSC101, colE1, and pUC.

Purple: communities were passaged daily at $1:2\times10^7$ dilution ratio in sponges immersed

in LB. Orange: communities were treated with additional antibiotic pulse between day 1

and 2 (grey bar) to select for the plasmid-carrying populations (pSC101 and colE1: 50

µg/mL Kanamycin, pUC: 50 µg/mL Spectinomycin). Green bar: mean ± SE GFP/OD for

E. coli MG1655 carrying the corresponding plasmid (n = 3). Error bars indicate

propagated measurement uncertainties (per-well SD).

(i – k). 16S-resolved community dynamics between day 0 (D0) and day 2 (D2) (n = 1 for

day 0, mean ± SE, n = 3 for other samples). After the antibiotic pulse, the communities

were dominated by *E. coli* and *P. aeruginosa*. Due to limited taxonomic resolution of

16S sequencing, *E. coli* was assigned as *Escherichia-Shigella*, here we denote it as

Escherichia. Taxa with maximum relative abundance < 1% across all samples are not

shown.