Title: Major cell type differences between larval and adult hemichordate body plans

Authors: Paul Bump[1,2], Carolyn Brewster[3], Laurent Formery[1,4], Lauren Lubeck[1], Catherine

Campbell[1,5,6], Maurizio Morri[7], Rene Sit[7], Daniel S. Rokhsar[4,7,8], Blair Benham-Pyle[9], Alejandro

Sánchez Alvarado[3], Christopher J. Lowe*[1,7]

Affiliations:

[1]Hopkins Marine Station of Stanford University; Pacific Grove, CA.

[2]Department of Organismic and Evolutionary Biology, Museum of Comparative Zoology,

Harvard University, Cambridge, MA.

[3]Stowers Institute for Medical Research; Kansas City, MO.

[4]Molecular and Cell Biology Department at the University of California; Berkeley, CA.

[5]Unit on Developmental Signaling, Division of Developmental Biology, Eunice Kennedy Shriver

National Institute of Child Health and Human Development, NIH; Bethesda, MD.

[6]Department of Biology, Johns Hopkins University; Baltimore, MD.

[7]CZ Biohub; San Francisco, CA.

[8]Okinawa Institute of Science and Technology, Onna, Okinawa, Japan.

[9]Baylor College of Medicine; Houston, TX.

*Corresponding author. Email: clowe@stanford.edu

Abstract:

A major gap in our understanding of animal development is how adult body plans arise in

animals with indirect development, where adults emerge from the transformation of a distinct

larval form during metamorphosis. We address this question by examining cellular changes in

the enteropneust hemichordate *Schizocardium californicum*, a species with a complex lifecycle

and dramatic metamorphosis. Employing whole-body single-cell RNA sequencing, we chart the

cellular composition and transcriptional dynamics of larval, metamorphosis, and adult stages.

Our tissue level atlas reveals that ectodermal and endodermal cell types in larvae and adults

occupy distinct transcriptional spaces, showing greater similarity to other cell types within the

same life stage than to their counterparts in the opposite stage. In contrast, mesodermal cell

types from both larvae and adults cluster closely together, indicating conserved transcriptional

profiles. These findings demonstrate that the extensive morphological reorganization during

metamorphosis is paralleled by profound shifts in cell-type specific transcriptional programs,

highlighting the complexity of the larva-to-adult transition.

<u>Introduction</u>

Our understanding of adult development is largely driven by the detailed study of a few select

species, most of which are direct developers, defined by the formation of the adult body plan

during embryogenesis (Held, 2005; McEdward & Janies, 1993; Raff, 2008). However, this type

of developmental strategy does not represent the full diversity of animal life histories, and in

many species the adult is formed through metamorphosis – the transformation of a larval body

plan into an adult – often weeks to months after embryogenesis (Figure 1A) (Bolker, 1995;

Hickman, 1999). Although species are largely categorized as either direct or indirect developers,

developmental strategy is far less categorical and instead developmental mode largely operates

along a continuum (Formery & Lowe, 2023; Wray, 2000). At one extreme are direct developers

where adults form straight from the embryo and there are no intervening larval structures

(Jäegersten, 1972; Nielsen, 1998; Pechenik, 1999). At the other end of the spectrum are

species where the larval and adult body plans are developmentally and morphologically

uncoupled with the adult formed by transformation of larval tissue during metamorphosis

(McEdward & Janies, 1993). Most developmental model species lie closer to the direct end of

the spectrum; even species with an ecologically significant larvae, like the frog *Xenopus laevis*,

form the main elements of the chordate adult body plan during embryogenesis. On the other

end of the continuum, sea urchins represent the extreme limit of indirect development: their

embryos give rise to a bilaterally symmetric, planktonic larva that is both morphologically and

axially very distinct from the radial adult body plan that forms by a drastic body reorganization

during metamorphosis (Hyman, 1955; Wray & Raff, 1990).

Indirect development has a broad phylogenetic distribution across bilaterians and has been

proposed to represent the ancestral bilaterian developmental strategy (Martín-Zamora et al.,

2023; Nielsen, 1998). However, our mechanistic understanding of animals defined by indirect

development is largely limited to embryogenesis and early larval patterning. How the adult body

plan forms by the transformation of the larval body plan rather than from embryonic

development is poorly characterized (Formery & Lowe, 2023; Wray, 2000).

Recently, single-cell sequencing approaches have expanded our understanding of

developmental trajectories and cell type diversification. However, the dynamics of cellular

changes during metamorphosis remain poorly understood. Comparative work in single-cell

genomics has either focused on direct developers or on embryonic and early larval stages,

excluding the later adult life history stages in animals with complex life cycles (Foster et al.,

2022; Massri et al., 2021; Paganos et al., 2021; Perillo et al., 2020; Piovani et al., 2023). For a

comprehensive understanding of developmental mechanisms through metamorphosis, it is

essential to have insights into the cellular composition of larvae and adults and how these may

change through metamorphosis. Recent studies are beginning to reveal that this perspective is

essential to fully evaluate the cellular diversity represented by particular lineages, to test

hypotheses of cell type evolution, and to understand the evolution of animal body plans (Link et

al., 2025; Paganos et al., 2025; Sebé-Pedrós, Chomsky, et al., 2018; Sebé-Pedrós, Saudemont,

et al., 2018).

The enteropneust hemichordate *Schizocardium californicum* is an excellent model for the study

of indirect development (Cameron & Perez, 2012) (Figure 1B-F). Like many enteropneusts, *S.*

californicum develops via a tornaria larva that is organizationally similar to echinoderm larvae,

with two primary ciliary bands used for swimming and feeding, and limited musculature (M.

Hadfield & Strathmann, 1996; Lacalli & Gilmour, 2002; Morgan, 1891). This type of larva has

been termed "head larvae" as their body plan is largely defined by regulatory networks that

define an anterior "head" territory in bilaterians and are missing a Hox-patterned trunk domain

(Gonzalez et al., 2016; Lacalli, 2005; R. R. Strathmann, 2020). The formation of the adult

through a radical metamorphosis transforms the planktonic larva into a burrowing benthic adult

and involves extensive remodeling of the ectoderm, rapid expansion of the mesoderm,

reorganization of the endoderm, and the formation and extension of the trunk territory. Adults

are adapted for a burrowing lifestyle using a muscular proboscis, and feed using both

mucociliary particle capture and filter feeding through their pharyngeal gill slits (Cameron, 2002;

Gonzalez & Cameron, 2009). The tripartite gut is structurally similar between larva and adult,

but the gill slits begin to develop in the late larva in the anterior gut and perforate the anterior

trunk of the adult after a posterior migration during metamorphosis (Nielsen & Hay-Schmidt,

2007). The larval nervous system is defined by an apical organ and neurons associated with the

feeding ciliary band, whereas the adult has an extensive basiepidermal plexus and both a dorsal

and ventral nerve cord (Andrade López et al., 2023; Bullock, 1945; Kaul-Strehlow et al., 2015;

Miyamoto et al., 2010).

While morphological changes during enteropneust metamorphosis are extensive, the axial

register between life histories is maintained along with the original mouth and anus. Previous

work suggests that metamorphosis does not involve extensive larval cell death and larval cells

may be carried over into the adult (Bump et al., 2022), raising the question of the fate of larval

cells in the adult body plan.

The goal of this study is to identify similarities and differences in cellular composition of the

larval and adult body plan of *S. californicum* using single-cell transcriptomics at time points

representing larval, metamorphosis, and juvenile stages. We assess the degree to which cell

types overlap between life histories and whether metamorphosis represents not only a major

morphological change in body plan organization, but also a change in cellular composition. We

find that larval and juvenile cells largely occupy distinct transcriptional states and show greater

similarity to other tissues within the same life-history stage than to their putative counterparts

across stages. Additionally, pulse-chase EdU experiments through metamorphosis demonstrate

persistence of labeled larval cells into juveniles, arguing against wholesale larval cell

replacement and supporting cellular reprogramming during metamorphosis.

Results:

Sampling distinct life history stages in the same organism

To understand the cell types that make up distinct larval and juvenile animals, we sampled five

distinct life history stages in *S. californicum*: early larva, late larva, metamorphosis, early

juvenile, and late juvenile based on the previous description of development (Gonzalez et al.,

2018) (Figure 1B-F) and which are generally representative of other, more distantly related

indirect developing enteropneust species such as *Ptychodera flava* and *Balanoglossus*

simodensis.

Early larva. The first larval stage was selected at 22 days post fertilization (Figure 1B, 1G). The larval body plan is fully formed but lacks any nascent adult structures. Tornaria larvae are small, simple planktonic organisms that feed and swim using cilia organized into two distinct ciliary bands (Figure 1B, 1G): the circumoral ciliary loop lines the anterior of the larva and is used primarily for feeding, while the circumferential telotroch (Figure 1G, 1H) in the posterior of the larva is defined by long compound cilia and is exclusively used for locomotion (Garstang, 1939; Lacalli & Gilmour, 2002; R. Strathmann & Bonar, 1976). The remaining ectoderm is squamous epithelium with a ciliated neurotroch that runs along the ventral midline between the anus and the posterior circumoral ciliary loop. At the most dorsal anterior there is an apical tuft of cilia and an associated underlying apical organ with neurons that project throughout the epithelium and two pigmented eyespots (Gonzalez et al., 2018). There is very little mesoderm in early larvae, as only a single anterior protocoel is present at this stage (Figure 1G). The larval muscle is localized around the pharynx and in a single apical, muscular strand that contracts and connects the protocoel to the apical plate (R. Strathmann & Bonar, 1976; Tagawa et al., 1998) (R. Strathmann & Bonar, 1976; Tagawa et al., 1998). The endoderm of the tornaria larva consists of a tripartite gut composed of the pharynx, stomach, and intestine with a terminal anus.

Late larva. The second larval stage was selected after two months of growth (Figure 1C, 1H), when the larva is competent to begin metamorphosis (M. G. Hadfield et al., 2001). While morphologically similar to the early tornaria, this stage is defined by a much larger size and more extensive feeding ciliary band and evidence of pre-patterning of adult structures. In the mesoderm the additional two pairs of posterior coeloms, the mesocoels and metacoels (Figure 1H), have formed, and nascent gill slits are present in the anterior endoderm. The posterior to the telotroch is larger and there is an expanded posterior region or anal field. The nervous

system is extensive, with an expansion of the serotonergic nervous system throughout the

ectoderm and the first signs of the development of the adult dorsal cord (Gonzalez et al., 2018).

Metamorphosis (Agassiz stage). The third developmental stage selected was the midpoint of

metamorphosis (Figure 1D, 1I). Metamorphosis is an approximately 48 hour long process that

begins with a thickening of the larval ectoderm, the initiation of trunk development, and an

expansion of all the mesodermal coeloms as the large blastocoelar space is reduced in an

anterior to posterior wave (Agassiz, 1873; Nielsen & Hay-Schmidt, 2007). The midpoint of

metamorphosis contains morphological elements of both the larval and adult body plans. It is

defined by a thickened prospective proboscis ectoderm resulting from the degeneration of the

circumoral ciliary band, and an expanded anterior coelom that becomes the musculature of the

proboscis. The stomochord has formed as an anterior projection from the gut at the base of the

proboscis supporting the heart/kidney complex responsible for coelomic fluid filtration and

circulation (Balser & Ruppert, 1990). At this stage trunk formation has been initiated and

elongation has begun, but the blastocoel remains prominent even as the mesocoel and

metacoels expand (Figure 1D, 1I). In the endoderm the nascent gill slits are migrating

posteriorly towards the trunk and the collagenous skeletons of the proboscis and gill slits are

beginning to develop. Metamorphosing *S. californicum* are still planktonic and use the telotroch

for locomotion.

Early and late juveniles. The definitive adult body plan is represented in our study by early

(Figure 1E, 1J) and late (Figure 1F, 1K) juveniles: benthic, burrowing worms that have not

reached sexual maturity. The tripartite adult body plan typical of enteropneusts is clearly defined

into proboscis, collar, and trunk regions: the mouth opens ventrally at the base of the proboscis,

the gill pores perforate the ectoderm in the anterior trunk, and both dorsal and ventral nerve

cords are present. Early juveniles are newly settled animals with well-developed proboscis and

collar that have not undergone substantial trunk elongation (Figure 1E, 1J). To ensure that we

captured the full cellular complement of the adult body plan, we also sampled late juveniles

(Figure 1F, 1K) defined by a significant elongation of the trunk.

Single-cell sequencing reveals the diversity of hemichordate cell types

To understand the cellular composition of each of these distinct life history stages whole

organism cell suspensions were prepared for single-cell RNA sequencing (scRNA-seq). Cell

suspensions were processed and a total of 87,021 single-cell profiles were sequenced: 15,403

cells for early larva, 22,027 for late larva, 13,771 during metamorphosis, 20,061 for early

juvenile, and 15,759 for late juvenile.

We generated an atlas of *S. californicum* across life history stages by clustering cellular profiles,

identifying 56 clusters containing between 68 and 8,392 cells (average 1,554 cells/cluster)

(Figure 2A, Supplemental Figure 1A, Supplemental Table 1). Cell clusters are defined by

markers obtained by analysis of differential expression (log fold change and p-value) and

specificity (Supplemental Table 2). Based on cluster-specific gene expression we further

organized these cell type clusters into 12 cell classes (cartilage, ciliary band, endoderm,

epidermis, immune cells, mesoderm, muscle, neural, pigment, proliferative, secretory, and

unknown) (Figure 2B, Supplemental Table 2).

We validated our cell type clusters using both newly discovered markers from our cluster

annotations (Supplemental Table 2, 3) and known tissue markers in hemichordates, by plotting

the relative expression of a gene of interest in each cell class (Figure 2B) and then performing

HCR (hybridization chain reaction) fluorescent *in situ* hybridization (Figures 2C-T, Supplemental

Table 4).

Endoderm derivatives. *Endoderm clusters: 0, 7, 8, 9, 21, 27, 35, 53* (Figure 2C-G,

Supplemental Figure 2A-C). Differentially expressed genes for endoderm include fatty acid

binding proteins and *foxA*, a known endoderm marker in hemichordates (Fritzenwanker et al.,

2014; Green et al., 2013). *FoxA* is found in all eight of our endoderm clusters, with limited

expression outside of this group. It is expressed throughout the tripartite gut of *S. californicum*

larvae in the mouth, stomach, and intestine (Figure 2C and 2D), consistent with prior

observations in *S. kowalevskii* (Fritzenwanker et al., 2014). As in *S. kowalevskii, S. californicum*

juveniles have a ring of *foxA* expression in the anterior collar groove, but unlike *S. kowalevskii*

there also seems to be a broader expression domain in the ectoderm (Figure 2D). We also

identified additional markers from our single-cell analysis that enabled finer-scale

characterization of endoderm. For example, the gut marker *fabp6*, which encodes a fatty acid

binding protein family member responsible for lipid balance and retinol- and retinoic acid-binding

across animals (Y. Zheng et al., 2013), is highly expressed in the larval intestine (Figure 2E)

while *fzd1/2/7* was expressed in the larval stomach (Supplemental Figure 2C).

Cartilage cluster: 19 (Figure 2F-G, Supplemental Figure 2D-F). A single cluster was enriched for

several cartilage markers, including *collagen alpha-1(IX) chain-like* (*co9a1*). Since cartilage in

enteropneust hemichordates is secreted by the endodermal pharyngeal epithelium (Rychel et

al., 2006) and the gill slits specifically are made up of fibrillar collagen (Pardos & Benito, 1988),

we expected *co9a1* expression would be localized to the gill slits, a characteristic deuterostome

feature. We found that *co9a1* is expressed in a small pocket of cells in the late larva in the

developing gill slits (Figure 2F), through metamorphosis (Supplemental Figure 2F), and in

juveniles in the gill bars and the wishbone-shaped proboscis skeleton (Figure 2G). This cluster

also expresses *foxI,* a marker implicated in gill pouch endoderm and in the posterior gut

(Supplemental Figure 2E) (Fritzenwanker et al., 2014).

Mesoderm derivatives. *Mesoderm clusters: 3 and 30* (Figure 2H-I, Supplemental Figure 3A-B).

Differentially expressed genes in the mesoderm clusters included *mesenchyme homeobox*

(*mox*), a gene expressed in the paired posterior coelomic cavities in *S. kowalevskii* (Lowe et al.,

2006). Using HCR we detected *mox*-positive cells in *S. californicum* during initiation of the

posterior coelom formation prior to any clear morphological outpocketing of mesoderm (Figure

2H) as well as in clearly defined coeloms of the late larva (Figure 2I).

Muscle clusters: 5 and 18 (Figure 2J-K, Supplemental Figure 3C-E). Enteropneust adults are

highly muscular, and muscle clusters were identified in our data set by the expression of muscle

components such as *titin, popeye domain-containing protein 3,* and *myosin light chain (myl)*

(Supplemental Figure 3D, E). Expression *myl* was detected in all life stages and was particularly

prevalent near the end of metamorphosis where longitudinal fibers are forming to give rise to the

musculature of the juvenile worm (Figure 2J, 2K).

Immune clusters: 15, 17, 20, 22, 23, 29, 37, 42, 46, 47 (Figure 2L, Supplemental Figure 4A-B).

scRNA-seq allowed us to characterize many previously undescribed cell types including

putative immune cells. We found a number of immune populations that expressed macrophage

mannose receptors including C-type lectins expressed by most tissue macrophages and

dendritic cells (Taylor et al., 2005; Vasta et al., 2017), as well as a specific population of cells

(cluster 17) marked by the expression of *collagen triple helix repeat-containing protein* (*cthr*)

(Figure 2L), a secreted glycoprotein implicated in tissue repair processes (Myngbay et al.,

2021).

Ectodermal derivatives. *Neural clusters: 1, 2, 10, 16, 26, 28, 33, 34, 38, 39, 40* (Figure 2M-O,

Supplemental Figure 5A-C). We identified neural cell clusters by the expression of

synaptotagmin (*syt1*), which is involved in synapse docking (Littleton et al., 1993) (Figure 2M)

as well as *elav*, a neuron-specific RNA binding protein expressed in differentiated bilaterian

neurons (Cunningham & Casey, 2014; Koushika et al., 1996; Robinow & White, 1991), but as

noted in other species, it does not mark all neurons (Berger et al., 2007; Pham & Hobert, 2019).

Neural subtypes were identified by the expression of enzymes involved in the synthesis of

neurotransmitters or transporters: *tyrosine hydroxylase* (*TH*), which marks dopaminergic

neurons (Figure 2N and 2O), the GABAergic marker *GAD*, and cholinergic marker *VAChT*

(Supplemental Figure 5C). We observed these neural cell types in both larvae and juveniles.

Ciliary band clusters: 6 and 11 (Figure 2P-Q, Supplemental Figure 5E-G). The tornaria larva is

characterized by two distinctive ciliary bands, the circumoral ciliary band and the telotroch

(Figure 1B, 1C, 1G, 1H). Clusters 6 and 11 expressed known ciliary markers. Cluster 6 showed

enrichment of *polycystic kidney disease protein 1-like 2* (*pk1l2*), which has been associated with

the primary apical cilium of renal epithelia (Marra et al., 2019; Pfeffer et al., 1998; Yoder et al.,

2002). When we examined the expression of *pk1l2* via HCR imaging we found that it labeled the

circumoral ciliary band but not the telotroch (Figure 2P). Cluster 11 was marked by *cyclic*

nucleotide-binding domain containing protein (*cnbd2*), which may be involved in ciliary beating

(Darszon et al., 2008; Hildebrandt et al., 1997). HCR imaging showed that *cnbd2* had a distinct

expression pattern in the telotroch (Figure 2Q).

Secretory clusters: 12, 32, 36, 49, 55 (Figure 2R-V, Supplemental Figure 6A-B). Hemichordate

epidermis is highly secretory, producing large amounts of mucus from the epidermis, and we

found several secretory clusters in both larvae and juveniles. We propose that cluster 12 is a

secretory cell population based on the high expression of *mucin2 (muc2)* (Supplemental Figure

6B). Mucins have been implicated as important for muco-ciliary feeding in hemichordates

(Ruppert et al., 1999). HCR imaging showed *muc2* expressed in a thin circumferential band of

cells in the collar of the juvenile (Figure 2R), similar to the expression of mucins in *S.*

kowalevskii (Simakov et al., 2015). Even for putative secretory clusters with a smaller number of

cells (374 for cluster 49) we were able to recover distinct patterns of cell type-specific gene

expression. Cluster 49 is characterized by expression of *alpha-amylase* (*amy1*). Alpha-

amylases hydrolyze 1,4-alpha-glucoside bonds in oligosaccharides and polysaccharides in the

salivary gland of humans (Samuelson et al., 1988). HCR imaging showed *amy1* expression in a

very distinctive pattern of cells surrounding the larval stomach (Figure 2S), so we propose that

this cell population may be involved in secretory function. Finally, we found a cluster of

secretory cells with a striking shift in abundance and distribution over developmental time. The

markers of a putative secretory cluster 32 included *arylsulfatase* (*arsi*), a protease that is

upregulated in the *Xenopus laevis* tadpole tail as an integral part of the resorption process

during metamorphosis (Das et al., 2006). In the late larva, these *arsi+* cells are found in clusters

throughout the ectoderm (Figure 2T), become broadly distributed during metamorphosis (Figure

2U), and finally are predominantly restricted to the most posterior of the early juvenile worm

(Figure 2V). Future work will be necessary to investigate the function of this intriguing cell type

during metamorphosis.

Divergence of transcriptional states between life histories

Given that we had sampled the single-cell composition of five life history stages with distinct

morphologies and ecological demands, we wanted to understand how cellular composition

relates to life history. We considered three possible scenarios: First, that larval cell types

represent a subset of adult cell types; second, that the cells of larval and adult cells share

transcriptional identity, while each stage also possesses unique cell types; and third, that larval

and adult body plans are composed of distinct cell types with limited overlap in transcriptional

identity.

To test these possibilities, we plotted life history on our scRNA-seq clusters (Figure 3A). We

observed a striking pattern when comparing the distribution of cells across the five different life

history stages: larval and juvenile stages largely clustered in distinct regions of transcriptional

space (Figure 3A). To confirm that this observation was not an artifact resulting from our chosen

methods for integrating cells from different stages into a single transcriptional space (See

Material and Methods, Supplemental Figure 1), we performed "Harmony" and "Portal"

291 integrations, which still resulted in spatially separated clustering of larval and juvenile life stages

292 (Supplemental Figure 1B and 1C).

293 To examine changes in cell type compositions between life stages, we looked at the proportion

294 of cells in a given cluster from each timepoint (Figure 3B). Broadly, these life stage contributions

295 matched our expectations of the morphology of the organism: some tissues are composed

296 almost entirely of juvenile cells, like cartilage (cluster 19), while others are predominantly larval

297 cells, such as the ciliary band (clusters 11 and 6). Consistent with the stage-related decrease in

298 the number of cells in ciliary band clusters, we showed that the expression of markers of cluster

299 6 such as *pk1l2* decrease over time and are restricted to the circumoral ciliary band and future

300 oral groove of the proboscis during metamorphosis (Supplemental Figure 5G). This aligns with

301 known morphology and with our previous study that found a higher level of cell death in the

302 circumoral ciliary band (Bump et al., 2022).

303 Based on our annotation, neural cells are one of most broadly distributed classes of cell types in

304 both larvae and juveniles. Some cell types were present in both life stages but varied in their life

305 stage composition, with a key example being the neural cell types (clusters 26, 16, 34, 1, 2

306 dashed boxes in Figure 3B with corresponding panels in Figure 3C-E).

307 There were neural clusters composed of predominately larval cells (Figure 3C), such as cluster

308 26 (Figure 3Ci) which was marked by expression of *nas15*, a zinc metalloproteinase, which is a

309 part of a large family of astacins, some of which are expressed in the nervous system (Park et

310 al., 2010). This cluster also expressed *lysine-specific demethylase 8* (*kdm8-7*), an important

311 regulator of neurite morphogenesis (Zibetti et al., 2010). We found *nas15* expressed throughout

312 the posterior of the larva in cells of the developing telotroch (Figure 3Ci). This neural cell type is

313 absent in juvenile life stages, likely due to the loss of ciliary band at metamorphosis. Neural

314 cluster 34 expressed *dopamine beta-hydroxylase* (*dopo*), which converts dopamine to

noradrenaline (Axelrod, 1972) and *neuronal acetylcholine receptor subunit alpha 6* (*acha6*),

which belongs to the ligand-gated ion channel superfamily, including GABA, glycine, and 5-HT3

receptors (Betz, 1990). We found *acha6*+ cells distributed around the ventral anterior of the

larva (Figure 3Cii). Other neural cell clusters were composed of cells that were predominantly

from the juvenile stage, such as clusters 1 and 2 (Figure 3D). Cluster 1 was marked by the

expression of *stum*, which has been implicated in mediating mechanical sensing in receptor

neurons and is a transducer for mechanical stimuli (Desai et al., 2014). HCR showed *stum*

expression along the dorsal and ventral nerve cords of the juvenile and projecting into the

proboscis on the dorsal surface (Figure 3Di). Neural cluster 2 is marked by *fibrilin-2-like*, a large

modular structural glycoprotein component of the extracellular matrix of peripheral nerves

(Chernousov et al., 2010). Visualizing *fibrilin-2-like* via HCR showed expression in the collar of

juveniles, with an enrichment of *fibrilin-2-like*+ cells with neural morphology at the base of the

collar (Figure 3Dii). An example of a neural cell type present in both life history stages, neural

cluster 16 (Figure 3E) is marked by *partner of bursicon* (*pburs*). HCR of *pburs* revealed

expression in the eyespots as part of the larval apical organ (Figure 3Ei) (Nakajima et al., 2004;

Nielsen, 2005). As development continued, expression was persistent in the eyespots

(Supplemental Figure 5D) and continued into the juvenile in a persistent expression territory in

two patches at the tip of the proboscis (Figure 3Eii). While the apical tuft of cilia associated with

the apical organ degenerates at metamorphosis, some of the neurons and the eyespots appear

to persist in the juvenile body, at least at juvenile stages (Gonzalez et al., 2018).

Finally, as described above, while many cell types had larval- or juvenile-specific transcriptional

signatures, muscle (clusters 5 and 18) was an interesting exception (Figure 3F). A highly

differentially expressed gene in these muscle clusters is *myosin heavy chain* (*mhc*). One of the

few muscle types that is present in the larva is the apical strand muscle, which is marked by the

expression of *mhc* (Figure 3Fi). During metamorphosis and into the early juvenile stage, *mhc*

expression expands into the distinctive musculature of the juvenile worm, one of its defining

morphological characters (Figure 3Fii, Supplemental Figure 3F). While the prevalence and

distribution of muscle cells change over developmental time in the animal, our single-cell

analysis suggests that transcriptional signature of these clusters remains essentially unchanged

as they do not show spatially separated clustering of larval and juvenile cells (Figure 3F).

Given the changes that we saw in the transcriptional profile, we considered two principal

possibilities: (1) extensive larval cell death during metamorphosis followed by proliferation of

adult stem cell populations, or (2) larval cells are carried over into the juvenile animal with

modified patterns of gene expression. Extensive larval cell death at metamorphosis is not

supported by our previous observations (Bump et al., 2022), so to further explore the possibility

of cells being carried over from larval to juvenile we performed a series of EdU (5-ethynyl-2'-

deoxyuridine) pulse-chase experiments through metamorphosis. We incubated early and late

larvae with EdU for 48 hours to label any cell in S-phase (Figure 3G, 3H), then washed out the

EdU label and let the animals continue to grow and develop until they completed

metamorphosis. In an early larva, a 48-hour pulse indiscriminately labels a number of different

cellular populations including those in the ectoderm (ciliary band), mesoderm (wrapped around

the gut), and endoderm (in the mouth and gut) similar to what we had observed with a short

EdU pulse previously (Bump et al., 2022). Forty-three days later, we found that some EdU+

cells were still present, particularly in the anterior proboscis and along the dorsal midline (Figure

3Gii), indicating that at least some cells from the larval body plan persist through to the juvenile

stage. In late larvae, closer to metamorphosis, 48-hour pulses label many cell populations within

all three germ layers (Figure 3Hi). These cell population include the ciliary bands, the epithelium

of the presumptive collar and proboscis ectoderm, the mesocoel and metacoel, and a large

enrichment in the gill bars, consistent with previous reports of shorter EdU pulse-chase

experiments at this particular stage (Bump et al., 2022). In this case, EdU+ cells persisted in the

juvenile, with a strong enrichment in proboscis, collar, and dorsal cord (Figure 3Hii). From this

we conclude that larval cells could be carried into the juvenile animal with possible changes in

transcriptional profile.

Differences between cells in larval and adult body plans

We further quantified and interrogated the overall transcriptional relationships between cell

types and life stages using our scRNA-seq dataset. While two- or three-dimensional

embeddings like UMAP plots (Figure 2A, Figure 3A) are convenient for visual analysis, low-

dimensional representations unavoidably distort complex relationships among large numbers of

clusters (Healy & McInnes, 2024). To more accurately quantify similarity (closeness) or

dissimilarity (remoteness) of clusters in 55-dimensional principal component (PC) space, we

calculated the centroid for the PC embeddings of the cells in each cluster and then calculated

pairwise distances between cluster centroids (Figure 4A). These distances were used to form a

hierarchical clustering, and we calculated the PC-space distance at each branch point (Figure

4B). We found that cluster 3 and 30 cluster together, which together defined the broad

mesodermal cell type, even though cluster 3 was primarily composed of juvenile cells and

cluster 30 was primarily composed of larval cells (Figure 4B). Larval gut clusters 9 and 35 are

closer to larval neural clusters 28 and 26 than to other adult clusters from the juvenile. This is

surprising as these cell types arise from separate germ layers, the endoderm and ectoderm.

Adult neural clusters 38, 1, 2, and 40 are transcriptionally more similar to adult gut clusters 21,

8, and 27 than they are to larval neural clusters (Figure 4B). These findings supported our

predictions based on the observations of the UMAP that some cell types are more similar

between life stages than they are between cell types.

We next quantified whether transcriptional shifts were larger between cell types in the same life

stage or between life stages of the same cell type. To do this, we grouped the cells first by

tissues of interest, then split those groups by life stage (all cells outside the tissue of interest are

grouped into "Other"). As before, we found the centroids of these groups in transcriptional space

and calculated pairwise distances between them, then used these to generate a hierarchical

clustering (Figure 4Ci, 4Di). To help evaluate how representative each centroid was of the

position of its cells in transcriptional space, we also calculated the distance from individual cells

in each group to each group centroid (Figure 4Cii, 4Dii).

Interestingly, we found that larval mesodermally-derived cells, either mesoderm or muscle, are

more transcriptionally similar to juvenile mesodermal cells than they are to other larval cell types

(Figure 4Ci-4Cii). This is counter to what we observed for many of the endodermally- and

ectodermally-derived cell types. For example, larval neurons and larval gut cells are generally

more transcriptionally similar than either is to adult neurons or adult gut cells, respectively

(Figure 4Di-Dii). This was suggested by the space they occupy in low-dimensional embedding

on the UMAP but bolstered by higher-dimensional analysis.

Genetic makeup of cell type diversity across life history stages

Next, we sought to understand what kind of genes were driving transcriptional differences

between larval and juvenile cell types and how those genes were being regulated. We first used

GO term analysis to find classes of genes enriched at different life stages in scRNA-seq

differential expression data (Figure 5A). These results recapitulated our understanding of

hemichordate biology: processes such as cilium organization, assembly, and movement were

highest in early and late larva, concomitant with the presence of the circumoral ciliary band and

the telotroch, as well as during metamorphosis when the circumoral band has degenerated but

the telotroch is still retained. In addition, we observed that GO terms were enriched on a cluster-

by-cluster basis (Figure 5B) which also reflected the possible function of these various cell types

– for example clusters 6 and 11 were enriched for ciliary components. Ordering the clusters by

life stage composition revealed temporal patterns with an enrichment of cellular differentiation

and developmental processes starting at metamorphosis. As development proceeds through

metamorphosis and into early juvenile and late juvenile stages, we observe an enrichment of

GO terms related to morphogenesis like neurogenesis (clusters 16, 1, 2) again reflecting the

biological processes occurring at these stages and their underlying cell types (Figure 5B).

Importantly, we also noted enrichment of GO terms associated with the Wnt signaling pathway

during metamorphosis and juvenile stages. It has been postulated that in hemichordates the

Wnt signaling pathway may be responsible for driving the patterning of posterior Hox-expressing

territory (Darras et al., 2018). The anteroposterior patterning program, including the Hox

complement (and body patterning more broadly) has previously been described in *S.*

californicum, with the larva being mostly "head" or anterior territory, while the trunk region that

starts to expand at metamorphosis expresses posterior markers (Gonzalez et al., 2016). We

investigated the expression of anteroposterior patterning genes across scRNA-seq, and

observed specific enrichment of posterior patterning genes in metamorphosis and juvenile

stages compared to larval stages (Figure 5C). More striking, these patterns of gene enrichment

also correlated with enrichment of cell types in each life stage (Figure 5D). Indeed, neural

clusters, 38, 40, 1, and 2 that are abundant in early and late juvenile stages also highly express

genes of the collar and trunk territories, such as *lim1/5*, *gbx*, *hox5*, and *hox6* (Figure 5D).

While these patterning genes may reflect the distribution of these cell types across life stages,

we next wanted to examine how different types of genes might explain the contrasts between

larval and juvenile cell types. To explore this, we ran a principal component (PC) analysis and

UMAP embedding on our scRNA-seq data using either exclusively predicted transcription

factors (Figure 5E, 5F) or non-transcription factor genes (Figure 5G, 5H). Both analyses

demonstrate some cell separation by life stage (Figure 5E, 5G) and recapitulate separation of

the broad cell types (Figure 5F, 5H) we originally identified (Figure 2A). This suggests that both

transcription factor and non-transcription factor genes reflect the large effect of physiology and

homeostasis between life history stages. This approach demonstrates that regulatory,

physiological, and homeostatic programs are all involved in determining cell type composition

and transcriptional signatures across disparate life stages. These differences are perhaps linked

to the differing ecological demands that body plan morphology must meet as a planktonic larva

versus as a mud-dwelling benthic juvenile.

Discussion

We examined the diversity of cell types in the hemichordate *S. californicum* across five

developmental stages: two larval stages, one during metamorphosis, and two juvenile stages.

This analysis identified 12 cell classes: cartilage, ciliary band, endoderm, epidermis, immune,

mesoderm, muscle, neural, pigment, proliferative, secretory, and unknown. Our findings

encompass all major tissue types and include previously unrecognized cell populations.

Some of our findings were unsurprising based on our existing knowledge of hemichordate

biology and the loss or gain of cell types associated with loss and gain of structures between life

history stages. For example, two ciliary clusters mark the two ciliary bands of the larva, and both

the cell clusters and morphological structures are lost following metamorphosis as the benthic

adult body plan is formed. Conversely, we observe the appearance of novel clusters associated

with the adult body plan, such as the cartilage that is used for structural support of the proboscis

and the gill slits.

Our primary objective was to assess how cell type composition changes throughout the

organism's complex life cycle. By characterizing broad cell types, we established new molecular

markers to track the distribution of major cell classes across developmental stages.

We considered three explicit hypotheses regarding how cellular composition maps to life history.

First, do larval cell types represent a subset of adult cell types, with metamorphosis leading to

an expansion of cellular diversity as the organism develops a more complex adult body plan?

Second, do larval and adult stages share a substantial portion of their cellular complements, but

each stage also possesses unique cell types? Third, are larval and adult body plans composed

of distinct cell types, exhibiting minimal overlap in their transcriptional programs?

Surprisingly, our data largely supports the third hypothesis: when life history is overlaid onto the

low-dimensional embeddings, there is a marked separation of cell states associated with life

history (Figure 3A), and this separation is borne out by higher-dimensional analysis (Figure 4B).

Larval cells cluster together to define a larval transcriptional state, largely to the exclusion of

juvenile states, with metamorphosis representing an intermediate state. In other words, cells

cluster primarily by life history stage rather than by cell type identity. For example, larval

neurons and larval gut cells are generally more transcriptionally similar to each other than either

is to adult neurons or gut, respectively (Figure 4B, 4Di-ii). This pattern is suggested in UMAP

embeddings and confirmed by hierarchical clustering.

Both endoderm and ectoderm-derived cell types follow this stage-specific clustering pattern:

larval cells clustering with other larval cells, and adult cells clustering with other adult cells.

Mesoderm and muscle are exceptions. These cell types maintain similar transcriptional profiles

in both larvae and adults, suggesting that mesoderm undergoes metamorphosis through

different mechanisms than the other two germ layers.

Developmental mechanisms of life history cellular transformation

We propose two hypotheses to explain these observations. First, larval cells may undergo large-

scale apoptosis, with adult cells arising from sequestered stem cells. This largely fits with

Davidson and colleagues' "set aside" hypothesis inspired from sea urchin development, where

adult structures would presumably originate from sequestered progenitor cells (Peterson et al.,

1997) and metamorphosis would represent a total cellular turnover between larvae and adults.

The transcriptional state discontinuity observed in endodermally and ectodermally derived cells

between larval and adult stages (Figures 3, 4) and the loss of larval ciliary bands at

metamorphosis could be consistent with this hypothesis. However, previous work did not reveal

extensive apoptosis during metamorphosis, with TUNEL staining restricted to ciliary band cells

at the start of metamorphosis (Bump et al., 2022). Furthermore, the extensive carryover of EdU-

labeled larval cells through metamorphosis into adulthood that we observed (Figure 3Gi-Hii) is

inconsistent with extensive larval cell turnover between life histories.

In the second hypothesis, larval cells may undergo transcriptional reprogramming during

metamorphosis via dedifferentiation and redifferentiation, or via transdifferentiation, which has

been postulated to play a key role during adult body plan development by transformation of a

larva (Arenas-Mena, 2010; Sánchez Alvarado & Yamanaka, 2014). Lineage and functional

studies are needed to test for evidence of reprogramming and to differentiate between the

possible alternative developmental explanations. However, evidence for transcriptional

reprogramming includes cluster 16 (neural cells), which spans all life stages and exhibits spatial

continuity in *pburs* expression. Larval *pburs* localizes to the eyespots in the apical organ (Figure

3Ei) and during metamorphosis it is carried over into adult apical ectoderm (Figures 3Eii,

Supplemental Figure 5D), suggesting transcriptional programming changes in retained cells.

In contrast, mesoderm does not follow this pattern. Muscle and mesoderm cells retain similar

transcriptional states irrespective of life history, as has also been described in a scRNA-seq

comparisons between larvae and adult body plans of two different sea urchin species (Paganos

et al., 2025). Additionally, similar to the findings of Paganos et al. (2025), some immune cells

exhibit overlapping transcriptional states or close interstage relationships (clusters 17, 20,

Figure 2A, 3A). In summary, the cellular composition of the transition from larval to adult body

plans in the enteropneust *S. californicum* likely involves transcriptional reprogramming of

ectodermal and endodermal cells and conservation of the mesodermal cellular program. Future

work combining cellular profiling and functional assays will clarify these mechanisms.

Life history complexities for the construction of cell atlases

Our data suggests that the striking differences in body plan organization of larvae and adults is

primarily based on fundamental differences in cell type composition and transcriptional state. As

we continue to build a better understanding of metazoan cell type evolution, it is essential that

we consider how to integrate life history complexity into a broader synthesis of animal cell type

evolution. Currently, the sampling of bilaterian phyla in comparative cell type studies is mostly

limited to direct developers or to early larval stages of animals with complex life cycles.

Much of what we understand of the changes in the cellular composition during metamorphosis

comes from seminal work in insects. In holometabolous insects many of the larval tissues

undergo programmed cell death, but some larval neurons are carried over and reprogrammed

during metamorphosis (Gilbert, 2009; Truman et al., 2023). In sea urchins and nemerteans it

has been proposed that most larval cells do not persist into adulthood (Davidson et al., 1995;

Maslakova, 2010) but this has never been explicitly tested. More broadly in bilaterians, the

process of deferred development (Bishop & Hall, 2020), such as temporal shifts in development

of the trunk programs (Gonzalez et al., 2016; Martín-Zamora et al., 2023), may help to explain

the diversity of animal life cycles. However, the cellular composition differences in relationship to

these heterochronies has not been explored in detail. Clearly our understanding of the dynamics

of cellular composition in complex life cycles requires further analysis.

Our view of cellular differentiation has been heavily influenced by the findings from model

species, and in most cases differentiation in somatic cells is terminal. Some of the earliest

branching animals are highly plastic and cell fate changes are very common (Brunet et al.,

2021; Sogabe et al., 2019). However, there are also many exceptions to the view that most cells

are terminally differentiated – even in vertebrates – and that cellular plasticity is more common

than generally acknowledged (Sánchez Alvarado & Yamanaka, 2014).

A growing range of single-cell RNA-seq projects are beginning to provide insights into how cell

types change between life history stages (Link et al., 2025; Paganos et al., 2025; Sebé-Pedrós,

Chomsky, et al., 2018; Sebé-Pedrós, Saudemont, et al., 2018). Our work demonstrates that

specific cellular complements have evolved to support the vastly different ecologies and

behaviors of larvae and adults, and that larval cells may have stable rather than terminally

differentiated cell states (Sánchez Alvarado & Yamanaka, 2014). This plasticity may be

common in species with a similar life history to *S. californicum* but that needs to be explored in

further studies. Ultimately, complex life cycles are phylogenetically broadly dispersed and an

essential component of animal biodiversity that should be investigated in more depth and

integrated into our understanding of cell type differentiation and evolution.

Methods

Collecting, spawning, and larval rearing

Adult *Schizocardium californicum* were collected in Morro Bay State Park, California, in a

mudflat located at 35°20′56.7″N 120°50′35.6″W with appropriate state permitting. Animals were

spawned as described in Gonzalez et al. 2018 with individual females transferred in bowls of

filtered seawater (FSW) and placed in an illuminated incubator at 24–26°C. Once embryos

hatched, larvae were transferred to 1 gallon glass jars with continuous stirring and fed with a 1:1

mix of *Dunaliella tertiolecta* and *Rhodomonas lens*. Every two to four days, containers were

washed, water was replaced with clean FSW, and fresh algae was added. To grow larger

numbers of animals through metamorphosis, some larvae were placed on a continuous flow-

through system by being transferred into diffusion tubes (Patry et al., 2020). Once animals

began metamorphosis, they were transferred into glass bowls with terrarium sand.

Single-cell sequencing

Cell preparation. Early and late juveniles were first diced into fine pieces with a razor blade on a

petri dish. All stages were suspended in 1 mg/mL Dispase II (Sigma D4693) in 3.3× PBS, 2%

Fetal Calf Serum, and 20 mM HEPES. Cells were pipetted intermittently with Pasteur pipettes

passed over a flame to create narrower openings. Cells were then passed through a 100 μm

mesh and centrifuged at 4°C, 150 g, for 5 minutes with slow braking. Cells were resuspended in

fresh media and passed through a 40 μm strainer on ice. Cell viability was checked with

Hoechst and counted on a hemocytometer. Importantly, cells were kept in a 3.3× PBS solution

to maintain osmolarity to seawater.

Sequencing. Cells were loaded into the 10x Chromium Controller for cell capture and libraries

were processed after SPRI size-selection with the 10x Genomics V2.0 workflow. Final libraries

were analyzed on an Agilent Bioanalyzer and sequenced using an Illumina Novaseq.

Data processing. Once reads were acquired, they were demultiplexed, mapped, and

subsequently converted into count matrices using 10x Genomics Cell Ranger 3.0.0 (G. X. Y.

Zheng et al., 2017). Cell quality filtering, clustering, and dimensionality reduction was done

using Seurat v3 in R (Hao et al., 2021; Satija et al., 2015; Stuart et al., 2019). Cells were filtered

to those with at least 500 and less than 57,000 unique molecular identifiers (UMIs) and >90 and

no more than 980 genes. We normalized each batch's gene expression using Seurat's

SCTransform, then combined the batches using reciprocal principal component analysis (rPCA)

integration (Stuart et al., 2019). This technique was chosen both for its efficiency on large

datasets, and because rPCA integration by batch resulted in good overlap of the same life stage

from different batches without forcing overlap of different life stages. After this we performed

principal component analysis (PCA) on the combined dataset with 55 principal components

(PCs), generated a shared nearest neighbor graph, used clustering to identify 56 cell clusters,

and determined which genes are most differentially expressed across the dataset.

Alternative integration approaches

To integrate data sets and test for batch effects we used the Portal algorithm (Zhao et al.,

2022). Each batch was read in from the Seurat file and integrated incrementally. The datasets

were preprocessed and integrated with default parameters and 1000 training steps. We

recovered the harmonized expression matrix in log-normalized level. We then processed this

expression matrix with the Self-Assembling-Manifold (SAM) algorithm (Tarashansky et al.,

2019), with default parameters except for without log normalization or filtering genes. Under

these parameters we recovered no significant batch effects.

To integrate data sets and test for batch effects we ran "Harmony" on our principle component

embeddings which projects cells into a shared embedding where cells group by cell type rather

than dataset-specific conditions (Korsunsky et al., 2019). Independently, we took the cell count

matrices and used the Portal algorithm where each batch was read in and integrated

incrementally (Zhao et al., 2022). The datasets were preprocessed and integrated with default

parameters and 1000 training steps. We recovered the harmonized expression matrix in log-

normalized level. We then processed this expression matrix with the Self-Assembling-Manifold

(SAM) algorithm (Tarashansky et al., 2019) with default parameters except for without log

normalization or filtering genes. Under both approaches we recovered distinct differences

between life history stage and no significant batch effects.

GO Analysis

We first performed differential gene expression of positive markers as calculated across the

relevant groups (cluster, life stage) using Seurat's FindMarkers on the SCTransform-normalized

single-cell expression data. We then filtered these results to genes with 1) average log2 fold-

change of at least 1.25; 2) adjusted p-value less than 0.0001; 3) which had a reciprocal best

BLASTp hit between *Schizocardium californicum* and *Mus musculus* GRCm39 predicted

peptide sequences. These differential expression results were then sorted by log2 fold-change

and fed into gprofiler2's gost function with organism="mmusculus" and

ordered_query=TRUE(Kolberg et al., 2023). GO Terms were manually curated for plotting.

High-dimensional analysis

To perform high-dimensional analysis of overall transcriptional similarity of clusters and cell

types, the centroid of each cluster was calculated in 55-dimensional PC space, inter-cluster

distances were found for all pairs of clusters, and these distances were scaled as the

percentage of the largest inter-cluster distance. These distances were then used to build a

hierarchical clustering of all clusters in the dataset. For more detailed analysis, cells were

grouped by the combination of cell type and life stage, group centroids and inter-group

distances calculated, and these distances scaled by the same factor as the inter-cluster

distances to facilitate direct comparison. Then the distribution of distances from each group's

individual cells to each group centroid was determined to assess the accuracy of inter-centroid

distances.

Ortholog identification

Gene orthology and naming was determined by collecting sequences of interest from related

species and then building gene trees. Sequences were aligned with MUSCLE (Edgar, 2004)

and trees were calculated with Bayesian inference trees using MrBayes version 3.1.2

(Huelsenbeck & Ronquist, 2001) in 1,000,000 generations with sampling of trees every 100

generations and a burn-in period of 25%.

In situ HCR

Probe design and preparation. Complementary DNA sequences specific to genes of interest

were submitted to the in situ probe generator from the Ozpolat Lab (Kuehn et al., 2021). The

sequences generated by the software were used to order DNA oligo pools (50 µmol DNA

oPools™ Oligo Pools) from Integrated DNA Technologies, resuspended to 1 µmol/µL in 50 mM

Tris buffer, pH 7.5. HCR amplifiers with fluorophores B1-Alexa Fluor-546, B2-Alexa Fluor-488,

and B3-Alexa Fluor-647 were ordered from Molecular Instruments, Inc.

HCR protocol. HCRs were performed based on Choi et al., 2018 and the Hybridization Chain

Reaction (HCR) in situ Protocol from the Patel Lab (Bruce et al., 2021; Choi et al., 2018).

Animals were fixed with 3.7% paraformaldehyde in MOPS fix buffer (0.1 M MOPS, 0.5 M NaCl,

2 mM EGTA, 1 mM MgCl2, 1× PBS) overnight at 4°C. Fixed samples were then dehydrated in

methanol and stored at -20°C. Samples were rehydrated in a gradual series of PBS containing

0.1% Tween-20 (PBST). They were permeabilized in detergent solution (1.0% SDS, 0.5%

Tween-20, 150 mM NaCl, 1 mM EDTA (pH 8), 50 mM Tris-HCl at pH 7.5) for 30 minutes. The

samples were then extensively washed in PBST, before being pre-hybridized in hybridization

buffer (Molecular Instruments) for 1 h at 37°C. The probes were then added to the hybridization

buffer at a final concentration of 0.05 µM and the samples were allowed to hybridize at 37°C

overnight under gentle agitation. Following hybridization, the samples were washed 4 times for

15 minutes in probe wash buffer (Molecular Instruments) at 37°C and then in 5× SSCT at room

temperature. They were then pre-amplified in amplification buffer (Molecular Instruments) for 30

minutes. Amplifiers were separately incubated at 95°C for 90 seconds and then pooled together

before being added to the amplification buffer at a final concentration of 60 nM. Amplification

was performed overnight, and the samples were subsequently washed 4 times for 30 minutes in

5× SSCT and incubated in 5× SSCT containing 1:1000 Hoechst overnight.

Pulse-chase EdU labeling

Labeling and detection of proliferating cells were performed using the Click-iT Plus EdU 488

Imaging Kit (Life Technologies), using the standard protocol with the following modifications.

Larvae were cultured in FSW supplemented with 10 µM EdU diluted from a 10 mM stock in

DMSO. Animals were pulsed for 48 hours with EdU then either fixed with 3.7%

paraformaldehyde in MOPS fix buffer (0.1 M MOPS, 0.5 M NaCl, 2 mM EGTA, 1 mM MgCl2, 1×

PBS) overnight at 4°C or transferred to fresh FSW and allowed to grow and were fixed at either

43 or 26 days later when they had completed metamorphosis. For detection of EdU

incorporation, labeled tissue was transferred to a solution of PBS and the detection was

performed following the manufacturer's protocol with an increased permeabilization time in 0.5%

Triton X-100 in PBS of 40 minutes and an increased detection time of 45 minutes.

Imaging

Samples were mounted in PBS (larva) or fructose (juveniles) between two glass coverslips

raised slightly with small clay feet. Imaging was performed on a Zeiss LSM 700. For large

samples, stacks were stitched together using the Tile tool (Zen2 software, Zeiss).

Acknowledgements

We would like to thank the staff of the Hopkins Marine Station and the staff of Morro Bay State

Park, in particular Vince Cicero, John Sayers, and Katie Drexhage, for facilitating our

collections. We would like to thank Norma Neff from Biohub for supporting the development of

genomic resources for *Schizocardium*. We thank members of the Lowe Lab, specifically Auston

Rutledge who reared many *Schizocardium* larvae and was an invaluable partner in collections.

We also thank Greg Wray, Bo Wang, Dominique Bergmann, Stephen Palumbi, Lowe Lab

members Nat Clarke, Paul Minor, Veronica Pagowski, José Andrade-Lopez, and Srivastava Lab

members Mansi Srivastava, Katy Loubet-Senear, Amber Rock, Carlos Rivera-Lopez and Catie

Breen for their helpful discussions. We finally thank previous Lowe Lab member Paul Gonzalez

for his pioneering work on *Schizocardium*. This work was supported by a collaborative CZ

Biohub Intercampus Research Award to C.J.L and a NSF predoctoral fellowship (DGE–

1147470), the Myers Trust Award, and Haderlie Memorial Award to P.B.

Data availability

The sequencing raw reads and data generated in this study have been deposited in the NCBI

BioProject database under the accession code PRJNA1297089.

Figure Legends:

Figure 1. Indirect developing hemichordates as a model to study the transition between

larval and adult body plans. A) Phylogenetic distribution of bilaterians highlighting the

prevalence of larval types (Adapted from Formery and Lowe 2023). Light microscopy of the

major life history stages of *S. californicum*: B) early larva, C) late larva, D) during

metamorphosis, E) early juvenile, F) late juvenile. Schematic diagrams of the major life history

stages of *S. californicum*: G) early larva, H) late larva, I) metamorphosis, J) early juvenile, K)

late juvenile. In the schematics, ectoderm derivatives are shown in blue, endoderm derivatives

in yellow, and mesoderm derivatives in red.

Figure 2. Single-cell sequencing reveals the broad diversity of hemichordate cell types.

A) UMAP embedding of the cell classes of *S. californicum;* 55 clusters were identified and 12

cell classes were described. B) Markers of cell classes of *S. californicum*, percentage of gene

max was calculated by taking the mean normalized expression of the gene of interest and

dividing by the max for any cell class. C-T) HCR signal in magenta, nuclei are in grey, scale

bars are 100 μm. C) *foxA* HCR expression in early larva. D) *foxA* HCR expression in early

juvenile. E) *fabp6* HCR expression in early larva. F) *co9a1* HCR expression in late larva. G)

co9a1 HCR expression in early juvenile. H) *mox* HCR expression in early larva. I) *mox* HCR

expression in late larvae. J) *myl* HCR expression during metamorphosis. I) *myl* HCR expression

in early juvenile. L) *cthr* expression in early larva. M) *syt1* HCR expression in early larva. N) *TH*

HCR expression in early larva. O) *TH* HCR expression in early juvenile. P) *pk1l2* HCR

expression in early larva. Q) *cnbd2* HCR expression in early larva. R) *muc2* HCR expression in

early juvenile. S) *amy1* HCR expression in early larvae. T) *arsi* HCR expression in late larvae.

U) *arsi* HCR expression in during metamorphosis. V) *arsi* HCR expression in early juvenile.

Figure 3. Divergence of transcriptional states is consistent with morphological changes

between life history stages

A) UMAP embedding of *S. californicum* single-cell RNA-seq data with cells colored by life stage.

Yellow: early larva, dark gold: late larva, grey: metamorphosis, light blue: early juvenile, dark

blue: late juvenile. B) Bar chart of life stage composition of clusters. C) UMAP embedding with

larval neural clusters (26, 28, 33, 34) colored by life stage. Ci) *acha6* HCR expression in early

larvae. Cii) *nas15* HCR expression in early larvae. D) UMAP embedding with juvenile neural

clusters (1, 2) colored by life stage. Di) *stum* HCR expression in early juvenile. Dii) *fibrilin-2-like*

HCR expression in early juvenile. E) UMAP embedding with neural cluster 16 colored by life

stage. Ei) *pburs* HCR expression in early larvae. Eii) *pburs* HCR expression in early juvenile. F)

UMAP embedding with muscle clusters (5, 18) colored by life stage. Fi) *mhc* HCR expression in

early larvae. Fii) *mhc* HCR expression in early juvenile. Gi) EdU pulse of early larva for 48

hours. Gii) EdU chase after 43 days once larva have completed metamorphosis, inset is of

proboscis region of the juvenile worm, boxed. Hi) EdU pulse of late larva for 48 hours. Hii) EdU

chase after 26 days once larva have completed metamorphosis, inset is of proboscis region of

the juvenile worm, boxed.

Figure 4. Mesodermal tissues share transcriptional similarity while ectodermal and

endoderm tissues reflect life stage divergence A) Schematic diagram of approach used to

accurately quantify similarity or dissimilarity of clusters in 55-dimensional PC space. B)

Dendrogram shows a hierarchical clustering of scRNA-seq clusters using the centroid of each

cluster in 55-dimensional PC space, with bars colored by cluster life stage composition.

Numbers at nodes indicate the distance between the centroids of the two branches merging at

that node, as a percent of the largest inter-cluster distance. Colored dots highlight tissues that

are further analyzed in Figures Ci, Cii, Di, Dii. C) Comparison of larval mesoderm and larval

muscle versus juvenile mesoderm and juvenile muscle, with "Other" denoting all other cells from

that life stage. Ci) Heatmap indicates inter-centroid distances as a percent of the largest inter-

cluster distance, a larger value means centroids are farther apart while a smaller value means

centroids are closer. Cii) Box plots show the distribution from the cells of a group (panel) to the

centroids of each other group (bars). D) Comparison of larval neurons and larval endoderm

versus juvenile neurons and juvenile endoderm, with "Other" denoting all other cells from that

life stage. Di) Heatmap indicates inter-centroid distances as a percent of the largest inter-cluster

distance Dii) Box plots show the distribution from the cells of a group (panel) to the centroids of

each other group (bars). In box plots, the upper whisker extends from the hinge to the largest

value no further than 1.5 * IQR from the hinge (where IQR is the inter-quartile range, or distance

between the first and third quartiles). The lower whisker extends from the hinge to the smallest

value at most 1.5 * IQR of the hinge.

Figure 5. Genetic makeup of cell type diversity across life history stages. A) GO Term

analysis of scRNA-seq differential expression data by life stage. B) GO Term analysis of

scRNA-seq differential expression data by cluster, ordered by cluster life stage composition per

Figure 3D. C) Analysis of anterior-posterior patterning genes in scRNA-seq data by life stage.

D) Mean expression of anteroposterior patterning genes in scRNA-seq data by cluster, ordered

by cluster composition chart from Figure 3D. E,F) UMAP embedding of scRNA-seq data based

on PCA using only transcription factors, colored by life stage (E) or broad cell type (F). G,H)

UMAP embedding of scRNA-seq data based on PCA using only non-transcription factors,

colored by life stage (G) or broad cell type (H).

Supplemental Figure Legends:

Supplemental Figure 1. A) Flowchart of single-cell analysis including alternative integration

approaches. B) UMAP embedding of alternative integration approaches, "Harmony" and "Portal"

plotted by life stage.

Supplemental Figure 2. A) Endoderm clusters (*0, 7, 8, 9, 21, 27, 35, 53)* plotted in global

UMAP embedding. B) Endoderm cluster expression of marker genes. C) HCR expression in

early larva of *fzd1/2/7* in cyan, *fabp6* in magenta, nuclei in grey, scale bars are 100 μm. D)

Cartilage cluster (*19)* plotted in global UMAP embedding. E) Cartilage cluster expression of

marker genes. F) HCR expression during metamorphosis of *co9a1* in magenta, nuclei grey,

scale bar 100 μm.

Supplemental Figure 3. A) Mesoderm clusters (*3, 30)* plotted in global UMAP embedding. B)

Mesoderm cluster expression of marker genes. C) Muscle clusters (*5, 18)* plotted in global

UMAP embedding. D) Muscle cluster expression of marker genes. E) HCR expression of

muscle markers *titin-4, popd3-2, myl1* in magenta, nuclei in grey, scale bars are 100 μm. F)

HCR expression during metamorphosis and in late juvenile of *mhc* in magenta, nuclei in grey,

scale bars are 100 μm.

Supplemental Figure 4. A) Immune clusters (*15, 17, 20, 22, 23, 29, 37, 42, 46, 47)* plotted in

global UMAP embedding. B) Immune cluster expression of marker genes.

Supplemental Figure 5. A) Neural clusters (*1, 2, 10, 16, 26, 28, 33, 34, 38, 39, 40)* plotted in

global UMAP embedding. B) Neural cluster expression of marker genes. C) HCR expression of

neural markers *GAD* (green), *VAChT* (cyan), nuclei in grey, scale bars are 100 μm. D) HCR

expression in late larva of *pburs* (magena), nuclei in grey, scale bars are 100 μm. E) Ciliary

clusters (*6, 11)* plotted in global UMAP embedding. F) Ciliary cluster expression of marker

genes. G) HCR expression in early juvenile proboscis inset of *pk1l2* in magenta, nuclei in grey,

scale bars are 100 μm.

Supplemental Figure 6. A) Secretory clusters (*1, 2, 10, 16, 26, 28, 33, 34, 38, 39, 40)* plotted in

global UMAP embedding. B) Secretory cluster expression of marker genes. C) Epidermis

clusters (*4, 13, 25, 52)* plotted in global UMAP embedding. D) Epidermis cluster expression of

marker genes.

Supplemental Figure 7. A) Proliferative clusters (*14, 24)* plotted in global UMAP embedding. B)

Proliferative cluster expression of marker genes. C) HCR expression in early larva of *pif1-7* in

cyan, *fgf9-1* in magenta, nuclei in grey, scale bars are 100 μm. D) Pigment cluster (*51)* plotted in

global UMAP embedding. E) Pigment cluster expression of marker genes.

Supplemental Figure 8. Phylogenetic trees of *Schizocardium californicum* orthologs.

Supplemental Tables:

Supplemental Table 1. scRNA-seq statistics for 56 clusters identified and cell class annotation

Supplemental Table 2. Cell cluster markers obtained by analysis of differential expression (log

fold change and p-value) and specificity (how frequently a marker appears).

Supplemental Table 3. Cell class markers including gene nomenclature

Supplemental Table 4. HCR probe sequences with corresponding amplifier design (B1, B2, or

B3).

References

Agassiz, A. (1873). The History of Balanoglossus and Tornaria. *Memoirs of the American*

Academy of Arts and Sciences, *9*(2), 421–436. JSTOR.

https://doi.org/10.2307/25058009

Andrade López, J. M., Pani, A. M., Wu, M., Gerhart, J., & Lowe, C. J. (2023). Molecular

characterization of nervous system organization in the hemichordate acorn worm

Saccoglossus kowalevskii. *PLOS Biology*, *21*(9), e3002242.

https://doi.org/10.1371/journal.pbio.3002242

Arenas-Mena, C. (2010). Indirect development, transdifferentiation and the macroregulatory

evolution of metazoans. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *365*(1540), 653–669. https://doi.org/10.1098/rstb.2009.0253

Axelrod, J. (1972). Dopamine-β-hydroxylase: Regulation of Its Synthesis and Release from

Nerve Terminals. *Pharmacological Reviews*, *24*(2), 233–243.

Balser, E. J., & Ruppert, E. E. (1990). Structure, Ultrastructure, and Function of the Preoral

Heart-Kidney in *Saccoglossus kowalevskii* (Hemichordata, Enteropneusta) Including

New Data on the Stomochord. *Acta Zoologica*, *71*(4), 235–249.

https://doi.org/10.1111/j.1463-6395.1990.tb01082.x

Berger, C., Renner, S., Lüer, K., & Technau, G. M. (2007). The commonly used marker ELAV is

transiently expressed in neuroblasts and glial cells in the *Drosophila* embryonic CNS.

Developmental Dynamics, *236*(12), 3562–3568. https://doi.org/10.1002/dvdy.21372

Betz, H. (1990). Ligand-gated ion channels in the brain: The amino acid receptor superfamily.

Neuron, *5*(4), 383–392. https://doi.org/10.1016/0896-6273(90)90077-s

Bishop, C., & Hall, B. (2020). *Deferring Development: Setting Aside Cells for Future Use in*

Development and Evolution. CRC Press/Taylor & Francis Group.

Bruce, H., Jerz, G., Kelly, S., McCarthy, J., Pomerantz, A., Senevirathne, G., Sherrard, A., Sun,

D., Wolff, C., & Patel, N. (2021). Hybridization Chain Reaction (HCR) In Situ Protocol.

Protocols.Io. https://dx.doi.org/10.17504/protocols.io.bunznvf6

Brunet, T., Albert, M., Roman, W., Coyle, M. C., Spitzer, D. C., & King, N. (2021). A flagellate-

to-amoeboid switch in the closest living relatives of animals. *eLife*, *10*, e61037.

https://doi.org/10.7554/eLife.61037

Bullock, T. H. (1945). The Anatomical Organization of the Nervous System of Enteropneusta.

Journal of Cell Science, *s2-86*(341), 55–111. https://doi.org/10.1242/jcs.s2-86.341.55

Bump, P., Khariton, M., Stubbert, C., Moyen, N. E., Yan, J., Wang, B., & Lowe, C. J. (2022).

Comparisons of cell proliferation and cell death from tornaria larva to juvenile worm in

the hemichordate *Schizocardium californicum*. *EvoDevo*, *13*(1), 13.

https://doi.org/10.1186/s13227-022-00198-1

Cameron, C. B. (2002). Particle Retention and Flow in the Pharynx of the Enteropneust Worm

Harrimania planktophilus: The Filter-Feeding Pharynx May Have Evolved Before the

Chordates. *The Biological Bulletin*, *202*(2), 192–200. https://doi.org/10.2307/1543655

Cameron, C. B., & Perez, M. (2012). Spengelidae (Hemichordate: Enteropnuesta) from the

Eastern Pacific including a new species, *Schizocardium californicum*, from California.

Zootaxa, *3569*, 79–88.

Chernousov, M. A., Baylor, K., Stahl, R. C., Stecker, M. M., Sakai, L. Y., Lee-Arteaga, S.,

Ramirez, F., & Carey, D. J. (2010). Fibrillin-2 is dispensable for peripheral nerve

development, myelination and regeneration. *Matrix Biology : Journal of the International*

Society for Matrix Biology, *29*(5), 357–368. https://doi.org/10.1016/j.matbio.2010.02.006

Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J.,

Cunha, A., & Pierce, N. A. (2018). Third-generation *in situ* hybridization chain reaction:

Multiplexed, quantitative, sensitive, versatile, robust. *Development*, *145*(12), dev165753.

https://doi.org/10.1242/dev.165753

Cunningham, D., & Casey, E. S. (2014). Spatiotemporal development of the embryonic nervous

system of *Saccoglossus kowalevskii*. *Developmental Biology*, *386*(1), 252–263.

https://doi.org/10.1016/j.ydbio.2013.12.001

Darras, S., Fritzenwanker, J. H., Uhlinger, K. R., Farrelly, E., Pani, A. M., Hurley, I. A., Norris, R.

P., Osovitz, M., Terasaki, M., Wu, M., Aronowicz, J., Kirschner, M., Gerhart, J. C., &

Lowe, C. J. (2018). Anteroposterior axis patterning by early canonical Wnt signaling

during hemichordate development. *PLoS Biology*, *16*(1).

https://doi.org/10.1371/journal.pbio.2003698

Darszon, A., Guerrero, A., Galindo, B. E., Nishigaki, T., & Wood, C. D. (2008). Sperm-activating

peptides in the regulation of ion fluxes, signal transduction and motility. *The International*

Journal of Developmental Biology, *52*(5–6), 595–606.

https://doi.org/10.1387/ijdb.072550ad

Das, B., Cai, L., Carter, M. G., Piao, Y.-L., Sharov, A. A., Ko, M. S. H., & Brown, D. D. (2006).

Gene expression changes at metamorphosis induced by thyroid hormone in *Xenopus*

laevis tadpoles. *Developmental Biology*, *291*(2), 342–355.

https://doi.org/10.1016/j.ydbio.2005.12.032

Davidson, E. H., Peterson, K. J., & Cameron, R. A. (1995). Origin of Bilaterian Body Plans:

Evolution of Developmental Regulatory Mechanisms. *Science*, *270*(5240), 1319–1325.

https://doi.org/10.1126/science.270.5240.1319

Desai, B. S., Chadha, A., & Cook, B. (2014). The stum Gene Is Essential for Mechanical

Sensing in Proprioceptive Neurons. *Science*, *343*(6176), 1256–1259.

https://doi.org/10.1126/science.1247761

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Research*, *32*(5), 1792–1797. PubMed.

https://doi.org/10.1093/nar/gkh340

Formery, L., & Lowe, C. J. (2023). *Integrating Complex Life Cycles in Comparative*

Developmental Biology.

Foster, S., Oulhen, N., Fresques, T., Zaki, H., & Wessel, G. (2022). Single-cell RNA-sequencing

analysis of early sea star development. *Development*, *149*(22), dev200982.

https://doi.org/10.1242/dev.200982

Fritzenwanker, J. H., Gerhart, J., Freeman, R. M., & Lowe, C. J. (2014). The Fox/Forkhead

transcription factor family of the hemichordate <i>Saccoglossus kowalevskii<i/>.

EvoDevo, *5*(1), 17. https://doi.org/10.1186/2041-9139-5-17

Garstang, W. (1939). Spolia Bermudiana. II. The ciliary feeding mechanism of Tornaria. *Q. Jl*

Microsc. Sci., *81*, 347–366.

Gilbert, L. I. (2009). *Insect development: Morphogenesis, molting and metamorphosis*.

Academic Press.

Gonzalez, P., & Cameron, C. (2009). The gill slits and pre-oral ciliary organ of *Protoglossus*

(Hemichordata: Enteropneusta) are filter-feeding structures. *Biological Journal of the*

Linnean Society, *98*(4), 898–906. https://doi.org/10.1111/j.1095-8312.2009.01332.x

Gonzalez, P., Jiang, J. Z., & Lowe, C. J. (2018). The development and metamorphosis of the

indirect developing acorn worm *Schizocardium californicuma* (Enteropneusta:

Spengelidae). *Frontiers in Zoology*, 1–24.

Gonzalez, P., Uhlinger, K. R., & Lowe, C. J. (2016). The Adult Body Plan of Indirect Developing

Hemichordates Develops by Adding a Hox-Patterned Trunk to an Anterior Larval

Territory. *Current Biology*, *27*(1), 1–9. https://doi.org/10.1016/j.cub.2016.10.047

Green, S. A., Norris, R. P., Terasaki, M., & Lowe, C. J. (2013). FGF signaling induces

mesoderm in the hemichordate *Saccoglossus kowalevskii*. *Development*, *140*(5), 1024–

1033. https://doi.org/10.1242/dev.083790

Hadfield, M. G., Carpizo-Ituarte, E. J., del Carmen, K., & Nedved, B. T. (2001). Metamorphic

Competence, a Major Adaptive Convergence in Marine Invertebrate Larvae. *American*

Zoologist, *41*(5), 1123–1131. https://doi.org/10.1093/icb/41.5.1123

Hadfield, M., & Strathmann, M. (1996). *Variability, flexibility and plasticity in life histories of*

marine invertebrates. 12.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J., Wilk, A.

J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain,

J., Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., … Satija, R. (2021). Integrated

analysis of multimodal single-cell data. *Cell*, *184*(13), 3573-3587.e29.

https://doi.org/10.1016/j.cell.2021.04.048

Healy, J., & McInnes, L. (2024). Uniform manifold approximation and projection. *Nature*

Reviews Methods Primers, *4*(1), 82. https://doi.org/10.1038/s43586-024-00363-x

Hildebrandt, F., Otto, E., Rensing, C., Nothwang, H. G., Vollmer, M., Adolphs, J., Hanusch, H.,

& Brandis, M. (1997). A novel gene encoding an SH3 domain protein is mutated in

nephronophthisis type 1. *Nature Genetics*, *17*(2), 149–153.

https://doi.org/10.1038/ng1097-149

Huelsenbeck, J. P., & Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogenetic

trees. *Bioinformatics*, *17*(8), 754–755. https://doi.org/10.1093/bioinformatics/17.8.754

Kaul-Strehlow, S., Urata, M., Minokawa, T., Stach, T., & Wanninger, A. (2015). Neurogenesis in

directly and indirectly developing enteropneusts: Of nets and cords. *Organisms Diversity*

& Evolution, *15*(2), 405–422.

Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., & Peterson, H. (2023). g:Profiler—

Interoperable web service for functional enrichment analysis and gene identifier mapping

(2023 update). *Nucleic Acids Research*, *51*(W1), W207–W212.

https://doi.org/10.1093/nar/gkad347

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner,

M., Loh, P., & Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of

single-cell data with Harmony. *Nature Methods*, *16*(12), 1289–1296.

https://doi.org/10.1038/s41592-019-0619-0

Koushika, S. P., Lisbin, M. J., & White, K. (1996). ELAV, a Drosophila neuron-specific protein,

mediates the generation of an alternatively spliced neural protein isoform. *Current*

Biology : CB, *6*(12), 1634–1641. https://doi.org/10.1016/s0960-9822(02)70787-2

Kuehn, E., Clausen, D. S., Null, R. W., Metzger, B. M., Willis, A. D., & Özpolat, B. D. (2021).

Segment number threshold determines juvenile onset of germline cluster expansion in

Platynereis dumerilii. *Journal of Experimental Zoology Part B: Molecular and*

Developmental Evolution, jez.b.23100. https://doi.org/10.1002/jez.b.23100

Lacalli, T. C. (2005). Protochordate body plan and the evolutionary role of larvae: Old

controversies resolved? *Canadian Journal of Zoology*, *83*(1), 216–224.

https://doi.org/10.1139/z04-162

Lacalli, T. C., & Gilmour, T. H. J. (2002). Locomotory and feeding effectors of the tornaria larva

of *Balanoglossus biminiensis*: Tornaria structure and feeding. *Acta Zoologica*, *82*(2),

117–126. https://doi.org/10.1046/j.1463-6395.2001.00075.x

Link, O., Jahnel, S. M., Janicek, K., Kraus, J., Montenegro, J. D., Zimmerman, B., Wick, B.,

Cole, A. G., & Technau, U. (2025). *Changes of cell-type diversity in the polyp-to-medusa*

metagenesis of the scyphozoan jellyfish Aurelia coerulea *(formerly sp.1)*.

https://doi.org/10.1101/2023.08.24.554571

Littleton, J. T., Stern, M., Schulze, K., Perin, M., & Bellen, H. J. (1993). Mutational analysis of

Drosophila synaptotagmin demonstrates its essential role in Ca(2+)-activated

neurotransmitter release. *Cell*, *74*(6), 1125–1134. https://doi.org/10.1016/0092-8674(93)90733-7

Lowe, C. J., Terasaki, M., Wu, M., Freeman, R. M., Runft, L., Kwan, K., Haigo, S., Aronowicz, J., Lander, E., Gruber, C., Smith, M., Kirschner, M., & Gerhart, J. (2006). Dorsoventral Patterning in Hemichordates: Insights into Early Chordate Evolution. *PLoS Biology*, *4*(9), e291. https://doi.org/10.1371/journal.pbio.0040291

Marra, A. N., Adeeb, B. D., Chambers, B. E., Drummond, B. E., Ulrich, M., Addiego, A., Springer, M., Poureetezadi, S. J., Chambers, J. M., Ronshaugen, M., & Wingert, R. A. (2019). Prostaglandin signaling regulates renal multiciliated cell specification and maturation. *Proceedings of the National Academy of Sciences*, *116*(17), 8409–8418. https://doi.org/10.1073/pnas.1813492116

Martín-Zamora, F. M., Liang, Y., Guynes, K., Carrillo-Baltodano, A. M., Davies, B. E., Donnellan, R. D., Tan, Y., Moggioli, G., Seudre, O., Tran, M., Mortimer, K., Luscombe, N. M., Hejnol, A., Marlétaz, F., & Martín-Durán, J. M. (2023). Annelid functional genomics reveal the origins of bilaterian life cycles. *Nature*, *615*(7950), 105–110. https://doi.org/10.1038/s41586-022-05636-7

Maslakova, S. A. (2010). Development to metamorphosis of the nemertean pilidium larva. *Frontiers in Zoology*, *7*(1), 30. https://doi.org/10.1186/1742-9994-7-30

Massri, A. J., Greenstreet, L., Afanassiev, A., Berrio, A., Wray, G. A., Schiebinger, G., & McClay, D. R. (2021). Developmental single-cell transcriptomics in the *Lytechinus variegatus* sea urchin embryo. *Development*, *148*(19), dev198614. https://doi.org/10.1242/dev.198614

Miyamoto, N., Nakajima, Y., Wada, H., & Saito, Y. (2010). Development of the nervous system in the acorn worm *Balanoglossus simodensis*: Insights into nervous system evolution: Development of hemichordate nervous system. *Evolution & Development*, *12*(4), 416–424. https://doi.org/10.1111/j.1525-142X.2010.00428.x

Morgan, T. H. (1891). The growth and metamorphosis of Tornaria. *Journal of Morphology*, *5*,

407–358.

Myngbay, A., Manarbek, L., Ludbrook, S., & Kunz, J. (2021). The Role of Collagen Triple Helix

Repeat-Containing 1 Protein (CTHRC1) in Rheumatoid Arthritis. *International Journal of*

Molecular Sciences, *22*(5). https://doi.org/10.3390/ijms22052426

Nakajima, Y., Humphreys, T., Kaneko, H., & Tagawa, K. (2004). Development and Neural

Organization of the Tornaria Larva of the Hawaiian Hemichordate, *Ptychodera flava*.

Zoological Science, *21*(1), 69–78. https://doi.org/10.2108/0289-

0003(2004)21%255B69:DANOOT%255D2.0.CO;2

Nielsen, C. (1998). Origin and evolution of animal life cycles. *Biological Reviews*, *73*(2), 125–

155. https://doi.org/10.1111/j.1469-185X.1997.tb00027.x

Nielsen, C. (2005). Larval and adult brains. *Evolution & Development*, *7*(5), 483–489.

https://doi.org/10.1111/j.1525-142X.2005.05051.x

Nielsen, C., & Hay-Schmidt, A. (2007). Development of the enteropneust *Ptychodera flava*:

Ciliary bands and nervous system. *Journal of Morphology*, *268*(7), 551–570.

https://doi.org/10.1002/jmor.10533

Paganos, P., Ullrich-Lüter, J., Almazán, A., Voronov, D., Carl, J., Zakrzewski, A.-C., Zemann,

B., Rusciano, M. L., Sancerni, T., Schauer, M., Akar, O., Caccavale, F., Cocurullo, M.,

Benvenuto, G., Croce, J. C., Lüter, C., & Arnone, M. I. (2025). *Single Nucleus Profiling*

Highlights the All-Brain Echinoderm Nervous System.

https://doi.org/10.1101/2025.03.24.644250

Paganos, P., Voronov, D., Musser, J. M., Arendt, D., & Arnone, M. I. (2021). Single-cell RNA

sequencing of the Strongylocentrotus purpuratus larva reveals the blueprint of major cell

types and nervous system of a non-chordate deuterostome. *eLife*, *10*, e70416.

https://doi.org/10.7554/eLife.70416

Pardos, F., & Benito, J. (1988). Blood Vessels and Related Structures in the Gill Bars of

Glossobalanus minutus (Enteropneusta). *Acta Zoologica*, *69*(2), 87–94.

https://doi.org/10.1111/j.1463-6395.1988.tb00905.x

Park, J.-O., Pan, J., Möhrlen, F., Schupp, M.-O., Johnsen, R., Baillie, D. L., Zapf, R., Moerman,

D. G., & Hutter, H. (2010). Characterization of the astacin family of metalloproteases in

C. elegans. *BMC Developmental Biology*, *10*(1), 14. https://doi.org/10.1186/1471-213X-

10-14

Patry, W. L., Bubel, M., Hansen, C., & Knowles, T. (2020). Diffusion tubes: A method for the

mass culture of ctenophores and other pelagic marine invertebrates. *PeerJ*, *8*, e8938.

https://doi.org/10.7717/peerj.8938

Perillo, M., Oulhen, N., Foster, S., Spurrell, M., Calestani, C., & Wessel, G. (2020). Regulation

of dynamic pigment cell states at single-cell resolution. *eLife*, *9*, e60388.

https://doi.org/10.7554/eLife.60388

Peterson, K. J., Cameron, R. A., & Davidson, E. H. (1997). Set-aside cells in maximal indirect

development: Evolutionary and developmental significance. *BioEssays*, *19*(7), 623–631.

https://doi.org/10.1002/bies.950190713

Pfeffer, P. L., Gerster, T., Lun, K., Brand, M., & Busslinger, M. (1998). Characterization of three

novel members of the zebrafish Pax2/5/8 family: Dependency of Pax5 and Pax8

expression on the Pax2.1 (noi) function. *Development*, *125*, 3063–3074.

Pham, K., & Hobert, O. (2019). Unlike Drosophila elav, the C. elegans elav orthologue exc-7 is

not panneuronally expressed. *microPublication Biology*, *2019*.

https://doi.org/10.17912/micropub.biology.000189

Piovani, L., Leite, D. J., Guerra, L. A. Y., Simpson, F., Musser, J. M., Salvador-Martínez, I.,

Marlétaz, F., Jékely, G., & Telford, M. J. (2023). Single-cell atlases of two

lophotrochozoan larvae highlight their complex evolutionary histories. *Science*

Advances.

Robinow, S., & White, K. (1991). Characterization and spatial distribution of the ELAV protein

during Drosophila melanogaster development. *Journal of Neurobiology*, *22*(5), 443–461.

https://doi.org/10.1002/neu.480220503

Ruppert, E. E., Cameron, C. B., & Frick, J. E. (1999). Endostyle-like Features of the Dorsal

Epibranchial Ridge of an Enteropneust and the Hypothesis of Dorsal-Ventral Axis

Inversion in Chordates. *Invertebrate Biology*, *118*(2), 202–212. JSTOR.

https://doi.org/10.2307/3227061

Rychel, A. L., Smith, S. E., Shimamoto, H. T., & Swalla, B. J. (2006). Evolution and

development of the chordates: Collagen and pharyngeal cartilage. *Molecular Biology*

and Evolution, *23*(3), 541–549.

Samuelson, L. C., Wiebauer, K., Gumucio, D. L., & Meisler, M. H. (1988). Expression of the

human amylase genes: Recent origin of a salivary amylase promoter from an actin

pseudogene. *Nucleic Acids Research*, *16*(17), 8261–8276. PubMed.

https://doi.org/10.1093/nar/16.17.8261

Sánchez Alvarado, A., & Yamanaka, S. (2014). Rethinking Differentiation: Stem Cells,

Regeneration, and Plasticity. *Cell*, *157*(1), 110–119.

https://doi.org/10.1016/j.cell.2014.02.041

Satija, R., Farrell, J. A., Gennert, D., Schier, A. F., & Regev, A. (2015). Spatial reconstruction of

single-cell gene expression data. *Nature Biotechnology*, *33*(5), 495–502.

https://doi.org/10.1038/nbt.3192

Sebé-Pedrós, A., Chomsky, E., Pang, K., Lara-Astiaso, D., Gaiti, F., Mukamel, Z., Amit, I.,

Hejnol, A., Degnan, B. M., & Tanay, A. (2018). Early metazoan cell type diversity and the

evolution of multicellular gene regulation. *Nature Ecology & Evolution*.

https://doi.org/10.1038/s41559-018-0575-6

Sebé-Pedrós, A., Saudemont, B., Chomsky, E., Plessier, F., Mailhé, M.-P., Renno, J., Loe-Mie,

Y., Lifshitz, A., Mukamel, Z., Schmutz, S., Novault, S., Steinmetz, P. R. H., Spitz, F.,

Tanay, A., & Marlow, H. (2018). Cnidarian Cell Type Diversity and Regulation Revealed by Whole-Organism Single-Cell RNA-Seq. *Cell*, *173*(6), 1520-1534.e20. https://doi.org/10.1016/j.cell.2018.05.019

Simakov, O., Kawashima, T., Marlétaz, F., Jenkins, J., Koyanagi, R., Mitros, T., Hisata, K., Bredeson, J., Shoguchi, E., Gyoja, F., Yue, J. X., Chen, Y. C., Freeman, R. M., Sasaki, A., Hikosaka-Katayama, T., Sato, A., Fujie, M., Baughman, K. W., Levine, J., … Gerhart, J. (2015). Hemichordate genomes and deuterostome origins. *Nature*, *527*(7579), 459–465. https://doi.org/10.1038/nature16150

Sogabe, S., Hatleberg, W. L., Kocot, K. M., Say, T. E., Stoupin, D., Roper, K. E., Fernandez-Valverde, S. L., Degnan, S. M., & Degnan, B. M. (2019). Pluripotency and the origin of animal multicellularity. *Nature*, *570*(7762), 519–522. https://doi.org/10.1038/s41586-019-1290-4

Strathmann, R., & Bonar, D. (1976). Ciliary feeding of tornaria larvae of *Ptychodera flava* (Hemichordata: Enteropneusta). *Marine Biology*, *34*(4), 317–324. https://doi.org/10.1007/BF00398125

Strathmann, R. R. (2020). Multiple origins of feeding head larvae by the Early Cambrian. *Canadian Journal of Zoology*, *98*(12), 761–776. https://doi.org/10.1139/cjz-2019-0284

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y., Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell Data. *Cell*, *177*(7), 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031

Tagawa, K., Humphreys, T., & Satoh, N. (1998). Novel pattern of Brachyury gene expression in hemichordate embryos. *Mechanisms of Development*, *75*(1–2), 139–143. https://doi.org/10.1016/S0925-4773(98)00078-1

Tarashansky, A. J., Xue, Y., Li, P., Quake, S. R., & Wang, B. (2019). Self-assembling manifolds in single-cell RNA sequencing data. *eLife*, *8*, e48994. https://doi.org/10.7554/eLife.48994

Taylor, P. R., Martinez-Pomares, L., Stacey, M., Lin, H.-H., Brown, G. D., & Gordon, S. (2005).

Macrophage receptors and immune recognition. *Annual Review of Immunology*, *23*,

901–944. https://doi.org/10.1146/annurev.immunol.23.021704.115816

Truman, J. W., Price, J., Miyares, R. L., & Lee, T. (2023). Metamorphosis of memory circuits in

Drosophila reveals a strategy for evolving a larval brain. *eLife*, *12*, e80594.

https://doi.org/10.7554/eLife.80594

Vasta, G. R., Amzel, L. M., Bianchet, M. A., Cammarata, M., Feng, C., & Saito, K. (2017). F-

Type Lectins: A Highly Diversified Family of Fucose-Binding Proteins with a Unique

Sequence Motif and Structural Fold, Involved in Self/Non-Self-Recognition. *Frontiers in*

Immunology, *8*, 1648–1648. PubMed. https://doi.org/10.3389/fimmu.2017.01648

Wray, G. A. (2000). The evolution of embryonic patterning mechanisms in animals. *Seminars in*

Cell and Developmental Biology, *11*(6), 385–393.

https://doi.org/10.1006/scdb.2000.0191

Yoder, B. K., Hou, X., & Guay-Woodford, L. M. (2002). The Polycystic Kidney Disease Proteins,

Polycystin-1, Polycystin-2, Polaris, and Cystin, Are Co-Localized in Renal Cilia. *Journal*

of the American Society of Nephrology, *13*(10), 2508.

https://doi.org/10.1097/01.ASN.0000029587.47950.25

Zhao, J., Wang, G., Ming, J., Lin, Z., Wang, Y., The Tabula Microcebus Consortium, Agarwal,

S., Agrawal, A., Al-Moujahed, A., Alam, A., Albertelli, M. A., Allegakoen, P., Ambrosi, T.,

Antony, J., Artandi, S., Aujard, F., Awayan, K., Baghel, A., Bakerman, I., … Yang, C.

(2022). Adversarial domain translation networks for integrating large-scale atlas-level

single-cell datasets. *Nature Computational Science*, *2*(5), 317–330.

https://doi.org/10.1038/s43588-022-00251-y

Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B.,

Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L.,

Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W.,

1100   Hindson, C. M., … Bielas, J. H. (2017). Massively parallel digital transcriptional profiling

1101    of single cells. *Nature Communications*, *8*(1), 14049.

1102    https://doi.org/10.1038/ncomms14049

1103 Zheng, Y., Blair, D., & Bradley, J. E. (2013). Phyletic Distribution of Fatty Acid-Binding Protein

1104    Genes. *PLoS ONE*, *8*(10), e77636. https://doi.org/10.1371/journal.pone.0077636

1105 Zibetti, C., Adamo, A., Binda, C., Forneris, F., Toffolo, E., Verpelli, C., Ginelli, E., Mattevi, A.,

1106    Sala, C., & Battaglioli, E. (2010). Alternative splicing of the histone demethylase

1107    LSD1/KDM1 contributes to the modulation of neurite morphogenesis in the mammalian

1108    nervous system. *The Journal of Neuroscience : The Official Journal of the Society for*

1109    *Neuroscience*, *30*(7), 2521–2532. https://doi.org/10.1523/JNEUROSCI.5500-09.2010

1110

Figure 1

A

Cnidaria | Echinodermata | Hemichordata | Cephalochordata | Vertebrata | Tunicata | Pancrustacea | Mollusca | Brachiopoda | Phoronida | Annelida | Nemertea

B **C** **D** **E** **F**

G
Apical Organ
Protocoel
Circumoral Ciliary Band
Telotroch

H
Protocoel
Mesocoel
Metacoel

I

J
Probocis
Gill slits
Collar
Trunk

K

☐ Early larva   ☐ Late larva   ☐ Metamorphosis   ☐ Early juvenile   ☐ Late juvenile

Figure 2

A

UMAP 1 / UMAP 2

B

Cell Class

% gene max
1.00
0.75
0.50
0.25
0.00

Cell Classes (B, y-axis): Cartilage, Ciliary Band, Endoderm, Epidermis, Immune, Mesoderm, Muscle, Neural, Pigment, Proliferative, Secretory, Unknown

Marker Genes, Grouped by Cell Class

Marker gene groups: Cartilage, Ciliary Band, Endoderm, Epidermis, Immune, Mesoderm, Muscle, Neural, Pigment, Proliferative, Secretory

Marker genes: foxl, co9a1, pk1l2-32, pk1l2, cnbd2, foxA, sap-4, sata4, fzd1/2/7, fabp6, foxb2, ftbp3-5, cntn4, cana-1, cptk1-4, casp1-5, cthr, catz-1, mkx, mox, unc22-1, mhc, mylk-3, myl1, th, syt1, sturn, plurs, nas15, fbn1-like, fez2, elav, acha6, white-1, opn4-4, nckx5-3, plf1-7, lbr-1/2, dlg5, muc2, jhamt-6, arsi, amy1

Cell Classes

- ● Cartilage (19)
- ● Ciliary Band (6,11)
- ● Endoderm (0,7,8,9,21,27,35,53)
- ● Epidermis (4,13,25,52)
- ● Immune (15,17,20,22,23,29,37,42,46,47)
- ● Mesoderm (3,30)
- ● Muscle (5,18)
- ● Neural (1,2,10,16,26,28,33,34,38,39,40)
- ● Pigment (51)
- ● Proliferative (14,24)
- ● Secretory (12,32,36,49,55)
- ● Unknown (31,41,43,44,45,48,50,54)

C Endoderm — *foxA - early larva*

D *foxA - early juvenile*

E *fapb6 - early larva*

F Cartilage — *co9a1 - late larva*

G *co9a1 - early juvenile*

H Mesoderm — *mox - early larva*

I *mox - late larva*

J Muscle — *myl - metamorphosis*

K *myl - early juvenile*

L Immune — *cthr - early larva*

M Neural — *syt1 - early larva*

N *TH - early larva*

O *TH - early juvenile*

P Ciliary Band — *pk1l2 - early larva*

Q *cnbd2 - early larva*

R Secretory — *muc2 - early juvenile*

S *amy1 - early larva*

T *arsi - late larva*

U *arsi - metamorphosis*

V *arsi - early juvenile*

Figure 3

A

UMAP 1
UMAP 2

B

Lifestage Contribution

100%
75%
50%
25%
0%

Cluster Number

0 17 11 4 48 35 33 9 22 28 26 6 30 46 49 47 51 16 14 34 32 44 24 20 21 8 45 39 43 50 29 31 19 3 10 36 42 40 5 23 12 7 18 31 15 27 2 52 41 53 54 55

Life Stage
Early larva | Late larva | Metamorphosis | Early juvenile | Late juvenile

C

Ci Cluster 26
acha6 - early larva

Cii Cluster 34
nas15 - early larva

D

Di Cluster 1
stum - early juvenile

Dii Cluster 2
fibrilin-2-like - early juvenile

E

Ei Cluster 16
pburs - early larva

Eii
pburs - early juvenile

F

Fi
mhc - early larva

Fii
mhc - early juvenile

Gi 48 hour EdU Pulse
Nuclei, EdU

Gii 43 day EdU Chase
Nuclei, EdU

Early Larva

Hi 48 hour EdU Pulse
Nuclei, EdU

Hii 26 day EdU Chase
Nuclei, EdU

Late Larva

Figure 4

Figure 5

Supplemental Figure 1

Supplemental Figure 2

A
Endoderm

B

C
Cluster 9
Cluster 0
fzd1/2/7, fapb6

D
Cartilage

E

F
co9a1- metamorphosis

Supplemental Figure 3

A   Mesoderm

B

		mox	mkx	mys-2
3	Mesoderm			
30				
other	other			

C   Muscle

D

		popd3-2	calms-13	tnni4	myl1	mlrm	mylk-3	mhc	tpm	acm3-3	titin-4	titin-5	titin-3	titin-1	speg	titin-2	smtn	unc22-1
5	Muscle																	
18																		
other	other																	

E

titin-4   *popd3-2*   *myl1*

F

mhc - metamorphosis   *mhc - late juvenile*

Supplemental Figure 4

Supplemental Figure 5

A
Neural

B

C
GAD   Vacht

D
pburs

E
Ciliary Band

F
Ciliary Band
other

G
plk1l2 - juvenile inset

Supplemental Figure 6

A

Secretory

B

D

Epidermis

E

Supplemental Figure 7

Supplemental Figure 8

Mhc

XP_035686563.1
myosin-7-like
[Branchiostoma floridae]

XP_017208743.1
myosin heavy chain, fast skeletal muscle
[Danio rerio]

AAI50740.1
Myosin, heavy polypeptide 1, skeletal muscle, adult
[Mus musculus]

NP_005954.3
myosin-1
[Homo sapiens]

XP_038064641.1
myosin heavy chain, striated muscle-like isoform X1
[Patiria miniata]

XP_022093558.1
myosin heavy chain, striated muscle-like
[Acanthaster planci]

SCA03015
Mhc
[Schizocardium californicum]

NP_001161591.1
myosin heavy chain protein
[Saccoglossus kowalevskii]