Authors

Eveliina Hanski, Alise J Ponsero, Kaija-Leena Kolho, Willem M de Vos, Anne Salonen

Title

Infant gut microbiota in multi-child families converges with adults without a sibling-specific

signature

Abstract

The human gut microbiota undergoes rapid development during early life. Sibling presence has

been associated with altered infant microbiota composition and accelerated maturation, but

distinguishing direct microbial transmission between siblings from confounding by maternal

reproductive history (parity) remains challenging. We analysed >5,000 faecal samples from >800

mother-father-infant triads in the Finnish HELMi birth cohort to characterise microbiota patterns

that could inform underlying mechanisms. While sibling presence was associated with

accelerated microbiota maturation and modest compositional differences, infants who both had

siblings (or both lacked siblings) were no more similar to each other than mixed pairs. Instead,

infants with siblings exhibited non-specific convergence with adult microbiota profiles and

contributed fewer unique genera to their family metacommunities (infant–mother–father triads).

Multi-child families harboured smaller but compositionally distinct metacommunities. These

patterns challenge simple models of direct sibling transmission and suggest that multiple factors

associated with family structure may jointly influence infant microbiota development, though

specific mechanisms remain unclear.

Introduction

The human gut microbiota is a complex and dynamic community of microorganisms that plays a crucial role in health through its involvement in nutrient absorption, immune system functioning, and metabolic regulation[1]. The early-life microbiota is particularly important because it actively contributes to the development of critical host systems[2], such that alterations during this period can have lasting health effects that persist even after initial microbiota differences have been resolved[3]. As such, understanding how multiple factors influence microbiota development in early life is critical for both immediate and long-term health outcomes.

The establishment of the gut microbiota begins at birth and continues through early life, driven by microbial transmission as well as selective forces posed by environmental factors (e.g., delivery mode, diet, antibiotics[4]) and host physiology (e.g., immune system[5]). While initial colonisation occurs through maternal transmission from vaginal and gut microbiota[6], ongoing microbial exchange between individuals becomes increasingly important as infants develop and interact more broadly with their environment. This social transmission process is facilitated by the highly individualised nature of the human gut microbiota: each person hosts a unique microbial community that can both influence and be influenced by the microbiota of others[7–9]. Growing evidence demonstrates that such social transmission can significantly shape microbial community development[6,10,11].

Within this framework of social transmission, sibling presence represents a particularly important influence on infant gut microbiota development[4,12–15]. Studies have documented that sibling presence is associated with accelerated microbiota maturation[4,12]. This might be because

siblings are still undergoing microbiota development themselves, creating a temporally shifting

pool of microbes, which may expose the index infant to microbes from later stages of microbiota

development and as such accelerate the process of microbiota maturation. Additional factors may

also explain 'sibling effects' on infant microbiota development. For example, siblings who attend

day care or school expose the index infant to microbes from a wider circle of children beyond

just their immediate household[11]. Further, the social dynamics of infant–infant microbial

transmission may differ from infant–adult interactions: children may share toys and engage in

close physical play, creating opportunities for microbial exchange that may be distinct from

interactions with adults. Finally, siblings may also influence the index child's behaviour in ways

that could affect the gut microbiota. For instance, the infant may imitate their older siblings'

behaviours, such as trying new foods, which could shape the infant's diet[16] and subsequently

influence their gut microbiota.

However, in many populations, the presence of siblings is closely correlated with parity (number

of previous late-stage pregnancies), creating a fundamental challenge in interpreting these

findings. Parity and broader aspects of maternal reproductive history (including miscarriages and

type of previous childbirths) may independently influence infant microbiota through distinct

biological pathways. Previous pregnancies and deliveries could alter maternal immune function

and hormonal profiles, potentially changing maternal microbiota composition that is then

transmitted to the infant. Supporting this, parity has been shown to significantly alter both

maternal vaginal and gut microbiota composition[17–19], suggesting previous pregnancies may

induce 'ecological memory' that modulates how the maternal microbiota responds to subsequent

pregnancies. Beyond these biological pathways, parenting behaviours may differ between first-

and subsequent-born children in ways that shape microbial exposures, such as more cautious

hygiene practices among first-time parents[20].

While studies have identified both maternal parity and sibling presence as potential factors

influencing infant gut microbiota development, the extent to which these factors operate

independently or interactively remains unclear. This distinction is particularly important because

sibling-associated microbiota changes have been linked to health outcomes, namely reduced risk

of allergic diseases. However, findings have been inconsistent across different allergic outcomes

and ages of assessment: studies report protective effects of sibling presence against food allergy

at 1 year[12] and asthma at 6 years[15] while others find no significant associations between sibling

presence and food allergy at 1–2 years[14] or multiple allergic conditions at 5 years[21] despite

documented microbiota changes. This conflicting evidence may partly reflect the difficulty in

separating these interrelated pathways, heterogeneity in outcomes, and developmental timing,

and/or varying sample sizes and statistical power, highlighting the need for approaches that can

distinguish direct sibling effects from confounding by maternal reproductive history.

In this study, we use a large longitudinal birth cohort to investigate patterns of infant microbiome

development associated with parity and sibling presence. Using triadic sampling (mother–father–

infant) from over 800 families in the Finnish Health and Early Life Microbiota (HELMi)

cohort[22], we characterise compositional differences, maturation patterns, and family-level

microbiota structures associated with parity/sibling presence. We examine signatures consistent

with social transmission (such as within-family microbiota similarity and associations with both

parental microbiota) alongside patterns that could reflect maternal reproductive history (such as

timing-dependent effects in mothers). By combining >1,200 cross-sectional parental and >6,000

longitudinal infant microbiota profiles from the first two years of life with comprehensive family

and health metadata, we aim to characterise the nature and extent of parity/sibling -associated

microbiota differences across the family unit.

Methods

Cohort data

A total of 1,055 families were recruited to the Health and Early Life Microbiota (HELMi) cohort

(NCT03996304) during 2016–2018 from the general population in the Helsinki metropolitan

area, Finland. Only healthy, singleton babies born healthy on gestational weeks 37–42 with a

body weight >2.5 kg were included in the cohort. Recruitment and data collection are described

in detail in Korpela et al., 2019[22]. Briefly, faecal samples were collected from both parents at a

single timepoint shortly before or after delivery and at several timepoints from infants during the

first two years (at 3, 6, and 12 weeks, as well as 6, 9, 12, 18, and 24 months). Electronic

questionnaires were used to collect extensive data on nutrition, growth, health, environment,

lifestyle, and parental factors at one or multiple timepoints (depending on the question),

including information on maternal parity and sibling household composition. Information related

to delivery (mode of delivery, administration of intrapartum antibiotics, maternal group B

streptococcus status, and timing of membrane rupture) was acquired from the hospital records.

Additional detailed data on reproductive history and sibling characteristics (including delivery

details and potential changes in living arrangements) were collected in May 2023 using an online

questionnaire. Out of all 1,055 families, 346 families participated in this additional data

collection. In the present study, for analyses involving infants only vaginally delivered infants

(*n*=844) were included. Out of these families, 293 participated in the additional data collection.

Collected data is described in detail in Jokela et al., 2023[13].

Ethics statement

The HELMi study was approved by The Hospital District of Helsinki and Uusimaa and

conducted in accordance with the principles of the Helsinki Declaration. Participating parents

signed an informed consent at enrolment and retained the right to withdraw at any point.

DNA extraction, library preparation, and amplicon sequencing

Faecal sample processing of HELMi samples for microbiota analysis is described in detail in

Jokela et al. 2023[13]. In brief, DNA was extracted from the faecal samples using a repeating bead-

beating step, followed by amplicon sequencing targeting the V3–V4 region of the 16S ribosomal

RNA gene. Amplicons were sequenced using Illumina MiSeq and HiSeq sequencing technology

at the Functional Genomics Unit and Institute for Molecular Medicine Finland, University of

Helsinki (Helsinki, Finland).

Data processing

Data processing and subsequent analyses were conducted using R v4.1.2. Demultiplexed

sequencing reads were processed through the DADA2 pipeline[23]. Taxonomy was assigned using

the SILVA v.138.1 database[24]. Only ASVs assigned to kingdom Bacteria were included, and

ASVs assigned to order Chloroplast or family Mitochondria were excluded. The presence of any

potential contaminants using the R package *decontam*[25] using the `prevalence` method, in

which the prevalence of each sequence in biological samples is compared to that in negative

controls. An ASV was considered a contaminant if it reached a probability of 0.1 in the Fisher's

exact test used in *decontam*. 15 ASV contaminants were identified and filtered from the data

before subsequent analysis. The package *iNEXT* was used to generate sample type and

sequencing platform -specific sample completeness and rarefaction curves, the plateau points of

which were used to determine a threshold for minimum read depth for each sample type across

the two different sequencing platforms. Samples with a sequencing depth under this threshold

were excluded, resulting in a dataset of 1009 unique families. Taxa were then aggregated at

genus level to account for variable sequencing quality and platform differences, followed by

measurement of asymptotic genus-level richness and Shannon diversity using *iNEXT*[26], which is

applicable across taxonomic resolutions, including OTUs, ASVs, and higher taxa, and thus suited

our genus-level data structure. The R package *phyloseq*[27] was then used to normalise genus

counts to proportional abundances.

Analysis

All analyses were conducted in R version 4.3.1. Parity/sibling presence (highly correlated) was

assessed qualitatively unless stated otherwise. To assess overall compositional differences, we

performed permutational (*n*=999) multivariate analysis of variance (PERMANOVA) using

genus-level Jaccard dissimilarity matrices using `adonis2` function from R package *vegan*[28].

Models were fitted separately for each timepoint, adjusting for relevant technical and biological

covariates. Homogeneity of dispersion between groups was tested using `betadisper`

function. To identify taxa distinguishing infants from different family structures (siblings vs no

siblings), we trained Random Forest classification models for each timepoint using genus-level

relative abundances (*randomForest* R package[29]). Model performance was evaluated using

classification accuracy. Variable importance was assessed using mean decrease Gini index.

To test whether infants with similar family structures showed greater microbiota similarity, we

calculated pairwise Jaccard dissimilarities between all infant pairs at each timepoint. Bayesian

regression models (using function `brm` from R package *brms*[30]) tested whether dyadic sibling

concordance (both with siblings, both without siblings, or discordant) predicted microbiota

similarity, adjusting for covariates.

Microbiota maturation

To assess whether sibling presence was associated with the maturation of the gut microbiota in

early life, we predicted infant age using a Random Forest regression model (R package

randomForest[29]) trained on genus-level relative abundances. Sequencing platform was strongly

confounded with infant age (early timepoints predominantly sequenced on HiSeq, later

timepoints on MiSeq). To account for this technical variation, sequencing platform was included

as a binary predictor variable (MiSeq vs HiSeq) in the Random Forest model, allowing the model

to learn age-related patterns while adjusting for batch effects. The model's predictive

performance was assessed using 5-fold cross-validation, where each sample was used for testing

exactly once. To test parity effects on microbiota maturation, we fitted a generalised additive

model (GAM) using the *mgcv* R package. Here, estimated microbiota age was used as the

response variable. A smooth term for chronological age was allowed to vary by sibling presence,

testing whether developmental trajectories differed between groups. The model also included

other predictors, such as sequencing platform and type of feeding. Family ID was included as a

random effect to account for repeated measures. Additional models tested dose-response

relationships by restricting analyses to families with 1–2 older siblings and testing whether

sibling number or youngest sibling age predicted microbiota age.

To identify when during development parity effects on microbiota age were most pronounced,

we fitted `brm` models (*brms* package[30]) separately for each timepoint, with microbiota age as the

outcome and sibling presence (yes/no) as the predictor, adjusting for covariates. To identify key

taxa underlying parity-associated maturation differences, we used a 'drop-one-taxon' approach

where each genus was dropped at a time and the age prediction model was refit[31]. Genera whose

removal substantially degraded model performance were considered key contributors to age

prediction. Age trajectories of these key genera were then visualised using GAMs, with smooth

terms for chronological age allowed to vary by sibling presence.

Microbiota convergence with adult samples

Microbiota similarity between infant–parent dyads was calculated using Jaccard dissimilarity at

each infant timepoint (parental samples were collected around the time of the birth of the index

child such that infant–parent sample pairs were not temporally matched). Bayesian regression

models (using function `brm` from *brms*[30]) tested whether parity (continuous: number of

previous deliveries/siblings) predicted infant–parent similarity, adjusting for covariates. Models

were fitted separately for each infant timepoints and for infant–mother and infant–father dyads.

To test whether convergence reflected similarity to own parents versus adults generally, we

compared infant–adult dyads from the same vs different families, testing whether family

membership and family composition concordance (both multi-child, both single-child, or mixed)

predicted similarity).

For all `brm` models, posterior checks were conducted to confirm reliable model performances.

Here, we ensured that the chains had converged, Rhat values were <1.05, bulk effective sample

sizes at least 100 times the number of chains and no smaller than 10% of total posterior draws,

and that the sampler took small enough steps to avoid excess (>10) divergent transitions after

warm-up. Additionally, posterior predictive checks were performed to assess model fit,

confirming that the simulated posterior distributions closely matched the observed data, with

only minor deviations in certain regions.

Metacommunity analyses

For each infant–mother–father triad, we calculated the total number of unique genera (family

metacommunity size) and the proportion of genera unique to each family member (this analysis

was repeated for each infant sample type). Differences between multi-child and single-child

families were tested at each timepoint using Wilcoxon rank sum tests with FDR correction across

timepoints. To test overall effects across the developmental trajectory, we fitted GAMs with the

proportion of unique genera as the outcome, including smooth terms for infant age (allowing

trajectories to vary by parity group) and a parametric term for parity, with family ID as a random

effect (*mgcv* R package). Separate models were fitted for infant, mother, and father proportions.

To test whether multi-child family metacommunities represent subsets of single-child family

metacommunities, pairwise Jaccard dissimilarity between triads was decomposed into nestedness

and turnover components using the *betapart* R package[32]. For each timepoint, dissimilarities

were compared within vs across parity groups using Wilcoxon rank sum tests (FDR corrected

across timepoints). To identify specific genera contributing to parity-associated metacommunity

differences, genus-level presence frequencies were compared between parity groups at each

timepoint using Fisher's exact tests. P-values were adjusted using FDR correction across all

genus–timepoint combinations.

Results

Sibling presence is associated with subtle compositional differences but does not create

microbial homogenisation among infants

During the first weeks of life (at 3 and 6 weeks), sibling presence (highly correlated with

maternal parity; assessed qualitatively) showed no significant association with genus richness (as

indicated by credible intervals (CIs) overlapping zero in Bayesian regression models (brm)).

Positive associations were observed at 3 months (posterior mean 0.27, CIs 0.13 to 0.42), 6

months (posterior mean 0.14, CIs 0.04 to 0.25), 18 months (posterior mean 0.09, CIs 0.02 to

0.16), and 2 years (posterior mean 0.09, CIs 0.01 to 0.16), with associations at the latter

timepoints being marginal. No significant associations were observed at 9 months or 1 year

(Table SX). Shannon diversity (reflecting community evenness) showed no significant

associations with sibling presence at any timepoint.

To assess whether sibling presence was associated with overall shifts in community composition,

we performed multivariate marginal permutational ANOVA (PERMANOVA) analyses on

Jaccard distance across infant timepoints, adjusting for technical and biological covariates

(covariates were determined separately for each infant timepoint based on whether they predicted

microbiota variation at that timepoint). Sibling presence showed significant associations with

genus-level community composition at all timepoints, though the proportion of explained

variance was small ($R^2<1.0\%$ and $p<0.005$ for all) and significantly different within-group

variances were observed for several timepoints (3 weeks, 18 months, 24 months). This suggests

that observed results at these timepoints may partly reflect heterogeneity in dispersion rather than

true differences in community structure in infants with or without siblings.

To assess whether specific taxa could reliably distinguish between infants from different family

structures, we trained Random Forest classification models for each timepoint using genus-level

microbiota composition. These models achieved modest accuracy (56–71% across timepoints) in

distinguishing infants from single-child vs multi-child families. Although classification

performance was limited, certain taxa – particularly *Bifidobacterium, Flavonibacterium,* and

Clostridium sensu stricto 1 – consistently ranked highly in importance in early, mid-, and late

timepoints, respectively, suggesting a potential temporal shift in the taxa most strongly

associated with siblings.

Whilst PERMANOVA indicated subtle compositional differences between groups, this does not

establish whether infants with siblings share a distinctive 'sibling microbiota' signature. To test

whether infants from families with similar structures showed greater microbial similarity to each

other than to infants from different family structures, we compared microbiota composition

between infant pairs across families at each timepoint, examining whether dyads with concordant

sibling background (where both had or both lacked siblings) were more similar than dyads with

discordant sibling background. Across all timepoints, dyadic sibling background (yes–yes / no–

no / yes–no) was not a significant predictor of microbial similarity (brm models).

To assess whether parity/sibling -associated microbiota patterns extended beyond infants, we

tested for compositional differences in maternal and paternal samples. In fathers, sibling

presence was significantly associated with microbiota composition (PERMANOVA; R^2=0.004,

F=1.793 p=0.011; beta dispersion F=2.274, p=0.139; n=552). In mothers, however, sibling

presence effects depended on sampling time relative to birth (no such effect observed in fathers).

When pre- and post-birth maternal samples were analysed together, no significant association

was detected (R^2=0.002, F=1.417, p=0.071; beta dispersion F=1.827, p=0.196; n=701), though

significant differences in within-group dispersion were observed between pre- and post-birth

samples (beta dispersion F=7.848, p=0.004). Stratifying by sampling time revealed that sibling

presence effects were only detectable in samples collected after giving birth (R^2=0.008, F=1.780

p=0.008; beta dispersion F=0.783, p=0.381; n=488) but not before birth (R^2=0.003, F=1.453,

p=0.045; beta dispersion F=9.042, p=0.005; n=213).

Parity accelerates microbiota maturation

To assess whether parity was associated with the rate of gut microbiota maturation in early life,

we predicted infant age using a Random Forest regression model trained on genus-level relative

abundances across eight timepoints from birth to 2 years. The predicted 'microbiota age' was

strongly correlated with chronological age (cross-validated R^2=0.853, RMSE=6.092 months; Fig.

1A). Sequencing platform was included as a predictor to account for strong confounding with

infant age (Table S1) and ranked 19[th] out of the 77 features in variable importance, with the top

predictors being genera associated with microbiota maturation (*Anaerostipes, Faecalibacterium,*

Blautia; Table S1).

We hypothesised that infants with siblings would experience acceleration of microbiota

maturation (where microbiota age is higher than expected from chronological age) due to early

exposure to microbiota from older siblings, effectively acquiring microbes representative of later

stages of microbiota development. In support of this, a generalised additive model (GAM)

showed distinct microbiota age trajectories for infants with and without siblings ($p<0.001$ for

both smooth terms; Fig. 1B), indicating a modest acceleration of maturation in infants from

multi-child families (estimate=0.140, $p=0.002$; $n=4,305$ from 808 infants; 1–8 samples per

infant; Table S2). To test whether this effect was modified by the number of older siblings or the

age of the youngest sibling, we fitted additional models. When restricting to infants from

families with one or two other children ($n=1,657$ samples from 304 infants; 249 infants with one

sibling, 55 infants with two siblings), the number of siblings did not significantly predict

microbiota age (estimate=0.107, $p=0.219$). Further, in a subset with data on sibling age, the age

of the youngest older sibling did not significantly predict microbiota age ($n=851$ samples from

147 infants with 1–2 siblings; estimate=0.003, $p=0.123$).

To examine when parity differences in microbiota maturation were most pronounced, we ran

brm models separately for each time point and found that sibling presence (yes/no) was

significantly associated with elevated microbiota age at 6 (posterior mean 0.03, CIs 0.01 to 0.05),

9 (posterior mean: 0.05, CIs 0.01 to 0.09) and 18 months (posterior mean: 0.02, CIs 0.01 to 0.03;

Fig. 1B), although the effects were small and only marginally significant. Other available

measures of social transmission (number of caregivers, day care attendance) did not predict

microbiota age (these predictors were only included in models with infants of 1–2 years). Early-

life feeding was significantly associated with microbiota age at several timepoints such that lack

of breastfeeding (be that use of formula as an alternative or absence of either milk in diet)

accelerated microbiota maturation, with effects consistently larger than those of presence of

siblings.

To better understand which microbial taxa underlie microbiota age predictions, we investigated

the age trajectories of the five genera whose removal most increased prediction error. Several

taxa revealed distinct age trajectories between infants with and without siblings (Fig. 1C).

Notably, *Faecalibacterium* – the second most important genus in predicting microbiota age –

exhibited an earlier and more rapid increase in relative abundance among infants with siblings

compared to those without. While the *Faecalibacterium* trajectory appeared to plateau at around

20% relative abundance in infants from multi-child families, the single-child group showed a

continued upward trend without a clear plateau within the observed sampling window (Fig. 2C).

Figure 1. (**A**) Relationship between chronological age and microbiota-predicted age in infants

aged 3 weeks to 2 years, modelled using a Random Forest regression model trained on genus-

level relative abundance data. Cross-validated R^2=0.853, RMSE=6.092 months. Model

performance was assessed using 5-fold cross-validation, where each sample was used for testing

exactly once. Model stability was further supported by Out-of-Bag (OOB) performance metrics

(average OOB R^2=0.879 ± 0.002, OOB RMSE=8.06 ± 0.09 months). (**B**) Generalised additive

model predictions showing microbiota age trajectories for infants with (blue) and without (red)

older siblings. Lines show fitted values with 95% confidence intervals; points show observed

data. (**C**) Generalised additive model visualising top 5 genera with highest importance scores in

microbiota age prediction in infants from multi-child (solid lines) or single-child (dashed lines)

families.

Microbiota of infants with older siblings converge more with adults

Across infant–parent dyads (infant–mother or infant–father), higher parity/number of siblings

(modelled as a continuous predictor) was significantly associated with greater microbiota

similarity from 3 months of infant age onwards (Fig. 2A, Fig. S1; parental samples were

collected shortly before or after index child's birth, such that samples in dyads were not

temporally matched). At 3 months of age, having 1–2 in contrast to >2 caregivers increased

similarity in infant–parent dyads (brm models; infant–father: posterior mean 0.54, CIs 0.21 to

0.92; infant–mother: posterior mean 0.28, CIs 0.02 to 0.56; families with 1–2 caregivers; families

with >2 caregivers) as expected when fewer competing microbial sources are present. In infant–

father dyads, this smaller number of caregivers increased microbiota similarity also at 3 and 6

weeks of age (3 weeks: posterior mean 0.31, CIs 0.03 to 0.62; 6 weeks: posterior mean: 0.38, CIs

0.05 to 0.74). Day care attendance nor day care group size (only assessed at 1.5 and 2 years of

age when sufficient variation was available) predicted infant–parent similarity.

To assess whether this convergence reflects greater similarity to one's own parent vs adult

microbiotas more generally, we compared infant–adult dyads from the same and different

families. Family composition similarity (both multi-child, both single-child, or mixed)

significantly predicted microbiota similarity at several timepoints across dyads (brm models),

whereas family membership had a detectable effect only at 18 months, and this effect was

weaker than that of parity. This suggests that parity promotes convergence with adult-like

microbial profiles broadly, rather than uniquely increasingly resemblance to the specific parent.

In contrast to infant–parent dyads, parity did not increase microbial similarity in mother–father

dyads (posterior mean: 0.05, CIs -0.02 to 0.12).

We next investigated whether the accelerated microbiota maturation in infants from multi-child

families was associated with acquisition of a broader range of taxa, which could manifest as

larger metacommunities within infant–mother–father triads. Contrary to this, multi-child families

consistently showed smaller metacommunities than single-child families across all timepoints,

although this difference was statistically significant only at 9 months of infant age (Wilcoxon

rank sum test, FDR-corrected $p=0.029$; $p>0.05$ at remaining timepoints; Fig. 2B). The proportion

of genera unique to infants (relative to the family-level metacommunity) was marginally lower in

multi-child families across the developmental trajectory (GAM parametric coefficient=-0.004,

$p=0.050$). A similar pattern was observed for the proportion of genera unique to mothers

(coefficient=-0.008, $p=0.035$), whereas father showed no significant difference

(coefficient=0.002, $p=0.634$). These proportions were calculated at each infant timepoint relative

to the metacommunity of that infant–mother–father triad.

To investigate whether metacommunities in multi-child families represent subsets of the larger

metacommunities observed in single-child families, we decomposed pairwise Jaccard

dissimilarity among infant–mother–father triads into components of nestedness (asymmetric

subset relationships) and turnover (taxa replacement). For each infant age group, we compared

dissimilarities within the same parity group ('within') and across different groups ('between').

Turnover was significantly higher in cross-parity comparisons than within-parity comparisons at

7 of 8 timepoints (Wilcoxon rank sum test, FDR-corrected $p<0.05$; exception at 6 months,

$p=0.876$; Fig. S2A), indicating that triads from different sibling contexts harbour taxonomically

distinct sets of genera throughout development. Turnover declined with infant age in both

comparison types, from approximately 65% at 3 weeks to 50% at 24 months (Fig. S2A),

reflecting developmental convergence toward more similar genus compositions across families.

However, the taxonomic distinctiveness between parity groups persisted despite this overall

convergence. In contrast, nestedness was generally low (median<0.1) and did not differ

significantly between groups, except at 9 months, where a small but statistically significant

difference was detected (FDR-corrected $p=0.033$; Fig. S2B). The higher turnover between

groups indicates that multi-child and single-child family metacommunities contain different sets

of genera rather than one being a subset of the other.

To explore which genera showed the largest differences in occurrence between parity groups, we

examined genus-level presence across infant–mother–father triads, comparing frequency

between parity groups at each timepoint. After correction for multiple comparisons across all

genus-timepoint combinations, only 6 of 3,504 tests remained significant (FDR $p<0.05$). We

visualised the top 30 genera showing the largest average differences in presence frequency

across ages (Fig. 2C). *Akkermansia* was significantly more frequent in multi-child family triads

at 12 months (50% vs 22%, FDR-corrected $p=0.008$), while *Flavonifractor* was significantly less

 frequent in multi-child families at both 12 months (66% vs 91%, *p*=0.007) and 24 months (73%

 vs 93%, *p*=0.007).

 Figure 2. (**A**) Parity effects on infant-mother similarity across infant ages. Points show the

 estimated effect of parity (posterior means with 95% credible intervals (CI)) from separate

 Bayesian regression models fitter at each timepoint. The y-axis represents the change in

 similarity per additional previous pregnancy (i.e., number of older siblings). (**B**) Number of

unique genera in infant–mother–father triads from multi-child (solid boxes) vs single-child

(dashed boxes) families. Number of triads varied from 224 to 319 across the 8 timepoints.

Statistical differences between multi-child and single-child families were tested with Wilcoxon

rank sum tests (FDR corrected). Ns, $p>0.05$. **(C)** Genus-level differences in presence frequency

between multi-child and single-child families across infant ages. For each infant age group,

presence frequency of each genus was compared between triads from multi-child and single-

child families using Fisher's exact tests. Genera shown represent the 30 with the highest average

absolute frequency difference across timepoints. Tile colour indicates the magnitude and

direction of the difference in presence frequency (multi-child – single-child). After correction for

multiple comparisons across all genus-timepoint combinations ($n=3,504$ tests), only 6

comparisons remained significant (FDR $p<0.05$). **, $p<0.01$.

Discussion

In this study, we characterised the effects of parity and sibling presence on infant gut microbiota

development using triadic sampling from over 800 families. While parity/sibling presence

showed significant associations with microbiota composition across infants, mothers, and

fathers, the effects were modest. Infants from multi-child families exhibited accelerated

microbiota maturation and enhanced convergence with adult microbiota, yet infants from similar

family structures did not show increased microbial similarity to one another. Multi-child families

harboured smaller but compositionally distinct metacommunities, with differences driven by

taxonomic turnover rather than loss of diversity. Together, these findings suggest that family

structure influences infant microbiota throughout multiple pathways that may involve both

maternal reproductive biology and household-level factors.

Direct biological effects of parity are expected in mothers, yet parity/sibling-related patterns

were detected in both mothers and fathers. This suggests that at least some parity/sibling-

associated microbiota patterns reflect household-level factors (such as changes in diet[16] or

hygiene practices[20], or social transmission of microbes[10]) that affect the entire family rather than

maternal biology alone. The timing-dependent pattern in mothers (effects only detectable in post-

birth sampled mothers) could reflect pregnancy-related changes in the maternal gut microbiota[33],

though longitudinal sampling would be needed to confirm this.

The modest compositional associations we observed are difficult to reconcile with a

straightforward sibling transmission model. While sibling presence explained small but

significant variance in community composition at all timepoints, suggesting subtle directional

shifts rather than distinctive community structures, these effects did not translate to increased

microbial similarity amongst infants from similar sibling backgrounds. Existing literature has

interpreted associations between sibling presence and infant microbiota composition as evidence

of direct sibling–to–infant microbial transmission[14]. However, without microbiota profiles from

the siblings themselves, such conclusions remain speculative. Observing that infants with

siblings have different microbiota than only children does not establish that siblings are the

source of those microbes – it is equally consistent with confounding by maternal reproductive

history, differences in household practices, or other family-level factors associated with having

multiple children. If siblings were a major source of shared microbes, we might expect infants

with siblings to resemble each other more than they resemble only children. The lack of such

homogenisation in our data suggests either that sibling transmission occurs but involves highly

individualised taxa (reflecting heterogeneity in siblings' own microbiota across different ages

and developmental stages), or that parity/sibling -associated factors beyond direct sibling contact

drive these compositional differences.

We observed modest but significant acceleration of microbiota maturation in infants with

siblings, aligning with previous studies[4,12]. Infants from multi-child families showed elevated

microbiota age, particularly at 6, 9 and 18 months, with *Faecalibacterium* – a key predictor of

microbiota age – exhibiting earlier and more rapid increases. Consistent with accelerated

maturation, these infants also showed increased genus richness at several timepoints. Shannon

diversity showed minimal associations, suggesting compositional shifts involve changes in

presence/absence of taxa rather than substantial restructuring of community evenness. Sibling

transmission appears a plausible mediator of these effects, as older siblings would provide

exposure to both more developmentally advanced microbiota (potentially accelerating

maturation) and a broader range of taxa (increasing richness). However, when we examined

whether this effect scaled with the number of siblings (comparing infants with one vs two

siblings) or with the age of the youngest older sibling, neither factor significantly predicted

microbiota age. While these analyses were limited by sample size and may have lacked statistical

power, the lack of dose-response or age-dependent effects raises questions about direct sibling

transmission. Several alternative mechanisms could contribute to this pattern. Parity-related

changes in maternal gut microbiota[18,19] could facilitate earlier establishment of taxa like

Faecalibacterium through vertical transmission at birth or ongoing post-birth maternal

transmission. Thus, while maturation acceleration and associated richness increases are

consistently associated with sibling presence, the mechanisms remain ambiguous.

Consistent with accelerated maturation, infants from multi-child families also showed increased convergence with adult microbiota profiles. Parity/sibling presence was associated with increased infant–parent similarity from 3 months onwards, yet this convergence occurred with adult microbiota broadly rather than being specific to the infant's own parents. Importantly, mother–father similarity did not increase with parity/sibling presence, suggesting that the observed pattern reflects changes in infant microbiota rather than household-level effects. Adult microbiota is considerably more stable than infant microbiota due to mature immune selection and established gut physiology, making it more resistant to colonisation by exogenous microbes[34]. The non-specific nature of this convergence (where infants become more similar to adults generally rather than their own parents specifically) reinforces that these infants are maturing more rapidly towards adult-like profiles. This pattern could arise if sibling presence increases infant exposure to adult microbiota through expanded social networks (visitors, other parents at activities), facilitating transmission of adult-type taxa that the infant's developing gut can accommodate. Alternatively, parity-related changes in maternal physiology, immunity, or microbiota composition could alter the infant gut environment in ways that favour establishment and persistence of adult-type taxa, either through modified vertical transmission or through programming of selective pressures in the infant gut. These mechanisms are not mutually exclusive, and both could contribute to the observed acceleration of maturation and convergence with adult profiles.

The family-level metacommunity analysis provides additional insight into how family structure shapes microbial communities. Multi-child families harboured smaller metacommunities than

single-child families, with differences driven by taxonomic turnover rather than nestedness,

meaning these families support compositionally different assemblages rather than simply

reduced subsets of single-child family taxa. This reduction in metacommunity size was driven by

infants contributing fewer unique genera relative to their family's metacommunity, while

parental contributions remained similar across family types. This pattern indicates that infants in

multi-child families harbour microbiota that overlap more extensively with their parents'

communities, consistent with the accelerated convergence towards adult-like profiles described

above. Whether this increased overlap reflects more efficient transmission of parental taxa or

these infants developing selective pressures that favour adult-type taxa remains unclear.

However, the observation that specific (even if only a few) genera show substantially different

presence frequencies between family types (e.g., *Akkermansia* more common in multi-child

families, *Flavonifractor* in single-child families) suggests that family structure creates distinct

selective pressures favouring different taxa. If sibling presence primarily increased microbial

diversity through transmission, we might expect multi-child families to harbour additional taxa

on top of those found in single-child families, rather than the compositional turnover and reduced

metacommunity size we observed.

Our findings have implications for interpreting epidemiological associations between sibling

presence and health outcomes. Previous studies have attributed protective effects against

allergies to increased microbial exposure from siblings[12,15], yet our results suggest that such

effects, if real, may operate through pathways beyond direct sibling transmission. If maternal

reproductive history independently shapes infant microbiota – through alterations in maternal

microbial communities, immune factors in breastmilk, or parenting practices – then the causal

pathways to health outcomes may differ substantially from those assumed. This could explain

inconsistent findings regarding sibling presence and allergic outcomes[14,21], as different studies

may capture different aspects of this complex relationship depending on their ability to account

for maternal factors.

Several limitations warrant consideration. While our triadic design provides leverage for

distinguishing household-level from maternal-specific effects, parity and sibling presence remain

correlated, limiting our ability to fully separate their influences. As we lack microbiota profiles

from siblings, we cannot directly assess sibling-to-infant microbial transmission. Our analyses

used 16S rRNA amplicon sequencing data, which provides genus-level resolution but cannot

detect strain-level transmission patterns. Our study is observational, precluding definitive causal

inference about the mechanisms generating observed patterns. While the data comprises >800

family triads, sample sizes for detailed reproductive history analyses were modest, limiting

statistical power for detecting associations with specific pregnancy outcomes such as known

miscarriage. Our cohort is relatively homogenous in socioeconomic status and findings may not

generalise to populations with different environmental and cultural contexts.

In conclusion, while sibling presence is associated with infant microbiota differences, our

findings suggest these effects likely reflect multiple interacting factors rather than primarily

direct sibling transmission. The lack of increased similarity amongst from similar sibling

backgrounds, the non-specific convergence with adult microbiota, and the compositionally

distinct metacommunities in different family contexts point towards more complex mechanisms

that may maternal reproductive history, household practices, and family-level dynamics.

Author information

Affiliations

Eveliina Hanski: Human Microbiome Research Program, University of Helsinki, Helsinki,

Finland; University of Oxford, Oxford, UK; Collegium Helveticum, Zurich, Switzerland

Alise Ponsero: Human Microbiome Research Program, Faculty of Medicine, University of

Helsinki, Helsinki, Finland; Core bioinformatics, Quadram Institute of Biosciences, Norwich UK

Kaija-Leena Kolho: Children's Hospital HUS, University of Helsinki and Faculty of Medicine,

University of Helsinki, Helsinki, Finland

Willem M de Vos: Human Microbiome Research Program, Faculty of Medicine, University of

Helsinki, Helsinki, Finland; Laboratory of Microbiology, Wageningen University, The

Netherlands

Anne Salonen: Human Microbiome Research Program, University of Helsinki, Finland

Contributions

EH and AS conceptualised the presented study. EH and AJP processed the sequencing data. EH

and AJP curated the metadata. EH analysed the data. EH wrote the manuscript. AS, KLK and

WMdV conceived the HELMi cohort, obtained funding, and provided data access. All authors

contributed to the final version of the manuscript.

Data availability

The 16S rRNA amplicon sequencing data from the HELMi birth cohort is publicly available at

https://www.ebi.ac.uk/ena/browser/view/PRJEB55243. Metadata, aerotolerance data, taxonomy

table, ASV relative abundance table as well as R code used in this study will be made publicly

available on GitHub upon publication.

References

1. Krautkramer, K. A., Fan, J. & Bäckhed, F. Gut microbial metabolites as multi-kingdom

intermediates. *Nat. Rev. Microbiol.* **19**, 77–94 (2021).

2. Gensollen, T., Iyer, S. S., Kasper, D. L. & Blumberg, R. S. How colonization by microbiota

in early life shapes the immune system. *Science* **352**, 539–544 (2016).

3. Tamburini, S., Shen, N., Wu, H. C. & Clemente, J. C. The microbiome in early life:

implications for health outcomes. *Nat. Med.* **22**, 713–722 (2016).

4. Stewart, C. J. *et al.* Temporal development of the gut microbiome in early childhood from

the TEDDY study. *Nature* **562**, 583–588 (2018).

5. Sprockett, D., Fukami, T. & Relman, D. A. Role of priority effects in the early-life assembly

of the gut microbiota. *Nat. Rev. Gastroenterol. Hepatol.* **15**, 197–205 (2018).

6. Ferretti, P. *et al.* Mother-to-Infant Microbial Transmission from Different Body Sites Shapes

the Developing Infant Gut Microbiome. *Cell Host Microbe* **24**, 133-145.e5 (2018).

7. Brito, I. L. *et al.* Transmission of human-associated microbiota along family and social

networks. *Nat. Microbiol.* **4**, 964–971 (2019).

8. Valles-Colomer, M. *et al.* The person-to-person transmission landscape of the gut and oral

microbiomes. *Nature* **614**, 125–135 (2023).

9. Zoetendal, E. G., Akkermans, A. D. & De Vos, W. M. Temperature gradient gel

electrophoresis analysis of 16S rRNA from human fecal samples reveals stable and host-

specific communities of active bacteria. *Appl. Environ. Microbiol.* **64**, 3854–3859 (1998).

10. Dubois, L. *et al.* Paternal and induced gut microbiota seeding complement mother-to-infant

transmission. *Cell Host Microbe* **32**, 1011-1024.e4 (2024).

11. Amir, A. *et al.* Gut microbiome development in early childhood is affected by day care

attendance. *Npj Biofilms Microbiomes* **8**, 2 (2022).

12. Gao, Y. *et al.* Gut microbiota maturity mediates the protective effect of siblings on food

allergy. *J. Allergy Clin. Immunol.* **152**, 667–675 (2023).

13. Jokela, R. *et al.* Sources of gut microbiota variation in a large longitudinal Finnish infant

cohort. *eBioMedicine* **94**, 104695 (2023).

14. Devotta, H. *et al.* Microbial Sharing Between Siblings Supports Metabolic Functions

Protective Against Allergy. *Allergy* **n/a**,.

15. Christensen, E. D. *et al.* The developing airway and gut microbiota in early life is influenced

by age of older siblings. *Microbiome* **10**, 106 (2022).

16. Fisk, C. M. *et al.* Influences on the quality of young children's diets: the importance of

maternal food choices. *Br. J. Nutr.* **105**, 287–296 (2011).

17. Kervinen, K. *et al.* Parity and gestational age are associated with vaginal microbiota

composition in term and late term pregnancies. *EBioMedicine* **81**, 104107 (2022).

18. Berry, A. S. F. *et al.* Remodeling of the maternal gut microbiome during pregnancy is shaped

by parity. *Microbiome* **9**, 146 (2021).

19. Kennedy, K. M. *et al.* Parity modulates impact of BMI and gestational weight gain on gut

microbiota in human pregnancy. *Gut Microbes* **15**, 2259316 (2023).

20. Kalomiris, A. E. & Kiel, E. J. Maternal Anxiety and Physiological Reactivity as Mechanisms

to Explain Overprotective Primiparous Parenting Behaviors. *J. Fam. Psychol. JFP J. Div.*

Fam. Psychol. Am. Psychol. Assoc. Div. 43 **30**, 791–801 (2016).

21. Hoskinson, C. *et al.* Delayed gut microbiota maturation in the first year of life is a hallmark

of pediatric allergic disease. *Nat. Commun.* **14**, 4785 (2023).

22. Korpela, K. *et al.* Cohort profile: Finnish Health and Early Life Microbiota (HELMi)

longitudinal birth cohort. BMJ Open 9 (6): e028500. (2019).

23. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon

data. *Nat. Methods* **13**, 581–583 (2016).

24. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing

and web-based tools. *Nucleic Acids Res.* **41**, D590-596 (2013).

25. Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A. & Callahan, B. J. Simple

statistical identification and removal of contaminant sequences in marker-gene and

metagenomics data. *Microbiome* **6**, 226 (2018).

26. Hsieh, T. C., Ma, K. H. & Chao, A. iNEXT: an R package for rarefaction and extrapolation

of species diversity (Hill numbers). *Methods Ecol. Evol.* **7**, 1451–1456 (2016).

27. McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis

and graphics of microbiome census data. *PloS One* **8**, e61217 (2013).

28. Oksanen, J. *et al.* vegan: Community Ecology Package. (2024).

29. Liaw, A. & Wiener, M. Classification and Regression by randomForest. **2**, (2002).

30. B\" urkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat.*

Softw. **80**, 1–28 (2017).

31. Raulo, A. *et al.* Social and environmental transmission spread different sets of gut microbes

in wild mice. *Nat. Ecol. Evol.* **8**, 972–985 (2024).

32. Baselga, A. & Orme, C. D. L. betapart: an R package for the study of beta diversity. *Methods*

Ecol. Evol. **3**, 808–812 (2012).

33. Nuriel-Ohayon, M., Neuman, H. & Koren, O. Microbial Changes during Pregnancy, Birth,

and Infancy. *Front. Microbiol.* **7**, (2016).

34. Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K. & Knight, R. Diversity,

stability and resilience of the human gut microbiota. *Nature* **489**, 220–230 (2012).

Supplementary Material

Supplementary Figure 1. Parity effects on infant-father similarity across infant ages. Points

show the estimated effect of parity (posterior means with 95% credible intervals) from separate

Bayesian regression models fitter at each timepoint. The y-axis represents the change in

similarity per additional previous pregnancy (i.e., number of older siblings).

Supplementary Figure 2. Decomposition of Jaccard dissimilarity into turnover and

nestedness components for infant–mother–father triads across infant development.

Pairwise dissimilarities between triads were calculated at each infant timepoint and partitioned

into (**A**) turnover (taxa replacement) and (**B**) nestedness (asymmetric subset relationships)

components. Comparisons are categorised as cross-parity (red; one triad from a multi-child

family and one from a single-child family) or within-parity (blue; both triads from the same

parity group). Violin plots show the distribution of values; black dots indicate medians. Asterisks

denote significant differences between comparisons (Wilcoxon rank sum tests; *<0.05, **<0.01;

***<0.001 FDR).

Supplementary Table 1. Importance scores from Random Forest regression model trained on

genus-level relative abundances across predicting microbiota age. Sequencing platform was

included as a binary predictor variable (MiSeq or HiSeq) as it was strongly confounded with

infant age. The model's predictive performance was assessed using 5-fold cross-validation,

where each sample was used for testing exactly once (cross-validated R^2=0.853, RMSE=6.092

months). Out-of-bag (OOB) performance metrics across folds: mean OOB R^2=0.879 ± 0.002;

mean OOB MSE=8.065 ± 0.085.

Genus	Importance
Anaerostipes	24792.7808
Faecalibacterium	22562.7705
Blautia	18336.567
Butyricicoccus	13773.9189
Roseburia	13613.013
Ruminococcus	12928.0235
Agathobacter	10481.7678
Lachnoclostridium	9895.53696
Monoglobus	9360.7023
`[Eubacterium] hallii group`	7253.84338
`[Ruminococcus] torques group`	5729.91155
`Lachnospiraceae NK4A136 group`	5541.21619
Fusicatenibacter	5324.73032
Flavonifractor	5310.93201
`Bacteria_Firmicutes_Clostridia_Oscillospirales_Ruminococcaceae_Incertae Sedis`	5152.76277
`Unknown genus from family [Eubacterium] coprostanoligenes group`	5138.04813
`[Eubacterium] eligens group`	4968.74245
Veillonella	4789.57779
Batch_MiSeq	4774.33042
`[Ruminococcus] gnavus group`	3708.49854
Subdoligranulum	3505.89252
Intestinibacter	3274.57521
Lachnospira	3121.72337
Dorea	3023.26626
`Unknown genus from family Lachnospiraceae`	2871.53016
`Lachnospiraceae UCG-004`	2702.9021
`Unknown genus from family Bifidobacteriaceae`	2230.60357
Megasphaera	1976.78083
Bifidobacterium	1806.7414
`Escherichia-Shigella`	1647.47675

Dialister	1593.0316
Romboutsia	1464.33577
UBA1819	1403.65915
`Erysipelotrichaceae UCG-003`	1367.34213
Erysipelatoclostridium	1363.47996
Staphylococcus	1360.53057
Akkermansia	1272.60348
Alistipes	1238.96869
Streptococcus	1206.11548
Sellimonas	1184.00218
Oscillibacter	1107.03506
`Unknown genus from family Enterobacteriaceae`	1095.84791
Eggerthella	1039.83355
`Clostridium sensu stricto 1`	970.428173
Bacteroides	968.140337
Coprococcus	858.876681
Haemophilus	844.088842
Lacticaseibacillus	838.474667
Hungatella	704.950626
`Unknown genus from family Ruminococcaceae`	700.697005
Enterococcus	687.115382
`Lachnospiraceae ND3007 group`	666.015682
Actinomyces	621.055534
Lactococcus	607.998665
Parasutterella	591.845494
`Unknown genus from family Oscillospiraceae`	579.019746
`Lachnospiraceae UCG-008`	544.83232
Collinsella	533.8195
Parabacteroides	524.000078
Rothia	518.463166
`[Eubacterium] ventriosum group`	500.481888
Aquamonas	492.481062
`UCG-005`	394.803628
`Lachnospiraceae NC2004 group`	391.973064
Klebsiella	381.773428
`UCG-002`	374.486725
`[Clostridium] innocuum group`	366.56486
Enterobacter	338.753234
`Unknown genus from family Clostridiaceae`	308.745643
`Christensenellaceae R-7 group`	294.080338
Sutterella	256.325613

Lactobacillus	233.984117
`[Eubacterium] siraeum group`	224.041835
Gemella	197.697795
Phascolarctobacterium	196.085618
`Unknown genus from family Lactobacillaceae`	185.458901
Bilophila	183.250592

Supplementary Table 2. Generalised additive model predicting microbiota age from

chronological age and covariates. Parametric coefficients represent linear effects on predicted

microbiota age (in months). Smooth terms represent non-linear age trajectories, with estimated

degrees of freedom (EDF) indicating curve complexity. The model included random intercepts

for Family ID to account for repeated measures (n=4,305 samples from 808 families; 1–8

samples per infants). Reference categories: no previous deliveries (siblings), both breastmilk and

formula, male infant, HiSeq sequencing platform, brown stool, no intrapartum antibiotics.

Adjusted R^2=0.918.

Parametric coefficients

Term	Estimate	SE	t-value	p-value
Intercept	12.772	0.435	29.37	<0.001
Previous deliveries (Yes)	0.140	0.045	3.14	0.002
Milk: Breastmilk	-0.459	0.055	-8.29	<0.001
Milk: Formula	0.403	0.092	4.36	<0.001
Milk: None	-0.039	0.083	-0.47	0.639
Infant sex (Girl)	0.008	0.043	0.19	0.849
Sequencing platform (MiSeq)	0.923	0.047	19.80	<0.001
Stool color: Clay gray	0.229	0.523	0.44	0.662
Stool color: Dark brown	-0.011	0.433	-0.03	0.980
Stool color: Green	-0.368	0.441	-0.83	0.404
Stool color: Light brown	-0.063	0.432	-0.15	0.885
Stool color: Yellow	-0.280	0.435	-0.64	0.520
Received intrapartum antibiotics	0.165	0.051	3.21	0.001

Smooth terms

Term	EDF	Ref. DF	F-value	p-value
Age trajectory: No previous deliveries	6.97	8	864.57	<0.001
Age trajectory: Previous deliveries	6.63	7	819.36	<0.001
Family (random effect)	377.68	804	0.93	<0.001