A single-cell atlas of *Toxoplasma* sexual development identifies regulators of

gametogenesis

Hisham S Alrubaye[1], Sarah M Reilly[1], Rafaela da Silva[1], NyJaee Washington[1] and Jon P

Boyle[1]*

[1] Department of Biological Sciences, Dietrich School of Arts and Sciences, University of
Pittsburgh
7
*To whom correspondence should be addressed:
University of Pittsburgh
4249 Fifth Avenue
Pittsburgh, PA. 15260
boylej@pitt.edu

Abstract

Toxoplasma gondii undergoes sexual development exclusively in the feline intestine, a process

critical for genetic diversity and population expansion. Recent studies have identified genes critical

in suppressing presexual development[1] and metabolic differences in felines that may promote

sexual development[2], but to date the gene regulatory networks driving development in the cat are

unknown. To investigate this, we performed single-cell transcriptomics on parasites isolated from

cat intestines, using fluorescent reporter strains and flow cytometry. From 15,068 cells across two

experiments, we identified rare populations, including cells that bear all of the hallmarks of

gametes. Candidate genes emerging from this study were tested via CRISPR-Cas9 Perturb-seq,

identifying AP2X6 as a regulator of macrogametocyte development. Our single-cell data extends

what is known about gene expression changes throughout sexual development and should be useful

to those in the field working towards inducing gametogenesis, mating, and oocyst production *in*

vitro.

Main

T. gondii has a complex life cycle that includes both asexual and sexual reproduction.

While asexual stages can develop in nearly all cells of warm-blooded animals, sexual development

is restricted to the epithelial cells of the felid small intestine. The sexual phase of *T. gondii* is

essential for generating genetic diversity through recombination and for large scale environmental

dissemination. A single feline host is capable of shedding almost a billion oocysts after a single

infection [3–6]. Shed oocysts mature further after completion of meiosis and one round of mitosis,

resulting in the production of infectious sporozoites.

Among the developmental stages of *T. gondii*, the tachyzoite and bradyzoite have been

studied the most due to their relative tractability *in vitro*, and a substantial amount of information

is available on the mechanisms of tachyzoite to bradyzoite stage transition[7,8] as well as stage-

specific cell cycle regulation[9,10], metabolism[11,12], and host-parasite interactions[13]. In contrast, due

to challenges associated with cultivating pre-sexual and ultimately sexual stages, there is far less

known about how this developmental program is triggered and executed by the parasite during

infection of the feline host. Early foundational ultrastructural studies by Dubey and Frenkel in the

1970s began to differentiate between the asexual and sexual enteric stages of *T. gondii*[3,14]. These

studies identified five morphologically distinct pre-gamete stages that develop following infection

of cat enterocytes[14]. While these investigations filled critical gaps in knowledge and provided a

developmental framework for understanding enteric development, the molecular mechanisms

underlying these processes remained unknown until recent advances. First, global transcriptomic

studies showed that *T. gondii* undergoing enteric development in the feline host have unique

transcriptional profiles compared to other developmental stages. These bulk transcriptomic studies

have been instrumental in uncovering significant transcriptional differences between

developmental stages in intermediate and definitive hosts, offering a comprehensive overview of

stage-specific transcriptional programs[15–17]. Furthermore, metabolic cues from the feline may help

trigger the transition to from asexual to presexual development in *T. gondii,* highlighting that host-

specific signals are essential in orchestrating parasite differentiation[18]. In addition, recent studies

have identified parasite transcription factors AP2XII-1 and AP2XI-2 as key suppressors of *T.*

gondii pre-sexual development, and their genetic depletion *in vitro* permits presexual development

[add citation]. These studies have paved the way for targeted investigations into the regulation of

this developmental program[1]. However, the depletion of those suppressors did not result in

parasites that were capable of forming gametes *in vitro*, indicating that additional factors are

required for gametogenesis.

Here, we interrogated the transcriptional program underlying the enteric stages of *T. gondii*

in the cat using single-cell RNA sequencing (scRNA seq) to identify previously unknown

regulators of enteric development. This approach enabled us to capture the transcriptional

heterogeneity of *T. gondii* parasites developing in the feline intestine at unprecedented resolution.

We identified multiple transcriptionally distinct clusters within the enteric stages, including

putative male and female gametocyte populations based on sex-specific expression signatures. To

functionally characterize key regulators of parasite development, we designed a targeted perturb-

seq screen of 28 candidate genes, leading to the identification of AP2X6 as a regulator of

macrogametocyte differentiation. Our data provides the field with novel gene markers for studying

distinct subpopulations of *T. gondii*, offering new targets for individual investigation to enhance

our understanding of the molecular mechanisms underlying the parasite development in the

definitive host.

                                                      Results

                      Generation of a cat-competent *Toxoplasma gondii* reporter strain

Transcriptomic analyses of *T. gondii* enteric stages in the feline intestine have thus far been

restricted to bulk RNA sequencing approaches, which lacks the resolution to capture the cellular

heterogeneity and obscures rare subpopulations such as transitional forms and gametes[16,17]. To

address this limitation, we interrogated the transcriptional program underlying enteric

development of *T. gondii* by generating single-cell transcriptomic profiles from parasites isolated

from the feline small intestine. To facilitate this, a *TgVEG* reporter line was engineered to express

GFP from a highly active tachyzoite promoter (GRA1) and mCherry from a hybrid promoter

(GRA11A + GRA1 upstream sequences) that resulted in a strain expressing GFP during tachyzoite

development and mCherry during both asexual and sexual development (Fig. 1A; Extended Data

Fig. 1).

To determine if our engineered strain retained the ability to complete its entire life cycle,

five brains from chronically infected CBA/J mice were fed to a cat. Tissue cysts from those brains

expressed both GFP and mCherry (Fig. 1B). During feline infection, fecal samples were collected

daily between days 3 and 17 post-infection. Unsporulated oocysts of TgVEG:GFP:mCh lacked

detectable GFP or mCherry fluorescence (Fig. 1C), whereas sporulated oocysts exhibited mCherry

fluorescence but no GFP fluorescence (Fig. 1D). Sporozoites were excysted from oocysts via

mouse infection[3], and parasites were recovered from the peritoneum by lavage and as expected

GFP and mCherry fluorescence were detected in TgVEG:GFP:mCh parasites growing as

tachyzoites *in vitro* (Fig. 1D). Because mCherry expression was observed in tachyzoites,

bradyzoites, and sporozoites, we hypothesize that constitutive expression results from the absence

of regulatory elements in the truncated GRA11A promoter, which has been proposed to be

occupied by MORC[19] , along with the physical proximity of this truncated GRA11A promoter to

GRA1 sequences in the plasmid driving GFP (see Extended Data Fig.1)

A single cell atlas of *T. gondii* enteric development

To identify the transcriptional program underlying the enteric development, we used

TgVEG:GFP:mCh parasites in feline infections. Thirty mice were infected with 1,000 low passage

TgVEG:GFP:mCh parasites, and 21 Days post-infection (DPI) mice were euthanized, and

deskinned skulls were divided into two groups and fed to two cats. Tissues were consumed within

30 min and infected cats began shedding oocysts on D4 PI. On D5 or D6 PI, feline intestinal

enterocytes were isolated from the small intestine by manual scraping, mechanically lysed, and

parasites were isolated by flow cytometry using expression of fluorescent proteins as proxies.

Single-cell transcriptomic profiles were generated for a total of 16,966 cells across two

independent experiments (Fig. 2A).

The single-cell RNA sequencing (scRNA-seq) transcriptomic profiles of *T. gondii* isolated

from feline enterocytes on 5 and 6 DPI differed significantly from *T. gondii* tachyzoites cultured

in vitro for 8 days post-excystation (Fig. 2B). As expected, the *in vitro* excysted parasites showed

high transcript abundance for canonical tachyzoite markers like SAG1 and SAG2A (Fig. 2C; top

panels)[20,21]. In contrast, parasites derived from the cat intestine exhibited significantly higher levels

of known merozoite markers like GRA80 and PNP (Fig. 2C; bottom panels) compared to

tachyzoites[1,16]. Overall, the gene expression profiles of intestine-derived parasites show many

unique transcripts that were distinct from those of tachyzoites as well as overlapping transcripts

(Fig. 2D), underscoring both the differences between life stages and the remarkable adaptability

of *T. gondii*, which enables it to thrive across various host environments. To identify any

differences between the cat samples, we downsampled to account for the differences in the cell

counts and performed differential gene expression analysis (Extended data Fig. 2). The enteric

samples from both cats are highly correlated (Fig. 2E)

Identifying distinct clusters of *T. gondii* in enteric stages

To interrogate the developmental program and identify distinct clusters in cat enteric

stages, we first identified and filtered out doublets using the scDblFinder algorithm[22]. The optimal

neighborhood size parameter (pK) for doublet detection was empirically determined as 0.08 to

maximize classification accuracy[23]. Artificial doublets were simulated from 25% of cells (pN =

0.25), and dimensionality reduction was performed using the top 25 principal components as

observed by the elbow method (Extended Fig. 3B). Predicted doublets were excluded prior to

downstream analysis. In addition, cells with fewer than 500 unique molecular identifiers (UMIs)

were removed based on the knee point detection in EmptyDrops (Extended Fig. 2A)[24]. After

filtering 15,068 of 16,966 cells passed these filters. Gap statistic analysis was then applied to

determine the optimal number of clusters in the scRNA-seq dataset[25,26]. This analysis identified K

= 10 as the optimal cluster number, which was used to partition cells into transcriptionally distinct

subpopulations (Fig. 3A). Feature plots of reported enteric stage-specific *T. gondii* genes show

variation in spatial distribution of transcript abundance (Fig. 3B).

To assess if any of the clusters were associated with previously reported canonical marker

genes, we examined the differentially expressed genes across all clusters and associated them with

cell-type based on previous studies. Cluster 1 showed enriched expression of canonical tachyzoite

markers like SAG2A and SAG1 (Fig. 3D). Both genes were found to be significantly enriched and

uniquely expressed in this cluster ($\log_2$FC >15; adj. p = 0.0). Therefore, we annotated cells in

cluster 1 as 'enteric tachyzoites' (Fig. 3C). It is possible that these parasites were in the process of

dissemination from the intestinal lining to other tissues within the host, since extraintestinal

development of *T. gondii* occurs even in the definitive host. Cluster 2 was defined by high

expression of genes like *Bradyzoite Rhoptry Protein 1* (*BRP1*; $\log_2$FC >9; adj. *p* = 1.43e-93) a

bradyzoite and merozoite marker, and AMA2 which is a protein involved bradyzoite invasion

($\log_2$FC >9; adj. *p* = 1.43e-93)[7,27,28]; Extended Data Fig. 2C and 2D). While BRP1 expression was

expected in merozoites[29], it was actually restricted to cluster 2 rather than being broadly distributed

across all parasites. TGME49_275790 is a merozoite-associated gene that is also enriched in this

cluster ($\log_2$FC >11.5; adj. *p* = 0), along with other tachyzoite-associated genes like ALV3 and

ALV12 ($\log_2$FC >12; adj. *p* = 0; Fig. 3E). The presence of bradyzoite/merozoite and tachyzoite

markers in these cells distinguishes this cluster from the rest of the enteric parasite population and

suggests that these parasites may represent an intermediate state between known canonical stages.

Therefore, this cluster was annotated as "BRP1+" based on one of its defining markers (Fig. 4E).

We compared the transcript abundance of the top 100 genes in the BRP1+ cluster to *in vivo* bulk

RNAseq transcriptome data from *T. gondii* cat infections[17] and found that these genes were most

characteristic of EES2 and EES3 stages (Fig. 3F). This confirms that this cluster is not an anomaly

of this particular infection as it is consistent with this prior dataset from infected definitive host

stages.

Clusters 3, 4, and 5 were enriched for multiple known merozoite-associated transcripts

(Fig. 3G) including SRS15B, SRS22E, SRS22D, GRA11A, and GRA11B[1,16,17,19,30]. Cluster 3

comprised 1,317 cells and was defined by high expression of TGME49_287040 ($\log_2$FC = 5.91;

adj. *p* = 0) and TGME49_200270 ($\log_2$FC = 5.87; adj. *p* = 0), both previously detected in enteric-

stage parasites[17]. Cluster 4 was the largest cluster, containing 12,051 cells (~80% of all cells), and

was characterized by the expression of genes like IMC7 ($\log_2$FC = 1.04; adj. p = 0) and

TGME49_244440 ($\log_2$FC = 1.73; adj. p = 0)[17]. Cluster 5 is made up of 1,199 cells with multiple

differentially expressed transcripts including TGME49_311720 ($\log_2$FC = 1.8; adj. p = 0) and

TGME49_293870 ($\log_2$FC = 1.8; adj. p = 0)[17]. Although all three clusters shared core merozoite

markers like GRA11A, GRA11B, and GRA82, each exhibited unique gene signatures that

distinguish them from one another. The top 10 differentially expressed genes for each cluster are

shown in Fig. 3H. Based on these profiles, clusters 3, 4, and 5 were annotated as "Merozoite-A,"

"Merozoite-B" and "Merozoite-C" respectively.

Cluster 6 was enriched for transcripts associated with late-stage development, as defined

by comparative expression analysis with data from Ramakrishnan et al[17]. Among the top

differentially expressed genes were TGME49_266960 ($\log_2$FC = 3.38; adj. p = 0) and

TGME49_290200 ($\log_2$FC = 4.41; adj. p = 0; Fig. 4B). When compared to Ramakrishnan et al[17],

the top 100 genes in this cluster exhibited increased expression in late EES samples (EES4 and 5;

Fig. 4C). Based on data presented immediately below, we hypothesizes that cluster 6 represents a

pre-gamete stage of development. Therefore, this cluster was annotated as "Pre-gamete" (Fig. 4A).

To determine if any of the clusters were associated with canonical markers of female

gametocytes, we used the genes HOWP1 and AO2 which are known female gamete markers in T.

$gondii$[31]. We found transcripts for these genes to be exclusively present in cells of cluster 7

(Extended Data Fig 4A and 4B)[31]. To further confirm this cluster as being made up of female

gametocytes, we used comparative transcriptomics between recently published $Cryptosporidium$

$parvum$ datasets, $Plasmodium$ and our data to identify conserved female-specific markers[32]. Out

of 108 female gametocyte-specific genes identified in $T.$ $gondii$, $C.$ $parvum$ and $P.$ $berghei$[31-33],

we found that 28 of them were enriched in Cluster 7 (Fig. 4D; Extended Data Fig 4C ;

Supplementary Table 2), including *TGME49_272900*, whose *Plasmodium* ortholog

(*PF3D7_0816800)* encodes the female-specific meiotic recombination protein DMC1[34], and

TGME49_294820, which encodes a type I fatty acid synthase, a protein found in high abundance

in oocysts from *T. gondii* and *Eimeria* and is also enriched in *Eimeria tenella* gametocytes [35]. The

expression of reported female-specific genes supported the annotation of this cluster as 'Female',

thereby providing the first detailed transcriptional view of *T. gondii* macrogametocytes. Among

the top 20 genes in the female cluster, three belong to the family D protein family

(TGME49_236975, TGME49_316670, and TGME49_277572), a large family of tandemly

duplicated genes of mostly unknown function. Additionally, we found that a putative surface

antigen, SRS46, is uniquely expressed in the cells of the Female cluster (Extended Data Fig. 3D),

making SRS46 a candidate surface marker of macrogametocytes. Furthermore, we examined the

top 100 differentially expressed genes with the cluster and found that their transcript abundance

was increased in late developmental samples from Ramakrishnan *et al*[17] (Fig. 4E). Thus, the

transcriptional signature of the top genes in Cluster 7 exhibited a late temporal expression profile,

consistent with the timing of macrogamete development.

Similar to macrogametes, very little is known about gene expression in *T. gondii*

microgametes. To identify these cells in our scRNAseq-derived clusters, we used putative

microgamete markers from Ramakrishnan *et al.* including genes involved in axoneme assembly,

flagellar transport, cell motility, and fertilization[17]. Mapping these genes onto our scRNAseq

projection (Fig. 4F; Supplementary Table 3) revealed a small number of cells across two clusters

(9 cells in cluster 8 and 3 cells in cluster 9) that express microgamete marker genes. We recognize

the limitation with the number of cells, however, the relative rarity of microgametocytes compared

to macrogametes is consistent with the literature[36,37]. The enrichment of previously identified

male-specific genes in these clusters supported their annotation as 'Males-1' and 'Males-2',

respectively[17] (see below). Importantly, we observed a similar female-biased ratio in a subsequent

scRNAseq experiment, where we identified 11 male and 106 female cells (Extended Data Fig. 5A

and 5B).

The Male-1 cluster expresses Hapless 2 (HAP2), a male-specific gene essential for gamete

fusion and subsequent fertilization in both *T. gondii and Plasmodium* (Extended Data Fig.

4B)[17,38,39]. In addition, Male-1 are predominantly in in S phase of the cell cycle which may suggest

that these are the replicating gametocytes (Extended Data Fig. 7). In contrast, the Male-2 cluster

exhibits higher expression of genes related to motility, such as *sperm associated antigen 6* (or

PF16; TGME49_297820), as well as genes encoding for flagellum-associated proteins (Extended

Data Fig. 4C). The *PF16* protein has been observed to associate with the growing cytoplasmic

axonemes along the emerging flagella in *T. gondii*[19], and in *Plasmodium berghei*, PbPF16 is

expressed in the male gametes[40]. PbPF16 is also essential for maintaining the correct microtubule

structure in the axoneme's central apparatus. This shows that PbPF16 plays a critical role in

flagellar formation, a process essential for male motility. Hence, it is possible to speculate that the

Male-1 cluster represents intracellular stages (either from a developing macrogamont or a

microgamete that has invaded a cell in search of the macrogamete), and that the Male-2 cluster

consists of extracellular microgametes, potentially searching for host cells with macrogametes.

When we compared the top 100 genes from clusters 8 and 9 to the transcriptomic data reported by

Ramakrishnan *et al*[17]*.,* we observed a strong concordance with genes enriched in samples from

later developmental stages (Fig. 4G and 4H), consistent with the expected timing of

microgametocyte development.

Another branch emerging from the trifurcation point of the pre-gamete cluster contains

cluster 10. Marker genes that are significantly enriched within cluster 10 include SRS48Q ($\log_2$FC

> 6.7; adj. p = 9.22E-164) and SAG2D ($\log_2$FC > 8.23; adj. p = 0), which are known to be

associated with merozoites[17]. Although cluster 10 shares some of the same genes with the Female

and Males-1 clusters, such as TGME49_294640 and TGME49_211030, it lacks the expression of

sex-specific genes. Specifically, SRS46, AP2X6, and TGME49_293368 are not detected in this

cluster. The expression profile of cluster 10 may indicate that these are cells re-entering the

merogony cycle after failing to differentiate into macrogametes or microgametes. This could

suggest a dynamic process where cells have the potential to shift between pre-gamete and

merogony depending on specific regulatory signals. Therefore, we have named this cluster

"Transitioning-merozoites" reflecting its intermediate state between pre-gametes and merozoites

(Fig. 4A). Furthermore, the top 100 genes from cluster 10 do not show an obvious association with

EES1-5 from Ramakrishnan *et al.* (Fig. 4L), suggesting that this is potentially a transient

population.

We expanded our analysis to investigate the biological processes distinguishing identified

male and female clusters by using GO term enrichment analysis on ToxoDB. The top 50 genes in

the female cluster were enriched in GO terms such as 'carbohydrate catabolic process'

(GO:0016052), 'small molecule metabolic process' (GO:0044281), and 'carboxylic acid metabolic

process' (GO:0019752) suggesting unique changes in specific metabolic pathways (Fig. 4I). It is

possible that female gametes may rely more on carbohydrate and carboxylic acid catabolism, as

well as small-molecule metabolic pathways, potentially to support ATP production and

biosynthetic processes required for fertilization and oocyst development. In contrast, the top 50

genes in the Male-1 cluster revealed overrepresentation of GO terms related to 'microtubule-based

movement' GO:0007018' 'microtubule-based process' (GO:0007017), and 'protein polymerization'

(GO:0051258) (Fig. 4J). The prominent GO term, 'microtubule-based process,' similar enrichment

was reported in *P. falciparum*[41] microgametes, suggesting a potential role for these genes in

shaping microgametes or driving the motility of flagellated male gametes. Furthermore, GO term

enrichment analysis of the top 50 genes from the Male-2 cluster revealed enrichment in terms such

as 'cilium assembly' (GO:0060271), 'plasma membrane-bounded cell projection assembly'

(GO:0120031), and 'cell projection assembly' (GO:0030031), emphasizing the important role of

structural and motility-related processes in this male cluster (Fig. 4K).

***In vivo* Perturb-seq screening identified AP2X6 as a repressor of asexual enteric**

development in the cat.

To genetically identify novel regulators of merogony and sexual commitment, we

conducted a targeted genetic screen, selecting 28 candidate genes based on their distinct

transcriptional profiles observed in our scRNA-seq dataset (Extended Fig. 4A). These genes

encode DNA- or RNA-binding proteins, surface antigens, and hypothetical proteins, and, more

importantly, 23 out of 28 exhibit unique expression profiles in sexual stages. Three different

protospacer sequences per gene were cloned into a *T. gondii* Perturb-seq vector using Gibson

assembly and validated by whole plasmid sequencing. The validated plasmids were pooled and

transfected into TgVEG:GFP:mCh parasites, generating a knockout pool that was used to infect

HFFs (Fig. 5A). To minimize bottleneck effects, each gene knockout was propagated as an

independent population, resulting in 28 distinct knockout parasite populations. These populations

were individually inoculated into mice at doses of 1,000 or 10,000 parasites per mouse (6–8 mice

per population). To control parasite growth, sulfadiazine treatment was administered to the mice

from 4 to 10-DPI. At 25 DPI, brain tissue containing bradyzoites in tissue cysts was harvested

from the chronically infected mice, and equal amounts (~16 g) were fed to two cats. Parasite

shedding was monitored, and upon detection parasites were isolated from the small intestine by

scraping and cell sorting for fluorescent parasites at 6 and 7 DPI (Fig. 5A).

To evaluate the complexity of the library prior to the *in vivo* infection, we performed

scRNA sequencing on the transfected parasites after drug selection, detecting protospacers in

23,285 of the 37,729 sequenced cells (~62%) and observing a sequencing saturation of 22.3%,

consistent with a highly complex library even at the low sequencing depth we used as our goal

was only to characterize library complexity and confirm the efficacy of gRNA capture using the

10X genomics system. To identify whether the targeted genes induced significant transcriptional

changes *in vitro*, we performed differential gene expression (DGE) analysis to compare the

expression profiles of each target gene across distinct populations. No significant transcriptional

differences were observed based on the presence of any single gRNA in tachyzoites (Extended

Data Fig. 8B). This lack of observable significance is consistent with the expression profile of

the targeted genes, which are predominantly active during the feline enteric stages and poorly

expressed in tachyzoites.

In the cat Perturb-Seq sample, 5,060 cells harboring single gRNAs targeting 18 different

genes were recovered (~19.5% of cells). This is lower than the expected recovery of cells

harboring gRNAs, which could be influenced by multiple factors including a bottleneck within the

cat small intestine. Cells expressing multiple gRNAs were filtered out, leaving 4,741 cells with

single gRNA sequences for downstream analysis. We found 5 specific perturbations in individual

cells that resulted in significant transcriptional changes in multiple genes (Fig. 5B). *AP2X6*

perturbation significantly increased transcript abundance for 121 genes relative to parasites with

other gRNAs, causing the largest perturbation of the parasite transcriptome across all detected

gRNAs. In cells harboring the SRS32-1 gRNA, we observed a significant decrease in transcript

abundance in 79 genes (Fig. 5B). Cells carrying the TGME49_311070-2 gRNA, a homolog of *fd1*

in *Plasmodium falciparum* 3D7[33] and associated with a female gametocytes (Extended Data Fig.

6A), had significant transcriptional changes in 36 genes. Furthermore, cells captured with

TGME49_297410 and SRS15B-2 gRNAs were found to have changes in transcript abundance in

73 and 5 genes, respectively. Cells harboring the gRNA corresponding to significant perturbations

are mapped on the UMAP (Fig. 5C). Some perturbations were observed to impact the same group

of genes, exposing a potential link in their function (Fig. 5E). However, follow-up experiments are

required to investigate the functions of these genes during enteric development and to confirm that

they participate in similar or connected gene regulatory networks.

AP2X6 is of a special interest due to its unique expression in the Female cluster.

Interrogating the top 50 differentially expressed genes in cells with the gRNAs targeting the female

specific *AP2X6* gene shows that the dysregulated genes are expressed primarily during the earlier

stages of merozoite development and particularly genes that are characteristic of the Merozoite-A

and BRP+ clusters (Extended Fig. 6B). This suggests that AP2X6 acts as a repressor of genes

express prior to gametogenesis in females and is potentially required to initiate and/or maintain

female gametocyte differentiation.

321   **Discussion**

The enteric stages of *T. gondii* are notoriously difficult to investigate due to their restriction

to feline hosts and the challenges associated with isolating them from cat intestinal epithelial cells.

Traditional bulk RNA sequencing approaches have been used on cat enteric stages[16,17] but these

bulk studies do not capture the heterogeneity and asynchronous development of the parasite. In

recent years, single-cell transcriptomics has emerged as a powerful tool, addressing these

challenges and enabling characterization of complex developmental trajectories in a wide array of

organisms including apicomplexan parasites[32,41–43]. In this study, we developed a constitutively

active reporter system using a hybrid promoter in *T. gondii* that enabled efficient sorting of enteric-

stage parasites from feline intestinal tissue using FACS, followed by scRNA-seq and

computational analysis to interrogate the transcriptional program underlying feline-specific

development. Using previously reported canonical markers, we identified 10 transcriptionally

distinct clusters, including intestinal tachyzoites, merozoites, transitional forms including pre-

gametes, and gametocytes. Evaluation of the top 100 genes from gametocyte clusters revealed

enrichment for genes found to have higher expression in later stages of development (i.e., EES5

samples in Ramakrishnan *et al.*) and supporting the late developmental identity of these clusters[17].

One of the most striking observations from our dataset is the disproportionate

representation of parasites undergoing merogony compared to those committed to sexual

development. Gametocytes were significantly outnumbered by merozoites, suggesting that only a

small fraction of parasites proceed toward gametogenesis. Our data support a model in which some

pre-gametes may re-enter the merogony cycle, forming what we term "transitioning merozoites."

Our observations suggest that not all cells within the cat enteric environment progress to

gametogenesis. Based on clustering and marker expression profiles, we hypothesize that the

majority of cells remain in an asexual enteric replication state rather than committing to sexual

development. This indicates that gamete formation may be a restricted or regulated process,

potentially requiring specific cues or conditions that are not detected by the majority of *T. gondii*

cells during development in cat enterocytes.. Such plasticity highlights the dominance of asexual

replication in apicomplexan life cycles and raises important questions about the regulatory

mechanisms that govern stage transitions. This skewed distribution is consistent with observations

in *P. falciparum*, where in controlled human malaria infections, fewer than 10% of parasites

commit to gametocyte formation[44,45]. It is reasonable to speculate that by maintaining a large

comparatively large asexual population it can sustain infection and prolong the window of

opportunity for mating, maximizing transmission potential. This strategy may be evolutionarily

conserved across apicomplexans.

Although our dataset constitutes significant progress towards gaining an understanding of

the gene expression dynamics in *T. gondii* enteric development, the comparatively low number of

male gametes indicate that we have yet to fully determine the spectrum of transcriptional profiles

in this life stage. This low number compared to female gametes is consistent with expectations

based on observations in *P. falciparum*[46,47]. Some male gametes may also be motile and either

invade host cells rapidly or be swept away in the cat gut and/or when we wash the intestine prior

to scraping. However, there is little doubt that these small clusters of cells represent male gametes.

For example, the centrin family protein (TGME49_237490) is associated with the basal structure

and function of flagella[48], sperm-associated antigen 6 (TGME49_297820) is a conserved marker

of male gametes, and *HAP2* is a well-characterized protein required for gamete fusion during

fertilization[49,40,19,17,31]. The specific and exclusive expression of these genes in the annotated male

gamete clusters supports their annotation, despite their comparative low numbers. To address the

challenge of low cell abundance, future studies could use male-specific markers identified in this

study to enrich for male gametocytes during isolation, allowing for deeper sequencing and

potentially functional studies with these poorly understood life stages.

Through scRNA-seq, we identified 28 candidate genes with potential contribution to various aspects of enteric development and used Perturb-Seq to determine their impact on parasite transcriptional profiles during enteric development. Among the genes with significant transcriptional impact was TGME49_311070, which is a female-specific gene. TGME49_311070 encodes a putative RNA-binding protein and is a homolog of the fd1 gene in *P. berghei*, which, when knocked out, results in infertile females lacking transcripts for a unique set of core female markers[33]. AP2X6-targeted cells showed increased transcript abundance for genes that are expressed earlier in development than AP2X6, suggesting that it functions as a repressor of specific genes during cat enteric development. This is consistent with the fact that multiple AP2 factors have been found to be repressors of transcription in *T. gondii*[1,50–52] rather than activators of transcription. The *in vivo* Perturb-seq data provide compelling evidence that *AP2X6* functions as a repressor of merozoite-specific genes in *T. gondii* during its enteric development. While we recognize the low number of cells detected with the *AP2X6* protospacer, the observed transcriptional differences between AP2X6-targeted cells and cells with other protospacers are robust. AP2X6-targeted cells induced upregulation of cells associated mostly with Merozoite-A and BRP1+ clusters. The expression of those genes decreased as we move down the developmental trajectory into Merozoite-B, which aligns with AP2X6 expression which was detected initially in Merozoite-B. The change in the transcriptional profile suggests that the loss of AP2X6 potentially leads to a failure in properly repressing asexual gene expression within the female lineage. This discovery constitutes a step forward in investigating gamete formation, even with the limited cell numbers.

Although this screen constitutes progress toward gaining a functional understanding of

enteric development, it has some limitations. First, the gRNA capture rate was lower than expected

at ~19%. Therefore, we were unable to detect all the gRNAs from the input library. Second, the

resources on gene essentiality and lethality are not available for the enteric cycle. gRNA targeting

of any number of these genes could have decreased or increased parasite fitness relative to others

in the population, leading to poor detection of those gRNAs.

In summary, our single-cell data provide valuable insights into the gene expression

dynamics during the enteric development of *T. gondii* in cats. This work holds the potential to

uncover previously unknown genetic drivers of differentiation, particularly those involved in

gametogenesis. These findings contribute to ongoing efforts in the field to induce gametogenesis,

mating, and oocyst formation *in vitro*, paving the way for future progress in manipulating the

parasite's life cycle.

407   **Materials and methods**

408   **Parasite strain and growth conditions:**

TgVEG parasites were used in this study. These were early passage parasites (P6) that were

excysted from sporulated oocysts as described[53], and maintained in human foreskin fibroblast

(HFF) cultures incubated at 37°C and 5% CO in Dulbecco's Modified Eagle Medium (DMEM;

Gibco) supplemented with 10% fetal bovine serum (FBS) (Atlas Biologicals), 2 mM L-glutamine

(Thermo, Fisher Scientific, Waltham, MA), and 50 mg/mL penicillin-streptomycin and

cryopreserved in 10% DMSO.

                                        Plasmid construction

          The TgVEG:GFP:mCh reporter strain was generated by introducing a cassette containing

   a truncated pGRA11A promoter upstream of mCherry into the pClick-GFP-Luc vector[54]. All

   primers used in this study are listed in Supplementary Table 7. The pTrunGRA11A was amplified

   from *TgVEG* genomic DNA, and mCherry was amplified from pCas9.T2A.HXGPRT[55]. The

   amplified products of pTrunGRA11A, mCherry, and linearized pClick-GFP-Luc were combined

   in a Gibson assembly reaction master mix (NEB E2621S) at a molar ratio of 0.014 pmol. The

   Gibson reaction was incubated at 50°C for 1 hour. Plasmid sequences were verified by Oxford

   Nanopore sequencing.

                          Generation of a *T. gondii* VEG reporter strain

          TgVEG was transfected with the reporter cassette using standard approaches. Briefly,

   parasites were mechanically released by needle passage (25 and 27-gauge needles) and then

   pelleted (10 minutes at 800 x g). Approximately 2×10^7 parasites were re-suspended in 100µl

   Amaxa Basic Parasite Nucleofector 1 (Lonza) along with 3µg of circular pSAG1::CAS9-

   U6::sgUPRT (Addgene plasmid #54467)[56] and 5µg of the pTrunGRA11A-mCherry-pGRA1-GFP

   cassette flanked by 20 bp of homology regions to the UPRT locus were co-transfected by

   electroporation (Supplementary Table 7). Transgenic and FUDR-resistant parasites were obtained

   by selection with 10 µM FUDR (Sigma-Aldrich, St. Louis, MO). We performed a diagnostic PCR

   to verify homologous integration and gene deletion.

                          Cat infections, oocyst isolation, and excystations

          For cat TgVEG:GFP:mCh parasites were released from HHF culture mechanically using

   25 and 27-gauge needles. The released parasites were then pelleted at 800 x g for 10 minutes, then

resuspended in PBS. Ten Mice (CBA/J from Jackson lab) were infected with 1,000 parasites in

200 ul PBS intraperitoneally (IP). The levels of IFN-γ were used as an indication of parasite

infection and burden early in infection. Blood serum from infected mice was obtained via

submandibular bleed, centrifuged at $100 \times g$ for 5 minutes, and then diluted at a 1:20 ratio. IFN-γ

concentrations were determined using BD OptEIA Mouse IFN-γ (AN-18) ELISA Set (Cat. #

551866) according to the manufacturer's instructions.

Mice were treated with Sulfadiazine (0.4g/L *ad libitum* in drinking water) from 6-10 DPI

to increase survival during the acute phage of infection. Throughout the course of infection, the

mice were monitored and weighed regularly. At 21DPI, a mouse was euthanized to determine the

cyst burden. The brain was harvested and homogenized in 1 ml of PBS (Sigma) and stained with

Biotin-conjugated Dolichos biflorus agglutinin (DBA) (1:500; Vector Laboratories, cat. no. RL-

1032) for 1 hour at room temperature. Subsequently, the sample was incubated with streptavidin

conjugated to Alexa Fluor 647. Cysts were visualized and counted using a 40x objective on an

Olympus IX83 microscope using cellSens software.

The remaining mice were euthanized, and the deskinned skulls and brains were fed to cats.

Feces were collected between days 3 and 17 post-infection and examined for oocyst production

using sucrose fecal floats. The collected feces were submerged in water and mixed with a wooden

tongue depressor to break down the materials. The emulsified mixture was then filtered through

840 μm and 250 μm sieves to further break down the particles. The filtered liquid was poured into

225 mL tubes and centrifuged at $2,000 \times g$ for 10 minutes. The resulting pellet was resuspended

in a 1.56 M sucrose solution (30 mL sucrose per tube per 10 g of feces). The sucrose mixture was

centrifuged at 1,180 × g for 10 minutes, and the top 10 mL of the supernatant was collected for

oocyst quantification using hemocytometer. This oocyst collection step was repeated at least twice

to minimize oocyst loss during processing. The isolated sucrose pellet containing oocysts was

mixed with water at five times the volume and centrifuged at 1,180 × g for 10 minutes. The pellet

was then resuspended in 2% H_2SO_4 and kept at room temperature with shaking for 7 days to allow

for sporulation. These samples were stored at 4°C.

To excyst *T. gondii* oocysts, sporulated oocysts were treated with 1N NaOH (at a ratio of

3:5 oocyst volume) to neutralize the 2% H_2SO_4 solution, and then washed multiple times with

Hanks' balanced salt solution (HBSS; Life Technologies, 14175145) until the pH was ~7.2. pH

of the solution was measured using pH indicator strips. The oocysts were then pelleted at 1,000 x

g for 8 minutes at 24°C and resuspended in PBS to prepare for excystation in mice. CBA/J mice

were infected with 100 oocysts orally or IP. On days 4-7 post-infection, mice were euthanized,

and peritoneal lavage fluids were collected in PBS. Lavage fluid was palleted by centrifugation,

resuspended in cDMEM, and then placed on confluent HHFs monolayers for 18 hours. Media was

changed as necessary, and parasites were visualized using light and fluorescence microscopy.

477   **Single-cell transcriptomics from *T. gondii* cultured *in vitro***

For this experiment, we used parasites that were cultured *in vitro* for 8 days post-

excystation. The sporozoites were excysted from oocysts as previously described [57,58]. Post-

excystation, the sporozoites were resuspended in culture media and used to infect confluent

monolayers of HHFs, grown at 37°C and 5% CO_2. The parasites were passaged as necessary. At

8 days post-excystation, the parasites were counted and processed for library preparation using

10x Genomics Chromium technology. The libraries were then sequenced on a HiSeq 4000

platform (Illumina) with a targeted depth of approximately 30,000 reads per cell for the 3' mRNA

library.

486   **Single cell transcriptomics of *T. gondii* tachyzoites and cat enteric stages.**

For single-cell transcriptomics, cats were euthanized at 5 and 6-DPI. The small intestine

was isolated from the euthanized cats and divided into three segments. The small intestine was

then washed with PBS and opened. The parasites were isolated from the small intestine of the cat

by scraping the intestinal lining with glass microscope slide (Fisherbrand: cat#12-550-112). The

scraped materials were syringe-passaged through 18, 22, 25, and 27-gauge needles, followed by

sequential filtration through cell strainers (100 μm and 40 μm, Corning). The strained materials

were pelleted at 1,000 x g for 8 minutes and resuspended in PBS with 1% FBS. The

*Tg*VEG:GFP:mCh parasites in the scraped materials were then sorted based on mCherry-derived

fluorescence expression. Sorting was performed using BD FACSAria. The sorted parasites cells

were concentrated by centrifugation at 1,000 x g for 8 minutes at 4°C and resuspended in PBS

containing 1% FBS (Sigma). The cells were then processed for library preparation using 10x

Genomics Chromium technology. The libraries were then sequenced on a HiSeq 4000 platform

(Illumina) with a targeted depth of approximately 50,000 reads per cell for the 3' mRNA library.

500   **scRNA-seq data processing**

Sequencing reads from the 3' mRNA libraries were processed using Cell Ranger v8.0.1

(10x Genomics). Reads were aligned to a custom *Toxoplasma gondii* reference genome (ME49,

based on release 56), including intronic reads, while retaining all other settings at their default

values. The custom library also included mCherry fused to the 3' end of DHFR and GFP fused to

the 3' end of GRA2, as these were components of the reporter cassette. We sequenced 16,695 cells

with 42,871 mean reads per cell; with a median UMI of 561 counts per cell. Cell calling was

performed using the two-stage algorithm implemented in Cell Ranger, which distinguishes cell-

associated barcodes from background noise such as ambient RNA from lysed or dead cells. The

barcode rank plot was used to visualize this separation, and the knee point guided the selection of

a UMI threshold to exclude low-quality cells and empty droplets. EmptyDrops was additionally

applied to refine cell calling. The resulting UMI count matrices were imported into Seurat v5.1.0

or Loupe Browser v8.0.0 for downstream analysis. A UMI threshold of 500 UMIs per cell was

applied to filter low-quality cells. Doublets were identified and removed using the scDblFinder

algorithm. The neighborhood size parameter (pK) was empirically optimized to maximize

classification accuracy, and artificial doublets were simulated from a subset of cells.

Dimensionality reduction was performed using principal component analysis, with the number of

components selected by the elbow method. Clustering analysis was performed using the gap

statistics method to determine the optimal number of clusters in R.

Generation of perturb-seq vector

Capture sequence 1 (GCTTTAAGGCCGGTCCTAGCAAG) was inserted into the

pCAS9-T2A-DHFR vector using Q5 mutagenesis (NEB) to generate pCAS9-T2A-DHFR-CS1

(TgPerturb-seq vector) and the perturb-seq vector was verified by nanopore sequencing. To

generate the pool of perturb-seq vectors used in the screen, single-stranded DNA (ssDNA)

oligonucleotides encoding the protospacer sequences for each gene of interest were generated

using EuPaGDT[59] . Three ssDNA were synthesized for each gene with 20 bases of homology to

either side of the perturb-seq vector integration site (Supplementary Table 5). Each ssDNA was

integrated into the linearized TgPerturb-seq vector (NcoI and PacI) by Gibson assembly (vector:

insert = 1:5). The Gibson reaction was incubated for 30 mins at 50 °C.

Perturb-seq experiment

*Tg*VEG:GFP:mCh parasites were transfected with NotI-HF-digested (NEB) perturb-seq

vectors by electroporation using Amaxa Basic Parasite Nucleofector 1 (Lonza). For each

transfection, approximately 10^7 parasites were electroporated with 15 μg of pooled Perturb-seq

vectors, composed of three vectors expressing different gRNA targeting the same gene (5 μg each).

Twenty-four hours post-transfection, 3 μM pyrimethamine (Sigma-Aldrich, St. Louis, MO) was

added to the media to select for integration of the perturb-seq vectors into the genome for 12 days.

The diversity of perturb-seq vectors was validated by amplifying the gRNAs, followed by

sequencing the amplicons.

Mice were infected IP with 1,000 or 10,000 parasites in 200 μl PBS. The infected mice

were treated with Sulfadiazine (0.2g/L) from 4DPI to 10DPI. The treatment was employed to

reduce the parasite burden. They were monitored and weighed regularly throughout the

experiments. To assess the cyst burden in the brains of infected mice, two mice were euthanized

at 21DPI for analysis. The brains were homogenized in 1 ml PBS and subsequently stained with

Biotin-conjugated *Dolichos biflorus* agglutinin (DBA) (1:500; Vector Laboratories, cat. no. RL-

1032) for 1 hour at room temperature, followed by incubation with Biotin-conjugated to Alexa

Fluor 647. Cysts were visualized and counted using a 40x objective on an Olympus IX83

microscope using cellSens software.

Upon cyst verification, brain tissues from the remaining mice were collected and fed to

two pathogen-free cats. Feces were collected from 3 DPI until oocysts were detected at 5 DPI. On

6 and 7 DPI, the cats were euthanized, and parasites were isolated from the small intestine and

sorted as previously described above.

The pooled sorted cells were concentrated by centrifugation at 1,000 x g for 8 minutes at

4°C and resuspended in PBS containing 1% FBS (Sigma). The cells were then processed for library

preparation using 10x Genomics Chromium technology. The libraries were then sequenced on a

HiSeq 4000 platform (Illumina) with a targeted depth of approximately 45,000 reads per cell for

the 3' mRNA library and 10,000 reads per cell for the CRISPR sgRNA library.

557   **Perturb-seq data processing**

The sequencing reads from the 3' mRNA and CRISPR sgRNA libraries were processed

using Cell Ranger v7.2.0 (10x Genomics). The reads were aligned to both a custom *T. gondii*

transcriptome reference and another reference containing protospacer sequences. Intronic reads

were included in the alignment, while all other parameters were left at their default settings. Cell

Ranger assigned the reads to specific cell barcodes, mapped them to the corresponding genes or

protospacers, and counted the unique molecular identifiers (UMIs) for each gene or protospacer

within each cell barcode. This generated UMI × cell barcode count matrices, which were

downloaded for downstream analysis to identify gRNA-specific perturbations using in the single-

cell transcriptome Loupe Browser v8.0.0 and Seurat v5.1.0. In total, we sequenced 26,000 cells

with a median depth of 1,996 UMI. 5060 cells were detected with one or more protospacer. For

analysis, we filtered cells with multiple protospacers to retains cells with a single protospacer

detected, 4741 cells, for further analysis.

Animal statement

The animal experiments were performed according to the guidelines of the American Veterinary Association. Euthanasia for mice was conducted using controlled CO_2 exposure. Euthanasia for cats was conducted using Beuthanasia. Animal protocols were approved under the Institutional Animal Care and Use Committee (IACUC) protocol no. 22061169.

References

1. Antunes, A. V. *et al.* In vitro production of cat-restricted Toxoplasma pre-sexual stages. *Nature* **625**, 366–376 (2024).

2. Di Genova, B. M., Wilson, S. K., Dubey, J. P. & Knoll, L. J. Intestinal delta-6-desaturase activity determines host range for Toxoplasma sexual reproduction. *PLoS Biol.* **17**, e3000364 (2019).

3. Frenkel, J. K. & Dubey, J. P. Toxoplasmosis and its prevention in cats and man. *J. Infect. Dis.* **126**, 664–673 (1972).

4. Dubey, J. P. Oocyst Shedding by Cats Fed Isolated Bradyzoites and Comparison of Infectivity of Bradyzoites of the VEG Strain Toxoplasma gondii to Cats and Mice. *J. Parasitol.* **87**, 215 (2001).

5. Dubey, J. P. TACHYZOITE-INDUCED LIFE CYCLE OF TOXOPLASMA GONDII IN CATS. *J. Parasitol.* **88**, 713–717 (2002).

6. Dubey, J. P. Unexpected oocyst shedding by cats fed Toxoplasma gondii tachyzoites: In vivo stage conversion and strain variation. *Vet. Parasitol.* **133**, 289–298 (2005).

7. Sokol-Borrelli, S. L. *et al.* A transcriptional network required for bradyzoite development in Toxoplasma gondii is dispensable for recrudescent disease. *Nat. Commun.* **14**, 6078 (2023).

8. Waldman, B. S. *et al.* Identification of a Master Regulator of Differentiation in Toxoplasma.
Cell **180**, 359-372.e16 (2020).

9. Radke, J. Defining the cell cycle for the tachyzoite stage of Toxoplasma gondii. *Mol. Biochem.*
Parasitol. **115**, 165–175 (2001).

10. Radke, J. R., Guerini, M. N., Jerome, M. & White, M. W. A change in the premitotic period
of the cell cycle is associated with bradyzoite differentiation in Toxoplasma gondii. *Mol.*
Biochem. Parasitol. **131**, 119–127 (2003).

11. Blume, M. & Seeber, F. Metabolic interactions between Toxoplasma gondii and its host.
F1000Research **7**, 1719 (2018).

12. Gallego-López, G. M., Contreras Guzman, E., Desa, D. E., Knoll, L. J. & Skala, M. C.
Metabolic changes in *Toxoplasma gondii* -infected host cells measured by autofluorescence
imaging. *mBio* **15**, e00727-24 (2024).

13. Hakimi, M., Olias, P. & David, L. Toxoplasma Effectors Targeting Host Signaling and
Transcription. *Clin. Microbiol. Rev.* **30**, 615-645p38 (2017).

14. Dubey, J. P. & Frenkel, J. K. Cyst-Induced Toxoplasmosis in Cats*. *J. Protozool.* **19**, 155–
177 (1972).

15. Behnke, M. S., Zhang, T. P., Dubey, J. P. & Sibley, L. D. Toxoplasma gondii merozoite gene
expression analysis with comparison to the life cycle discloses a unique expression state during
enteric development. *BMC Genomics* **15**, (2014).

16. Hehl, A. B. *et al.* Asexual expansion of Toxoplasma gondii merozoites is distinct from
tachyzoites and entails expression of non-overlapping gene families to attach, invade, and
replicate within feline enterocytes. *BMC Genomics* **16**, 1–16 (2015).

17. Ramakrishnan, C. *et al.* An experimental genetically attenuated live vaccine to prevent

transmission of Toxoplasma gondii by cats. *Sci. Rep.* **9**, 1–14 (2019).

18. Martorelli Di Genova, B., Wilson, S. K., Dubey, J. P. & Knoll, L. J. Intestinal delta-6-

desaturase activity determines host range for Toxoplasma sexual reproduction. *PLOS Biol.* **17**,

e3000364 (2019).

19. Farhat, D. C. *et al.* A MORC-driven transcriptional switch controls Toxoplasma

developmental trajectories and sexual commitment. *Nat. Microbiol.* **5**, 570–583 (2020).

20. Ferguson, D. J. P. Use of molecular and ultrastructural markers to evaluate stage conversion

of Toxoplasma gondii in both the intermediate and definitive host. *Int. J. Parasitol.* **34**, 347–

360 (2004).

21. Manger, I. D., Hehl, A. B. & Boothroyd, J. C. The Surface of *Toxoplasma* Tachyzoites Is

Dominated by a Family of Glycosylphosphatidylinositol-Anchored Antigens Related to

SAG1. *Infect. Immun.* **66**, 2237–2244 (1998).

22. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet

identification in single-cell sequencing data using scDblFinder. *F1000Research* **10**, 979

(2022).

23. Sun, R. *et al.* TREM2 inhibition triggers antitumor cell activity of myeloid cells in

glioblastoma. *Sci. Adv.* **9**, eade3559 (2023).

24. participants in the 1st Human Cell Atlas Jamboree *et al.* EmptyDrops: distinguishing cells

from empty droplets in droplet-based single-cell RNA sequencing data. *Genome Biol.* **20**, 63

(2019).

25. Tibshirani, R., Walther, G. & Hastie, T. Estimating the Number of Clusters in a Data Set Via

the Gap Statistic. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **63**, 411–423 (2001).

26. Guo, J. *et al.* Integrating imaging and genomic data for the discovery of distinct glioblastoma

subtypes: a joint learning approach. *Sci. Rep.* **14**, 4922 (2024).

27. Schwarz, J. A., Fouts, A. E., Cummings, C. A., Ferguson, D. J. P. & Boothroyd, J. C. A novel

rhoptry protein in Toxoplasma gondii bradyzoites and merozoites. *Mol. Biochem. Parasitol.*

144, 159–166 (2005).

28. Najm, R. *et al.* Invasion of *Toxoplasma gondii* bradyzoites: Molecular dissection of the moving

junction proteins and effective vaccination targets. *Proc. Natl. Acad. Sci.* **120**, e2219533120

(2023).

29. Schwarz, J. A., Fouts, A. E., Cummings, C. A., Ferguson, D. J. P. & Boothroyd, J. C. A novel

rhoptry protein in Toxoplasma gondii bradyzoites and merozoites. *Mol. Biochem. Parasitol.*

144, 159–166 (2005).

30. Ramakrishnan, C., Walker, R. A., Eichenberger, R. M., Hehl, A. B. & Smith, N. C. The

merozoite-specific protein, TgGRA11B, identified as a component of the Toxoplasma gondii

parasitophorous vacuole in a tachyzoite expression model. *Int. J. Parasitol.* **47**, 597–600

(2017).

31. Walker, R. A. *et al.* RNA Seq analysis of the Eimeria tenella gametocyte transcriptome reveals

clues about the molecular basis for sexual reproduction and oocyst biogenesis. *BMC Genomics*

16, 1–20 (2015).

32. Walzer, K. A. *et al.* Transcriptional control of the Cryptosporidium life cycle. *Nature* **630**,

174–180 (2024).

33. Russell, A. J. C. *et al.* Regulators of male and female sexual development are critical for the

transmission of a malaria parasite. *Cell Host Microbe* **31**, 305-319.e10 (2023).

34. Mlambo, G., Coppens, I. & Kumar, N. Aberrant Sporogonic Development of Dmc1 (a Meiotic
Recombinase) Deficient Plasmodium berghei Parasites. *PLoS ONE* **7**, e52480 (2012).

35. Bushkin, G. G. *et al.* Evidence for a Structural Role for Acid-Fast Lipids in Oocyst Walls of
Cryptosporidium , *Toxoplasma* , and *Eimeria*. *mBio* **4**, e00387-13 (2013).

36. Ferguson, D. J. P. Toxoplasma gondii and sex: essential or optional extra? *Trends Parasitol.*
18, 351–355 (2002).

37. Reece, S. E. & Read, A. F. Malaria sex ratios. *Trends Ecol. Evol.* **15**, 259–260 (2000).

38. Liu, Y. *et al.* The conserved plant sterility gene *HAP2* functions after attachment of fusogenic
membranes in *Chlamydomonas* and *Plasmodium* gametes. *Genes Dev.* **22**, 1051–1068 (2008).

39. Angrisano, F. *et al.* Targeting the Conserved Fusion Loop of HAP2 Inhibits the Transmission
of Plasmodium berghei and falciparum. *Cell Rep.* **21**, 2868–2878 (2017).

40. Straschil, U. *et al.* The Armadillo Repeat Protein PF16 Is Essential for Flagellar Structure and
Function in Plasmodium Male Gametes. *PLoS ONE* **5**, e12901 (2010).

41. Mohammed, M. *et al.* Single-cell transcriptomics reveal transcriptional programs underlying
male and female cell fate during Plasmodium falciparum gametocytogenesis. *Nat. Commun.*
15, 7177 (2024).

42. Xue, Y. *et al.* A single-parasite transcriptional atlas of Toxoplasma Gondii reveals novel
control of antigen expression. *eLife* **9**, e54129 (2020).

43. Real, E. *et al.* A single-cell atlas of Plasmodium falciparum transmission through the mosquito.
Nat. Commun. **12**, 3196 (2021).

44. Collins, K. A. *et al.* A controlled human malaria infection model enabling evaluation of
transmission-blocking interventions. *J. Clin. Invest.* **128**, 1551–1562 (2018).

45. Reuling, I. J. *et al.* A randomized feasibility trial comparing four antimalarial drug regimens

to induce Plasmodium falciparum gametocytemia in the controlled human malaria infection

model. *eLife* **7**, e31549 (2018).

46. Schall, J. J. The sex ratio of *Plasmodium* gametocytes. *Parasitology* **98**, 343–350 (1989).

47. Gruenberg, M. *et al.* qRT-PCR versus IFA-based Quantification of Male and Female

Gametocytes in Low-Density Plasmodium falciparum Infections and Their Relevance for

Transmission. *J. Infect. Dis.* **221**, 598–607 (2020).

48. Francia, M. E., Dubremetz, J.-F. & Morrissette, N. S. Basal body structure and composition in

the apicomplexans Toxoplasma and Plasmodium. *Cilia* **5**, 3 (2015).

49. Smith, E. F. & Lefebvre, P. A. PF16 encodes a protein with armadillo repeats and localizes to

a single microtubule of the central apparatus in Chlamydomonas flagella. *J. Cell Biol.* **132**,

359–370 (1996).

50. Radke, J. B. *et al.* ApiAP2 transcription factor restricts development of the *Toxoplasma* tissue

cyst. *Proc. Natl. Acad. Sci.* **110**, 6871–6876 (2013).

51. Fan, F., Xue, L., Yin, X., Gupta, N. & Shen, B. AP2XII-1 is a negative regulator of merogony

and presexual commitment in Toxoplasma gondii. *mBio* https://doi.org/10.1128/mbio.01785-

23 (2023) doi:10.1128/mbio.01785-23.

52. Srivastava, S., Holmes, M. J., White, M. W. & Sullivan, W. J. Toxoplasma gondii AP2XII-2

Contributes to Transcriptional Repression for Sexual Commitment. *mSphere* **8**, e00606-22

(2023).

53. Sokol, S. L. *et al.* Dissection of the in vitro developmental program of Hammondia hammondi

reveals a link between stress sensitivity and life cycle flexibility in Toxoplasma gondii. *eLife*

7, e36491 (2018).

54. English, E. D. & Boyle, J. P. Impact of Engineered Expression of Mitochondrial Association

Factor 1b on *Toxoplasma gondii* Infection and the Host Response in a Mouse Model. *mSphere*

3, e00471-18 (2018).

55. Young, J. *et al.* A CRISPR platform for targeted in vivo screens identifies Toxoplasma gondii

virulence factors in mice. *Nat. Commun.* **10**, 3963 (2019).

56. Shen, B., Brown, K. M., Lee, T. D. & Sibley, L. D. Efficient Gene Disruption in Diverse

Strains of Toxoplasma gondii Using CRISPR/CAS9. https://doi.org/10.1128/mBio.01114-14

(2014) doi:10.1128/mBio.01114-14.

57. Sokol, S. L. *et al.* Dissection of the in vitro developmental program of Hammondia hammondi

reveals a link between stress sensitivity and life cycle flexibility in Toxoplasma gondii. *eLife*

7, e36491 (2018).

58. Sokol, S. L., Wong, Z. S., Boyle, J. P. & Dubey, J. P. Generation of Toxoplasma gondii and

Hammondia hammondi Oocysts and Purification of Their Sporozoites for Downstream

Manipulation. in *Toxoplasma gondii* (ed. Tonkin, C. J.) vol. 2071 81–98 (Springer US, New

York, NY, 2020).

59. Peng, D. & Tarleton, R. EuPaGDT: a web tool tailored to design CRISPR guide RNAs for

eukaryotic pathogens. *Microb. Genomics* **1**, (2015).

Fig. 1: Generation of a cat-competent *Toxoplasma gondii* reporter strain.
A) A well-characterized oocyst-forming strain, TgVEG, was used in this study. A dual-reporter cassette was constructed with GFP driven by the *GRA1* promoter and mCherry under the control of a truncated *GRA11A* promoter. Both promoters were active in *T. gondii* tachyzoites. **B)** Tissue cysts derived from the TgVEG reporter strain were stained with Dolichos biflorus agglutinin (DBA; blue) and expressed both mCherry and GFP. **C)** TgVEG oocysts isolated from a cat experimentally infected with TgVEG reporter strain tissue cysts exhibited strong UV autofluorescence but lacked detectable GFP or mCherry expression. **D)** Sporulated oocysts of the TgVEG reporter strain expressed mCherry but not GFP. **E)** Sporulated oocysts were used to infect mice. Parasites excysted in the peritoneal cavity and were placed into culture, where they again expressed both mCherry and GFP.

Fig. 2: scRNA-seq transcriptome landscape of *Toxoplasma gondii* sexual and asexual development. A) Experimental workflow: Mice were infected with the TgVEG reporter strain for 25 days. Brain tissue containing bradyzoite cysts from 28 chronically infected mice was fed to cats (n = 2). Enteroepithelial stages were isolated from the small intestine of a cat euthanized at 5–6 days post-infection (DPI), sorted by mCherry fluorescence, and subjected to single-cell RNA sequencing (10x Genomics). **B)** UMAP visualization of scRNA-seq data from cat enteric stages (D5 and D6 DPI) and tachyzoites cultured for 8 days post-excystation. Distinct clustering reflects transcriptional differences between developmental stages. **C)** Expression profiles of canonical tachyzoite markers (top) and merozoite markers (bottom), highlighting stage-specific transcript abundance. **D)** Venn diagram showing overlap of expressed transcripts among 5 DPI, 6 DPI, and 8 DPI samples. Shared and unique transcripts are indicated by overlapping and non-overlapping regions. **E)** Correlation analysis of transcriptomes across samples, illustrating the degree of similarity between *in vitro* tachyzoites and enteric stages at 5 and 6 DPI.

Figure 3. Single-cell transcriptomic profiling of *T. gondii* enteric stages.
A) UMAP visualization of scRNA-seq data from cat enteroepithelial stages at 5 and 6 days post-infection (DPI) (n = 15,068 cells). Predicted doublets were removed using scDblFinder. Dimensionality reduction was performed with the top 25 principal components prior to UMAP embedding, and cluster number was estimated using the gap statistic. **B)** Feature plots showing expression of canonical markers for enteroepithelial stages. **C)** Cluster annotation based on top differentially expressed genes. **D)** Feature plot of Cluster 1 highlighting expression of canonical tachyzoite markers.**E)** Cluster 2 annotated as BRP1+ based on enrichment of BRP1-associated genes. **F)** Line plot of median z-scores for the top 100 genes defining the BRP1+ cluster across EES samples from Ramakrishnan *et al.* Shaded ribbons indicate ±1 SD. **G)** Venn diagram of significantly expressed genes shared and unique to Merozoite-A, -B, and -C clusters. **H)** Dot plot of the top 10 differentially expressed genes for each cluster. Dot size indicates the proportion of cells expressing the gene; color intensity reflects average expression. **I–K)** Line plots of median z-scores for the top 100 genes defining Merozoite-A, -B, and -C clusters across EES samples. Shaded ribbons represent ±1 SD.

Fig. 4: Transcriptomic signatures of pre-gamete and gamete-associated clusters in *T. gondii*.

A) Annotation of Clusters 6–10 based on top differentially expressed genes. **B)** Feature plots showing genes enriched in the pre-gamete cluster. **C)** Median z-scores of the top 100 pre-gamete cluster genes across EES samples from Ramakrishnan *et al.* Lines represent median expression per sample; shaded ribbons indicate ±1 SD. **D)** 106 female-specific genes identified in *T. gondii*, *Plasmodium berghei*, and *Cryptosporidium parvum* exhibit transcriptional enrichment in cluster 7. **E)** Median z-scores of the top 100 female cluster genes, showing increased transcript abundance in late EES samples. **F)** Distribution of male-specific genes across clusters. Feature plot of 46 male-specific genes identified by Ramakrishnan *et al.*, enriched primarily in clusters 8 and 9. **G–H)** Median z-scores of the top 100 genes defining Male-1 and Male-2 clusters, respectively. Both clusters show increased expression in later EES samples. **I–K)** Gene Ontology (GO) enrichment analysis of female (cluster 7), Male-1 (cluster 8), and Male-2 (cluster 9) clusters. Female cluster terms are associated with metabolic processes; male clusters are enriched for microtubule-based movement and organization. **L)** Median z-scores of the top 100 genes defining the transitioning merozoite cluster.

Fig. 5: Perturb-CRISPR screen and transcriptomic impact of candidate gene disruptions in *Toxoplasma gondii.*

A) Schematic of the perturb-CRISPR workflow. For each gene in the 18-candidate list, three distinct gRNAs targeting the same gene were transfected into TgVEG parasites to minimize bottleneck effects. At 24 h post-infection, cultures were switched to cDMEM containing 3 μM pyrimethamine for 12 days. Each knockout population was injected into 6–8 mice. After 25 days, brains were harvested for DNA extraction and for feeding to two cats. At 6–7 days post-infection, enteric stages were isolated from cat intestines, syringe-lysed, filtered (40 μm), and sorted by fluorescence before perturb-seq analysis. **B)** Heatmap of gRNAs associated with significant transcriptomic perturbations. **C)** UMAP visualization of single cells, highlighting those harboring gRNAs that caused significant transcriptional changes. **D)** Feature plot showing expression of AP2X6, a gene associated with the female cluster. **E)** Venn diagram illustrating shared and unique transcripts among cells harboring gRNAs that caused significant transcriptional changes. Only transcripts with consistent directional changes (all upregulated or all downregulated across compared groups) were included. Transcripts with opposing regulation in any group were excluded (e.g., *SRS32-1*). **F)** Expression dynamics of AP2X6 across EES samples. Blue points represent log$_2$-normalized read counts; the red line indicates the smoothed trend.