Single recipient cell tracking of tellurium-labeled extracellular vesicle proteomes (TeLEV) identifies EV-driven immunomodulation

Daniel Bachurski[1,2]*, Rahil Gholamipoorfard[1,3]*, Yong-Jia Bu[4]*, Patrick Hoelker[1,2], Lisa Wessendorf[1,2], Hendrik Jestrabek[1,2], Luca D. Schreurs[1,2], David Stahl[1], Amin Mokhlesi[1,3], Philipp Gödel[1], Felix Gaedke[5], Elias Ranjbari[6], Selen Seyhan[7], Ulrike Resch[8], Luisa Marie Schmidt[2,9], Tobias Tertel[10], France Rose[11,12], Cláudio Pinheiro[13,14], Maribel L. Corona[15], Anton von Lom[1,2], Alexander Frederik vom Stein[1,2], Phuong-Hien Nguyen[1,2], Katrin S. Reiners[16], An Hendrix[13,14], Guillaume van Niel[15], Marcus Krüger[2], Katarzyna Bozek[2,11,12], Lydia Meder[17], Per Malmberg[6], Paula Cramer[1], Barbara Eichhorst[1], Martin Peifer[3], Roland T. Ullrich[1,12], Astrid Schauss[5], Christian Pallasch[1,2], Paul J. Bröckelmann[1,18], Ron D. Jachimowicz[1,2,18], Christian Preußer[19], Bernd Giebel[10], Elke Pogge von Strandmann[19], Mark Nitz[4#], Michael Hallek[1,2,12#] and Nima Abedpour[1,3,20#]

*These authors contributed equally: Daniel Bachurski, Rahil Gholamipoorfard, Yong-Jia Bu
#These authors jointly supervised this work: Mark Nitz, Michael Hallek, Nima Abedpour

[1] University of Cologne, Faculty of Medicine and University Hospital Cologne, Department I of Internal Medicine, Cologne, Germany
[2] Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, Cologne, Germany
[3] University of Cologne, Faculty of Medicine and University Hospital Cologne, Department of Translational Genomics, Cologne, Germany
[4] Department of Chemistry, University of Toronto, Toronto, ON M5S 3H6, Canada
[5] University of Cologne, Faculty of Mathematics and Natural Sciences, Cologne Excellence Cluster for Aging-Associated Diseases (CECAD), Cologne, Germany
[6] Chalmers University of Technology, Gothenburg, Sweden
[7] Dokuz Eylül University, Izmir, Turkey
[8] University of Vienna, Vienna, Austria
[9] University of Copenhagen, Copenhagen, Denmark
[10] University of Duisburg-Essen, Essen, Germany
[11] Institute for Biomedical Informatics, Faculty of Medicine and University Hospital Cologne, Cologne, Germany
[12] Center for Molecular Medicine Cologne (CMMC), Faculty of Medicine and University Hospital Cologne, Cologne, Germany
[13] Laboratory of Experimental Cancer Research, Department of Human Structure and Repair, Ghent University, Ghent, Belgium
[14] Cancer Research Institute Ghent, Ghent, Belgium
[15] Université Paris Cité, Institute of Psychiatry and Neuroscience of Paris (IPNP), INSERM U1266, van Niel's Team, Paris, France
[16] Institute of Clinical Chemistry and Clinical Pharmacology - Immunology in Translation, University Hospital Bonn, Bonn, Germany
[17] Chair of Experimental Medicine I, Medical Faculty, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany
[18] Max Planck Institute for Biology of Ageing, Cologne, Germany
[19] Philipps University of Marburg, Marburg, Germany
[20] Cancer Research Center Cologne-Essen (CCCE), Cologne, Germany

Corresponding author:
Daniel Bachurski
Department I of Internal Medicine, University Hospital Cologne
Kerpener Str. 62, 50937 Cologne, Germany
E-Mail: daniel.bachurski@uk-koeln.de
Phone: +49 221 478 84118

Abstract

Single recipient cell tracking of tellurium-labeled extracellular vesicle proteomes
(TeLEV) identifies EV-driven immunomodulation

Extracellular vesicles (EVs) mediate tumor-immune cell communication by carrying protein
cargo that can immediately modulate signaling and antigen presentation. Yet mapping the
uptake of primary EV proteomes by human immune cells at single-cell resolution has been
constrained by a lack of labeling strategies. We show here that TeLEV, a tellurium-based
metabolic mass tagging approach that incorporates L-2-tellurienylalanine (TePhe) into EV
proteomes, can produce a biologically rare monoisotopic signal, which is detectable by mass
cytometry, imaging mass cytometry, and nanoscale SIMS, without perturbing EV morphology,
yield, or proteome composition. We applied TeLEV to label primary malignant B-cell-derived
EVs (MBC-EVs) from chronic lymphocytic leukemia (CLL) patients and could follow EV uptake
by distinct cell populations of healthy donor peripheral blood mononuclear cells. MBC-EV
uptake occurred predominantly in cells of myeloid lineages. In direct control experiments with
matched secreted proteins, a machine learning approach identified CD123, CD127, and CD25
as key markers distinguishing primary MBC-EV recipients from matched secreted protein
recipient cells. Nanoscale imaging enabled localization of EV-delivered proteins within
heterochromatin, whereas Te-labeled secreted proteins accumulated in the cytoplasm of
recipient cells. We then generated a pan-immune EV uptake atlas by tracing the uptake of
primary and cell-line EVs from nine cell lines and six donors with chronic lymphocytic leukemia
into 2,977,094 recipient cells across 43 cell types and subpopulations. We found that the
uptake of MBC-EVs by myeloid recipients induced monocyte-derived dendritic-cell polarization
characterized by the co-expression of the interleukin-receptor triad (IL-RT: CD123, CD127,
CD25) identified above. Time-resolved EV uptake analysis showed a rapid, transient
expression of CD123/CD127 followed by CD25, both tightly coupled to MBC-EV uptake by
myeloid cells. The intensity of IL-RT expression correlated with that of PD-L1 and BCL-2. Using
different STAT degraders to bidirectionally modify the EV-induced STAT5 signal, we observed
that MBC-EV uptake and IL-RT, PD-L1, and BCL-2 expression increased with STAT3
degradation and decreased with STAT5 degradation. To investigate the functional
consequences of the MBC-EV-induced changes, we showed that MBC-EVs in the presence
of IL-2 induced a high-CD25 immune state with low cytotoxic and high B cell proliferation.
Taken together, TeLEV represents a novel tool for single-cell tracking of EV proteomes,
revealing STAT5-dependent immune remodeling of recipient cells.

Introduction

Tumor-derived extracellular vesicles (EVs) and their protein cargo are potent immune
modulators[1–6]. Yet, research on EVs has been limited by tremendous difficulties in identifying
the target cells of the EV proteomes and studying their destiny[7–11]. Moreover, functional
investigations into the consequences of EV uptake remained limited, as cargo uptake could
not be followed at the single-cell level. The central hurdle for this problem was the detection of
EVs or the released proteins at single-cell resolution, as most labeling strategies stain material
that is not EV, alter the EV biogenesis or function, or conflate EVs with contaminating secreted
proteins, thereby disturbing functional studies linking cargo delivery with functional changes of
recipient cells[12–15]. An effective solution should provide a minimally disruptive, proteome-
sensitive label compatible with various cytometry and imaging methods, and an internal control
to distinguish EV-specific from general protein uptake[9,10,16].

To address this problem, we developed TeLEV, a tellurium-based mass tag that labels EV
proteomes with minimally perturbing, non-toxic L-2-tellurienylalanine (TePhe) at working
concentrations[16,17]. This allowed the creation of matched primary MBC-EV and secreted
protein controls based on TePhe, which mimics phenylalanine in both uptake and protein
incorporation[18]. The biologically rare monoisotopic tellurium signal enabled cross-platform
readout by mass cytometry, imaging mass cytometry, and correlative TEM-nanoSIMS[19–21],
allowing detection across multiple platforms and preserving EV morphology and composition.

When applied to MBC-EVs taken up by human peripheral blood mononuclear cells, TeLEV
revealed a predominantly myeloid MBC.EV uptake, which induced the coordinated expression
of interleukin receptors (CD123, CD127, CD25). EV proteomes upon uptake localized in the
heterochromatin of the recipient, while the matched secreted Te-labeled protein control
accumulated in the cytoplasm of recipient cells.

To causally interrogate the STAT5-dependence of EV-driven myeloid interleukin receptor
programs, we finally paired TeLEV recipient cell analysis with selective degraders of STAT3
and STAT5. For STAT3, we used SD-36, a selective PROTAC that degrades STAT3 and
increases STAT5 signaling in dendritic cells, as evidenced by increased pSTAT5[22]. For STAT5,
the dual isoform PROTAC AK-2292 degrades STAT5A/B with high selectivity over other
STATs[23]. Using both degraders, which bidirectionally modify STAT5 signaling—enhancing it via
STAT3 degradation and suppressing it via STAT5 degradation—allowed us to test that EV-
driven STAT3/5 programs were necessary to induce the observed myeloid phenotypes in
primary recipient cells. We discovered that MBC-EVs activate STAT5 signaling in myeloid cells,
orchestrating critical processes such as uptake mechanisms, interleukin receptor programs,
and monocytic polarization—highlighting a key role in immune regulation.

Main

Cross-platform recipient cell tracking with matched controls and nanoscale localization

We established TeLEV, a tellurium-based metabolic mass tag in which primary cells or cell
lines are cultured with L-2-tellurienylalanine (TePhe), the conditioned medium is fractionated
by size-exclusion chromatography, and Te-labeled MBC-EVs (F3–F5) are analyzed alongside
matched Te-labeled secreted proteins (F9–F11) from the same secretome (Figure 1a)[24]. This
matched-fractions design allows direct attribution of monoisotopic tellurium to EV-mediated
cargo versus freely secreted proteins and enables readout across suspension mass cytometry,
imaging mass cytometry, and correlative TEM-nanoSIMS (Figure 1b,c,d). At single-cell
resolution, the approach produced robust Te signals in peripheral blood mononuclear cells
(PBMCs) exposed to MBC-EVs for 16 h with negligible background in PBS-HAT controls, and
it cleanly distinguished uptake of EV cargo from uptake of matched Te-labeled soluble proteins
by high-dimensional analysis of recipient cell phenotypes (Figure 1b and Figure S1a,b,c).
Imaging mass cytometry on sister cultures reproduced heterogeneous recipient-cell intensities
with the expected per-pixel sensitivity difference relative to suspension mass cytometry
(Figure 1c)[20]. Correlative TEM-nanoSIMS localized the Te tag derived from Te-labeled primary
MBC-EVs predominantly to electron-dense heterochromatin in myeloid recipients[25]. In
contrast, the signal from Te-labeled secreted proteins was found to be concentrated in the
cytoplasm (Figure 1d). Together, Figure 1 demonstrates that TeLEV allows for the sensitive,
specific tracking of EV proteomes across diverse technologies, from population-level uptake
down to nanoscale localization, and to separate it from non-EV mediated protein transfer.

TeLEV preserves EV morphology, abundance, and proteome composition

Because labeling approaches involving modifications to EV-secreting cells could potentially distort the generation of primary EVs or their destiny, we investigated the effects of the TeLEV procedure on EV morphology, abundance, and protein composition[26]. Triplex immunogold TEM for CD9, CD63, and CD81 revealed indistinguishable labeling and preserved cup-shaped morphology in primary Te-labeled MBC-EVs and unlabeled controls. In the absence of uranyl-acetate contrast, primary Te-labeled MBC-EVs remained visible, consistent with intrinsic tellurium-derived contrast that reveals fine ultrastructural features (Figure 2a). Moreover, nanoparticle tracking analysis (NTA) of MBC-EVs generated from 10 primary CLL donors showed overlapping size distributions. Fractionation by size-exclusion chromatography in 3 donors produced concordant particle and protein profiles, with MBC-EV-enriched fractions peaking at F3–F5 and soluble proteins enriched in later fractions (Figure 2b,c)[27]. Pooled MBC-EV fractions (F3–F5) exhibited no difference in particle counts or modal diameters between TeLEV and control preparations (Figure 2d,e). Marker distribution and global proteomes were likewise unaffected. Immunoblots placed canonical EV markers (CD81, Flotillin-1, Syntenin) in F3–F5 and excluded GM130 from MBC-EV fractions in both conditions, with total protein enriched in late soluble fractions as expected (Figure 2f). Shotgun proteomics recapitulated these patterns and revealed no systematic shift with TeLEV. For example, samples clustered by donor rather than labeling status in Pearson-correlation matrices and principal-component analysis, global $\log_2$-intensity distributions, and the numbers of identified proteins were comparable between conditions, and differential testing detected no significantly changed proteins at FDR < 0.05 (s_0 > 0.1), including GO-annotated EV proteins (Figure 2g–m). Together, these data suggested that TeLEV did not measurably perturb MBC-EV morphology, yield, size distribution, or proteome composition across donors, justifying downstream single-cell recipient analyses.

Preferential MBC-EV uptake by myeloid cells

We next mapped primary MBC-EV uptake after 16 h by different immune cells at single-cell resolution (Data S2). After data cleaning, 785,260 mass cytometry events were integrated to construct a reference PBMC landscape of primary MBC-EV uptake[21]. Unsupervised FlowSOM clustering followed by manual curation identified 43 immune populations and subpopulations, including rare plasmacytoid dendritic cells (pDCs), myeloid dendritic cells (mDCs), γδ T cells, regulatory T cells, and residual low-density neutrophils (Figure 3a,b). Within this scope, TeLEV quantification revealed MBC-EV-protein uptake across all major cell lineages. However, TeLEV signal was clearly enhanced in myeloid cells, distributed across classical, intermediate, and non-classical monocytes, residual neutrophils, mDCs, and pDCs, but showed lower levels in all lymphoid subsets (Figure 3c, left and middle columns)[28]. Cell density distributions showed condition-specific redistributions in myeloid and T-cell compartments relative to vehicle controls, as indicated by shifts in density maxima in the t-SNE space. (Figure 3c, right column). In comparison, matched Te-labeled soluble proteins from the same secretomes were also preferentially taken up by myeloid cell populations and exhibited similar uptake patterns. Te-labeled proteins also resulted in two separate density peaks in classical monocytes and naïve CD4 T cells. Additionally, Te-labeled soluble protein uptake was overall 4.3 times higher (median Te signal all cells primary MBC-EVs 0.5199 vs. matched Te-labeled secreted proteins 2.2486), reflecting the greater abundance of labeled proteins in matched samples (Figure 2c,f and Figure 3c, middle column). To characterize primary MBC-EV-specific phenotypes, we trained separate random-forest classifiers on cells in the top quintile of Te signal for MBC-EVs

or soluble proteins (each relative to control)[29]. We evaluated the EV-to-protein log-ratio derived
from permutation-based feature importance values (Figure 3d). Interleukin-receptor (IL-R)
markers dominated this ratio, CD127 (IL7Rα), CD123 (IL3Rα) (IL-R triad, IL-RT), and CD25
(IL2Rα), with CD25 particularly highlighting TeLEV-positive monocytes and mDCs, including
IL-RT positive cells (Figure S2). These findings suggested that primary MBC-EV uptake
enhanced specifically IL-R expression in different myeloid recipient cells.

MBC-EVs induce an interleukin-receptor triad and a CD197$^+$ mo-derived DC-like state in
myeloid cells

To create a comprehensive immune recipient atlas of EVs from different sources, we
investigated 2,977,094 PBMC recipient cells across sixteen conditions (vehicle plus fifteen EVs
from nine cell lines and six primary CLL donors, Data S1). EV input was normalized to the
amount of conditioned medium rather than particle count (reported in Supplemental Table 2)
due to inaccuracies, methods limitations and detection thresholds of all current EV
quantifications methods[9]. A unified t-SNE embedding served as a reference for all comparisons
and revealed widespread EV uptake after 16 h (Figure 4a,b). We modeled cell-type proportions
using a Poisson generalized linear mixed model (GLMM) and identified condition-specific
frequency shifts in PBMC recipients, mainly affecting the myeloid compartment (Figure 4c). In
PBMC recipients, we observed that certain immune cells, such as IL-R+ and IL-R+/CD197+
monocytes and mDCs, showed an inverse relationship with the frequency of other immune
cells, such as non-classical monocytes and neutrophils. This pattern was observed in EV
conditions in two of six primary CLL donors (Patients 1 and 4). Additionally, similar cell
relationships were found in two cell lines, Ramos (a Burkitt lymphoma) and Su-DHL-4 (a
DLBCL line). A high-resolution re-embedding of 276,867 myeloid cells confirmed a recipient
cell signature only observed in MBC-EVs but not in malignant non-B cell EVs with enrichment
of mDCs, and a CD197$^+$ (CCR7+) monocyte subset with myeloid and DC markers, suggesting
a mo-DC differentiation induced by MBC-EVs (Figure 4d,e). This signature was absent with
malignant EVs from non-B cells, but was not elicited by all malignant B cell EVs. Unexpectedly
EVs from the CLL lines OSU-CLL and MEC-1 did not reproduce it, indicating dependence on
EV source and patient context (Figure 4f). Across B cell EV sources, the global responses to
Ramos and to CLL patient 1 and 4 were most similar, whereas the two CLL cell lines diverged
and lacked the MBC signature (Figure 4f). Particle number did not account for the signature,
as CLL patient 1 and Ramos EV preparations contained fewer particles than all other cell lines
tested (Supplemental Table 2). In the shared response pattern, myeloid recipient cells
increased levels of CD25, CD123, and CD127 (the interleukin receptor triad, IL-RT), with the
initial co-expression observed in Patient 1 (Figure S3). Some IL-RT-positive monocytes also
showed increased CD197 expression, indicating a transition toward a monocyte-derived
dendritic cell-like state. These cells had lower HLA-DR than conventional myeloid dendritic
cells, reduced CD14, and lacked CD183 (CXCR3)[30]. Ternary plots showed that MBC-EVs
increased CD25 expression of myeloid and lymphoid populations, while some non-B-cell EVs
(such as those from HEK293T and HS-5) increased CD127 (Figure 4g). Two controls
confirmed that these effects were specific to EVs, as the feature-importance analysis identified
CD25 as a distinguishing marker for EVs compared to soluble proteins (Figure 3d and Figure
S2). Unlabeled EVs from Patient 1 produced the same IL-RT pattern (Figure S3).

Biphasic IL-RT coupled to early myeloid uptake in EV uptake kinetics

We then characterized dynamics by exposing healthy-donor PBMCs to primary MBC-EVs
across eleven time points (0–48 h) and jointly analyzing 1,512,100 single cells in a common
embedding (Figure 5a). Tellurium signal became detectable by fifteen minutes, with myeloid
cells showing the earliest and strongest uptake (Figure 5b). Comparing the uptake dynamics
of secreted Te-labeled proteins at 2 hours to primary Te-labeled MBC-EV uptake, we observed
that MBC-EV uptake was more heterogeneous than the relatively uniform protein uptake in the
monocyte compartment (Figure S4). Early cell-composition profiles were stable for roughly the
first two hours and then progressively rebalanced after ~4 h, including an increase in CD197[+]
monocytes consistent with the emergence of a monocyte-derived DC-like state (Figure 5c). By
sixteen hours, uptake patterns converged with those observed in the broader study:
monocytes, mDCs, and pDCs dominated primary MBC-EV acquisition, and mDCs/CD197[+] IL-
R[+] monocytes displayed the highest Te levels (Figure 5d). Quantifying IL-RT trajectories within
myeloid lineages revealed a biphasic program tightly coupled to uptake. In monocytes, CD123
and CD127 rose within 15–30 minutes, dipped toward baseline by ~1 hour, and then entered
a second phase in which CD127 peaked near eight hours while CD123 continued to increase
through 48 hours; CD25 exhibited a delayed induction that lagged CD127 by ~8 hours and
remained elevated thereafter (Figure 5e-f). pDCs and mDCs showed analogous but lineage-
specific kinetics, with earlier and higher CD123 in pDCs and more sustained CD25 in mDCs
(Figure 5g). Taken together, primary MBC-EV uptake induced a transient early IL3Rα/IL7Rα
rise followed by IL2Rα expression and changes of the immunophenotype of myeloid cells.

Interleukin priming links EV exposure to sustained CD25 induction and increased IL-2
responsiveness in myeloid cells

To further explore and validate the functional effects of MBC-EVs on the myeloid compartment,
we conducted live-cell immunocytochemistry and (phospho)flow cytometry after exposure to
MBC-EVs. To confirm our results on a larger scale and to account for heterogeneity and limited
patient material, we chose Ramos-EVs as the MBC cell line model because it most closely
resembles Pat. 1 EVs in terms of IL-TR expression and monocyte-derived DC induction (Figure
4). In purified human monocytes, live-cell immunocytochemistry across MBC-EV dose titration
showed a dose-dependent increase in CD25; EV-treated monocytes displayed higher viability
(low Cytotox NIR signal) and frequent polarized/elongated morphologies, with the
highest-CD25 cells typically non-polarized but adjacent to polarized neighbors (Figure 5h,i).
Early flow cytometry at sixteen hours confirmed MBC-EV-dependent increases in CD25 and
CD127 across MBC-EV titration, with CD123 decreasing at the highest MBC-EV dose
(Figure 5j). The stimulation by various interleukin combinations revealed MBC-EV-dependent
priming in CD14[+] purified monocytes. IL-2 plus IL-7 in the presence of MBC-EVs produced
pronounced polarization and maximal CD25 by day 7, whereas IL-2 plus IL-3 or IL-3 plus IL-7
did not produce comparable effects; adding IL-2 + IL-3 + IL-7 increased viability and CD25,
with stronger responses when combined with EVs (Figure 5k). Consistent with direct
contributions from EV cargo to receptor–ligand landscapes, immunogold TEM detected IL-3,
CD123, and CD127 on MBC-EVs (Figure 5l). Intracellular phospho-flow cytometry linked
receptor remodeling to signaling: EV exposure increased pSTAT5 (MBC-EVs vs. Ctrl one-
sided Mann–Whitney U test, $p < 0.0001$); a STAT3 degrader further enhanced EV-induced
pSTAT5 (Stat3d + MBC-EVs vs. EVs one-sided Mann–Whitney U test, $p < 0.0001$); a STAT5
degrader reduced EV-induced pSTAT5 (Stat5d + MBC-EVs vs. MBC-EVs one-sided Mann–
Whitney U test, $p < 0.0001$); and IL-3 alone partially reproduced the MBC-EV-induced pSTAT5
increase (IL3 vs. Ctrl one-sided Mann–Whitney U test, $p < 0.0001$ and MBC-EVs vs. IL3 one-
sided Mann–Whitney U test, $p < 0.0001$) (Figure 5m). In PBMC co-cultures, MBC-EVs induced

CD25+ clusters and enhanced IL-2 responsiveness, while IL-only controls showed the
expected down-modulation of CD127 by IL-7 and the unexpected up-modulation by IL-3,
underscoring that MBC-EVs reorganize IL-R programs and their downstream signaling
properties rather than simply mirroring IL-only behavior (Figure 5n)[31]. These findings align with
the time-resolved biphasic IL-RT and help explain why MBC-EVs preferentially drive a
high-CD25 state in myeloid recipients.

STAT5 is required for CD25 induction and polarization and amplifies EV-driven
programs

In healthy donor PBMCs imaged for seven days, a STAT5 degrader (AK-2292) abolished MBC-
EV-driven monocyte polarization and prevented maximal CD25 induction, whereas anti-CD25
(Basiliximab) had little effect on MBC-EV-induced morphology; neither intervention affected the
MBC-EV-associated viability advantage in our in vitro setting (Figure 5o). In CD14+ purified
monocytes, phosphor flow cytometric analysis revealed that STAT3 degradation
augmented (Stat3d + MBC-EVs vs. MBC-EVs one-sided Mann–Whitney U test, p <
0.0001) and STAT5 degradation suppressed MBC-EV-induced CD25 (Stat5d + MBC-EVs vs.
MBC-EVs one-sided Mann–Whitney U test, p < 0.0001); STAT5 degradation also lowered
CD123 under both control (Stat5d vs. Ctrl one-sided Mann–Whitney U test, p<0.0001) and EV
conditions (Stat5d + MBC-EVs vs. MBC-EVs one-sided Mann–Whitney U test, p<0.0001). IL-
3 increased CD123 and CD127 (IL-3 vs. Ctrl, one-sided Mann–Whitney U test, both
p<0.0001), but not CD25 (IL-3 vs. Ctrl, one-sided Mann–Whitney U test, p=1). CD123 and
CD127 correlation evident in the controls was diminished in the presence of MBC-EVs
(Figure 5p). These experiments suggest that STAT5 may act as a signaling node for CD25
induction and MBC-EV-dependent morphological polarization of myeloid recipients.

STAT signaling modulates the uptake of MBC-EVs and tunes the IL-RT with checkpoint
and survival co-regulation

To assess how STAT signaling influences EV responses, TeLEV tracking was combined with
pharmacological degradation of STAT3 or STAT5 (± IL-2), and 3,590,520 PBMCs were profiled
across sixteen conditions. Cell-type frequencies remained consistent, allowing direct
comparison (Figure 6a,b). STAT3 degradation globally increased Te-labeled MBC-EV uptake
across most lineages, whereas STAT5 degradation decreased it (Figure 6c,d). The uptake
differences were not attributable to generalized activation, as CD69 levels were similar across
IL-2 conditions—EV alone, STAT3-degrader + MBC-EVs, and STAT5-degrader + MBC-EVs
(Figure S5). In myeloid cells, MBC-EVs increased CD25, CD123, and CD127; STAT5
degradation reduced their expression, and STAT3 degradation further amplified EV-driven IL-
R expression (Figure 6e,f). EV-induced CD127 was less affected by STAT5 loss than CD123,
and CD123 was reduced by STAT5 degradation even without MBC-EVs. STAT3 degradation
alone did not induce the IL-RT program but enhanced the EV-induced response (Figure 6g–k).
MBC-EVs also increased PD-L1 and BCL-2 in myeloid and dendritic cell subsets, monocytes,
mo-derived DCs, mDCs, and pDCs, with these changes heightened by STAT3 degradation
and diminished by STAT5 degradation (Figure 6l,m). At the PBMC level, IL-2 plus MBC-EVs
led to high CD25 and lower Ki-67 in cytotoxic lineages, but higher Ki-67 in B cells, even with
elevated CD69. STAT3 and STAT5 modulated both EV cargo uptake and interleukin receptor
expression, impacting checkpoint and survival markers

Discussion

We established TeLEV, a tellurium-based metabolic label that preserved EV morphology and
proteome composition. Across a total of 8.9M cells, it traced primary MBC-EV-proteomic
transfer into primary single-recipient cells using mass cytometry, imaging mass cytometry, and
nanoSIMS[32]. In PBMCs, uptake was myeloid-dominant. MBC-EVs induced a coordinated
interleukin receptor triad (CD123, CD127, CD25) with kinetics that aligned with EV uptake.
Two design elements guided interpretation. First, TeLEV labeled the MBC-EV proteome while
generating matched soluble protein fractions as control conditions, allowing attribution of
recipient cell phenotypes to EV-encapsulated proteins rather than co-secreted proteins[9,10,33].
Second, labeling did not change EV size distributions, yield, or proteome composition;
unlabeled MBC-EVs produced the same recipient-cell patterns.

Across nine cell lines and six primary CLL donors, MBC-EVs reproducibly increased CD123,
CD127, and CD25 in monocytes and dendritic cells, with source-dependent differences.
Importantly, differences in EV count did not explain these MBC-EV-induced interleukin receptor
co-expression. MBC-EVs pushed populations toward higher CD25, whereas few non-B
sources favored CD127. A subset of triad-positive monocytes acquired CD197 and features of
a monocyte-derived dendritic-like state with lower HLA-DR and absent CD183. Unlabeled
MBC-EVs reproduced the triad pattern seen with Te-labeled MBC-EVs, supporting EV
specificity. Time-resolved analyses showed that myeloid cells rapidly take up MBC-EVs,
triggering a rapid but transient increase in CD123 and CD127, followed by a later, longer-
lasting increase in CD25[35]. This process was observed across various myeloid cell types and
depended on EV dose and the surrounding cytokines[36]. The presence of IL-2 and IL-7, together
with MBC-EVs, led to the highest CD25 expression and cell polarization in purified monocytes.
In agreement with previous reports, MBC-EVs may deliver IL-3, which initiates STAT5
signaling[37–40]. In addition to IL-3, MBC-EVs also expressed CD123 and CD127. Exposure to
MBC-EVs boosted pSTAT5, linking receptor changes to signaling activity. Functionally, PBMCs
exposed to MBC-EVs and IL-2 became highly activated but did not proliferate as much,
especially in cytotoxic cells, while B cells showed increased proliferation. Overall, these results
outline a sequence where rapid MBC-EV uptake leads to early receptor priming and later CD25
upregulation, resulting in myeloid cells with enhanced survival and PD-L1 expression.

Our findings also demonstrate that STAT5 signaling is essential for mediating myeloid cell
responses to MBC-EVs. Specifically, loss of STAT5 abolishes the capacity of myeloid cells to
maximally upregulate CD25 expression and to acquire characteristic monocyte polarization,
indicating that STAT5 is indispensable for the terminal stages of the EV-driven response[41]. In
contrast, STAT3 degradation enhances EV uptake and amplifies the IL-RT response in myeloid
cells, but this effect depends on the presence of MBC-EVs. This observation suggests that
STAT3 functions as a negative regulator of the response, effectively acting as a brake on
STAT5 activity[22]. Notably, STAT3 degradation further increases pSTAT5 levels following EV
exposure, supporting the role of STAT3 in modulating STAT5 signaling during this process. A
novel finding of this study is that the recipient cell's STAT signaling state regulates EV uptake
itself, suggesting a feed-forward regulatory loop. A high STAT5/low STAT3 basal state may
prime myeloid cells to be more receptive to MBC-EVs, and the EV cargo then further amplifies
this state. This is further supported by the reduced CD123 expression observed in the baseline
condition exposed to a STAT5 degrader.

Pharmacological targeting of STAT3 (leading to increased pSTAT5) or STAT5 (resulting in
decreased pSTAT5) enables fine-tuning of both EV- and IL-RT–driven responses[22,23,42].
Immune checkpoint and survival markers such as PD-L1 and BCL-2 exhibit parallel regulation:
their expression is elevated in response to MBC-EVs, further enhanced by STAT3 loss, and

attenuated by STAT5 loss, mirroring the regulatory patterns seen for CD25 and MBC-EV uptake[22,43,44]. In PBMCs, the combination of IL-2 and MBC-EVs increases CD25 expression, reduces proliferation of cytotoxic lineages, and enhances B cell proliferation compared with IL-2 alone. Collectively, these results indicate that the interplay between EVs and the STAT pathway orchestrates recipient cell adaptation, with STAT5 serving as a requisite driver of late CD25 induction and monocyte polarization, and STAT3 acting as a context-dependent suppressor by regulating STAT5 signaling. This regulatory framework is consistently observed for PD-L1 and BCL-2 across myeloid cell subsets.

Our findings have important implications. First, they help clarify why myeloid remodeling and immune evasion often accompany high EV levels in B-cell malignancies: MBC-EVs appear to shift monocytes toward a CD197[+], PD-L1 high, BCL-2 high phenotype that supports cell migration, survival, and engagement of T-cell checkpoints, while also altering IL-2 signaling[44–47]. Second, the comprehensive EV uptake immune atlas developed here supplies quantitative metrics—such as EV uptake, IL-RT marker profiles, and EV-regulated immune profiles—that can be used to assess EV samples or guide experimental designs. Third, the discovery that STAT3 and STAT5 interact to fine-tune cell responses opens the door to experiments combining EV exposure with STAT pathway modulators or defined cytokine mixes to test specific mechanisms, and these strategies can be translated to animal models or patient samples to explore broader applications and therapeutic potential[5,48,49]. Finally, using a minimally disruptive, proteome-sensitive labeling strategy alongside matched soluble protein controls addresses ongoing uncertainty in the field, making it easier to determine whether observed effects are truly caused by MBC-EVs or by other secreted factors[50,51].

Taken together, TeLEV represents a breakthrough technology for high-resolution analysis of EV proteomes at the single-cell level. Using this innovative platform revealed that MBC-EVs actively shape myeloid immune landscapes, orchestrating a sequence of receptor changes that promote survival and immune checkpoint readiness. The STAT3/STAT5 signaling axis emerges as a key regulator, with targeted STAT3 degradation modulating STAT5 activity to finely tune immune responses. By combining nanoscale imaging with large-scale cellular profiling and targeted pathway interventions, TeLEV offers a powerful framework for dissecting EV-driven immunomodulation in both healthy and disease contexts. This technology holds promise for guiding future therapeutic strategies and for controlling EV-mediated effects in B-cell malignancies.

Methods

Primary leukemic samples

Primary CLL cells and PBMCs isolated from the peripheral blood of CLL patients and healthy donor PBMCs were collected at the University Hospital of Cologne after written and informed consent according to the Declaration of Helsinki and with Institutional Review Board approvals at the University of Cologne, no. 13-091 (BioMaSOTA), no. 19-1559 (Buffy Coats), no 21-1317 (SFB1530) and 21-1472. PBMC isolation and CLL cell purification procedures were performed as previously described[52]. CLL cells were freshly isolated to produce cell-conditioned medium. Clinical information relative to patient samples is listed in Supplemental Table 1.

Cell lines and cell culture

HS-5, THP-1, Su-DHL-4, OSU-CLL, MEC-1, Ramos, Jurkat, U-937, HEK293T were cultured in RPMI 1640 (Gibco, #11875093) supplemented with GlutaMAX (Gibco, #61870036), 10%

fetal bovine serum (Gibco, #10270106), and 1% penicillin–streptomycin (Gibco, #15140122).
All cell lines were tested for Mycoplasma contamination by PCR. All cells were maintained at
37°C in a humidified incubator with 5% CO_2. In all mass cytometry experiments, 1 x 10^6 cells
were analyzed per condition. For live-cell immunocytochemistry, cultures comprised 25,000
monocytes and 50,000 PBMCs. Flow cytometric analyses utilized a total of 100,000 cells per
condition.

Stimulation and treatment of recipient cells

Primary purified monocytes or PBMCs were stimulated with 20 ng/mL IL-2 (BioLegend,
#589102), 5 ng/mL IL-3 (BioLegend, #578002), and 25 ng/mL IL-7 (BioLegend, #581902)[53–55].
STAT3 (SD-36, MedChemExpress, #HY-129602) and STAT5 degraders (AK-2292,
MedChemExpress, #HY-148813) were used at 500 nM and compared to DMSO (PanReac
AppliChem, #A3672) controls[22,23]. Anti-CD25 antibody (Basiliximab, MedChemExpress, #HY-
10885) was used at 10 µg/mL and compared to an isotype control (Invitrogen, #MA5-55090).
All stimulation, inhibition, and degradation conditions were simultaneously added with MBC-
EVs.

Healthy donor CD14$^+$ monocyte and PBMC isolation

PBMCs were isolated from healthy donor buffy coats. Cells were washed twice in PBS
containing 2% FCS and 1 mM EDTA, then counted by flow cytometry. Cell counts were
determined by gating on singlets and DAPI-negative events. The necessary number of PBMCs
was resuspended to a final concentration of 1 × 10^8 cells/mL. CD14$^+$ monocytes were purified
by magnetic bead positive selection (EasySep Human CD14 Positive Selection Kit II,
STEMCELL Technologies, #17818) with an EasySep Magnet (STEMCELL Technologies,
#18001) using 100 µL of selection cocktail per 1 × 10^8 cells. Labeled cells were incubated for
6 minutes at room temperature, magnetically separated, and the CD14- fraction was
transferred to a new tube. The remaining CD14$^+$ cells were washed twice with 10 mL of buffer,
incubated for 3 minutes each time, and the supernatant was decanted. The previously
collected CD14$^-$ fraction was again incubated on the magnet for 6 minutes before collecting
the final supernatant. Viability was confirmed by DAPI exclusion using flow cytometry. Purified
cells were cryopreserved in FBS containing 10% DMSO, cooled at a controlled rate to –80°C,
then transferred to -150°C storage for long-term preservation.

EV separation

Primary MBC-EVs were isolated from EDTA-anticoagulated peripheral blood collected by
venipuncture from patients with chronic lymphocytic leukemia who presented with leukocytosis
≥ 15,000 leukocytes per mL, and samples were processed immediately (Sarstedt, #003.1524).
Leukemic B cells were enriched as untouched cells via immunodensity-based negative
selection according to the manufacturer's instructions using the RosetteSep Human B Cell
Enrichment Cocktail (STEMCELL Technologies, #15064).

For EV production, the primary cells were adjusted to 1.0 × 10^7 cells/mL, and the cell lines
were adjusted to 1.0 × 10^6 cells/mL, with viability greater than 95% as determined by DAPI
exclusion. The cells were washed twice and cultured in serum-free medium for 48 hours.
Conditioned supernatants were harvested and clarified by centrifugation at 350 × g for 5
minutes at room temperature, followed by 2,000 × g for 10 minutes at 4 °C. The clarified
medium was passed through a 0.22 µm PES membrane filter (Filtropur S, Sarstedt,
#83.1826.001) after thawing and then concentrated in 10 kDa centrifugal ultrafiltration units
(Amicon Ultra-15 10 kDa, Merck Millipore, #UFC901024) to a final volume of 0.5 mL before

being applied to a qEVoriginal 35 nm size-exclusion column (qEVoriginal 35 nm Gen2, Izon,
#ICO-35). The concentration step was performed at 4 °C.

Fractions were collected in 1 mL increments once the 0.5 mL load had fully entered the column.
Fractions F3–F5 were retained as EV-enriched fractions, and fractions F9–F11 were collected
as protein controls. Particle concentrations of the EV-enriched fractions were quantified by
nanoparticle tracking analysis as described in the dedicated NTA section, and the samples
were then reconcentrated. For functional assays, EV preparations were supplemented with in-
house-prepared PBS-HAT as previously reported[56]. All EV samples were stored at −80 °C
before use.

TePhe synthesis

472   125TePhe and 130TePhe (L-2-tellurienylalanine) were synthesized by adapting previously
473   reported methods for the synthesis of TePhe with natural-abundance Te[17,18]. Isotopically
474   enriched Te was sourced from Trace Sciences (Lot 134-1, 92.8% isotopic enrichment for Te-
475   125 and Lot 130Te/9, 99.40% isotopic enrichment for Te-130).

Tellurium-based metabolic labeling of the extracellular vesicle proteome

477   Immediately after purification, primary CLL cells or cell lines are seeded for media conditioning
478   (see EV separation). 50 µM of the isotopologues Te-125 or Te-130 L-2-tellurienylalanine
479   (TePhe) are added as medium supplements for metabolic labeling of nascent secretomes.

EV assay normalization

Current challenges in EV assay normalization stem from the lack of standardized methods for
quantifying EVs across different systems[9,10]. Co-isolated secreted proteins often confound
protein yields from EV preparations, and preparations deemed 'pure' typically exhibit very low
protein concentrations. Quantification techniques such as nanoparticle tracking analysis
(NTA), nanoflow cytometry, and resistance pulse sensing are limited by device-specific
detection thresholds, which may exclude smaller EVs (<60 nm) and introduce variability across
cell systems. Additionally, TeLEV normalization cannot rely on TePhe incorporation signals, as
these are strongly modulated by cellular protein synthesis and secretion rates. To provide a
pan-immune EV uptake atlas, we normalized EV input using 25 mL of conditioned medium
collected over 48 hours from 25 million cells, thereby facilitating cross-cell-line comparisons.
Supplemental Table 2 details EV counts from NTA, volumes, and specifies the input material.
We acknowledge that EV input normalization remains an unresolved challenge within current
methodological constraints.

Immunoblotting

Cells or MBC-EVs were lysed using RIPA buffer (Cell Signaling Technology, # 9806S). From
the cell lysates, 10 µg of protein was loaded per lane. The secretome fractions were adjusted
to a cut-off of 10 µg, whereas the EV fractions did not reach this level. The samples were
separated on NuPAGE 4-12% Bis-Tris Gel (1.5mm, 15 wells, Invitrogen, # NP0336BOX) under
reducing conditions and transferred to a nitrocellulose membrane (Amersham, Protran, #
GE10600016). Protein expression was visualized using Revert 700 Total Protein Stain for
Western Blot Normalization (LI-COR, # 926-11011). The membranes were blocked for 1 hour
at room temperature with Intercept (TBS) Blocking Buffer (LI-COR, #927-60001) and then
incubated overnight at 4°C with unlabeled primary antibodies against GM130, Alix, CD19, IgM,
HSP90, HSP70, Syntenin-1, and CD81. After washing, the blots were incubated with an IRDye
800CW Donkey anti-Rabbit IgG secondary antibody (1:15,000, LI-COR, #926-32213) for 1

hour at room temperature and imaged with a LI-COR Odyssey CLx Imager (Lincoln, Nebraska, USA). To allow re-probing with additional primary antibodies, membranes were stripped using NewBlot Nitro Stripping Buffer (5X, LI-COR, #928-40030) according to the manufacturer's instructions and subsequently re-incubated with alternative primary antibodies.

Nanoparticle tracking analysis

Measurements were performed on a ZetaView PMX-120-12F-R5 (Particle Metrix, Germany) equipped with a 488 nm, 40 mW laser and a CMOS camera. Samples were diluted in PBS to a final volume of 1 mL. Optimal concentrations were determined by pre-testing to achieve 100-300 particles per frame. Each measurement involved one cycle, during which 11 cell positions were scanned at a frame rate of 30 Hz. Instrument settings were: autofocus enabled, camera sensitivity 80.0, shutter 100, and cell temperature 25 °C. Videos were analyzed using ZetaView Software v8.06.01 SP1 with analysis thresholds of maximum particle size 1000 nm, minimum size 10 nm, and minimum brightness 20.

Immunogold transmission electron microscopy

Formvar-coated copper grids (Science Services, Munich) were prepared with 5 µl of undiluted purified EV samples. The grids were incubated with the samples for 20 minutes, blocked for 30 minutes (Aurion, #905.001), and washed three times for 2 minutes in PBS containing 0.1% BSA-c (Aurion, #900.099), pH 7.4. Three gold-coupled antibodies or primary unlabeled antibodies (Supplemental Table 3) were diluted 1:25 in PBS with 0.1% BSA-c: CD9, 10 nm (ABCAM, #STN-273562), CD63, 20 nm (ABCAM, #STN-273564), and CD81, 40 nm (ABCAM, #AB273949) combined in one mix. Grids were incubated in drops of antibody solution for 60 minutes, washed 6 times for 5 minutes in drops of PBS with 0.1% BSA-c, followed by 3 2-minute washes in PBS. Unlabeled antibodies were detected with a Protein G Immungold detection step at 1:20 dilution (Aurion, #810.133). After washing, grids were fixed for 5 minutes with 1% Glutaraldehyde (Sigma, #G5882-100ML) in PBS and washed three times for 2 minutes in ddH2O. Grids were stained for 4 minutes with 1.5% Uranyl acetate (Agar Scientific, #R1260A) and blotted dry. For each sample, an unstained duplicate was prepared. Images were captured with a Gatan OneView 4K camera on a JEM-2100Plus (Jeol) at 200 kV and analyzed with ImageJ (NIH).

Cytospin and Imaging Mass Cytometry

Fully stained cells (see "Mass cytometric surface and intracellular protein detection" and Figure 1b) were subjected to Cytospin on a Cytospin 4 centrifuge (Epredia, #A78300002) using Shandon EZ Single Cytofunnel assemblies (Epredia, #11972345) and Superfrost Plus glass slides (Epredia, #J1800AMNZ). Optimal cell density was determined by titration to 0.5×10^6 cells in 0.5 mL PBS, yielding a single cell layer on the deposition area. After exposure to EVs, the PBMCs were washed and resuspended in PBS, then immediately loaded into the cytofunnel assemblies according to the manufacturer's instructions. Assemblies were mounted with Superfrost Plus slides and centrifuged for 7 min at 800 rpm. Slides were removed and allowed to air dry at room temperature before further processing. IMC of Cytospin preparations was performed on a Hyperion imaging mass cytometer (Standard BioTools), ablating 750 µm x 750 µm representative regions of interest with a laser power of 3 (laser frequency 200 Hz).

The imaging mass cytometric images were converted from the mcd format to tiff files, and hot pixels were removed using a previously reported method[57]. We used the pretrained mesmer2 model from DeepCell to segment the cells[58]. Namely, images of the nuclear marker DNA and the cytoplasmic marker CD45 were combined into an RGB image (red = DNA, green = CD45,

blue = none) and cut into 512x512-pixel patches with a 256-pixel overlap. The Mesmer2 model is called for each patch, yielding nuclear and whole-cell probability maps. The probability maps from all the patches are merged using averaging, and a watershed algorithm is then applied, as in the original DeepCell code. For each detected cell (whole-cell segmentation map), the averaged signal intensity for the marker 125TeLEV is computed and arcsinh-corrected with cofactor 5. For display, aggregates were removed using morphological operations. After otsu thresholding, the connected components are processed with a closing operation with a disk element of radius 1 and with an opening operation with a disk element of radius 2. If the connected component has a radius over 150 pixels, it is considered an aggregate and removed from the detections.

Mass spectrometry sample preparation

Secreted proteins and EV samples collected in PBS were diluted with 1:1 (v:v) 8% SDS in PBS, homogenized by heating for 10 min at 95 °C, and sonicated with a Bioraptor sonicator; the sonication settings were 21 °C water with 10 cycles of 30 on and 30 off. 20 µg protein extract was reduced and alkylated for 10 min at 70 °C using 5 mM TCEP and 15 mM CAA. Protein digestion was performed following the SP3 protocol (Hughes, 2019 #117). In brief, 20 µg of both hydrophobic and hydrophilic beads were added to the sample, and the beads were bound by adding 1:1 volume of acetonitrile (ACN). After 8 minutes of incubation time, magnetic beads were immobilized and washed 2x with 70% ethanol and acetonitrile. Proteins were digested with trypsin (substrate:enzyme ratio 100:1) and LysC (substrate:enzyme ratio 200:1) overnight at RT. Samples were acidified to 1% formic acid (FA), then cleaned up using house-made SDB-RPS tips. Desalted peptides were loaded onto Evotips, and separation was performed on an Evosep ONE LC system (Evosep, Denmark) equipped with 8 cm PepSep columns (EvoSep, Denmark) at 45 °C with a gradient length of 30SPD (44 min). Mobile phases were composed of 0.1% FA as solvent A and 0.1% FA in ACN as solvent B. The LC system was coupled to a timsTOF pro 2 using a CaptiveSpray source (both Bruker, Germany). Samples were measured in dia-PASEF mode, with ion mobility calibrated using three ions from the Agilent ESI-Low Tuning Mix, following vendor specifications. The dia-PASEF window ranged in dimension 1/k0 0.7-1.45, with 16x2 Th windows and in dimension m/z from 350 to 1250.

LC-MS/MS sample analysis

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD069581[59]. Acquired spectra were analyzed with DIA-NN (V1.9.1) using library-free search against UniProt Homo Sapiens database (Apr. 2024)[60]. Mass ranges were set according to the settings of the mass spectrometer. Data were further processed using R (v. 4.2.2) with the libraries tidyverse, diann, and data.table, magrittr, FactoMineR, factoextra, and ggplot2, gprofiler, ggplot2. Data input was filtered for unique peptides, q-values < 0.01, and Lib.Q.Value <0.01, PG.Q.Value < 0.01, Global.Q.Value < 0.01, Quantity.Quality > 0.7, Fragment.count >= 4.

Further analysis was performed in Perseus (V 1.6.5.0) and InstantClue (V 0.10.10.20211105). Data were filtered to ensure 100% data completeness. Unpaired Student T-tests were performed on s0 = 0.1, FDR < 0.05, and 500 randomizations. Identified GO terms and GMI extracellular vesicle annotation were imported and compared for each group with an unpaired t-test.

Correlative transmission electron microscopy and NanoSIMS analysis of EV recipients

PBMCs were centrifuged at 350 xg and fixed for 1 h in 2% Glutaraldehyde with 2.5 % Sucrose
(Roth, #4621.1) and 3mM CaCl2 (Sigma, #C7902-500G) in 0.1M HEPES buffer (Sigma,
#C7902-500G) pH 7,4. The supernatant was removed, and the pellets were mixed with 3%
low-melting agarose in 0.2M HEPES buffer, incubated for 10 minutes at 37°C, and hardened
at 4°C for 30 minutes. Agarose blocks were cut into pieces of 1 mm³, washed four times for
15min with 0.1M HEPES buffer, and incubated with 1% Osmium tetroxide (Science Services,
#E19190) and 1% Potassium hexacyanoferrate (Sigma, #P8131) for 1 h at 4°C. After 3x5 min
washes with 0.1 M Cacodylate buffer (Applichem, #A2140,0100), samples were dehydrated at
4°C using an ascending ethanol series (50%, 70%, 90%, 3x100%) for 15 min each. The sample
was incubated in a mixture of 50% ethanol/propylene oxide, then twice in pure propylene oxide
for 15 minutes each. Infiltration was performed with a mixture of 50% Epon/propylenoxide and
75% Epon/propylenoxide for 2h each step and with pure Epon (Sigma, #45359-1EA-F)
overnight at 4°C. Samples were embedded into TAAB capsules (Agar Scientific, #G3744) and
cured for 48 h at 60°C. Ultrathin sections of 70 nm were cut using an ultramicrotome (Leica
Microsystems, UC6) and a diamond knife (Diatome, Biel, Switzerland) and picked up using
200 mesh finder Grids without film (Plano, #7FGC200) Sections were stained with 1.5 % uranyl
acetate (Agar Scientific, #R1260A) for 15 min at 37°C and 3% Reynolds lead citrate solution
made from Lead (II) nitrate (Roth, #HN32.1) and tri-Sodium citrate dehydrate (Roth #4088.3)
for 4 min. Images were acquired using a JEM-2100 Plus Transmission Electron Microscope
(JEOL) operating at 80kV equipped with a OneView 4K camera (Gatan). Mosaics of regions
of interest were acquired using SerialEM [61]. Location of the cell of interest was noted using the
finder pattern. After TEM imaging, sections were transferred onto small 5x5 mm silicon wafer
pieces (Plano, #G3390). For this, a small drop of water was added to the wafer, and the Grid
was placed on the drop with the section facing the wafer. After the water had evaporated, the
grid was removed. Wafers were shipped by placing them onto double-sided sticky tape (Tesa,
#05338) inside a petry dish. Images of the same cells were acquired using nanoSIMS.
Registration of images obtained by electron microscopy and nanoSIMS was done using nuclei
as fiducials with the plugin EC-CLEM from the software ICY[62].

Analysis was performed at NanoSIMS Sweden, University of Gothenburg, using a NanoSIMS
50L (CAMECA, Gennevilliers, France). Samples were gold-coated, and a 16 keV Cs^+ primary
ion beam of 0.2 pA (D1-5) was used to raster the sample surface. The detectors were tuned to
measure 12C2-, 12C14N-, and 130Te- secondary ions. Before imaging, the areas of interest
were implanted to achieve a fluence of $5×10^5$ $Cs^+.cm^{-2}$ to enhance the ionization yield. Images
were acquired on a field size of $10×10$ μm^2 with an image resolution of $512×512$ pixels and a
dwell time of 3 $ms.pixel^{-1}$. 50 planes were accumulated for each sample. NanoSIMS images
were visualized and analyzed with the WinImage software (CAMECA, Gennevilliers, France).
Sequential image planes were collected, drift corrected, and accumulated to increase ion
counts.

Incucyte live cell immunocytochemistry

Live-cell immunocytochemistry on an Incucyte system was performed to quantify surface
expression of CD25 and CD127 on primary human cells under standard incubator conditions,
with a near-infrared (NIR) cytotoxicity readout. PBMCs were plated at 50,000 cells per well,
and purified monocytes were plated at 25,000 cells per well in 100 μL of 0.22 μm-filtered,
complete medium in Falcon 96-well black-wall, clear-bottom, tissue culture-treated imaging
microplates (Corning Falcon, #353219). Plates were incubated for 1 h at 37 °C and 5% CO_2
before kinetic imaging was initiated. Imaging was conducted on an Incucyte instrument
equipped with green, red, and NIR channels using a 20× objective. Phase-contrast and

fluorescence images were acquired every 2 h for 48 – 7 days from four non-overlapping fields
per well with hardware autofocus enabled. Alexa Fluor 488 was detected in the green channel,
phycoerythrin (PE) in the red channel, and Cytotox NIR in the NIR channel. Exposure times
were kept within the linear range across the time course using the vendor's recommended
presets and were held constant within experiments.

For live-cell immunocytochemistry, non-IL-2/IL-7-blocking anti-CD25-Alexa Fluor 488 and anti-
CD127-PE were added at 1 µg mL^{-1} each directly to cultures (Supplementary Table 3).
Incucyte Cytotox NIR was added at a final concentration of 0.6 µM to report membrane
compromise. Image analysis was performed in Incucyte software. Total cell masks were
generated from phase-contrast images to estimate object counts, and fluorescence masks
were generated in the green and red channels to calculate the fraction of CD25^{+} and CD127^{+}
cells and mean integrated fluorescence per cell at each time point. Local background
subtraction was applied for visualization.

Flow cytometry (surface and intracellular phospho-staining)

For flow cytometric analysis, cells were cultured for 16 hours or 2 days in V-bottom 96-well
plates (100 µL/well). Cells were first incubated with 0.5 µL TrueStain FcX (BioLegend,
#422302) per well, for 15 min at 4°C. Subsequently, surface staining was performed using a
pre-titrated antibody cocktail (see Table SX1) for 1 h at 4°C in the dark. Following incubation,
cells were washed twice with Cell Staining Buffer (CSB, BioLegend, #420201) and immediately
processed for acquisition.

For intracellular flow cytometric analysis of phosphoproteins, cells were fixed by adding an
equal volume of Fixation Buffer (BioLegend, #420801) directly to the culture medium and
incubating for 15 min at 37°C. Fixed cells were washed three times in PBS by centrifugation
at 900 × g for 2 min to remove residual fixative. Fixed cells were surface-stained as described
above, then permeabilized for intracellular staining.

Permeabilization was performed using True-Phos Perm Buffer (BioLegend, #763344) by
resuspending cells in 200 µL per well and incubating for at least 1 h at -20°C. Cells were then
washed four times with CSB (900 × g, 2 min) and subjected to FcR blocking (as above),
followed by overnight staining with an intracellular antibody cocktail at 4°C.

After staining, cells were washed twice with CSB and resuspended in the same buffer for flow
cytometric acquisition. Data were collected on a Miltenyi MACS Quant X and analyzed using
FlowJo software (version 10.10, BD Biosciences).

Antibodies

Antibodies for mass cytometry were either purchased pre-conjugated with heavy metals
(Standard BioTools) or conjugated in-house using the Maxpar X8 chelating polymer kit
(Standard BioTools) or the Maxpar MCP9 polymer kit (Standard BioTools) according to the
manufacturer's instructions. Details on antibody clones, corresponding heavy metal tags, and
suppliers are provided in Supplementary Table 3.

Mass cytometric surface and intracellular protein detection

Cells were exposed to EVs in 2 mL deep-well 96-well plates (Eppendorf, #0030502310). After
EV incubation, the samples were washed and then incubated in the same plates with IdU at
50 µM (Standard BioTools, #201127) for 1 h at 37 °C with 5 % CO_2, followed by two washes
before antibody staining. TrueStain FcX Fc-receptor block was added, and the samples were

incubated for 15 min at 4 °C. For surface staining, the antibody master mix was added to each
well, and the cells were incubated for 45 minutes at 4 °C with intermittent gentle vortexing.
They were then washed twice in CSB at 350 × g for 5 minutes at 4 °C (BioLegend, #420201).
For intracellular antigen detection, the cells were fixed and permeabilized for 45 min at 4 °C
using the FoxP3 fixation/permeabilization buffer set (Invitrogen eBioscience, #00-5523-00),
then washed twice in the corresponding permeabilization buffer at 900 × g for 5 min at 4 °C.
Fc-receptor block was reapplied for 15 minutes at 4 °C. The intracellular or cytoplasmic
antibody master mixes were then added, and the samples were incubated for 45 min at 4 °C
with intermittent gentle vortexing, followed by two washes in permeabilization buffer at 900 ×
g for 5 min at 4 °C. The cells were fixed in 1.6% methanol-free paraformaldehyde for 10 min
at room temperature (ThermoFisher, #28906), then washed and incubated overnight at 4 °C
with MaxPar Intercalator-Ir 191/193 at a final concentration of 125 nM prepared in MaxPar
fix/perm buffer (Standard BioTools, Intercalator #201192B, Fix and Perm #201067). On the
following day, the samples were washed twice in CSB and prepared for acquisition.

Mass cytometric data acquisition

Data were acquired on a Helios mass cytometer (Standard BioTools) with daily instrument
quality control and tuning. Manual gating using FCSExpress7 (DeNovo Software) was applied
to identify live single cells in the sample convolute based on 140Ce, event length, center, width,
DNA (191Ir and 193Ir), and live/dead (103Rh) channels.

Mass cytometry data preprocessing

Signal intensity measurements in mass cytometry (CyTOF) are often influenced by neighboring
channels, leading to spillover effects due to inherent technological limitations. These
interferences, known as spillover effects, can significantly reduce the accuracy of cell-type
identification. To compensate for spillover effects, we use CytoSpill [63], a tool that quantifies and
corrects spillover in CyTOF data without requiring single-stained controls. We used the control
condition dataset to estimate the spillover matrix with the R package *CytoSpill* (v1.0.1), which
was then applied to compensate for the other datasets. Following spillover correction, the
single-cell data undergo an inverse hyperbolic sine transformation with a co-factor of 5
(arcsinh(x/5)).

FlowSOM clustering of mass cytometry data and cell type annotation

Preprocessed mass cytometry data from multiple experiments were integrated for clustering.
The lineage-defining markers 89Y-CD45, 141Pr-CD196, 143Nd-CD123, 144Nd-CD19, 145Nd-
CD4, 146Nd-CD8a, 147Sm-CD11c, 148Nd-CD16, 149Sm-CD45RO, 150Nd-CD45RA, 151Eu-
CD161, 152Sm-CD194, 153Eu-CD25, 154Sm-CD27, 155Gd-CD57, 156Gd-CD183, 158Gd-
CD185, 160Gd-CD28, 161Dy-CD38, 163Dy-CD56, 164Dy-TCRgd, 166Er-CD294, 167Er-
CD197, 168Er-CD14, 170Er-CD3, 171Yb-CD20, 172Yb-CD66b, 173Yb-HLA-DR, 174Yb-IgD,
176Yb-CD127 were normalized using the 99.9th percentile prior to clustering. Cells were
initially grouped into 125 clusters based on lineage marker expressions using the R package
FlowSOM (v2.10.0)[64]. These clusters were subsequently manually merged and annotated
according to biological relevance, yielding 43 distinct cell populations and subpopulations.

Cell type composition analysis

A generalized linear mixed model (GLMM) with a Poisson outcome[65] was used to model the
impact of conditions on cell–type–specific counts, with conditions included as a covariate. The
effect of this factor on cell-type composition was estimated through interaction terms with cell

type. Models were fitted using the glmer function from the R package *lme4* (v1.1.37). The fold
change is relative to the grand mean and adjusted. Statistical significance was evaluated using
the local true sign rate (LTSR), which represents the probability that the estimated direction of
effect is correct.

Random Forest analysis

A random forest model was used to evaluate the impact of Te-labeled MBC-EVs and Te-labeled
MBC-secreted proteins on recipient cells relative to control cells. For robust comparisons, cells
with EV and protein signal levels in the top 20% (highest quintile) were selected. The primary
MBC-EV (or Te-labeled protein) and control data were combined, and features were
normalized using z-score transformation. This integrated dataset was used to train a random
forest classifier with the RandomForestClassifier() function from the Python package scikit-
learn (v1.4.0). To address class imbalance, class weights were applied, and hyperparameters
were optimized via 5-fold cross-validation. Feature importance was calculated as the average
across 10 nested cross-validation runs, with 5 permutations per feature, to estimate each
feature's contribution to group differentiation.

Visualization

746 Visualizations were carried out in R (v.4.3.2) and Python (v.3.10) using scanpy (v.1.11.0)[66],
747 which is a scalable toolkit for analyzing single-cell data. For data visualization, high-
748 dimensional single-cell data were reduced to two dimensions using the nonlinear
749 dimensionality reduction algorithm t-SNE (t-distributed stochastic neighbor embedding)[67]. t-
750 SNE plots were created and visualized using scanpy.

EV-TRACK

752 We have submitted all relevant data of our experiments to the EV-TRACK knowledge database
753 (EV-TRACK ID: EV250125)[68].

Statistics

755 All statistical differences were calculated in GraphPad Prism, R, or Python. Statistical tests are
756 indicated in the corresponding figure legends. The immunoblot analyses were repeated twice
757 and showed consistent results. p values below 0.05 were considered statistically significant
758 and are indicated in the figures as follows: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$
759 0.0001

Conflict of Interest statement:

761 DB, MH, MN, and YJB are inventors of "Mass-tag labeling of the cellular secretome", as
762 described in PCT patent application PCT/EP2023/062049. The remaining authors declare no
763 conflicts of interest.

Acknowledgments

DB and MH are funded by the German Research Foundation (DFG) SFB1530-455784452
(subprojects B01, Z01, Z02) and the José Carreras Leukemia Foundation (Grant DJCLS
04R/2021). NA is supported by the Cancer Research Center Cologne Essen (CCCE) as well
as by the German Research Foundation (DFG) SFB1530-455784452 (subproject Z03) and
Project-ID 497777992. DB, DS, RTU, AVL, and HJ are supported by the Mildred Scheel
Nachwuchszentrum Grant 70113307 from the German Cancer Aid. RDJ is funded by the

German Research Foundation (DFG) FOR5504-496650118 (subproject TP05), JA2439/4-1, and SFB1530-455784452 (subproject C02), by the Ministry for Culture and Science of North Rhine-Westphalia (NW21-062A CANTAR), and by the Behrens-Weise-Foundation (for the Improvement of Human Health). DS was supported by an MD Research Stipend of the Else Kröner Forschungskolleg Clonal Evolution in Cancer, University Hospital Cologne, Cologne, Germany. DB is supported by the Exzellenz Initiieren (E.I.) – Stiftung Kölner Krebsforschung. DB and BE are supported by the German Federal Ministry of Education and Research (BMBF; project 01KU2108) within the ERA PerMed framework. We thank Dr. Dirk Strunk (Paracelsus Medical University) for his scientific insights and guidance. We also thank Dr. Jan van Deun (University of Erlangen-Nuremberg) and Dr. Anna Kashkanova (Max Planck Institute for the Science of Light, Erlangen) for their experimental support. Additionally, we appreciate Dr. Christian Jüngst and Beatrix Martiny (CECAD Imaging Facility, University of Cologne) for their experimental assistance. (Imaging) Mass cytometric data acquisition was carried out by Petra Hofmann (Department I of Internal Medicine) and Marion Müller (Institute of Pathology) at the CyTOF Facility of the Medical Faculty, University of Cologne, supported by the DFG-funded CRC1530 (Project Z02). Immunogold transmission electron microscopy was performed at the CECAD Imaging Facility using a Jeol JEM 2100 Plus, funded by DFG-INST 216/793-1 FUGG. We additionally thank the Regional Computing Center of the University of Cologne for providing computing time on the DFG-funded High Performance Computing system CHEOPS (Funding number: DFG-INST 216/512-1 FUGG) and for support. The NanoSIMS measurements were performed at the NanoSIMS Sweden Facility at the University of Gothenburg, SciLifeLab Swedish infrastructure. Graphical representations were created with BioRender (https://BioRender.com/hlo8aeq).

References

1. Buzas, E. I. The roles of extracellular vesicles in the immune system. *Nat Rev Immunol* **23**, 236–250 (2023).

2. Kalluri, R. The biology and function of extracellular vesicles in immune response and immunity. *Immunity* **57**, 1752–1768 (2024).

3. Alashkar Alhamwe, B. *et al.* BAG6 restricts pancreatic cancer progression by suppressing the release of IL33-presenting extracellular vesicles and the activation of mast cells. *Cell Mol Immunol* **21**, 918–931 (2024).

4. van Niel, G., D'Angelo, G. & Raposo, G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol* **19**, 213–228 (2018).

5. Kalluri, R. & McAndrews, K. M. The role of extracellular vesicles in cancer. *Cell* **186**, 1610–1626 (2023).

6. Tran, H. L. *et al.* Extracellular Vesicles for Clinical Diagnostics: From Bulk Measurements to Single-Vesicle Analysis. *ACS Nano* **19**, 28021–28109 (2025).

7. Maas, S. L. N. *et al.* Possibilities and limitations of current technologies for quantification of

biological extracellular vesicles and synthetic mimics. *Journal of Controlled Release* **200**, 87–96

(2015).

8. Melling, G. E. Fluorescent labeling of EVs: strengths and limitations. *Sci Rep* (2022).

9. Welsh, J. A. *et al.* Minimal information for studies of extracellular vesicles (MISEV2023): From basic

to advanced approaches. *J Extracell Vesicles* **13**, e12404 (2024).

10.   Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a

position statement of the International Society for Extracellular Vesicles and update of the

MISEV2014 guidelines. *J Extracell Vesicles* **7**, 1535750 (2018).

11.   Seliger, B. & Massa, C. CyTOF as a suitable tool for stratification and monitoring of cancer

patients. *Journal of Translational Medicine* **23**, 724 (2025).

12.   Li, Y.-J., Wu, J.-Y., Wang, J.-M., Hu, X.-B. & Xiang, D.-X. Emerging strategies for labeling and

tracking of extracellular vesicles. *J Control Release* **328**, 141–159 (2020).

13.   Haines, D. S. Confounding lipid dye particles complicate EV labeling studies. *J Extracell*

Vesicles (2025).

14.   Tertel, T. *et al.* Imaging flow cytometry challenges the usefulness of classically used

extracellular vesicle labeling dyes and qualifies the novel dye Exoria for the labeling of

mesenchymal stromal cell-extracellular vesicle preparations. *Cytotherapy* **24**, 619–628 (2022).

15.   Dehghani, M., Gulvin, S. M., Flax, J. & Gaborski, T. R. Systematic Evaluation of PKH Labelling

on Extracellular Vesicle Size by Nanoparticle Tracking Analysis. *Sci Rep* **10**, 9533 (2020).

16.   Bachurski, D. & Hallek, M. Biological and technical complexities in analyzing extracellular

vesicle immune interactions in B-cell malignancies. *FEBS Lett* https://doi.org/10.1002/1873-

3468.70102 (2025) doi:10.1002/1873-3468.70102.

17.   Jia Bu, Y. & Nitz, M. Incorporation of TePhe into Expressed Proteins is Minimally Perturbing.

Chembiochem **22**, 2449–2456 (2021).

18.   Bassan, J. *et al.* TePhe, a tellurium-containing phenylalanine mimic, allows monitoring of

protein synthesis in vivo with mass cytometry. *Proc Natl Acad Sci U S A* **116**, 8155–8160 (2019).

19.   Lange, I. Correlative TEM–nanoSIMS for subcellular isotopic localization. *PLOS One* (2021).

20.   Giesen, C. Imaging mass cytometry by multiplexed metal-tagged antibodies. *Nat Methods*

(2014).

21.   Bendall, S. C. Single-cell mass cytometry of immune system. *Science* (2011).

22.   Zhou, J. *et al.* STAT5 and STAT3 balance shapes dendritic cell function and tumour immunity.

Nature **643**, 519–528 (2025).

23.   Kaneshige, A. *et al.* A selective small-molecule STAT5 PROTAC degrader capable of achieving

tumor regression in vivo. *Nat Chem Biol* **19**, 703–711 (2023).

24.   Monguió-Tortajada, M., Gálvez-Montón, C., Bayes-Genis, A., Roura, S. & Borràs, F. E.

Extracellular vesicle isolation methods: rising impact of size-exclusion chromatography. *Cell Mol*

Life Sci **76**, 2369–2382 (2019).

25.   Lovrić, J. *et al.* Correlative High-Resolution Imaging of Iron Uptake in Lung Macrophages. *Anal*

Chem **94**, 12798–12806 (2022).

26.   Boudna, M. *et al.* Strategies for labelling of exogenous and endogenous extracellular vesicles

and their application for in vitro and in vivo functional studies. *Cell Communication and Signaling*

22, 171 (2024).

27.   Bachurski, D. *et al.* Extracellular vesicle measurements with nanoparticle tracking analysis -

An accuracy and repeatability comparison between NanoSight NS300 and ZetaView. *J Extracell*

Vesicles **8**, 1596016 (2019).

28.   Loconte, L. *et al.* Detection of the interactions of tumour derived extracellular vesicles with

immune cells is dependent on EV-labelling methods. *J Extracell Vesicles* **12**, e12384 (2023).

29.   Breiman, L. Random Forests. *Machine Learning* **45**, 5–32 (2001).

30.   Liu, J., Zhang, X., Cheng, Y. & Cao, X. Dendritic cell migration in inflammation and immunity.

Cell Mol Immunol **18**, 2461–2471 (2021).

31.   Carrette, F. & Surh, C. D. IL-7 signaling and CD127 receptor regulation in the control of T cell

homeostasis. *Semin Immunol* **24**, 209–217 (2012).

32.   Willis, L. M. Tellurium-based mass cytometry barcode for live and fixed cells. *Cytometry A*

(2018).

33.   Jeppesen, D. K. *et al.* Reassessment of Exosome Composition. *Cell* **177**, 428-445.e18 (2019).

34.   Liam-Or, R. *et al.* Cellular uptake and in vivo distribution of mesenchymal-stem-cell-derived

extracellular vesicles are protein corona dependent. *Nat. Nanotechnol.* **19**, 846–855 (2024).

35.   Zhang, B. *et al.* CD127 imprints functional heterogeneity to diversify monocyte responses in

inflammatory diseases. *J Exp Med* **219**, e20211191 (2022).

36.   Jurgielewicz, B. J., Yao, Y. & Stice, S. L. Kinetics and Specificity of HEK293T Extracellular Vesicle

Uptake using Imaging Flow Cytometry. *Nanoscale Res Lett* **15**, 170 (2020).

37.   Lombardo, G. *et al.* Activated Stat5 trafficking Via Endothelial Cell-derived Extracellular

Vesicles Controls IL-3 Pro-angiogenic Paracrine Action. *Sci Rep* **6**, 25689 (2016).

38.   Lopatina, T. *et al.* IL-3 signalling in the tumour microenvironment shapes the immune

response via tumour endothelial cell-derived extracellular vesicles. *Pharmacol Res* **179**, 106206

(2022).

39.   Poveda, E. *et al.* Interleukin 18 (IL-18) and IL-3 in Extracellular Vesicles: Biomarkers for

Durable Elite Control of HIV-1. *J Infect Dis* **227**, 1381–1385 (2023).

40.   Bunn, K. E., Giese-Byrne, B. G., Blatt, A. M., Newcomb, D. C. & Pua, H. H. Mouse TH2 cell

extracellular vesicles promote eosinophil survival through the surface cytokine cargo IL-3. *Journal*

of Allergy and Clinical Immunology **156**, 1053–1065 (2025).

41.   Monaghan, K. L. *et al.* Tetramerization of STAT5 regulates monocyte differentiation and the

dextran sulfate sodium-induced colitis in mice. *Front Immunol* **14**, 1117828 (2023).

42.   Bai, L. *et al.* A Potent and Selective Small-molecule Degrader of STAT3 Achieves Complete

Tumor Regression in vivo. *Cancer Cell* **36**, 498-511.e17 (2019).

43.   Huang, Z.-W. *et al.* STAT5 promotes PD-L1 expression by facilitating histone lactylation to drive

immunosuppression in acute myeloid leukemia. *Sig Transduct Target Ther* **8**, 391 (2023).

44.   Haderk, F. *et al.* Tumor-derived exosomes modulate PD-L1 expression in monocytes. *Sci*

Immunol **2**, eaah5509 (2017).

45.   Stahl, D. *et al.* CSF1R+ myeloid-monocytic cells drive CAR-T cell resistance in aggressive B cell

lymphoma. *Cancer Cell* **43**, 1476-1494.e10 (2025).

46.   Grześ, K. M. IL-3/JAK2-STAT5 axis programs human pDC/myeloid responses. *Immunity* (2021).

47.   Zeng, H. *et al.* Exosomal PD-L1 promotes the formation of an immunosuppressive

microenvironment in gastric diffuse large B-cell lymphoma. *Oncol Rep* **49**, 88 (2023).

48.   Kumar, M. A. *et al.* Extracellular vesicles as tools and targets in therapy for diseases. *Sig*

Transduct Target Ther **9**, 27 (2024).

49.   Mentkowski, K. I., Snitzer, J. D., Rusnak, S. & Lang, J. K. Therapeutic Potential of Engineered

Extracellular Vesicles. *AAPS J* **20**, 50 (2018).

50.   Vergauwen, G. *et al.* Confounding factors of ultrafiltration and protein analysis in extracellular

vesicle research. *Sci Rep* **7**, 2704 (2017).

51.   Wallis, R., Mizen, H. & Bishop, C. L. The bright and dark side of extracellular vesicles in the

senescence-associated secretory phenotype. *Mechanisms of Ageing and Development* **189**,

111263 (2020).

52.   Vom Stein, A. F. *et al.* LYN kinase programs stromal fibroblasts to facilitate leukemic survival

via regulation of c-JUN and THBS1. *Nat Commun* **14**, 1330 (2023).

53.   Chen, C.-C., Kobayashi, T., Iijima, K., Hsu, F.-C. & Kita, H. IL-33 dysregulates regulatory T (Treg)

cells and impairs established immunological tolerance in the lungs. *J Allergy Clin Immunol* **140**,

1351-1363.e7 (2017).

54.   Peng, X., Zhao, M., Gao, L., Sen, R. & MacGlashan, D. Identifying regulatory pathways of

spleen tyrosine kinase expression in human basophils. *Journal of Allergy and Clinical Immunology*

145, 947–957 (2020).

55.   Singh, K. *et al.* IL-7–mediated expansion of autologous lymphocytes increases CD8+ VLA-4

expression and accumulation in glioblastoma models. *J Clin Invest* **135**, e181471.

56.   Görgens, A. *et al.* Identification of storage conditions stabilizing extracellular vesicles

preparations. *J Extracell Vesicles* **11**, e12238 (2022).

57.   Xiao, X. *et al.* Dice-XMBD: Deep Learning-Based Cell Segmentation for Imaging Mass

Cytometry. *Front. Genet.* **12**, (2021).

58.   Greenwald, N. F. *et al.* Whole-cell segmentation of tissue images with human-level

performance using large-scale data annotation and deep learning. *Nat Biotechnol* **40**, 555–565

(2022).

59.   Perez-Riverol, Y. *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids Research*

53, D543–D553 (2025).

60.   Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: neural networks

and interference correction enable deep proteome coverage in high throughput. *Nat Methods* **17**,

41–44 (2020).

61.   Mastronarde, D. N. Automated electron microscope tomography using robust prediction of

specimen movements. *Journal of Structural Biology* **152**, 36–51 (2005).

62.   Paul-Gilloteaux, P. *et al.* eC-CLEM: flexible multidimensional registration software for

correlative microscopies. *Nat Methods* **14**, 102–103 (2017).

63.   Chevrier, S. *et al.* Compensation of Signal Spillover in Suspension and Imaging Mass

Cytometry. *Cell Syst* **6**, 612-620.e5 (2018).

64.   Van Gassen, S. *et al.* FlowSOM: Using self-organizing maps for visualization and interpretation

of cytometry data. *Cytometry A* **87**, 636–645 (2015).

65.   Yoshida, M. *et al.* Local and systemic responses to SARS-CoV-2 infection in children and

adults. *Nature* **602**, 321–327 (2022).

66.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data

analysis. *Genome Biology* **19**, 15 (2018).

67.   Maaten, L. van der & Hinton, G. Visualizing Data using t-SNE. *Journal of Machine Learning*

Research **9**, 2579–2605 (2008).

68.   EV-TRACK Consortium *et al.* EV-TRACK: transparent reporting and centralizing knowledge in

extracellular vesicle research. *Nat Methods* **14**, 228–232 (2017).

Figure legends

Figure 1 Single-cell tracking of tellurium-labeled EV proteomes.

a, Schematic of the TeLEV (tellurium-based labeling of the extracellular vesicle proteome) mass-tag labeling workflow yielding Te-labeled MBC-EVs. Primary chronic lymphocytic leukemia cells or cell lines were cultured for 48 h in chemically defined, serum-free medium supplemented with 50 µM L-2-tellurienylalanine (TePhe). Conditioned medium was collected, 0.22 µm filtered, concentrated by 10 kDa ultrafiltration (Amicon 10 kDa RC), and fractionated to separate EVs from soluble secreted proteins by size-exclusion chromatography (35 nm resin; IZON qEV original 2).

b, Mass cytometry of PBMC recipient cells showing a two-dimensional t-SNE embedding colored by the TeLEV signal originating from primary CLL donors. Columns show PBS-HAT vehicle control (left), CLL-derived MBC-EVs (middle), and Te-labeled soluble proteins (right). *HAT* denotes HEPES, human albumin, and trehalose. Data were acquired on a Helios mass cytometer. Primary MBC-EVs and secreted proteins shown in Figure 1 were isolated from 25 mL of cell-conditioned medium (from 25M primary CLL cells over 48 hours), resulting in 1.4 x 10^{10} MBC-EVs used per EV condition.

c, Imaging mass cytometry of PBMC recipients exposed to PBS–HAT control, CLL-derived MBC-EVs, or Te-labeled secreted proteins. Representative fields are shown (acquired on a Hyperion imaging mass cytometer). Scale bar, 200 µm.

d, Correlative ultrastructural and isotopic mapping of PBMC recipients. Transmission electron microscopy (TEM) on 70 nm sections was followed by nanoSIMS of the same regions of the same grids (NanoSIMS 50L; CAMECA, Gennevilliers, France) to detect tellurium enrichment. Arrows indicate TeLEV signal (heat-map overlay) localized predominantly in heterochromatin of myeloid recipients after Te-EV exposure (middle) and in the cytoplasm after exposure to Te-labeled soluble proteins (right). Scale bar, 2 µm.

Figure 2 TeLEV does not perturb EV proteomes.

a, Triplex immunogold labeling of tetraspanins on MBC-EVs. Primary CLL-derived MBC-EVs and unlabeled control MBC-EVs were stained with anti-tetraspanin antibodies coupled to distinct gold sizes: CD9 (10 nm), CD63 (20 nm), and CD81 (40 nm), with and without uranyl-acetate contrasting. Representative micrographs are shown. Scale bars, 100 nm (100,000×) and 200 nm (30,000×). MBC-EVs and secreted proteins shown in Figure 2 were isolated from 50 mL of cell-conditioned medium (from 50M primary CLL cells over 48 hours).

b, Particle sizing of purified primary MBC-EVs. Nanoparticle tracking analysis (NTA) reported particle size distributions. Statistical significance was evaluated using the Kolmogorov-Smirnov test. Data are represented as mean + SEM.

c, Size-exclusion chromatography (SEC) fractionation of conditioned medium. 1 mL fractions were collected and analyzed by NTA (particle counts) and BCA (protein). F1–F2, void volume; F3–F5, EV-enriched (MBC-EVs) fractions; F9–F11, secreted-protein fractions. SEC used 35 nm IZON resin following 10 kDa ultrafiltration. Data are represented as mean ± SEM.

d, NTA particle counts for pooled primary MBC-EV fractions (F3–F5). No significant changes were identified based on Wilcoxon matched pairs signed rank test (p value = 0.8457).

e, NTA particle diameters for pooled primary MBC-EV fractions (F3–F5). No significant changes were identified based on Wilcoxon matched pairs signed rank test (p value = 0.3008).

f, Immunoblot analysis of SEC fractions from primary-leukemic donors. Fractions were concentrated and loaded at equal volumes; total-protein staining shows higher abundance in soluble-protein fractions. EV and non-EV markers are shown for TeLEV and control conditions. To avoid overloading of the gel, a 10 µg cut-off of protein fractions was chosen, which was not exceeded by the EV fractions.

g, Proteomics comparison of isolated primary MBC-EVs and soluble secreted proteins with and without TeLEV treatment. Heatmap shows log_2 intensity values for selected marker proteins across conditions (n = 3 paired CLL donors).

h, Pearson correlation of protein intensities from isolated primary MBC-EVs (left) and secreted proteins (right), showing no clustering based on TeLEV treatment.

i, Bar plots of log_2 intensities from isolated primary MBC-EVs (left) and secreted proteins (right) in control and TeLEV-treated samples.

j, Mean number of total identified proteins (left) and EV proteins (right) in control and TeLEV-treated samples of isolated primary MBC-EVs and secreted proteins. Data are represented as mean ± SEM.

k, Principal component analysis of isolated primary MBC-EVs (left) and secreted proteins (right), showing no grouping based on TeLEV treatment.

l, Volcano plot showing differences of log_2 intensities between TeLEV-treated and control samples in isolated primary MBC-EVs. No significantly abundant proteins were identified (Student's t test: FDR < 0.05, s0 > 0.1). GO term-annotated "extracellular vesicle" (GO:1903561) proteins are marked in light blue.

m, Boxplot of mean log_2 intensities for control and TeLEV-treated samples of isolated primary MBC-EVs and secreted proteins based on the GO term "vesicle membrane" (GO:1903561) and MGI GO EV annotation (GO:1903561). Data are represented as mean ± SEM. No significantly changed proteins were identified based on unpaired t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Figure 3 Single-cell mapping of PBMC uptake of Te-labeled EVs and soluble proteins secreted by CLL cells.

a, t-SNE projection of PBMC recipient cells (758,260 cells in total) of PBS-HAT control (329,774 cells), primary MBC-EVs (226,157cells), and matched Te-labeled secreted proteins (229,329 cells), coloured according to cell type. Numerical annotations indicate the relative proportions of each cell lineage. Primary MBC-EVs and secreted proteins shown in Figure 3 were isolated from 25 mL of cell-conditioned medium (from 25M primary CLL cells over 48 hours), resulting in 1.4×10^{10} MBC-EVs used per EV condition.

b, Dot plot depicting the expression of selected markers across major cell types and their subtypes from three conditions, PBS-HAT control, primary Te-labeled MBC-EVs, and matched Te-labeled secreted proteins. Dot size corresponds to the proportion of cells expressing each marker within a given cell type, and color intensity indicates the mean expression level.

c, Comparison of TeLEV signal distribution across major cell types and their subtypes in each indicated condition, with colors representing the average TeLEV signal level in each population. Bars above the plots show the relative proportions of each population in the corresponding condition (left). White lines denote the median, edges the IQR, and whiskers either 1.5 × IQR or minima/maxima (if no point exceeded 1.5 × IQR; minima/maxima are indicated by the violin plot range). t-SNE projection of cells for each condition showing TeLEV signal (middle). Two-dimensional density distributions of cells in t-SNE embedding space, stratified by conditions (right).

d, Feature importance of EV-to-protein log ratios as determined by the random forest classifier, where higher positive values indicate stronger discriminative power in primary MBC-EVs compared to matched Te-labeled secreted proteins.

Figure 4 Pan-immune single-cell atlas identifies an interleukin-receptor triad induced by B-cell malignant EVs.

a, t-SNE projection of PBMC recipient cells across 16 conditions (PBS-HAT control plus 15 EV sources, in total 2,977,094 cells), colored according to cell type. Numerical annotations indicate the relative proportions of each cell lineage. EVs shown in Figure 4 were isolated from 25 mL of cell-conditioned medium (from 25M primary CLL cells or cell lines over 48 hours). The EV uptake conditions were normalized according to the input volume of cell-conditioned media (25 mL, 25M cells over 48 hours), and EV counts and buffer volumes are reported in Supplementary Table 2.

b, t-SNE projection of PBMC recipients across 16 conditions showing TeLEV signal.

c, Fold changes in cell type proportions across conditions, estimated using a Poisson generalized linear mixed model that accounts for confounding factors (Methods). Dot size represents the probability of change, measured as the local true sign rate (LTSR).

d, t-SNE projection of myeloid cells across 16 conditions (276, 867 cells), colored according to subtypes.

e, t-SNE plots showing expression of monocytic and DC lineage markers and interleukin receptors on myeloid cells.

f, Two-dimensional density distributions of myeloid cells in t-SNE embedding space, stratified by conditions.

g, A cell type–color–coded ternary plot depicting expression levels of CD25, CD123, and CD127. Circle size reflects the average TeLEV signal level within each cell type. Colors correspond to those in the legend shown in (a)

Figure 5 Temporal dynamics and functional consequence of the interleukin receptor triad.

a, t-SNE projection of PBMC recipient cells across 11 time points (0-48h, in total 1,512,100 cells) of EV uptake, colored according to cell type. Numerical annotations indicate the relative proportions of each cell lineage. Primary MBC-EVs and secreted proteins shown in Figure 5a-g were isolated from 4,2 mL of cell-conditioned medium (from 4.2M primary CLL cells over 48 hours), resulting in 2.7×10^9 EVs used per time point condition.

b, t-SNE plots showing TeLEV signal, CD25, CD127, and CD123 expression at selected time points (columns).

c, Stacked bar plot depicting changes in cell type composition over 48h. Colors correspond to those in the legend shown in (a).

d, Comparison of TeLEV signal distribution across major cell types and their subtypes at 16 h, with colors representing the average TeLEV signal level in each population. Bars above the plots show the relative proportions of each population. White lines denote the median, edges the IQR, and whiskers either 1.5 × IQR or minima/maxima (if no point exceeded 1.5 × IQR; minima/maxima are indicated by the violin plot range).

e, Comparison of TeLEV signal, CD25, CD123, and CD127 distributions within the Monocyte population (including Non-classical Monocytes, Intermediate Monocytes, Classical Monocytes, IL2R/IL7R$^+$ Monocytes, and CD197$^+$ IL2R/IL7R$^+$ Monocytes) over 48 h. Statistical significance across all time points for each marker was evaluated using the Kruskal-Wallis test. Pairwise comparisons between 00 h and 00 h 15 min were performed using a one-sided Mann–Whitney U test. *p-values* < 0.05 were considered significant and are indicated in the figures as follows: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

f, Line plots depicting the mean expression levels of TeLEV signal, CD25, CD123, and CD127 within the Monocytes during the first 2 hours of EV uptake. Error bars represent mean ± SEM.

g, The same analysis as in (e) for mDCs (left) and pDCs (right).

h, Live-cell immunocytochemistry of purified monocytes exposed to increasing doses of Ramos-derived EVs. Representative frames from day 2 and day 7 are shown. CD25 (Alexa Fluor 488; green) and Cytotox NIR (blue; loss of membrane integrity) are displayed after automated single-cell segmentation (Incucyte built-in). Cells were seeded at 25,000 per well and imaged every 2 h for 7 days using an Incucyte SX5 (20× objective). Scale bar, 200 µm. EVs and secreted proteins shown in Figure 5h-p were isolated from 1.5 mL of cell-conditioned medium (from 1.5M Ramos cells over 48 hours), resulting in 9 x 10^8 EVs used per EV condition (3X condition, Figure 6h,i,j).

i, Single-cell quantification of CD25 (Alexa Fluor 488) fluorescence over 7 days corresponding to (h). Values are unnormalized arbitrary units; data shown as mean ± SEM; n = 4 fields of view.

j, Flow cytometric contour plots of purified monocytes after 16 h of EV exposure across a dose titration (see Figure legend 6i). CD25 (APC), CD123 (PE), and CD127 (FITC) were measured on a Miltenyi MACSQuant X with compensation using BioLegend compensation beads. DAPI was used as a surface-flow viability dye.

k, Live-cell immunocytochemistry of EV-exposed monocytes stimulated with cytokines. Representative images from day 7 are shown for IL-2 (20 ng mL⁻¹), IL-3 (5 ng mL⁻¹), or, IL-7 (25 ng mL⁻¹), or IL-2 + IL-3, IL-2 + IL-7, IL-3 + IL-7, or IL-2 + IL-3 + IL-7 in RPMI with serum. CD25 (Alexa Fluor 488; green) and Cytotox NIR (blue) are shown after automated segmentation and background subtraction. Imaging on Incucyte SX5 (20×, 2 h intervals). Scale bar, 200 µm.

l, Immunogold transmission electron microscopy of Ramos-EVs labeled for IL-3, CD123, and CD127 alongside an isotype control. Scale bar, 100 nm.

m, Intracellular phospho-flow cytometry of purified monocytes incubated with EVs, showing
total STAT5 (Alexa Fluor 488), total STAT3 (PE), pSTAT5 (PE-Cy7), and pSTAT3 (BV421). EV
exposure was terminated by immediate formaldehyde fixation after 16 h prior to staining.
Cleaved caspase was included as an apoptosis marker. Acquisition on a Miltenyi MACSQuant
X with bead-based compensation of all markers.

n, Live-cell immunocytochemistry of EV-exposed healthy-donor PBMCs stimulated for 2 days
with IL-2 (20 ng mL^{-1}), IL-3 (5 ng mL^{-1}), or IL-7 (25 ng mL^{-1}) in RPMI with serum.
Representative images are shown (Incucyte SX5, 20×, 2 h intervals). Scale bar, 200 μm.

o, Live-cell immunocytochemistry of EV-exposed healthy-donor PBMCs treated for 7 days with
a STAT5 degrader (AK-2292) or anti-CD25 (Basiliximab), compared with DMSO and isotype
controls. CD25 (Alexa Fluor 488; green), CD127 (PE; red), and Cytotox NIR (blue) are shown
after automated segmentation and background subtraction. Cells were seeded at 50,000 per
well and imaged on an Incucyte SX5 (20×, every 2 h). Scale bar, 200 μm.

p, Flow-cytometric contour plots of healthy-donor PBMCs after 48 h of EV exposure across a
dose titration (1× = 3×10^8 EVs per 10^5 cells), showing CD25 (APC), CD123 (PE), and CD127
(FITC). Acquisition and compensation as in (m).

1123

Figure 6 EV-driven immunomodulation by STAT5-regulated interleukin receptor triad.

a, t-SNE projection of 16 conditions (in total 3,590,520 cells) coloured according to cell type.
Numerical annotations indicate the relative proportions of each cell lineage. MBC-EVs shown
in Figure 6 were isolated from 8 mL of cell-conditioned medium (from 8M Ramos cells over 48
hours), resulting in 5 x 10^9 EVs used per EV condition.

b, Stacked bar plot depicting changes in cell type composition. Colors correspond to those in
the legend shown in (a).

c, Comparison of TeLEV signal distribution across major cell types and their subtypes. Shown
are EVs (217,211 cells), STAT5d + EVs (226,884 cells), and STAT3d + EVs (206,018 cells)
conditions, with colors representing the average TeLEV signal level in each population. Bars
above the plots show the relative proportions of each population in the corresponding condition
(left). White lines denote the median, edges the IQR and whiskers either 1.5 × IQR or
minima/maxima (if no point exceeded 1.5 × IQR; minima/maxima are indicated by the violin
plot range). t-SNE projection of cells for each condition showing TeLEV signal (middle). Two-
dimensional density distributions of cells in t-SNE embedding space, stratified by conditions
(right).

d, Comparison of TeLEV signal distribution across three conditions shown in (c). Pairwise
comparisons were performed using a two-sided Mann–Whitney U test. ****p < 0.0001.

e, t-SNE projection of myeloid population of 16 conditions (865,413 cells), colored according
to subtypes.

f, t-SNE plots showing expression of monocytic and DC lineage markers and interleukin
receptors on myeloid cells.

g-k, t-SNE plots showing CD25, CD127, CD123, BCL-2, and PD-L1 expressions across the
indicated conditions.

l, Comparison of PD-L1 expression distribution across the indicated conditions in Monocytes, mo-derived DCs, mDCs, and pDCs. For each cell type, statistical significance across all conditions was evaluated using the Kruskal-Wallis test.****$p < 0.0001$.

m, Same as in (l) for BCL-2.

n-p, Comparison of CD25 (n), Ki-67 (o), and CD69 (p) expression distributions across major cell types and their subtypes, with colors representing the average respective marker expression level in each population. Shown are IL-2 (238,611 cells) and IL-2 + EVs (207,300 cells) conditions.

Supplemental figure legends

Figure S1 High-dimensional profiling of MBC-EV recipients, related to Figure 1.

a-c, t-SNE plots showing selected marker expressions for the three conditions. PBS-HAT control, primary Te-labeled MBC-EVs, and matched Te-labeled secreted proteins.

Figure S2 Characterization of Te-labeled CLL-derived EVs and secreted proteins across immune cell populations, related to Figure 3.

a, Dot plot depicting the expression of selected markers across major cell types and their subtypes for three conditions: PBS-HAT control (top, 329,774 cells), primary Te-labeled MBC-EVs (middle, 226,157 cells), and Te-labeled secreted proteins (bottom, 229,329 cells). Dot size corresponds to the proportion of cells expressing each marker within a given cell type, and color intensity indicates the mean expression level.

b, Feature importance of surface markers for distinguishing the primary MBC-EV from control data, as determined by a random forest classifier (left). The same analysis as in (b) is shown for matched Te-labeled secreted proteins (right).

c, TeLEV signal histograms for primary MBC-EVs (top) and matched secreted proteins (bottom). The dashed lines indicate the 80th percentile cut-off.

d, t-SNE plots showing CD25, CD127, and CD123 expressions for the three conditions.

Figure S3 Comparison of EV uptake from primary and cell line-derived EV sources.

EV uptake, related to Figure 4.

a, t-SNE projection of PBMC recipient cells for 16 conditions (PBS-HAT control plus 15 EV sources) showing TeLEV signal.

b, Two-dimensional density distributions of cells in t-SNE embedding space, stratified by conditions.

c, t-SNE projection of PBMC recipients of PBS-HAT control (200,468 cells), unlabeled control EVs (200,629 cells), and Te-labeled MBC-EVs (191,964 cells) from primary CLL cells coloured according to cell type (column 1). Two-dimensional density distributions of cells in t-SNE embedding space, stratified by conditions (column 2). t-SNE plots showing CD25, CD123, and CD127 expression, and TeLEV signal (columns 3-6).

1186 d, As in c showing the myeloid compartment. Shown are PBS-HAT control (17,445 cells),
1187 unlabeled control MBC-EVs (19,003 cells), and Te-labeled MBC-EVs (15,300 cells) from
1188 primary CLL cells.

1189

Figure S4 Time-resolved uptake of Te-labeled primary MBC-EVs and MBC-secreted
proteins, related to Figure 5.

a, t-SNE projection of PBMC recipient cells showing uptake dynamics across 11 time points
for primary Te-labeled MBC-EVs and 11 time points for matched secreted proteins (in total
2,900,351 cells), coloured according to cell type. Numerical annotations indicate the relative
proportions of each cell lineage.

b, t-SNE plots showing TeLEV expressions for 11 time points of MBC-EV uptake.

c, t-SNE plots showing TeLEV expressions for 11 time points of Te-labeled secreted proteins
uptake.

d, t-SNE plots showing TeLEV expression for primary MBC-EV uptake (left, 1,512,100 cells)
and matched Te-labeled secreted protein uptake (right, 1,388,251 cells) across concatenated
time points.

e, Two-dimensional density distributions of cells in t-SNE space for MBC-EV (left) and
matched secreted protein conditions (right).

f, Fold changes in cell type proportions across 11 time points of EV uptake, estimated using a
Poisson generalized linear mixed model. Dot size represents the probability of change,
measured as the local true sign rate (LTSR).

1207

Figure S5 Global immune cell expression of the interleukin receptor triad, related to
Figure 6.

a, t-SNE projection of cells showing CD69 expression. Shown are MBC-EVs (217,211 cells),
STAT5d + MBC-EVs (226,884 cells), and STAT3d + MBC-EVs (206,018 cells) conditions.

b, A cell type–color–coded ternary plot depicting expression levels of CD25, CD123, and
CD127. Circle size reflects the average TeLEV level within each cell type.

c, t-SNE projection of the non-myeloid population of 16 conditions (2,725,107 cells), colored
according to cell types. Numerical annotations indicate the relative proportions of each cell
lineage.

d-f, t-SNE plots showing CD25 (d), CD123 (e), and CD127 (f) expressions on the non-
myeloid population across the indicated conditions.

g-i, Comparison of CD25 (g), CD123 (h), and CD127 (i) expression distributions across the
indicated conditions in Monocytes, mo-derived DCs, mDCs, and pDCs. For each cell type,
statistical significance across all conditions was evaluated using the Kruskal-Wallis test. ****p
< 0.0001. White lines denote the median, edges the IQR, and whiskers either 1.5 × IQR or
minima/maxima (if no point exceeded 1.5 × IQR; minima/maxima are indicated by the violin
plot range).

1225

1226 **Data S1 Pan-immune cell atlas of EV uptake from cell lines and primary EVs, related to**
1227 **Figure 4.**

1228   t-SNE projection of major cell types per condition. Shown are cell type overlays, two-
1229   dimensional density distributions in t-SNE space, and overlays of selected marker
1230   expressions.

1231

1232   **Data S2 Immune recipient cell mapping of primary MBC-EV and MBC-secreted protein**
1233   **uptake, related to Figure 3.**

1234   t-SNE projection of major cell types per condition. Shown are cell type overlays, two-
1235   dimensional density distributions in t-SNE space, and overlays of selected marker
1236   expressions of primary Te-labeled MBC-EV and matched secreted protein uptake.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6