Differential Control of HIV-1 Replication by IFN-α14 Compared to IFN-α2 Relates to

Differences in the Modulation of Host Antiretroviral Restriction Factors

Saurav S. Rout[1], Madeline T.E. Stewart[1], Nathan B. Seidel[1], Ulf Dittmer[2], Kathrin Sutter[2] and

Kerry J. Lavender[1#]

[1]Department of Biochemistry, Microbiology and Immunology, College of Medicine, University

of Saskatchewan, Saskatoon, SK, S7N 5E5, Canada

[2]Institute of Virology, University Hospital Essen, University of Duisburg-Essen, Essen, Germany.

[#]Send correspondence to Kerry J. Lavender kerry.lavender@usask.ca

Running Head: IFN-α subtype specific control of restriction factors

Keywords: HIV-1, restriction factor, interferon-alpha, subtypes

Abstract word count: 250

Word count: 4966

ABSTRACT

Type I IFN, including IFN-α, induces the expression of antiviral restriction factors that can

interfere with multiple steps of the HIV-1 replication cycle. Humans have 13 IFN-α genes which

encode 12 different IFN-α subtypes. Our previous work in HIV-1 infected humanized mice showed

that IFN-α14 treatment more potently controlled HIV-1 than treatment with the clinically approved

IFN-α2 subtype. However, the mechanisms behind the more potent control of HIV-1 by IFN-α14

are unknown. The IFN-α14 subtype is known to more potently induce the expression of the

restriction factors MX2 and ISG15 and increased APOBEC3G signature mutations *in vivo*

compared to IFN-α2. To study the importance of each of these restriction factors in mediating the

potent control of HIV-1, we used a CRISPR-Cas9 lentivirus system to create stable knockouts in

the MT4C5 cell line that is susceptible to HIV-1 but does not produce measurable amounts of

endogenous IFN-α. Knock out of ISG15, but not MX2, eliminated differences in viral suppression

after IFN-α14 and IFN-α2 treatment. Similarly, APOBEC3G deletion eliminated differences in

viral suppression and the number of infectious particles produced after IFN-α14 and IFN-α2

treatment. Furthermore, APOBEC3G deletion resulted in significantly fewer GG→AG mutations

in viral DNA isolated from target cells incubated with supernatant from IFN-α14 treated groups.

However, APOBEC3G knock out did not result in significant increases in vDNA compared to the

wild type in any experimental group. Overall, elimination of APOBEC3G and ISG15 impaired

IFN-α14–mediated suppression of HIV-1, highlighting them as downstream effectors of IFN-α14's

more potent anti-HIV-1 activity.

IMPORTANCE

This study uncovers the molecular basis for the more potent antiviral activity of IFN-α14 compared

to the clinically used IFN-α2 subtype against HIV-1. Although interferons are known to induce

numerous restriction factors, the mechanisms underlying subtype-specific antiviral potency

remained unclear. By using CRISPR-Cas9 knockout MT4C5 cell lines, the study identifies ISG15

and APOBEC3G as key effectors mediating IFN-α14's enhanced suppression of HIV-1 replication.

Loss of either ISG15 or APOBEC3G abolished the differential antiviral effect between IFN-α14

and IFN-α2, demonstrating their essential roles in IFN-α14 driven viral restriction. These findings

highlight that individual IFN-α subtypes engage distinct downstream pathways and that subtype

diversity encodes functional specialization rather than redundancy. Overall, this work advances our

understanding of innate immune control of HIV-1 and provides a foundation for developing

targeted interferon-based therapies that exploit the unique mechanisms of potent subtypes like IFN-

α14.

INTRODUCTION

Type I interferon (IFN) is a crucial component of the innate immune system, playing an essential role in defending against a wide range of viruses(1). In humans, type I IFN includes 12 IFN-α subtypes along with IFN-β, IFN-κ, IFN-ε, IFN-o, IFN-τ, and IFN-δ(2). All type I IFN signal through the widely expressed IFN-α receptor (IFNAR)(3), triggering the activation of interferon regulated genes (IRG)(4, 5). Interferon-α plays a role in controlling many viral diseases including HIV-1. Produced primarily by plasmacytoid dendritic cells, IFN-α engages the IFNAR and activates JAK–STAT and MAPK pathways(6), inducing hundreds of IRG that interfere with viral replication at multiple stages, from entry to assembly(7, 8). It is now known that IFN-α subtypes induce IRG, including viral restriction factors, in a differential manner(9, 10, 11, 12). Many antiretroviral restriction factors have been identified that can inhibit HIV-1, including classical and well documented apolipoprotein B mRNA editing catalytic polypeptide-like 3G (APOBEC3G), SAM and HD domain-containing protein 1 (SAMHD1), tetherin, ISG15 and tripartite motif-containing protein 5 (TRIM5α) and those of more recent characterization: MX Dynamin Like GTPase 2 (MX2), serine incorporator (SERINC) 3/5, interferon-induced transmembrane proteins (IFITMs), schlafen 11, and membrane-associated RING-CH (MARCH) 2/8(13, 14, 15, 16, 17, 18, 19, 20, 21).

Restriction factors can impede HIV-1 replication in many ways. For example, APOBEC3G is incorporated into HIV-1 virions from a producer cell and exerts its antiviral activity in newly infected cells. The antiviral activity of APOBEC3G includes inhibition of reverse transcription, impairment of proviral integration and deamination of the viral RNA resulting in G to A hypermutation of the nascent viral DNA(22, 23, 24) that can result in defective provirus. The MX2 protein inhibits HIV-1 by interfering with the nuclear import of the viral pre-integration complex,

which is a crucial step after viral DNA synthesis but before its integration into the host cell's

genome(25, 26). Additionally, MX2 inhibits viral capsid uncoating, thereby blocking HIV-1

infection(27). ISG15, a ubiquitin-like protein, is the most highly expressed IRG(28). It inhibits

both the assembly and release of HIV-1 virions from cells(29, 30).

It has been well documented that IFN-α subtypes differentially control HIV-1 replication. IFN-

α14 is the more potent subtype both *in vitro* and *in vivo* when compared to clinically used IFN-

α2(12, 31). However, whether the more potent control of HIV-1 by IFN-α14 is due to differential

modulation of restriction factors is unknown. Lavender et al. and Harper et al.(31, 32) showed that

transcription of *MX2* was significantly increased by IFN-α14 but not IFN-α2. In contrast,

APOBEC3G was not transcriptionally upregulated but there was a significant increase in

APOBEC3G-mediated signature mutations in the proviral HIV-1 DNA (vDNA) from IFN-α14

compared to IFN-α2 treated mice. In 2021, Tauzin et al.(12) showed that IFN-α14 also induced

greater *ISG15* expression in MT4C5 cells. Since different IFN-α subtypes appear to modulate

different restriction factors and control HIV-1 in a differential manner, we wanted to investigate

whether the MX2, ISG15 and APOBEC3G restriction factors are important in IFN-α14's more

potent control of HIV-1. We used a CRISPR-Cas9 system to knock out each restriction factor in

MT4C5 cells to study their importance in the more potent control of HIV-1 by IFN-α14 compared

to IFN-α2.

RESULTS

IFN-α14 more potently reduced HIV-1 replication than IFN-α2 when used at a

biologically relevant activity. The relationship between protein concentrations and the functional

activity of cytokines such as IFN-α is not strictly linear, as functional activity is influenced not

only by expression levels but also by factors such as folding efficiency when using recombinant

proteins and ligand-receptor interactions. Different IFN-α subtypes have been shown to bind with

different affinities and kinetics to the interferon alpha receptor (IFNAR)(33, 34, 35, 36, 37). For

example, the IFN-α14 subtype binds more tightly to IFNAR compared to IFN-α2(38) and this

difference in affinity could potentially influence the intensity of the downstream signaling

cascade(39) leading to differential expression of IRGs. Also, doses of biologics like interferons

should be expressed in activity units rather than mass units to account for the presence of inactive

protein fractions in recombinant formulations(40). Since the IFN-α2 and -α14 subtypes in our

study were produced recombinantly and have different affinities for IFNAR, we first needed to

normalize their concentrations based on signaling activity in order to better understand their

potentially different antiviral effects(31, 41). To normalize the activity between IFN-α2 and IFN-

α14 we used a reporter cell line containing an interferon-stimulated response element (ISRE)

promoter/enhancer element driving a luciferase reporter gene**.** Using the reporter cell line, we

titrated each IFN-α subtype over a range of protein concentrations and normalized the activities of

each subtype based on their EC_{50} values to that of an IFN-α standard with a known activity prior

to use in experiments (Fig. 1A).

To evaluate the importance of antiretroviral factors in mediating more potent control of HIV-

1 by IFN-α14 compared to IFN-α2 we utilized the MT4C5 cell line. The MT4C5 cell line is a

derivative of MT4 cells that additionally expresses CCR5 and can be infected by CXCR4-tropic

(X4) or CCR5-tropic (R5) HIV-1 viruses. Importantly, following HIV-1 infection, MT4C5 cells

do not produce measurable amounts of endogenous IFN-α in the cell culture supernatant(42) that

may otherwise confound findings. We first established our system in MT4C5 cells to test the

contributions of each restriction factor to the more potent control of HIV-1 by IFN-α14. Wild-

type (WT) MT4C5 cells were infected with HIV-1$_{JR-CSF}$ at 0.1 MOI for 2 hours and then treated

with a biologically relevant activity (100U/ml)(43) or two supraphysiological doses (500U/ml or

1000U/ml)(44) of IFN-α2 or IFN-α14. Three days post infection, fresh medium with 100U/ml,

500U/ml or 1000U/ml of IFN-α2 or IFN-α14 was replenished. Six days post infection,

supernatant was collected and a p24 ELISA performed to quantify HIV-1 replication through the

production and release of HIV-1 virions into the supernatant. Both IFN-α subtypes significantly

reduced p24 in supernatants (Fig. 1B). However, supernatants from cells cultured in 100U/ml of

IFN-α2, contained significantly higher amounts of viral p24 compared to cells cultured in

100U/ml of IFN-α14 (Fig. 1B). In cultures containing 500U/ml and 1000U/ml of IFN-α, there

were no significant differences observed between subtypes, although a trend toward lower levels

of virus production could be detected in the cultures containing IFN-α14 (Fig. 1B). The results

indicated that greater suppression of HIV-1 replication in MT4C5 cells by IFN-α14 compared to

IFN-α2 occurred at the biologically relevant activity(43) of 100U/mL, but less so at

supraphysiological concentrations.

Potent suppression of HIV-1 replication by IFN-α14 occurs intracellularly. We first

investigated if lower levels of virus in supernatants from IFN-α14 treated cells might have resulted

from an accumulation of virions within the cell with inhibition of HIV-1 release into the

supernatant. We performed a p24 immunoblot using cell lysates from untreated HIV-1 infected

MT4C5 cells and cells treated for 6 days with 100U/ml of either IFN-α2 or -α14. The immunoblot

indicated that lysates from cells treated with IFN-α14 had no detectable cell-associated p24

whereas p24 was clearly visible in lysates from cells treated with IFN-α2 (Fig. 2A). Furthermore,

quantification of the p24 bands in relation to the GAPDH loading control on the western blot

confirmed there was some reduction in cellular p24 in the cells treated with 100U/ml of IFN-α2

compared to untreated cells as expected based on the p24 ELISA results in Fig. 1B, although not

nearly to the degree as in IFN-α14 treated cells (Fig. 2B). The results indicated that the more potent

suppression of HIV-1 replication by IFN-α14 than IFN-α2 at the biologically relevant

concentration of 100U/mL(43) did not primarily originate due to accumulation of cell-associated

virions that were not being released into the supernatant.

IFN-α14 treatment more potently reduced cellular HIV-1 viral DNA (vDNA) and viral

infectivity compared to IFN-α2. Having established that IFN-α14 potently suppressed levels of

HIV-1 p24 within the cell we next evaluated whether the reduction in p24 was due to reductions

in viral HIV-1 DNA (vDNA) within cells and/or the amount of infectious virus being produced

over the 6-day infection. First, we evaluated levels of vDNA in HIV-1 infected MT4C5 cells that

had been left untreated or treated with 100U/ml of either IFN-α2 or IFN-α14 for 6 days. To

quantify the levels of vDNA we isolated genomic DNA from the cells at day 6 and performed

qPCR targeting the HIV-1 POL region. We found that cells treated with 100U/ml of IFN-α14

contained relatively lower levels of total vDNA compared to untreated and IFN-α2 treated cells

(Fig. 3A). Cellular input levels were normalized to copies of *GAPDH*. Reductions in cellular p24

and vDNA over 6 days of infection could also relate to antiretroviral restriction factor-mediated

production and release of defective virions. Therefore, we assessed if IFN-α14 treatment of

MT4C5 cells resulted in the production of fewer infectious HIV-1 virions. Supernatants from cells

treated with 100U/ml of IFN-α2 or IFN-α14 were isolated and tested for levels of infectivity per

pg of p24 using TZM-bl reporter cells, which contain a Tat-sensitive LTR promotor driving the β-

galactosidase gene. Individually infected cells can thus be quantified through the presence of β-

galactosidase activity. The infectivity of virus-containing supernatants isolated from cells treated

with 100U/ml IFN-α14 was significantly lower than the infectivity of HIV-1 in supernatants after

IFN-α2 treatment (Fig. 3B). The data show that IFN-α14 more potently reduced both HIV-1 vDNA

and viral infectivity than IFN-α2 in MT4C5 cells.

CRISPR-Cas9-mediated knock out of MX2, APOBEC3G and ISG15 in MT4C5 cells. At this

point in our investigations the antiretroviral restriction factors MX2, APOBEC3G and ISG15

previously shown to be either more highly expressed or activated by IFN-α14, remained as

candidates factors mediating IFN-α14's more potent control of HIV-1 infection(12, 31, 32). To

investigate the role of each factor further, we used the CRISPR-Cas9 method to knock out each

restriction factor in MT4C5 cells. Briefly, two to three guide oligos were designed for each gene

(Table 1) and individually cloned into a lentiCRISPRv2 backbone. Lentiviral vectors carrying the

guide sequences were then made by transfecting lentiCRISPRv2, psPAX and pMD2.G plasmids

into HEK293FT cells. The MT4C5 cells were then transduced with the each of the resultant

lentiviruses for 48h before plating them into selection medium. Successful MX2 and ISG15 gene

knock out (KO) was determined using immunoblots and APOBEC3G KO confirmed by PCR due

to a specific antibody being unavailable. A concentration of 1000U/ml of IFN-α2 was needed to

induce detectable levels of MX2 in WT MT4C5 cells whereas MX2 could be detected in cells

treated with either 100 or 1000U/ml of IFN-α14, albeit less so after stimulation with 100U/ml of

the cytokine. However, MX2 expression was not detected after stimulation with either IFN-α in

the MX2 KO cells (Fig. 4A). To detect ISG15 protein in MT4C5 cells 1000U/ml of both IFN-α

subtypes was used. However, no ISG15 was detected in KO cells after IFN-α stimulation (Fig.

4B). The APOBEC3 family is highly homologous at the protein level and thus there are no

commercially available antibodies that could specifically detect the APOBEC3G KO in MT4C5

cells. Thus, we used APOBEC3G-specific sequences in genomic DNA to detect the APOBEC3G

KO. Amplification of genomic DNA using APOBEC3G sequence-specific primers showed

amplification of a shorter band in APOBEC3G KO cells compared to genomic DNA from WT

MT4C5 cells. In summary, we were able to successfully knock out MX2, ISG15 and APOBEC3G

in MT4C5 cells in order to further evaluate the contribution of each antiretroviral restriction factor

to the more potent control of HIV-1 infection by IFN-α14 compared to IFN-α2.

Deletion of APOBEC3G and ISG15 but not MX2 eliminated IFN-α14's more potent

control of HIV-1 compared to IFN-α2. We next used our MX2, APOBEC3G and ISG15 KO cells

to evaluate their importance in IFN-α14's more potent control of HIV-1 compared to IFN-α2. To

evaluate whether IFN-α14's potency was dependent on APOBEC3G, ISG15 or MX2 each of the

KO cells as well as WT MT4C5 cells were treated with 100U/mL of either IFN-α2 or IFN-α14 as

performed earlier. As expected, IFN-α14 reduced levels of HIV-1 p24 in the cell supernatants

compared to IFN-α2 in the WT MT4C5 cells. Unexpectedly, IFN-α14 still reduced p24 levels in

supernatants significantly better than IFN-α2 in the MX2 KO cells (Fig. 5). However, supernatants

collected from APOBEC3G KO and ISG15 KO cells no longer contained different amounts of p24

regardless of which IFN-α treatment they received (Fig. 5). Therefore, the data suggested that the

restriction factors APOBEC3G and ISG15 have a role in the more potent control of HIV-1 by IFN-

α14 compared to IFN-α2, whereas MX2 does not.

IFN-α14 more potently augmented APOBEC3G cytidine deaminase activity and reductions

in viral infectivity than IFN-α2 but did not affect vDNA levels. Since HIV-1 vDNA levels and

viral infectivity were significantly reduced by IFN-α14 but not IFN-α2 treatment in MT4C5 WT

cells, we tested whether these effects were being mediated through the restriction factor

APOBEC3G. First, we assessed levels of vDNA in HIV-1 infected MT4C5 WT or APOBEC3G

KO cells treated for 6 days with either 100U/ml IFN-α14 or IFN-α2 by amplifying the HIV-1 POL

region from isolated genomic DNA as already described. Fold change was calculated from levels

of vDNA in untreated MT4C5 WT cells. As we had observed previously, IFN-α14 treatment

resulted in significantly lower amounts of vDNA in cells compared to IFN-α2 in MT4C5 WT cells.

However, knock out of APOBEC3G did not alleviate the effect (Fig. 6A). In contrast, when we

collected day 6 supernatants from MT4C5 WT and APOBEC3G KO cells and evaluated the

infectivity of HIV-1 per pg of p24 we observed that APOBEC3G KO restored the infectivity of

HIV-1 virions produced from the IFN-α14 treated cells to comparable levels as IFN-α2 treated

cells (Fig. 6B). Since the cytidine deaminase activity of APOBEC3G can contribute to reductions

in viral infectivity through deleterious mutagenesis of the vDNA we also assessed the frequency

of GG → AG signature mutations. We evaluated day 6 supernatants from MT4C5 WT and

APOBEC3G KO cells to induce signature mutations in target cells. Interestingly, the frequency of

GG → AG signature mutations in the target cells incubated with supernatants from MT4C5 WT

cells treated with IFN-α2 was not altered in the absence of APOBEC3G. Conversely, when treated

with IFN-α14, knock out of APOBEC3G resulted in a significant reduction in the frequency of

signature mutations in target cells (Fig. 6C). Overall, IFN-α14 more potently controlled HIV-1

than IFN-α2 through augmentation of APOBEC3G cytidine deaminase activity leading to the

production of more non-infectious HIV-1 virions.

DISCUSSION

Greater expression of restriction factors such as *ISG15*, and *MX2* and increased activity of

APOBEC3G in IFN-α14 treated humanized mice(12, 31) suggested that IFN-α14's potency may

be, at least in part, due to the contribution of specific restriction factors. The data presented in the

current study indicated that the potent HIV-1 suppression mediated by IFN-α14 is driven, at least

in part, cell intrinsically by ISG15 and APOBEC3G. We observed that the MT4C5 cells harboring

genomic deletion of *ISG15* and *APOBEC3G* showed no difference in viral p24 production when

treated with 100U/mL of IFN-α2 or IFN-α14 suggesting that ISG15 and APOBEC3G are

responsible for the more potent control of HIV-1 by IFN-α14. It further demonstrated that IFN-

α14 treatment not only reduced viral load but also resulted in the production of fewer infectious

particles. However, when we increase the dosage of both IFN-αs, the difference ceased to exist. A

study done by Schlaepfer and Fahrny et al. showed that at higher doses of IFN-α subtypes, reporter

gene induction in response to different IFN-α subtypes waned(45). One of the reasons could be

that at higher doses of IFN-α, the IFNAR1/2 receptors become saturated, meaning nearly all

receptor complexes are occupied regardless of subtype affinity. When that happens, downstream

signaling (e.g., STAT phosphorylation, ISG induction) and antiviral effects may reach a maximal

plateau, minimizing observable subtype-specific differences(46). Therefore, we used a

biologically relevant dose of 100U/ml(43) where differences in IFN-α responses between subtypes

become detectable.

IFN-α14 suppressed virus replication in HIV-1 infected MT4C5 cells. Although p24 was

detectable in the supernatant, a complete loss of p24 was observed in the cell lysates of the IFN-

α14–treated cells by day 6. This indicated that HIV-1 replication initially occurred following

infection, leading to viral production in the supernatant, but IFN-α14 treatment subsequently

slowed within cells over the 6 days. This was confirmed by amplifying the POL region of HIV-1

from genomic DNA isolated from cells. However, given that ELISA and PCR are more sensitive

than western blot, the absence of p24 detection by western blot does not rule out the presence of

cellular virions.

Restriction factors like MX2, APOBEC3G and ISG15 hinder HIV-1 replication at different

stages of HIV-1's lifecycle. MX2 and APOBEC3G act during earlier stages whereas ISG15 acts

during the later stages of HIV-1 replication. Since we saw a decrease in viral infectivity in the IFN-

α14 treated group compared to IFN-α2 we anticipated that the decrease in infectivity could be a

result of increased APOBEC3G activity, therefore we investigated APOBEC3G more thoroughly.

ISG15 on the other hand, is known to hinder HIV-1 by preventing ubiquitination of Gag and

Tsg101 which is important for viral budding and release(47). ISG15 also modifies a broad range

of cellular proteins, many of which are key components of the antiviral innate immune signaling

pathway(48). In future, experiments could be performed to determine if ISG15 is restricting HIV-

1 more potently by preventing ubiquitination of Gag and Tsg101 or by modifying other cellular

proteins after IFN-α14-treatment. A previous study(31), along with our own findings, concluded

that APOBEC3G activity is enhanced during IFN-α14 treatment. However, the mechanism

underlying the increased frequency of GG $\rightarrow$ AG hypermutations in the absence of elevated

APOBEC3G expression levels remains unclear. Others have shown that treatment with IFN-α

increases *APOBEC3G* mRNA levels 36 folds in a different cell line(49). However, in MT4C5 cell

line there was no elevation in *APOBEC3G* mRNA levels(12). Deletion of APOBEC3G did result

in comparable levels of HIV-1 cell supernatants between IFN-α treatment groups suggesting that

APOBEC3G has a role in IFN-α14's increased anti-HIV-1 potency. Apart from APOBEC3G,

281 deletion of ISG15 eliminated subtype specific differences in viral load in MT4C5 cells. However,

282 deletion of MX2 did not eliminate the subtype specific differences in HIV-1 control in cells.

283  A study showed that inactivation of multiple ISGs induced by IFN-β revealed a polygenic and

284 synergistic mechanism underlying HIV-1 control**(50).** Notably, deletion of MX2 did not reduce

285 IFN-α14's capacity to suppress HIV-1 replication, suggesting that other IFN-α14 induced

286 restriction factors can maintain antiviral activity. In polygenic systems, the effect of one protein

287 can be influenced by the abundance or activity of others, depending on their mode of interaction.

288 ISG15 functions late in the viral life cycle, blocking assembly and release(29, 30), whereas

289 APOBEC3G and MX2 act at earlier stages. When early acting factors such as APOBEC3G or

290 MX2 substantially limit the production of competent virions, ISG15 has fewer targets, making its

291 impact appear smaller; however, when early restrictions are weak, ISG15's role in blocking release

292 becomes more critical. Therefore, deletion of ISG15 may lead to a greater increase in viral

293 replication compared to deletion of MX2 under IFN-α14 treatment, as the antiviral effect of MX2

294 can be partially compensated by ISG15, whereas loss of ISG15 is less likely to be offset by other

295 restriction factors. It is plausible that the expression level of a compensatory restriction factor

296 influences HIV-1 replication. However, in our experiments, we did not measure the expression of

297 other restriction factors when a single factor was knocked out during HIV-1 infection. Future

298 studies involving the simultaneous knock out of ISG15, APOBEC3G, and MX2 could help reveal

299 synergistic interactions among restriction factors regulated by IFN-α14. Since the MT4C5 cell line

300 does not produce endogenous IFN-α, it served as a useful model to examine the exogenous effects

301 of IFN-α subtypes. However, not all IRG responses could be evaluated, as MT4C5 cells may

302 respond differently from primary CD4+ T cells. Therefore, future validation using primary CD4+

303 T cells would be useful to assess the physiological relevance of IFN-α subtype responses.

Here we demonstrated that IFN-α14, in contrast to IFN-α2, exerted a stronger effect on HIV-

1 replication and restriction factors like ISG15 and APOBEC3G play significant roles in the

differential effect between IFN-α2 and IFN-α14. However, the contribution of other restriction

factors like MX2 and BST2 cannot be ignored in the context of HIV-1 replication. Although in our

experimental setup these restriction factors (MX2 and BST2) may seem redundant, but their

contribution to control HIV-1 cannot be disregarded. In future, short-term experiments for 2 or 3

days with a fully established infection and IFN-α treatment could be done to evaluate the effect of

MX2 and BST2 on IFN-α14 potency.

MATERIALS AND METHODS

Determination of IFN-α units

A stable reporter cell line using human retinal pigment epithelial cells (ATCC CRL2302)

transfected with a plasmid containing interferon-stimulated response element promoter/enhancer

elements driving a luciferase reporter gene. Cells were grown for 24 h before adding serial

dilutions of recombinant IFN-α subtypes and a predetermined IFN-α lab standard for 6 h. The cells

were lysed with Pierce Firefly Luc One-Step Assay Buffer (Thermofisher, Waltham, MA), and

luciferase activity was measured 10 min later. The relative light units were plotted against the IFN-

α dilutions using non-linear fit with three parameters and EC50 was determined. Units of IFN-α

were calculated using EC50 values and known units/ μL of the standard.

HIV-1 infection and IFN-α treatment in MT4C5 WT and KO cells

MT4C5 cells were kindly donated by Dr. Oliver Schwartz (Institut Pasteur, Paris, France). MT4C5

is a $CD4^+$ T cell line expressing CCR5 and has little to no endogenous production of IFN-α(42).

MT4C5 WT cells were maintained in RPMI-1640 (Hyclone, Logan, UT) supplemented with 10%

FCS (Hyclone) and 1X PenStrep (Hyclone) (R-10) while KO made from MT4C5 cells were

maintained with R-10 supplemented with 5μg/mL puromycin (Invivogen, San Diego, CA). Cells

were either left untreated or treated with IFN-α2 or IFN-α14 (100U/mL, 500U/mL and

1000U/mL) for 24h and then infected with 0.1 MOI of HIV-1$_{JR-CSF}$ (credit proper people off AIDS

Reagent) for 2 hours. Cells were then washed 2 times with complete media and plated in a 24 well

plate. Post-infection all the cells were maintained in either IFN-α2 or IFN-α14 at the initial

treatment dose.

Guide RNA selection and cloning into lentiCRISPRv2

Each guide sequence was designed using Synthego's CRISPR Design Tool. Guide sequences for

MX2 (ENST00000330714), APOBEC3G (ENST00000407997) and ISG15 (ENST00000379389)

were chosen from the list provided by Synthego's CRISPR Design Tool

(https://www.synthego.com/products/bioinformatics/crispr-design-tool). Location for cut site for

MX2 (41.376.931, 41.377.150 and 41.377.009), APOBEC3G (39.080.934, 39.080.990 and

39.081.088) and ISG15 (1.014.247 and 1.014.006) were chosen such that the fragment that would

be deleted would not be a multiple of three. DNA oligos (**Table 1**) for the guide sequences were

ordered from Integrated DNA Technologies.

The protocol for cloning was adapted from Zhang Lab(51). The complementary oligos were

phosphorylated using T4 PNK enzyme (New England Biolabs, Massachusetts, USA) and T4

Ligase buffer (Invitrogen, Carlsbad, California, USA) at 37°C for 1h. After phosphorylation, the

complementary oligos were heated to 95°C in water bath and cooled down gradually to room

temperature (22°C-25°C). LentiCRISPR v2 was a gift from Feng Zhang (Addgene plasmid #

52961 ; http://n2t.net/addgene:52961 ; RRID:Addgene_52961). LentiCRISPRv2 was digested

using *BsmBI-v2* and the 12kb backbone was gel extracted using QIAquick Gel Extraction Kit Gel

Purification (QIAGEN, Hilden, Germany). Oligos and the LentiCRISPRv2 backbone were ligated

overnight at 16°C. Each of the ligated plasmids were transformed into *Stbl2* cells (Invitrogen) and

were grown in freshly made LB Agar plates with 100μg/mL of ampicillin. Single colonies were

grown in 5mL of LB media supplemented with 100μg/mL of ampicillin. Plasmids were isolated

from transformed *Stbl2* cells using QIAprep Spin Miniprep Kit (QIAGEN) and PCR used to

confirm the presence of the insert.

Lentivirus production and transduction

HEK293 FT cells were maintained in DMEM (Gibco, Waltham, MA) supplemented with 10%

FCS, 1X PenStrep, 1X Sodium Pyruvate and 1X NEAA (DMEM complete media). Cells were

transfected with cloned LentiCRISPRv2 plasmid carrying oligos, pMD2.G and psPAX2 at a ratio

of (1:2:1) using MEM (Gibco) and Lipofectamine 2000 (Invitrogen). After 72h of transfection,

supernatant containing lentiviruses were collected and spun down at 1000 x g for 5mins to separate

cell debris. Lentiviruses for same restriction factor were then incubated with MT4C5 cells for 24h.

After 72h of transduction, freshly prepared selection medium (R-10 + 5 μg/mL puromycin) was

added to the culture. After two weeks of selection, cells were treated with IFN-αs to confirm the

KOs.

Western blots

Cells were counted and 2X SDS buffer was added to cells. The mixture was then heated at 95°C

for 10 mins and then spun at 13,000xg for 1 min. Samples were loaded onto a 12% SDS-PAGE

gel for 1.2 hrs. After protein transfer was done onto the nitrocellulose membrane, blocking was

done using 1X clear milk (Thermofisher). Membrane was then incubated with primary antibodies

(Table 2) overnight at 4°C and washed 5 times with 0.1% tween-20 containing 1XPBS for 5mins

each. Secondary antibodies **(Table 2)** were added and incubated at room temperature for one hour

and then washed with 5 times with 0.1% tween-20 containing 1XPBS for 5mins each.

Genomic DNA isolation and PCR amplification

Genomic DNA from different experimental groups was extracted using the QIAamp DNA minikit

(QIAGEN) kit, following the manufacturer's instructions. The APOBEC3G sequence flanking the

region targeted by guide RNAs was amplified using forward primer 5'-

GGATGAAGCCTCACTTCAGAAA-3' and reverse primer 5'-

CACAGGTAAGTCTCATGCCGT-3'. Determination of total HIV-1 DNA was performed by

quantitative real-time PCR using SsoAdvanced Universal SYBR Green Supermix (BIORAD,

Hercules, CA) targeting the POL region of HIV-1 with forward primer 5'-

ACTCCAGTATTTGCCATAAAGAAAAAA-3' and reverse primer 5'-

CTGTCTTTTTCTGGCAGCACTAT-3' and GAPDH with forward primer 5'-

GCTCTTAAAAAGTGCAGGGTCT-3' and reverse primer 5'-

GATGGTACATGACAAGGTGCG-3'.

p24 enzyme linked immunosorbent assay

Cell supernatants were collected at day 6 for p24 ELISA (Advanced Bioscience Laboratories,

Maryland, USA) following the manufacturer's instructions using a laboratory optimized

modification that halved the sample and reagent volumes used at each step. Plates were read in a

96 well plate reader at 450nm. Data acquisition was done using Softmax Pro (San Jose, CA).

Infectivity Assay

TZM-bl reporter cells (NIH AIDS Research and Reference Reagent Program, from John Kappes,

Xiaoyun Wu, and Tranzyme Inc.) were used to determine the number of infectious particles in the

supernatant. Briefly, cells were plated 24h prior to the assay in a 96-well plate. Supernatants were

10X serially diluted and incubated with cells for 2h. Cells were then washed and cultured with

fresh media for 48h. Cells were then fixed with 2% gluteraldehyde and X-gal solution was added

and plate was incubated at 37°C at 5% CO_2 for 24h. Spots were then counted at 4X magnification

and the infectivity was normalised to the p24 of the supernatants.

Detection of APOBEC3G signature mutations

Supernatants from MT4C5 WT or A3G KO treated with different IFN-α subtypes for 6 days were

incubated with target cells (A3G KO) for 48h. The POL region was amplified, gel purified using

QIAquick Gel Extraction Kit – Gel Purification (QIAGEN) and sent for sequencing

(Plasmidsaurus, South San Francisco, CA) to determine the ratio of GG→ AG mutations vs total

mutations.

Statistics

Data were plotted with mean and standard deviation. Normality of each dataset was evaluated by

Student's unpaired t-test or one-way ANOVA with Tukey's post-tests were performed accordingly

and are specified within figure legends for each data set. $P < 0.05$ was considered statistically

significant. All statistics were performed using Prism 10 (GraphPad, San Diego, CA).

419   **ACKNOWLEDGEMENTS**

REFERENCES

1.   Kalliolias GD, Ivashkiv LB. Overview of the biology of type I interferons. Arthritis Res Ther. 2010;12 Suppl 1(Suppl 1):S1.

2.   Pestka S, Krause CD, Walter MR. Interferons, interferon-like cytokines, and their receptors. Immunological Reviews. 2004;202(1):8-32.

3.   Uzé G, Schreiber G, Piehler J, Pellegrini S. The Receptor of the Type I Interferon Family. Interferon: The 50th Anniversary. Current Topics in Microbiology and Immunology2007. p. 71-95.

4.   Der SD, Zhou A, Williams BR, Silverman RH. Identification of genes differentially regulated by interferon alpha, beta, or gamma using oligonucleotide arrays. Proc Natl Acad Sci U S A. 1998;95(26):15623-8.

5.   O'Garra A, Trinchieri G. Are dendritic cells afraid of commitment? Nature Immunology. 2004;5(12):1206-8.

6.   Platanias LC. Mechanisms of type-I- and type-II-interferon-mediated signalling. Nature Reviews Immunology. 2005;5(5):375-86.

7.   Lin Y, Zhu Y, Jing L, Lei X, Xie Z. Regulation of viral replication by host restriction factors. Frontiers in Immunology. 2025;16.

8.   Yang E, Li MMH. All About the RNA: Interferon-Stimulated Genes That Interfere With Viral RNA Processes. Frontiers in Immunology. 2020;11.

9.   Guo K, Shen G, Kibbie J, Gonzalez T, Dillon SM, Smith HA, et al. Qualitative

Differences Between the IFNalpha subtypes and IFNbeta Influence Chronic Mucosal HIV-1

Pathogenesis. PLoS Pathog. 2020;16(10):e1008986.

10.   Schuhenn J, Meister TL, Todt D, Bracht T, Schork K, Billaud JN, et al. Differential

interferon-alpha subtype induced immune signatures are associated with suppression of SARS-

CoV-2 infection. Proc Natl Acad Sci U S A. 2022;119(8).

11.   Lavender KJ, Gibbert K, Peterson KE, Van Dis E, Francois S, Woods T, et al. Interferon

Alpha Subtype-Specific Suppression of HIV-1 Infection In Vivo. J Virol. 2016;90(13):6001-13.

12.   Tauzin A, Espinosa Ortiz A, Blake O, Soundaramourty C, Joly-Beauparlant C, Nicolas A,

et al. Differential Inhibition of HIV Replication by the 12 Interferon Alpha Subtypes. Journal of

Virology. 2021;95(15).

13.   Zhang H, Yang B, Pomerantz RJ, Zhang C, Arunachalam SC, Gao L. The cytidine

deaminase CEM15 induces hypermutation in newly synthesized HIV-1 DNA. Nature.

2003;424(6944):94-8.

14.   Van Damme N, Goff D, Katsura C, Jorgenson RL, Mitchell R, Johnson MC, et al. The

interferon-induced protein BST-2 restricts HIV-1 release and is downregulated from the cell

surface by the viral Vpu protein. Cell Host Microbe. 2008;3(4):245-52.

15.   Stremlau M, Perron M, Lee M, Li Y, Song B, Javanbakht H, et al. Specific recognition

and accelerated uncoating of retroviral capsids by the TRIM5alpha restriction factor. Proc Natl

Acad Sci U S A. 2006;103(14):5514-9.

16.   Stremlau M, Owens CM, Perron MJ, Kiessling M, Autissier P, Sodroski J. The

cytoplasmic body component TRIM5alpha restricts HIV-1 infection in Old World monkeys.

Nature. 2004;427(6977):848-53.

17.   Neil SJ, Zang T, Bieniasz PD. Tetherin inhibits retrovirus release and is antagonized by

HIV-1 Vpu. Nature. 2008;451(7177):425-30.

18.   Mangeat B, Turelli P, Caron G, Friedli M, Perrin L, Trono D. Broad antiretroviral defence

by human APOBEC3G through lethal editing of nascent reverse transcripts. Nature.

2003;424(6944):99-103.

19.   Lahouassa H, Daddacha W, Hofmann H, Ayinde D, Logue EC, Dragin L, et al. SAMHD1

restricts the replication of human immunodeficiency virus type 1 by depleting the intracellular

pool of deoxynucleoside triphosphates. Nat Immunol. 2012;13(3):223-8.

20.   Laguette N, Sobhian B, Casartelli N, Ringeard M, Chable-Bessia C, Segeral E, et al.

SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1 restriction factor counteracted by

Vpx. Nature. 2011;474(7353):654-7.

21.   Hrecka K, Hao C, Gierszewska M, Swanson SK, Kesik-Brodacka M, Srivastava S, et al.

Vpx relieves inhibition of HIV-1 infection of macrophages mediated by the SAMHD1 protein.

Nature. 2011;474(7353):658-61.

22.   Harris RS, Hultquist JF, Evans DT. The restriction factors of human immunodeficiency

virus. J Biol Chem. 2012;287(49):40875-83.

23.   Luo K, Wang T, Liu B, Tian C, Xiao Z, Kappes J, et al. Cytidine deaminases APOBEC3G

and APOBEC3F interact with human immunodeficiency virus type 1 integrase and inhibit

proviral DNA formation. J Virol. 2007;81(13):7238-48.

24.   Iwatani Y, Chan DS, Wang F, Stewart-Maynard K, Sugiura W, Gronenborn AM, et al.

Deaminase-independent inhibition of HIV-1 reverse transcription by APOBEC3G. Nucleic Acids

Res. 2007;35(21):7096-108.

25.   Goujon C, Moncorgé O, Bauby H, Doyle T, Ward CC, Schaller T, et al. Human MX2 is

an interferon-induced post-entry inhibitor of HIV-1 infection. Nature. 2013;502(7472):559-62.

26.   Kane M, Yadav SS, Bitzegeio J, Kutluay SB, Zang T, Wilson SJ, et al. MX2 is an

interferon-induced inhibitor of HIV-1 infection. Nature. 2013;502(7472):563-6.

27.   Fricke T, White TE, Schulte B, de Souza Aranha Vieira DA, Dharan A, Campbell EM, et

al. MxB binds to the HIV-1 core and prevents the uncoating process of HIV-1. Retrovirology.

2014;11(1).

28.   Scagnolari C, Monteleone K, Selvaggi C, Pierangeli A, D'Ettorre G, Mezzaroma I, et al.

ISG15 expression correlates with HIV-1 viral load and with factors regulating T cell response.

Immunobiology. 2016;221(2):282-90.

29.   Pincetic A, Kuang Z, Seo EJ, Leis J. The Interferon-Induced Gene ISG15 Blocks

Retrovirus Release from Cells Late in the Budding Process. Journal of Virology.

2010;84(9):4725-36.

30.   Okumura A, Lu G, Pitha-Rowe I, Pitha PM. Innate antiviral response targets HIV-1

release by the induction of ubiquitin-like protein ISG15. Proceedings of the National Academy of

Sciences. 2006;103(5):1440-5.

31.   Lavender KJ, Gibbert K, Peterson KE, Van Dis E, Francois S, Woods T, et al. Interferon

Alpha Subtype-Specific Suppression of HIV-1 InfectionIn Vivo. Journal of Virology.

2016;90(13):6001-13.

32.   Harper MS, Guo K, Gibbert K, Lee EJ, Dillon SM, Barrett BS, et al. Interferon-alpha

Subtypes in an Ex Vivo Model of Acute HIV-1 Infection: Expression, Potency and Effector

Mechanisms. PLoS Pathog. 2015;11(11):e1005254.

33.   Brierley MM, Fish EN. Review: IFN-α/βReceptor Interactions to Biologic Outcomes:

Understanding the Circuitry. Journal of Interferon & Cytokine Research. 2002;22(8):835-45.

34.   Severa M, Remoli ME, Giacomini E, Ragimbeau J, Lande R, Uzé G, et al. Differential

responsiveness to IFN-α and IFN-β of human mature DC through modulation of IFNAR

expression. Journal of Leukocyte Biology. 2006;79(6):1286-94.

35.   Langer JA. Interferon at 50: New Molecules, New Potential, New (and Old) Questions.

Science's STKE. 2007;2007(405).

36.   Jaks E, Gavutis M, Uzé G, Martal J, Piehler J. Differential Receptor Subunit Affinities of

Type I Interferons Govern Differential Signal Activation. Journal of Molecular Biology.

2007;366(2):525-39.

37.   Jaitin DA, Roisman LC, Jaks E, Gavutis M, Piehler J, Van der Heyden J, et al. Inquiring

into the Differential Action of Interferons (IFNs): an IFN-α2 Mutant with Enhanced Affinity to

IFNAR1 Is Functionally Similar to IFN-β. Molecular and Cellular Biology. 2023;26(5):1888-97.

38.   Lavoie TB, Kalie E, Crisafulli-Cabatu S, Abramovich R, DiGioia G, Moolchan K, et al.

Binding and activity of all human alpha interferon subtypes. Cytokine. 2011;56(2):282-9.

39.   Marles JA, Dahesh S, Haynes J, Andrews BJ, Davidson AR. Protein-Protein Interaction

Affinity Plays a Crucial Role in Controlling the Sho1p-Mediated Signal Transduction Pathway in

Yeast. Molecular Cell. 2004;14(6):813-23.

40.   Robinson CJ. Bioassays for growth factors. Dev Biol Stand. 1999;97:21-7.

41.   Rout SS, Di Y, Dittmer U, Sutter K, Lavender KJ. Distinct effects of treatment with two

different interferon-alpha subtypes on HIV-1-associated T-cell activation and dysfunction in

humanized mice. Aids. 2022;36(3):325-36.

42.   Lepelley A, Louis S, Sourisseau M, Law HK, Pothlichet J, Schilte C, et al. Innate sensing

of HIV-infected cells. PLoS Pathog. 2011;7(2):e1001284.

43.   Hardy GA, Sieg S, Rodriguez B, Anthony D, Asaad R, Jiang W, et al. Interferon-alpha is

the primary plasma type-I IFN in HIV-1 infection and correlates with immune activation and

disease markers. PLoS One. 2013;8(2):e56527.

44.   Nicot F, Legrand-Abravanel F, Lafont T, Dubois M, Saune K, Pasquier C, et al. Serum

concentrations of ribavirin and pegylated interferon and viral responses in patients infected with

HIV and HCV. J Med Virol. 2008;80(9):1523-9.

45.   Schlaepfer E, Fahrny A, Gruenbach M, Kuster SP, Simon V, Schreiber G, et al. Dose-

Dependent Differences in HIV Inhibition by Different Interferon Alpha Subtypes While Having

Overall Similar Biologic Effects. mSphere. 2019;4(1).

46.   Moll HP, Maier T, Zommer A, Lavoie T, Brostjan C. The differential activity of

interferon-α subtypes is consistent among distinct target genes and cell types. Cytokine.

2011;53(1):52-9.

47.   Okumura A, Lu G, Pitha-Rowe I, Pitha PM. Innate antiviral response targets HIV-1

release by the induction of ubiquitin-like protein ISG15. Proc Natl Acad Sci U S A.

2006;103(5):1440-5.

48.   Zhao C, Denison C, Huibregtse JM, Gygi S, Krug RM. Human ISG15 conjugation targets

both IFN-induced and constitutively expressed proteins functioning in diverse cellular pathways.

Proceedings of the National Academy of Sciences. 2005;102(29):10200-5.

49.   Chen H, Wang LW, Huang YQ, Gong ZJ. Interferon-alpha induces high expression of

APOBEC3G and STAT-1 in vitro and in vivo. Int J Mol Sci. 2010;11(9):3501-12.

50.   Itell HL, Humes D, Overbaugh J. Several cell-intrinsic effectors drive type I interferon-

mediated restriction of HIV-1 in primary CD4+ T cells. Cell Reports. 2023;42(6).

51.   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries for

CRISPR screening. Nature Methods. 2014;11(8):783-4.

FIGURE LEGENDS

Figure 1. IFN-α14 more potently suppressed HIV-1 replication than IFN-α2 at similar activity levels. (A) The activities of IFN-α2 and IFN- α14 were determined by titration of each subtype on an IFN-α responsive reporter cell line in comparison to an IFN-α stock of known activity. (B) HIV-1 p24 ELISA of supernatants collected from MT4C5 cells treated with 0, 100, 500 or 1000U/ml of IFN-α2 or IFN-α14 for 6 days starting 2 hours post infection. One-way ANOVA with Tukey's multiple comparison test. **** $p<0.0001$

Figure 2. IFN-α14 more potently reduced cellular p24 levels in MT4C5 cells than IFN-α2. (A) HIV-1 p24 immunoblot of MT4C5 cell lysates previously treated for 6 days with IFN-α2 or IFN-α14. GAPDH loading control. (B) Quantification of the p24 immunoblot normalizing the intensity of the p24 bands to GAPDH.

Figure 3. IFN-α14 more potently reduced HIV-1 viral DNA (vDNA) and HIV-1 infectivity in MT4C5 cells than IFN-α2. (A) Fold change in HIV-1 vDNA levels in MT4C5 cells differentially treated with IFN-α upon amplification of the POL region from isolated genomic DNA. Cellular input was normalized to *GAPDH*. One-way ANOVA with Tukey's multiple comparison test ** $p<0.01$; **** $p<0.0001$. (B) Tissue culture infectious units (TCIU) per pg of input HIV-1 p24 was determined by TZM-bl infectivity assay using supernatants from MT4C5 cells treated with 100U/mL of IFN-α2 or IFN-α14. Student's unpaired t-test * $p<0.05$.

Figure 4. The MX2, APOBEC3G and ISG15 antiretroviral restriction factors were successfully knocked out of MT4C5 cells using CRISPR-Cas9. MT4C5 cell lysates of IFN-α

treated wild-type (WT), MX2 knockout (KO) and ISG15 KO cells were run on a 14% SDS PAGE

gel. The proteins were then transferred to a nitrocellulose membrane and immunoblotted for (A)

MX2 and (B) ISG15. GAPDH was used as a loading control. (C) PCR amplification product of a

461bp region of APOPEC3G (A3G) expected to be disrupted in A3G KO cells compared to WT

using APOBEC3G sequence specific primers.

Figure 5. Knock out of APOBEC3G (A3G KO) and ISG15 in MT4C5 cells eliminated

differential control of HIV-1 by IFN-α subtypes. HIV-1 p24 ELISA of supernatants collected

from HIV-1 infected MT4C5 (WT), A3G KO, ISG15 KO and MX2 KO cells after 6 days of

treatment with either 100U/ml of IFN-α2 or IFN-α14. Student's unpaired t-test * $p < 0.05$.

Figure 6. Deletion of APOBEC3G did not alter vDNA in cells but restored infectious particles

and frequency of GG→ AG signature mutations. (A) Fold change in HIV-1 vDNA levels in

MT4C5 WT and A3G KO cells differentially treated with IFN-α upon amplification of the POL

region from isolated genomic DNA. Cellular input was normalized to *GAPDH*. (B) TCIU per pg

of input HIV-1 p24 was determined by TZM-bl infectivity assay using supernatants from MT4C5

WT and A3G KO cells treated with 100U/mL of IFN-α2 or IFN-α14. (C) Amplified POL region

from vDNA of target cells were amplified and sequenced and GG→AG mutations were plotted

against total mutations. Student's unpaired t-test * $p < 0.05$; ** $p < 0.01$.

TABLES

Table 1. Guide DNA oligo sequences

Gene	Location	Oligo Sequence (5'-3')
MX2	41.376.931	**CACCG**CTTCTCCTCCGGTAGGGCCA
		AAACTGGCCCTACCGGAGGAGAAGC
	41.377.150	**CACCG**ACCGGGCTTACCATTGCCCT
		AAACAGGGCAATGGTAAGCCCGGTC
	41.377.009	**CACCG**GGCACTGTGCCTAATGGCGG
		AAACCCGCCATTCGGCACAGTGCCC
APOBEC3G	39.080.934	**CACCG**CTTAAGTTCGGAATACACCT
		AAACAGGTGTATTCCGAACTTAAGC
	39.080.990	**CACCG**TCCACTGGTTCAGCAAGTGG
		AAACCCACTTGCTGAACCAGTGGAC
	39.081.088	**CACCG**TAACCTTCGGGTCCTCGGCC
		AAACGGCCGAGGACCCGAAGGTTAC
ISG15	1.014.247	**CACCG**ATCCTGGTGAGGAATAACAA
		AAACTTGTTATTCCTCACCAGGATC
	1.014.006	**CACCG**GGACCTGACGGTGAAGATGC
		AAACGCATCTTCACCGTCAGGTCCC

605 **Table 2. Antibody dilutions**

Sl. no.	Antibody	Host	Dilution	Primary/Secondary
1.	Hu-MX2	Mouse	1:100	Primary
2.	Hu-ISG15	Mouse	1:1000	Primary
3.	Hu-GAPDH	Rabbit	1:10000	Primary
4.	m-IgG	Mouse	1:10000	Secondary
5.	m-IgG	Rabbit	1:10000	Secondary

FIGURES

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6