Title: Decoding pathogen ecological memory from chemical-stress phenotypes

Running title: Decoding fungal ecological memory

Insuck Baek[1], Seunghyun Lim[2], Amelia Lovelace[2], Sookyung Oh[1], Masoud Kazem-Rostami[3], Helen Ngo[3], Moon S. Kim[1], Lyndel W. Meinhardt[2], Lalit Kandpal[1], Minhyeok Cha[1], Chansong Hwang[1], Richard D. Ashby[3], Ezekiel Ahn[2,*]

[1] Environmental Microbial and Food Safety Laboratory, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, 20705, USA

[2] Sustainable Perennial Crops Laboratory, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, 20705, USA

[3] Sustainable Biofuels and Co-Products Research Unit, Agricultural Research Service, United States Department of Agriculture, Eastern Regional Research Center, Wyndmoor, PA, 19038, USA

* Author to whom correspondence should be addressed (email: ezekiel.ahn@usda.gov; ORCID: https://orcid.org/0000-0003-4447-410).

Abstract

Can standardized chemical stress reveal an ecological "memory" of host origin in plant pathogens? We profiled six coffee-associated *Colletotrichum* isolates using four phenolic-branched compounds and compared them with a previously characterized cacao panel. Quantitative morphology, hyperspectral imaging (HSI), and supervised machine learning (ML) yielded pathosystem-specific fingerprints under uniform, isotropic *in vitro* conditions. Circularity, a measure of edge symmetry, was the most informative morphological feature, and ML classified host origin with 87% accuracy in cross-validation. HSI detected dose-dependent physiological shifts, including changes near 1930 nm in the short-wave infrared consistent with water-associated bands. Multi-locus phylogeny showed the coffee isolates are polyphyletic, indicating that these fingerprints likely reflect convergent adaptation rather than shared ancestry. We propose a "chemical priors" framework in which long-term environmental exposure imprints stress-response

35   pathways that become legible under simple, standardized probes. This integrative workflow

36   supports scalable screening of eco-friendly antifungals and precision disease management.

38   **Keywords:** Ecological Memory, Machine Learning, Fungal Adaptation, Pathosystem-specificity,

39   Hyperspectral Imaging, Morphological Fingerprint, *Colletotrichum*, Phenolic Antifungals

1. Introduction

Fungal pathogens like *Colletotrichum* threaten global food security, creating an urgent need for

novel, sustainable control strategies that require a deeper understanding of pathogen adaptation [1–

5]. A fundamental challenge is decoding an organism's adaptive history from its observable traits,

a problem we address here using an integrated spectroscopy and machine learning approach. To

this end, we turned to the anthracnose pathogen *Colletotrichum*. While notorious for its broad host

range, we selected its interaction with two globally significant perennial crops, cacao (*Theobroma*

cacao) and coffee (*Coffea arabica*). Their distinct chemistries and ecologies make them ideal

natural experiments for testing the generality of our core hypotheses. For decades, the primary

defense against such pathogens has been the application of synthetic chemical fungicides.

However, this strategy is increasingly challenged by the evolution of fungicide-resistant pathogen

strains and growing concerns about the environmental and toxicological effects of chemical

residues [6–8]. This creates an urgent need for novel antifungal strategies that are not only effective

but also ecologically sustainable and are often sought from natural sources [9, 10]. This led us to a

central hypothesis: that a pathogen's observable response to chemical stress is an echo of its

ecological past. To test this, we developed a novel, high-throughput workflow combining

quantitative morphology, hyperspectral imaging (HSI), and machine learning (ML). We contend

that this integrative platform provides a powerful new lens for studying microbial adaptation and

has significant practical applications in phytopathology.

The distinct ecologies of coffee and cacao pathogens provide an ideal natural experiment to deploy

this workflow. Here, we applied our integrative approach to decode what we call 'chemical

priors'—stress-response phenotypes that reveal a pathogen's ecological memory. To elicit these

phenotypes, we utilized a 2x2 matrix of novel, stabilized phenolic-branched antifundals derived

from sustainable feedstocks [11–15]. These compounds, with systematic variations in their

65 phenolic headgroups and backbones, allow us to probe the structural basis of antifungal activity,

66 building upon our initial findings in the cacao pathosystem where specific derivatives induced

67 reproducible morphological fingerprints [16].

69 This observation prompted a central question: do pathogens from different agroecosystems carry

70 a 'memory' of their origin, one that is legible in their response to chemical stress? From this

71 question arise two competing hypotheses. This memory could be imprinted by either the

72 immediate host-pathogen interaction or by larger-scale biogeographical factors. The distinction is

73 critical, as it determines whether a morphological response indicates a universal chemical

74 mechanism or a pathosystem-specific adaptation. Given the contrasting biochemical environments

75 of cacao and coffee plantations, each defined by distinct host metabolites, microbial communities,

76 and abiotic pressures, it is plausible that fungal pathogens from these systems have evolved

77 divergent physiological strategies for coping with xenobiotic stress (i.e., exposure to biologically

78 foreign compounds, including our phenolic derivatives). For instance, cacao agroecosystems are

79 typically characterized by the humid, shaded conditions of their native tropical understory [17],

80 while arabica coffee is frequently grown in more open, higher-altitude environments, often above

81 1,000 meters [18, 19]. Furthermore, the host plants themselves produce distinct classes of defensive

82 secondary metabolites that create unique chemical landscapes. Cacao relies on a defense chemistry

83 rich in methylxanthines, primarily theobromine and caffeine, as well as compounds from the

84 phenylpropanoid pathway [20]. In contrast, coffee leaves maintain high concentrations of caffeine

85 and chlorogenic acid to suppress the growth of microbial pathogens [21]. To test this hypothesis,

86 we utilized the identical four phenolic-branched compounds previously tested in cacao: a phenol-

87 soy oil fatty acid (PhSOFA) and a beechwood creosote-branched grease fatty acid (BCBGFA),

88 both consisting of a phenolic headgroup covalently linked to a fatty acid backbone, and their

89 respective amide analogs, PhSOAM and a beechwood creosote-branched grease amide

90 (BCBGAM), which replace the terminal carboxylic acid with an ethylenediamine-derived amide

91 group. PhSOFA contains a simple, unsubstituted phenol, while BCBGFA incorporates a complex

92 mixture of substituted phenols derived from beechwood creosote. Their amide counterparts

93 (PhSOAM and BCBGAM) preserve the phenolic identity but differ in their increased polarity and

94 nitrogen content. These systematic structural variations allow us to probe how differences in

95 phenolic complexity and headgroup chemistry influence antifungal activity and the resulting

morphological response. Thus, the four compounds represent a 2×2 matrix of structural variation:

simple vs. complex phenol and acid vs. amide headgroup, allowing direct inference of both

phenolic and backbone contributions to antifungal activity.

Our primary goal was to establish and validate an integrated workflow that can decode these

signatures. By applying this platform to the contrasting coffee and cacao pathosystems, we aimed

to: (1) quantify compound-induced morphological fingerprints; (2) use ML to test their predictive

power for pathogen origin; and (3) use HSI to explore the underlying physiological basis of these

predictive phenotypes.

106 2. Materials and methods

107 2.1 Biobased compounds tested as antifungal agents

108   Four previously described phenolic-branched compounds (PhSOFA, PhSOAM, BCBGFA, and

109   BCBGAM), synthesized from biobased feedstocks, were used in this study [13, 15]. Stock solutions

110   (10 mg/mL) were prepared in pure ethanol for all assays.

112 Fungal isolates and culture conditions

113   Six coffee-pathogenic *Colletotrichum* spp. isolates were used, with origins and host details

114   provided in Table S4. For comparative analyses, data from three previously characterized cacao-

115   pathogenic isolates were also included. All isolates were maintained on Potato Dextrose Agar

116   (PDA) at 24°C in the dark.

118 2.2 Phylogenetic analysis

To determine the evolutionary relationships among isolates, we performed a multi-locus

phylogenetic analysis using the internal transcribed spacer (ITS), glyceraldehyde-3-phosphate

dehydrogenase (GAPDH), and actin (ACT) gene regions. Phylogenetic trees were constructed in

MEGA 11 using the Neighbor-Joining method with nodal support assessed by 1,000 bootstrap

replications. Detailed procedures for DNA extraction, PCR amplification, sequencing, and

alignment are provided in the Supplementary Methods.

126 2.3 Antifungal activity assay and morphological Analysis

Antifungal activity was assessed using a modified agar plate assay. Each compound (150 µg) was
spread on PDA plates, with ethanol-treated plates serving as controls. A 5-mm mycelial plug of
each isolate was placed at the center and incubated for 96 hours. Digital images of the colonies
were captured under consistent lighting conditions. Colony morphology was quantified using
SmartGrain software to measure seven traits: area, perimeter, length, width, length-to-width-ratio
(LWR), circularity, and an asymmetry distance ((the distance between the intersection of length
and width and the center of gravity (IS & CG)) [22].

2.4 Statistical and machine learning analysis

All statistical analyses were performed using JMP Pro 17 [23]. For the comparative analysis, the
coffee pathogen dataset was merged with the previously published cacao pathogen dataset [16].
One-way ANOVA followed by Tukey's HSD test was used for pairwise comparisons. For
multivariate analyses, all morphological data were log-transformed. We used Principal Component
Analysis (PCA) to visualize overall relationships and Permutational Multivariate Analysis of
Variance (PERMANOVA) to test for group differences. To predict a pathogen's host of origin, we
evaluated a suite of machine learning classification models using a repeated K-Fold cross-
validation strategy (k=5) on a class-balanced dataset. Full details on data transformation and model
parameters are available in the Supplementary Methods.

2.5 HSI and data analysis

HSI was conducted on select isolates cultured on PDA with varying compound concentrations (0,
150, 300, and 450 µg per plate). Three imaging modalities were employed using custom-built line-
scan systems: fluorescence (450–720 nm), visible and near-infrared (VNIR) reflectance (430–1000
nm), and short-wave infrared (SWIR) reflectance (1000–2400 nm). Spectral data were extracted
from standardized regions of interest (ROIs), calibrated, and analyzed using one-way ANOVA F-
statistics to identify wavelengths with significant treatment-induced variation [24]. Detailed system
specifications are provided in the Supplementary Methods.

3 Results
3.1 Potent but distinct effects of amide derivatives PhSOAM and BCBGAM

The two ethylenediamine-carrying amide derivatives, PhSOAM and BCBGAM, were among the

most potent growth inhibitors tested. PhSOAM exhibited potent but selective antifungal activity,

reducing colony area in the most susceptible isolates, P24-192 and P24-85, by 19.4% ($p < 0.001$)

and 20.7% ($p < 0.001$), respectively (Fig. S1A). This reduction is visually illustrated for P24-192

(Fig. S1B, C). In contrast, BCBGAM induced slightly stronger inhibition in certain isolates (e.g.,

20.1% in P24-192) but was less effective against others (Fig. 1A). While both amides reduced

size metrics, their shape metrics (circularity, LWR, IS&CG) diverged, underscoring a size–shape

decoupling central to this study. PhSOAM triggered diverse morphological responses, in some

cases inducing irregular expansion (e.g., a decrease in circularity in isolate P24-193) (Fig. S1A).

Conversely, BCBGAM promoted a more compact, rounded morphology in three of the six

isolates (Fig. 1A). The most striking response was observed in isolate P24-192, where BCBGAM

treatment induced a significant increase in colony elongation (LWR), as illustrated in the

representative images (Fig. 1).

Fig. 1. Representative morphological disruption by phenolic antifungal compounds, illustrated using BCBGAM-treated isolates, including P24-192. (A) Box plots quantifying the morphological response of six *Colletotrichum* sp. isolates to one representative phenolic antifungal (BCBGAM), which exhibited isolate-specific effects. BCBGAM is featured because it elicited the most visually distinct morphological changes in isolate P24-192. While PhSOAM showed stronger overall inhibition (Fig. S1), the response to BCBGAM better illustrates the complex 'morphological fingerprint' central to this study, highlighting our focus on decoding characteristic stress responses rather than solely quantifying growth inhibition. Measured traits include colony area (mm²), perimeter (mm), length (mm), width (mm), LWR, circularity, and an asymmetry distance (IS & CG, mm). Significant differences between a treated group and its corresponding control (Tukey's HSD; *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$) are marked with asterisks. (B-C) Representative images of isolate P24-192 after 96 hours on (B) a control plate and (C) a plate treated with BCBGAM. Sample size, n = 10 for all groups except P24-192 and P24-193 controls (n = 11). Scale bars = 1 cm.

3.2 Variable and opposing effects of fatty acid analogs

The two phenol-branched fatty acids, PhSOFA and BCBGFA, exhibited overall weaker antifungal
activity than their corresponding amide derivatives, yet each triggered highly distinctive and
isolate-specific morphological responses (Fig. 2; Fig. S2–S3).

PhSOFA demonstrated selective growth inhibition, significantly reducing colony size in two of
the six isolates: P24-83 and P24-192 (Fig. S2A). For example, PhSOFA treatment reduced the
colony area of isolate P24-83 by 18.8% ($p < 0.01$), a change illustrated in the representative images
(Fig. S2B–C). In contrast, BCBGFA's effects were highly variable. It had no significant inhibitory
effect on any isolate but paradoxically promoted the growth (assessed by area size measurement)
of P24-85 by 16.5% ($p < 0.05$), as detailed in Fig. S3. The remaining isolates were largely resistant
to either compound. Morphological shape metrics highlighted further, though more sporadic,
divergence between the two treatments. The effects on shape metrics were more sporadic:
significant increases in colony circularity were rare, observed only for BCBGFA in P24-83 and
PhSOFA in P24-193, while PhSOFA was the only compound to cause significant increases in both
colony elongation (LWR) and asymmetry (IS & CG), effects that were limited to isolate P24-192
(Figs. S2 & S3).

When morphological data were pooled across all isolates (Fig. 2), a clear hierarchy of antifungal
potency, based on the reduction in colony area, was observed. PhSOAM remained the most potent
growth inhibitor. It was followed by PhSOFA. The other amide, BCBGAM, showed no significant
difference from the control, while BCBGFA was the least effective, showing no significant
inhibition. Despite variable effects on colony size, the compounds induced reproducible shifts in
shape. PhSOFA uniquely elevated the LWR. A significant increase in colony circularity was
observed for three compounds: PhSOFA, BCBGAM, and BCBGFA. Furthermore, only PhSOFA
caused a significant increase in the asymmetry distance (IS & CG), pointing to non-lethal but
geometry-altering cellular effects. Together, these data show isolate- and structure-dependent
effects across compounds (Fig. 2; Figs. S2–S3).

212
213

Fig. 2. Comparative analysis of the antifungal effects of four phenolic-branched compounds on coffee-pathogenic *Colletotrichum* isolates. Box plots show the pooled morphological data from six *Colletotrichum* isolates under four treatment conditions (PhSOAM, BCBGAM, PhSOFA, and BCBGFA) and an untreated control. Measured traits include colony area (mm²), perimeter (mm), length (mm), width (mm), LWR, circularity, and an asymmetry distance (IS & CG, mm). Different letters denote statistically significant groupings ($p < 0.05$) determined by Tukey's HSD.

3.3 Multivariate analysis reveals a dominant treatment effect on fungal morphology

PCA of the morphological traits showed that the first two PCs captured 76.2% of the total variance (Fig. S4). However, both chemical treatments and fungal isolates showed considerable overlap in PCA space, indicating that linear ordination provided limited separation between groups. This necessitated the use of supervised machine learning to test for nonlinear, diagnostic patterns. Permutational multivariate analysis of variance (PERMANOVA) within the coffee dataset indicated a small but significant treatment effect and a stronger isolate effect (Compound: pseudo-$F = 4.06$, $R^2 = 0.0518$, $p = 0.001$; Strain: pseudo-$F = 13.00$, $R^2 = 0.180$, $p < 0.0001$). This partial grouping in PCA, combined with substantial overlap, suggests that linear ordination provides limited separation. Because PCA did not fully resolve isolates or pathosystem identity, we next used supervised ML to test for nonlinear, diagnostic patterns.

Further analyses were conducted to explore the relationships between the seven morphological traits (Fig. S5). The PCA loading plot (Fig. S5A) and partial contributions plot (Fig. S5B) confirmed that the primary axis of variation, PC1, was driven by the four main size metrics (area, perimeter, length, and width). Shape-related traits had greater impacts on subsequent components: LWR and IS & CG contributed most to PC2, while circularity was a major driver of PC3. This clear distinction between size and shape was corroborated by hierarchical clustering, which grouped the four size metrics into one tight cluster and the three shape metrics into another (Fig. S5C). Pearson's correlation analysis further quantified these relationships, revealing strong positive correlations among all size-related traits ($r > 0.9$ for most pairs, detailed in Supplementary Data 1).

3.4 Phylogenetic analysis reveals coffee-pathogenic isolates are polyphyletic and intermixed with cacao-associated isolates

To determine the evolutionary relationships among the isolates, we first conducted a multi-locus phylogenetic analysis using a concatenated dataset of the internal transcribed spacer (ITS), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and actin (ACT) gene regions (Fig. 3). The analysis revealed the six coffee-pathogenic isolates were polyphyletic, distributing across several distinct clades rather than forming a monophyletic group.

Specifically, isolate P24-88 formed a sister group with *C. karstii* (79% bootstrap support), and this

clade in turn was a sister group to isolate P24-193. Isolate P23-55 was placed in a separate, distant

clade with *C. pseudoboninense*. The remaining three coffee isolates were located in different parts

of the tree: P24-192 clustered with *C. siamense* (99% bootstrap), P24-83 was a sister taxon to *C.*

rhizophorae and *C. fructicola*, and P24-85 was placed near *C. gloeosporioides* and *C. kahawae*.

This polyphyletic distribution indicates that the coffee pathogens used in this study represent

multiple distinct lineages within the *Colletotrichum* genus.

An analysis based on the standard fungal barcode, the ITS region, was also performed to directly

compare the coffee isolates with cacao-associated isolates (Fig. S6). This analysis further

highlighted the complex relationships between isolates from the two pathosystems. Notably, a

single, strongly supported clade (99% bootstrap) contained an admixture of isolates from different

hosts and origins. This inclusive clade grouped coffee pathogens P24-192, P24-85, and P24-83

together with cacao isolates (CGH17, CGH53) and shade tree isolates (CGH34, CGH38). Overall,

based on the ITS region, some coffee pathogens are more closely related to cacao-associated

isolates than to other coffee pathogens, such as P24-88 and P24-193, which formed a separate,

well-supported clade (96% bootstrap).

Fig. 3. Multi-locus phylogeny reveals the polyphyletic nature of coffee-pathogenic *Colletotrichum* isolates

The phylogenetic tree was inferred by the Neighbor-Joining method using a concatenated alignment of the ITS, GAPDH, and ACT sequences. The evolutionary distances were computed using the Tamura-Nei (TN+G) model. Numbers at the nodes represent bootstrap support values based on 1,000 replications; only values of 75% or greater are shown. The six coffee-pathogenic isolates (highlighted in blue) are shown to be polyphyletic, distributing across multiple distinct clades and indicating affiliations with different known *Colletotrichum* species.

3.5 Comparative multivariate analysis of coffee and cacao pathosystems

To investigate whether the chemical compounds induced environment-specific morphological responses, the coffee pathogen dataset was combined with a previously generated dataset from cacao pathogens[16], and a comparative PCA was performed on the combined, log-transformed data.

The first two principal components accounted for 78.6% of the total variance (PC1: 56.8%, PC2: 21.8%) (Fig. 4). When the data were colored by the pathogen's system of origin (Fig. 4A), no clear separation was evident. The coffee (black) and cacao (red) samples were heavily intermixed. PERMANOVA detected a weak host effect (Host: pseudo-F = 8.49, R^2 = 0.0174, p = 0.0017) and a small but significant treatment effect (Compounds: pseudo-F = 5.57, R^2 = 0.0447, p < 0.001), consistent with the minimal variance explained by these factors.

When visualized by treatment type, the analysis showed significant overlaps between the groups (Fig. 4B). The untreated control samples clustered closely with those treated with BCBGFA. A separate cluster was formed by samples treated with PhSOAM and PhSOFA, while the BCBGAM-treated samples were positioned between these two main groupings. This strong overlap indicates that the morphological signatures distinguishing the two pathosystems, if they exist, are not linearly separable and thus are too subtle for PCA to resolve.

Fig. 4. Comparative PCA of coffee and cacao pathosystems reveals treatment-driven separation but pathosystem overlap. PCA of morphological traits from the combined coffee and cacao pathogen datasets (log-transformed). (A) PCA scores plot with samples color-coded by pathosystem: Cacao pathogens (red) and Coffee pathogens (black). Note the significant overlap between the two pathosystems, indicating the first two principal components cannot separate them. (B) The identical PCA scores plot with samples color-

303   coded by chemical treatment: Control (black), PhSOAM (orange), PhSOFA (green), BCBGAM (blue), and
304   BCBGFA (purple). The red squares indicate the mean of each pathosystem group in (A) and each chemical
305   treatment group in (B). PERMANOVA indicated a weak host effect (Host: pseudo-F = 8.49, R^2 = 0.0174,
306   p = 0.0017) and a small but significant treatment effect (Compound: pseudo-F = 5.57, R^2 = 0.0447, $p <$
307   0.001), both explaining only a small proportion of total variance.

3.6 Machine learning models successfully classify pathogen host origin

Following the PCA, which could not clearly separate the two pathosystems, a suite of ML models

was employed to quantitatively test if a predictive morphological signature of host origin existed

within the combined dataset. The analysis revealed that several models could classify a pathogen's

host of origin with high accuracy, confirming the existence of a robust 'fingerprint' based on the

response to the chemical compounds (Fig. 5). A comprehensive screening of nine different

algorithms identified the Neural Boosted model (an F1-score of 0.87, and a Matthews Correlation

Coefficient (MCC) of 0.73) as the top performer, achieving a validation accuracy of 87%, which

was followed closely by Nominal Logistic and Generalized Regression (Lasso) models (both

85%), as well as Support Vector Machines, Bootstrap Forest, and Fit Stepwise (all 84%) (Table

S1).

To understand which traits were driving this successful classification, feature importance was

analyzed using the top-performing Neural Boosted model (Table S2). The analysis revealed that

circularity was the single most dominant predictor of host origin, with a Total Effect score of

0.592, which was followed by the primary size-related traits of area (Total Effect = 0.405) and

perimeter (Total Effect = 0.168). This indicates that the model primarily learned to distinguish the

pathosystems by identifying a unique 'shape signature', driven by how circular a colony becomes

in response to chemical stress, complemented by information about the colony's overall size.

To further test the robustness of the classification, the analysis was repeated on the log-transformed

dataset after removing circularity, the top predictor (full details in Supplementary Data 1). Even

without this primary feature, the models maintained high predictive power. The Neural Boosted

model was again the top performer with a validation accuracy of 86.1%, followed by the Nominal

Logistic (85%), Lasso (85%), and Fit Stepwise (84%) models. Feature importance analysis for this

new model revealed a clear shift in predictive strategy; in the absence of circularity, the model

pivoted to a 'size signature', with area size (Total Effect = 0.825) and perimeter (Total Effect =

0.467) emerging as the most dominant predictors (Supplementary Data 1), demonstrating that the

morphological 'fingerprint' distinguishing the pathosystems is robust and encoded in multiple features.

Fig. 5. Machine learning classification of pathogen host origin based on chemical-induced morphology. Decision thresholds plot showing the performance of nine ML models in classifying a

3.7 Dose-dependent hyperspectral fingerprints in CGH5 under PhSOAM treatment

To probe the physiological basis of these morphological fingerprints, we first employed three modalities of HSI: fluorescence HSI, visible and near-infrared (VNIR) reflectance imaging, and short-wave infrared (SWIR) reflectance imaging. We began by testing the response of CGH5, a cacao-derived isolate known to exhibit a dramatic 48.8% reduction in fungal growth upon PhSOAM treatment [16]. HSI revealed distinct, dose-dependent responses of *Pestalotiopsis* sp. CGH5 to PhSOAM treatment across all three spectral modalities (Fig. 6).

Fig. 6. Hyperspectral imaging of *Pestalotiopsis* sp. CGH5 treated with the biobased fungicide PhSOAM across three spectral modalities. (A) Fluorescence intensity profiles of CGH5 colonies

358 treated with 0, 150, 300, and 450 µg of PhSOAM; color indicates dose (control=blue, 150 µg = orange,
359 300 µg = green, 450 µg= red). (B) F-value curve corresponding to the fluorescence data. (C) Reflectance
360 intensity curves under broadband illumination. (D) F-value profile corresponding to the reflectance data.
361 (E) SWIR reflectance curves showing spectral shifts. (F) F-value curve corresponding to the SWIR data.
362 All data represent the mean of four biological replicates per treatment (n = 4).
363

3.8 Attenuated hyperspectral response in a tolerant coffee isolate
In contrast to the strong response of CGH5, the more tolerant coffee isolate P24-85 exhibited an

attenuated and nonlinear hyperspectral response to PhSOAM (Fig. 7). This included a non-

monotonic fluorescence signal and muted shifts in the SWIR water-associated bands. These

attenuated spectral changes are consistent with a more resilient physiological state capable of

active damage management.

Fig. 7. Hyperspectral imaging of *Colletotrichum* sp. P24-85 treated with the biobased fungicide
PhSOAM across three spectral modalities. (A) Fluorescence intensity profiles of P24-85 colonies
treated with 0 µg (control, blue), 150 µg (orange), 300 µg (green), and 450 µg (red) of PhSOAM. (B) F-
value curve corresponding to the fluorescence data. (C) Reflectance intensity curves under broadband

illumination. (D) F-value profile corresponding to the reflectance data. (E) SWIR reflectance curves. (F) F-value curve corresponding to the SWIR data. All data represent the mean of four biological replicates per treatment (n = 4).

3.9 Compound-specific hyperspectral fingerprints of *Colletotrichum* sp. P24-192

Different compounds also induced unique spectral fingerprints in the same isolate, *Colletotrichum* sp. P24-192, suggesting divergent modes of action (Fig. S7). For example, in the SWIR modality, BCBGAM produced a focused diagnostic band near ≈1950 nm, whereas PhSOAM yielded a broader separation across multiple bands. These distinct spectral signatures are consistent with compound-specific biochemical interactions with the fungus.

4 Discussion

Here, we established that a microbe's response to a uniform stressor can reveal its ecological history. More importantly, we developed and validated a powerful workflow to decode this "ecological memory." By integrating quantitative morphology, HSI, and ML, we showed that pathosystem-specific fingerprints could predict an isolate's host of origin with high accuracy. Our work demonstrates that this scalable, phenotype-driven platform is a powerful tool for studying how ecological history shapes microbial adaptation, with direct applications in screening new antifungals and decoding complex morphological signatures.

The varied fungal responses underscored a clear structure-activity relationship dependent on both the headgroup and backbone of the four phenolic compounds. Consistent with previous reports, amide derivatives generally showed stronger antifungal activity than their fatty acid analogs. Specifically, PhSOAM, with its simple phenol and amide moiety, was the most potent growth inhibitor. This suggests that both the increased polarity from the amide group and the steric accessibility of the unsubstituted phenol contribute to efficacy [12, 25, 26].

An unexpected observation was that BCBGFA appeared to promote growth in isolate P24-85. While this may reflect experimental variability, an alternative explanation is that certain fungal isolates possess enzymatic adaptations that allow them to tolerate or even metabolize specific phenolic compounds. This possibility is supported by earlier studies demonstrating that some fungi can grow on phenolic acids and related aromatic compounds [27, 28].

This divergence in morphological response suggests a deeper biological basis rooted in ecological

adaptation. We propose that long-term exposure to a native environment, whether a specific host,

soil habitat, aquatic niche, or built environment, imprints an ecological memory in stress-response

pathways. When challenged with a novel stressor, this legacy is expressed as what we term a

'chemical prior': a reproducible, system-specific phenotypic fingerprint. In this study, the coffee–

cacao comparison serves as one example, but the same principle could apply to other fungi,

bacteria, protists, or mixed microbial communities [29, 30]. While our previous work showed that

PhSOAM induces a dramatic increase in colony elongation (LWR) in susceptible cacao pathogens

[16], this response was attenuated and inconsistent among coffee isolates. Notably, P24-88

exhibited moderate elongation, but this phenotype was not broadly observed across the coffee

panel. Instead, circularity emerged as the more robust and pathosystem-distinctive morphological

trait. Intriguingly, this response is not limited to chemical insults; it is further reinforced by our

earlier findings where physical (UVC) stress also induced increased circularity in multiple cacao

isolates [24]. Circularity may therefore be a generalizable indicator of stress, as disparate pressures

from chemical and UVC treatments appear to converge on a similar morphological outcome.

While linear methods like PCA failed to separate the pathosystems, supervised machine learning

successfully classified host origin with 87% accuracy, validating the existence of a predictive

'interaction phenotype' [31, 32]. Feature-importance analysis consistently revealed colony

circularity, a measure of isotropic growth, as the most robust biomarker of this latent ecological

memory. The model's reliance on this shape metric, even after its removal from the feature set,

indicates the fingerprint is robustly encoded across multiple morphological traits [33, 34]. The

importance of circularity as a stress indicator is also consistent with recent work on biotic

antagonism [35–37].

We interpret this circularity shift as a direct readout of the pathogen's pre-programmed growth

machinery responding to a uniform, isotropic stress field. An alternative hypothesis, that these

shifts reflect rapid selection of resistant mutants, is unlikely. The 96-hour incubation is insufficient

for the multi-generational selection typically required for such changes[38, 39], and the consistent,

pathosystem-wide nature of the fingerprint is inconsistent with the stochastic nature of mutation

[40]. Therefore, we conclude these phenotypes represent plastic responses reflecting an evolutionarily imprinted memory [41].

HSI provided physiological evidence for this ecological memory, revealing stressor- and isolate-specific spectral fingerprints. Previous studies show that near- and shortwave-infrared reflectance and fluorescence spectroscopy can distinguish fungal physiological responses, including hydration level, pigmentation, and cell-wall composition [42–45]. In our study, strong isolate- and dose-dependent signals emerged across all three imaging modalities. HSI provided a physiological basis for these fingerprints, revealing two distinct stress-response archetypes. The susceptible cacao isolate CGH5 exhibited strong spectral shifts, including increased fluorescence (470–520 nm) and prominent absorption near water-associated SWIR bands ($\approx$1930 nm), potentially indicating a loss of homeostasis or 'metabolic collapse' [24, 42, 46, 47]. In contrast, the more tolerant coffee isolate P24-85 showed a muted or attenuated spectral response at these same bands, suggesting an active 'damage management' strategy that maintains physiological stability under stress. This attenuated spectral response, particularly at water-associated bands, may reflect a distinct and potentially more resilient physiological state under chemical stress. Consistent with the SWIR water band and the circularity shift, this resilience may arise from rapid water-balance control (osmolyte accumulation, aquaporin gating) coupled with cell-wall remodeling at the colony boundary. These mechanisms are hypotheses consistent with our readouts and will require direct molecular validation.

The spectral contrast observed between PhSOAM and BCBGAM treatments further suggests divergent biochemical interactions (Fig. S7, Table S3). These compound-specific signatures, likely reflecting distinct impacts on cell hydration, membrane lipids, and phenolic compounds, require future molecular validation to elucidate precise mechanisms [24, 48]. Because the $\approx$1930 nm band is highly moisture-sensitive, future work should explicitly model or filter water effects (e.g., variable band weighting or EMSC) to isolate biochemical signals [47]. Interestingly, some of these morphological and spectral stress markers, particularly increased circularity and shifts in SWIR bands, converge with those we previously observed under physical (UVC) stress, suggesting a potentially generalizable stress-response program in these pathogens [24].

These parallels between xenobiotic chemical exposure and UVC irradiation suggest that HSI can capture both compound-specific and isolate-specific physiological responses. The recurrence of CGH53's specific spectral features under both conditions raises the possibility that its native agroecosystem has shaped a stress-response pathway that is preferentially activated when homeostasis is disrupted. This pathway, once triggered by a sufficiently strong stressor, may lead to a cascade of metabolic and physiological changes, though the precise molecular mechanisms remain to be determined.

Crucially, the convergence of findings from macroscopic colony morphology and physiological hyperspectral signatures suggests the observed "ecological memory" is a robust biological signal. This multi-modal evidence allows us to propose a conceptual model of two contrasting physiological archetypes: a 'metabolic collapse' in susceptible isolates like CGH5, characterized by spectral shifts indicative of lost homeostasis, and a buffered 'damage management' in tolerant isolates like P24-85. This framework, while requiring future molecular validation, provides a powerful new model for understanding the diverse strategies pathogens employ under chemical stress.

The polyphyletic nature of the coffee isolates provides critical evolutionary context for our findings (Fig. 3, Fig. S6). The fact that these isolates do not form a single clade, and are sometimes more closely related to cacao pathogens, argues that the 'coffee fingerprint' is not a trait inherited from a single ancestor. Instead, this suggests convergent adaptation, where distinct lineages independently evolved similar stress-response strategies when faced with the unique chemical landscape of the coffee agroecosystem [49]. This interpretation is supported by the ability of our ML models to classify isolates by their host of origin, detecting a true, host-imprinted ecological signal that transcends the complex phylogenetic backdrop and the limited resolution of ITS-based phylogenies.

While our data suggest host adaptation as the primary driver, we do not rule out the influence of biogeography and lineage-specific traits, although direct biogeographical effects were not explicitly tested in this study. The coffee isolates' distribution across multiple clades, some of which are shared with isolates from a geographically distinct cacao agroecosystem, underscores

the complex interplay between evolutionary history and environmental adaptation [50, 51].
Disentangling these factors presents a critical and now-testable question for the field. Ultimately,
our work demonstrates that a pathogen's observable response to chemical stress is a rich,
multi-layered phenotype, reflecting a powerful ecological memory forged at the intersection of its
lineage and its environment.

To synthesize these findings, we propose a generalized conceptual model that links a microbe's
long-term ecological history to its observable stress-response phenotype (Fig. 8). We term this the
"chemical priors" framework, which posits that persistent exposure to a native environment,
whether a specific host, soil, or aquatic niche, imprints a latent "ecological memory" in
stress-response pathways. When challenged with a novel stressor, such as the phenolic compounds
used in this study, this embedded memory is expressed as a reproducible, system-specific
phenotypic fingerprint, such as the shifts in colony circularity observed here. This fingerprint can
then be decoded by supervised machine learning to accurately infer the pathogen's origin,
providing a quantitative readout of its ecological history. While demonstrated with the coffee–
cacao pathosystem, this framework offers an adaptable approach for studying microbial adaptation
across diverse biological contexts. This "chemical priors" framework suggests a form of heritable,
environmentally-induced memory, potentially rooted in epigenetic mechanisms that warrant future
investigation.

Fig. 8. Conceptual model of ecological memory and chemical priors in microbial response to environmental stress. (A) Long-term exposure of a microorganism to a specific native environment, whether a host–pathogen agroecosystem (e.g., coffee or cacao), soil, aquatic, or built environment, imprints an "ecological memory" shaped by environmental chemistry and associated microbial communities. (B) When challenged *in vitro* with a novel environmental stressor (chemical, physical, or biological), the fungus (spore; center in the panel for representative image) expresses a distinct, reproducible phenotypic fingerprint (e.g., circularity shift, hyperspectral change) shaped by this memory. (C) Supervised machine learning classifies these fingerprints to

infer the organism's ecological origin or adaptation history with high accuracy, quantitatively linking past ecological exposure to present-day phenotypic stress responses. (D) This framework, termed the "chemical priors" model, provides a testable hypothesis for how historical selective pressures are embedded in microbial stress physiology and can be leveraged to predict responses in novel environments. While illustrated here for coffee–cacao *Colletotrichum* under phenolic antifungal stress, the same conceptual workflow could be adapted to other microorganisms and environmental contexts, including bacterial, algal, and mixed microbial systems exposed to chemical, physical, or biological challenges.

Our study establishes a scalable, phenotype-driven platform for decoding microbial ecological memory. The immediate value of this methodological advance is its application in high-throughput screening of eco-friendly fungicides and precision disease management. While future work should aim to validate these findings in planta and resolve their molecular bases, the "chemical priors" framework and the integrative workflow presented here provide a transformative new tool for the phytopathology and microbial ecology communities.

Our study establishes a scalable, phenotype-driven platform for decoding microbial ecological memory. The immediate value of this methodological advance is its application in high-throughput screening of eco-friendly fungicides and precision disease management. Building on this foundation, future work will prioritize three key directions: (1) validating these phenotypic fingerprints *in planta* to connect them to real-world disease outcomes; (2) resolving the underlying molecular and genomic bases of these stress responses to move from prediction to mechanism; and (3) testing the generality of the "chemical priors" model across diverse pathogen systems to establish it as a universal principle.

Acknowledgements

Mention of any trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U. S. Department of Agriculture. USDA is an equal opportunity provider and employer, and all agency services are available without discrimination. This work was supported by the U.S. Department of Agriculture, Agricultural Research Service, In-House Projects Nos. 8042-21220-258-000-D and

562   8072-41000-113-000-D, and by the National Institute of Food and Agriculture (NIFA) under Grant

563   No. 2021-67021-34502. We thank Lisa Keith (Daniel K. Inouye U.S. Pacific Basin Agricultural

564   Research Center, USDA-ARS, Hilo, Hawaii) for kindly providing the *Colletotrichum* isolates used

565   in this study.

Competing interests

568   The authors declare no competing interests.

Author contributions

571   E.A. conceptualized and supervised the project. A.L., C.H., and M.C. developed the

572   methodology. L.K., C.H., S.O., and E.A. conducted the experiments. I.B., S.L., and E.A.

573   performed the formal analysis. M.K-R., H.N., R.D.A., and A.L. provided resources. E.A.,

574   L.W.M., and M.S.K. acquired funding. E.A. wrote the original manuscript draft, and all authors

575   contributed to reviewing and editing the final version.

Data and materials availability

578   All raw and processed data, including colony morphology measurements, per-wavelength mean

579   spectra, and F-statistics for HSI, are provided in Supplementary Data 1. Multi-locus sequence data

580   (ITS, GAPDH, ACT) generated for this study are openly available at figshare:

581   https://figshare.com/articles/dataset/29924960.

Code availability

584   Not applicable.

Supplementary Materials

587   This PDF file includes: Figs. S1 to S7

588   Table S1 to S4

589   Supplementary methods

References

1.   Ferrucho RL, Marín-Ramírez GA, Gaitan A. Integrated Disease Management for the Sustainable

Production of Colombian Coffee. *Agronomy* 2024;**14**. https://doi.org/10.3390/agronomy14061286

2.   Ploetz R. The impact of diseases on cacao production: a global overview. *Cacao Dis Hist Old*

Enemies New Encount 2016;33–59.

3.   Kifle Belachew KB, Demelash Teferi DT. Climatic variables and impact of coffee berry diseases

(Colletotrichum kahawae) in Ethiopian coffee production. *J Biol Agric Healthc* 2015;**5**:55–64.

4.   Mohali Castillo SR, Miller S, Stewart J. Colletotrichum spp. and other fungi associated with

anthracnose on Coffea arabica L. in Mérida State, Venezuela. *Summa Phytopathol* 2022;**48**:99–111.

https://doi.org/10.1590/0100-5405/245876

5.   Ploetz RC. Cacao diseases: important threats to chocolate production worldwide. *Phytopathology*

2007;**97**:1634–1639.

6.   Cortaga CQ et al. Mutations associated with fungicide resistance in Colletotrichum species: A

review. *Phytoparasitica* 2023;**51**:569–592.

7.   Schnabel G et al. Inherent tolerance of Colletotrichum gloeosporioides to fludioxonil. *Pestic*

Biochem Physiol 2021;**172**:104767.

8.   Tsalidis GA. Human Health and Ecosystem Quality Benefits with Life Cycle Assessment Due to

Fungicides Elimination in Agriculture. *Sustainability* 2022;**14**. https://doi.org/10.3390/su14020846

9.   Cantrell CL, Dayan FE, Duke SO. Natural Products As Sources for New Pesticides. *J Nat Prod*

2012;**75**:1231–1242. https://doi.org/10.1021/np300024u

10. Thind TS. Role of Fungicides in Crop Health Management: Prospects and Challenges. In:

Satyanarayana T, Deshmukh SK, Johri BN (eds.), Developments in Fungal Biology and Applied

Mycology. Singapore: Springer Singapore, 2017, 433–447.

11. Borges DF et al. Formulation of botanicals for the control of plant-pathogens: A review. *Crop Prot*

2018;**110**:135–140. https://doi.org/10.1016/j.cropro.2018.04.003

12. Du S et al. Synthesis, antifungal activity and QSAR of some novel carboxylic acid amides.

Molecules 2015;**20**:4071–4087.

13.   Huang K et al. Structure-activity relationship of antibacterial bio-based epoxy polymers made from

phenolic branched fatty acids. *Prog Org Coat* 2021;**155**:106228.

14.   Kazem-Rostami M et al. Antibacterial agents from waste grease: arylation of Brown grease fatty

acids with beechwood creosote and derivatization. *ACS Sustain Chem Eng* 2023;**11**:17760–17768.

15.   Lew HN et al. Methods for preparing phenolic branched chain alkyl fatty acids or esters thereof and

methods for killing microorganisms. 2018.

16.   Ahn E et al. Structure-Activity Relationship of Novel Phenolic-Branched Amides Reveals Potent,

Isolate-specific Antifungal Activity and Morphological Disruption of Cacao Fungal Pathogens.

2025. ChemRxiv, 2025.

17.   Lahive F, Hadley P, Daymond AJ. The physiological responses of cacao to the environment and the

implications for climate change resilience. A review. *Agron Sustain Dev* 2019;**39**:5.

18.   Nugroho D, Basunanda P, Mw S. Physical bean quality of Arabica coffee (Coffea arabica)

cultivated at high and medium altitude. *Pelita Perkeb* 2016;**32**:151–161.

19.   Cassamo CT et al. Shade and altitude implications on the physical and chemical attributes of green

coffee beans from Gorongosa Mountain, Mozambique. *Agronomy* 2022;**12**:2540.

20.   Winters NP et al. A combination of conserved and diverged responses underlies Theobroma cacao's

defense response to Phytophthora palmivora. *BMC Biol* 2024;**22**:38.

https://doi.org/10.1186/s12915-024-01831-2

21.   de Sousa LP, Guerreiro Filho O, Costa Mondego JM. Differences between the leaf mycobiome of

Coffea arabica and wild coffee species and their modulation by caffeine/chlorogenic acid content.

Microorganisms 2021;**9**:2296.

22.   Tanabata T et al. SmartGrain: high-throughput phenotyping software for measuring seed shape

through image analysis. *Plant Physiol* 2012;**160**:1871–1880.

23.   Klimberg R. Fundamentals of predictive analytics with JMP. Sas institute, 2023.

24.   Baek I et al. Unveiling differential responses to UVB (305 nm) and UVC (275 nm) in cacao-

infecting Colletotrichum gloeosporioides and Pestalotiopsis sp. *bioRxiv* 2025;2025.05.21.655292.

644 https://doi.org/10.1101/2025.05.21.655292

645 25. Dembitsky VM. Microbiological aspects of unique, rare, and unusual fatty acids derived from

646  natural amides and their pharmacological profile. *Microbiol Res* 2022;**13**:377–417.

647 26. Ōmura S et al. Relationship between the structures of fatty acid amide derivatives and their

648  antimicrobial activities. *Antimicrob Agents Chemother* 1974;**6**:207–215.

649 27. Middelhoven WJ. Catabolism of benzene compounds by ascomycetous and basidiomycetous yeasts

650  and yeastlike fungi: A literature review and an experimental approach. *Antonie Van Leeuwenhoek*

651  1993;**63**:125–144.

652 28. Middelhoven WJ et al. Degradation of some phenols and hydroxybenzoates by the imperfect

653  ascomycetous yeasts Candida parapsilosis and Arxula adeninivorans: evidence for an operative

654  gentisate pathway. *Antonie Van Leeuwenhoek* 1992;**62**:181–187.

655 29. Brown AJ et al. Stress adaptation in a pathogenic fungus. *J Exp Biol* 2014;**217**:144–155.

656 30. Gasch AP. Comparative genomics of the environmental stress response in ascomycete fungi. *Yeast*

657  2007;**24**:961–976.

658 31. Carobene A et al. The multicenter European Biological Variation Study (EuBIVAS): a new glance

659  provided by the Principal Component Analysis (PCA), a machine learning unsupervised algorithms,

660  based on the basic metabolic panel linked measurands. *Clin Chem Lab Med CCLM* 2022;**60**:556–

661  568.

662 32. Tarca AL et al. Machine learning and its applications to biology. *PLoS Comput Biol* 2007;**3**:e116.

663 33. Lyou ES et al. Single-cell phenotypes revealed as a key biomarker in bacterial–fungal interactions: a

664  case study of Staphylococcus and Malassezia. *Microbiol Spectr* 2023;**11**:e00437-23.

665 34. Talento AF et al. A prospective study of fungal biomarkers to improve management of invasive

666  fungal diseases in a mixed specialty critical care unit. *J Crit Care* 2017;**40**:119–127.

667 35. Baek I et al. Dissecting Trichoderma antagonism: Role of strain identity, volatiles, biomass, and

668  morphology in suppressing cacao pathogens. *Biol Control* 2025;**207**:105807.

669  https://doi.org/10.1016/j.biocontrol.2025.105807

36. Zhang F et al. The stress response regulator AflSkn7 influences morphological development, stress

response, and pathogenicity in the fungus Aspergillus flavus. *Toxins* 2016;**8**:202.

37. Sterflinger K, Krumbein W. Multiple stress factors affecting growth of rock-inhabiting black fungi.

Bot Acta 1995;**108**:490–496.

38. Fisher KJ, Lang GI. Experimental evolution in fungi: An untapped resource. *Fungal Genet Biol*

2016;**94**:88–94.

39. Schoustra SE et al. Reducing the cost of resistance; experimental evolution in the filamentous

fungus Aspergillus nidulans. *J Evol Biol* 2006;**19**:1115–1127.

40. Mani G, Clarke BC. Mutational order: a major stochastic process in evolution. *Proc R Soc Lond B*

Biol Sci 1990;**240**:29–37.

41. Alster CJ et al. Phenotypic plasticity of fungal traits in response to moisture and temperature. *ISME*

Commun 2021;**1**:43.

42. Kong W et al. Estimating leaf water content at the leaf scale in soybean inoculated with arbuscular

mycorrhizal fungi from in situ spectral measurements. *Int J Agric Biol Eng* 2019;**12**:149–155.

43. Neurauter M et al. Exploring near-infrared spectroscopy and hyperspectral imaging as novel

characterization methods for anaerobic gut fungi. *FEMS Microbes* 2024;**5**:xtae025.

44. Teena M et al. Near infrared (NIR) hyperspectral imaging to classify fungal infected date fruits. *J*

Stored Prod Res 2014;**59**:306–313.

45. Valenzuela-Gloria M et al. Molecular characterization of fungal pigments. J Fungi (Basel) 7. 2021.

46. Melo D et al. Characterization of fungal melanins from black stains on paper artefacts. *Heritage*

2022;**5**:3049–3065.

47. Jochemsen A, Alfredsen G, Burud I. Hyperspectral imaging as a tool for profiling basidiomycete

decay of Pinus sylvestris L. *Int Biodeterior Biodegrad* 2022;**174**:105464.

48. Park E et al. Hyperspectral reflectance imaging for nondestructive evaluation of root rot in Korean

ginseng (Panax ginseng Meyer). *Front Plant Sci* 2023;**14**:1109060.

49. Rojas EI et al. Colletotrichum gloeosporioides sl associated with Theobroma cacao and other plants

in Panama: multilocus phylogenies distinguish host-associated pathogens from asymptomatic

endophytes. *Mycologia* 2010;**102**:1318–1338.

50.   Fumagalli M et al. Signatures of environmental genetic adaptation pinpoint pathogens as the main

selective pressure through human evolution. *PLoS Genet* 2011;**7**:e1002355.

51.   Kroymann J. Natural diversity and adaptation in plant secondary metabolism. *Curr Opin Plant Biol*

2011;**14**:246–251.

702