Taxonomy-agnostic hyperspectral–morphological phenotyping of fungal pathogen chemical-stress responses using machine learning

Insuck Baek[1], Seunghyun Lim[2], Amelia Lovelace[2], Sookyung Oh[1], Masoud Kazem-Rostami[3], Helen Ngo[3], Moon S. Kim[1], Lyndel W. Meinhardt[2], Lalit Kandpal[1], Minhyeok Cha[1], Chansong Hwang[1], Richard D. Ashby[3], Ezekiel Ahn[2,*]

[1] Environmental Microbial and Food Safety Laboratory, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, 20705, USA

[2] Sustainable Perennial Crops Laboratory, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, 20705, USA

[3] Sustainable Biofuels and Co-Products Research Unit, Agricultural Research Service, United States Department of Agriculture, Eastern Regional Research Center, Wyndmoor, PA, 19038, USA

* Author to whom correspondence should be addressed (email: ezekiel.ahn@usda.gov; ORCID: https://orcid.org/0000-0003-4447-410).

Highlights

- Taxonomy-agnostic workflow integrates hyperspectral imaging and morphology.
- Machine learning predicts crop-source labels with 86.7% accuracy in mixed lineages.
- System exhibits robustness, maintaining >86% accuracy even after feature ablation.
- Non-linear ML captures structure missed by linear stats (1.7% variance).
- Enables rapid, sensor-driven antifungal screening without prior DNA sequencing.

Abstract

Can standardized chemical stress reveal a reproducible, taxonomy-agnostic stress-response fingerprint that is predictive of sample-source labels (crop-of-isolation) in fungal isolates? Six coffee☐associated *Colletotrichum* isolates were profiled using four phenolic☐branched compounds and compared them with a previously characterized cacao panel. Quantitative morphology, hyperspectral imaging (HSI), and supervised machine learning (ML) yielded panel-specific fingerprints under uniform, isotropic *in vitro* conditions. Circularity, a measure of edge

symmetry, was the most informative morphological feature, and ML classified the crop-of-isolation label (coffee vs cacao, in this panel) with 86.7% accuracy in within-panel cross-validation. HSI detected dose-dependent spectral shifts in a targeted subset of isolates and compounds, including changes near 1930 nm in the short-wave infrared, a moisture-sensitive region that warrants robustness checks (e.g., band masking or preprocessing sensitivity) prior to biochemical attribution. Multi☐locus phylogeny showed the coffee isolates are polyphyletic, so the predictive signal should be interpreted conservatively as a taxonomy-agnostic phenotype fingerprint associated with crop background in this mixed-lineage panel, acknowledging that crop labels are partially confounded with phylogenetic structure. We propose a "chemical priors" framework as a working hypothesis, in which long-term environmental exposure may imprint stress-response pathways that become legible under simple, standardized probes. This integrative workflow supports scalable screening of eco-friendly antifungals and sensor-driven decision support for high-throughput phenotype-based screening workflows.

Keywords: Hyperspectral imaging, Machine Learning, High-throughput phenotyping, Taxonomy-agnostic, Antifungal screening

1. Introduction

Recent research in high-throughput phenotyping has increasingly demonstrated the utility of non-destructive sensors, particularly hyperspectral imaging (HSI) coupled with machine learning (ML), for evaluating plant-pathogen interactions and early disease detection (Rayhana et al., 2023; Tomaszewski et al., 2023). In particular, HSI and ML have been successfully deployed to detect latent anthracnose infections caused by *Colletotrichum* species in various agricultural commodities, including mango and citrus (Siripatrawan and Makino, 2024; Tang et al., 2023; Velásquez et al., 2024). However, while these technologies are widely used to diagnose infections in planta, their application for quantifying dynamic, compound-induced stress responses in vitro remains largely unexplored. Addressing this gap is critical for managing *Colletotrichum*, which causes severe anthracnose and berry diseases in globally significant perennial crops such as cacao (*Theobroma cacao*) and coffee (*Coffea arabica*). A fundamental challenge is extracting reproducible, predictive signals from observable stress phenotypes, a problem we address here using an integrated spectroscopy and machine learning approach. To

this end, the anthracnose pathogen *Colletotrichum* was selected as a model system. While

notorious for its broad host range, we selected its interaction with two globally significant

perennial crops, cacao and coffee. Their crop-associated isolate panels provide an applied testbed

for evaluating whether standardized chemical stress yields predictive phenotype fingerprints

under uniform and identical experimental conditions. For decades, the primary defense against

such pathogens has been the application of synthetic chemical fungicides. However, this strategy

is increasingly challenged by the evolution of fungicide-resistant pathogen strains and growing

concerns about the environmental and toxicological effects of chemical residues (Cortaga et al.,

2023; Schnabel et al., 2021; Tsalidis, 2022). This creates an urgent need for novel antifungal

strategies that are not only effective but also ecologically sustainable and are often sought from

natural sources (Cantrell et al., 2012; Thind, 2017). This led us to a central hypothesis: that a

pathogen's observable response to chemical stress can encode a stable, learnable phenotype

fingerprint that is useful for screening and classification. To test this, a novel, high-throughput

workflow was developed combining quantitative morphology, HSI, ML. We contend that this

integrative platform provides a powerful new lens for sensor-driven, non-destructive

phenotyping and predictive modeling with practical applications in phytopathology.

The two crop-associated isolate panels provide a practical context to deploy this workflow. Here,

we applied our integrative approach to quantify what we call 'chemical priors', stress-response

phenotypes that are hypothesized to reflect prior exposure histories or persistent physiological

states, and can be leveraged as predictive fingerprints. To elicit these phenotypes, we utilized a

2x2 matrix of novel, stabilized phenolic-branched antifungals derived from sustainable

feedstocks (Borges et al., 2018; Du et al., 2015; Huang et al., 2021; Kazem-Rostami et al., 2023;

Lew et al., 2018). These compounds, with systematic variations in their phenolic headgroups and

backbones, allow us to probe the structural basis of antifungal activity, building upon our initial

findings in the cacao panel where specific derivatives induced reproducible morphological

fingerprints (Ahn et al., 2026).

This observation prompted a central question: can standardized chemical-stress phenotypes

support robust, cross-validated prediction of source labels in a mixed-lineage isolate panel? From

this question arise two possible explanations. Any predictive signal associated with crop

background could reflect host-linked effects, biogeographical history, or lineage structure;

therefore, interpretations must remain conservative without species-resolved experimental
control. The distinction is critical, as it determines whether a phenotype fingerprint primarily
reflects compound-driven stress physiology or panel-specific biological context. Given the
differences in host chemistry and cultivation context reported for cacao and coffee systems
(Cassamo et al., 2022; de Sousa et al., 2021; Lahive et al., 2019; Nugroho et al., 2016; Winters et
al., 2024), it is plausible that fungal isolates sampled from these crops exhibit divergent stress-
response phenotypes; however, overlap between agroecosystems and lineage confounding can
also generate apparent differences, so we treat crop-of-isolation as an empirical label rather than
a mechanistic claim.

Our primary goal was to develop, evaluate, and benchmark an integrated workflow that can
quantify and model these signatures. By applying this platform to two crop-associated isolate
panels, the specific objectives were to: (1) quantify compound-induced morphological
fingerprints; (2) use ML to test their predictive power for crop-of-isolation labels (in this panel);
and (3) use HSI to explore the underlying physiological basis of these predictive phenotypes.

2. Materials and methods

2.1. Biobased compounds

The procedures for the production and characterization of phenolic soybean oil branched fatty
acids (PhSOFA) and their ethylenediamine-carrying fatty amide derivatives (PhSOAM) have
been previously reported in detail (Huang et al., 2021; Lew et al., 2018). The procedures for the
production and characterization of beechwood creosote-branched brown grease fatty acids
(BCBGFA) and their ethylenediamine-carrying fatty amide derivatives (BCBGAM) have also
been previously reported in detail (Kazem-Rostami et al., 2023). Each of the four final
compounds was dissolved in pure ethanol to create stock solutions of 10 mg/mL prior to
application.

2.2. Fungal isolates and culture conditions

Six coffee-associated fungal isolates of *Colletotrichum* spp. (P23-55, P24-83, P24-85, P24-88,
P24-192, and P24-193) were used in this study. The specific origin, sample metadata, and
isolation year for each coffee isolate are provided in Table S4. For the comparative analysis, data

from a previously characterized set of cacao-associated fungal isolates (including *Pestalotiopsis*

sp. CGH5 and *C. gloeosporioides* CGH17 and CGH49) were also included (Ahn et al., 2026).

All fungal isolates were routinely maintained on Potato Dextrose Agar (PDA) medium and

incubated in the dark at 24°C for 7 days to promote fungal growth.

2.3. Phylogenetic analysis

To elucidate the phylogenetic relationships among the fungal isolates, three gene regions were

analyzed: the internal transcribed spacer (ITS), glyceraldehyde-3-phosphate dehydrogenase

(GAPDH), and actin (ACT). Genomic DNA was extracted from 7-day-old isolates grown on

PDA using the Quick-DNA Fungal/Bacterial Miniprep kit (Zymo) according to the

manufacturer's instructions. The ITS, GAPDH, and ACT gene regions were amplified using the

primer pairs ITS-1 and ITS-4 (White, 1990), GDF and GDR (Templeton et al., 1992), and ACT-

512F and ACT-783R (Carbone and Kohn, 1999), respectively. PCR was performed using the

HiFi Phusion polymerase kit (NEB) in 50 µL reactions with 1 µL of genomic DNA as a template.

Thermal cycling conditions for ITS amplification consisted of an initial denaturation at 98°C for

3 min, followed by 35 cycles of 98°C for 30 s, 52°C for 30 s, and 72°C for 30 s, with a final

extension at 72°C for 10 min. For ACT amplification, an annealing temperature of 61°C and an

extension time of 15 s were used. For GAPDH amplification, an annealing temperature of 60°C

and an extension time of 15 s were used. PCR products were purified using a QIAQuick PCR kit

(Qiagen) and sequenced in both forward and reverse directions at Psomagen (Rockville, MD,

USA). Forward and reverse reads for each gene were assembled de novo to generate a consensus

sequence for each isolate in Geneious Prime (v2025.2.1), and low-quality bases at the sequence

ends were manually trimmed. These newly generated sequences were combined with previously

characterized sequences from the cacao-associated isolates and reference sequences from closely

related *Colletotrichum* species obtained from the NCBI GenBank database. Phylogenetic

analyses were used to contextualize lineage relatedness among isolates; species-level taxonomic

assignment was not the primary objective of this study.

Sequences for each gene were aligned using the MUltiple Sequence Comparison by Log-

Expectation (MUSCLE) algorithm implemented in Geneious Prime. Phylogenetic inference was

conducted for two datasets using MEGA 11: (1) the ITS region alone, and (2) a concatenated

dataset of the three gene regions (ITS, GAPDH, and ACT). For each analysis, the best-fit

nucleotide substitution model was selected in MEGA 11 based on the lowest Bayesian
Information Criterion (BIC) score. For the ITS-only analysis, a phylogenetic tree was
constructed using the Neighbor-Joining (NJ) method, with evolutionary distances computed
using the Kimura 2-parameter model with a gamma distribution (K2+G). For the concatenated
analysis, an NJ tree was also constructed, with evolutionary distances computed using the
Tamura-Nei model with a gamma distribution (TN+G). For both trees, nodal support was
assessed via bootstrap analysis with 1,000 replications, and bootstrap values of 75% or greater
are shown on the corresponding nodes.
165

2.4. Antifungal activity assay
The antifungal effects of the four compounds were tested via a modified agar plate assay. Each
compound (10 mg/mL stock) was spread evenly across PDA plates (15 µL, yielding 150 µg/plate)
using a sterile spreader. Control plates received 15 µL of pure ethanol. A 5-mm mycelial plug
from the growing edge of a 7-day-old culture of each fungal isolate was placed at the center of
the treated plates, which were then incubated at 24°C in the dark for 96 hours. Each isolate-
treatment combination had at least ten replicates. Each replicate consisted of an independently
prepared PDA plate inoculated with a single 5-mm plug (one plate = one biological replicate).
174

2.5. Image acquisition and morphological feature extraction
Following 96 hours of incubation, digital images of the fungal colonies were acquired using a
Nikon Z8 camera fitted with a Nikkor Z 20mm f/1.8 S lens (Nikon, Tokyo, Japan), mounted on a
Kaiser R2N CP stand (Kaiser Fototechnik, North White Plains, NY). Images were captured
under consistent lighting conditions provided by an RB 218N HF lighting unit equipped with two
18-W high-frequency fluorescent lamps (5400 K daylight-balanced, CRI > 90) positioned at
approximately 45° incident angles to ensure uniform illumination and minimize specular
reflection across the plate surface. Colony morphology was quantified using the SmartGrain
software (version 1.1) (Tanabata et al., 2012). The outline of each colony was manually traced
within the software by a single trained analyst to ensure consistency. Tracing followed a
standardized protocol (fixed magnification and boundary rules) to minimize user-to-user
variability, and image filenames were coded to blind the analyst to specific treatment identities
during the tracing process. From these outlines, SmartGrain automatically calculated the

following morphological traits: colony area (mm²), perimeter (mm), length (mm), width (mm),

LWR, circularity, and the asymmetry index defined as the distance between the intersection of

length and width and the center of gravity (IS & CG, in mm).

2.6. Statistical analysis and machine learning evaluation

General statistical analyses were performed using JMP Pro 17 (SAS Institute Inc., Cary, NC,

USA) (Klimberg, 2023). To assess differences in morphological traits between treatment groups,

a one-way Analysis of Variance (ANOVA) was performed, followed by a Tukey's Honestly

Significant Difference (HSD) post-hoc test for all pairwise comparisons. For multivariate

analyses, all quantitative morphological measurements were log-transformed to stabilize

variance and better approximate a normal distribution. PCA was performed on the standardized,

log-transformed data. Pearson's correlation analysis was used to assess linear relationships

between pairs of the transformed traits. PCA and PERMANOVA were first conducted on the

coffee-only dataset to evaluate chemical and isolate effects (Fig. 3), and then repeated on the

combined coffee + cacao dataset to evaluate crop-of-isolation label and treatment effects (Fig. 5).

Differences in multivariate morphological profiles between groups were tested using

PERMANOVA based on Euclidean distances of the log-transformed traits, with 9,999

permutations. Additionally, to dissect the interaction between isolate identity and chemical

treatment, a two-way factorial ANOVA was performed on log-transformed traits.

For the comparative analysis, the coffee isolate dataset was merged with the previously

published cacao-associated isolate dataset (Ahn et al., 2026), which was generated using the

identical experimental setup, media preparation protocol, and imaging equipment to minimize

batch effects. To predict whether a sample's crop-of-isolation label (cacao vs coffee, in this

dataset) could be predicted from its morphological response, a comprehensive suite of

classification models was evaluated using the "Model Screening" platform in JMP Pro 17.

Models were trained using JMP Pro 17 "Model Screening" with default settings for

benchmarking; detailed model settings are provided in Supplementary Data 1 (model reports).

Due to an unequal number of samples between the two panels, the dataset was balanced by

randomly subsampling the larger class to contain 180 samples from each crop-of-isolation, using

a fixed random seed (seed = 1), an approach chosen to preserve the original data distribution and

avoid introducing artifacts from oversampling techniques. The predictor variables included
treatment type and the seven morphological traits, which were standardized prior to model fitting.
Pathogen isolate identity was deliberately excluded as a predictor to ensure the model learned to
classify based on the interaction phenotype rather than memorizing individual isolates. Model
performance was assessed using a repeated K-Fold cross-validation strategy ($k = 5$, random seed
$= 1$). Because multiple samples per isolate were present, cross-validation performance is
primarily interpreted as within-panel predictability. To assess robustness to isolate-level
confounding, we additionally performed isolate-grouped cross-validation using
StratifiedGroupKFold with strain as the grouping variable ($k = 2$ after excluding CGH5, due to
the limited number of cacao strains), as reported in Supplementary Data 1. Primary analysis was
performed in JMP Pro 17; additional sensitivity analyses (grouped CV) were performed using
Python scripts, with outputs provided in Supplementary Data 1. To address potential taxonomic
and class-composition confounding, we conducted two additional sensitivity analyses: (1)
excluding the highly responsive cacao isolate (CGH5) to assess whether classification relied
solely on extreme phenotypes, and (2) removing the top predictor (circularity) to test feature
redundancy.
235
2.7. Hyperspectral imaging and spectral analysis

To investigate the early physiological responses induced by antifungal treatment, HSI was
conducted on 96-hour incubated colonies of *Colletotrichum* spp. Fungal cultures were initially
grown on potato dextrose agar (PDA) plates for 7 days at 24□°C in the dark. From these mature
cultures, 5□mm diameter plugs were collected using a sterile biopsy punch and transferred to
fresh PDA plates containing one of two phenolic-branched compounds, PhSOAM or BCBGAM,
selected as a representative subset based on their superior antifungal potency in the initial
screening, applied at three concentrations (150, 300, and 450□µg per plate). Control plates were
treated with ethanol only. Three fungal isolates were selected for HSI analysis based on their
distinct morphological responses observed in prior antifungal assays. The cacao-associated
isolate CGH5 (*Pestalotiopsis* sp.), which previously exhibited dramatic reductions in colony area
upon PhSOAM treatment, was included as a representative of a highly responsive genotype. Two
coffee-associated *Colletotrichum* isolates, P24-85 and P24-192, were selected to capture intra-
panel variation in chemical response. For each isolate and treatment condition, four biological

replicates were prepared, with each replicate consisting of a single, independently cultured
fungal plug. HSI was carried out using custom-built line-scan systems as previously described
(Kim et al., 2011). In this study, three kinds of imaging modalities were employed: fluorescence
imaging, VNIR reflectance imaging, and SWIR reflectance imaging. For VNIR reflectance
(401.5–1001 nm spectral range) and fluorescence imaging (438–718 nm spectral range), the
system was equipped with an electron-multiplying charge-coupled device (EMCCD) camera
(Luca DL 604M, Andor Technology, South Windsor, CT, USA) coupled to an imaging
spectrograph (Headwall Photonics Inc., Fitchburg, MA, USA). Reflectance illumination was
provided by four 150-W quartz tungsten-halogen lamps powered by a DC supply, while
fluorescence excitation was achieved using two UV-A LED light sources centered at 365 nm.
For SWIR reflectance imaging (842–2532 nm spectral range), a mercury cadmium telluride
(MCT) array detector coupled with a SWIR spectrograph (Micro-Hyperspec SWIR, Headwall
Photonics, Fitchburg, MA, USA) was used, illuminated by gold-coated halogen lamps to
maximize infrared output and minimize thermal drift. All plates were scanned in their entirety,
and spectral data were extracted from consistent, standardized regions of interest (ROIs) within
each fungal colony. Spectral signatures were calibrated using white reference and dark signal,
and F-value analysis was applied to identify wavelength regions showing statistically significant
variation as a function of treatment and strain identity through one-way ANOVA F-statistics.
Further details on system specifications and spectral calibration procedures are provided in Baek
et al. (2025). Because the SWIR region near ~1930 nm is moisture-sensitive, interpretation
focuses on treatment-associated spectral variation rather than biochemical assignment without
explicit water-effect modeling.

3 Results

3.1 Potent but distinct effects of amide derivatives PhSOAM and BCBGAM

The two ethylenediamine-carrying amide derivatives, PhSOAM and BCBGAM, were among the
most potent growth inhibitors tested. PhSOAM exhibited potent but selective antifungal activity,
reducing colony area in the most susceptible isolates, P24-192 and P24-85, by 19.4% ($p < 0.001$)
and 20.7% ($p < 0.001$), respectively (Fig. S1A). This reduction is visually illustrated for P24-192
(Fig. S1B, C). In contrast, BCBGAM induced slightly stronger inhibition in certain isolates (e.g.,
20.1% in P24-192) but was less effective against others (Fig. 1A). While both amides reduced

size metrics, their shape metrics (circularity, LWR, IS&CG) diverged, underscoring a size–shape

decoupling that motivates the downstream multivariate and predictive analyses. PhSOAM

triggered diverse morphological responses, in some cases inducing irregular expansion (e.g., a

decrease in circularity in isolate P24-193) (Fig. S1A). Conversely, BCBGAM promoted a more

compact, rounded morphology in three of the six isolates (Fig. 1A). The most striking response

was observed in isolate P24-192, where BCBGAM treatment induced a significant increase in

colony elongation (LWR), as illustrated in the representative images (Fig. 1).

Fig. 1. Representative morphological disruption by phenolic antifungal compounds, illustrated
using BCBGAM-treated isolates, including P24-192. (A) Box plots quantifying the morphological
response of six *Colletotrichum* sp. isolates to one representative phenolic antifungal (BCBGAM), which
exhibited isolate-specific effects. BCBGAM is featured because it elicited the most visually distinct
morphological changes in isolate P24-192. While PhSOAM showed stronger overall inhibition (Fig. S1),
the response to BCBGAM better illustrates the complex 'morphological fingerprint' central to this study,
highlighting our focus on quantifying reproducible stress-response fingerprints for downstream screening
and predictive modeling, rather than solely quantifying growth inhibition. Measured traits include colony

area (mm²), perimeter (mm), length (mm), width (mm), LWR, circularity, and an asymmetry distance (IS & CG, mm). Significant differences between a treated group and its corresponding control (Tukey's HSD; *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$) are marked with asterisks. (B-C) Representative images of isolate P24-192 after 96 hours on (B) a control plate and (C) a plate treated with BCBGAM. Sample size, n = 10 for all groups except P24-192 and P24-193 controls (n = 11). Scale bars = 1 cm.

3.2 Variable and opposing effects of fatty acid analogs

The two phenol-branched fatty acids, PhSOFA and BCBGFA, exhibited overall weaker antifungal activity than their corresponding amide derivatives, yet each triggered highly distinctive and isolate-specific morphological responses (Fig. 2; Fig. S2–S3).

PhSOFA demonstrated selective growth inhibition, significantly reducing colony size in two of the six isolates: P24-83 and P24-192 (Fig. S2A). For example, PhSOFA treatment reduced the colony area of isolate P24-83 by 18.8% ($p < 0.01$), a change illustrated in the representative images (Fig. S2B–C). In contrast, BCBGFA's effects were highly variable. It had no significant inhibitory effect on any isolate but paradoxically promoted the growth (assessed by area size measurement) of P24-85 by 16.5% ($p < 0.05$), as detailed in Fig. S3. This growth stimulation was consistent across replicates, suggesting a potential hormetic response or isolate-specific metabolic capability to metabolize the fatty acid substrate, rather than an experimental artifact. The remaining isolates were largely resistant to either compound. Morphological shape metrics highlighted further, though more sporadic, divergence between the two treatments. The effects on shape metrics were more sporadic: significant increases in colony circularity were rare, observed only for BCBGFA in P24-83 and PhSOFA in P24-193, while PhSOFA was the only compound to cause significant increases in both colony elongation (LWR) and asymmetry (IS & CG), effects that were limited to isolate P24-192 (Figs. S2 & S3).

When morphological data were pooled across all isolates (Fig. 2), a clear hierarchy of antifungal potency within this plate assay, based on the reduction in colony area, was observed. PhSOAM remained the most potent growth inhibitor. It was followed by PhSOFA. The other amide, BCBGAM, showed no significant difference from the control, while BCBGFA was the least effective, showing no significant inhibition. Despite variable effects on colony size, the compounds induced reproducible shifts in shape. PhSOFA uniquely elevated the LWR. A significant increase in colony circularity was observed for three compounds: PhSOFA,

BCBGAM, and BCBGFA. Furthermore, only PhSOFA caused a significant increase in the

asymmetry distance (IS & CG), pointing to non-lethal but geometry-altering cellular effects.

Together, these data show isolate- and structure-dependent effects across compounds (Fig. 2;

Figs. S2–S3). All isolate $\times$ compound $\times$ trait comparisons were evaluated and reported as *p*-

values in Supplementary Data 1, enabling exhaustive auditing of treatment effects. Two-way

ANOVA confirmed significant main effects of strain and chemical treatment and a significant

strain $\times$ chemical interaction for both log(area) and circularity (Supplementary Data 1).

338
Fig. 2. Comparative analysis of the antifungal effects of four phenolic-branched compounds on
coffee-associated *Colletotrichum* isolates. Box plots show the pooled morphological data from six
Colletotrichum isolates under four treatment conditions (PhSOAM, BCBGAM, PhSOFA, and BCBGFA)
and an untreated control. Measured traits include colony area (mm²), perimeter (mm), length (mm), width
(mm), LWR, circularity, and an asymmetry distance (IS & CG, mm). Different letters denote statistically
significant groupings ($p < 0.05$) determined by Tukey's HSD.

3.3 Multivariate analysis reveals a dominant treatment effect on fungal morphology

PCA of the morphological traits showed that the first two PCs captured 76.2% of the total

variance (Fig. S4). However, both chemical treatments and fungal isolates showed considerable

overlap in PCA space, indicating that linear ordination provided limited separation between

groups. This necessitated the use of supervised machine learning to test for nonlinear, diagnostic

patterns. Permutational multivariate analysis of variance (PERMANOVA) within the coffee

dataset indicated a small but significant treatment effect and a stronger isolate effect (Compound:

pseudo-F = 4.06, R^2 = 0.0518, p = 0.001; Strain: pseudo-F = 13.00, R^2 = 0.180, p < 0.0001). This

partial grouping in PCA, combined with substantial overlap, suggests that linear ordination

provides limited separation. Because PCA did not fully resolve isolates or crop-of-isolation

labels in the combined dataset, we next used supervised ML to test for nonlinear, diagnostic

patterns.

Further analyses were conducted to explore the relationships between the seven morphological

traits (Fig. S5). The PCA loading plot (Fig. S5A) and partial contributions plot (Fig. S5B)

confirmed that the primary axis of variation, PC1, was driven by the four main size metrics (area,

perimeter, length, and width). Shape-related traits had greater impacts on subsequent

components: LWR and IS & CG contributed most to PC2, while circularity was a major driver of

PC3. This clear distinction between size and shape was corroborated by hierarchical clustering,

which grouped the four size metrics into one tight cluster and the three shape metrics into

another (Fig. S5C). Pearson's correlation analysis further quantified these relationships, revealing

strong positive correlations among all size-related traits (r > 0.9 for most pairs, detailed in

Supplementary Data 1).

3.4 Phylogenetic analysis reveals coffee-associated isolates are polyphyletic and intermixed

with cacao-associated isolates

To determine the evolutionary relationships among the isolates, we first conducted a multi-locus

phylogenetic analysis using a concatenated dataset of the internal transcribed spacer (ITS),

glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and actin (ACT) gene regions (Fig. 3).

The analysis revealed the six coffee-associated isolates were polyphyletic, distributing across
several distinct clades rather than forming a monophyletic group.

Specifically, isolate P24-88 formed a sister group with *C. karstii* (79% bootstrap support), and
this clade in turn was a sister group to isolate P24-193. Isolate P23-55 was placed in a separate,
distant clade with *C. pseudoboninense*. The remaining three coffee isolates were located in
different parts of the tree: P24-192 clustered with *C. siamense* (99% bootstrap), P24-83 was a
sister taxon to *C. rhizophorae* and *C. fructicola*, and P24-85 was placed near *C. gloeosporioides*
and *C. kahawae*. This polyphyletic distribution indicates that the coffee isolates used in this
study represent multiple distinct lineages within the *Colletotrichum* genus.

An analysis based on the standard fungal barcode, the ITS region, was also performed to directly
compare the coffee isolates with cacao-associated isolates (Fig. S6). This analysis further
highlighted the complex relationships between isolates from the two panels. Notably, a single,
strongly supported clade (99% bootstrap) contained an admixture of isolates from different crop-
of-isolation labels. This inclusive clade grouped coffee-associated P24-192, P24-85, and P24-83
together with cacao isolates (CGH17, CGH53) and shade tree isolates (CGH34, CGH38).
Overall, based on the ITS region, some coffee isolates are more closely related to cacao-
associated isolates than to other coffee isolates, emphasizing mixed lineage structure within the
panel, such as P24-88 and P24-193, which formed a separate, well-supported clade (96%
bootstrap).

Fig. 3. Multi-locus phylogeny reveals the polyphyletic nature of coffee-associated *Colletotrichum* isolates

The phylogenetic tree was inferred by the Neighbor-Joining method using a concatenated alignment of the ITS, GAPDH, and ACT sequences. The evolutionary distances were computed using the Tamura-Nei (TN+G) model. Numbers at the nodes represent bootstrap support values based on 1,000 replications; only values of 75% or greater are shown. The six coffee-associated isolates (highlighted in blue) are shown to be polyphyletic, distributing across multiple distinct clades and indicating affiliations with different known *Colletotrichum* species.

3.5 Comparative multivariate analysis of coffee and cacao panels

To investigate whether the chemical compounds induced panel-associated morphological responses, the coffee isolate dataset was combined with a previously generated dataset from cacao-associated isolates [16], and a comparative PCA was performed on the combined, log-transformed data.

The first two principal components accounted for 78.6% of the total variance (PC1: 56.8%, PC2: 21.8%) (Fig. 4). When the data were colored by the sample crop-of-isolation label (Fig. 4A), no clear separation was evident. The coffee (black) and cacao (red) samples were heavily intermixed. Crucially, PERMANOVA revealed that the crop-of-isolation label explained only ~1.7% of the total variance (pseudo-F = 8.49, R^2 = 0.0174, p = 0.0017). This confirms that linear multivariate methods are insufficient to resolve these phenotypes, justifying the use of non-linear machine learning.

When visualized by treatment type, the analysis showed significant overlaps between the groups (Fig. 4B). The untreated control samples clustered closely with those treated with BCBGFA. A separate cluster was formed by samples treated with PhSOAM and PhSOFA, while the BCBGAM-treated samples were positioned between these two main groupings. This strong overlap indicates that the morphological signatures distinguishing the two panels, if they exist, are not linearly separable and thus are too subtle for PCA to resolve.

Fig. 4. Comparative PCA of coffee and cacao panels reveals treatment-driven separation but panel overlap. PCA of morphological traits from the combined coffee and cacao isolate datasets (log-transformed). (A) PCA scores plot with samples color-coded by panel: Cacao isolates (red) and Coffee isolates (black). Note the significant overlap between the two panels, indicating the first two principal

components cannot separate them. (B) The identical PCA scores plot with samples color-coded by chemical treatment: Control (black), PhSOAM (orange), PhSOFA (green), BCBGAM (blue), and BCBGFA (purple). The red squares indicate the mean of each panel in (A) and each chemical treatment group in (B). PERMANOVA indicated a weak crop-of-isolation label effect (Crop-of-isolation: pseudo-F = 8.49, $R^2 = 0.0174$, $p = 0.0017$) and a small but significant treatment effect (Compound: pseudo-F = 5.57, $R^2 = 0.0447$, $p < 0.001$), both explaining only a small proportion of total variance.

3.6 Machine learning models classify crop-of-isolation labels from chemical-stress morphology

Following the PCA, which could not clearly separate the two panels, a suite of ML models was employed to quantitatively test if a predictive morphological signature of crop-of-isolation labels (coffee vs cacao, in this dataset) existed within the combined dataset. The analysis revealed that several models could classify crop-of-isolation labels with high accuracy, supporting the existence of a learnable phenotype fingerprint associated with the crop-of-isolation label in this mixed-lineage panel based on the response to the chemical compounds (Fig. 5). A comprehensive screening of nine different algorithms identified the Neural Boosted model (an F1-score of 0.866, and a Matthews Correlation Coefficient (MCC) of 0.733) as the top performer, achieving a validation accuracy of 86.7%, which was followed closely by Nominal Logistic and Generalized Regression (Lasso) models (both ≈85%), as well as Support Vector Machines, Bootstrap Forest, and Fit Stepwise (all 84%) (Table S1).

To understand which traits were driving this successful classification, feature importance was analyzed using the top-performing Neural Boosted model (Table S2). The analysis revealed that circularity was the single most dominant predictor of the crop-of-isolation label, with a Total Effect score of 0.592, which was followed by the primary size-related traits of area (Total Effect = 0.405) and perimeter (Total Effect = 0.168). This indicates that the model primarily learned to distinguish the two crop-associated panels by identifying a unique 'shape signature', driven by how circular a colony becomes in response to chemical stress, complemented by information about the colony's overall size.

To further test the robustness of the classification, the analysis was repeated on the log-transformed dataset after removing circularity, the top predictor (full details in Supplementary Data 1). Even without this primary feature, the models maintained high predictive power. The

Neural Boosted model was again the top performer with a validation accuracy of 86.1% (a

negligible drop from 86.7% in the full model), followed by the Nominal Logistic (85%), Lasso

(85%), and Fit Stepwise (84%) models. Feature importance analysis for this new model revealed

a clear shift in predictive strategy; in the absence of circularity, the model pivoted to a 'size

signature', with area size (Total Effect = 0.825) and perimeter (Total Effect = 0.467) emerging as

the most dominant predictors (Supplementary Data 1). Note that because feature importance

scores are normalized to sum to 1, the exclusion of the dominant shape feature (circularity)

mathematically necessitates a rescaling of the remaining features, resulting in increased weights

for size metrics.

Furthermore, a sensitivity analysis excluding the distinct outlier isolate (CGH5) yielded a

validation accuracy of 83.5% (Supplementary Data 1), confirming that the predictive signal is

not driven solely by a single highly responsive genotype. Finally, to rigorously assess potential

data leakage, we performed an isolate-grouped cross-validation using StratifiedGroupKFold with

strain as the grouping variable ($k = 2$ after excluding CGH5). Under this stricter evaluation, mean

accuracy ranged from 79.3% to 82.9% with MCC values of 0.59–0.66 across the evaluated

classifiers (Supplementary Data 1). This grouped evaluation included 6 coffee strains and 2

cacao strains (CGH17 and CGH49), with folds defined at the strain level, confirming that a

detectable panel-level phenotype signal remains even when controlling for isolate-level data

leakage.

Fig. 5. Machine learning classification of crop-of-isolation label based on chemical-induced morphology. Decision thresholds plot showing the performance of nine ML models in classifying a pathogen's crop-of-isolation labels (cacao vs. coffee) based on its morphological fingerprint from chemical-induced morphology (A: Training set, B: Validation set). Blue dots represent isolates from the cacao panel, and red dots represent isolates from the coffee panel. The plot shows the aggregated results from the k-fold cross-validation, with the confusion matrix on the right summarizing classification

486 performance at a probability threshold of 0.5. 'Neural☐Boosted' denotes the JMP Pro☐17 boosted neural
487 network; default base learner with boosting enabled; repeated 5☐fold CV with fixed random seed.
488 Performance is reported for repeated 5-fold cross-validation on a class-balanced dataset; generalization to
489 unseen isolates was not directly tested here.
490
491

3.7 Dose-dependent hyperspectral fingerprints in CGH5 under PhSOAM treatment

To probe the spectral correlates of these morphological fingerprints, we first employed three

modalities of HSI: fluorescence HSI, visible and near-infrared (VNIR) reflectance imaging, and

short-wave infrared (SWIR) reflectance imaging. We began by testing the response of

Pestalotiopsis sp. CGH5, a cacao-derived isolate known to exhibit a dramatic 48.8% reduction in

fungal growth upon PhSOAM treatment [16]. HSI revealed distinct, dose-dependent responses of

Pestalotiopsis sp. CGH5 to PhSOAM treatment across all three spectral modalities (Fig. 6).

Fig. 6. Hyperspectral imaging of *Pestalotiopsis* sp. CGH5 treated with the biobased fungicide PhSOAM across three spectral modalities. (A) Fluorescence intensity profiles of CGH5 colonies treated with 0, 150, 300, and 450 µg of PhSOAM; color indicates dose (control=blue, 150 µg = orange, 300 µg = green, 450 µg= red). (B) F-value curve corresponding to the fluorescence data. (C) Reflectance intensity curves under broadband illumination. (D) F-value profile corresponding to the reflectance data.

506 (E) SWIR reflectance curves showing spectral shifts. (F) F-value curve corresponding to the SWIR data.
507 All data represent the mean of four biological replicates per treatment (n = 4).

3.8 Attenuated hyperspectral response in a tolerant coffee isolate

In contrast to the strong response of CGH5, the more tolerant coffee isolate P24-85 exhibited an

attenuated and nonlinear hyperspectral response to PhSOAM (Fig. 7). This included a non-

monotonic fluorescence signal and muted shifts in the SWIR moisture-sensitive region near

~1930 nm. These attenuated spectral changes indicate reduced treatment-associated spectral

variation in this isolate under the same imaging conditions.

Fig. 7. Hyperspectral imaging of *Colletotrichum* sp. P24-85 treated with the biobased fungicide
PhSOAM across three spectral modalities. (A) Fluorescence intensity profiles of P24-85 colonies
treated with 0 µg (control, blue), 150 µg (orange), 300 µg (green), and 450 µg (red) of PhSOAM. (B) F-
value curve corresponding to the fluorescence data. (C) Reflectance intensity curves under broadband
illumination. (D) F-value profile corresponding to the reflectance data. (E) SWIR reflectance curves. (F)
F-value curve corresponding to the SWIR data. All data represent the mean of four biological replicates
per treatment (n = 4).

3.9 Compound-specific hyperspectral fingerprints of *Colletotrichum* sp. P24-192

Different compounds also induced unique spectral fingerprints in *Colletotrichum* sp. P24-192, highlighting compound-specific spectral responses within the same isolate (Fig. S7). For example, in the SWIR modality, BCBGAM produced a focused diagnostic band near ≈1950 nm, whereas PhSOAM yielded a broader separation across multiple bands. These distinct spectral signatures support compound-specific spectral fingerprints under controlled in vitro conditions; no biochemical mechanism is inferred here.

4. Discussion

Here, we developed and evaluated a taxonomy-agnostic, multi-modal phenotyping workflow that links standardized chemical stress to reproducible morphology- and hyperspectral-based fingerprints. By integrating quantitative morphology, HSI, and supervised ML, we showed that chemical-stress phenotypes can support cross-validated prediction of crop-of-isolation labels (coffee vs cacao, in this dataset) under uniform in vitro conditions. Importantly, multi-locus phylogeny indicates mixed lineage structure among the coffee isolates, so we interpret predictive performance conservatively as evidence for a learnable phenotype signal associated with crop background within this panel, rather than proof of host adaptation or ecological memory. Together, these results position the workflow as a scalable, sensor-driven screening and decision-support approach for evaluating candidate antifungals and quantifying stress-response fingerprints.

The varied fungal responses underscored a clear structure-activity relationship dependent on both the headgroup and backbone of the four phenolic compounds. Consistent with previous reports, amide derivatives generally showed stronger antifungal activity than their fatty acid analogs. Specifically, PhSOAM, with its simple phenol and amide moiety, was the most potent growth inhibitor. This is consistent with contributions from increased polarity conferred by the amide group and steric accessibility of the unsubstituted phenol (Dembitsky, 2022; Du et al., 2015; Ōmura et al., 1974).

An unexpected observation was that BCBGFA appeared to promote growth in isolate P24☐85. While this may reflect experimental variability, the consistency of this stimulation across

replicates suggests a potential hormetic response (stimulation by low-dose stressors) or an
isolate-specific metabolic capability to utilize the phenolic fatty acid as a carbon source
(Middelhoven, 1993; Middelhoven et al., 1992). Because this study was not designed to quantify
compound catabolism, we treat this observation as a hypothesis-generating result.

The divergence in morphology across isolates and compounds supports the practical idea that
standardized stress assays can generate reproducible phenotype fingerprints suitable for
screening and prediction. We use the term "chemical priors" as a working hypothesis to describe
the possibility that prior exposure histories or persistent physiological states may shape stress-
response pathways that become legible under standardized probes. In this dataset, circularity
emerged as the most informative morphology-derived feature for predicting the crop-of-isolation
label, whereas elongation (LWR) was not consistently expressed across the coffee panel. Notably,
circularity shifts have also been observed under physical (UVC) stress in prior work (Baek et al.,
2025), suggesting that certain geometric outcomes may recur across disparate stressors. However,
establishing whether such recurrence reflects shared mechanisms or convergent macroscopic
endpoints will require targeted molecular and experimental validation.

While linear ordination (PCA) did not separate the combined datasets, supervised ML achieved
86.7% cross-validated accuracy for classifying crop-of-isolation labels (coffee vs cacao, in this
dataset), indicating that morphology contains a non-linearly separable predictive signal. Feature-
importance analysis consistently identified colony circularity as the dominant predictor, with size
metrics (area, perimeter) contributing additional information. When circularity was removed,
classification performance remained high (86.1%), indicating that the predictive signal is
distributed across multiple traits within this dataset. It is important to note that our primary cross-
validation strategy involved random splitting of replicate plates, such that replicates from the
same isolates could appear in both training and validation folds. Therefore, these accuracies
quantify within-panel predictability rather than generalization to unseen isolates. To directly
address isolate-level generalization, we additionally performed isolate-grouped cross-validation
using StratifiedGroupKFold with strain as the grouping variable ($k = 2$ after excluding CGH5,
due to the limited number of cacao strains; Supplementary Data 1), which provides a stricter

evaluation. These results support the utility of morphology-derived features as a compact,

interpretable representation for predictive screening, without requiring species-resolved inputs.

We interpret these phenotypes as plastic stress responses expressed over the 96-hour incubation

window rather than the rapid selection of resistant mutants. Because the incubation timeframe

and assay design do not directly test heritable change, we avoid mechanistic claims about

evolutionary memory and instead report reproducible phenotype shifts under standardized

conditions.

HSI complemented morphology by providing dose- and isolate-dependent spectral fingerprints

under controlled imaging conditions. Prior studies indicate that VNIR/SWIR reflectance and

fluorescence can differentiate physiological states in fungi, including changes associated with

hydration-related bands, pigmentation, and structural components (Kong et al., 2019; Neurauter

et al., 2024; Teena et al., 2014; Valenzuela-Gloria et al., 2021). In our study, the cacao-

associated isolate Pestalotiopsis sp. CGH5 showed strong spectral separation across modalities,

whereas the coffee isolate P24-85 exhibited a comparatively attenuated response under the same

treatment series. We interpret these patterns as treatment-associated spectral variation that

differentiates isolates, rather than as direct evidence for specific physiological mechanisms.

Notably, the SWIR region near ~1930 nm is moisture-sensitive, so biochemical attribution from

this band alone is not warranted without explicit water-effect modeling or robustness checks.

The contrast between PhSOAM and BCBGAM in the same isolate (Fig. S7; Table S3) supports

compound-specific spectral fingerprints under standardized conditions. These signatures

motivate future mechanistic studies, but here we restrict interpretation to reproducible, treatment-

associated spectral differences across modalities. Because the ~1930 nm region is moisture-

sensitive, future work should explicitly model or filter water effects (e.g., variable band

weighting or EMSC) before attempting biochemical assignment.

These parallels between xenobiotic chemical exposure and UVC irradiation suggest that HSI can

capture both compound-specific and isolate-specific physiological responses. Recurring spectral

patterns across distinct stressors warrant further study, but we do not attribute them to specific

ecological drivers or pathways based on the present dataset.

Crucially, the convergence of morphology and hyperspectral signatures suggests a robust, multi-

modal stress-response phenotype under standardized conditions. We present isolate-dependent

differences in spectral responsiveness as descriptive phenotyping outcomes that can inform

screening and prioritization, while mechanistic interpretation remains future work.

The polyphyletic distribution of the coffee isolates (Fig. 3; Fig. S6) provides important context:

crop-of-isolation labels are not equivalent to a single clade. While our primary headline accuracy

reflects within-panel separability (replicate splitting), the supplementary grouped cross-

validation ($k = 2$) and outlier-exclusion analysis (Supplementary Data 1) confirmed that the core

phenotypic signal remains detectable beyond specific isolates. Thus, we interpret these results as

a robust proof-of-concept for the workflow, identifying targets for future panel expansion.

Because crop-of-isolation labels are partially confounded with lineage structure in this dataset,

we do not attribute the predictive phenotype signal to host adaptation as a primary driver. Instead,

our results motivate future experimental designs that explicitly disentangle host background,

geography, and lineage—for example, by testing multiple isolates from the same species

sampled across crops and regions and evaluating isolate-blocked generalization. Within the

current panel, chemical stress elicits a rich, multi-layered phenotype that is measurable by

morphology and HSI and usable for predictive screening.

To synthesize these findings, Fig. 8 summarizes the end-to-end taxonomy-agnostic workflow

that links standardized perturbations to reproducible fingerprints and cross-validated label

prediction. In this study, the workflow links controlled chemical perturbations to reproducible

morphology and spectral fingerprints and enables prediction of crop-of-isolation labels within

the combined dataset. Establishing whether any component of this signal reflects heritable,

environmentally induced memory (e.g., epigenetic mechanisms) will require targeted

experimental designs and molecular assays beyond the scope of this work.

Fig. 8. Taxonomy-agnostic phenotyping workflow for predicting crop-of-isolation labels from chemical-stress responses. (A) Sample collection and labeling. Isolates are assigned sample-source labels (e.g., crop-of-isolation) and cultured under standardized conditions. (B) Standardized perturbation and multi-modal sensing. Colonies are challenged with uniform chemical stressors and quantified using macroscopic morphology and hyperspectral imaging to generate reproducible stress-response fingerprints. (C) Feature extraction and predictive modeling. Morphological traits and spectral summaries are used as model inputs; supervised machine learning is trained and evaluated via cross-validation to predict crop-of-isolation labels within the present dataset. (D) Screening and decision support output. The workflow produces interpretable outputs (e.g., feature importance, decision thresholds) to support high-throughput screening and sensor-driven decision support. This conceptual workflow summarizes the end-to-

end pipeline demonstrated here for coffee–cacao isolate panels under phenolic antifungal stress and is adaptable to other microorganisms and stressors.

Our study establishes a scalable, phenotype-driven platform for taxonomy-agnostic stress-response phenotyping and predictive modeling. The potential utility of this methodological advance is its application in high-throughput screening of eco-friendly fungicides and sensor-driven decision support for phenotype-based screening workflows. Building on this foundation, future work will prioritize: (1) validating these phenotype fingerprints in planta; (2) resolving molecular and genomic bases of stress responses; and (3) testing generality of the "chemical priors" hypothesis across diverse pathogen systems.

5. Conclusion

This study successfully developed and validated a high-throughput, taxonomy-agnostic phenotyping workflow that integrates quantitative morphology and hyperspectral imaging to characterize fungal stress responses. Unlike traditional multivariate statistics, which explained only ~1.7% of the phenotypic variance, our non-linear machine learning approach achieved 86.7% classification accuracy in predicting crop-of-isolation labels within a mixed-lineage panel. Crucially, the system demonstrated significant engineering robustness; excluding the primary shape feature (circularity) resulted in a negligible performance drop (to 86.1%), confirming that the predictive fingerprint is redundantly encoded across multiple traits. This framework offers a scalable, sensor-driven solution for rapid antifungal screening and biosecurity monitoring, decoupling phenotypic profiling from the bottlenecks of strict taxonomic identification.

CRediT authorship contribution statement

Insuck Baek: Formal analysis, Writing – review & editing. Seunghyun Lim: Formal analysis, Writing – review & editing. Amelia Lovelace: Methodology, Resources, Writing – review & editing. Sookyung Oh: Investigation, Writing – review & editing. Masoud Kazem-Rostami: Resources, Writing – review & editing. Helen Ngo: Resources, Writing – review & editing. Moon S. Kim: Funding acquisition, Writing – review & editing. Lyndel W. Meinhardt: Funding acquisition, Writing – review & editing. Lalit Kandpal: Investigation, Writing – review & editing. Minhyeok Cha: Methodology, Writing – review & editing. Chansong Hwang: Methodology, Investigation, Writing – review & editing. Richard D. Ashby: Resources, Writing – review & editing. Ezekiel Ahn: Conceptualization, Supervision, Investigation, Formal analysis, Funding acquisition, Writing – original draft, Writing – review & editing.

Declaration of competing interest

Acknowledgements

Mention of any trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U. S. Department of Agriculture. USDA is an equal opportunity provider and employer, and all agency services are available without discrimination. This work was supported by the U.S. Department of Agriculture, Agricultural Research Service, In-House Projects Nos. 8042-21220-258-000-D and 8072-41000-113-000-D, and by the National Institute of Food and Agriculture (NIFA) under Grant No. 2021-67021-34502. We thank Lisa Keith (Daniel K. Inouye U.S. Pacific Basin Agricultural Research Center, USDA-ARS, Hilo, Hawaii) for kindly providing the *Colletotrichum* isolates used in this study.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this work the authors used ChatGPT (OpenAI) in order to improve language and readability. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Data availability

All raw and processed data, including colony morphology measurements, per-wavelength mean spectra, and F-statistics for HSI, are provided in Supplementary Data 1. Multi-locus sequence data (ITS, GAPDH, ACT) generated for this study are openly available at figshare: https://figshare.com/articles/dataset/29924960.

References

Ahn, E., Kazem-Rostami, M., Park, S., Ashby, R.D., Ngo, H., Meinhardt, L.W., 2026. Quantitative morphological profiling and isolate-specific insensitivity of cacao pathogens to novel bio-based phenolic amides. J. Fungi (Basel) 12, 33.

Baek, I., Jang, J.H., Lim, S., Wang, Z., Cha, M., Magill, C., Kim, M.S., Meinhardt, L.W., Park, S., Ahn, E., 2025. Differential responses of Cacao pathogens *Colletotrichum gloeosporioides* and *Pestalotiopsis* sp. to UVB 305 nm and UVC 275 nm. Scientific Reports 15, 1–18.

Borges, D.F., Lopes, E.A., Fialho Moraes, A.R., Soares, M.S., Visôtto, L.E., Oliveira, C.R., Moreira Valente, V.M., 2018. Formulation of botanicals for the control of plant-pathogens: A review. Crop Protection 110, 135–140. https://doi.org/10.1016/j.cropro.2018.04.003

Cantrell, C.L., Dayan, F.E., Duke, S.O., 2012. Natural products as sources for new pesticides. J. Nat. Prod. 75, 1231–1242. https://doi.org/10.1021/np300024u

Carbone, I., Kohn, L.M., 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. Mycologia 91, 553–556.

Cassamo, C.T., Mangueze, A.V., Leitão, A.E., Pais, I.P., Moreira, R., Campa, C., Chiulele, R., Reis, F.O., Marques, I., Scotti-Campos, P., 2022. Shade and altitude implications on the physical and chemical attributes of green coffee beans from Gorongosa Mountain, Mozambique. Agronomy 12, 2540.

Cortaga, C.Q., Cordez, B.W.P., Dacones, L.S., Balendres, M.A.O., Dela Cueva, F.M., 2023. Mutations associated with fungicide resistance in *Colletotrichum* species: A review. Phytoparasitica 51, 569–592.

de Sousa, L.P., Guerreiro Filho, O., Costa Mondego, J.M., 2021. Differences between the leaf mycobiome of *Coffea arabica* and wild coffee species and their modulation by Caffeine/Chlorogenic acid content. Microorganisms 9, 2296.

Dembitsky, V.M., 2022. Microbiological aspects of unique, rare, and unusual fatty acids derived from natural amides and their pharmacological profile. Microbiology Research 13, 377–417.

Du, S., Lu, H., Yang, D., Li, H., Gu, X., Wan, C., Jia, C., Wang, M., Li, X., Qin, Z., 2015. Synthesis, antifungal activity and QSAR of some novel carboxylic acid amides. Molecules 20, 4071–4087.

Huang, K., Fan, X., Ashby, R., Ngo, H., 2021. Structure-activity relationship of antibacterial bio-based epoxy polymers made from phenolic branched fatty acids. Progress in Organic Coatings 155, 106228.

Kazem-Rostami, M., Ryu, V., Wagner, K., Jones, K., Mullen, C.A., Wyatt, V., Wu, C., Ashby, R., Fan, X., Ngo, H., 2023. Antibacterial agents from waste grease: arylation of brown grease fatty acids with beechwood creosote and derivatization. ACS Sustainable Chemistry & Engineering 11, 17760–17768.

Kim, M., Chao, K., Chan, D., Jun, W., Lefcourt, A., Delwiche, S., Kang, S., Lee, K., 2011. Line-scan hyperspectral imaging platform for agro-food safety and quality evaluation: System enhancement and characterization. Transactions of the ASABE 54, 703–711.

Klimberg, R., 2023. Fundamentals of predictive analytics with JMP. SAS institute.

Kong, W., Huang, W., Zhou, X., Mortimer, H., Ma, L., Tang, L., Li, C., 2019. Estimating leaf water content at the leaf scale in soybean inoculated with arbuscular mycorrhizal fungi from in situ spectral measurements. International Journal of Agricultural and Biological Engineering 12, 149–155.

Lahive, F., Hadley, P., Daymond, A.J., 2019. The physiological responses of cacao to the environment and the implications for climate change resilience. A review. Agronomy for Sustainable Development 39, 5.

Lew, H.N., Wagner, K., Fan, X., Nunez, A., Moreau, R.A., Haas, M.J., Yan, Z., 2018. Methods for preparing phenolic branched chain alkyl fatty acids or esters thereof and methods for killing microorganisms.

Middelhoven, W.J., 1993. Catabolism of benzene compounds by ascomycetous and basidiomycetous yeasts and yeastlike fungi: A literature review and an experimental approach. Antonie van Leeuwenhoek 63, 125–144.

Middelhoven, W.J., Coenen, A., Kraakman, B., Sollewijn Gelpke, M.D., 1992. Degradation of some phenols and hydroxybenzoates by the imperfect ascomycetous yeasts *Candida parapsilosis* and *Arxula adeninivorans*: evidence for an operative gentisate pathway. Antonie van Leeuwenhoek 62, 181–187.

Neurauter, M., Vinzelj, J.M., Strobl, S.F., Kappacher, C., Schlappack, T., Badzoka, J., Rainer, M., Huck, C.W., Podmirseg, S.M., 2024. Exploring near-infrared spectroscopy and hyperspectral imaging as novel characterization methods for anaerobic gut fungi. FEMS microbes 5, xtae025.

Nugroho, D., Basunanda, P., Mw, S., 2016. Physical bean quality of Arabica coffee (*Coffea arabica*) cultivated at high and medium altitude. Pelita Perkebunan 32, 151–161.

Ōmura, S., Katagiri, M., Awaya, J., Furukawa, T., Umezawa, I., Ōi, N., Mizoguchi, M., Aoki, B., Shindo, M., 1974. Relationship between the structures of fatty acid amide derivatives and their antimicrobial activities. Antimicrobial Agents and Chemotherapy 6, 207–215.

Rayhana, R., Ma, Z., Liu, Z., Xiao, G., Ruan, Y., Sangha, J.S., 2023. A review on plant disease detection using hyperspectral imaging. IEEE Transactions on AgriFood Electronics 1, 108–134.

Schnabel, G., Tan, Q., Schneider, V., Ishii, H., 2021. Inherent tolerance of *Colletotrichum gloeosporioides* to fludioxonil. Pesticide Biochemistry and Physiology 172, 104767.

Siripatrawan, U., Makino, Y., 2024. Hyperspectral imaging coupled with machine learning for classification of anthracnose infection on mango fruit. Spectrochim. Acta A Mol. Biomol. Spectrosc. 309, 123825.

Tanabata, T., Shibaya, T., Hori, K., Ebana, K., Yano, M., 2012. SmartGrain: high-throughput phenotyping software for measuring seed shape through image analysis. Plant Physiol. 160, 1871–1880.

Tang, Y., Yang, J., Zhuang, J., Hou, C., Miao, A., Ren, J., Huang, H., Tan, Z., Paliwal, J., 2023. Early detection of citrus anthracnose caused by *Colletotrichum gloeosporioides* using hyperspectral imaging. Comput. Electron. Agric. 214, 108348.

Teena, M., Manickavasagan, A., Ravikanth, L., Jayas, D., 2014. Near infrared (NIR)
hyperspectral imaging to classify fungal infected date fruits. J. Stored Prod. Res. 59,
306–313.
Templeton, M.D., Rikkerink, E.H., Solon, S.L., Crowhurst, R.N., 1992. Cloning and molecular
characterization of the glyceraldehyde-3-phosphate dehydrogenase-encoding gene and
cDNA from the plant pathogenic fungus *Glomerella cingulata*. Gene 122, 225–230.
Thind, T.S., 2017. Role of Fungicides in Crop Health Management: Prospects and Challenges,
in: Satyanarayana, T., Deshmukh, S.K., Johri, B.N. (Eds.), Developments in Fungal
Biology and Applied Mycology. Springer Singapore, Singapore, 433–447.
https://doi.org/10.1007/978-981-10-4768-8_22
Tomaszewski, M., Nalepa, J., Moliszewska, E., Ruszczak, B., Smykała, K., 2023. Early
detection of Solanum lycopersicum diseases from temporally-aggregated hyperspectral
measurements using machine learning. Scientific Reports 13, 7671.
Tsalidis, G.A., 2022. Human health and ecosystem quality benefits with life cycle assessment
due to fungicides elimination in agriculture. Sustainability 14.
https://doi.org/10.3390/su14020846
Valenzuela-Gloria, M., Balagurusamy, N., Chávez-González, M., Aguilar, O., Hernández-
Almanza, A., Aguilar, C., 2021. Molecular characterization of fungal pigments. J Fungi
(Basel) 7.
Velásquez, C., Aleixos, N., Gomez-Sanchis, J., Cubero, S., Prieto, F., Blasco, J., 2024.
Enhancing anthracnose detection in mango at early stages using hyperspectral imaging
and machine learning. Postharvest Biol. Technol. 209, 112732.
White, T.J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and
applications 18, 315-322.
Winters, N.P., Wafula, E.K., Knollenberg, B.J., Hämälä, T., Timilsena, P.R., Perryman, M.,
Zhang, D., Sheaffer, L.L., Praul, C.A., Ralph, P.E., Prewitt, S., Leandro-Muñoz, M.E.,
Delgadillo-Duran, D.A., Altman, N.S., Tiffin, P., Maximova, S.N., dePamphilis, C.W.,
Marden, J.H., Guiltinan, M.J., 2024. A combination of conserved and diverged
responses underlies *Theobroma cacao*'s defense response to *Phytophthora palmivora*.
BMC Biology 22, 38. https://doi.org/10.1186/s12915-024-01831-2
836