Dormant viral pathways underlie space-induced neural senescence

Aline M.A. Martins[1,2†], Hideyuki Nakashima[3†], Angela Macia[4†], Taichi Umeyama[3], Satoshi Miyashita[5], Livia Luz de Souza Nascimento[1], Luisa Bulcao Vieira Coelho[1], Pinar Mesci[4], Erik Viirre[6], Nicole Suarez[4], Aurian Saleh[4], Alex Savchenko[4], Beatriz Freitas[4], Aline L. Leite[7], Diego M. Assis[7], Masataka Ishizu[3], Tomoo Owa[5], Mikio Hoshino[5], John Yates III[2], Kinichi Nakashima[3*] and Alysson R. Muotri[1,4,8*]

[1]Integrated Space Stem Cell Orbital Research Center, University of California San Diego, La Jolla, CA, USA.

[2]The Scripps Research Institute, La Jolla, CA, USA

[3]Department of Stem Cell Biology and Medicine, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka, Fukuoka 812-8582, Japan.

[4]Department of Pediatrics and Cellular & Molecular Medicine, School of Medicine, University of California San Diego, La Jolla, CA, USA.

[5]Department of Biochemistry and Cellular Biology National Institute of Neuroscience, National Center of Neurology and Psychiatry (NCNP) Tokyo 187-8502 Japan

[6]Department of Neurosciences, School of Medicine, University of California San Diego, La Jolla, CA, USA.

[7]Bruker Daltonics Corporation, Billerica, MA, USA.

[8]Kavli Institute for Brain and Mind, Center for Academic Research and Training in Anthropogeny, Archealization Center, La Jolla, CA, USA.

†These authors contributed equally to this work

*Correspondence: muotri@ucsd.edu or nakashima.kinichi.718@m.kyushu-u.ac.jp

SUMMARY

Long-duration spaceflight is associated with neurological symptoms in astronauts, yet the
underlying molecular mechanisms remain unclear. Using human brain organoids cultured
aboard the International Space Station, we analyzed three independent spaceflights to
demonstrate that exposure to the space environment triggers Space-Induced Neural
Senescence (SINS), characterized by chromatin remodeling, mitochondrial dysfunction, and
activation of viral-like transcriptional programs in the absence of infection. Multi-omics
analyses identified upregulation of endogenous LINE-1 (L1) retroelements, whose activity was
markedly enhanced in organoids lacking MECP2, a known L1 repressor implicated in Rett
syndrome. The resulting accumulation of cytoplasmic L1 DNA elicited an IL-6–mediated
inflammatory and neurotoxic response, which was reversed by reverse transcriptase inhibitors
(RTi) such as lamivudine or stavudine. Parallel preclinical experiments in *Mecp2*-deficient
mice confirmed that RTi treatment restored neuronal morphology, synaptogenesis, function,
cognition, and survival. These findings reveal that the space environment reactivates dormant
genomic retroelements, providing an unexpected mechanistic insight into astronaut
neurobiology and identifying a potential therapeutic strategy for both space-induced and
terrestrial neurological conditions. Our pioneering study demonstrates the value of space-
enabling research in accelerating drug discovery and disease treatment on Earth.

KEYWORDS

Brain organoids, space environment, proteomics, transcriptomics, L1, retroelements

INTRODUCTION

In recent years, human space exploration efforts have increased substantially, including the Artemis
mission to land on the Moon in preparation for missions to Mars, as well as numerous private companies
preparing to build the next generation of space stations in Low-Earth Orbit (LEO) to replace the
International Space Station (ISS) [1]. As humans push farther into space, they face hazards like cosmic
radiation and prolonged microgravity, which can cause a spectrum of physiological issues, including
DNA damage, musculoskeletal degeneration, and neurological problems [2]. Space exposure is known
to affect the astronaut's brain, as evidenced by changes such as increased ventricular volume [3,4],
altered global white matter volume [5,6], and persistent cognitive decline post-flight [2-4,7-9]. However, the
exact mechanisms remain unknown, largely due to limitations of current noninvasive brain imaging
technologies. Therefore, successful long-duration exploration requires a comprehensive understanding
of spaceflight's impact to develop effective countermeasures [10]. The SpaceX Inspiration4 mission
highlighted the value of biomedical research in private space missions by collecting extensive health
data, including multi-omic analyses. This contributes to the expanding biomedical database [11] and
supports precision astronaut medicine. This "Second Space Age," characterized by commercial
participation [12], provides diverse participant data from non-professional astronauts. Recent studies
utilize multi-omics and systems biology to characterize the effects of spaceflight. Authors aggregated
transcriptomic, proteomic, metabolomic, and epigenetic profiles, revealing alterations converging on
mitochondrial and metabolic processes, innate immunity, chronic inflammation, cell cycle regulation,
circadian rhythm, and olfactory functions [1-13]. Confirmed by NASA's Twin Study, mitochondrial
dysregulation emerged as a consistent phenotype across multiple biological layers [14-16].
67

Although microgravity is the most recognized abiotic event in the space environment, radiation is among
the foremost health risks facing astronauts during space missions, particularly long-duration exploration
missions [14]. The chronic low-dose radiation exposure increases their risk of developing cancer, induces
DNA damage, and triggers oxidative stress [14,17]. Radiation also contributes to neurological disruptions,
such as neurovascular damage and neurodegeneration, in simulated models [18,19], potentially
accelerating aging [18].
74
75 To understand how the space environment affects the brain, Masarapu and colleagues used single-
76 nucleus multi-omics and spatial transcriptomics on mouse brains flown on the ISS for around 40 days
77 [20]. This study revealed neuroinflammation, oxidative stress, and impaired synaptic signaling, consistent
78 with findings in neurodegenerative disorders and radiation studies. However, similar studies cannot be
79 conducted in humans; thus, the translatability of these observations is unclear.
80
Due to the inability to study astronaut brains at the cellular level, we used human brain organoids as a
model system. These organoids mimic the functional and cellular features of the human brain [21],
providing a unique experimental model for investigating the consequences of space exposure [22]. We
developed an automated device to sustain organoid culture on the ISS and return live samples, with an
Earth replica serving as ground control. Next, both proteomics and transcriptomics analyses were
performed on the brain organoid samples returned from the ISS. One of the most representative
molecular pathways upregulated in the flight samples relative to the ground control was associated with
viral infection. In the absence of any additional viral infections, we hypothesized that this activation
resulted from an increase in endogenous Long Interspersed Nuclear Elements-1 (LINE-1, or L1)
retroelements. L1 elements are a class of retrotransposons comprising approximately 20% of the
mouse and human genomes [23,24]. Several neurological and psychiatric diseases exhibited variation in
L1 expression, suggesting L1s could be detrimental when de-regulated [25-31], especially in age-related
neurodegenerative diseases [32,33]. The upregulation of this pathway is linked to a series of histone
modifications, relaxing the chromatin, and alterations in MECP2, a well-known L1 repressor [34-38].
95
To translate our observations in space to a clinically relevant model on Earth, we studied L1
upregulation in Rett syndrome (RTT, OMIM#312750), caused by MECP2 mutations [39]. We show that
increased L1 expression in MECP2-mutant human astrocytes triggers an Interleukin 6 (IL6)-mediated
inflammatory neurotoxic response. Blocking L1 reverse transcription using reverse transcriptase
inhibitors (RTi) ameliorates neurotoxicity, improving cognition and survival. Our results contribute to
understanding the molecular mechanisms underlying the neurological consequences of spaceflight and
offer potential neuroprotective countermeasures for astronauts and therapeutic opportunities for
neurological conditions on Earth.
104
105

RESULTS

An autonomous platform for human brain organoids at ISS

To investigate how the space environment affects human brain organoids, we used a multi-modal
approach, combining proteomics (mass spectrometry-based) and transcriptomics in two different levels
of information (bulk and single-nuclei analysis). We used human brain organoids generated using a
semi-guided protocol, a robust approach that yields consistent results across different iPSC lines [1,2].
Derivation of brain organoids using this approach yields clear neural rosettes with a spatially organized
arrangement of proliferative cells, intermediate progenitor cells, glial cells, and lower- and upper-layer
cortical neurons. Based on cell population annotation, 1-month organoids consisted of approximately
70% progenitor and radial glial cells (expressing markers such as SOX2) that would mature into
functional neurons (NeuN2-positive) and astrocytes (GFAP-positive). Over the course of organoid
maturation, neurons exhibit a characteristic pyramidal morphology, and mature synaptic ultrastructure

develops, enabling the formation of functional populations of inhibitory interneurons, which in turn
facilitate the establishment of complex neural circuitry. One-month-old human brain organoids from 3
iPSC lines derived from healthy individuals with different genetic backgrounds, plus a cell line with
mutated *MECP2* (MECP2-KO), were loaded into a 9U CubeLab™ hardware (Space Tango) on three
independent Commercial Resupply Services (CRS) using SpaceX (SpX) rockets missions (CRS-
SpX18, CRS-SpX30; CRS-SpX31), each lasting around 30 days at the ISS (**Fig. 1**), returning with two
months old. Space Tango's flight and ground hardware supported 12 wells, each equipped with
impellers and automated operations (fluid conveyance, fluid routing, thermal control, environmental
monitoring, imaging, and sampling), containing approximately 30 organoids (**Fig. 1** and **Fig. S1A**). The
organoids were monitored in real time in each well in both environments, the ground and the ISS units
(**Fig. S1A**). CubeLab environmental telemetry (CO_2 levels, culture medium temperature, and radiation)
was recorded during operations (**Fig. S1B and E,** and **Table S1**). As they return from the ISS at 2
months old, brain organoids are self-organized in a structure containing five major cell types –
progenitors, intermediate progenitors, glial cells, glutamatergic neurons, and GABAergic neurons, as
previously described (**Fig. S1C**). We also visually tracked the organoids' viability and overall health in
both environments, measuring their diameter during the experimental weeks (**Fig. S1D**). ISS and
ground samples were run synchronously, and scheduled operations were initiated simultaneously. Upon
return to Earth, the samples were processed for proteomic and transcriptomic analyses.

Space-induced neural senescence - SINS

The effects of the space environment on brain organoids revealed significant alterations in both
proteomic and transcriptomic profiles. A summary of all experiments performed in this study, including
flight missions and details of their statistical analyses, is provided in **Table S2**. In the proteomics
analysis, we identified an average of 6000 proteins per run and 4059 differentially expressed proteins
between brain organoids on Earth (ground control) and those exposed to the space environment at the
ISS ($p \leq 0.01$; fold change 1.2) (**Table S3**). Proteomic analysis of samples at the ISS-exposed to space
demonstrated a distinct set of upregulated proteins (**Fig. 2A**). Proteins such as tenascin (TNC);
neurofilament light polypeptide (NEFL); mitochondrial import inner membrane translocase (TIMM8A),
and subunit of cytochrome c oxidase (HIGD1A), show significant fold changes and p-values, implying
alterations on the metabolism of the brain organoids. Reactome pathway analysis identified key affected
pathways, including RNA metabolism, intracellular protein transport, and viral infection (**Fig. 2B**).

To evaluate the oxidation/phosphorylation (OxPHOS) state of the experiment, we performed an
enrichment analysis and observed the downregulation of the mitochondrial-related proteins in the ISS
(**Fig. 2C**). We found 629 downregulated proteins are significantly enriched in functions and pathways
crucial for cellular energy production (respiration) and the synthesis of mitochondrial proteins,
demonstrating that space environment causes a deep alteration in the energetic machinery in the brain
organoids. The most significantly affected molecular functions are related to the electron transport chain
and oxidoreductase activity. The most enriched biological pathway is associated with the citric acid
(TCA) cycle and respiratory electron transport, indicating that multiple aspects of mitochondrial protein
synthesis, including translation initiation, elongation, and termination, are significantly downregulated,
suggesting reduced capacity for building and repairing mitochondrial components (**Fig. 2C**).

We also observed alterations in epigenetic marks. Histones play a crucial role in regulating gene
expression by influencing chromatin structure and accessibility. Reduced histone levels can alter
chromatin dynamics, making DNA more accessible to transcription factors and enzymes involved in
DNA repair and replication [3]. Such alterations might reflect cellular responses to stress conditions and
cellular senescence[4]. Histones are globally downregulated in the ISS when compared to ground control
brain organoids (**Fig. 2D**). Specifically, the histone acetyltransferases (HAT1 - histone acetyltransferase
type B; KAT8 - histone acetyltransferase KAT8; KAT7 - histone acetyltransferase KAT7), usually linked
to open chromatin, were upregulated at the ISS (**Fig. S2A**, upper panel). At the same time, we observed
downregulation of deacetylases at the ISS, indicating a withdrawal of acetyl groups (**Fig. S2A,** bottom
panel). Together, these observations suggest that cells in brain organoids exposed to the ISS undergo
chromatin remodeling, thereby facilitating genomic accessibility and overall transcription.

To further complement our proteomics analysis, we evaluated the secretome panel on a subsequent
flight (Spx31), aiming to understand how the space environment altered the pattern of secreted effectors
(**Fig. 2E**). The conditioned media of a batch of control brain organoids were analyzed using Olink
Reveal® panel technology, a Proximity Extension Assay technology (PEA). The readouts were
compared with an aliquot of unconditioned medium (blank) used to grow the organoids in the ISS, and
this negative control was compared with each of the tested conditioned media. The data analysis
revealed that the secretome of brain organoids exposed to a space environment exhibited a coordinated
response of protein effectors in the ISS relative to ground control. The secretome panel highlight: *i)*
modulation of neural development activity: the upregulation of secreted Glial Cell Line-Derived
Neurotrophic Factor (GDNF) and sustained expression of Fibroblast Growth Factor 19 (FGF19)
suggests an intrinsic adaptive mechanism where supportive glial cells respond to the unique demands
of the space environment by enhancing trophic support and developmental cues, against space-
induced cellular insults [5]; *ii)* alteration of proliferative state, impacting epigenetic regulation:
downregulation in the ISS brain organoids of DPY30 (Dpy30 histone methyltransferase complex
subunit), SMARCA4 (SWI/SNF related, matrix associated, actin dependent regulator of chromatin,
subfamily A, member 4), TOP2B (DNA Topoisomerase II Beta) and MKI67 (marker of proliferation Ki-
67) emphasizes changes in chromatin remodeling and gene expression regulation in the space
environment, and a coordination of a robust epigenetic response[6]; *iii)* alterations in cytoskeletal
dynamics and trafficking: together, the downregulation compared to ground samples of BAIAP2 (brain-
specific angiogenesis inhibitor 1-associated protein 2), CAMSAP1 (calmodulin regulated spectrin
associated protein 1) and ABL1 (ABL proto-oncogene 1, non-receptor tyrosine kinase) indicates a
significant disruption or alteration in actin cytoskeleton organization and dynamics, that could lead to
changes in cell shape, migration, and the formation of cellular protrusions essential for neuronal
connectivity[7,8].

The bulk transcriptomic data also showed significant differential gene expression between the ground
control and organoids at the ISS. We identified 150 differentially expressed genes (DEGs) with a 1.5-
fold change and p-values <0.05. (**Table S4**). Most DEGs are involved in viral RNA translation, followed
by selenocysteine synthesis and peptide chain elongation (**Fig. 2F**). Consistent with the proteomic
pathway analyses, the transcriptional Reactome pathway analyses also uncovered activation of viral
response pathways (**Fig. 2G**). Moreover, WikiPathway analyses revealed significant alterations in
MECP2-related gene expression under space environment conditions (**Fig. 2H**). To determine whether
the viral pathway could also be detected in the blood of astronauts, we compared our DEGs with those
from the NASA Twins Project using Ingenuity Pathway Analysis (IPA). As expected, pathways related
to the hematopoietic system (e.g., T-cell activation) are present only in the blood astronaut samples
(**Fig. S2B**). The top hits across biological and disease pathways include several viral-related pathways,
such as viral infection, retrovirus replication, and HIV infection, among others (**Fig. S2B**).

To determine which cell-type-specific responses were affected by the space environment at the ISS,
we performed a single-nucleus transcriptomics experiment (snRNA-seq). We identify 10 cell
populations in the brain organoids: lower-layer neurons, upper-layer neurons, radial glial cells,
intermediate progenitors, glial cells, astrocytes, dividing neural progenitors, early neurons, pan-cortical
neurons, and oligodendrocyte progenitor cells (**Fig. 2I** and **Fig. S2C**). Changes in cell populations were
observed when we compared ground control and brain organoids exposed to the space environment
(**Fig. 2J**). Mainly, we observed a higher number of active astrocytes (glial fibrillary acidic protein (GFAP)-
expressing cells and transforming growth factor beta 2 (TGFβ2)-expressing cells) in organoids exposed
to the space environment (**Figs. 2K-L**). We generate a heatmap of differentially expressed key marker
genes across cell types. We observed distinct expression patterns across cell clusters, with GFAP
expression with a higher intensity in astrocytes; forkhead-box G1 (FOXG1), B-cell lymphoma/leukemia
11b (BCL11b) and (glutamic acid decarboxylase) GAD1, with high intensity in upper- and lower-layer
neurons and intermediate progenitors (**Fig. S2C**).

We evaluated human brain organoids from another independent spaceflight (SpX30) to validate our
previous findings using bulk transcriptomics. The subsequent spaceflight also revealed similar signaling
pathways altered in the ISS, complementing the original observations from the SpX18 flight (**Fig. 2M**).
The analysis of the gene expression differences between brain organoids in the ground experiments

vs. brain organoids exposed to the ISS revealed: *i)* downregulation of non-coding RNAs: the
downregulation of LINC RNAs (Long Interspersed Non-Coding RNAs), such as LINC01873,
LINC02654, LINC01911. Interestingly, the primary function of these LINC RNAs is to suppress the
activity of retrotransposons, such as L1 elements; *ii)* cellular stress: the decreased expression of
CAPN13 (calpain 13), PNPLA1 (patatin-like phospholipase 1), and SLC22A14 (transmembrane solute
carrier family 22 member 14) indicates impairment of neurogenesis and reduction in cell proliferation
and long-term survival of neurons; suppressed oxidative phosphorylation, specially through
mitochondria deficiency in ATP production, leading to impaired in fatty acid beta-oxidation (FAO) and
reduction in acyl-carnitines and TCA cycle; *iii)* altered neurodevelopment and impaired plasticity: the
coordinated downregulation of key developmental genes – NPPB (natriuretic peptide B), MYO7B
(myosin VIIB), and CDH13-AS2 (long non-coding RNA (lncRNA) that plays a role in regulating the
expression of the CDH13 gene, also known as T-cadherin) indicates that the space environment may
hinder the brain's developmental genesis **(Fig. 2M).**

Collectively, all the changes in transcriptomics and proteomics described above represent a novel
phenomenon not observed on Earth. Thus, we coined the term "Space-Induced Neural Senescence"
(SINS) to distinguish it from the linear aging process observed on the ground. These findings provide
insights into the neurological effects of spaceflight, advancing our understanding of the molecular
mechanisms underlying the negative impact on the human brain caused by the space environment.

Space environment activates retroelement transcription in mice

To independently validate our main observations in a different model system, we analyzed publicly
available RNA-seq data from the retinas of mice exposed to spaceflight[10]. Although there was no overall
significant change in global gene expression between the ISS samples and ground controls **(Fig. 3A-**
C), gene set enrichment analysis (GSEA) revealed enrichment of genes associated with
retrotransposon silencing, transposition, IL6-type cytokine receptor interaction, inflammasome
activation, and phagocytosis in the ISS samples. In addition, mitochondrial dysfunction in ISS was
suggested by downregulation of the aerobic electron transport chain and mitochondrial biogenesis, and
by upregulation of cytochrome c release **(Fig. 3D, bottom panel)**, corroborating our previous
mitochondrial findings in the human model. Based on these observations, we inferred that
retrotransposons are upregulated in the mouse retina during spaceflight. Supporting this, we found an
increased expression of class I retrotransposable elements: LINE, SINE, and LTR **(Fig. S3A)**. Notably,
multiple L1 subfamilies were significantly upregulated in the ISS samples **(Fig. S3B)**. Retrotransposable
elements are also upregulated following exposure to simulated 5-ion galactic cosmic radiation (GCR),
which models the effect of cosmic rays **(Fig. S3C-D)**. Furthermore, both evolutionary young and ancient
L1 subfamilies have been previously shown to be upregulated in the mouse lung under combined GCR
and microgravity exposure[11]. Together, these findings suggest that environmental factors specific to
space, including microgravity and cosmic radiation, can trigger the activation of endogenous L1
retroelements in both mouse and human brains.

MECP2-KO brain organoids exacerbate the viral pathway at the ISS

In the absence of external viral infections and based on our previous results, we hypothesized that
activation of endogenous L1 retroelements led to the upregulation of the viral pathway observed upon
space exposure in human brain organoids. L1s can reverse-transcribe their RNA sequences and
accumulate in the cytoplasm of astrocytes, thereby triggering an inflammatory response [12,13]. To test
this hypothesis, we used brain organoids lacking functional MeCP2 (MECP2-KO[9]), a known L1
repressor[14-16]. Moreover, our transcriptomic results indicate alterations in MECP2-related gene
expression under spaceflight conditions **(Fig. 2H)**. MECP2-KO and control brain organoids were
exposed to the space environment and subjected to bulk proteomics analysis **(Fig. 4A** and **Table S3)**.
The Reactome analysis showed that the most upregulated proteins are related to cellular stress
responses, including viral pathways **(Fig. 4B** and **Fig. S4)**. The viral pathway was highly enriched (-
$\log_{10}(p)$ = -9.975, ratio = 0.044328553), comprising 34 proteins **(Table S5)**. The viral pathway was
further increased in the absence of the MECP2 gene **(Fig. 4B** and **Fig. S4A)**, confirming MeCP2's role
in silencing this pathway. We evaluate the bulk transcriptome of the same MeCP2-KO model brain

organoids in a subsequent flight (SpX30). The heatmap shows that the genes listed are overall more
highly expressed in the MeCP2-KO ISS samples and show significantly lower expression in the healthy
brain organoids samples (ISS control) **(Fig. 4C)**. A functional analysis of the most highly upregulated
genes reveals a pro-inflammatory and immune-active state in the MeCP2-KO ISS samples,
corroborating the first spaceflight (SpX18). Genes related to *i)* inflammation: SELE (E-selectin), OSMR
(oncostatin M Receptor); *ii)* immune activation: CCR7 (C-C motif chemokine receptor 7); *iii)* tissue stress
and remodeling: TGFBI; and *iv)* neurological stress response: NPY (neuropeptide Y) are highlighted in
the ISS condition. To reinforce our findings, we also compared, from the same flight, the ground MeCP2-
KO brain organoids *vs.* ISS MeCP2-KO brain organoids using a bulk transcriptomics approach **(Fig.
S4B)**. In this evaluation, we highlight the following findings: *i)* upregulation of excitatory and inhibitory
circuits in the brain: upregulation of NPAS4 (neuronal PAS domain-containing protein 4) in the ISS
samples strongly suggests that the combined stress of the MeCP2-deficient background and spaceflight
triggers a state of significant neurological distress or aberrant neuronal activity. This highlights the
brain's vulnerability in a system where a key epigenetic safeguard (MeCP2) is already absent; *ii)*
activation of inflammatory and tissue remodeling: LRRC15 (leucine rich repeat containing 15) is
implicated in tissue remodeling and its upregulation is a response to stress or injury and has been
described with an essential role in response to viral infections, including as a key determinant of host's
susceptibility/resistance to SARS-CoV-2[17], IL36B (interleukin 36 beta): a pro-inflammatory cytokine
belonging to the IL-1 family, indicating an active inflammatory response, when compared to the ground
samples.

Translating to Earth: accumulated L1 transcripts in astrocytes promote neurotoxicity

To translate the observations in space to Earth, we decided to conduct a series of groundwork studies
in a clinically relevant human condition. A lack of functional MECP2 in the nervous system can cause
Rett syndrome (RTT) and other neurodevelopmental conditions[19]. Accumulating L1-derived single-
stranded DNA (ssDNA) in human astrocytes can trigger a neurotoxic inflammatory response in another
neurodevelopmental disorder[12]. Using the MECP2-KO and control lines, we checked for L1 expression
and found that endogenous L1 transcripts and L1-derived ssDNA are enriched in iPSC-derived
astrocytes lacking MECP2 **(Fig. 5A, B** and **Fig. S5A)**. Cytokines are important signaling molecules
produced by glial cells and are responsible for cell maturation and inflammation during
neurodevelopment[18]. We found several cytokines that were upregulated in astrocytes lacking MECP2
(Fig. S5B), including an increase in secreted IL6 **(Fig. 5C** and **Figs. S5C-E)**. High levels of secreted
astrocytic IL6 have recently been implicated in RTT, affecting synaptogenesis in a mouse model [21]. We
validated these observations in our human system, using astrocyte-conditioned media (ACM) and by
adding recombinant human IL6 (rhIL-6) to the culture media of iPSC-derived control neurons at the
same concentration found in MECP2-KO astrocytes (2 ng/mL) for two weeks **(Fig. 5D** and **Figs. S5F,
G)**. Moreover, we also found a two-fold increase in the expression of the IL6 family cytokine receptor's
signal transducer, gp130 receptor (IL6ST) in postmortem cortical RTT samples compared with age-
matched control samples **(Fig. S5E)**. More importantly, postmortem cortical tissues showed a fivefold
increase in MECP2-KO IL6 protein levels compared with controls, consistent with the in vitro
observat*ions* **(Fig. S5C)**. We observed a significant decrease in levels of synapsin1 puncta in control
iPSC-derived neurons exposed to MECP2-KO ACM **(Fig. 5D)**. These alterations were associated with
functional synaptic defects in neurons exposed to conditioned media from MECP2-KO astrocytes. We
also observed a significant reduction in synapsin1 puncta and total dendritic length in neurons when
rhIL-6 was added to the medium **(Figs. S5F-G)**. Moreover, control neurons treated with rhIL-6 that
matured in multi-electrode array (MEA) plates decreased their spontaneous activity **(Fig. S5H)**. Finally,
we co-cultured independently generated human neurons and astrocytes. MECP2-KO astrocytes
negatively affected the dendritic morphology of control neurons, reducing total dendritic length, the
number of segments per neuron, and the spine density **(Figs. S5I-L)**.

L1 inhibition rescues MECP2-KO astrocytes defects

To confirm that the deleterious effect of MECP2-KO astrocytes is due to the accumulation of L1
elements, we treated our cultures with FDA-approved antiretroviral reverse transcriptase (RTi) drugs
Lamivudine (3TC) and Stavudine (d4T), known to inhibit L1 reverse transcription[40]. We additionally used

another antiretroviral drug, Nevirapine (NVP), as a negative control since no effect on L1 reverse
transcription has been previously shown[40,41]. When MECP2-KO astrocytes were chronically treated with
RTi, cytoplasmic L1 copies, and IL6 expression decreased to control levels (**Figs. 5B-C**). Similarly, we
observed reductions in other pro-inflammatory cytokines and related factors, including CXCL10
(interferon-gamma-induced protein 10), CCL2 (monocyte chemoattractant protein 1), and TMEM173
(stimulator of interferon genes) (**Figs. S5M-O**). Previous studies have reported elevated levels of the
excitatory neurotransmitter glutamate in the cerebrospinal fluid of patients with MECP2 mutations [22,24].
MeCP2 has been shown to modulate glutamate clearance by regulating various astroglial genes[23].
Indeed, we found that several genes related to glutamate metabolism were misregulated in MECP2-
KO astrocytes (**Fig. S5P**). MECP2-KO astrocytes treated with RTi restored glutamate, reactive oxygen
species (ROS), and glutathione levels to control cells (**Figs. 5E-G**).

L1 inhibition rescues MECP2-KO neuronal defects in mixed cultures

We also tested if RTi could restore neuronal function in mixed brain organoid cultures containing
MECP2-KO neurons and astrocytes. Treating MECP2-KO-derived cultures with RTi, rescued neuronal
cell body area and dendritic complexity (**Figs. 5H-I** and **Figs. S5Q-S**). To determine if the control of L1
activity by RTi can ameliorate synapse formation in MECP2-KO neurons, co-localization of SYN1
(synapsin I) and PSD95 (postsynaptic density protein 95) on a MAP2 (microtubule-associated protein
2) positive neurite was quantified. MECP2-KO neurons in the RTi-treated culture displayed a significant
increase in co-localized puncta compared to those without RTi-treatment, which is not seen in the NVP-
treated culture (**Fig. 5J**). At the functional level, neuronal network activity measured by the number of
bursts and firing rate in MECP2-KO-derived organoids was also rescued in cultures treated with RTi
(**Figs. 5K-L**).

L1 reverse transcriptase inhibition rescued survival and behavior in a Mecp2-KO mouse model

Hemizygous Mecp2-KO males have a median lifespan of approximately 10 weeks and exhibit
pronounced behavioral and physiological abnormalities [25,29,30,42]. We treated these animals with 3TC or
d4T in the drinking water (physiological dose of 2 mg/ml[26] starting at 4 weeks old. Both 3TC and d4T
ameliorated the overall aspect of the Mecp2-KO animals and extended their lifespan (**Fig. 6A-B**). Next,
we tested whether any behavior would be rescued following the animals' extended lifespan. The open-
field test revealed that 3TC treatment improves the mobility of Mecp2-KO mice (**Figs. 6C-D** and **Fig.
S6A**). Previous studies have demonstrated that this Mecp2-KO mouse model exhibits decreased
anxiety behavior [27]. We found no effect of 3TC treatment on the anxiety-related behavior of Mecp2-KO
mice (**Fig. 6E** and **Fig. S6B, Supp. Videos 1-3**). There are conflicting results regarding the sociability
of these rodent models [31,33]. In our hands, Mecp2-KO mice have impaired social memory, and this
phenotype can be rescued with 3TC treatment (**Fig. 6F**). Following behavioral analysis, the animals'
brains were fixed and subjected to Golgi staining for morphological analysis. We focus on the prefrontal
cortex, as this region is critical for social behavior. We found that the 3TC treatment rescued neuronal
soma size and spine density of Mecp2-KO neurons in the cingulate cortex (**Fig. 6G-H** and **Fig. S6C-
D**).

**L1 reverse transcriptase inhibition recovered abnormal gene expression in both astrocytes and
neurons in the brains of Mecp2-KO mice**

Given that behavioral impairments and aberrant neuronal morphologies in Mecp2-KO mice were
restored by 3TC treatment, we next examined whether these restorations are also detected at the
molecular level. To this end, we conducted spatial transcriptomics analysis (Xenium, 5k) on forebrain
sections from 8–9-week-old WT, Mecp2-KO, and Mecp2-KO mice treated with 3TC. We applied
unsupervised clustering on the combined dataset of 270,741 cells that passed quality control to define
cell clusters. We identified 10 unique clusters and annotated each cluster (excitatory neurons, inhibitory
neurons, astrocytes, oligodendrocyte progenitor cells (OPC), oligodendrocytes, microglia, endothelial
cells, pericytes, fibroblasts, smooth muscle cells) with the expression profiles of canonical cell type
marker genes and spatial information (**Fig. 7A-C**). Since we have demonstrated that astrocytes play
critical roles in neuronal morphology and function (**Figs. 5 and S5**), we focused on astrocytes and

reanalyzed the astrocyte cluster using a different clustering method. We found that the cell number and
spatial distribution of each astrocyte subcluster in the forebrain were not substantially altered in Mecp2-
KO compared to WT **(Figure 7D-E)**. However, a comparison of gene expression across the entire
astrocyte cluster between WT and Mecp2-KO identified 173 genes as significantly up-regulated and
173 as significantly down-regulated under Mecp2-KO conditions **(Table S6)**. Gene ontology (GO)
analysis of the upregulated genes revealed enrichment of terms related to oxidative stress and
interleukin signal **(Fig. 7F)**. Intriguingly, significant overlap was observed between the genes
upregulated in these Mecp2-KO astrocytes and the 30 genes included in the gene list (5006 genes) in
the Xenium panel of 150 genes **(Table S4)** that were upregulated under ISS conditions **(Fig. 7G)**.
Among the 4 overlapping genes, ALDOA (aldolase A), an enzyme involved in cellular metabolism and
previously implicated in inflammation [32] and DNA damage responses [43], was significantly
downregulated in Mecp2-KO mouse brains following 3TC treatment. Furthermore, we found that 3TC
treatment restored the impaired expression of inflammation-related genes implicated in these GO terms
(Fig. 7H). In addition, consistent with these transcriptomic findings, immunostaining results showed that
complement component 3 (C3), a key immune marker associated with reactive astrocytes, was
upregulated in Mecp2-KO astrocytes and reduced by 3TC treatment **(Fig. 7I-J)**.

Based on the recovery of gene expression in astrocytes, we next compared gene expression in clusters
of excitatory neurons in WT, Mecp2-KO, and Mecp2-KO mice treated with 3TC and performed GO
analysis. We found that downregulated genes in Mecp2-KO relative to WT were enriched for GO terms
associated with synaptic signaling **(Fig. S7A)**. In addition, genes upregulated with 3TC treatment in
Mecp2-KO mice were enriched in similar GO categories **(Fig. S7B)**. Moreover, when we examined
genes in a specific GO term (synaptic signaling), expression of those genes was reduced in Mecp2-KO
but recovered with 3TC treatment **(Fig. S7C)**, suggesting that 3TC treatment recovered impaired gene
expression not only in astrocytes but also in neurons.

DISCUSSION

The modern cell has evolved to coexist with autonomous L1 sequences by activating defense
mechanisms within the cell's intrinsic immune system [44]. Moreover, DNA hypermethylation responses
were significantly pronounced in evolutionarily younger retrotransposons, suggesting that recent
genomic insertions might be particularly sensitive indicators of space exposure[11]. By employing a multi-
omics approach that integrates proteomics, bulk transcriptomics, and single-nucleus transcriptomics,
we provided a comprehensive view of the molecular alterations that affect human brain cells across
space. This led us to reveal a complex interplay between viral pathways, astrocyte reactivity, and
inflammatory responses, a novel phenomenon that we call Space-Induced Neural Senescence (SINS).

Multiple studies on neurological and psychiatric disorders have revealed increased L1 expression in
the brain [39]. We previously reported that the accumulation of L1-derived ssDNA in human iPSC-derived
astrocytes triggered pathological inflammation in a severe progeroid neurological condition, Aicardi-
Goutieres Syndrome (AGS) [12]. There is substantial evidence that transposable elements, particularly
L1 retrotransposons, are aberrantly reactivated in the aging brain and across various
neurodegenerative diseases, including Amyotrophic Lateral Sclerosis (ALS), Alzheimer's Disease (AD),
Parkinson's Disease (PD), and Huntington's Disease[45-49]. Our findings demonstrated that the histone
proteins and deacetylation processes are altered by the space environment, likely contributing to an
"epigenetic rewiring" and the activation of viral pathways in space. Histone downregulation is frequently
observed under conditions of cellular stress or in disease states, indicating alterations in chromatin
dynamics that affect the accessibility and stability of genomic DNA [50]. Reduced histone expression
impairs nucleosome assembly, leading to chromatin relaxation and aberrant activation of otherwise
silenced genomic regions, such as transposable elements[51]. This chromatin disruption often results in
compromised epigenetic regulation, enabling the transcriptional activation of repetitive sequences,
including L1s. Histone depletion thus facilitates L1 mobilization, reflecting a critical link between

chromatin remodeling and potential genome instability[52]. The activation of L1s following histone
downregulation has profound biological implications, particularly in the context of aging and
neurodegeneration. Aberrant L1 activation is closely associated with age-related neurodegenerative
diseases such as Alzheimer's disease, where increased retrotransposition events correlate with
neuronal loss and cognitive decline[53]. It is interesting to note that our re-analyses of the blood
transcriptomic data from the NASA Twins Project [54] have also revealed an increase in viral molecular
pathways, suggesting that the reactivation of endogenous retroviruses is likely not specific to brain
tissue. However, due to the brain's limited regenerative capacity, the impact of this phenomenon is
expected to be more pronounced. Our findings are in agreement with the space medicine literature[7,55-
57] that dysregulation in OxPHOS proteins corroborates the idea that mitochondrial impairment is a
shared pathological mechanism underlying diverse neurodegenerative processes, including
Parkinson's disease (PD), Huntington's, and Alzheimer's diseases[58,59]. Also, corroborated with the
genomic findings from the Inspiration4 mission, which also evaluated the OxPHOS genomic framework,
revealing a sustained downregulation of PD-associated OxPHOS genes in astronauts' peripheral blood
mononuclear cells (PBMCs)[60,61].

We and others have previously revealed that brain tissue isolated from RTT patients is more susceptible
to L1 expression[14,16]. Here, we found that the accumulation of L1 transcripts in the absence of MeCP2
induces IL6 release by astrocytes. Several groups have previously reported interleukin abnormalities
and cytokine release in RTT [18,21,62,63]. However, in the absence of a stimulating agent, this observation
was not linked to a potential disease contribution. Activation of transposable elements exacerbates
aging and neurodegeneration [45,64,65], promoting genomic instability, neuroinflammation, and aberrant
immune signaling, as observed in our experiments in LEO. Moreover, recent studies have revealed a
significant upregulation of transposable elements, including L1s, in excitatory neurons and astrocytes
in AD brains[66,67], demonstrating that an increase in TE expression in AD is related to immune activation,
synaptic dysfunction, and cellular senescence, similar to the findings associated to the SINS
phenomenon, described in this study.

Thus, we hypothesized that the cytokine abnormalities in RTT are not a direct consequence of MECP2
mutations, but rather a pathological condition exacerbated by the accumulation of cytoplasmic L1
copies in astrocytes, which triggers the IL6-mediated neurotoxic response. Our findings linked L1
expression in RTT astrocytes to disease phenotypes, thereby enabling us to test whether L1 inhibition
by RTi could reduce inflammation and mitigate neurotoxicity. The analyses from independent
spaceflights corroborate that, in LEO, brain organoids activate sterile-mediated cellular responses,
resulting in the activation of an immune response and epigenetic derangement that can activate viral
pathways, inducing SINS, and immune dysfunction, offering a new framework for understanding how
"dormant" genomic elements become pathogenic under certain conditions. Notably, increased levels of
IL6 are also associated with neuropsychiatric conditions such as depression, Alzheimer's Disease, and
aging [68,69]. Thus, we compared *in vitro* and *in vivo* non-treated and RTi-treated models in MECP2-KO
brain organoids and Mecp2-KO mice, respectively. Our findings are summarized in **Fig. 8**. *In vitro*, the
loss of MeCP2 results in increased transcription of L1 elements, leading to elevated inflammation,
characterized by increased levels of IL6, ROS, glutamate, and inflammatory markers such as CXCL10,
CCL2, and TMEM173. The molecular findings in the secretome panel and mitochondrial framework
analysis indicate that these signatures are consistent with known space-environment stressors,
including activation of the viral pathway, leading to oxidative stress and impairment of mitochondrial
homeostasis. Importantly, they also align with recent literature describing spaceflight-induced activation
of retrotransposons in mice [10] and the cellular senescence pathway in hematopoietic stem cells [70-72].
These findings collectively suggest a potentially space-environment-sensitive state, characterized by
shifts in differentiation toward an inflammatory and viral-responsive pattern of transcription and
translation.

MECP2-KO brain organoids treated with RTi restore neuronal function, promoting astrocytic
homeostasis and indicating a significant recovery in cellular and synaptic function. *In vivo*, untreated

Mecp2-KO mice exhibited severe neurological deficits, including reduced survival, sociability, mobility,
and speed. When Mecp2-KO mice were subjected to RTi treatment, we observed improved
neurological symptoms and increased survival, as well as increased sociability, mobility, and speed,
compared with untreated controls. Furthermore, we confirmed these therapeutic effects also at the
molecular level. We demonstrated that RTi treatment suppresses cytokine responses and reduces
inflammatory markers typically associated with senescent cells and aged mice. These effects
underscore a causal link between retroelement activation and space-related inflammation, as observed
in our experiments on the ISS during independent spaceflights.

Radiation and microgravity stressors do not act in isolation but rather appear to have additive or even
synergistic effects on brain function. Since we cannot isolate the various factors acting at the ISS, we
refer to the combined effects of cosmic radiation, microgravity, and other unmeasured factors as the
"space environment". Due to its nature, space radiation - especially GCR (galactic cosmic radiation) -
plays a crucial role in biological systems. Rodents subjected to a complex 33-beam GCR simulation
exhibited deficits in attentional processing, along with disrupted dopaminergic neurotransmission in the
prefrontal cortex, despite maintaining reward-related behaviors [73], illustrating that cosmic radiation
inflicts structural and functional damage on neural circuits. In this study, the radiation measured in all
experiments at ISS, specifically the GCR dose recorded, averaged 0.119 mGy/day (absorbed dose)
and 0.126 mGy/day (dose equivalent). In contrast, the South Atlantic Anomaly (SAA) doses were about
0.084 mGy/day (absorbed dose) and 0.086 mGy/day (dose equivalent). Given these values, the
recorded doses from our spaceflights fall within the lower end of the expected radiation exposure range
for ISS missions. For shorter missions, such as typical CRS resupply flights, the biological impact is
likely negligible, as the total absorbed dose remains low and below critical thresholds that can trigger
acute health effects. Moreover, we have previously investigated the role of radiation in brain organoids
submitted to proton radiation (0.5 and 2 Gy)[74]. We found dose- and time-dependent increases in DNA
damage, oxidative stress, and activation of DNA damage checkpoint genes, particularly at 48 hours
post-irradiation. Other significant findings included declines in mitochondrial function and neural identity,
corroborating the results of this study; however, we acknowledge the limitations of attributing the
phenotypic findings solely to this stressor.

The collective findings on the secretome panel suggest that the space environment may be triggering
a "recalibration" of the brain organoids' developmental program, causing these models to strengthen
their cellular defenses to address the unique challenges of space. This may serve as a mechanism to
achieve a more resilient, faster-differentiated state, leading to a SINS state. The combined findings of
the three spaceflights validate the results described in this study, demonstrating that the space
environment leads to biological alterations in the brain organoids model, resulting in a phenotype that
displays profound cellular stress, impaired genome regulation, and inflammation, potentially mediated
by cGAS/STING pathway, a mechanism previously described by our group[12], and which is activated by
presence of viral DNA[75] and retroelements[75,76], especially in neurodegenerative diseases[77,78]. It is well
established in the literature that transposable element activation is not merely a bystander effect in
aging and senescence but a potentially causal mechanism in disease progression, contributing to cell-
to-cell variability and neuronal vulnerability [79,80].

The successful preclinical data highlight RTi's immediate utility in therapeutic regimens for RTT and
other neuroinflammatory disorders, revealing a potential novel neuroprotective treatment for astronauts
on long spaceflight missions as a translational venue. The proven safety and tolerability profile of 3TC
makes it an attractive candidate for drug repurposing strategies aimed at preventive treatment in
astronauts, particularly given the altered immune function and potential viral reactivation that can occur
during extended space missions [81,82]. Prophylactic and therapeutic administration of 3TC drugs could
theoretically mitigate latent retroviral or endogenous retroviral elements whose activation may be
potentiated by the stress-induced immunosuppression associated with spaceflight[83], and its known
efficacy against viral replication during spaceflight has already been documented in the literature[84]. The
broad anti-inflammatory and immunomodulatory effects of RTis, particularly their capacity to mitigate

inflammation driven by endogenous retroviruses, offer a novel and powerful mechanism to counter the
pervasive immune dysregulation observed in microgravity.

Our study's findings align with recent astronaut [60,85] and mouse [20] data on the space environment,
underscoring the critical need for a comprehensive approach to protecting astronaut health during long-
duration space missions. This could include implementing antiviral and anti-inflammatory strategies to
ensure the well-being of future generations of explorers.

LIMITATIONS OF THE STUDY

Limitations of the study are attributable to inherent difficulties in the space environment and in LEO.
The experiments are extremely precious and constrained by the payload capacity of each spaceflight,
making it difficult and costly to send multiple biological replicates on a single flight.

RESOURCE AVAILABILITY

Lead Contact

Requests for further information and resources should be directed to and will be fulfilled by the lead
contact, Dr. Alysson R. Muotri (muotri@ucsd.edu).

Materials Availability

This study did not generate new unique reagents or materials

Data and code availability

• The mass spectrometry proteomics data have been deposited to the ProteomeXchange
Consortium via the PRIDE (https://www.ebi.ac.uk/pride/) partner repository with the dataset
identifier   PXD055001   and   also   in   the   NASA   GeneLab   repository,
https://osdr.nasa.gov/bio/submission/rdsa/RDSA-939 and are publicly available as of the date
of publication.

• The transcriptomics data is deposit in the NASA GeneLab repository under the study number
OSD-765 and also in the Zendo (10.5281/zenodo.16388786 and 10.5281/zenodo.16389553)
and are publicly available as of the date of publication.
551
• This paper does not report original code.

• Any additional information required to reanalyze the data reported in this paper is available
from the lead contact upon request.

ACKNOWLEDGMENTS

We would like to thank the invaluable efforts of the ISSCOR team: Natalia Chermont, Isabelle Luz,
Paulo Carvalho, Juliana Fisher, and Blake Tsu. Roberto Herai, Luiz Pedro Petroski, and Adriano
Ferrasa for all the insights on RNAseq data. We acknowledge the invaluable efforts of Space Tango,
our implementation partner, in providing telemetric data from all missions. We acknowledge the Space

Radiation Analysis Group (SRAG) for providing the radiation data used in this work. We acknowledge
Daniel McClatchy and Claire Delahunty from Yates's lab for their experimental advice, Richard Waite,
Narayanaganesh "Ganesh" Balasubramanian, Leslie Sandoval, Michael Krawitzky, and Matt Willetts
from Bruker for their support in this work. This work was supported by grants from the NSF/CASIS grant
(CBET 2126309) and NASA (80JSC018M0005) to A.R.M.; AMED (JP20gm1310008,
JP24wm0625306), JSPS KAKENHI (JP23H00391) and an Intramural Research Grant (3-9) for
Neurological and Psychiatric Disorders of NCNP Grant to K.N.; JSPS KAKENHI (JP22K15201,
JP23H04170, JP25H01866) and AMED (23gm1310008) to H.N. This research was also supported by
NPO the Rett Syndrome Support Organization to H.N. (2019-2-10) and an NIH grant (#S10 OD026929)
from Dr. K. Jepsen at the UCSD Institute of Genomic Medicine. A.R.M., H.N., and K.N. are inventors
on a pending U.S. patent application for the use of reverse transcriptase inhibitors for Rett syndrome
(U.S. Application No. US18/569,721).

AUTHOR CONTRIBUTIONS

Conceptualization, A.M., H.N., A.M., P.M., E.V., K.N., A.R.M.; Methodology, A.M., K.N., J.Y., A.M.,
P.M., A.R.M.; Investigation, A.M., H.N., A.M., T.U., S.M., P.M., L.L., L.C., E.V., N.S., A.S., A.S., B.F.,
A.L., D.A., M.I., J.Y.,T.O., M.H., K.N., A.R.M. Visualization, A.M., P.M., L.L., L.C., A.S., K.N., A.R.M.
Funding acquisition, K.M., H.N., A.R.M.; Project administration, K.N., A.R.M.; Supervision, K.N., A.R.M.
Writing – original draft, A.R.M.; Writing – review & editing, A.M., H.N., A.M., T.U., P.M., L.L., L.C., E.V.,
N.S., A.S., A.S., B.F., A.L., D.A., M.I., J.Y., K.N., A.R.M.

DECLARATION OF INTERESTS

The authors declare no competing interests.

SUPPLEMENTAL INFORMATION

Figures S1-S7, Tables S1-6, Supplementary Videos 1-3.

589
Fig. 1. Schematic brain organoids experiments sent to the International Space Station (ISS). This
schematic depicts the process and experimental setup for studying brain organoids under space
environment conditions aboard the International Space Station (ISS). Induced pluripotent stem cells
were differentiated into brain organoids. These brain organoids were launched on SpaceX CRS-SpX18,
CRS-SpX30; CRS-SpX31 missions (on average 30 days) and maintained aboard the ISS in an
automated system (CubeLab), exposed to the space environment. A synchronous ground control group
of brain organoids was maintained in parallel under identical conditions in another CubeLab to enable
comparative studies. Following their return from ISS, comprehensive molecular analyses, including
proteomic and transcriptomics assessments, were conducted on both ISS and Earth-bound brain
organoids to evaluate the impact of space environment conditions on molecular and cellular processes
and development.
601

602
603

Fig. 2. Proteomics and transcriptomics analysis data from healthy brain organoids at the International Space Station (ISS). (A) Volcano plot of proteomics data showing proteins upregulated (in blue) and downregulated (in red) in brain organoids at ISS, when compared to ground control. **(B)** Reactome pathway analysis of the proteins upregulated in the proteomics experiment. **(C)** Gene Ontology (GO) analysis for the mitochondria framework. 629 downregulated proteins are significantly enriched in functions and pathways crucial for cellular energy production (respiration) and the synthesis of mitochondrial proteins. **(D)** Heatmap of a framework of general downregulation in histone showing global chromatin remodeling. **(E)** Secretome panel on a subsequent flight (Spx31). The media of a batch of healthy brain organoids with 2 months at arrival were evaluated using Olink Reveal® panel technology (Thermo Fisher Scientific). Sample controls were used to determine precision within and between plates, triplicate negative controls samples set the limit of detection (LOD) and triplicate samples were used to standardize protein levels within a plate. Secretome sample run 1:1. Counts of known sequences were translated into Normalized Protein eXpression (NPX) values as their log-2 scaled quantification. **(F)** Heatmap of the top 25 Differentially Expressed Genes (DEGs) in ISS *vs.* ground conditions. The colors represent fold changes, with red indicating genes that are downregulated and blue indicating genes that are upregulated. **(G** Reactome pathway analysis of the upregulated genes from the transcriptomics experiment (Panel D). **(H)** WikiPathway enrichment analysis of the transcriptomic data (Panel D). **(I)** The UMAP plot shows the distribution of various neural cell types identified across both ground and ISS conditions. Each cluster is color-coded to represent a specific cell population: lower layer neurons (red), upper layer neurons (orange), radial glial cells (yellow-green), intermediate progenitors (green), glial cells (teal), astrocytes (cyan), dividing neural progenitors (light blue), early neurons (purple), pan cortical neuron (pink), and oligodendrocytes progenitor cells (magenta). The plot highlights the spatial organization of cell clusters, emphasizing the diversity of cell types within the dataset. **(J)** UMAP analysis comparing brain organoids samples from ground and ISS. **(K)** Stacked bar plots representing the relative proportions of different cell types in the ground and ISS. Each bar represents a specific cell type, with the dashed line indicating equal proportion mark. **(L)** *GFAP* and *TGFB2* expression in cells across ground and ISS conditions. Violin plots show the distribution of gene expression levels for GFAP and TGFB2 in cells from the ground and ISS environments. Each violin represents the kernel density estimation of log expression values, indicating higher levels of expression in the ISS condition compared to the ground condition. **(M)** Heatmap of healthy brain organoids, using bulk transcriptomics analysis, in a subsequent flight (SpX30). For proteomics analysis: $p \leq 0.01$; $fc \geq 1.2$. Test T Student for comparisons. For transcriptomics bulk analysis: $p \leq 0.05$; $fc \geq 1.2$. For pathway analysis: *Homo sapiens* as the reference organism, minimum overlap of 3 genes, enrichment factor > 1.5, and $p < 0.01$.

643
644 **Fig. 3. Reanalysis of publicly available RNAseq data from ISS and ground control mouse retinas.**
645 **(A)** Heatmap of ISS and ground control retinas. **(B)** Principal coordinates analysis (PCoA) of ISS and
646 ground control retinas**. (C)** Volcano plot of differentially expressed genes. **(D)** GSEA analysis of
647 spaceflight and ground control retinas.

649
Fig. 4. Proteomics and transcriptomics analysis overview of MECP2-KO brain organoids at the
International Space Station (ISS). (A) Volcano plot displaying proteins upregulated in MECP2-KO
cortical brain organoids (in blue) and downregulated (in red), when compare to control brain
organoids, both at ISS. **(B)** Reactome pathway analysis from proteins upregulated in MECP2-KO
brain organoids at ISS. This panel includes an overall pathway visualization followed by a focusing on
the "cellular response to stress" pathway, highlighting the viral pathway. **(C)** Heatmap from a
subsequent spaceflight (SpX30) demonstrating a robust and consistent difference between the
MECP2-KO brain organoids *vs.* healthy brain (control) organoids, both submitted to the space
environment, at ISS. The hierarchical clustering indicates a distinct gene expression signature
associated with the MeCP2-KO condition at ISS. Each experiment was done in biological duplicates.
For proteomics bulk analysis: p≤0.01; fc≥1.2. Test T Student for comparisons. For pathway analysis:
Homo sapiens as the reference organism, minimum overlap of 3 genes, enrichment factor > 1.5, and
p < 0.01.

664
Fig. 5. L1 expression in MECP2-KO astrocytes causes neurotoxicity that can be rescued with
reverse transcriptase inhibitors (RTi), but not nevirapine (NVP). (A) Schematic figure on how
ssDNA accumulates in the astrocytes, leading to increased neurotoxicity in neurons. **(B)**
Quantification of extrachromosomal L1-derived single-stranded DNA (ssDNA) on MECP2-KO
astrocytes. **(C)** Quantification of released IL-6 from astrocytes by enzyme-linked immunosorbent
assay (ELISA). **(D)** Quantification of synaptic puncta per 20 μm on control and MECP2-KO neurons
treated with astrocyte conditioned media (ACM). **(E)** Quantification of glutamate buffering in MECP2-
KO astrocytes. Data are represented as mean ±SEM. Levels are normalized to control equaling one.
(F) Quantification of reactive oxygen species (ROS) levels in MECP2-KO astrocytes. Data are
represented as mean ±SEM. Levels are normalized to control without treatment equaling one. **(G)**
Quantification of intracellular glutathione in MECP2-KO astrocytes. Data are represented as mean
±SEM. Levels are normalized to control without treatment equaling one. **(H)** Quantification of neuronal
cell body area. **(I)** Quantification of neuronal dendritic complexity by number of branch points. **(J)**
Quantification of co-localized synaptic puncta (defined by SYN1+/PSD95+ on MAP2+ dendrite) per 20
μm on neurons from each condition. **(K)** Quantification of the total number of spikes of control
neurons matured on MEA plates. **(L)** Quantification of weighted mean firing rate from organoids in
each condition. Reverse-transcriptase inhibitor (RTi) or nevirapine (NVP) treatment is indicated. Each
color represents one condition: control (black), MECP2-KO (blue), MECP2-KO+RTi (orange), and
MECP2-KO+NVP (green). Each experiment was done in biological triplicates. The number of
measurements (*n*) for these experiments is indicated directly by the symbols (o) on each bar graph.
Data are represented as mean ±SEM. $p<0.5$ (*), $p<0.01$ (**), $p<0.001$ (***) and $p<0.0001$ (****);
ordinary one-way ANOVA.
687

Fig. 6. Treatment of Mecp2-KO with reverse transcriptase inhibitors (RTis) in a mouse model.
(A) Time plot of scores from a neurological symptoms assessment of Mecp2-KO mice with and without
treatment of RTis lamivudine (3TC) and stavudine (d4T) at 2 mg/mL. The time plot of neurological
symptoms scores depicts 5 weeks after RTi treatment was introduced in drinking water (weeks 5-10).
(B) Time plot of percent survival depicts mice's lifespan in a number of days. **(C)** Quantification of total
distance traveled and average speed of WT, Mecp2-KO, and 3TC-treated Mecp2-KO mice in the open-
field test. **(D)** Quantification of average speed (cm/seconds) by WT, Mecp2-KO, and 3TC-treated
Mecp2-KO mice in an open arm chamber during the elevated plus maze test. **(E)** Quantification of time
spent (in percentage of total time) by WT, Mecp2-KO, and 3TC-treated Mecp2-KO mice in an open arm
chamber during the elevated plus maze test. **(F)** Quantification of social novelty index of WT, Mecp2-
KO, and 3TC-treated Mecp2-KO mice in a three-chamber test. Social novelty index is measured as
time spent in the chamber with an intruder divided by the total time spent in both chambers. **(G)**
Quantification of soma size and spine density. **(H)** Quantification of neurons from the cingulate cortex
in brains fixed from WT, Mecp2-KO, and 3TC-treated Mecp2-KO mice. Spine density is measured by
number of spines per 10 μm of dendrite. The number of measurements (*n*) for these experiments is
indicated directly by the symbols (o) on each bar graph. Data are represented as mean ±SEM. $p<0.5$
(*), $p<0.01$ (**), $p<0.001$ (***) and $p<0.0001$ (****); ordinary one-way ANOVA.
707

Fig. 7. Spatial transcriptomics analysis of WT, Mecp2-KO and 3TC-treated Mecp2-KO astrocytes.
(A) UMAP plot showing 270741 cells annotated into 10 clusters from the brains, including PFC region,
of WT (n=3), Mecp2-KO (n=3) and 3TC-treated Mecp2-KO (n=3) mice. **(B)** Scaled dot plot of canonical
cell type marker genes used to define each cluster. **(C)** Spatial representative image plots of cell types
in (A) across coronal hemisection of the WT, Mecp2-KO and 3TC-treated Mecp2-KO mice forebrains.
(D) Spatial representative image plots of astrocyte subtypes in coronal hemisections of WT, Mecp2-KO
and 3TC-treated Mecp2-KO mouse forebrains. **(E)** UMAP analysis comparing astrocyte subtypes
across WT, Mecp2-KO and 3TC-treated Mecp2-KO mouse forebrains. **(F)** GO analysis of upregulated
genes in Mecp2-KO astrocytes compared to WT astrocytes. (G) Overlap between ISS upregulated
genes and Mecp2-KO astrocyte upregulated genes. **(H)** Heatmap of differentially expressed response
to interleukin-7 (GO:0098760) and cellular response to external stimulus (GO:0071496) genes that
were upregulated in Mecp2-ko astrocytes compared to WT astrocytes and downregulated in 3TC-

treated Mecp2-KO astrocytes. *(I)* Immunofluorescence staining for C3 (red) and S100β (green) in the
PFC regions of 8-week-old WT, Mecp2-KO and 3TC-treated Mecp2-KO mouse brains. Nuclei are
counterstained with Hoechst (blue). Scale bar, 5 μm. **(J)** Quantification of C3 signal intensity within the
soma of S100β+ astrocytes. The number of individual measurements (n) for each experimental group
is indicated by the symbols (o) on the bar graph. Data are presented as mean ± SEM. ****p<0.001 by
ordinary one-way ANOVA.
728
729

730
Fig. 8. Summary comparing non-treated and RTi-treated models in MECP2-KO brain organoids
and Mecp2 mice. Top Panel: Brain Organoids (MeCP2-KO). The left side depicts non-treated
MeCP2-KO brain organoids, where the loss of MeCP2 results in increased transcription of LINE-1
elements, which are recognized as infection triggers. This leads to elevated inflammation,
characterized by increased levels of IL-6, ROS, glutamate, and inflammatory markers such as
CXCL10, CCL2, and TMEM173. In parallel, neuronal dysfunction is observed, including reductions in
cell body size, dendritic branching, dendrite length, the formation of synaptic puncta, and firing rates.
The right side represents organoids treated with RTi (reverse transcriptase inhibitor). Treatment
restores neuronal function and promotes astrocytic homeostasis. Neuronal characteristics, such as
cell body size, number of dendritic branches, dendrite length, synaptic puncta formation, and firing
rates, are restored to levels comparable to the control condition, indicating a significant recovery in
cellular and synaptic function. **Bottom Panel: Mecp2-KO Mice.** The left side shows non-treated
Mecp2 mice, where a decrease in soma size and spine density is observed, accompanied by severe
neurological symptoms, including reduced survival rate, sociability, mobility, and speed. The right side
illustrates the effects of RTi treatment in Mecp2 mice. Treatment leads to increased soma size and
spine density, accompanied by improvements in neurological symptoms. Treated mice show
enhanced survival rate, sociability, mobility, and speed compared to non-treated controls.
748

754
Fig. S1. CubeLab and environmental conditions and functionality. (A) Illustration representation
of the CubeLab capabilities monitored during the space environment experiment on the International
Space Station (ISS). The outer view and internal setup of the CubeLab, which houses the wells
containing the organoids, are depicted, as well the organoids inside the well in the ISS. The CubeLab
is designed to control temperature, humidity, and gas composition, automatically exchanging the cell
medium. Additionally, the CubeLab is equipped with cameras that allow real-time observation of the
organoids from Earth during spaceflight. **(B)** Environmental parameters monitored during the space
environment experiment on the International Space Station (ISS) compared to ground control. The X-
axis represents the days of the mission, while the Y-axis shows values for CO_2 concentration,

organoid incubator temperature and fresh culture medium temperature, in both environments. The
CO_2 concentration was maintained at around 5%; the organoid incubator temperature was kept

around 37°C; and the fresh culture medium temperature at 4°C until it was used to feed the organoids
every two days. Duplicate measurements were taken for each temperature to ensure accurate
readings. **(C)** Schematic view of the brain organoid used in this work. On the left, a confocal image of
a human cortical organoid stained for stem cell marker SOX2 and neural progenitor marker nestin.
Cells are organized in radial units that resemble early cortical plate layering. Neural stem and
progenitor cells (magenta nuclei) densely populate the rosette region. Neural progenitors (green)
extend long basal processes outward, guiding the migration of newly generated neurons. DAPI (blue)
marks all nuclei. Scale bar = 70 μm. On the right, a schematic representation of the cellular
architecture observed in the organoid, depicting the spatial arrangement and lineage progression from
undifferentiated stem cells to mature neurons and astrocytes. **(D)** Comparison of average organoid
diameter over five weeks between the Ground and ISS units. The graphs show the diameters of
organoids in control and MECP2-KO groups. **(E)** REM – Radiation environmental monitoring readings
for the three spaceflights used in this work. Each point represents the mean diameter of multiple
organoids within a single well ± standard error of the mean (SEM) for organoids cultured under each
condition (ISS and Ground). A significant difference in organoid diameter is observed in the MECP2-
KO group at week 5 between the microgravity (ISS) and Earth gravity (Ground) conditions, suggesting
an impact of microgravity on MECP2-KO organoid growth. *p≤ 0.05.
783

Fig. S2. Proteomics and transcriptomics approach for viral pathway evaluation. (A) Heatmap of a framework of histone proteins showing upregulation of acetyltransferases and downregulation of deacetylases at ISS. **(B)** Differentially expressed genes (DEG) comparison between this study and NASA Twin study using Ingenuity Pathway Analysis (IPA), showing canonical pathways (on the left) and diseases and biological pathways (on the right). The colors correspond to activation z-score (negative score in blue, positive in orange). For pathway analysis: *Homo sapiens* as the reference organism, minimum overlap of 3 genes, enrichment factor > 1.5, and p < 0.01.
(C) Heatmap of differentially expressed genes across cell types. The heatmap shows the normalized expression levels of key marker genes across different neural cell populations. Warmer colors (red) indicate higher expression levels, while cooler colors (blue) represent lower expression. Key genes exhibit distinct expression patterns across cell clusters.

Fig. S3. Retroelement expression analysis using publicly available RNA-seq data. (A) Expression levels of transposable elements in the retina of mice exposed to space flight (GSE159771). **(B)** Subfamily-level expression of L1 elements in the spaceflight retina**. (C)** Expression of transposable elements in mouse lung following simulated five-ion galactic cosmic radiation (GSE199680) **(D)** L1 subfamily expression profiles in mouse lung under five-ion radiation condition. (**B, D**) The color of L1 subfamily members represents evolutionary age: ancient members in gray, old members in black, middle members in blue, and young members in red.

Fig. S4. Proteomics and transcriptomics analysis of MECP2-KO brain organoids experiments.
(A) Reactome from the proteomic analysis comparing control cells to MECP2-KO cells in ground conditions, emphasizing that MeCP2-KO cells demonstrate upregulation of the viral pathway. This analysis highlights the intrinsic alterations in MECP2-KO cells that predispose them to enhanced viral pathway activity and nervous system development even without the influence of the space environment. **(B)** Heatmap of bulk transcriptomics comparing ground MECP2-KO brain organoids vs. ground MECP2-KO brain organoids. For proteomics bulk analysis: p≤0.01; fc≥1.2. Test T Student for comparisons. For pathway analysis: *Homo sapiens* as the reference organism, minimum overlap of 3 genes, enrichment factor > 1.5, and p < 0.01.

Fig. S5. MECP2-KO astrocytes phenotypes and interplay with neurons. (A) Increase L1
endogenous retroelement expression in MECP2-KO astrocytes measured by qPCR. **(B)**
Quantification of TNFα, IL-4, IL-13, and IL-10 released in the media from control and MECP2-KO
astrocytes by enzyme-linked immunosorbent assay (ELISA). **(C)** IL-6 protein level in post-mortem
RTT brain sample compared to an age-matched control sample by ELISA. **(D)** Increased IL-6
released from RTT-derived astrocytes was rescued when MECP2 was re-expressed in the cells. The
dotted red line indicates the background buffer signal for the ELISA assay. **(E)** Expression levels of
the IL-6 family cytokine receptor's signal transducer, gp130 receptor (IL6ST in post-mortem RTT brain
samples compared to an age-matched control sample by qPCR. **(F)** Reduction of synapsin1 puncta in
control neurons upon IL-6 treatment. **(G)** Reduction of Total Dendritic Length (TDL) in control neurons
upon IL-6 treatment. **(H)** Reduction of neuronal activity measured by the number of control neurons
upon IL-6 treatment, measured through MEA over a 260-second time course. **(I)** Schematic diagram
of the experimental design. iPSC-derived NPCs were used to produced enriched populations of
neurons and astrocytes for co-cultured experiments. **(J)** Co-culture experiments of neurons and
astrocytes from the two genetic backgrounds (control and MECP2-KO) showing the negative impact
of MECP2-KO astrocytes in neurons regarding total dendritic length (TDL), number of segments **(K)**,
and number of spines **(L)**. **(M)** Quantification of CXCL10, CCL2 **(N)**, and TMEM173 **(O)** expression in

MECP2-KO astrocytes treated with RTis by enzyme-linked immunosorbent assay (ELISA). **(P)**
Quantification of GS, GLT1, GLAST, and SLC26A7 gene expression in control and MECP2-KO
astrocytes compare to control. **(Q) Top:** Representative image of neurons. SYN::EGFP positive cells
co-stained with a cortical marker CTIP2 and EGFP to be further traced. Scale bar, 1000 μm for
neurons; Scale bar, 100 μm for CTIP+ and EGFP+ neurons. **Bottom:** Representative images of
tracings from control, MECP2-KO, RTi-treated MECP2-KO, and NVP-treated MECP2-KO neurons.
(R) Quantification of dendritic length and complexity **(S)** in mixed cultures of neurons and astrocytes.
Reverse-transcriptase inhibitor (RTi) or nevirapine (NVP)-treatment is indicated. Each color
represents one condition: control (black), MECP2-KO (blue), MECP2-KO+RTi (orange), and MECP2-
KO+NVP (green). Each experiment was done in biological triplicates. The number of measurements
(*n*) for these experiments is indicated directly by the symbols (o) on each bar graph. Data are
represented as mean ±SEM; $p<0.5$ (*), $p<0.01$ (**), $p<0.001$ (***) and $p<0.0001$ (****).
851

852
Fig. S6. RTT-like phenotypic analysis of reverse transcriptase inhibitor-treated Mecp2-KO mice.
(A) Digital tracking of WT, Mecp2-ko and 3TC-treated Mecp2-KO mice exposed to the open-field test.
Representative traces of mice activity obtained from video tracking. **(B)** Digital tracking of WT, Mecp2-
ko and 3TC-treated Mecp2-KO mice exposed to the elevated plus maze test. Representative traces of
mice activity obtained from video tracking. **(C)** Representative images of somata of Golgi-stained layer
2/3 pyramidal neurons in PFC of WT, Mecp2-ko and 3TC-treated Mecp2-KO mice. Scale bar, 10 μm.
(D) Representative images of dendrites of Golgi-stained layer 2/3 pyramidal neurons in PFC of WT,
Mecp2-ko and 3TC-treated Mecp2-KO mice. Scale bar, 5 μm.
861

863
Fig. S7. Spatial transcriptomics analysis of WT, Mecp2-KO and 3TC-treated Mecp2-KO excitatory
neurons. (A) GO analysis of downregulated genes in Mecp2-KO excitatory neurons compared to WT
excitatory neurons. **(B)** GO analysis of upregulated genes in 3TC-treated Mecp2-KO excitatory neurons
compared to Mecp2-KO excitatory neurons. **(C)** Gene expression profiles of synaptic signaling
(GO:0099536) genes that were significantly downregulated in Mecp2-KO excitatory neurons compared
to WT excitatory neurons and whose expression was significantly improved by 3TC treatment.
870

871 Supplementary Video 1. Representative video of WT mice performing the open field test.
872
873 Supplementary Video 2. Representative video of Mecp2-KO mice performing the open field
874 test.
875
876 Supplementary Video 3. Representative video of Mecp2-KO+3TC mice performing the open
877 field test.
878

 MATERIAL AND METHODS

IPSC tissue culture

All iPSCs were previously characterized[86]. In addition, we also used MECP2 isogenic cell lines using
CRISPR/Cas9 genome editing (RTT/C9)[86]. Two isolated clones for each cell line were selected and
expanded for further differentiation and experimentation. Mycoplasma and CNV analyses were
performed regularly in all cultured cells. For consistency, "Control" or "MECP2-KO" were used in the
manuscript. All iPSCs were cultured and manually passaged onto Matrigel-coated (Corning) and fed
daily with mTESR1 (StemCell Technologies). iPSCs were differentiated into NPCs as described[87].
NPCs were cultured on the plate coated with 10 µg/mL of poly-L-ornithine (Sigma-Aldrich) and 2.5
µg/mL laminin (Life Technologies). Media was changed every other day with NG medium (DMEM/F12
50/50 1X (Corning Cellgro), 1% HEPES (VWR International), 0.5% Penicillin Streptomycin (Pen-Strep)
(Life Technologies), 1X GlutaMAX (Life Technologies), 0.5% N2 NeuroPlex (Gemini Bio-Products) and
1% Gem21 NeuroPlex (Gemini Bio-Products) was supplemented with 20 ng/mL basic human fibroblast
growth factor (bFGF) (R&D Systems, 414-TE).

Neural differentiation

For neural differentiation, bFGF was withdrawn from the NPCs containing NG media, and Rock inhibitor
(Y27632) was added at a concentration of 10ng/mL for 48 hours. Cells matured for 5 to 8 weeks,
replacing the media every 3 to 4 days. IPSC-derived cortical organoids were differentiated following[1].
Briefly, iPSCs were cultured, dissociated using a 1:1 Dulbecco's phosphate-buffered saline (DPBS,
Fisher Scientific) and StemPro Accutase solution, transferred to 6-well plates, and kept in suspension.
Neural induction media consisted of DMEM/F12, 1% Glutamax, 1% N2 Neuroplex, 1% non-essential
amino acids (Gibco), 1% Pen-Strep, 1µM of Dorsomorphin (R&D Systems), and 10µM SB431542 (SB,
Stemgent). Neural proliferation media consisted of Neurobasal media (Life Technologies), 2% Gem21
Neuroplex, 1% non-essential amino acids, 1% Glutamax, 20 ng/mL epidermal growth factor (EGF,
Peprotech), and 20 ng/mL bFGF. For neuronal maturation, the cells were kept in the same media, but
the growth factors were withdrawn. Organoid results are combined from four separate batches of
differentiation.

Astrocytes were differentiated as described[88]. Briefly, monolayer NPCs are dissociated into single-cell
suspension using StemPro Accutase. To form astrospheres, cells were seeded at a density of 4-5
million cells per well in a 6-well plate (Genesee, 25-105) shaking at 95 r.p.m. at 37°C in NSCR EL
medium containing DMEM/F12 (Gibco) with 1% N2 supplement (Gibco) and 0.1% SM1 supplement,
1% nonessential amino acids (Gibco), 1% penicillin/streptomycin (Gibco), 1% GlutaMAX solution
(Gibco), supplemented with 20 ng/ml epidermal growth factor 1 (EGF-1, Peprotech) and 20 ng/mL
leukemia inhibitory factor (LIF, Peprotech). Following 2-4 weeks in the NSCR EL media, the
astrospheres were expanded in NSCR media now supplemented with 20 ng/mL EGF and 20 ng/mL
FGF (NSCR EF) for 2-4 weeks. Following this expansion phase, astrospheres were plated on 20 µg/mL
poly-L-ornithine and 5 µg/mL laminin and remained in NSCR EF media, allowing astrocyte progenitor
cells (APCs) to migrate outward until 70% confluency. For astrocyte assays, progenitors were expanded
and seeded at 50,000 cells/well in a 96-well plate. These progenitor cells can be expanded and frozen
in cryovials for later use. For differentiation into mature astrocytes, media was changed to contain half
of the NSCR basal media and half Astromed CNTF media. Astromed media contained Neurobasal
media (Gibco) with 0.2% SM1, 1% NEAA, 1% penicillin/streptomycin, 1% Glutamax and 10ng/mL of
ciliary neurotrophic factor (CNTF, Peprotech). After 48 hours, cells were transitioned into full Astromed
CNTF media for 14 days.

CubeLab operations (flight and ground)

Space Tango's flight and ground hardware support 12 wells containing impellers and automated
operations (fluid conveyance, fluid routing, thermal control, environmental monitoring, imaging, and
sampling) within each 9U CubeLab™. All wetted materials selected within the CubeLab were certified
biocompatible and sterilizable. Wetted materials that come in contact with fluids or cells were sterilized,
assembled, maintained sterile, tested, and confirmed to be free of microbial contamination prior to
loading experimental fluids and biology. Science and the Flight system CubeLab were prepared at the
Kennedy Space Center (KSC) Space Station Processing Facility (SSPF) in Brevard County, FL.
Scheduled operations began shortly after science loading, continuing through launch, ascent, and on-
orbit, and ended upon fixation and transfer to cold stowage. Space Tango's Powered Ascent Utility
Locker (PAUL) allows CubeLabs to be fully powered while installed in the SpaceX Cargo Dragon
vehicle. The PAUL provides the optimal culture environment and capabilities for the entire time the
science remains within the CubeLab, including while on the launch pad. Flight and Ground systems
were run synchronously. The Ground system scheduled operations were initiated 9 days after the Flight
system.

The Flight system CubeLab culture environment was maintained at 37°C and 5–7% CO_2, and culture
media was held at 2–8°C, providing optimal culture conditions. Media exchanges were performed every
3 – 4 days using microfluidic routing and pumping. Organoids were exposed to the CubeLab
environmental gases through a 10:1 (base:curing agent) polydimethylsiloxane (Sylgard 184, Dow)
interface within each well cap. Impeller-driven agitation was required for the Flight system on Earth,
through ascent and installation, and during media exchanges. During all other times, no agitation was
performed. Imaging of wells was performed twice per week for 30 days, beginning after installation and
ending prior to cold stowage transfer. SpaceX mission CRS-25 was 30 days. On the 30th day of on-orbit
operations, 4% paraformaldehyde fixation of 6 wells and RNALater fixation of the remaining 6 wells was
performed. The Flight system was stored at approximately 4°C for 6 days until science removal at the
SSPF. Additionally, Space Tango's cryovial flight hardware contains 6 stainless steel boxes, each
supporting up to 98 1 mL cryovials (Nunc, 374088) for up to 588 1 mL cryovials per 9U CubeLab™.
The Flight system CubeLab culture environment was maintained at 37°C and 5% CO_2, providing optimal
environmental culture conditions. SpaceX CRS-25 mission was 30 days. Ground control samples were
loaded into 1x stainless steel cryovial box and maintained within a humidified incubator set to 37°C and
5% CO_2, located at the SSPF, until the flight samples returned to Earth. Flight and ground samples
were run synchronously, and scheduled operations were initiated at the exact same time.

The Ground system experiment was performed at the Space Tango Headquarters in Lexington, KY.
The CubeLab culture environment was maintained at 37°C, and 5–7% CO_2, and culture media was held
at 2–8°C, providing optimal culture conditions. Media exchanges were performed every 3–4 days using
microfluidic routing and pumping. Impeller-driven agitation was required for the Ground system
throughout the entire experiment on Earth. Imaging of wells was performed during media exchanges
only due to impeller movement. On the 30th day of matched on-orbit operations, 4% paraformaldehyde
fixation of 6 wells and RNALater fixation of 6 wells was performed. The Ground system was stored at
approximately 4°C for 17 days until scientific removal. CubeLab environmental telemetry (culture
environment temperature, temperature of culture media, CubeLab CO_2, CubeLab humidity) was
recorded during powered operations. Telemetry data was available in near real-time through Space
Tango's customer dashboard.

Imaging operations and post-processing (flight and ground)

During the ascent phase, agitation was enabled continuously on all wells. 13-second videos were
recorded once per hour for each well during this time. Once installed into the ISS, the constant agitation
period was discontinued. This was changed to record videos during the agitation event during the media
exchanges, which occurred every 3.5 days. To get more insight into how well the organoids were

growing, a health check was added after the first 2 media exchanges. The health check involved
capturing a photo of each well once per hour for the remainder of the mission. Post-processing included
adding a scale bar onto each image that represents the size of 1mm in the image. Images were
uploaded to Space Tango's data portal. The ground study payload had constant agitation enabled from
the experiment start until fixation. During this time, 13-second videos were recorded once per hour for
each well. Additionally, health check images were added before and after each media exchange at an
interval of 3.5 days. Post-processing included adding a scale bar onto each image that represents the
size of 1mm in the image. Images were uploaded Space Tango's data portal.

Mass spectrometry bulk analysis

A total of 500 ng of each sample was randomized loaded onto individual Evotips trap columns following
manufacture's protocol. The peptides were separated using an Evosep One LC system (Evosep,
Denmark) in 44 min gradient (30 samples per day predefined Evosep proprietary method) on a PepSep
8 (8 cm × 150 μm X 1.5 μm C18 column, Bruker Daltonics) at 50°C. The mobile phases were 0.1% FA
in LC-MS grade water (buffer A) and 0.1% FA in 99.9% ACN (buffer B). The LC system was coupled
with timsTOF Pro 2, a hybrid trap ion mobility spectrometer and quadrupole time-of-flight instrument
(Bruker Daltonics) operated in positive mode previously calibrated for mass (m/z) and ion mobility
dimension (1/K0), using three ions of Agilent ESI-Low Tuning Mix (m/z 622.0289; 922.0097 and
1221.9906). The nano-electrospray source (CaptiveSpray source, Bruker Daltonics) was operated at
1600 V of Capillary voltage, 3.0 L/min of dry gas, and 180 °C of dry temperature. Data was acquired in
dia-PASEF acquisition mode with a cycle time estimate of 1.8 s46. To cover the mass range of 400 to
1200 m/z, 32 isolation windows of 25 m/z width were used. Ramp Time and accumulation time were
set to 100 ms, and collision energy ramped linearly as a function of the mobility from 59 eV at 1/K0 =
1.6 Vs. cm−2 to 20 eV at 1/K0 = 0.6 Vs. cm−2.

Data processing for mass spectrometry-based bulk proteomics

All diaPASEF data was imported in Spectronaut (v.19, Biognosys) and processed using directDIA
workflow which does not require spectral library from DDA runs. In brief, Spectronaut Pulsar search
engine firstly generates DDA-like pseudospectra from the fasta database and then performs searches.
The human protein reference database (UP000005640 reviewed canonical proteins) was used with
Pulsar and the following settings: Trypsin/P as specific enzyme; peptide length from 7 to 52; max missed
cleavages 2; toggle N-terminal M turned on; Carbamidomethyl on C as fixed modification; Oxidation on
M and Acetyl at protein N-terminus as variable modifications; FDR at PSM, peptide and protein group
level set to 0.01; minimum fragment relative intensity 1%; 3–6 fragments kept for each precursor; the
extraction of the ion chromatograms and ion mobility window was set as "dynamic" with correction factor
1.

Functional analysis for mass spectrometry-based bulk proteomics

The interpretation of systems-level studies was done on Metascape, an online platform that provides a
comprehensive resource for annotating and analyzing gene/protein lists[53]. In this study, we used
enrichment analysis to compare thousands of protein sets defined by their involvement in specific
biological processes, protein localization, enzymatic function, pathway membership, and protein
network analysis and interactions. All data was post-processed to generate Kappa similarities, and all
pairs of enriched terms were computed into clusters of comparable terms[54].

The proteins quantified in at least 70% of the samples within each group were subjected to statistical
and graphical analyses using the R programming environment. Abundance values were log10-
transformed prior to analysis. Boxplots were generated using the ggplot2, dplyr, and tidyr packages[89].
Heatmaps were constructed using the heatmap REF package, based on ANOVA p-values to highlight
differential expression patterns. Volcano plots were created using the ggplot2 and ggrepel REF

packages. Gene Ontology (GO) enrichment analyses were performed using Cytoscape[90] and
STRING[91].

Protein identifiers and associated statistics ($\log_2$ fold change and adjusted p-values) were submitted to
Metascape (www.metascape.org) for pathway enrichment analysis. Parameters were set to default
values: *Homo sapiens* as the reference organism, minimum overlap of 3 genes, enrichment factor >
1.5, and p < 0.01. The resulting output was used for downstream processing, focusing on the table
containing pathway description, enrichment ratio, $-\log_{10}$(p-value), and the number of proteins
contributing to each term. Data visualization was performed in R (version 4.3.2) using the readxl,
ggplot2, and viridis packages. The enriched pathways were ordered according to their enrichment ratio,
and a dot plot was generated to display the top biological processes. In the plot, the x-axis represents
the enrichment ratio (GeneRatio), the y-axis lists pathway descriptions, dot size corresponds to the
number of input proteins (Count), and dot color encodes statistical significance based on $-\log_{10}$(p-
value).

Secretome panel evaluation by Proximity Extension Assay (PEA) Olink® Reveal (Thermo Fisher
Scientific)

The Olink Reveal technology uses Proximity Extension Assay (PEA)[92], where a matched pair of
antibodies labelled with unique complementary oligonucleotides (proximity probes) bind to the
respective target protein in a sample. As a result, the probes come into close proximity and hybridize
to each one, enabling DNA amplification of the protein signal, which is quantified on a next generation
sequencing read-out which has been described in detail previously. Antibodies targeting 1034 unique
proteins are distributed across 384-plex panel. Assays are run in a 96-well format. Each plate consisted
of 32 secretome samples (in triplicates) from brain organoids that belong to the ground experiment and
the ISS experiment, using 20 μL of media sample. In brief, pairs of oligonucleotide-labeled antibody
probes designed for each protein bind to their target, bringing the complementary oligonucleotides in
close proximity and allowing for their hybridization. The addition of a DNA polymerase leads to the
extension of the hybridized oligonucleotides, generating a unique protein identification "barcode". Next,
library preparation adds sample identification indexes and the required nucleotides for Illumina
sequencing. Prior to sequencing using the Illumina® NovaSeq™ 6000/NextSeq™ 550/NextSeq™
2000, libraries go through a bead-based purification step and the quality is assessed using the Agilent
2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA). The raw output data is quality controlled,
normalized and converted into NPX®, values as their log-2 scaled quantification unit of relative
abundance. Data normalization is performed using an internal extension control and an external plate
control to adjust for intra- and inter-run variation. All assay validation data (detection limits, intra- and
inter-assay precision data, predefined values, etc.) are available on manufacturer's website
(www.olink.com). Data was processed by Olink's MyData Cloud Software.

RNA sequencing and transcriptomic analysis

For transcriptomic analyses, we combined brain organoids from the three healthy donors for ground
processing for RNA sequencing. Ribodepletion RNA sequencing was performed at the UC San Diego
IGM core facility. Data was analyzed by ROSALIND® (https://rosalind.bio/), with a HyperScale
architecture developed by ROSALIND, Inc. (San Diego, CA). Briefly, reads were trimmed using
cutadapt. Quality scores were assessed using FastQC. Reads were aligned to the Homo sapiens
genome build hg19 using STAR. Individual sample reads were quantified using HTseq and normalized
via Relative Log Expression (RLE) using DESeq2 R library. Read Distribution percentages, violin plots,
identity heatmaps, and sample MDS plots were generated as part of the QC step using RSeQC.
DEseq2 was also used to calculate fold changes and p-values and perform optional covariate
correction. The clustering of genes for the final heatmap of differentially expressed genes was done
using the PAM (Partitioning Around Medoids) method using the fpc R library. Hypergeometric
distribution was used to analyze the enrichment of pathways, gene ontology, domain structure, and

other ontologies. The topGO R library was used to determine local similarities and dependencies
between GO terms to perform Elim pruning correction. Several databases were referenced for
enrichment analysis, including Interpro, NCBI, MSigDB, REACTOME, WikiPathways. Enrichment was
calculated relative to a set of background genes relevant to the experiment. Functional enrichment
analysis of pathways, gene ontology, domain structure, and other ontologies was performed using
HOMER.

In our study, we employed the DESeq2 package in R for differential expression analysis, utilizing a
significance threshold of $p < 0.05$. Raw count data, obtained from experimental samples representing
ground and ISS conditions, were processed and normalized. Subsequently, differential expression
analysis was performed, allowing for the identification of genes significantly altered between the two
conditions, with a fold change threshold of 1.5. Additionally, to gain insights into the biological pathways
underlying these expression changes, Reactome pathway analysis was conducted on the differentially
expressed genes, with a significance threshold of $p < 0.05$. Furthermore, we performed additional
pathway analysis using Wiki Pathways library, employing a false discovery rate (FDR) adjusted p-value
cutoff of 0.1, to further elucidate the enriched biological processes associated with the observed gene
expression alterations.

Xenium experiment

Mice were anesthetized and perfused trans-cardiac with PBS followed by 4% buffered
paraformaldehyde (PFA). Brains were removed and fixed with 4% PFA overnight at 4°C. The fixed
tissue was cryoprotected with 30% sucrose and frozen in optimal cutting temperature medium (OCT).
Coronal sections were obtained by sectioning the tissue at 10 µm using Leica CM3050S cryostat.
Sections were washed with PBS and placed onto Xenium slides.

The Xenium 5K Mouse Pan-Tissue and Pathways Panel (10x Genomics) was used for transcriptomic
profiling. Probe hybridization, RNase treatment, ligation, amplification, and cell staining were performed
according to the manufacturer's protocol. Data acquisition and on-board analysis were conducted using
the Xenium Analyzer (v3.3).

Xenium data processing

Regions of interest (ROIs) in the WT, Mecp2-KO, and 3TC-treated Mecp2-KO brains were manually
selected using Xenium Explorer 3 (v3.1.1). Only cells within ROIs with more than 250 detected genes
and more than 350 transcripts were included in downstream analysis. Data normalization, principal
component analysis (PCA), UMAP-based dimensionality reduction, and cell clustering were performed
using the Seurat package (v5.0.1). Cell type annotations were manually assigned based on the
expression of established marker genes. Differential gene expression analysis was performed using
the FindAllMarkers function in Seurat. Over-representation analysis of genes included in the predefined
gene list of the Xenium panel in the transcriptome data of brain organoid samples returned from the
ISS was performed using Fisher's Exact Test. Gene Ontology (GO) analysis of Biological Processes
was conducted using the clusterProfiler package (v4.4.4).

Reverse Transcriptase inhibitors

The reverse-transcriptase inhibitors for cultured cells were prepared as follows: Lamivudine (3TC,
Sigma-Aldrich) was prepared in dimethyl sulfoxide (DMSO, Sigma-Aldrich), and 10 µM of 3TC was
used in the media. Stavudine (d4T, Sigma-Aldrich) was re-suspended in water, and 1µM of d4T was
used in the media in combination with 3TC. Nevirapine (Sigma Aldrich, SML0097) was re-suspended
in DMSO and 400 nM nevirapine was used in media. Control cells contained 10 µM of DMSO as a
vehicular control.

Immunofluorescence

Cells cultured in monolayer were washed with DPBS, fixed in 4% paraformaldehyde (PFA, Core Bio) for 20 minutes at room temperature, and permeabilized with a solution of DPBS and 0.1% Triton X-100 (Promega) for 10 minutes at room temperature. Cells were incubated at room temperature with blocking solution for 30 minutes (DPBS, 10% donkey or goat serum (Fisher Scientific), and 0.005% Triton X-100). Primary and secondary antibodies were diluted in a solution containing DPBS, 1% donkey or goat serum, and 0.005% Triton X-100. Cells were incubated with the primary antibody overnight at 4°C. The next day, cells were washed with DPBS three times for 5 minutes and incubated with secondary antibodies for 30 minutes at room temperature. Cells were washed twice with DPBS, and nuclei were stained using DAPI (VWR International) for 10 minutes at room temperature. Finally, cells were washed once with DPBS for 5 minutes, and the coverslips were mounted using Prolong Gold antifade mounting (Life Technologies) onto glass slides (Fisher Scientific).

For cortical brain organoids, cells were washed with DPBS and placed in a 1.5 mL microcentrifuge tube with 4% PFA overnight at 4°C. Cortical organoids were washed with DPBS and incubated in 30% sucrose (Sigma-Aldrich) for 48 hours at 4°C. The tissue was removed from the sucrose solution, placed with optimal cutting temperature compound (OCT, VWR) within an embedding mold (Fisher Scientific), and incubated at -80°C overnight. Using a cryostat (Leica), the molds containing the organoids were sliced at 10.0 μm and placed onto a glass slide which was left to dry at room temperature for at least 4 hours. The slides were lined with a hydrophobic barrier pen (Vector Laboratories). Organoid slides were permeabilized, stained, and mounted as explained above.

Immunohistochemistry

Mice were anesthetized and perfused trans-cardiac with PBS followed by 4% buffered paraformaldehyde (PFA). Brains were removed and fixed with 4% PFA overnight at 4°C. The fixed tissue was cryoprotected with 30% sucrose and frozen in optimal cutting temperature medium (OCT). Coronal sections were obtained by sectioning the tissue at 40 μm using Leica CM3050S cryostat. Sections were washed with PBS to remove the OCT, permeabilized and blocked with blocking buffer at RT for 1 h, and then incubated with primary antibody diluted in blocking buffer overnight at 4°C. After three washes with PBS, the sections were incubated with corresponding secondary antibodies at RT for 2 h. The following primary and secondary antibodies were used in this experiment: rabbit anti-hC3D (1:500; DAKO, Cat.# A0063), mouse anti-S100β (1:500; Sigma, Cat.#S2532); CF488 donkey anti-mouse IgG (H+L), highly cross-adsorbed (1:500; Biotium), CF555 donkey anti-rabbit IgG (H+L), highly cross-adsorbed (1:500; Biotium). Hoechst 33258 (1:500) was used for nuclear staining. After three washes with PBS, the sections were mounted on glass slides. Fluorescence images of astrocyte were acquired using Zeiss LSM 800 confocal microscope.

Magnetic cell sorting

Briefly, iPSC-derived cells were dissociated using the neurosphere dissociation kit (MACS), incubated with the CD44-PE and CD184-PE (both antibodies were from BD Biosciences) for 15 minutes in the dark at 4°C and washed twice with dPBS to remove unbound antibodies before sorting. Next, the cells were incubated with PE antibodies and conjugated with magnetic beads (MACS). Following the antibody incubation, cells are loaded into a magnetic column, and the cells that are CD44 and CD184 positive are retained in the column. The remaining enriched neuronal population is obtained and counted for viability and seeding.

Astrocyte calcium wave propagation

Calcium waves were induced by focal mechanical stimulation of single astrocytes in the center of the field of view. A 175 W Xenon lamp and a wavelength switcher (Lambda DG-4, Sutter Instrument)

provided 485 nm fluorescence excitation. The emission from Fluo-4 loaded cells is detected at a
wavelength of 520 nm using a F-Fluar 40X/1.3 NA oil immersion objective and an attached 12-bit CCD
camera. Changes in Fluo-4 fluorescence intensities emitted at 520 nm were acquired at 0.67 Hz. The
calcium imaging data were visualized and analyzed offline using MetaMorph software (Universal
Imaging Corporation). The fluorescent intensity of Fluo-4 was determined in individual cells in every
astrocyte soma in the field of view. The distance that calcium waves propagated within a certain time
was determined by measuring the distance from the stimulating point to the farthest responding cell in
the field of view.

Morphological analysis of iPSC-derived neurons

Cultured NPCs were transduced with a lentivirus backbone containing Synapsin 1 (SYN) promoter
upstream of an EGFP reporter[93]. The transduction multiplicity of infection (M.O.I.) was equal to 2. Cells
transduced with a SYN::GFP lentivirus were differentiated into neurons for 5-6 weeks, fixed, and stained
with GFP and CTIP2 antibodies. Differentiated neurons were traced using Neurolucida v.2017 (MBF
Bioscience, Williston, VT) connected to a Zeiss Axio Imager 2 microscope at a 40X oil objective. Only
neurons that were both GFP- and CTIP2-positive were included in this analysis. The morphology of the
neurons was quantified using Neurolucida Explorer v.11 (MBF Bioscience, Williston, VT), and the
results reflect the summed length of all neurites and dendrites per traced neuron. Sholl analysis was
performed using Neurolucida Explorer's Sholl analysis. This analysis specified a center point within the
soma and created a grid of concentric rings around it with radii increasing in 10 μm increments.
Neuronal complexity was determined by recording the number of intersections within each ring.

Synaptic puncta quantification

5-6-week-old neurons were fixed and stained for the following markers: SYN1, postsynaptic density
protein 95 (PSD-95), and MAP2 (refer to **Immunocytochemistry** for protocol). Using a fluorescence
microscope (Z1 Axio Observer Apotome Zeiss), the slides were imaged by compiling Z stack images
taken from incremental focus distances. Co-localized SYN1 (presynaptic) and PSD-95 (postsynaptic)
were quantified using the compiled Z stack image. Only co-localized puncta that were in proximity of a
MAP2-positive neurite were included in this analysis.

Measurement of cortical organoid diameter

1182 Cortical organoids were imaged on an Evos FL Imagine System (Thermo Fisher Scientific) using a 4X
1183 magnification. The images were uploaded to Image J, and the diameter of each organoid was
1184 determined by the program's length measurement.

RNA extraction and qPCR

RNA was extracted using the RNAse Plus Mini Kit (Qiagen). Briefly, adherent cells were washed with
DPBS and further lysed using RLT buffer with B-mercaptoethanol (Thermo Fisher Scientific) and
dissociated using a 1mL syringe (Fisher Scientific, 1482330) and a needle (VWR, BD- 305196). RNA
from lysates was extracted following the manufacturer's instructions. cDNA was generated using the
Qiagen Quantitect Reverse Transcription Kit. Protocol was followed on 1 μg of the RNA samples. qPCR
was performed using selected primers and SYBR Green master mix (BioRad). Diluted cDNA was used
in a 96-well or 384-well plates (BioRad) and ran on a BioRad CFX machine. PCR conditions were as
follows: initial cycle of 95ºC for 32 min, followed by 40 cycles of 15s at 95ºC, 30s at 58ºC, 30s at 72ºC,
with a final step of 5s at 57ºC and 5s at 95ºC for the melting curve. 3 plates were run to obtain triplicates
per condition. The L1 primers used were: ORF1:
ATGGGGAAAAAACAGAACAGAAAAACTGGAAACTCTAAAACGCAGAGCGCCTCTCCTCCTCCAA
AGGAACGCAGTTCCTC; ORF2:
GCTCATGGGTAGGAAGAATCAATATCGTGAAAATGGCCATACTGCCCAAGGTAATTTACAGATTC

AATGCCATCCCCATC; ORF2-3′UTR:

TGGAAACCATCATTCTCAGTAAACTATCGCAAGAACAAAAAACCAAACACCGCATATTCTCACTCA

TAGGTGGGAATTGA

1202 PCR array

A PCR array was performed using RT^2 Profiler PCR Array 384 format (Qiagen), following the
manufacturer's instructions. Briefly, 400 ng of total RNA was used for reverse transcription using the
RT^2 First Strand Kit (Qiagen). Genomic DNA was eliminated prior to reverse transcription. Real-time
PCR was performed using RT^2 SYBR Green Mastermix and cDNA mix, divided into 384 wells. Plates
were sealed with Optical Adhesive Film and ran into a CFX384 (BioRad). Run conditions: 1 cycle of 10
min at 95°C, followed by 40 cycles of 15s at 95°C and 1 minute at 60°C, where data collection was
performed. For data analysis, GeneGlobe Data Analysis Center was used. Reference genes used:
peptidyl-prolyl cis-trans isomerase H (PPIH), glyceraldehyde-3-phosphate dehydrogenase (GAPDH),
Non-POU Domain Containing Octamer Binding (NONO), Lactate dehydrogenase A (LDHA), Actin Beta
(ACTB). Significant changes in gene expression were set to a min of 2fold differences.

1213 Multi-electrode array (MEA)

12-well MEA plates (Axion Biosystems) were coated with 100 µg/mL of poly-L-ornithine (Sigma-Aldrich)
and 5 µg/mL laminin (Life Technologies). Cultures were placed in the center of the MEA well to fully
cover the electrodes. Media was changed twice a week, and recordings were measured one day after
the media was changed. Recordings were performed once a week using a Maestro MEA System and
AxIS Software (Axion Biosystems). A band-pass filter with 10 Hz and 2.5 kHz cutoff frequencies was
used, and a spike detector threshold was set at 5.5 times the standard deviation. For each recording,
the plate was left untouched in the Maestro for 3 minutes prior to recording, then 3 minutes of data were
recorded. Analysis was performed using Axion Biosystems Neural Metrics Tool. The criteria to detect
an active electrode was set to 5 spikes per minute, and the criteria to detect a burst electrode was set
to 5 bursts per minute.

1224 TUNEL assay

Cells were cultured with the different treatments on 96-well plates, and once confluent, the cells were
fixed with 4% PFA and permeabilized with 0.1% Triton X-100 for 15 minutes. Cells were blocked with
3% bovine serum album (Gemini) for 1 hour. Cells were incubated with primary antibodies of Nestin
and TUNEL (Click-iT TUNEL assay kit, Life Technologies) overnight at 4°C. The following day, cells
were incubated with secondary antibodies for 30 minutes at room temperature and with DAPI for 5
minutes at room temperature. Cells were mounted using Prolong Gold Antifade mounting, and images
were taken and manually quantified.

1232 IL6 Elisa

Astrocyte progenitor cellss were dissociated using 1:1 Accutase:PBS for three minutes and seeded at
50.000 cells/well in a 96-well plate. Following 14 days of differentiation with CNTF, astrocyte media was
conditioned for 48 hours and tested for the presence of IL6 without exogenous stimulation (R&D
Systems). All samples were assayed in triplicate following the manufacturer's instructions for cell culture
supernatant. To confirm the functionality of the iPSCs-derived astrocytes, selected wells were
stimulated with 10ng/ml interleukin-1-β (IL-β) during 48 hours prior to IL-6 analysis. Optical density was
measured at 450 nm with wavelength correction set to 540 nm using the Tecan Infinite Pro 200 plate
reader. Standards were assayed in duplicates, and samples were assayed in triplicates. The
corresponding plate was fixed and stained for normalization. Images were taken using Z1 Axio
Observer Apotome Zeiss, and 8-10 pictures were taken per condition. GFAP-positive cells were

manually counted, and DAPI was quantified using Fiji-ImageJ imaging. Recombinant human IL6 was
added as a positive control.

ROS Assay

Astrocyte progenitors were plated at a density of 50,000 cells/well in a white-walled, clear bottomed 96
well plate (Corning) and differentiated for 14 days with CNTF. ROS-Glo H2O2 (Promega) Homogenous
(Lytic) protocol was followed according to the manufacturer's instructions. Astrocytes and RTI
treatments were conditioned in the Astromed media for 48 hours. The H_2O_2 substrate solution was
added to the plate and treated for 4 hours before ROS Glo detection solution was added. The relative
luminescence was recorded using a Tecan Infinite Pro 200 plate reader. Six replicates were run per
condition.

Glutathione Assay

GSH-GLO Glutathione Assay (Promega V6911) protocol for adherent mammalian cells was followed
according to the manufacturer's instructions to measure glutathione levels present in astrocytes. Cells
were seeded at 50,000 cells/well on a white-walled 96 well plate. The medium was removed, 1X GSH-
Glo reagent was added to the cells and incubated for 30 minutes at room temperature. Following the
addition of Luciferin Detection reagent, relative luminescence was recorded using Tecan Infinite Pro
200 luminometer. Values were normalized to GFAP+ expressing cells.

Glutamate Assay

1261 The glutamate assay (ab83389) was used to detect extracellular glutamate levels. Astrocytes
1262 differentiated with CNTF for 14 days were plated in 96-well plates to confluency. The media was
1263 conditioned for 48 hours.

Ca^{2+} Flux Assay

Rhod-4 No Wash Calcium Assay Kit was performed following the manufacturer's instructions. iPSCs-
derived neurons were grown in a 96-well plate and cultured in NG media. Media was changed 48 hours
prior to the experiment.100ul of Rhod-4 Dye-Loading Solution was added per well and incubated at
room temperature for 1 hour. Calcium flux assay was run by monitoring the fluorescence intensity at
Ex/Em = 540/590 nm. Images were acquired for 20 seconds (approximately 600 frames) per field. For
analysis, calcium fluorescence traces were retrieved using customized software, Netcal (Orlandi et al.,
2017) based on MatLab software®. First, a manual selection of Regions of Interest (ROIs) was carried
out to track the activity of all the cells in the culture, both neurons and astrocytes. After analyzing the
recording, further post-processing allows the division of the traces into 2 groups with different firing
behavior: neurons that display a sharp and fast increase of fluorescence followed by a rapid decline in
the wave function. These astrocytes have a slow progression in the fluorescence intensity and a similar
slow decrease in intensity. The classification of the traces was done by training the Netcal program with
the same parameters for each experiment. The analysis of each group highlights specific characteristics
that all the traces in the group share together: Average fluorescence trace, Average number of bursts,
Burst intensity, and Duration of bursts.

Animals

Mecp2-KO mice[25] (C57BL/6 background) were obtained from Mutant Mouse Resource and Research
Centers (MMRRC). They were maintained on a 12-hour light/dark cycle with free access to food and
water. All aspects of animal care and treatment were carried out according to the guidelines of the
Experimental Animal Care Committee of Kyushu University.

3TC and d4T *in vivo* treatments

Mecp2-KO mice were administered 2 mg/mL 3TC (QA-4615, Combi-Blocks) or d4T (D3580, TCI) in
drinking water starting at 4 weeks. All mice were observed daily and weighed once per week.

Scoring of symptoms

Neurological symptoms of mice were scored as previously described[94]. In brief, mice were scored blind
to genotype and treatment status, focusing on mobility, gait, hindlimb clasping, tremor, breathing, and
general condition. Each of the 6 symptoms was scored from 0 to 2; 0 corresponds to the symptom being
absent or the same as in the WT, 1 to the symptom being present, and 2 to the symptom being severe.

Open-field test

The open-field test was performed with 8-week-old WT, Mecp2-KO, and 3TC-treated Mecp2-KO male
mice. Each mouse was placed in the corner of the open-field apparatus (50 cm × 50 cm × 30 cm), which
was illuminated at 100 lux. Total distance traveled (cm) and average speed (cm/s) were calculated
using the automated video tracking software TimeOFCR2 (O'Hara & Co., Tokyo, Japan). Data were
collected for 10 min.

Elevated plus maze test

The elevated plus maze consisted of two opposite open arms (25 cm × 5 cm × 0.25 cm) and two
enclosed arms (25 cm × 5 cm × 15 cm), illuminated at 30 lux and placed 50 cm above the floor. Each
mouse was placed in the central square of the maze (5 × 5 cm), facing one of the closed arms. Mouse
behavior was recorded for 10 min, and the time spent on open arm was analyzed using the automated
video tracking software TimeEP1 (O'Hara & Co., Tokyo, Japan). The percentages of time spent in open
arm over total time were calculated.

Three-chambers social interaction

The three-chambers apparatus is a rectangular non-transparent gray Plexiglas box (60 × 40 × 22 cm),
partitioned into three equal-sized (20 × 40 × 22 cm) chambers by transparent Plexiglas plates with small
openings (5 × 3 cm) allowing access into each chamber. One quarter-cylinder cage made of wire was
placed in the corner of each of the side chambers. The box was illuminated at 20 lux. The subject
mouse was placed into the central chamber and allowed to freely explore the three chambers for 5
minutes. In the first testing section, an age-matched stranger mouse (S1) of the same strain was placed
in one of the cages, and the subject mouse was allowed to explore the three chambers without
restrictions for 10 min. In the second section of preference for social novelty, a second stranger mouse
(S2) was placed in the opposite cage, and the subject mouse was allowed to explore again for 10
minutes. All tests were video-recorded, and the time spent in the quadrant around the cage was
analyzed using the automated video tracking software TimeCSI (O'Hara & Co., Tokyo, Japan). The
social novelty index was calculated by the following formula: [Time spent interacting with stranger 2 in
the second section] / [Time spent interacting with stranger 1+stranger 2 in the second section].

Golgi staining

Golgi staining was carried out using the FD Rapid GolgiStain Kit (FD NeuroTechnologies, Cat.# PK401)
according to the manufacturer's instructions. Briefly, mice at 8-9 weeks of age were euthanized, and
brains were quickly removed and rinsed in double distilled water to remove blood from the surface.
Brains were immersed in impregnation solution prepared by mixing equal volumes of Solutions A and
B, and stored at RT for 2 weeks in the dark. They were then transferred into Solution C and stored at
RT for 1 week in the dark, subsequently embedded in TFM tissue freezing medium (General Data

Healthcare, Cat.# 15-183-1) and frozen at -80°C. Brains were sectioned serially at a thickness of
100 µm using a Cryostat (Leica, Cat.# CM1950). The sections were mounted on gelatin-coated glass
slides and dried naturally overnight at RT. The dry-mounted sections were rinsed with double distilled
water twice for 4 min. They were placed in the staining solution consisting of one part Solution D, one
part Solution E, and 2 parts double distilled water for 10 min in the dark. They were further rinsed in
double distilled water two times for 4 min and dehydrated in sequential rinses of 50, 75, and 95% ethanol
for 4min each rinse. After further dehydration by 100% ethanol, the sections were cleaned in xylene
three times for 4 min and finally coverslipped with Permount. Brightfield images of Golgi-stained
neurons were acquired using a KEYENCE BZ-X700 fluorescence microscope.

Measurement of dendritic soma size *in vivo*

Three mice of each group (WT, Mecp2-KO, and 3TC-treated Mecp2-KO) were examined. For images
of Golgi-stained neurons, Z-series brightfield images of layer 2/3 pyramidal neurons in PFC were
acquired at 20X objective using a KEYENCE BZ-X700 fluorescence microscope. To profile entire cell
morphology in the Z direction, Z-series images were acquired every 1 µm. Sequential images of each
field were reconstructed using the maximum intensity projection method. Soma size was quantified by
tracing the outline of the cell body using ImageJ.

Measurement of dendritic spine density *in vivo*

Three mice of each group (WT, Mecp2-KO, and 3TC-treated Mecp2-KO were examined. Z-series
brightfield images of layer 2/3 pyramidal neurons in PFC were acquired every 0.2 µm at 100X -oil
objective using KEYENCE BZ-X700 fluorescence microscope. Sequential images of each field were
reconstructed using the maximum intensity projection method. The number of spines of secondary
apical dendrites was manually counted, and then spine density per 10 µm was calculated.

Quantification and Statistical analysis

Error bars for figures are standard error of the mean (S.E.M.) using GraphPad Prism v6 (GraphPad
Software Inc). Technical replicates were used to determine the standard error.

Publicly available RNA-seq data analysis

Publicly available RNA-seq data were obtained from the Gene Expression Omnibus (GSE159771 for
spaceflight mouse retina; GSE199680 for mouse lung exposed to simulated GCR). The GCR exposure
included five ion species: 1 GeV 1H (34.8%), 600 MeV/U 28Si (1.1%), 250 MeV/U 4He (18.0%), 350
MeV/U 16O (5.8%), 600 MeV/U 56Fe (1.0%), and 250 MeV 1H (39.2%). Raw FASTQ files were
retrieved using SRA Toolkit. Adapter trimming and quality filtering were performed using Trimmomatic
and Cutadapt. Reads were aligned to the mouse T2T genome (mhaESC) using STAR-2.7.11b, and
transposable elements expression was quantified using TEtranscripts, with annotations from
RepeatMasker. Differential expression analysis was conducted using DESeq2, and gene set
enrichment analysis was performed with GSEA (gene set database:
msigdb.v2023.2.Mm.symbols.gmt).

References

Trujillo, C. A. *et al.* Complex Oscillatory Waves Emerging from Cortical Organoids Model Early Human Brain Network Development. *Cell stem cell* (2019). https://doi.org/10.1016/j.stem.2019.08.002

Fitzgerald, M. Q. *et al.* Generation of 'semi-guided' cortical organoids with complex neural oscillations. *Nature protocols* **19**, 2712–2738 (2024). https://doi.org/10.1038/s41596-024-00994-0

Corujo, D. & Buschbeck, M. Post-Translational Modifications of H2A Histone Variants and Their Role in Cancer. *Cancers (Basel)* **10** (2018). https://doi.org/10.3390/cancers10030059

Hauer, M. H. & Gasser, S. M. Chromatin and nucleosome dynamics in DNA damage and repair. *Genes Dev* **31**, 2204–2221 (2017). https://doi.org/10.1101/gad.307702.117

Azevedo, M. D. *et al.* Oxidative stress induced by sustained supraphysiological intrastriatal GDNF delivery is prevented by dose regulation. *Mol Ther Methods Clin Dev* **31**, 101106 (2023). https://doi.org/10.1016/j.omtm.2023.09.002

Morrison, A. J. Chromatin-remodeling links metabolic signaling to gene expression. *Mol Metab* **38**, 100973 (2020). https://doi.org/10.1016/j.molmet.2020.100973

Gross, S. R. & Kinzy, T. G. Improper organization of the actin cytoskeleton affects protein synthesis at initiation. *Mol Cell Biol* **27**, 1974–1989 (2007). https://doi.org/10.1128/MCB.00832-06

Lai, W. F. & Wong, W. T. Roles of the actin cytoskeleton in aging and age-associated diseases. *Ageing Res Rev* **58**, 101021 (2020). https://doi.org/10.1016/j.arr.2020.101021

Trujillo, C. A. *et al.* Pharmacological reversal of synaptic and network pathology in human MECP2-KO neurons and cortical organoids. *EMBO Mol Med* **13**, e12523 (2021). https://doi.org/10.15252/emmm.202012523

Chen, Z. *et al.* Spaceflight decelerates the epigenetic clock orchestrated with a global alteration in DNA methylome and transcriptome in the mouse retina. *Precis Clin Med* **4**, 93–108 (2021). https://doi.org/10.1093/pcmedi/pbab012

Clement, K. *et al.* Long-term effects of combined exposures to simulated microgravity and galactic cosmic radiation on the mouse lung: sex-specific epigenetic reprogramming. *Radiat Environ Biophys* **64**, 17–27 (2025). https://doi.org/10.1007/s00411-025-01108-4

Thomas, C. A. *et al.* Modeling of TREX1-Dependent Autoimmune Disease using Human Stem Cells Highlights L1 Accumulation as a Source of Neuroinflammation. *Cell stem cell* **21**, 319–331 e318 (2017). https://doi.org/10.1016/j.stem.2017.07.009

Crow, M. K. Long interspersed nuclear elements (LINE-1): potential triggers of systemic autoimmune disease. *Autoimmunity* **43**, 7–16 (2010). https://doi.org/10.3109/08916930903374865

Muotri, A. R. *et al.* L1 retrotransposition in neurons is modulated by MeCP2. *Nature* **468**, 443–446 (2010). https://doi.org/10.1038/nature09544

Yu, F., Zingler, N., Schumann, G. & Stratling, W. H. Methyl-CpG-binding protein 2 represses LINE-1 expression and retrotransposition but not Alu transcription. *Nucleic Acids Res* **29**, 4493–4501 (2001). https://doi.org/10.1093/nar/29.21.4493

16 Zhao, B. *et al.* Somatic LINE-1 retrotransposition in cortical neurons and non-brain
tissues of Rett patients and healthy individuals. *PLoS genetics* **15**, e1008043 (2019).
https://doi.org/10.1371/journal.pgen.1008043
17 Loo, L. *et al.* Fibroblast-expressed LRRC15 is a receptor for SARS-CoV-2 spike and
controls antiviral and antifibrotic transcriptional programs. *PLoS Biol* **21**, e3001967
(2023). https://doi.org/10.1371/journal.pbio.3001967
18 Deverman, B. E. & Patterson, P. H. Cytokines and CNS development. *Neuron* **64**, 61–
78 (2009). https://doi.org/10.1016/j.neuron.2009.09.002
19 Moretti, P. & Zoghbi, H. Y. MeCP2 dysfunction in Rett syndrome and related
disorders. *Curr Opin Genet Dev* **16**, 276–281 (2006).
https://doi.org/10.1016/j.gde.2006.04.009
20 Masarapu, Y. *et al.* Spatially resolved multiomics on the neuronal effects induced by
spaceflight in mice. *Nat Commun* **15**, 4778 (2024). https://doi.org/10.1038/s41467-
024-48916-8
21 Albizzati, E. *et al.* Mecp2 knock-out astrocytes affect synaptogenesis by interleukin 6
dependent mechanisms. *iScience* **27**, 109296 (2024).
https://doi.org/10.1016/j.isci.2024.109296
22 Lappalainen, R. & Riikonen, R. S. High levels of cerebrospinal fluid glutamate in Rett
syndrome. *Pediatr Neurol* **15**, 213–216 (1996). https://doi.org/10.1016/s0887-
8994(96)00218-4
23 Okabe, Y. *et al.* Alterations of gene expression and glutamate clearance in astrocytes
derived from an MeCP2-null mouse model of Rett syndrome. *PLoS One* **7**, e35354
(2012). https://doi.org/10.1371/journal.pone.0035354
24 Pan, J. W., Lane, J. B., Hetherington, H. & Percy, A. K. Rett syndrome: 1H
spectroscopic imaging at 4.1 Tesla. *J Child Neurol* **14**, 524–528 (1999).
https://doi.org/10.1177/088307389901400808
25 Chen, R. Z., Akbarian, S., Tudor, M. & Jaenisch, R. Deficiency of methyl-CpG binding
protein-2 in CNS neurons results in a Rett-like phenotype in mice. *Nat Genet* **27**,
327–331 (2001). https://doi.org/10.1038/85906
26 De Cecco, M. *et al.* L1 drives IFN in senescent cells and promotes age-associated
inflammation. *Nature* **566**, 73–78 (2019). https://doi.org/10.1038/s41586-018-0784-
9
27 Pelka, G. J. *et al.* Mecp2 deficiency is associated with learning and cognitive deficits
and altered gene activity in the hippocampal region of mice. *Brain* **129**, 887–898
(2006). https://doi.org/10.1093/brain/awl022
28 Pitcher, M. R. *et al.* Insulinotropic treatments exacerbate metabolic syndrome in
mice lacking MeCP2 function. *Human Molecular Genetics* **22**, 2626–2633 (2013).
https://doi.org/10.1093/hmg/ddt111
29 Samaco, R. C. *et al.* Female Mecp2(+/-) mice display robust behavioral deficits on
two different genetic backgrounds providing a framework for pre-clinical studies.
Hum Mol Genet **22**, 96–109 (2013). https://doi.org/10.1093/hmg/dds406
30 Ward, C. S. *et al.* MeCP2 Is Critical within HoxB1-Derived Tissues of Mice for Normal
Lifespan. *Journal of Neuroscience* **31**, 10359–10370 (2011).
https://doi.org/10.1523/jneurosci.0057-11.2011
31 Wu, Y. *et al.* Characterization of Rett Syndrome-like phenotypes in Mecp2-knockout
rats. *J Neurodev Disord* **8**, 23 (2016). https://doi.org/10.1186/s11689-016-9156-7

32 Bai, D. *et al.* ALDOA maintains NLRP3 inflammasome activation by controlling AMPK
activation. *Autophagy* **18**, 1673–1693 (2022).
https://doi.org/10.1080/15548627.2021.1997051
33 Castro, J. *et al.* Functional recovery with recombinant human IGF1 treatment in a
mouse model of Rett Syndrome. *Proc Natl Acad Sci U S A* **111**, 9941–9946 (2014).
https://doi.org/10.1073/pnas.1311685111
34 Lilley, C. E., Chaurushiya, M. S. & Weitzman, M. D. Chromatin at the intersection of
viral infection and DNA damage. *Biochimica et Biophysica Acta (BBA) - Gene*
Regulatory Mechanisms **1799**, 319–327 (2010).
https://doi.org/10.1016/j.bbagrm.2009.06.007
35 Nehme, Z., Pasquereau, S. & Herbein, G. in *Histone Modifications in Therapy* Vol. 20
(eds Pedro Castelo-Branco & Carmen Jeronimo) 255–292 (Academic Press, 2020).
36 Vuu, Y. M., Roberts, C.-T. & Rastegar, M. MeCP2 Is an Epigenetic Factor That Links
DNA Methylation with Brain Metabolism. *International Journal of Molecular Sciences*
24, 4218 (2023). https://doi.org/10.3390/ijms24044218
37 You, E. *et al.* LINE-1 ORF1p Mimics Viral Innate Immune Evasion Mechanisms in
Pancreatic Ductal Adenocarcinoma. *Cancer Discovery* **15**, 1063–1082 (2025).
https://doi.org/10.1158/2159-8290.cd-24-1317
38 Zhang, X., Zhang, R. & Yu, J. New Understanding of the Relevant Role of LINE-1
Retrotransposition in Human Disease and Immune Modulation. *Front Cell Dev Biol* **8**
(2020). https://doi.org/10.3389/fcell.2020.00657
39 Evans, T. A. & Erwin, J. A. Retroelement-derived RNA and its role in the brain. *Semin*
Cell Dev Biol **114**, 68–80 (2021). https://doi.org/10.1016/j.semcdb.2020.11.001
40 Jones, R. B. *et al.* Nucleoside analogue reverse transcriptase inhibitors differentially
inhibit human LINE-1 retrotransposition. *PLoS One* **3**, e1547 (2008).
https://doi.org/10.1371/journal.pone.0001547
41 Dai, L., Huang, Q. & Boeke, J. D. Effect of reverse transcriptase inhibitors on LINE-1
and Ty1 reverse transcriptase activities and on LINE-1 retrotransposition. *BMC*
Biochem **12**, 18 (2011). https://doi.org/10.1186/1471-2091-12-18
42 Pitcher, M. R. *et al.* Insulinotropic treatments exacerbate metabolic syndrome in
mice lacking MeCP2 function. *Hum Mol Genet* **22**, 2626–2633 (2013).
https://doi.org/10.1093/hmg/ddt111
43 Sobanski, T. *et al.* The fructose-bisphosphate, Aldolase A (ALDOA), facilitates DNA-
PKcs and ATM kinase activity to regulate DNA double-strand break repair. *Sci Rep* **13**,
15171 (2023). https://doi.org/10.1038/s41598-023-41133-1
44 Bogerd, H. P. *et al.* Cellular inhibitors of long interspersed element 1 and Alu
retrotransposition. *Proc Natl Acad Sci U S A* **103**, 8780–8785 (2006).
https://doi.org/10.1073/pnas.0603313103
45 Tam, O. H., Ostrow, L. W. & Gale Hammell, M. Diseases of the nERVous system:
retrotransposon activity in neurodegenerative disease. *Mob DNA* **10**, 32 (2019).
https://doi.org/10.1186/s13100-019-0176-1
46 Roy, N. *et al.* Elevated expression of the retrotransposon LINE-1 drives Alzheimer's
disease-associated microglial dysfunction. *Acta Neuropathol* **148**, 75 (2024).
https://doi.org/10.1007/s00401-024-02835-6
47 Zhang, W. *et al.* Retrotransposon: an insight into neurological disorders from
perspectives of neurodevelopment and aging. *Transl Neurodegener* **14**, 14 (2025).
https://doi.org/10.1186/s40035-025-00471-y

1501	48	Peze-Heidsieck, E. *et al.* Retrotransposons as a Source of DNA Damage in
1502		Neurodegeneration. *Front Aging Neurosci* **13**, 786897 (2021).
1503		https://doi.org/10.3389/fnagi.2021.786897
1504	49	Ravel-Godreuil, C., Znaidi, R., Bonnifet, T., Joshi, R. L. & Fuchs, J. Transposable
1505		elements as new players in neurodegenerative diseases. *FEBS Lett* **595**, 2733–2755
1506		(2021). https://doi.org/10.1002/1873-3468.14205
1507	50	Reid, M. A., Dai, Z. & Locasale, J. W. The impact of cellular metabolism on chromatin
1508		dynamics and epigenetics. *Nat Cell Biol* **19**, 1298–1306 (2017).
1509		https://doi.org/10.1038/ncb3629
1510	51	Xu, Q. *et al.* SETD2 regulates the maternal epigenome, genomic imprinting and
1511		embryonic development. *Nat Genet* **51**, 844–856 (2019).
1512		https://doi.org/10.1038/s41588-019-0398-7
1513	52	Jachowicz, J. W. *et al.* LINE-1 activation after fertilization regulates global chromatin
1514		accessibility in the early mouse embryo. *Nat Genet* **49**, 1502–1510 (2017).
1515		https://doi.org/10.1038/ng.3945
1516	53	Terry, D. M. & Devine, S. E. Aberrantly High Levels of Somatic LINE-1 Expression and
1517		Retrotransposition in Human Neurological Disorders. *Front Genet* **10**, 1244 (2019).
1518		https://doi.org/10.3389/fgene.2019.01244
1519	54	Garrett-Bakelman, F. E. *et al.* The NASA Twins Study: A multidimensional analysis of a
1520		year-long human spaceflight. *Science* **364** (2019).
1521		https://doi.org/10.1126/science.aau8650
1522	55	Indo, H. P. *et al.* Changes in mitochondrial homeostasis and redox status in
1523		astronauts following long stays in space. *Sci Rep* **6**, 39015 (2016).
1524		https://doi.org/10.1038/srep39015
1525	56	Waisberg, E. *et al.* Mitochondrial dysfunction in Spaceflight Associated Neuro-Ocular
1526		Syndrome (SANS): a molecular hypothesis in pathogenesis. *Eye (Lond)* **38**, 1409–1411
1527		(2024). https://doi.org/10.1038/s41433-024-02951-3
1528	57	da Silveira, W. A. *et al.* Comprehensive Multi-omics Analysis Reveals Mitochondrial
1529		Stress as a Central Biological Hub for Spaceflight Impact. *Cell* **183**, 1185–1201 e1120
1530		(2020). https://doi.org/10.1016/j.cell.2020.11.002
1531	58	Hroudova, J., Singh, N. & Fisar, Z. Mitochondrial dysfunctions in neurodegenerative
1532		diseases: relevance to Alzheimer's disease. *Biomed Res Int* **2014**, 175062 (2014).
1533		https://doi.org/10.1155/2014/175062
1534	59	Area-Gomez, E., Guardia-Laguarta, C., Schon, E. A. & Przedborski, S. Mitochondria,
1535		OxPhos, and neurodegeneration: cells are not just running out of gas. *J Clin Invest*
1536		**129**, 34–45 (2019). https://doi.org/10.1172/JCI120848
1537	60	Jones, C. W. *et al.* Molecular and physiological changes in the SpaceX Inspiration4
1538		civilian crew. *Nature* **632**, 1155–1164 (2024). https://doi.org/10.1038/s41586-024-
1539		07648-x
1540	61	Ali, N., Beheshti, A. & Hampikian, G. Space exploration and risk of Parkinson's
1541		disease: a perspective review. *NPJ Microgravity* **11**, 1 (2025).
1542		https://doi.org/10.1038/s41526-024-00457-6
1543	62	Pecorelli, A., Cervellati, C., Cordone, V., Hayek, J. & Valacchi, G. Compromised
1544		immune/inflammatory responses in Rett syndrome. *Free Radic Biol Med* **152**, 100–
1545		106 (2020). https://doi.org/10.1016/j.freeradbiomed.2020.02.023

63   Pecorelli, A. *et al.* Altered inflammasome machinery as a key player in the
perpetuation of Rett syndrome oxinflammation. *Redox Biol* **28**, 101334 (2020).
https://doi.org/10.1016/j.redox.2019.101334
64   Frost, B. & Dubnau, J. The Role of Retrotransposons and Endogenous Retroviruses in
Age-Dependent Neurodegenerative Disorders. *Annu Rev Neurosci* **47**, 123–143
(2024). https://doi.org/10.1146/annurev-neuro-082823-020615
65   Mangoni, D. *et al.* From the genome's perspective: Bearing somatic
retrotransposition to leverage the regulatory potential of L1 RNAs. *Bioessays* **47**,
e2400125 (2025). https://doi.org/10.1002/bies.202400125
66   McEntee, C. M. & LaRocca, T. J. Single-Cell Transcriptome Patterns of Transposable
Elements in Alzheimer's Disease. *Mol Neurobiol* (2025).
https://doi.org/10.1007/s12035-025-05140-9
67   LaRocca, T. J., Cavalier, A. N. & Wahl, D. Repetitive elements as a transcriptomic
marker of aging: Evidence in multiple datasets and models. *Aging Cell* **19**, e13167
(2020). https://doi.org/10.1111/acel.13167
68   Baune, B. T. *et al.* Interleukin-6 gene (IL-6): a possible role in brain morphology in the
healthy adult brain. *J Neuroinflammation* **9**, 125 (2012).
https://doi.org/10.1186/1742-2094-9-125
69   Zhao, Z. *et al.* Plasma IL-6 levels and their association with brain health and dementia
risk: A population-based cohort study. *Brain Behav Immun* **120**, 430–438 (2024).
https://doi.org/10.1016/j.bbi.2024.06.014
70   Camera, A. *et al.* Aging and putative frailty biomarkers are altered by spaceflight. *Sci
Rep* **14**, 13098 (2024). https://doi.org/10.1038/s41598-024-57948-5
71   Mao, W. *et al.* Microgravity Inhibits Cell Proliferation and Promotes Senescence and
Apoptosis in Embryonic Stem Cells. *Space: Science & Technology* **4** (2024).
https://doi.org/10.34133/space.0104
72   Pham, T. P., Kremer, V. & Boon, R. A. RNA-based therapeutics in cardiovascular
disease. *Curr Opin Cardiol* **35**, 191–198 (2020).
https://doi.org/10.1097/HCO.0000000000000724
73   Desai, R. I. *et al.* Complex 33-beam simulated galactic cosmic radiation exposure
impacts cognitive function and prefrontal cortex neurotransmitter networks in male
mice. *Nat Commun* **14**, 7779 (2023). https://doi.org/10.1038/s41467-023-42173-x
74   Oyefeso, F. A. *et al.* Effects of acute low-moderate dose ionizing radiation to human
brain organoids. *PLoS One* **18**, e0282958 (2023).
https://doi.org/10.1371/journal.pone.0282958
75   Chen, Q., Sun, L. & Chen, Z. J. Regulation and function of the cGAS–STING pathway of
cytosolic DNA sensing. *Nature Immunology* **17**, 1142–1149 (2016).
https://doi.org/10.1038/ni.3558
76   Zhen, Z. *et al.* Nuclear cGAS restricts L1 retrotransposition by promoting TRIM41-
mediated ORF2p ubiquitination and degradation. *Nature Communications* **14** (2023).
https://doi.org/10.1038/s41467-023-43001-y
77   Gulen, M. F. *et al.* cGAS–STING drives ageing-related inflammation and
neurodegeneration. *Nature* **620**, 374–380 (2023). https://doi.org/10.1038/s41586-
023-06373-1
78   Dvorkin, S., Cambier, S., Volkman, H. E. & Stetson, D. B. New frontiers in the cGAS-
STING intracellular DNA-sensing pathway. *Immunity* **57**, 718–730 (2024).
https://doi.org/10.1016/j.immuni.2024.02.019

79   Colombo, A. R., Elias, H. K. & Ramsingh, G. Senescence induction universally activates
transposable element expression. *Cell cycle (Georgetown, Tex* **17**, 1846–1857 (2018).
https://doi.org/10.1080/15384101.2018.1502576
80   Merenciano, M., Larue, A., Garambois, C., Nunes, W. V. B. & Vieira, C. Exploring the
Relationship of Transposable Elements and Ageing: Causes and Consequences.
Genome Biol Evol **17** (2025). https://doi.org/10.1093/gbe/evaf088
81   Crucian, B. E. *et al.* Countermeasures-based Improvements in Stress, Immune System
Dysregulation and Latent Herpesvirus Reactivation onboard the International Space
Station - Relevance for Deep Space Missions and Terrestrial Medicine. *Neurosci*
Biobehav Rev **115**, 68–76 (2020). https://doi.org/10.1016/j.neubiorev.2020.05.007
82   Crucian, B. *et al.* Terrestrial stress analogs for spaceflight associated immune system
dysregulation. *Brain Behav Immun* **39**, 23–32 (2014).
https://doi.org/10.1016/j.bbi.2014.01.011
83   Stowe, R. P., Sams, C. F. & Pierson, D. L. Adrenocortical and immune responses
following short- and long-duration spaceflight. *Aviat Space Environ Med* **82**, 627–634
(2011). https://doi.org/10.3357/asem.2980.2011
84   Mehta, S. K. *et al.* Latent virus reactivation in astronauts on the international space
station. *NPJ Microgravity* **3**, 11 (2017). https://doi.org/10.1038/s41526-017-0015-y
85   McGregor, H. R. *et al.* Impacts of spaceflight experience on human brain structure.
Sci Rep **13**, 7878 (2023). https://doi.org/10.1038/s41598-023-33331-8
86   Zhang, Z. N. *et al.* Layered hydrogels accelerate iPSC-derived neuronal maturation
and reveal migration defects caused by MeCP2 dysfunction. *Proc Natl Acad Sci U S A*
113, 3185–3190 (2016). https://doi.org/10.1073/pnas.1521255113
87   Marchetto, M. C. *et al.* A model for neural development and treatment of Rett
syndrome using human induced pluripotent stem cells. *Cell* **143**, 527–539 (2010).
https://doi.org/10.1016/j.cell.2010.10.016
88   Serio, A. *et al.* Astrocyte pathology and the absence of non-cell autonomy in an
induced pluripotent stem cell model of TDP-43 proteinopathy. *Proc Natl Acad Sci U S*
A **110**, 4697–4702 (2013). https://doi.org/10.1073/pnas.1300398110
89   Wickham, H. *Ggplot2 : elegant graphics for data analysis*. (Springer, 2009).
90   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498–2504 (2003).
https://doi.org/10.1101/gr.1239303
91   Szklarczyk, D. *et al.* The STRING database in 2023: protein-protein association
networks and functional enrichment analyses for any sequenced genome of interest.
Nucleic Acids Res **51**, D638–D646 (2023). https://doi.org/10.1093/nar/gkac1000
92   Assarsson, E. *et al.* Homogenous 96-plex PEA immunoassay exhibiting high
sensitivity, specificity, and excellent scalability. *PLoS One* **9**, e95192 (2014).
https://doi.org/10.1371/journal.pone.0095192
93   Nageshappa, S. *et al.* Altered neuronal network and rescue in a human MECP2
duplication model. *Mol Psychiatry* **21**, 178–188 (2016).
https://doi.org/10.1038/mp.2015.128
94   Guy, J., Hendrich, B., Holmes, M., Martin, J. E. & Bird, A. A mouse Mecp2-null
mutation causes neurological symptoms that mimic Rett syndrome. *Nat Genet* **27**,
322–326 (2001). https://doi.org/10.1038/85899
1638

P

GS GLT1 GLAST SLC26A7 SLC38A1

**** ** ****

Fold change

□ Control □ MECP2-KO

Q

Neurons   CTIP2+   EGFP+

Control   MECP2-KO   MECP2-KO RTi   MECP2-KO NVP

100 μm

R

Dendrite Length (μm)

**** * **

Control MECP2-KO MECP2-KO + RTi MECP2-KO + NVP

S

Dendrite complexity

**** **

Control MECP2-KO MECP2-KO + RTi MECP2-KO + NVP

A   WT   Mecp2-KO   Mecp2-KO + 3TC

B   WT   Mecp2-KO   Mecp2-KO + 3TC

closed

open

C   WT   Mecp2-KO   Mecp2-KO + 3TC

D

WT

Mecp2-KO

Mecp2-KO + 3TC

F

carboxylic acid metabolic process
oxoacid metabolic process
carbohydrate metabolic process
negative regulation of intracellular signal transduction
Wnt signaling pathway
cell-cell signaling by wnt
generation of precursor metabolites and energy
hexose metabolic process
monosaccharide metabolic process
small molecule biosynthetic process
response to oxygen levels
response to hypoxia
positive regulation of small molecule metabolic process
small molecule catabolic process
cellular response to oxygen levels
cellular response to hypoxia
negative regulation of transmembrane transport
somatic stem cell population maintenance
gluconeogenesis
hexose biosynthetic process
glycolytic process
ATP generation from ADP
regulation of fatty acid metabolic process
nucleoside diphosphate phosphorylation
nucleotide phosphorylation
positive regulation of gluconeogenesis
positive regulation of transcription from RNA polymerase II promoter involved in cellular response to chemical stimulus
regulation of transcription from RNA polymerase II promoter in response to oxidative stress
response to interleukin-7
cellular response to interleukin-7

GeneRatio

Count
● 10
● 20
● 30

p.adjust
0.03
0.02
0.01

G

ISS Upregulated   26 | 4 | 169   MeCP2-KO Astrocyte Upregulated

p = 0.01871

H

Eno1
Jak1
Stip1
P4hb
Tcf7l2
Tfrc
Sox9
Nr4a2
Srebf1
Cbs
Cadps2
Sesn1
Tfeb
Atf4
Rxra
Nfe2l2
Akt1
Foxo1

Mecp2-KO-1
Mecp2-KO-2
Mecp2-KO-3
Mecp2-KO + 3TC-1
Mecp2-KO + 3TC-2
Mecp2-KO + 3TC-3
WT-1
WT-2
WT-3

I

WT   Mecp2-KO   Mecp2-KO + 3TC

C3

S100β / Hoechst

J

Normalized S100β fluorescence intensity

****   ****

WT   Mecp2-KO   Mecp2-KO + 3TC