Calibrated analysis framework for nanopore direct RNA sequencing uncovers

cell-specific m⁶A stoichiometry at conserved sites

Authors

Denise Ohnezeit*[1], Elene Loliashvili*[2], Gregory Putzel[3], Ruth Verstraten[2,4], Jianheng Liu[5], Luke S.

Nicholson[5], Alejandro Pironti[3], Samie R. Jaffrey[5], Daniel P. Depledge[#1,2,4,6], Angus C. Wilson[#1]

Affiliations

[1]Department of Microbiology, New York University School of Medicine, New York, NY, USA

[2]Institute of Virology, Hannover Medical School, Hannover, Germany

[3]Antimicrobial-Resistant Pathogens Program, New York University Grossman School of Medicine, New

York, NY, USA

[4]German Center for Infection Research (DZIF), partner site Hannover-Braunschweig, Hannover, Germany

[5]Department of Pharmacology, Weill Medical College, Cornell University, New York, NY, USA

[6]Cluster of Excellence RESIST (EXC 2155), Hannover Medical School, Carl-Neuberg-Straße 1, 30625

Hannover, Germany

* Joint first authors: contributed equally

[#] Joint senior authors: contributed equally

ORCIDs

DO = 0000-0002-2346-2545

EL = 0009-0006-4098-5010

GP = 0000-0002-5211-2608

RV = 0009-0008-6046-6335

JL = 0000-0003-0216-1951

LSN = 0009-0005-8203-7249

AP= 0000-0002-7759-6776

SRJ = 0000-0003-3615-6958

DPD = 0000-0002-4292-0599

ACW = 0000-0002-5016-4164

Abstract

Nanopore direct RNA sequencing (DRS) coupled with Dorado modification-aware basecalling enables

mapping of epitranscriptomic modifications including N^6-methyladenosine (m^{6}A) at the level of individual

RNAs. However, a lack of systematic benchmarking continues to raise questions regarding the sensitivity,

specificity, and reproducibility of this method. To address this and to establish a best-practice workflow,

we evaluated multiple Dorado versions using *in vitro* transcribed RNA and an m^{6}A methyltransferase

inhibitor as specificity controls. We established that stringent filtering is necessary to reduce false-positive

calls and found strong concordance at high-stoichiometry sites when compared to an orthogonal m^{6}A

mapping method (GLORI). Further, by applying DRS to human primary fibroblasts and HD10.6 neurons, we

uncovered cell type-specific differences in m^{6}A stoichiometry, indicating a finely tuned epitranscriptomic

regulation. Our study thus presents the first systematic comparison of Dorado and GLORI from the same

input RNA and expands characterization of the m^{6}A epitranscriptome to fibroblasts and neurons.

Introduction

Epitranscriptomics describes the occurrence of chemical modifications on RNA that regulate stability,

localization, splicing, translation or decay (Gilbert and Nachtergaele 2023). To date, over 170 RNA

modifications are known to occur across all classes of RNA including eukaryotic mRNA (Cappannini et al.

2023). Of these, at least thirteen different chemical modifications have been characterized in mRNAs,

comprising both cap-proximal and internal modifications (Anreiter et al. 2021). By contrast, current

evidence indicates that lncRNAs harbor a more limited repertoire of modifications, primarily N^6-

methyladenosine (m^{6}A), 5-methylcytosine (m^{5}C), and N7-methylguanosine (m^{7}G) (Yang et al. 2024a). One

of the best studied and most abundant internal mRNA modifications is m^{6}A. Functional studies of m^{6}A

methylases, demethylases and m^{6}A binding proteins, as well as sequencing-based mapping methods have

increased our understanding of m^{6}A as a reversible and dynamic modification that fine-tunes nearly all

aspects of mRNA metabolism (Zaccara et al. 2019; He and He 2021; Murakami and Jaffrey 2022; Gilbert

and Nachtergaele 2023; Sendinc and Shi 2023).

In addition to m^{6}A, other widespread and functionally important marks include m^{5}C, which is less

abundant but influences mRNA export and stability, pseudouridine (Ψ), which contributes to mRNA

stability, translation and splicing, and adenosine-to-inosine (A to I) editing, which is involved in splicing,

stability and recoding events that contribute to protein diversity (Sun et al. 2023).

Techniques based on short-read (Illumina) sequencing have advanced in recent years to achieve single-

nucleotide resolution and quantitative mapping of RNA modifications. The approaches that have emerged

to map m^{6}A can be broadly classified as antibody-based, enzyme-assisted or chemical conversion methods

(Moshitch-Moshkovitz et al. 2024).

Early antibody-based techniques (Dominissini et al. 2012; Meyer et al. 2012; Linder et al. 2015) provided

the first transcriptome-wide insights into m^{6}A architecture, albeit at limited resolution. Enzyme-assisted

methods (Meyer 2019; Xiao et al. 2023) have improved precision but can introduce enzymatic biases.

These biases are avoided with GLORI-Sequencing (GLORI), which selectively deaminates unmethylated

adenosines, enabling theoretically unbiased, quantitative, single-nucleotide mapping of m^{6}A (Liu et al.

2023).

Despite advances, the broad utility of these approaches is limited by a lack of isoform specificity, as well

as biases associated with amplification, specificity of antibodies or variable adenosine conversion rates

(Diensthuber and Novoa 2025). Oxford Nanopore Technologies (ONT) direct RNA sequencing (DRS) offers

an alternative to these methods by analyzing native RNA molecules that have retained their modifications

(Garalde et al. 2018). As single RNAs pass through nanopores, signal changes are interpreted to determine

canonical and modified residues at single-nucleotide resolution on individual transcripts (Jain et al. 2016;

Viehweger et al. 2019). For the now discontinued DRS SQK-RNA002 chemistry used with R9 flowcells,

multiple research groups developed algorithms for detecting RNA modifications. These methods can be

divided into error rate-based and current signal-based (Abebe et al. 2022). While the recent release of the

DRS SQK-RNA004 chemistry and RNA-specific flowcells reduced the utility of many of these tools,

introduction of a modification-aware basecaller (Dorado) provides an all-in-one toolkit for basecalling,

RNA modification detection, and poly(A) tail length estimation. Early versions of Dorado (e.g. v0.5.0 and

v0.6.0) enabled m^{6}A (DRACH-context only) detection, while version v0.7.0 enabled detection of Ψ and all-

context m^{6}A. More recent versions (v0.8.0 and v0.9.0) allow both DRACH- and all-context m^{6}A detection,

together with Ψ, inosine and m^{5}C (Zou et al. 2025).

In a recent overview of this emerging technology, Cruciani and Novoa highlighted the importance of

rigorous evaluation of the limitations of Dorado and advocated forcefully for the establishment of best

practices to ensure consistency and accuracy going forward (Cruciani et al. 2025). To date, only a handful

of studies of this kind have been published. A series of studies from Novoa and colleagues showed that

Dorado performs robustly on synthetic modified and unmodified RNA molecules but also found that

overlaps in the identification of specific modifications and false-positive calls remain a challenge but can

be mitigated by the inclusion of *in vitro* transcribed (IVT) controls (Cruciani et al. 2025; Diensthuber and

Novoa 2025). Independently, Esfahani and colleagues used poly(A) RNA from a B-lymphocyte cell line to

compare SQK-RNA002 and SQK-RNA004-derived DRS data. While they concluded that Dorado

outperformed other RNA modification detection tools, IVT controls were necessary to reduce false-

positive calls (Esfahani et al. 2025). Recent benchmarking efforts comparing Dorado and m6Anet on SQK-

RNA004 chemistry have shown that both tools achieve high recall and strong stoichiometry correlations

at DRACH motifs but exhibit higher false discovery rates on IVT controls and lower recalls of low-

stoichiometry genomic m^{6}A sites. However, these evaluations relied on previously published GLORI and

eTAM-Seq datasets, rather than matched input material (Diensthuber and Novoa 2025; Zou et al. 2025).

By directly comparing m^{6}A detection by DRS coupled with Dorado and GLORI on the same RNA inputs, our

study overcomes this limitation. Specifically, we systemically benchmark multiple versions of Dorado using

poly(A) RNA from primary (normal) human dermal fibroblasts (NHDFs) and functional human neurons

produced through terminal differentiation of HD10.6 cells (Raymon et al. 1999), along with IVT RNA. We

included treatments with STM2457, an enzymatic inhibitor of the primary m^{6}A methyltransferase METTL3

(Yankova et al. 2021), to robustly assess false-positive m^{6}A detection rates and also compared the DRS

outputs with those obtained from GLORI (Yang et al. 2024b) using the same RNA as input. While our

analysis is centered on the sensitivity and specificity of m^{6}A detection in DRS datasets, we also benchmark

the ability of Dorado to reliably detect m^{5}C, Ψ, and inosine. Together, our data provides a robust workflow

for DRS-based modification detection that reliably captures transcriptome-wide, isoform-level, site-

specific stoichiometries of RNA modifications.By expanding m^{6}A mapping on primary human fibroblasts

and functional human neurons, we uncover mostly conserved m^{6}A profiles but reveal increased site-

specific m^{6}A stoichiometries in neuronal cells.

Results

Systematic benchmarking of ONT's Dorado across human cell types.

To evaluate the performance of ONT's basecalling tool Dorado across different human cell types, we selected primary NHDFs and differentiated HD10.6 cells as a neuron-like model. Prior to isolation of total RNA, cells were treated for 48h with a small molecule inhibitor of METTL3 (STM2457) or with vehicle (DMSO). We then isolated poly(A) RNA and subjected the same input material to both DRS and GLORI. The same material was also used to generate IVT RNA (**Fig. 1A**). Dorado basecalling using version v.0.9.0 resulted in over 13M reads using poly(A) RNA from NHDFs treated with DMSO or STM2457, while we obtained over 5M and 10M reads from DMSO- and STM2457-treated HD10.6 cells, respectively. The rate of primary alignment to the human genome (hg38) was over 98% for all the sequenced samples. A detailed overview of read statistics generated by DRS can be found in **Supplementary Table 1**.

To systematically benchmark Dorado, we performed basecalling using versions v0.5.0 through v0.9.0. Reads were aligned to the human genome and transcriptome to obtain comprehensive information on isoform-level modification detection. Further, we explored different aspects of detected modifications using the outputs generated by ONT's Modkit tool. We assessed multiple filtering strategies using base and modification probabilities, as well as stoichiometry distributions, and evaluated them based on the IVT data and sensitivity to METTL3 inhibition. By filtering out false-positive calls using multiple control layers, we determined high-confidence m^{6}A sites (**Fig. 1B**). Finally, we compared these high-confidence predictions with sites identified by GLORI performed on the same input RNA. Using these results, we developed a benchmarked pipeline that filters m^{6}A sites predicted by Dorado to minimize false-positive detection without impacting sensitivity (**Fig. 1C**).

Stringent filtering of modification probabilities removes false positives

During modification-aware basecalling, each nucleotide is assigned a basecall probability score ranging from 0 to 1. Similarly, individual modified nucleotides are assigned a modification probability score in the same range. To establish a baseline for false-positive m^{6}A predictions, we plotted basecall and modification probability distributions for an IVT poly(A) RNA dataset processed using Dorado v0.9.0 and aligned to human genome (**Fig.2**). We binned basecall and modification probability values for A and m^{6}A (all-context and DRACH) and compared this data from DMSO- and STM2457-treated NHDF datasets.

We observed that 75-90% of A basecalls (in all datasets) have probability scores above 0.8 **(Fig. S1A)**. Since

all A bases are considered during all-context m^6A calling, the fraction of m^6A sites is substantially smaller

when compared to DRACH-context m^6A sites (**Fig.2A,B, Fig.S1B,C**). The modification probability

distributions are effectively identical for DMSO-treated, STM2457-treated and IVT datasets below 0.98,

while the profiles differ substantially at probabilities ≥ 0.98. This suggests that sites detected below 0.98

are likely false positives, indicating that only high-probability m^6A calls can be considered reliable. Here,

STM2457 treatment shows a marked reduction in the number of reported m^6A sites in both all-context

and DRACH-only datasets when compared to the DMSO treatment (**Fig. 2A,B**).

In addition to m^6A, we examined modification probability distributions for Ψ, inosine and m^5C, each of

which can be basecalled with Dorado v0.9.0 (**Fig. S2**). For Ψ and I, the modification probability distributions

for the STM2457 and DMSO datasets only diverge from the IVT dataset at ≥ 0.99 modification probability,

thus providing a relevant threshold for filtering out false positives (**Fig. S2A,B**). By contrast, m^5C

modification probability distributions were similar between all three datasets and no filter could be

adequately established (**Fig. S2C**). While this likely reflects a low rate of m^5C installation, it also indicates

that the m^5C model utilized by Dorado requires further refinement.

Modification probability distributions have similar profiles for transcriptome-aligned NHDF data (**Fig. S3**)

as well as genome- and transcriptome-aligned data from HD10.6 cells (**Fig. S4, Fig. S5**). Of note, the fraction

of high-probability inosine sites is increased in STM2457-treated cells, compared to DMSO-treated ones

in genome-aligned data (**Fig. S2B, Fig.S4D**), which points to increased numbers of retained introns in the

METTL3-inhibited sample. Accordingly, this effect is absent in transcriptome-aligned data (**Fig. S3D, Fig.**

S5D).

While IVT transcriptomes serve as excellent negative controls, we also considered whether the RNA

calibration strand (RCS) that is added during DRS library preparation could serve a similar function. The

RCS is a synthetic Enolase II RNA derived by IVT[13] that, similarly to the IVT transcriptomes, are devoid of

RNA modifications. Despite the reduced diversity of sequence in the RCS data, the overall results were

comparable to that from the IVT transcriptome data (**Fig. S6A,B, Fig. 2A,B**), demonstrating that the RCS

can be used as a "built-in" negative control for modification detection with DRS. Finally, we determined

that basecall and modification probability distributions remain generally consistent between Dorado

versions although more m^6A and Ψ sites are detected with high modification probability in Dorado versions

v0.8.0/v0.9.0 compared to v0.7.0 (**Fig. S6C-E**).

Filtering by stoichiometry further eliminates potential false positives

We next examined the stoichiometry distributions for all m^{6}A sites in both unfiltered and filtered NHDF

datasets (**Fig. S7**). Here, unfiltered datasets showed that the vast majority of m^{6}A sites, 92% for all-context

and 75% for DRACH-context, have stoichiometry values < 10% with less than 1% sites showing high (> 50%)

stoichiometry. Application of our filtering strategy removed over 70% of the sites (**Fig. 2E**), regardless of

the context, almost all of which showed stoichiometries of < 10%, without significantly changing the

numbers of high stoichiometry sites detected (**Fig. S7**). A similar analysis of the IVT transcriptome revealed

large numbers of false positive m^{6}A sites with stoichiometry values below 10%, most of which were

eliminated by application of our filters (**Fig. S8A,B**). However, the remaining presence of low stoichiometry

false-positive sites indicates that an additional layer of filtering to exclude sites with < 10% stoichiometry

is beneficial. An additional argument in support of our modification probability filtering strategy is the

observation that treatment with the METTL3 specific inhibitor STM2457 dramatically reduced m^{6}A

installation at sites with ≥ 10% stoichiometry in untreated datasets in a context-independent manner (**Fig.**

2C,D, Supplementary Table 2). Similar results were observed for the transcriptome aligned NHDF and

HD10.6 m^{6}A stoichiometry distributions as well (**Fig. S9, Fig. S10, Supplementary Table 2**).

To determine whether our filtering strategy could be applied to other RNA modifications detected by

Dorado, we generated stoichiometry distributions for Ψ and inosine in unfiltered and filtered (≥ 0.99

modification probability) human genome-aligned DMSO- and STM2457-treated NHDFs and HD10.6s. As

with m^{6}A (**Fig. S7**), 92% of detected Ψ and 96% of inosine have less than 10% stoichiometry (**Fig. S11, Fig.**

S12). A recent study that explores Ψ in human mRNAs using 2-bromoacrylamide-assisted cyclization

sequencing (BACS) corroborates our results, showing that 82% of Ψ modifications are detected at low

levels (below 20%) (Xu et al. 2024).

Above 10% stoichiometry, in our data, there are more inosine sites in STM2457-treated cells than DMSO-

treated cells, reflected also in the stoichiometry distributions (**Fig. S11, Fig. S12, Supplementary Table 2**).

Ψ and inosine stoichiometry distributions in our IVT data **(Fig.S8C,D)** further validates our filtering strategy

of ≥ 0.99 modification probability, as it substantially reduces the number of detected false-positive sites in

the unfiltered data even at higher (> 50%) stoichiometries in case of inosine (**Fig.S8D**).

We have thus demonstrated that setting thresholds for basecall modification probabilities provides a

robust method for identifying and filtering false-positive modification calls. For the purposes of this study,

we further evaluated the effect of placing the modification threshold on the modification data and

restricted the downstream analyses to m^{6}A sites with probability scores ≥ 0.98.

RNA modification detection rates differ between Dorado versions

Dorado basecalling relies on models generated using Remora (github.com/nanoporetech/remora) that are

frequently updated. This raises an important question as to whether these updates produce notably

different outputs that could impact biological interpretation. To address this, we utilized multiple Dorado

versions and Remora models to perform modification calling of our NHDF dataset. We restricted our

analyses to m^{6}A and Ψ as these are the only modifications that could be detected in previous versions.

Here, we observe that m^{6}A and Ψ stoichiometry distributions are generally consistent between different

Dorado versions (**Fig. S13**), albeit with greater numbers of low stoichiometry (< 10%) sites detected in

older Dorado versions. There is a large overlap of the detected ≥ 10%-stoichiometry m^{6}A (DRACH and all-

context) and Ψ sites between versions, although there are a lot of unique all-context m^{6}A and Ψ sites

identified with just v0.7.0 and not v0.8.0-v0.9.0, that are reduced after applying our modification

probability filter (**Fig. S14A,B**). Ψ sites uniquely detected with different versions are found in slightly

different sequence contexts (**Fig. S14C,D**). Thus, while RNA modification detection with Dorado is sensitive

to changes in the underlying models/versions, this can be effectively controlled through the application of

filters that only retain sites with high-confidence modification probability calls.

GLORI exhibits reduced specificity but converges with Dorado predictions at high stoichiometry sites

To assess the agreement of DRS Dorado-derived predictions with an orthogonal m^{6}A detection method,

we conducted GLORI on the same NHDF RNA preparations. To ensure a fair comparison of m^{6}A sites across

the two methods, we applied the following filtering strategies: For Dorado (DRACH model), this included

thresholds on modification probability (A ≥ 0.8, m^{6}A ≥ 0.98) and read coverage (≥ 20). For GLORI, we used

filters based on coverage (≥ 20) and statistical significance (p-value ≤ 0.05). Next, we extracted m^{6}A sites

that were present in both DMSO and STM2457 treated samples and removed all sites with ≤ 10%

stoichiometry in DMSO (**Fig. 3A**). This stepwise strategy reduced false positives and enabled a fair

comparison of the high confidence m^{6}A sites identified by both approaches.

As shown in the violin plots, both the all-context and DRACH-context Dorado models identify similar

numbers of m^{6}A sites (81,012 (all-context model) and 72,372 (DRACH model)), the stoichiometries of

which were strongly reduced upon STM2457 treatment (**Fig. 3B**). When we analyzed the motifs of the m^{6}A

sites detected with the all-context model, we observed nearly 9,000 sites outside of DRACH motifs.

Importantly, the total number of m^{6}A sites within DRACH motifs was nearly identical to the number of m^{6}A

sites reported using the DRACH model (**Fig.3B**). Unexpectedly, GLORI detects a substantially higher

number of sites when using all motifs (140,934) compared to DRACH motifs alone (27,926) (**Fig. 3C**). The

fact that most GLORI-identified sites fall outside DRACH motifs suggests these are false positives. This

interpretation is reinforced by the observation that the overlap between Dorado and GLORI datasets

remains virtually unchanged when comparing GLORI "all motif" and "DRACH motifs" (n=13,614) site sets,

underscoring that the additional non-DRACH sites in GLORI (n= 113,008) are largely spurious (**Fig. S15A**).

We obtained similar results when comparing m^{6}A sites obtained with Dorado and GLORI from HD10.6

cells. Overall, we detected a similar number of m^{6}A sites in HD10.6 cells using the all-context and DRACH-

specific Dorado models (77,728 and 67,222, respectively) (**Fig. S15B**). By contrast, we detected 174,282 vs

27,965 m^{6}A sites using GLORI without or with DRACH filtering, respectively (**Fig. S15C**).

To evaluate how stoichiometry thresholds affect the agreement between GLORI and Dorado, we compared

the overlap of high-confidence GLORI-identified sites with Dorado m^{6}A (DRACH context) calls across

increasing m^{6}A stoichiometry cutoffs (**Fig. 3D**). At the lowest stoichiometry threshold applied (≥ 10%),

13,614 sites overlapped, representing less than half (48.8%) of GLORI sites. However, as the stringency

increased, the relative overlap increased sharply: at ≥ 20%, nearly 70% of GLORI sites overlapped (11,536

sites), at ≥ 50% almost 90% overlapped (6,307 sites), and at the highest cutoff (≥ 80%), over 93% of GLORI

sites (3,119 sites) were also detected by Dorado. These results demonstrate that while GLORI detects a

broad set of potential m^{6}A sites, only the sites with high stoichiometries overlap with Dorado, indicating

high signal to noise ratios in the GLORI datasets.

GLORI relies on the high-efficiency conversion of adenosine to inosine with overall gene A to I conversion

rates needing to exceed 95-98% (Xu et al. 2024; Xie et al. 2025). Similarly, we observed the conversion

rates for our NHDF and HD10.6 datasets to be 95% (Supplementary **Table 3**), but we found that small

variations in gene conversion rates can strongly influence the sensitivity and specificity of GLORI, as only

high-stoichiometry m^{6}A sites overlapped with DRS-derived predictions (**Fig. 3D**).

DRS coupled with Dorado outperforms GLORI in isoform-specific m^{6}A detection

Next, we wanted to highlight the advantage of long-read sequencing in terms of mapping reads with

isoform-precision to the human transcriptome, which is not possible using GLORI due to its reliance on

short-read sequencing. We first confirmed that mapping Dorado-basecalled sites to the human

transcriptome results in the detection of similar numbers of m^{6}A sites in both NHDFs and HD10.6 cells (**Fig.**

S15D,E). To assess if DRS coupled with Dorado identifies distinct m^{6}A profiles on different transcript

isoforms, we analyzed the *SPEN* gene locus which was previously shown to contain more than 100 m^{6}A

sites (Liu et al. 2023). On a genomic level, we detected 42 m^{6}A sites using DRS, while GLORI only detected

5 m^{6}A sites (**Fig. 3E**). Out of the 7 total isoforms of *SPEN*, *SPEN-201* and *SPEN-202* were detectable in DRS-

derived data, considering a minimal coverage of 20 (**Supplementary Table 4**). When plotting the m^{6}A sites

detected by transcriptome-aligned DRS data, we found that the *SPEN-201* isoform exhibited more m^{6}A

sites, albeit with a lower median stoichiometry compared to the *SPEN-202* isoform (**Fig. 3F**). This example

highlights that DRS is superior compared with GLORI for isoform-level m^{6}A quantification.

Dorado profiling reveals cell type differences in m^{6}A distribution and stoichiometry

To assess the reproducibility of Dorado across DRS datasets derived from different cell types, we compared

the m^{6}A methylomes of NHDFs and HD10.6 cells. To ensure a fair comparison, we extracted genes that

were expressed in both cell types. Next, we compared the total numbers of high confidence m^{6}A sites in

NHDFs and HD10.6 cells within the genes expressed in both cell types and found that 87% of HD10.6-

derived m^{6}A sites overlap with those detected in NHDFs (**Fig. 4A**). Site-specific m^{6}A stoichiometries from

the shared gene set correlated strongly across cell types (Pearson r = 0.93), although we observed a global

shift in the HD10.6 dataset, where m^{6}A stoichiometries were generally higher than in NHDFs (**Fig. 4B**).

Plotting the m^{6}A stoichiometry distributions of the 38,217 m^{6}A sites shared between both cell types

showed a strong increase in the median m^{6}A stoichiometry of HD10.6 cells (70.2%), compared to NHDFs

(54%) (**Fig. 4C**).

We next aimed to conduct a comparative motif analysis of m^{6}A sites detected in NHDFs and HD10.6 cells.

Sequence logos confirmed strong enrichment of the canonical DRACH consensus motif for both cell types

in modification probability-filtered data, with the GGACU pentamer being the most prominent (**Fig. 4D,E,**

Fig.S16). Quantification of 5-mer distributions revealed highly similar motif preferences across datasets.

While the global motif architecture of m^{6}A deposition is conserved between NHDFs and HD10.6 cells, we

observed a trend toward higher stoichiometries in the HD10.6 dataset, particularly within the most

abundant DRACH motifs, suggesting enhanced modification efficiency in the neuron-like context (**Fig. 4F,**

G).

Metagene analysis showed, as known previously, that m^{6}A is enriched near the stop codons and in the 3'

UTRs (**Fig. S17A**). Several studies have examined the relationship between exon architecture and m^{6}A

distribution on mRNAs[1,2,3], and so we also examined m^{6}A sites in monoexonic transcripts separately and

found that monoexonic transcripts are more broadly m^{6}A-modified across entire CDS (**Fig. S17A**). m^{6}A

(both all-context and DRACH-context) distribution profiles look less typical for all-stoichiometry or ≤ 10%

stoichiometry sites in both NHDF and HD10.6 datasets (**Fig. S17B+C**). For all-context m^{6}A (**Fig. S17B**),

modification filter also proves to be deciding for the distribution profile, together with ≥ 10%

stoichiometry.

To connect shared m^{6}A sites with their functional context and illustrate their distribution on individual

transcripts, we conducted a KEGG pathway analysis with the genes that contain shared m^{6}A sites between

NHDFs and HD10.6 cells. We observed a significant enrichment in pathways related to neurodegeneration,

infection, and intracellular signaling, including axon guidance and mTOR signaling (**Fig. 4H**). To further

illustrate these findings, we examined representative genes contained in one or more of these pathways.

WNTB5, *MAPK1*, *AKT1*, *NCK1* and *NRP1* all showed robust methylation in both NHDFs and HD10.6 cells,

with consistently higher stoichiometries observed in the neuronal-like context. In contrast*, ACTG1*,

included as a housekeeping control not represented in the enriched pathways, displayed stable and

comparatively uniform methylation across both cell types (**Fig. 4I**).

Together, these results demonstrate that while the global architecture of m^{6}A deposition is largely

conserved between primary fibroblasts and neuronal-like cells, but unexpectedly the neuronal context is

characterized by a global shift toward higher m^{6}A stoichiometries. This difference is readily detectable in

single genes. These findings suggest that cell type-specific regulation of m^{6}A may fine-tune core cellular

pathways relevant to that cell type.

Discussion

High rates of false positives remain a major problem in epitranscriptomic mapping tools, even with the

advent of modification-aware basecalling models (Cruciani et al. 2025). Comparing the outputs of

orthogonal methods performed on the same input material, combined with use of IVT RNA and/or specific

methyltransferase inhibition, offers an effective mitigation strategy (Cruciani et al. 2025; Diensthuber and

Novoa 2025). The results presented here establish a robust framework that minimizes false positives and

enables isoform-specific and quantitative m^{6}A mapping. Through a direct comparison with GLORI, we

showed that combining DRS datasets with Dorado modification-aware basecalling provides greater

scalability, isoform resolution, and reproducibility across different cell types. Further, the simplicity of DRS

library preparation enhances consistency whereas GLORI is constrained by a laborious and technically

challenging protocol that is extremely sensitive to adenosine conversion rates (Shen et al. 2024). Despite

these limitations, we determined that the majority of m^{6}A sites in our filtered DRS datasets could be

confirmed by GLORI. By expanding our analysis across multiple human cell types, we further determine

that the sites of m^{6}A installation are conserved for the most part but that the stoichiometries show cell-

type specific differences.

Crucially, the inclusion of transcriptome-wide IVT datasets and the use of a selective METTL3 inhibitor

(STM2457) provides a robust approach to developing filters that dramatically decrease false-positive m^{6}A

detection. The impact of both approaches was readily visualized by plotting modification probability

distributions, enabling our determination that retaining only m^{6}A calls with confidence scores of 0.98 or

greater retained almost all STM2457-sensitive m^{6}A sites while eliminating the large numbers of false-

positive m^{6}A sites reported for the IVT datasets. This approach is supported by a recent study which

highlighted that Dorado introduces large numbers of false positives outside canonical DRACH contexts (Xie

et al. 2025). Our data confirms that in the unfiltered datasets, the number of false positives is higher in

the all-context model compared to the DRACH model. This may reflect the degenerate nature of the

canonical DRACH consensus and understudied differences in sequence preference between different cell

types and organisms. Retaining only sites at which m^{6}A stoichiometries exceeded 10% further reduced

false-positive rates while also focusing the results on identification of sites that are more likely to be

biologically significant. The combined effect of these filters was to provide a strong agreement between

the m^{6}A all-context and m^{6}A DRACH model in the detection of high confidence m^{6}A sites, as observed by

filtering for DRACH motifs from m^{6}A sites detected with the all-context model (**Fig.3B**).

Prior studies have emphasized the need for isoform-resolved m^{6}A quantification by demonstrating

widespread heterogeneity in m^{6}A levels between genes and their isoforms (Guo et al. 2025; Xie et al.

2025). Here, we showed that aligning DRS datasets to the human transcriptome reveals differential m^{6}A

installation patterns between isoforms, underscoring the importance of accurate, isoform-specific m^{6}A

quantification at single-molecule resolution and highlighting the advantage of DRS against short-read

sequencing approaches such as GLORI (**Fig. 3E,F**).

Recently, the ability of Dorado and m6Anet (Hendra et al. 2022) to identify m^{6}A in DRS datasets from 293T

and HeLa cells were systematically evaluated using IVT RNA and comparisons to published datasets

obtained by GLORI and eTAM-seq (Zou et al. 2025). In this case the authors found that Dorado (v0.7.2)

outperforms m6Anet but also advocated for the importance of applying filters to reduce false positive

calls. Likewise, we found that IVT RNA served as a valuable negative control to detect false positive calls.

Furthermore, using the RNA calibration strand (RCS) as a negative control provided a convenient and

simpler alternative to IVT RNA, yielding comparable probability distributions and thereby enabling a built-

in control for routine experiments (Viehweger et al. 2019). While m^{6}A detection was generally

reproducible across Dorado versions, model-specific differences are present, i.e. v0.7.0 produces

substantially more low-confidence all-context m^{6}A and Ψ calls than v0.8.0–0.9.0, highlighting that the

models remain under active development and that benchmarking should be repeated when switching to

a new model. Recent work further highlights that continued improvements in both nanopore sequencing

chemistry and basecalling models significantly enhance RNA modification detection accuracy, emphasizing

the importance of revisiting benchmarking and filtering strategies as models evolve (Hewel et al. 2025).

Focusing on other modifications that are detected by Dorado, we observe that inosine and Ψ detection

appears robust when applying similar filtering strategies as for m^{6}A, the latter supported by other studies

(Esfahani et al. 2025; Verstraten et al. 2025). For inosine, we observed an enrichment of high-probability

(> 0.99) inosine sites in genome-aligned data from STM2457-treated samples. Further studies are needed

to determine whether this reflects increased intron retention or misclassification of modifications by the

current inosine model. While DRS is the only approach that allows for robust isoform-specific m^{6}A

mapping, comparisons with orthogonal approaches remain essential to validate findings. When comparing

Dorado and GLORI using the same input RNA, we unexpectedly observed that only 50% of m^{6}A sites with

stoichiometries above 10% that were identified by Dorado were also present in the GLORI data although

increasing the stoichiometries led to increased overlap (**Fig.3D**). We attribute this to a sub-optimal gene-

wide adenosine conversion rate of 95% during GLORI, compared to the 98-99% rate reported in the original

study (Shen et al. 2024). Here, unconverted adenosines result in high signal-to-noise ratios which result in

lower sensitivity, explaining why the overlap with Dorado increases at higher-stoichiometry sites. This is

also reflected in the relatively high amount of false positive sites detected outside of DRACH motifs in the

GLORI datasets (**Fig. 3C**). The sensitivity of GLORI to small fluctuations thus limits its utility in comparison

to DRS. Whether this can be resolved with improvements to the GLORI methodology (Sun et al. 2025)

requires further study.

Prior analyses of m^{6}A stoichiometry distributions across immortalized human cell lines, including cancer

cell lines, have revealed broadly similar m^{6}A stoichiometry distributions (Guo et al. 2025). By contrast, our

analysis of primary fibroblasts (NHDFs) and neuronal cells (differentiated HD10.6 cells) revealed an overall

increase in m^{6}A stoichiometries in neurons compared to fibroblasts, especially in genes associated with

neuronal function. While further studies are warranted to establish the underlying mechanisms and

functional consequences of these differences, this aligns with the importance of m^{6}A in neuronal function
(Yu et al. 2021). This includes m^{6}A maps from synaptic and nuclear RNA isolated from mouse dorsal
hippocampus which showed that synapse-specific RNAs showed generally higher methylation levels (Sun
et al. 2025). Intriguingly, neuronal regulation pathways utilize differentially spliced isoforms (Su et al. 2018)
and a recent study revealed isoform-specific differences in m^{6}A sites and stoichiometries in the human
brain that are linked to m^{6}A machinery expression and cell type context (Gleeson et al. 2025).

In summary, using matched RNA samples from different experimental conditions and from different
human cell types analyzed with both Dorado and GLORI orthogonal methodologies, we establish a robust
pipeline for the precise and quantitative transcriptome-wide detection of m^{6}A in DRS datasets. Future
studies should aim at extending this approach using orthogonal sequencing approaches to benchmark
Dorado-based detection of Ψ, m^{5}C and inosine, as well as the newly released models for 2'O-methylation.

Methods

Cell culturing and STM2457 treatment

NHDFs (Lonza) were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco), supplemented with
5% heat-inactivated FBS (HI-FBS, Gibco) and 1% penicillin-streptomycin (PS, Lonza) under standard culture
conditions. The HD10.6 cell line was originally derived from fetal human dorsal root ganglia (DRG) and
immortalized by overexpression of the v-myc oncogene under tetracycline regulation. Upon addition of
tetracycline and neurotrophins, these cells can be differentiated into neuron-like cells that show typical
neuronal morphology and nociceptive properties(Raymon et al. 1999). For our experiments, we used
HD10.6 cells that have been matured for 10 days. For maintenance of undifferentiated HD10.6 cells (a gift
from Anna Cliffe, University of Virginia School of Medicine), T-75 flasks were coated with 17 µg/mL
fibronectin in PBS for 15 min at 37°C before adding the cell suspension. Cells were grown in Advanced
DMEM/F12 (Gibco), supplemented with 1x Neurocult SM1 (STEMCELL Technologies), 2 mM L-glutamine
(Gibco), 1x Primocin (Invivogen) and 10 ng/mL prostaglandin E1 (Sigma). 0.5 ng/mL bFGF (PeproTech) were
added freshly to each media change. For maturation, HD10.6 cells were seeded at a density of 25.000
cells/cm² in 6-well plates which were pre-coated overnight with 50 µg/mL poly-L-ornithine hydrobromide
(Sigma) in 0.5M borate buffer pH 8.5 (Boston Bioproducts), and 1 µg/mL fibronectin (overnight). After cell
attachment overnight, media was changed to Neurobasal media (Gibco), supplemented with 1x Neurocult
SM1, 2mM L-glutamine, 1x Primocin, 1 µg/mL doxycycline, 50 ng/mL NGF (Alomone Labs), and 25 ng/mL

CNTF, GDNF and NT-3 (PeproTech). Media was changed every three days until HD10.6 cells showed

complete neuronal morphology after 12 days. NHDFs and matured HD10.6 cells were treated with 30 µM

STM2457 (Sigma) or DMSO for 48h, harvested in TRIzol (Invitrogen), and stored at -80°C until RNA

isolation.

RNA isolation

RNA was extracted by the TRIzol method according to the manufacturer's instructions. Briefly, 0.2 mL

chloroform per 1 mL TRIzol were added and mixed by vortexing. After centrifugation for 30 min at 12,000xg

at 4°C, the aqueous phase was transferred to isopropanol (0.5 mL per 1 mL TRIzol), mixed and incubated

for 10 min at room temperature. Subsequently, the samples were centrifuged for 10 min at 12,000xg at

4°C and the RNA pellets were washed twice in 75% fresh ethanol. After air-drying for 5-10 min, RNA was

dissolved in nuclease-free H_2O and stored at -80°C until further processing. One third of each sample was

used for DRS and the remaining same-input material was processed for GLORI. For both DRS and GLORI,

poly(A) RNA was selected using the Dynabeads™ mRNA Purification Kit (Invitrogen), according to the

manufacturer's instructions.

GLORI-Sequencing

GLORI reactions were largely performed following the published protocol for RNA protection, deamination

and deprotection with several minor adjustments (Shen et al. 2024). Briefly, poly(A) RNA was fragmented

at 94°C for 2 min (NEB E6150S) and subjected to glyoxal protection, sodium-nitrite-mediated deamination

(750 mM $NaNO_2$ at 16°C for 8h and 4°C for 6h), and deprotection in triethylammonium acetate-formamide

buffer at 95°C for 10 min. Library construction was conducted using a ligation-based strategy as described

by Liu and colleagues incorporating 5'adapters with 11-nt UMIs (Liu et al. 2025).

Indexed libraries were amplified with NEBNext Dual Primers Set 2 (E7780S), purified using AMPure XP

beads, and validated on an Agilent TapeStation (D1000 High Sensitivity). Sequencing was performed on an

Illumina NovaSeq 6000 platform (150 bp paired-end mode; > 80 million reads per sample).

For data processing, we employed the same computational pipeline described in Liu et al (Liu et al. 2025),

but implemented within a Snakemake-based framework that automates the complete analysis starting

from raw fastq files (Köster and Rahmann 2012). Details on software versions, workflow availability, and

repository links are provided in the Data and code availability section below.

Generation of a transcriptome-wide IVT RNA library

442 We sequenced the IVT RNA from both NHDFs and HD10.6 cells along with the native poly(A) RNA to

443 evaluate potential false-positive predictions by Dorado. This was achieved by converting the poly(A) RNA

444 to cDNA using a poly(A) primer and a T7-incorporating strand switch primer and using the cDNA as a

445 template for IVT synthesis.

Nanopore Sequencing of poly(A) RNA

447 250-300 ng of isolated poly(A) RNA and 25 ng RNA calibration strand (RCS) were used as input for the

448 standard DRS SQK-RNA004 protocol. Resulting libraries were loaded onto individual PromethION RNA flow

449 cells before sequencing on the PromethION 2 Solo instrument for 48h.

Nanopore basecalling

Basecalling was performed using Dorado v0.5.0, v0.6.0, v0.7.0, v0.8.0 and v0.9.0 (ONT) in super accuracy

mode with modification detection enabled for m^6A (DRACH context) in v.0.5.0, 0.6.0, 0.8.0 and 0.9.0, m^6A

(all context) in v0.7.0, 0.8.0 and 0.9.0, Ψ in v0.7.0, v0.8.0 and v0.9.0, and inosine and m^5C in v.0.9.0. Each

dataset was processed with adapter trimming. Resulting unaligned BAM files were converted to FASTQ

files containing modification tags for every read in the FASTQ read headers, using SAMtools v1.18 (Li et al.

2009) [*samtools fastq -TMM,ML*].

Genome and transcriptome level alignments

For genome-level alignment, reads were aligned against a combined human genome reference

(GRCh38.p14), and the RCS gene sequence, using minimap2 v2.26 [*minimap2 -ax splice -k14 -y -uf --*

secondary=no]. For transcriptome-level alignment, the reads were aligned against the human

transcriptome (v47), obtained via GENCODE (Mudge et al. 2025) [*minimap2 -t 8 -ax map-ont -y -L -p 0.99*].

Using SAMtools v1.18, resulting genome- and transcriptome-aligned SAM files were converted to BAM

files retaining only primary alignments [*samtools view -b -F2308* and *samtools view -b -F2324,*

respectively]. BAM files were sorted [*samtools sort*] and indexed [*samtools index*].

Processing of RNA modification calls using Modkit

Probability distributions of the detected sites in human genome- and transcriptome-aligned reads were

generated using sample-probs subcommand from Modkit v0.4.1 [sample-probs --hist --only-mapped --

percentiles 0.1,0.2,0.3,0.4,0.5,0.6,0.7,0.8,0.85,0.9 -f 0.1]. For RCS reads -f value was increased to 0.25. The

resulting .tsv files were imported into RStudio (v. 2024.12.1+563) and plotted using a custom R(v. 4.4.2)

script (Dorado_mod_probability_distributions.R), with the following packages: dplyr (v1.1.4), ggplot2
(v3.5.1), reshape2 (v1.4.4), patchwork (v1.3.0).

To generate extended bedMethyl files with m^6A-modified and unmodified base counts from the reads
mapping against the potential modification sites, we used Modkit pileup subcommand with the thresholds
[--*filter-threshold A:0.8 --filter-threshold T:0.8 --filter-threshold C:0.8 --filter-threshold G:0.8 --motif A 0 --*
ref $REF] with and without modification threshold filter [--*mod-thresholds a:0.98*] to generate our
"filtered" and "unfiltered" datasets, respectively. Canonical nucleotide calls (A, C, G, U) were retained with
> 80% probability scores for all modification datasets. Modification thresholds were set at 98% for m^6A
and 99% for all the other modifications. --*motif* was adjusted to the base for which modification status
was examined, which further restricted the analysis to only the positions that matched canonical
nucleotides in the reference genome. The resulting sites were further filtered according to the read
number per site (≥ 20 reads) for all the downstream analysis.

Gene Ontology analysis

GO analysis was performed using DAVID (Huang da et al. 2009). Genes harboring m^6A sites were filtered
for genes that were generally present in both datasets. Next, they were divided into NHDF-specific,
HD10.6-specific, and shared categories, and enrichment was assessed for biological processes. GO terms
were considered significant at FDR ≤ 0.05 with a minimum of 20 genes per GO term. For plotting, the top
20 GO terms that contained the highest gene numbers were extracted from the analysis of the shared
gene sets.

Analysis and visualization of modification stoichiometries and distributions

Stoichiometry distribution histograms and metaplots were generated using Rstudio and R v4.4.2 with the
following libraries: data.table, patchwork, scales, tidyr, dplyr, ggplot2[4]. Access to scripts used for analysis
and visualization is detailed below.

Data and code availability

Raw fastq and pod5 datasets generated in this study are available via the European Nucleotide Archive
(ENA) under the accession number PRJEB101380.

R scripts used in the analysis and visualization of data presented herein are available in the
https://github.com/DepledgeLab/HOLDEN repository.

The Snakemake workflow implementing the GLORI analysis pipeline is available in the
https://github.com/gp-micro/glori repository.

Acknowledgements and funding

DPD is supported by a German Centre for Infection Research (DZIF) Associate Professorship and the NIAID

grants R01-AI170583 and R01-AI152543. DPD also receives funding from the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC

2155, project number 390874280. EL is supported by the NIAID grant R01-AI170583 as well as the

Hannover Biomedical Research School (HBRS) and the Center for Infection Biology (ZIB). DO is supported

by the Walter Benjamin Programme of the DFG, project 554758329. ACW is supported by NIAID grants

R01AI176335 and R01-AI170583.

References

https://github.com/nanoporetech/modkit.

Abebe JS, Verstraten R, Depledge DP. 2022. Nanopore-Based Detection of Viral RNA Modifications. *mBio* **13**: e03702-03721.

Anreiter I, Mir Q, Simpson JT, Janga SC, Soller M. 2021. New Twists in Detecting mRNA Modification Dynamics. *Trends Biotechnol* **39**: 72-89.

Cappannini A, Ray A, Purta E, Mukherjee S, Boccaletto P, Moafinejad SN, Lechner A, Barchet C, Klaholz Bruno P, Stefaniak F et al. 2023. MODOMICS: a database of RNA modifications and related information. 2023 update. *Nucleic Acids Research* **52**: D239-D244.

Cruciani S, Delgado-Tejedor A, Pryszcz LP, Medina R, Llovera L, Novoa EM. 2025. De novo basecalling of RNA modifications at single molecule and nucleotide resolution. *Genome Biology* **26**: 38.

Diensthuber G, Novoa EM. 2025. Charting the epitranscriptomic landscape across RNA biotypes using native RNA nanopore sequencing. *Mol Cell* **85**: 276-289.

Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, Cesarkas K, Jacob-Hirsch J, Amariglio N, Kupiec M et al. 2012. Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature* **485**: 201-206.

Esfahani NG, Stein AJ, Akeson S, Tzadikario T, Jain M. 2025. Evaluation of Nanopore direct RNA sequencing updates for modification detection. *bioRxiv* doi:10.1101/2025.05.01.651717.

Garalde DR, Snell EA, Jachimowicz D, Sipos B, Lloyd JH, Bruce M, Pantic N, Admassu T, James P, Warland A et al. 2018. Highly parallel direct RNA sequencing on an array of nanopores. *Nat Methods* **15**: 201-206.

Gilbert WV, Nachtergaele S. 2023. mRNA Regulation by RNA Modifications. *Annu Rev Biochem* **92**: 175-198.

Gleeson J, Madugalle SU, Wan CY, McLean C, Bredy TW, De Paoli-Iseppi R, Clark MB. 2025. Isoform-level profiling of m(6)A epitranscriptomic signatures in human brain. *Sci Adv* **11**: eadp0783.

Guo W, Ren Z, Huang X, Liu J, Shao J, Ma X, Wei C, Cun Y, He J, Zhang J et al. 2025. Single-molecule m6A detection empowered by endogenous labeling unveils complexities across RNA isoforms. *Molecular Cell* doi:https://doi.org/10.1016/j.molcel.2025.01.014.

He PC, He C. 2021. m(6) A RNA methylation: from mechanisms to therapeutic potential. *Embo j* **40**: e105977.

Hendra C, Pratanwanich PN, Wan YK, Goh WSS, Thiery A, Göke J. 2022. Detection of m6A from direct RNA sequencing using a multiple instance learning framework. *Nature Methods* **19**: 1590-1598.

Hewel C, Wierczeiko A, Miedema J, Friedrich J, Hofmann F, Weißbach S, Dietrich V, Holthöfer L, Haug V, Mündnich S et al. 2025. Direct RNA sequencing enables improved transcriptome assessment and tracking of RNA modifications for medical applications. *Nucleic Acids Research* **53**.

Huang da W, Sherman BT, Lempicki RA. 2009. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**: 44-57.

Jain M, Olsen HE, Paten B, Akeson M. 2016. The Oxford Nanopore MinION: delivery of nanopore sequencing to the genomics community. *Genome Biology* **17**: 239.

Köster J, Rahmann S. 2012. Snakemake--a scalable bioinformatics workflow engine. *Bioinformatics* **28**: 2520-2522.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078-2079.

Linder B, Grozhik AV, Olarerin-George AO, Meydan C, Mason CE, Jaffrey SR. 2015. Single-
nucleotide-resolution mapping of m6A and m6Am throughout the transcriptome. *Nature*
Methods **12**: 767-772.
Liu C, Sun H, Yi Y, Shen W, Li K, Xiao Y, Li F, Li Y, Hou Y, Lu B et al. 2023. Absolute quantification of
single-base m(6)A methylation in the mammalian transcriptome using GLORI. *Nat*
Biotechnol **41**: 355-366.
Liu JF, Hawley BR, Nicholson LS, Jaffrey SR. 2025. Decoding m(6)Am by simultaneous transcription-
start mapping and methylation quantification. *Elife* **13**.
Meyer KD. 2019. DART-seq: an antibody-free method for global m6A detection. *Nature Methods* **16**:
1275-1280.
Meyer Kate D, Saletore Y, Zumbo P, Elemento O, Mason Christopher E, Jaffrey Samie R. 2012.
Comprehensive Analysis of mRNA Methylation Reveals Enrichment in 3' UTRs and near
Stop Codons. *Cell* **149**: 1635-1646.
Moshitch-Moshkovitz S, Sevilla-Sharon M, Ashwal-Fluss R, Glick-Saar E, Rechavi G, Dominissini D.
2024. mRNA m6A detection. *Nature Reviews Methods Primers* **4**.
Mudge JM, Carbonell-Sala S, Diekhans M, Martinez JG, Hunt T, Jungreis I, Loveland JE, Arnan C,
Barnes I, Bennett R et al. 2025. GENCODE 2025: reference gene annotation for human and
mouse. *Nucleic Acids Res* **53**: D966-d975.
Murakami S, Jaffrey SR. 2022. Hidden codes in mRNA: Control of gene expression by m(6)A. *Mol*
Cell **82**: 2236-2251.
Raymon HK, Thode S, Zhou J, Friedman GC, Pardinas JR, Barrere C, Johnson RM, Sah DWY. 1999.
Immortalized Human Dorsal Root Ganglion Cells Differentiate into Neurons with
Nociceptive Properties. *The Journal of Neuroscience* **19**: 5420-5428.
Sendinc E, Shi Y. 2023. RNA m6A methylation across the transcriptome. *Mol Cell* **83**: 428-441.
Shen W, Sun H, Liu C, Yi Y, Hou Y, Xiao Y, Hu Y, Lu B, Peng J, Wang J et al. 2024. GLORI for absolute
quantification of transcriptome-wide m6A at single-base resolution. *Nature Protocols* **19**:
1252-1287.
Su C-H, D D, Tarn W-Y. 2018. Alternative Splicing in Neurogenesis and Brain Development. *Frontiers*
in Molecular Biosciences **Volume 5 - 2018**.
Sun H, Li K, Liu C, Yi C. 2023. Regulation and functions of non-m(6)A mRNA modifications. *Nat Rev*
Mol Cell Biol **24**: 714-731.
Sun H, Lu B, Zhang Z, Xiao Y, Zhou Z, Xi L, Li Z, Jiang Z, Zhang J, Wang M et al. 2025. Mild and
ultrafast GLORI enables absolute quantification of m6A methylome from low-input
samples. *Nature Methods* **22**: 1226-1236.
Verstraten R, Cetraro P, Fitzpatrick AH, Alwie Y, Frommeyer YN, Loliashvili E, Stein SC, Häussler S,
Ouwendijk WJD, Depledge DP. 2025. Defining expansions and perturbations to the RNA
polymerase III transcriptome and epitranscriptome by modified direct RNA nanopore
sequencing. *bioRxiv* doi:10.1101/2025.03.07.641986.
Viehweger A, Krautwurst S, Lamkiewicz K, Madhugiri R, Ziebuhr J, Hölzer M, Marz M. 2019. Direct
RNA nanopore sequencing of full-length coronavirus genomes provides novel insights into
structural variants and enables modification analysis. *Genome Res* **29**: 1545-1554.
Xiao YL, Liu S, Ge R, Wu Y, He C, Chen M, Tang W. 2023. Transcriptome-wide profiling and
quantification of N(6)-methyladenosine by enzyme-assisted adenosine deamination. *Nat*
Biotechnol **41**: 993-1003.
Xie YY, Zhong ZD, Chen HX, Ren ZH, Qiu YT, Lan YL, Wu F, Kong JW, Luo RJ, Zhang D et al. 2025.
Single-molecule direct RNA sequencing reveals the shaping of epitranscriptome across
multiple species. *Nat Commun* **16**: 5119.

Xu H, Kong L, Cheng J, Al Moussawi K, Chen X, Iqbal A, Wing PAC, Harris JM, Tsukuda S, Embarc-
Buh A et al. 2024. Absolute quantitative and base-resolution sequencing reveals
comprehensive landscape of pseudouridine across the human transcriptome. *Nature*
Methods **21**: 2024-2033.
Yang L, Tang L, Min Q, Tian H, Li L, Zhao Y, Wu X, Li M, Du F, Chen Y et al. 2024a. Emerging role of
RNA modification and long noncoding RNA interaction in cancer. *Cancer Gene Ther* **31**:
816-830.
Yang Y, Lu Y, Wang Y, Wen X, Qi C, Piao W, Jin H. 2024b. Current progress in strategies to profile
transcriptomic m(6)A modifications. *Front Cell Dev Biol* **12**: 1392159.
Yankova E, Blackaby W, Albertella M, Rak J, De Braekeleer E, Tsagkogeorga G, Pilka ES, Aspris D,
Leggate D, Hendrick AG et al. 2021. Small-molecule inhibition of METTL3 as a strategy
against myeloid leukaemia. *Nature* **593**: 597-601.
Yu J, She Y, Ji S-J. 2021. m6A Modification in Mammalian Nervous System Development, Functions,
Disorders, and Injuries. *Frontiers in Cell and Developmental Biology* **Volume 9 - 2021**.
Zaccara S, Ries RJ, Jaffrey SR. 2019. Reading, writing and erasing mRNA methylation. *Nature*
Reviews Molecular Cell Biology **20**: 608-624.
Zou Y, Ahsan MU, Chan J, Meng W, Gao S-J, Huang Y, Wang K. 2025. A comparative evaluation of
computational models for RNA modification detection using nanopore sequencing with
RNA004 chemistry. *Briefings in Bioinformatics* **26**.

Figure legends

Figure 1. Workflow for m^{6}A mapping in different human cell types using orthogonal sequencing
approaches.

(A) Schematic of the experimental setup. RNA was isolated from NHDFs and differentiated HD10.6 cells

treated with STM2457 or DMSO for 48h. Poly(A) RNA was subjected to DRS and GLORI from the same

input material. IVT RNA was generated in parallel to assess potential false-positive modification calls. **(B)**

Workflow for benchmarking Dorado basecalling. Reads were processed with Dorado versions v.0.5.0

through v0.9.0, aligned to the human genome or transcriptome, and analyzed with ONT's Modkit tool.

Multiple filtering strategies were tested against IVT controls and METTL3 inhibition (STM2457). Resulting

m^{6}A sites were compared with GLORI data from the same input RNA. **(C)** Final benchmarked Dorado

pipeline to detect high confidence m^{6}A sites.

Figure 2. Modification probability of 0.98 and stoichiometry score of ≥ 10% are necessary cutoffs to identify m⁶A sites accurately.

(A) Dorado v0.9.0-generated modification probability distributions for all-context m⁶A from genome-aligned DMSO- and STM2457-treated NHDFs plotted against the ones from IVT NHDF poly(A) RNA. DMSO probability score distribution is shown as a red solid line, STM2457 – red dashed line, IVT – grey solid line. **(B)** Same as in (A), but for DRACH-context m⁶A sites. **(C)** Stoichiometry (≥ 10%) distributions for all-context m⁶A from DMSO- (in red) or STM2457-treated (in pink) NHDF datasets (genome-aligned) are plotted using a 0.98 modification probability cutoff and a filter for ≥ 20 reads. **(D)** Same as in (C), but for DRACH-context m⁶A sites. **(E)** Overlap of m⁶A sites between all-context and DRACH models (A basecall accuracy ≥ 80%, coverage ≥ 20reads, stoichiometry ≥ 10%), comparing non-filtered to filtered (modification probability > 0.98) outputs. **(F)** Metagene plots showing the density of m⁶A (all-context – dark red, DRACH-context – bright red) sites across all annotated poly(A) transcripts (left) and monoexonic poly(A) transcripts (right) from genome-aligned DMSO-treated NHDF datasets filtered with 0.98 modification probability, ≥ 20 coverage and ≥ 10% stoichiometry.

Figure 3. Comparison of m⁶A sites detected by Dorado and GLORI.

(A) Filtering strategy for Dorado vs GLORI to obtain a high-confidence list of m⁶A sites in NHDFs. **(B)** Stoichiometry distributions are plotted from genome-aligned DMSO- or STM2457-treated NHDF datasets using the indicated Dorado models and the filtering strategy shown in (A). **(C)** Stoichiometry distributions from GLORI performed on DMSO- or STM2457-treated NHDFs are plotted, using the filtering strategy shown in (A). In addition, m⁶A sites were filtered for DRACH motifs (right plot). **(D)** Overlap of high confidence m⁶A sites detected by Dorado vs GLORI. GLORI datasets were filtered using increasing m⁶A stoichiometry cutoffs (from left to right). **(E)** m⁶A sites in the *SPEN* gene detected by GLORI and Dorado. Boxplots (right) represent the m⁶A stoichiometries per m⁶A site shown on the left. **(F)** *SPEN* isoforms with m⁶A sites, determined by Dorado. Boxplots (right) show the distribution of m⁶A stoichiometries on the m⁶A sites on *SPEN-201* and *SPEN-202*.

Figure 4. Comparative analysis of m⁶A modification patterns in NHDFs and HD10.6 cells using Dorado v0.9.0.

665 **(A)** Overlap of high confidence m^{6}A sites between NHDFs and HD10.6 cells. **(B)** Correlation of shared m^{6}A

666 stoichiometries (n=38,217). **(C)** Boxplots showing the m^{6}A stoichiometry distributions of the 38,217

667 detected in both NHDFs and HD10.6 cells. **(D+E)** Motif analysis showing m^{6}A (all-context and DRACH-

668 context) motif logos, 5-mer distributions and modification stoichiometries in NHDFs. **(F+G)** Same as in

669 (D+E) but for HD10.6 cells. **(H)** KEGG pathway analysis (DAVID Bioinformatics Resources) of shared m^{6}A-

670 containing genes. **(I)** m^{6}A stoichiometries of representative examples of individual genes from enriched

671 pathways (H). *ACTG1* serves as a housekeeping control.

673 **Figure S1. Evaluation of Dorado by base and modification probability distributions.**

674 **(A)** Dorado v0.9.0-generated basecall accuracy probability values for A (all-context and DRACH-context

675 basecaller models) from genome-aligned DMSO- and STM2457-treated NHDF samples against IVT poly(A)

676 NHDF RNA. A probability score distribution is shown as a black solid line for DMSO, STM2457 – black

677 dashed line, IVT – grey solid line. **(B)** Dorado v0.9.0-generated modification probability distributions,

678 visualized as histograms, for DRACH-context m^{6}A from genome-aligned DMSO- and STM2457-treated

679 NHDFs and IVT NHDFs. IVT modification probability score distribution is shown in grey, DMSO – bright red,

680 STM2457 – dark red. **(C)** Same as in (B), but for all-context m^{6}A sites.

682 **Figure S2. Modification probability cutoffs are necessary to identify Ψ and inosine sites more accurately.**

683 **(A)** Dorado v0.9.0-generated basecall accuracy probability values for U (left) and modification probability

684 distributions for Ψ (right) from genome-aligned DMSO- and STM2457-treated NHDF samples are plotted

685 against IVT poly(A) RNA. Ψ probabilities are visualized in blue solid and dashed lines for DMSO and

686 STM2457, respectively and in grey solid line for IVT. **(B)** Same as in (A) but showing A base accuracy

687 probabilities and inosine sites. Inosine probabilities are visualized in purple solid and dashed lines for

688 DMSO and STM2457, respectively and in grey solid line for IVT. **(C)** Same as in (A) but showing C base

689 accuracy probabilities and m^{5}C sites. m^{5}C probabilities are visualized in green solid and dashed lines for

690 DMSO and STM2457, respectively and in grey solid line for IVT. **(D+E)** DRACH-context and all-context m^{6}A

691 modification probability distributions (red) from genome-aligned DMSO- and STM2457-treated NHDFs are

692 plotted against RCS negative control (grey).

Figure S3. Modification probability cutoffs are necessary to identify modification sites accurately, shown
for transcriptome-aligned data.

Dorado v0.9.0-generated basecall accuracy probability values for A, U and C (left, **A-E**) and modification
probability distributions for m^{6}A (all-context), m^{6}A (DRACH-context), Ψ, inosine and m^{5}C (right, **A-E**) in
transcriptome-aligned DMSO-and STM2457-treated NHDFs, plotted against the values from IVT NHDF
poly(A) RNA.

Figure S4. Modification probability distribution profiles in genome-aligned DMSO-and STM2457-treated
HD10.6 cells.

Dorado v0.9.0-generated basecall accuracy probability values for A, U and C (left, **A-E**) and modification
probability distributions for m^{6}A (all-context), m^{6}A (DRACH-context), Ψ, inosine and m^{5}C (right, **A-E**) in
human genome-aligned DMSO-and STM2457-treated HD10.6 cells, plotted against the values from IVT
NHDF poly(A) RNA.

Figure S5. Modification probability distribution profiles in transcriptome-aligned DMSO-and STM2457-
treated HD10.6 cells.

Dorado v0.9.0-generated basecall accuracy probability values for A, U and C (left, **A-E**) and modification
probability distributions for m^{6}A (all-context), m^{6}A (DRACH-context), Ψ, inosine and m^{5}C (right, **A-E**) in
human transcriptome-aligned DMSO-and STM2457-treated HD10.6 cells, plotted against the values from
IVT NHDF poly(A) RNA.

Figure S6. Modification site probabilities detected with different versions of Dorado and compared to
RCS.

(A+B) DRACH-context and all-context m^{6}A modification probability distributions from genome-aligned
DMSO (solid red line)- and STM2457 (dashed red line)-treated NHDFs are plotted against RCS negative
control (grey). **(C)** Modification probability distributions for all-context m^{6}A (Dorado v0.7.0, v0.8.0 and
v0.9.0), the bottom panel showing the distributions in 0.95-1.00 modification range. **(D)** Same as in (C),
but for DRACH-context m^{6}A (Dorado v0.5.0, v0.6.0, v0.8.0 and v0.9.0). **(E)** Same as in (C), but for Ψ (Dorado

v0.7.0, v0.8.0, v0.9.0). All the plots in C-E are generated from genome-aligned DMSO-treated NHDF

datasets and are plotted against RCS negative controls.

Figure S7. Evaluation of Dorado by stoichiometry distributions.

(A) Stoichiometry distributions for all-context m^{6}A from DMSO- (in red) or STM2457-treated (in pink) NHDF

datasets (genome-aligned and filtered for ≥ 20 reads) are plotted, comparing 0.98 modification probability-

unfiltered (left) and filtered (right) sites. **(B)** Same as in (A), but for DRACH-context m^{6}A sites.

Figure S8. Stoichiometry distributions of m^{6}A, Ψ and inosine in IVT datasets.

(A) All-context m^{6}A stoichiometry distributions (all stoichiometries (top) and ≥ 10% stoichiometries

(bottom)) in (genome-aligned) IVT RNA from DMSO-treated NHDFs, comparing modification probability-

unfiltered and filtered data. The bottom panel shows stoichiometry distributions only for ≥ 10%

stoichiometry sites. **(B)** As in (A), but for DRACH-context m^{6}A. **(C)** As in (A), but for Ψ. **(D)** As in (A), but for

inosine.

Figure S9. Evaluation of Dorado by stoichiometry distributions in transcriptome-aligned NHDFs.

(A) Stoichiometry distributions for all-context m^{6}A from DMSO- (in red) or STM2457-treated (in pink) NHDF

datasets (transcriptome-aligned and filtered for ≥ 20 reads) are plotted, assessing the 0.98 modification

probability cutoff. The bottom panel shows stoichiometry distributions only for ≥ 10% stoichiometry sites.

(B) Same as in (A), but for DRACH-context m^{6}A sites.

Figure S10. Evaluation of Dorado by stoichiometry distributions in transcriptome-aligned HD10.6 cells.

(A) Stoichiometry distributions for all-context m^{6}A from DMSO- (in red) or STM2457-treated (in pink)

HD10.6 datasets (transcriptome-aligned and filtered for ≥ 20 reads) are plotted, assessing the 0.98

modification probability cutoff. The bottom panel shows stoichiometry distributions only for ≥ 10%

stoichiometry sites. **(B)** Same as in (A), but for DRACH-context m^{6}A sites.

Figure S11 Stoichiometry distributions of Ψ and inosine in NHDFs.

(A) Distributions of all (left) and ≥ 10% (right) Ψ stoichiometries in the 0.99 modification probability-

unfiltered and filtered data from genome-aligned DMSO- (darker color) and STM2457-treated (lighter

color) NHDFs (filtered for ≥ 20 reads). **(B)** Same as in (A), but for inosine.

Figure S12. Stoichiometry distributions of Ψ and inosine in HD10.6 cells.

(A) Distributions of all and ≥ 10% Ψ stoichiometries in the 0.99 modification probability-unfiltered and

filtered data from genome-aligned DMSO- (darker color) and STM2457-treated (lighter color) HD10.6 cells

(filtered for ≥ 20 reads). **(B)** Same as in (A), but for inosine.

Figure S13. m^{6}A and Ψ stoichiometry distributions in different versions of Dorado.

Stoichiometry distributions (all stoichiometries – left, ≥ 10% stoichiometries – right) in a line graph format

for **(A)** all-context m^{6}A detected in genome-aligned DMSO-treated NHDF dataset, with Dorado versions

0.7.0, 0.8.0 and 0.9.0 colored in gradients of red, darkness increasing with the later versions. **(B)** DRACH-

context m^{6}A detected with Dorado versions 0.5.0, 0.6.0, 0.8.0 and 0.9.0, colored in gradients of red,

darkness increasing with the later versions. **(C)** Ψ detected with Dorado versions 0.7.0, 0.8.0 and 0.9.0,

colored in gradients of blue, darkness increasing with the later versions. All the datasets were filtered for

≥ 20 reads.

Figure S14. Comparison of modification sites detected with different versions of Dorado.

Upset plots showing unique and shared detected modifications (filtered for ≥ 20 reads, ≥ 10%

stoichiometry) in modification probability-unfiltered and filtered genome-aligned DMSO-treated NHDF

data. The versions are ordered on the plot by the number of detected sites, in ascending order (on the

left). Results are shown for the following models and versions: **(A)** all-context m^{6}A detected with v.0.7.0,

v0.8.0 and v0.9.0, **(B)** DRACH-context m^{6}A detected with v.0.5.0, v0.6.0, v0.8.0 and v0.9.0, **(C)** Ψ detected

with v.0.7.0, v0.8.0 and v0.9.0. **(D)** Nucleotide frequencies in the 5-base motif of Ψ sites for uniquely

v0.7.0-, v0.8.0-0.9.0- and v0.7.0-0.8.0-0.9.0-detected Ψ sites. Position 0 marks the modification site.

Figure S15. Comparison of m^{6}A sites detected by Dorado and GLORI-Sequencing in NHDFs and HD10.6s.

(A) Upset plot showing the number of m^{6}A sites detected by GLORI and Dorado in NHDFs and their

overlaps. Dorado sites represent high-confidence m^{6}A sites obtained after filtering DRACH-model detected

data. Similarly, only high confidence m^{6}A sites detected by GLORI are shown but were additionally filtered

for DRACH sites. **(B)** Stoichiometry distributions are plotted from genome-aligned DMSO- or STM2457-

treated HD10.6 datasets using the indicated Dorado models. **(C)** Stoichiometry distributions from GLORI

performed on DMSO- or STM2457-treated NHDFs are plotted. In addition, m^{6}A sites were filtered for

DRACH motifs (right plot). **(D)** Stoichiometry distributions are plotted from transcriptome-aligned DMSO-

or STM2457-treated NHDF datasets using the indicated Dorado models. **(E)** Stoichiometry distributions

are plotted from transcriptome-aligned DMSO- or STM2457-treated HD10.6 datasets using the indicated

Dorado models.

Figure S16. m^{6}A (all-context and DRACH-context) motif logos for modification probability-unfiltered

NHDF and HD10.6 data.

(A) 5-mer motif logo (showing nucleotide frequences at each position) of m^{6}A-modified sites (position 0

= modified A) from genome-aligned DMSO-treated NHDFs (all-context – left, DRACH-context – right). All

the datasets were filtered for ≥ 20 coverage & ≥ 10% stoichiometry. **(B)** Same as (A), but for HD10.6 cells.

Figure S17. Metagene plots showing m^{6}A (all-context and DRACH-context) distribution profiles in NHDF

and HD10.6 data filtered at different stoichiometries.

(A) Metagene plots showing the density of m^{6}A (all-context – dark red, DRACH-context – bright red) sites

across all annotated poly(A) transcripts (upper plot) and monoexonic poly(A) transcripts (lower plot) from

genome-aligned DMSO-treated NHDF datasets filtered with 0.98 modification probability, ≥ 20 coverage

and ≥ 10% stoichiometry. **(B)** The density of all-context m^{6}A sites (0.98 modification probability-unfiltered

– left, filtered – right) is plotted across all annotated poly(A) transcripts of NHDF (dark red) and HD10.6

(bright red) datasets, filtered with different stoichiometries (descending: all stoichiometries, ≥ 50%, ≥ 10%

and < 10%. All datasets were filtered additionally for ≥ 20 coverage & ≥ 10% stoichiometry. **(C)** Same as

(A), but for DRACH-context m^{6}A sites.

A

B

ONT DRS (P2 solo) and Dorado

C

Benchmarked m⁶A Dorado pipeline

Figure 2

Figure 3

Figure 4

Figure S4

Figure S6

Figure S10

Figure S11

Figure S12

Figure S14

A all-context

m⁶A DRACH

NHDF

B all-context

m⁶A DRACH

HD10.6

Figure S17