Drought, thermal response and climate-patterning in cuticular conductance of the widespread C4 grass, *Themeda triandra*

Anu Middha[1], Vinod Jacob[1], Chris J. Blackman[2], Brendan Choat[1], Ian J. Wright[1]

[1]ARC Centre of Excellence for Plant Success in Nature and Agriculture, Hawkesbury Institute for the Environment, Western Sydney University, Richmond, NSW 2753, Australia

[2]ARC Centre of Excellence for Plant Success in Nature and Agriculture, School of Natural Sciences, University of Tasmania, Hobart 7001, Tasmania, Australia

Author emails:

a.middha@westernsydney.edu.au (corresponding author)

vinod.jacob@westernsydney.edu.au

christopher.blackman@utas.edu.au

b.choat@westernsydney.edu.au

ian.j.wright@westernsydney.edu.au

Type of article: Primary Research Paper (Original Article)

Running title: Cuticle responses to drought and temperature in *Themeda triandra.*

Abstract

- *Background and Aims:* Grasslands underpin global biodiversity, carbon storage, and ecosystem resilience, yet are increasingly threatened by rising temperatures and water scarcity. Understanding how key physiological traits respond to drought and heat is essential for ensuring grassland function under future conditions. Here, we investigated intraspecific variation in leaf cuticular traits—specifically minimum cuticular conductance after stomatal closure (g_{min}) and its response to temperature—in six Australian accessions of *Themeda triandra* spanning a wide climatic gradient. We asked whether g_{min} and its response to drought reflect climate-of-origin and whether the cuticle shows a thermal threshold (Tp) beyond which conductance rises sharply.

- *Methods:* Plants from six accessions were grown under well-watered (control) glasshouse conditions and then exposed to drought. We measured g_{min} in both control and drought treatments and assessed the response of gmin to increasing temperature (30–55 °C) using fresh fully hydrated leaves (control). Climatic data for each site of origin were used to explore trait–environment relationships.

- *Key results:* Under well-watered conditions, g_{min} showed no link to climate-of-origin. Under drought, however, g_{min} displayed clearer climate-linked patterns: accessions from cooler, wetter regions had lower values, contrary to expectations. Drought responses varied strongly among accessions, ranging from marked reductions to significant increases in $g_{min.}$ With increasing temperature, g_{min} declined gradually and no accession exhibited a distinct phase transition, indicating a thermally stable cuticle.

- *Conclusions: Themeda triandra* shows considerable intraspecific diversity in g_{min} and its response to drought. The clearer alignment between climate of origin and g_{min} measured under drought conditions suggests drought stress is an important filter for g_{min} expression. These findings provide a physiological basis for identifying genotypes with enhanced resilience for use in grassland conservation and restoration under a warming, drying climate.

31   **Keywords:** drought**,** leaf cuticle, minimum cuticular conductance, phase transition

32   temperature, *Themeda triandra*, water loss.

52   **Introduction**

Grasslands occupy 31% of earth's land area (Petri *et al.*, 2010) and are essential for preserving biodiversity, sequestering carbon and maintaining ecosystem balance (Blair *et al.*, 2014). However, they are highly vulnerable to climate change, particularly rising temperatures and altered rainfall patterns (Sala *et al.*, 2000, Teuling, 2018). These shifts threaten grassland resilience by reducing species diversity, productivity, and soil integrity (Overpeck, 2013, Edenhofer *et al.*, 2014), underscoring the need to understand their impacts for effective conservation and management.

Under well-watered daytime conditions, stomata remain open to facilitate gas exchange for photosynthesis while simultaneously losing water via transpiration. During drought, plants reduce water loss primarily by closing their stomata. However, even after full stomatal closure, plants continue to lose water through the leaf epidermis ("residual transpiration"; Billon *et al.* (2020)). This residual transpiration can be significant, particularly under prolonged drought or high temperatures, and is primarily attributed to the permeability of the leaf cuticle, although leaky stomata may also contribute (Muchow & Sinclair, 1989, Machado *et al.*, 2021). The rate of this residual water loss from the leaf surface when stomata are fully closed is often referred to as "minimum cuticular conductance" or "g_{min}". g_{min} is increasingly recognized as being ecologically important, indexing the effectiveness of the cuticle for slowing water loss during drought (Schuster *et al.*, 2016). In dryland and savanna species, for example, small differences in g_{min} can significantly affect drought survival (Burghardt & Riederer, 2003). Recent work has highlighted that g_{min} is not constant but dynamic throughout dehydration (Burlett *et al.*, 2025). This variability in g_{min} can potentially alter predictions of time to hydraulic failure (Martin☐StPaul *et al.*, 2017), thus highlighting the need to understand the response of g_{min} to drought stress, especially against the background of projected increasingly prolonged and severe droughts.

At moderate temperatures (e.g. 20–25 ☐), g_{min} of healthy intact leaves is relatively low, typically averaging around 10% of the transpiration through fully open stomata, depending on the species (Holmgren *et al.*, 1965, Boyer *et al.*, 1997). This is because cuticular waxes serve as an efficient barrier against water movement (Boyer *et al.*, 1997, Burghardt & Riederer, 2003, Riederer, 2006). However, g_{min} may vary with ambient temperature, displaying non-linear dynamics such that it remains relatively stable across these moderate temperatures, then increases sharply beyond some threshold temperature (Schreiber & Schönherr, 1990). This threshold temperature is commonly known as the

"phase transition temperature" or (T_p). At T_p, the structural integrity of the cuticle begins to
shift due to thermal disruption of wax and polymer organization, resulting in a dramatic
increase in g_{min} and hence, higher water loss (Schreiber & Schönherr, 1990). T_p can vary
among species and populations and is considered a marker of thermal stability of the
cuticle (Schuster *et al.*, 2016, Bueno *et al.*, 2019, Billon *et al.*, 2020). For example,
Schuster *et al.* (2016) reported that the desert shrub *Rhazya stricta* had a markedly high T_p,
suggesting an adaptation to hot, arid conditions. The temperature response of g_{min} is further
influenced by the chemical composition of cuticular waxes, as compounds such as alkanes,
fatty alcohols, aldehydes and cyclic triterpenoids differ in their melting behaviour and
structural stability under heat. Busta *et al.* (2021) showed that triterpenoid-rich sorghum
leaves possess a stronger, more heat-resistant cuticular barrier than maize leaves which
lacked triterpenoids. In the context of climate change and heatwaves, understanding T_p is
critical because it determines the threshold temperature at which the primary barrier to
plant water loss is compromised during drought. The extent to which these changes are
reversible is also unknown. For instance, Schönherr *et al.* (1979) observed that heating
cuticular membranes above their transition temperature irreversibly altered the distribution
of soluble cuticular lipids, leading to increased water permeability.

Substantial interspecific variation in g_{min} has been documented; e.g. Duursma *et al.*
(2019) compiled data for 221 species from 40 original studies, reporting values from ~ 0.1
to 50 mmol m^{-2} s^{-1}. By taxonomic order, Poales on average exhibited the highest values,
and conifers (Pinales, Araucariales) the lowest, but there was wide variation within any
given group. There is evidence showing that g_{min} variation reflects differences in the
chemical composition of cuticles; e.g. the richness of cuticular wax components (such as
long-chain alkanes) is known to reduce cuticle permeability (Huang *et al.*, 2017, Cheng *et*
al., 2019).

Climate-related *intraspecific* variation in g_{min} and cuticle traits has also been
observed. For example, some studies have reported g_{min} as lower in plants from arid
compared to mesic environments, presumably to minimize residual water loss during
drought; this was the case for one of two *Hakea* (Proteaceae) species studied by López *et*
al. (2021). Furthermore, Smith *et al.* (2006) compared *Digitaria californica* populations
from Texas (arid) and Arizona (less arid), as well as the invasive species *Eragrostis*
lehmanniana, in a common garden setting. They found that the arid-zone population and *E.*
lehmanniana both exhibited significantly lower g_{min} values. In *Nicotiana benthamiana*, two

accessions exhibited contrasting drought strategies: plants from arid central Australia relied on rapid growth and early flowering to escape drought, with minimal changes in cuticle properties or g_{min} under experimental drought (Asadyar *et al.*, 2024). In contrast, plants from tropical northern Australia reduced g_{min} in response to water limitation by altering wax composition, specifically increased very-long-chain alkanes, to minimize water loss under drought conditions (Asadyar *et al.*, 2024). Other studies have also shown acclimation of g_{min} to drought and elevated temperature conditions (Duursma *et al.*, 2019). However, other studies have shown a lack of response in g_{min} to changing growth conditions (Bueno *et al.*, 2020).

Most research on cuticle conductance has focused on woody species and a limited subset of herbaceous plants, primarily high-value annual crops such as wheat and sorghum (Muchow & Sinclair, 1989, Araus *et al.*, 1991, Qariani *et al.*, 2000, Lin *et al.*, 2020). In contrast, native and perennial grass species—despite their ecological significance—remain understudied. As a result, our understanding of how perennial grasses regulate cuticular water loss in response to drought remains limited, highlighting the need for broader investigation across diverse grass lineages.

This study focuses on *Themeda triandra*, a perennial C4 tussock grass endemic to Africa, Australia and Asia (Snyman *et al.*, 2013). It has great ecological significance due to its broad distribution across various ecosystems and geological substrates. It is a valuable source of food for both native and introduced herbivores, and plays a crucial role in supporting rural livelihoods, wildlife and livestock production (Snyman *et al.*, 2013). *T. triandra* is widely distributed across Australia and is believed to have been the primary dominant species in many Australian grasslands prior to European settlement (Mitchell & Miller, 1990). It grows in habitats as diverse as the hot, semiarid interior, the wet tropics, and cool temperate sub-alpine regions (Mitchell & Miller, 1990). This makes it an ideal model for exploring physiological responses to climate-induced stresses in grasses. A reduction in the occurrence of *T. triandra* within grasslands is often associated with a drop in grazing potential, a loss of biodiversity, reduced vegetation cover and a decline in overall health of the ecosystem (Snyman *et al.*, 2013). Despite its huge ecological and economic significance, to this point there has been no attempt to study its cuticle properties.

In this study, we measured g_{min} in plants under well-watered and droughted conditions and explored its response to increasing temperature across six distinct

accessions of *T. triandra* originating from diverse regions across Australia, spanning a wide range of temperature and rainfall regimes. We had two key aims. First, we sought to elucidate the relationships between climate conditions at the seed source ("climate-at-origin") and two aspects of cuticle permeability: 1) g_{min} and 2) its temperature response (T_p). Second, we sought to characterise the different accessions in terms of their variation in g_{min} and T_p under control and water-limiting conditions.

We tested the following hypotheses:

1. *T. triandra* accessions from hotter and drier environments would exhibit generally lower g_{min}, reflecting selection for reduced cuticular water loss under high evaporative demand and limited water availability.

2. Cuticle conductance would show a bi-phasic response to increasing temperature, remaining relatively steady at moderate temperatures, then rising sharply once temperatures exceeded a threshold, T_p, which itself would vary with climate-at-origin. Specifically, we expected plants from hotter environments to have higher T_p; i.e., to maintain cuticular stability up to higher temperatures.

3. Experimental drought would reduce g_{min} in all accessions. This expectation assumes that the lower gmin would reduce plant water loss when savings are needed most. We also expected the proportional change in g_{min} (from control to drought conditions) to be greater in accessions originating from arid or high-temperature regions. This is based on the assumption that plants from such environments have evolved greater plasticity or regulation of cuticle properties to limit water loss during droughts (Bueno *et al.*, 2019).

Materials and methods

Plant material and growth conditions

Six accessions of *T. triandra* (Dalby, QLD; Forbes, NSW; Mt. Fox National Park, Queensland, Pannawonica, Western Australia; Rainbow Valley, Northern Territory; and Virginia Gardens, South Australia) (Fig. 1 (a)) were grown from seeds sourced from the Australian Pastures Genebank, Native Seeds Pty Ltd and Nindethana Seed Service Pty Ltd (Albany, Western Australia). The selected accessions represent a range of climate-at-origin conditions (climate data from CHELSA (Karger *et al.*, 2017); Table S1), with mean summer temperatures ranging from 21.0 ☐ to 32.0 ☐, summer rainfall from 75.1 to 435.9

mm, annual temperatures (AT) from 16.2 □ and 22.1 □ and annual precipitation (AP) from

281 to 712 mm. Plants were propagated from seeds, germinated in seed trays, and

transplanted into 250 mm (~8 L) pots filled with sandy-loam soil (volumetric water holding

capacity of 15−20%, pH ≈ 5.6) collected from the Pastures and Climate Extremes (PACE)

field facility at Western Sydney University (Churchill *et al.*, 2022), supplemented with

slow-release native fertiliser (Osmocote® Native; 20 g pot^{-1})

Water deficit treatment

Each accession was represented by four replicates, all maintained under well-watered

conditions for two months. Plants were irrigated to full field capacity using a drip system.

Toward the end of this period, all four replicates from each accession were sampled for g_{min}

and Tp measurements under well-watered conditions. The plants were then subjected to an

acute water-deficit treatment by withholding irrigation until the leaves were visibly wilted

(3 days for all plants) to simulate a rapid decline in soil moisture. Soil water content was

monitored using a soil moisture probe, and leaf water potential was measured with a

pressure chamber to assess plant water status (Figs S1 and S2). At the end of the drying

period, the plants were re-sampled for determination of gmin under water-deficit-treated

conditions.

Leaf drying methods used for g_{min} determination

Initially, g_{min} measurements were conducted using the benchtop drying method (Sack &

Scoffoni, 2010) at a single temperature (30 □), which was well-suited for assessing the

steady state cuticular conductance under controlled conditions. However, as the study

progressed, the need to measure g_{min} across multiple temperatures became evident,

requiring a more efficient and automated approach. The benchtop method, though reliable,

was time- and labour-intensive and limited in its throughput. To overcome these limitations

and facilitate rapid measurements across a range of temperatures, we incorporated a

"DroughtBox" (Billon *et al.*, 2020) into our workflow, which provided an automated, high-

throughput system for assessing g_{min} under precisely controlled environmental conditions.

Description of the DroughtBox

The DroughtBox, originally described by Billon *et al.* (2020) consists of a 70 cm cubic box with two compartments for semi-automatically determining g_{min} under controlled environmental conditions, specifically temperature and humidity. The system includes strain gauges connected to a data acquisition board (for measuring the mass loss from leaves as they dehydrate), heating resistors for temperature control, and electronically controlled valves for adjusting relative humidity. The original design also incorporates a Raspberry Pi for overall control and monitoring.

In this study, we used a modified version of the DroughtBox to measure g_{min} and T_p. Our version consisted of a 60 cm × 140 cm × 70 cm cuboidal box made of 40 mm thick polystyrene boards with a front door and two compartments (Fig. S4). The upper compartment housed 8 strain gauges (Phidgets single point load cell 780g), fixed in an aluminium frame and directly connected to a Campbell Scientific CR300 datalogger for data acquisition. Leaves for g_{min} measurements were placed in the main lower compartment and attached to the strain gauges through holes in the roof board. An air temperature and relative humidity sensor (Vaisala HMP155) was suspended in the lower compartment to monitor environmental conditions, along with a fine-wire type T thermocouple for rapid response temperature control. Temperature regulation was achieved using two PTC heaters (Joyzan, 220 V, 300 W) mounted in galvanized steel conduits, with a 50 mm 12 V fan to push heated air into the box. A thermostat switch (MULTICOMP T23A150BSR2-15) was installed for safety, shutting off the heaters if they overheat. A solid-state relay (Sparkfun electronics, 40A, 3–32 V DC) was used to allow logger to regulate the heaters via pulse width modulation. Relative humidity in the lower compartment was adjusted by a water-filled plastic container with ultrasonic mist maker (Vaguelly mist maker, 350 mL, 24 V). A 12 V pump (DANXQ, 12 V DC) circulated air through the mist container to introduce fine mist into the box when needed. All environmental variables were monitored and controlled using a Campbell Scientific CR3000 datalogger with program written and compiled in Visual Basic.

Performance testing of the DroughtBox system

The ability of the DroughtBox to achieve and maintain stable temperature and humidity conditions was assessed. To do this, the system was programmed to run at each target

temperature and relative humidity required for our experiment over 24-hour periods, with
temperature, relative humidity, and VPD recorded every 10 seconds.

Cuticle conductance measurement (benchtop drying)

To assess leaf water loss, 4–5 healthy tillers with fully developed, mature leaves were
collected in the morning on each sampling day. The cut ends of the tillers were immediately
wrapped in damp paper towels, placed in plastic zip lock bags, and transported to the lab.
To relieve xylem tension, the tiller ends were recut under water and allowed to rehydrate in
falcon tubes for about 30 minutes. The cut ends were then sealed with parafilm. The tillers
were weighed using a balance to record their fully hydrated weight. The leaves were then
dehydrated in open plastic trays placed in a controlled growth cabinet set at 30 °C and 60%
relative humidity, in the dark, for 2–3 hours, with leaf weight (w) measured every 20
minutes until the rate of water loss stabilized. We considered changes in w to represent the
number of water molecules (n_w, mmol) transpiring from the branches. Based on this
assumption, transpiration rate, E (mmol m^{-2} s^{-1}) was calculated as

$$E = -\frac{slope\ (n_w;\ t)}{LA} \tag{1}$$

260 where, t is the time in seconds, and LA represents two-sided leaf area. LA was estimated
261 using the dry leaf mass of each tiller set and empirical relationships between leaf dry mass
262 and fresh leaf area (i.e., LMA). Subsequently, g_{min} was determined from the linear segment
263 of the water loss curve, that is, after stomatal closure, as following

$$g_{min} = -\frac{E}{VPD} \times 101.6 \tag{2}$$

where, E is transpiration in mmol m^{-2} s^{-1}, VPD refers to the vapour pressure deficit of the
air in kPa, and 101.6 denotes the atmospheric pressure within the cabinet (described below)
in kPa.

At the end of the experiment, the leaves were oven-dried at 70°C for 72 hours, and
the final dry weight was recorded.

The percentage change in g_{min} in response to drought ($\%\Delta g_{min}$) was determined

using the g_{min} data from control and drought-treated plants. It was calculated using the

formula

$$\%\Delta gmin = \frac{gmin(drought) - gmin(control)}{gmin(control)} \times 100 \qquad (3)$$

g_{min} measurement using the DroughtBox

The tillers were harvested and prepared for drying in a manner consistent with benchtop

drying procedure. Multiple leaf-drying runs were conducted in the DroughtBox at target

temperatures of 30, 35, 40, 45, 50 and 55 □ (using a distinct cohort of samples at each

temperature), with corresponding increases in relative humidity (RH) to maintain a

constant absolute humidity of 18 $g(H_2O)m^{-3}$. This standardization ensured consistent

conditions during g_{min} measurements. The mass data from the DroughtBox were similarly

used to calculate g_{min} from the linear portion of the mass loss curve, as was done following

benchtop drying procedure. After the DroughtBox measurements were complete, samples

were oven-dried at 70 □ for 72 hours to determine their dry mass.

Leaf mass per area (LMA)

286   LMA was calculated as the ratio of leaf dry mass to leaf area. For each replicate plant, fully

287   expanded leaves were harvested and scanned using a LICOR leaf area meter (model LI-

288   3100C, LI-COR Biosciences, USA) to determine the total leaf area. The scanned leaves

289   were then oven-dried at 70 °C for 72 hours to reach dry mass. LMA ($g\ m^{-2}$) was then

290   calculated by dividing dry mass by leaf area.

Statistical analyses

Differences in midday leaf water potentials were analysed using two-way ANOVA. To

assess treatment effects within each accession, assumptions of normality (Shapiro-Wilk

test) and homogeneity of variance (F-test, Levene's test) were evaluated. Based on these.

Appropriate statistical tests (paired t-test, Welch's t-test, or Wilcoxon signed-rank test) were

used. The relationship between leaf water potential and the direction and magnitude of the

g_{min} response across individuals was analysed using ordinary least squares linear regression.

The g_{min} data, including those obtained from both benchtop drying and from

multiple temperature runs in the DroughtBox, were visually examined for normality before

analysis using histograms and density plots, and log-transformed wherever the data were

right-skewed. Significant differences in g_{min} values between accessions and treatments were

assessed using a two-way analysis of variance (ANOVA), followed by Tukey's Honestly

Significant Difference (HSD) test and paired t-test for pairwise comparisons. The

relationships of g_{min} and %Δg_{min} with climate variables were evaluated through multiple

linear regression models using mixed effects models with accession treated as a random

effect. By including both temperature and rainfall in these models, we were able to identify

the rainfall-independent effect of temperature and the temperature-independent effect of

rainfall. Climate variables were treated both on a yearly basis (annual temperature, AT;

annual precipitation, AP), and on a seasonal basis (mean summer temperature; summer

rainfall).

To determine T_p, we used segmented regression applied to Arrhenius-transformed

data. Specifically, g_{min} values were natural log-transformed (ln g_{min}) and plotted against the

inverse of absolute temperature (1/T, in Kelvin). For each replicate, a segmented linear

regression model was fitted using the segmented package in R to estimate a potential

breakpoint along the temperature axis, corresponding to a putative T_p. A significance

threshold of 0.05 was applied for all statistical tests.

All analyses were conducted using R v.4.4.0 (R, **Core Team** 2025).

Results

DroughtBox performance under different temperature settings

322 When programmed to maintain constant temperatures of 30, 35, 40, 45, and 50 °C—with

323 corresponding increases in RH to keep absolute humidity consistent—the DroughtBox

324 successfully maintained stable temperature, relative humidity, and VPD throughout each

325 24-hour run (Fig. S5).

Analysis of drought exposure and effect on g_{min}

Although soil moisture declined at different rates among accessions and replicates (Fig. S1), indicating that plants experienced drought progression at slightly different rates, all droughted plants exhibited visible signs of leaf wilt and crossed their accession-specific TLP (Jacob *et al*, unpubl. res.) (Fig. S2), confirming that each plant experienced physiological drought. In all accessions, droughted plants exhibited a decline in leaf water potential relative to control conditions (ANOVA: treatment, $p < 0.001$, accession x treatment, $p < 0.05$; pairwise differences: all $p < 0.05$) and all crossed their respective TLP indicating loss of turgor. We found no significant relationship between percentage change in g_{min} from control to drought ($\%\Delta g_{min}$) and leaf water potentials ($p = 0.642$, $R^2 = 0.01$).

Drought-induced changes in cuticular conductance across accessions

Pooling data across the six accessions, g_{min} did not differ between control and droughted plants (Fig. 2). However, substantial variation among accessions was evident, both in mean g_{min} values and in their response of g_{min} to drought (Fig. 3). g_{min} of control plants varied among accessions (ANOVA, $p < 0.001$), being significantly higher in plants from Pannawonica and Mt. Fox than those from Dalby (Tukey HSD, both $p < 0.001$), Forbes (Tukey HSD, $p = 0.003$ and 0.015, respectively), and Virginia Gardens (Tukey HSD, $p = 0.004$ and 0.019, respectively).

For droughted plants g_{min} also varied among accessions (ANOVA, $p = 0.041$), albeit sufficiently weakly that no pairwise differences were statistically significant after post hoc correction (Tukey HSD, all $p > 0.05$). One comparison was very nearly statistically significant, with plants from Rainbow Valley tending to have higher g_{min} than those from Mt. Fox (Tukey HSD, $p = 0.054$).

Although there was no overall difference in g_{min} between control and droughted plants when pooled across all accessions, individual accessions differed in their g_{min} response to drought (accession $\times$ treatment interaction: ANOVA, $p < 0.001$), with two exhibiting pronounced reductions in g_{min} (Mt. Fox and Pannawonica; paired t-test, $p = 0.0162$ and $p = 0.0104$, respectively), three showing no significant difference in g_{min} with treatment, and one (Dalby) exhibiting increased g_{min} (paired t-test, $p < 0.001$). The asterisks in Figure 3 identify these accessions and the degree of significance of these shifts.

Influence of climate-at-origin on g_{min} and its response to drought

To evaluate how climate-at-origin influences g_{min}, we assessed relationships between g_{min} and both seasonal and annual climate variables. Our primary focus was on summer temperature and summer rainfall (Fig. 4), with relationships involving mean annual precipitation and mean annual temperature reported as supplementary results (Fig. S6).

Under control conditions, g_{min} showed no relationship with either summer rainfall (p = 0.484; Fig. 4a) or summer temperature (p = 0.286; Fig. 4b). Similarly, g_{min} showed no relationship to either AP or AT (both p > 0.12; Figs S6a,b). By contrast, clear climate-linked patterns emerged under drought conditions, g_{min} being on average higher in plants from drier sites (Figs 4c, S5c; p = 0.014 and 0.002 respectively), such that a reduction in summer rainfall from 436 mm to 75 mm corresponded to a 16.5% increase in g_{min}. On average, g_{min} of droughted plants was also higher in plants from sites with warmer summers (p = 0.003; Fig. 4d), such that an increase in summer temperature from 21 °C to 32 °C corresponded to a 23.5% increase in g_{min}. There was no relationship between g_{min} and MAT (p = 0.961; Fig. S6d).

To assess plasticity in cuticle conductance, we quantified the percentage change in g_{min} from control to drought (%Δg_{min}). Contrary to our expectation that accessions from hotter or drier environments would show greater reductions in g_{min}, %Δg_{min} was not statistically related to any of summer rainfall, summer temperature, AP or AT (Figs. 5a, b and S6a, b; all p > 0.05).

Temperature dependence of g_{min} across accessions

The "textbook" response of g_{min} to temperature was not observed in our study (i.e., we did not observe a consistent pattern of near-constant g_{min}, rapidly increasing at some threshold temperature, T_p). Overall, the response of g_{min} to increasing temperature was broadly similar across accessions (Fig. 6, a–f). While most accessions (Mt. Fox, Pannawonica, Rainbow Valley and Virginia Gardens) showed a relatively flat response with slightly decreasing g_{min} as temperature increased, plants from Dalby and Forbes displayed a modest rise in g_{min} around 35–40 °C before eventually declining at higher temperatures, like the other accessions. A segmented regression model applied to individual replicates revealed

389 no consistent evidence of a distinct breakpoint across accessions. This statistical outcome
390 aligns with the visual inspection of the plots, where g_{min} appeared largely stable or declined
391 gradually across the experimental temperature range.

Discussion

Intraspecific variation in g_{min} responses to drought

As the plants were well-watered before a single, short drought cycle, the treatment
primarily reflects acute physiological plasticity rather than long-term structural
acclimation. Brief water deficits can rapidly induce cuticular adjustments via abscisic acid
(ABA) signalling and reorganisation of the wax biosynthetic pathways in plants, often
within hours to days (Shepherd & Wynne Griffiths, 2006, Macková *et al.*, 2013, Yang *et*
al., 2020). Similar short term wax shifts have been documented in *Nicotiana glauca*
(Cameron *et al.*, 2006).

This study highlights the impact of short-term drought stress on g_{min} across six
accessions of *T. triandra,* revealing substantial within-species variability in responses.
Although the overall effect of drought on g_{min} was not statistically significant at the species
level (Fig. 2), the direction and magnitude of responses differed substantially among
accessions (Fig. 3), pointing to accession-specific strategies in cuticular water regulation.
The lack of association between leaf water potentials and $\%\Delta g_{min}$ suggests that differences
in cuticular conductance among accessions were not driven by unequal drought severity but
instead reflect inherent variation in cuticle function and plasticity.

The significant reductions in g_{min} in response to drought observed in accessions
from Mt. Fox (48%, p = 0.0162) and Pannawonica (36%, p = 0.0104) align closely with the
classical drought-adaptive responses. While the drought response of g_{min} was not directly
correlated with the accessions' climate-at-origin, these two regions share ecological
features—seasonally intense or prolonged drought, high evaporative demand, and
unpredictable rainfall. These climate features may favour physiological strategies that
reinforce the cuticle under water deficit. In these cases, the decrease in g_{min} may therefore
reflect inducible cuticular responses, such as increased wax deposition or structural
adjustments that strengthen the water barrier.

However, these patterns were not universal. The Dalby accession exhibited a 31%

increase in g_{min} (p < 0.001) under drought stress—a response that diverges from the typical

drought-induced reduction. This increase may reflect a plastic, rather than adaptive,

response to drought—potentially involving alterations in cuticular wax composition that do

not enhance water barrier function—as observed in grape berries under water deficit

(Dimopoulos *et al.*, 2020). Notably, Dalby had lower g_{min} than the other accessions under

control conditions and, even with this increase, its absolute g_{min} remained relatively low.

This suggests that plants from Dalby may possess inherently lower cuticular permeability

overall, and that the observed increase may still fall within a functionally conservative

range.

In contrast, in plants from Forbes, Rainbow Valley and Virginia Gardens, g_{min} did

not differ between control and drought treatments. This suggests a high degree of drought

tolerance or a mechanism of homeostasis that stabilizes physiological performance under

fluctuating water availability. Such accessions might possess robust cuticular barriers that

maintain consistent conductance levels regardless of external stressors. Alternatively, these

accessions may rely on non-cuticular mechanisms—such as deep or extensive roots

systems or osmotic adjustment—to cope with water stress. For instance, efficient root

systems can enhance water uptake (Wasson *et al.*, 2012), and osmotic adjustment can

maintain cell turgor (Turner, 2018) and help plants tolerate water limitation (DaCosta &

Huang, 2006). Osmotic adjustment involves the accumulation of solutes in plant cells

(Hsiao *et al.*, 1976), which aids in retaining water and sustaining physiological processes

during periods of limited water availability. Further research in *Themeda* is needed to

investigate these processes.

443   Climatic correlations with g_{min} and $\%\Delta g_{min}$

T. triandra accessions revealed nuanced climate relationships with g_{min}, underscoring the

complexity of adaptive trait variation across environmental gradients. Under control

conditions, g_{min} showed no association with either annual (AP, AT) or seasonal (summer

temperature, summer rainfall) climate variables. This finding contrasts with our hypothesis

that accessions from hotter and drier regions would exhibit inherently lower g_{min} to limit

water loss. Instead, it suggests that cuticular permeability in this species under control

conditions is not tightly linked to climate-at-origin, especially under control conditions. A

comprehensive review by Duursma *et al.* (2019) found that g_{min} varies widely among species, but this variation is not clearly related to climate-of-origin. Very few studies have identified significant relationships between g_{min} and environmental variables. In a study on 42 conifer species (Brodribb *et al.*, 2014) found a relationship between g_{min} and rainfall at the species' origin, specifically for the driest quarter of the year, but their methodology involved gas exchange measurements rather than mass loss from detached leaves.

Under drought conditions, however, g_{min} displayed clear climate-linked trends, being typically higher in plants from drier and hotter sites (Figs 4c,d). This runs counter to classical expectations of drought adaptation, which predict reduced g_{min} as a water saving mechanism (Smith *et al.*, 2006). Instead, it suggests that in hot, dry climates, maintaining moderate levels of cuticle permeability may serve alternative functions—such as leaf surface cooling via residual transpiration (Marchin *et al.*, 2023). However, it is important to note that the magnitude of these trends was relatively modest—across the range of observed summer rainfall (75–436 mm), g_{min} increased by only 16.5%, and across the summer temperature range (21–32 °C), by 23.5%. Notably, MAT was not a significant predictor of g_{min} under either control or drought conditions, emphasizing that seasonal climate variables may be more informative than annual averages for understanding trait-environment relationships in this species.

The climate-linked pattern of g_{min} under drought—specifically, its positive relationship with summer rainfall in fig. 4c—may reflect an evolved alignment of trait means with long-term climate. However, within individual accessions, the short-term drought responses were not always consistent with this trend. For example, Pannawonica—an arid origin accession—showed a reduction in g_{min} under drought. In contrast, Dalby—from a wetter site—showed an increase in g_{min} under drought. This suggests that trait means (shaped by climate adaptation), and trait plasticity (plastic responses to acute stress) are not always aligned. Trait means likely reflect long-term selective pressures at population level, whereas plastic responses to short-term drought may be shaped by local adaptive strategies, trade-offs, or even stress-related responses. Recognizing that trait means and plasticity are distinct and potentially uncorrelated axes of variation is important for interpreting plant responses to environmental change. Future work should therefore aim to disentangle the genetic, physiological and ecological bases of

both axes, ideally by pairing analyses of traits in relation to long-term climate with

experiments designed to capture short-term plasticity under controlled-stress scenarios.

We also assessed the plasticity of g_{min} by examining the percentage change from

control to drought %Δg_{min} , under the expectation that accessions from harsher

environments might show greater reductions to conserve water. However, %Δg_{min} was not

related to any climatic variable (Fig. S7), suggesting that responsiveness of cuticular

conductance to drought is not predictable by climate in this species. This could imply that

g_{min} plasticity is under weak environmental selection, for example, or perhaps that

traits—such as osmotic adjustment or rooting depth—buffer water loss during drought

more effectively.

Taken together, these findings highlight that drought-expressed g_{min} is more strongly

shaped by climate-at-origin than baseline g_{min}, particularly in relation to summer climate

extremes. Yet, the direction of this relationship is opposite to classical water conservation

models, pointing to the multifaceted roles of cuticle conductance in plant adaptation.

Temperature dependent behaviour of g_{min}

The temperature response of g_{min} is often described as biphasic, monotonically increasing,

where conductance rises sharply beyond a transition point—typically around 35–40 °C.

However, not all species follow this pattern. For instance, *Phoenix dactylifera* shows an

invariant relationship between g_{min} and temperature (Bueno *et al.*, 2019). Additionally, in a

broader survey of tropical trees, only seven out of twenty-four species displayed this

biphasic response, while most showed weakly decreasing, U-shaped, or invariant trends

(Slot *et al.*, 2021). Our investigation into temperature response of g_{min} in *T. triandra*

accessions revealed a general trend of gradual decline in g_{min} with increasing temperature

across most accessions. Studies report that g_{min} can decrease, remain invariant, or show a

U-shaped response with increasing temperature, depending on species and environmental

context (Duursma *et al.*, 2019, Slot *et al.*, 2021, Garen & Michaletz, 2025). In some cases,

acclimation to higher growth temperatures leads to a reduction in g_{min}. For example, Zailaa

et al. (2025) found that Quercus shows a reversible response to temperature, with g_{min}

decreasing from 25–45 °C and then returning back to original steady state. Similarly, *T.*

triandra shows a decreasing trend in g_{min}, although we do not know if this response is a

reversible one. While direct studies on *T. triandra* or grasses in general are lacking, there is
broad evidence among other species showing that g_{min} can show variable temperature-
dependent trends.

Two accessions (Dalby and Forbes) showed a modest increase in g_{min} at intermediate
temperatures, followed by a subsequent decrease—a pattern not observed in other
accessions. Such non-monotonic trends may arise from reversible changes in wax fluidity.
This atypical pattern suggests the involvement of temperature-sensitive physiological
pathways or distinct differences in cuticular lipid composition, which are known to play a
crucial role in regulating water permeability (Schönherr *et al.*, 1979).

Arrhenius plots of g_{min} did not reveal a clear biphasic pattern, indicating the absence
of a distinct phase transition across accessions and suggesting that cuticle remains
relatively stable across the temperature range tested. We used fresh cohorts from well-
watered plants for each temperature treatment and leaves from the higher temperature runs
(50 °C and 55 °C) did not appear extremely dry or noticeably different from those from
lower temperatures. This suggests that markedly reduced water status was unlikely to have
prevented an increase in g_{min}, although relative water content (RWC) was not measured and
subtle differences cannot be excluded. Such differences could potentially contribute to the
absence of a clear T_p. However, these findings imply that *T. triandra* may lack a distinct
thermal threshold for cuticle conductance, and that any structural adjustments in the cuticle
occur progressively or are buffered by other physiological or biochemical mechanisms. The
absence of a common temperature response pattern across accessions further highlights the
complexity of thermal strategies within the species.

Conclusion

Our findings provide new insights into the drought and thermal responses of *T. triandra*
leaf cuticles and their relationship with environmental conditions, highlighting potential
mechanisms driving these patterns. The observed variability in g_{min} could inform the
selection of resilient genotypes for ecological restoration or breeding programs aimed at
mitigating the impacts of climate change. Future work should focus on the biochemical
composition of cuticular waxes and their role in modulating g_{min} under stress conditions.
Comparative transcriptomics or metabolomic analyses may also elucidate the genetic

underpinnings of these diverse responses. Additionally, field studies assessing the interplay
between cuticular properties, drought, and temperature in natural populations would
provide valuable context for our controlled experiment findings. While this study provides
important insights, the controlled conditions of the experiments of do not fully replicate the
complexity of field environments. Further research is needed to validate these findings
across broader spatial and temporal scales. Incorporating long-term monitoring of *T.*
triandra populations under varying climatic conditions will help refine our understanding
of the evolutionary and ecological significance of cuticular adaptations in response to
environmental change.

As climate change drives more frequent droughts and heatwaves, understanding
how different species/accessions regulate water loss through their cuticles and withstand
high temperatures is key to strengthening ecosystem resilience. Additionally, examining
how cuticular permeability interacts with other drought responses—such as osmotic
regulation and root structure—could improve predictions of plant survival under future
climatic conditions. To turn these insights into practical conservation strategies, future
studies should expand to include other grassland species and assess their responses in long-
term field trials.

Supplementary information

Supplementary data consist of the following: Table S1: Climate and origin of the *Themeda*
triandra accessions used in the study. Figure S1: Soil moisture across three days of drought
stress. Figure S2: Midday leaf water potential (MPa) under control and drought. Figure S3:
Relationship between midday leaf water potential and $\%\Delta g_{min}$. Figure S4: Photograph of
DroughtBox. Figure S5: Performance of DroughBox in maintaining stable conditions.
Figure S6: Relationship between g_{min} and annual climatic variables. Figure S7: Relationship
between $\%\Delta g_{min}$ and annual climatic variables.

Author contributions

572 AM, VJ and IJW designed and conducted the experiment, analysed the data, and drafted the
573 initial manuscript. IJW, BC, VJ and CJB assisted with data interpretation and manuscript
574 revisions. All authors reviewed and approved the final manuscript.

Conflict of interest

577 The authors declare no conflict of interest.

Funding

580 This research was supported by funding from the Australian Research Council (ARC)
581 Centre of Excellence (CoE) for Plant Success in Nature and Agriculture (CE200100015).
582 Anu Middha was supported by a Postgraduate Research Scholarship from Western Sydney
583 University and a PhD top-up scholarship from the ARC CoE.

Acknowledgements

586 This work was supported by funding from the Australian Research Council and from
587 Hawkesbury Institute for the Environment, Western Sydney University (WSU). We are
588 very grateful to Dr Craig Barton, WSU, for constructing the DroughtBox and providing
589 training in its use. We also thank Erick Calderon-Morale for his help in using the
590 DroughtBox during the early stages of this work.

Data availability

594 The raw data supporting this study have been uploaded to Zenodo (doi:
595 10.5281/zenodo.16884594) in restricted mode and will be made publicly accessible upon
596 acceptance.

References

Araus J, Febrero A, Vendrell P. 1991. Epidermal conductance in different parts of durum wheat
grown under Mediterranean conditions: the role of epicuticular waxes and stomata. *Plant, Cell &*
Environment **14**: 545-558
Asadyar L, de Felippes FF, Bally J, Blackman CJ, An J, Sussmilch FC, Moghaddam L, Williams B,
Blanksby SJ, Brodribb TJ, et al. 2024. Evidence for within-species transition between drought
response strategies in *Nicotiana benthamiana*. *New Phytologist* **244**: 464-476

Billon LM, Blackman CJ, Cochard H, Badel E, Hitmi A, Cartailler J, Souchal R, Torres-Ruiz JM. 2020.
The DroughtBox: a new tool for phenotyping residual branch conductance and its temperature
dependence during drought. *Plant, Cell & Environment* **43**: 1584-1589
Blair J, Nippert J, Briggs J. 2014. Grassland ecology. *Ecology and the Environment* **389**: 389-423
Boyer JS, Wong SC, Farquhar GD. 1997. CO2 and water vapor exchange across leaf cuticle
(epidermis) at various water potentials. *Plant physiology* **114**: 185-191
Brodribb TJ, McAdam SA, Jordan GJ, Martins SC. 2014. Conifer species adapt to low-rainfall
climates by following one of two divergent pathways. *Proceedings of the National Academy of*
Sciences **111**: 14489-14493
Bueno A, Alfarhan A, Arand K, Burghardt M, Deininger A-C, Hedrich R, Leide J, Seufert P, Staiger
S, Riederer M. 2019. Effects of temperature on the cuticular transpiration barrier of two desert
plants with water-spender and water-saver strategies. *Journal of Experimental Botany* **70**: 1613-
1625
Bueno A, Sancho-Knapik D, Gil-Pelegrín E, Leide J, Peguero-Pina JJ, Burghardt M, Riederer M.
2020. Cuticular wax coverage and its transpiration barrier properties in *Quercus coccifera* L. leaves:
does the environment matter? *Tree Physiology* **40**: 827-840
Burghardt M, Riederer M. 2003. Ecophysiological relevance of cuticular transpiration of deciduous
and evergreen plants in relation to stomatal closure and leaf water potential. *Journal of*
Experimental Botany **54**: 1941-1949
Burlett R, Trueba S, Bouteiller XP, Forget G, Torres Ruiz JM, Martin-StPaul N, Parise C, Cochard H,
Delzon S. 2025. Residual leaf water loss changes during drought, impacting on future predicted
plant mortality. *New Phytologist* **246**: 1001-1014
Busta L, Schmitz E, Kosma DK, Schnable JC, Cahoon EB. 2021. A co-opted steroid synthesis gene,
maintained in sorghum but not maize, is associated with a divergence in leaf wax chemistry.
Proceedings of the National Academy of Sciences **118**: e2022982118
Cameron KD, Teece MA, Smart LB. 2006. Increased accumulation of cuticular wax and expression
of lipid transfer protein in response to periodic drying events in leaves of tree tobacco. *Plant*
Physiology **140**: 176-183
Cheng G, Huang H, Zhou L, He S, Zhang Y, Cheng Xa. 2019. Chemical composition and water
permeability of the cuticular wax barrier in rose leaf and petal: A comparative investigation. *Plant*
Physiology and Biochemistry **135**: 404-410
Churchill AC, Zhang H, Fuller KJ, Amiji B, Anderson IC, Barton CV, Carrillo Y, Catunda KL,
Chandregowda MH, Igwenagu C. 2022. Pastures and climate extremes: impacts of cool season
warming and drought on the productivity of key pasture species in a field experiment. *Frontiers in*
plant science **13**: 836968
DaCosta M, Huang B. 2006. Osmotic adjustment associated with variation in bentgrass tolerance
to drought stress. *Journal of the American Society for Horticultural Science* **131**: 338-344
Dimopoulos N, Tindjau R, Wong DC, Matzat T, Haslam T, Song C, Gambetta GA, Kunst L,
Castellarin SD. 2020. Drought stress modulates cuticular wax composition of the grape berry.
Journal of Experimental Botany **71**: 3126-3141
Duursma RA, Blackman CJ, Lopéz R, Martin-StPaul NK, Cochard H, Medlyn BE. 2019. On the
minimum leaf conductance: its role in models of plant water use, and ecological and
environmental controls. *New Phytologist* **221**: 693-705
Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S, Seyboth K, Adler A, Baum I,
Brunner S, Eickemeier P. 2014. Contribution of working group III to the fifth assessment report of
the intergovernmental panel on climate change. *Climate change* **5**: 1-1552
Garen JC, Michaletz ST. 2025. Temperature governs the relative contributions of cuticle and
stomata to leaf minimum conductance. *New Phytologist* **245**: 1911-1923
Holmgren P, Jarvis PG, Jarvis MS. 1965. Resistances to carbon dioxide and water vapour transfer in
leaves of different plant species. *Physiologia Plantarum* **18**: 557-573

Hsiao TC, Acevedo E, Fereres E, Henderson D. **1976**. Water stress, growth and osmotic
adjustment. *Philosophical Transactions of the Royal Society of London* **273**: 479-500
Huang H, Burghardt M, Schuster A-C, Leide J, Lara I, Riederer M. **2017**. Chemical composition and
water permeability of fruit and leaf cuticles of *Olea europaea* L. *Journal of Agricultural and Food*
Chemistry **65**: 8790-8797
Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP,
Kessler M. **2017**. Climatologies at high resolution for the earth's land surface areas. *Scientific data*
4: 1-20
Lin M, Matschi S, Vasquez M, Chamness J, Kaczmar N, Baseggio M, Miller M, Stewart EL, Qiao P,
Scanlon MJ. **2020**. Genome-wide association study for maize leaf cuticular conductance identifies
candidate genes involved in the regulation of cuticle development. *G3: Genes, Genomes, Genetics*
10: 1671-1683
López R, Cano FJ, Martin-StPaul NK, Cochard H, Choat B. **2021**. Coordination of stem and leaf
traits define different strategies to regulate water loss and tolerance ranges to aridity. *New*
Phytologist **230**: 497-509
Machado R, Loram-Lourenço L, Farnese FS, Alves RDFB, de Sousa LF, Silva FG, Filho SCV,
Torres-Ruiz JM, Cochard H, Menezes-Silva PE. **2021**. Where do leaf water leaks come from?
Trade-offs underlying the variability in minimum conductance across tropical savanna species with
contrasting growth strategies. *New Phytologist* **229**: 1415-1430
Macková J, Vašková M, Macek P, Hronková M, Schreiber L, Šantrůček J. **2013**. Plant response to
drought stress simulated by ABA application: changes in chemical composition of cuticular waxes.
Environmental and Experimental Botany **86**: 70-75
Marchin RM, Medlyn BE, Tjoelker MG, Ellsworth DS. **2023**. Decoupling between stomatal
conductance and photosynthesis occurs under extreme heat in broadleaf tree species regardless of
water access. *Global Change Biology* **29**: 6319-6335
Martin-StPaul N, Delzon S, Cochard H. **2017**. Plant resistance to drought depends on timely
stomatal closure. *Ecology letters* **20**: 1437-1447
Mitchell M, Miller M. **1990**. *The identification of some common native grasses in Victoria.*
Rutherglen Research Institute, Victoria, Australia.
Muchow R, Sinclair T. **1989**. Epidermal conductance, stomatal density and stomatal size among
genotypes of *Sorghum bicolor* (L.) Moench. *Plant, Cell & Environment* **12**: 425-431
Overpeck JT. **2013**. The challenge of hot drought. *Nature* **503**: 350-351
Peel MC, Finlayson BL, McMahon TA. **2007**. Updated world map of the Köppen-Geiger climate
classification. *Hydrology and earth system sciences* **11**: 1633-1644
Petri M, Batello C, Villani R, Nachtergaele F. **2010**. Carbon status and carbon sequestration
potential in the world's grasslands. In: Michael Abberton, Richard Conant and C. Batello, eds.
Grassland carbon sequestration: management, policy and economics. Rome: 19-31.
Qariani L, El Jaafari S, Dekkaki M, Araus J. **2000**. Cuticular conductance, water use efficiency and
drought tolerance of durum wheat isolines of differing glaucousness. *Options Méditerranéennes*
Série A, séminaires Méditerranéens **40**: 315-318
R Core Team 2025. *R: a language and environment for statistical computing*. Vienna, Austria: R
Foundation for statistical Computing. Retrieved 10 Nov. 2025, https://www.r-project.org/.10 Nov.
2025.
Riederer M. **2006**. Thermodynamics of the water permeability of plant cuticles: characterization of
the polar pathway. *Journal of Experimental Botany* **57**: 2937-2942
Sack L, Scoffoni C 2010. *Minimum epidermal conductance (gmin, aka cuticular conductance)*
protocol. Retrieved 24 Dec 2023, https://prometheusprotocols.net/function/gas-exchange-and-
chlorophyll-fluorescence/stomatal-and-non-stomatal-conductance-and-transpiration/minimum-
epidermal-conductance-gmin-a-k-a-cuticular-conductance/.24 Dec 2023.

Sala OE, Stuart Chapin F, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E,
Huenneke LF, Jackson RB, Kinzig A. **2000**. Global biodiversity scenarios for the year 2100. *science*
287: 1770-1774
Schönherr J, Eckl K, Gruler H. **1979**. Water permeability of plant cuticles: the effect of temperature
on diffusion of water. *Planta* **147**: 21-26
Schreiber L, Schönherr J. **1990**. Phase transitions and thermal expansion coefficients of plant
cuticles: the effects of temperature on structure and function. *Planta* **182**: 186-193
Schuster A-C, Burghardt M, Alfarhan A, Bueno A, Hedrich R, Leide J, Thomas J, Riederer M. **2016**.
Effectiveness of cuticular transpiration barriers in a desert plant at controlling water loss at high
temperatures. *AoB Plants* **8**: plw027
Shepherd T, Wynne Griffiths D. **2006**. The effects of stress on plant cuticular waxes. *New*
Phytologist **171**: 469-499
Slot M, Nardwattanawong T, Hernández GG, Bueno A, Riederer M, Winter K. **2021**. Large
differences in leaf cuticle conductance and its temperature response among 24 tropical tree
species from across a rainfall gradient. *New Phytologist* **232**: 1618-1631
Smith S, Fendenheim D, Halbrook K. **2006**. Epidermal conductance as a component of dehydration
avoidance in *Digitaria californica* and *Eragrostis lehmanniana*, two perennial desert grasses.
Journal of Arid Environments **64**: 238-250
Snyman HA, Ingram LJ, Kirkman KP. **2013**. *Themeda triandra*: a keystone grass species. *African*
Journal of Range & Forage Science **30**: 99-125
Teuling AJ. **2018**. A hot future for European droughts. *Nature Climate Change* **8**: 364-365
Turner NC. **2018**. Turgor maintenance by osmotic adjustment: 40 years of progress. *Journal of*
experimental botany **69**: 3223-3233
Wasson AP, Richards R, Chatrath R, Misra S, Prasad SS, Rebetzke G, Kirkegaard J, Christopher J,
Watt M. **2012**. Traits and selection strategies to improve root systems and water uptake in water-
limited wheat crops. *Journal of experimental botany* **63**: 3485-3498
Yang SU, Kim H, Kim RJ, Kim J, Suh MC. **2020**. AP2/DREB transcription factor RAP2. 4 activates
cuticular wax biosynthesis in Arabidopsis leaves under drought. *Frontiers in plant science* **11**: 895
Zailaa J, Scoffoni C, Brodersen CR. **2025**. Stomatal closure as a driver of minimum leaf
conductance declines at high temperature and vapor pressure deficit in Quercus. *Plant Physiology*
197: kiae551

Figures

Figure 1. (a) Köppen-Geiger climate classification map of Australia adapted from Peel *et al.* (2007) showing the distribution of various climate zones. The locations of six *Themeda triandra* accessions included in this study are marked on the map. (b) Scatterplot illustrating the annual temperature (AT) and annual precipitation (AP) of the selected accessions, highlighting the climatic differences across their native habitats. (c) Köppen-Geiger climate classifications represented by distinct colours corresponding to specific climate types based on temperature and precipitation patterns. (The Köppen-Geiger classification system utilizes a combination of letters and colours to denote different climate types. For instance, 'BWh' in red denotes a hot desert climate with high temperatures and minimal precipitation. These classifications are visually represented through distinct colour codes to facilitate easy identification on the map)

Figure 2. Box plot showing the distribution of g_{min} under control and drought treatments. The boxes represent the interquartile range (IQR), the black horizontal line within each box represents the median, and whiskers extend to $1.5 \times IQR$ (n = 24). Red line inside the boxes represent mean. The p-value (0.3163; paired t-test) reflects the statistical comparison between treatments.

Figure 3. Box plots showing the variation in g_{min} across six accessions of *Themeda triandra* under control and drought treatments. Significant differences between treatments within accessions are denoted by asterisks (*$p < 0.05$, ***$p < 0.001$; paired t-test). The boxes represent the interquartile range (IQR), the black horizontal line within each box represents the median, and whiskers extend to $1.5 \times$ IQR (n = 4). Red line inside the boxes represent mean.

Figure 4. Relationship of g_{min} (mmol m^{-2} s^{-1}) with climate-at-origin variables under control (a–b) and drought (c–d) conditions across *Themeda triandra* accessions. g_{min} is plotted against summer rainfall (a, c) and summer temperature (b, d). Data points represent individual measurements. Black lines and shaded areas depict fitted values and 95% confidence intervals from linear mixed effects models, with accession included as a random effect.

Figure 5. Relationship between $\%\Delta g_{min}$ (mmol m^{-2} s^{-1}) and climate-at-origin variables across *Themeda triandra* accessions. (a) Summer rainfall (mm) and (b) summer temperature (°C) are shown as predictors. Each point represents an individual observation. Relationships were evaluated using linear mixed-effects models, with accession included as a random effect.

Figure 6. Temperature dependence of g_{min} (a–f) and corresponding Arrhenius plots (g–l) across a temperature range of 30–55□ at a constant absolute humidity of 18 g(H_2O)m^{-3}