The Function of *Efhd1*[+] Telocytes in the Synovial Lymphatic System and Inflammatory-Erosive Arthritis

Running Title: Telocyte Function in the Synovial Lymphatic System

Yue Peng[1,2], H. Mark Kenney[1,2], Sean Lydon[2], Karen L. de Mesy Bentley[1,2], Lianping Xing[1,2], Benjamin D. Korman[1,3]

Christopher T. Ritchlin[1,3] & Edward M. Schwarz[1,2*]

[1]Center for Musculoskeletal Research; [2]Department of Pathology & Laboratory Medicine; [3]Department of Medicine, Division of Allergy, Immunology, Rheumatology; University of Rochester Medical Center

*Corresponding Author; Edward_Schwarz@URMC.Rochester.edu; (585) 276-5609; University of Rochester Medical Center, 601 Elmwood Ave, Box 665, Rochester, NY, 14642, USA.

Funding: F30AG076326, T32GM007356, R01AG059775, R01AR069000, R01AR056702, and P30AR069655

Conflict of interest: No conflicts of interest to disclose.

Keywords: Telocyte, Mast Cell, Lymphatic Vessel, Electron Microscopy, Inflammatory-Erosive Arthritis, Animal Models

Telocyte Function in the Synovial Lymphatic System

Abstract (247 Words of 250)

Resting collecting lymphatic vessels (cLVs) sense edema in distal joints and initiate contractions

via unknown mechanisms. Rheumatoid arthritis (RA) patients have lymphatic drainage

deficiencies from affected joints, and defects in the synovial lymphatic system exacerbate

inflammatory-erosive arthritis in animal models. To understand this, we generated *Efhd1*-

CreER^{T2} and *Myoc*-CreER^{T2} mice for cell-specific genetic gain and loss of function studies.

These mice were crossed with tdTomato reporter (Ai9) mice, and studies showed selective

tamoxifen-induced transgene expression in CD31^{-}/CD34^{+} telocyte-like cells in knee and ankle

synovium, and in networks physically associated with mast cells proximal to popliteal lymphatic

vessels (PLVs). Consistent with the known loss of CD31^{-}/CD34^{+} telocyte in RA synovium,

these cells were also decreased in TNF-tg knees and partially recovered by anti-TNF treatment.

Ultrastructural and gene expression studies confirmed a distinct telocyte phenotype versus

closely related fibroblasts. *In vivo* depletion studies in tamoxifen-treated *Efhd1*-CreER^{T2} and

Myoc-CreER^{T2} mice crossed to diphtheria toxin alpha-floxed (DTA^{flox}) mice demonstrated

telocyte requirements for physiologic lymphatic drainage and resolution of joint inflammation

and focal erosions from zymosan-induced arthritis in the knee. *In vitro* studies demonstrated

increased sensitivity to osmotic shock and decreased motility versus fibroblasts, and telocyte

potential to differentiate into myofibroblasts on stiff matrix. Collectively, these findings support

a model of joint homeostasis in which osmotic pressure-sensing telocyte networks extend from

the synovium into mast cells proximal to joint-draining cLVs, and telocyte loss is associated with

defects in the synovial lymphatic system and increased susceptibility to joint inflammation and

structural damage from arthritis.

Teaser

This study discovered a unique cell type (telocytes) responsible for draining inflammation from

arthritic joints.

INTRODUCTION

The lymphatic system reabsorbs excess interstitial fluid and protein for return to the central veins, and dysfunction is associated with lymphedema in affected tissues, exacerbation of chronic inflammatory diseases, and fibrosis (*1*). While some lymph migrates passively through unidirectional bicuspid lymphatic vessel valves secondary to interstitial fluid pressure and passive compression by adjacent tissue movement (*2*), the majority of lymph is transported by the contractile activity of collecting lymphatic vessels (cLVs) (*3*). Central to cLV functions are tightly regulated contraction frequency changes (*4*), known as 'pressure-induced lymphatic chronotropy' (*5*), which range from ~3 to 20 contractions/min as intraluminal pressure increases from 0.5 to 10 cmH$_2$O (*6, 7*) to enable coupled lymph transport with absorption via lymphatic capillaries (*8*). Recently, major advances have greatly enhanced our understanding of intraluminal pressure-induced lymphatic chronotropy and intrinsic cLV pacemaking by lymphatic muscle cells (LMCs) (*5, 6, 9, 10*). Interestingly, LMCs contain functional and molecular characteristics of both myocardium (cardiac muscle) and blood vessels (vascular smooth muscle), as evidenced by concomitant rapid cardiac-like phasic contractions to generate flow and smooth-muscle-like slow, long-lasting tonic contractions to regulate flow (*11*). However, several critical knowledge gaps remain including the identity of cells that sense acute edema in tissue distal to cLVs to trigger enhanced lymphatic clearance by initiating pacemaking signals in LMCs, and the mechanisms that underlie the loss of cLV contractility in lymphedema (*12*).

We previously demonstrated that in rheumatoid arthritis (RA), lymphatic dysfunction is a critical factor in joint inflammation, damage, and flare which can progress independent of autoimmunity (*13*). Evidence of synovial lymphatic system dysfunction in osteoarthritis (OA)

also exists (*14*). Studies on innate immune mechanisms identified increased expression of the

lymphangiogenic factor vascular endothelial growth factor C (VEGF-C) in monocytes and

osteoclasts, together with lymphangiogenesis in inflamed joints (*15*). Subsequent functional

studies demonstrated that loss of VEGF-C signaling together with decreased lymphangiogenesis

exacerbates inflammatory-erosive arthritis (*16*), while VEGF-C therapy ameliorates

inflammatory arthritis by increasing lymphangiogenesis (*17*). Longitudinal MRI (*18, 19*), power

Doppler ultrasound (PDUS) (*20*), and near-infrared (NIR) imaging of injected indocyanine green

(ICG) dye (*7, 21*), identified three distinct mechanisms of lymphatic dysfunction in murine

models of RA. The first is mediated by release of local vasodilators (e.g. nitric oxide) from

inflammatory cells that contribute to reduced lymphatic flow (*13, 22*). Another involves

accumulation and translocation of B cells into the sinuses of joint-draining lymph nodes,

obstructing passive lymphatic flow (*19, 21, 23*). Finally, we observed the loss of cLV

contractions during arthritic progression in the setting of chronic joint inflammation (*20*). In

support of the translational relevance of these findings are data demonstrating that RA patients

display similar lymphatic dysfunction, including reduced cLV contractility and lymphatic

clearance compared to healthy controls (*24-26*), which is now a high priority area of research.

Mast cells are recognized as effective regulators of LMC contractility due to their

anatomical proximity and ability to produce, store, and release various inflammatory and

vasoactive mediators (*27*). Additionally, mast cells are positioned throughout the RA synovial

sublining, where they comprise 5% or more of the expanded synovial cell population, and

microanatomic clusters of mast cells are present in pannus tissue proximal to cartilage and bone

erosion (*28*). Based on these findings and evidence from preclinical and clinical studies

supporting pro-inflammatory and catabolic roles for mast cells in RA (*28, 29*), we investigated

mast cell effects on joint-draining lymphatics and inflammatory-erosive arthritis in tumor

necrosis factor-transgenic (TNF-tg) mice (*30*) and their wild-type (WT) littermates. We found

that popliteal lymphatic vessels (PLVs) are surrounded by an expanded frequency of

$MCT^+/MCPT1^+/MCPT4^+$ mast cells in TNF-tg mice, and the percentage of peri-PLV

degranulating mast cells was inversely correlated with lymphatic clearance (*31*). We also

identified a unique population of $MCT^+/MCPT1^-/MCPT4^-$ mast cells embedded within the PLV

cellular architecture, but the function of these cells remains unknown (*31*). Genetic and

pharmacologic mast cell loss of function experiments in TNF-tg mice demonstrated severe

lymphatic defects, which were associated with exacerbated inflammatory-erosive arthritis (*31*).

Collectively, these findings led us to surmise that while $MCT^+/MCPT1^+/MCPT4^+$ mast cells

have a well-established role in inflammation, particular mast cell subtypes are also required for

normal lymphatic function, proposed to be mediated by $MCT^+/MCPT1^-/MCPT4^-$ mast cells

embedded within PLVs. However, several unanswered questions remain including: i) the

anatomy and physiology of mast cell integration into the synovial lymphatic system, ii) the

mechanisms that facilitate recognition of increased joint fluid by the distal cLV, iii) the resulting

increased contraction frequency during the early phase of inflammation, and iv) how this

compensatory response is lost during chronic arthritis, remain be elucidated.

To address these questions, we developed conditional-inducible transgenic mice for

genetic gain and loss of function studies initially considered to specifically target LMCs. Single-

cell RNA sequencing (scRNAseq) studies identified three genes (*Efhd1, Myoc & Pla1a*) in

stromal cell population selectively expressed in PLVs compared to adjacent blood vessels (*32*).

We then generated three transgenic mouse lines in which a *CreER^{T2}* cassette was knocked into

these genes, and the resulting animals were crossed to B6.Cg-*Gt(ROSA)26Sor^{tm9(CAG-tdTomato)Hze}*/J

(Ai9) (*33*) and B6.129P2-*Gt(ROSA)26Sor^{tm1(DTA)Lky}*/J (DTA) (*34*) mice for lineage tracing and

loss of function studies, respectively. Herein, we describe the construction of these models, and

the serendipitous finding of tamoxifen-inducible gene expression in peri-PLV spindle-shaped

cells with very long cytoplasmic extensions that were physically integrated into mast cells, while

there was no expression in the predicted LMC population (*32*), which has been further described

in more recent scRNAseq studies (*9*). We hypothesized the identified peri-PLV spindle-shaped

cells to be telocytes, which are unique mesenchymal cells found in most tissues and are

characterized by a small cell body and long, thin, cytoplasmic extensions called telopodes (*35*).

Of note are the various telocyte functions that include extracellular matrix (ECM) synthesis for

tissue structural support, cell-to-cell communication, regulation of immune responses, and

pacemaking of smooth muscle contractions (*36*). To confirm these characteristics supporting

telocyte identity and determine the function of telocytes within the synovial lymphatic system,

we performed various *in vivo* and *in vitro* studies of the tdTomato-positive (tdT$^+$) cells. Together,

we tested the hypotheses that: 1) networks of contiguous telocytes extend from ankle synovium

to PLVs, 2) telocytes are functionally distinct from lymphatic fibroblasts, and 3) synovial

telocytes differentiate into myofibroblasts when grown on stiff ECM.

RESULTS

Generation of Novel CreER^{T2} Mouse Models for Genetic Targeting of Telocytes in the

Synovial Lymphatic System

We recently described single-cell RNA sequencing (scRNAseq) analysis of lymphatic vessels in

which we identified *Efhd1, Myoc,* and *Pla1a* as PLV-selective genes (*32*). Based on these

findings, we generated three knock-in mouse lines in which a *CreER^{T2}* cassette was inserted into

these genes of embryonic stem cells (Fig. 1a), and the resulting mice were crossed to Ai9

reporter mice to assess tamoxifen-induced in vivo gene targeting (Fig. 1b & c). *Ex vivo* whole-

mount immunofluorescent microscopy (WMIFM) revealed the presence of elongated tdT$^+$ cells

that were preferentially clustered in networks around PLVs vs. blood vessels from tamoxifen-

treated *Efhd1*-CreER$^{+/-}$x Ai9$^{+/-}$ and *Myoc*-CreER$^{+/-}$ x Ai9$^{+/-}$ mice (Fig. 1d). Tamoxifen treatment

of younger mice (P10-P13) resulted in a similar pattern of PLV adventitial tdT$^+$ cells, suggesting

that they maintain their identity from early postnatal development (Supplemental Fig. 1a-b).

These tdT$^+$ cells were not found in tissues from tamoxifen-treated *Pla1a*-CreER$^{+/-}$ x Ai9$^{+/-}$ mice,

and these mice were not further analyzed (Supplemental Fig. 1b).

Interrogation of the PLV WMIFM revealed two distinct tdT$^+$ populations of fibroblastic

cells that contained very long dendritic processes phenotypically consistent with telopodes (Fig.

2a). One population extended longitudinally along the PLV (Fig. 2b), and the other existed in

telocyte arrays within adipose tissue adjacent to PLVs (Fig. 2c). To gain wholistic insight on the

extent of tdT$^+$ cell arrays along the PLV, we performed WMIFM on tissues from tamoxifen

treated *Efhd1*-CreER^{T2} x Ai9 mice with antibodies against αSMA and CD34 to label the vascular

smooth muscle and mast cells. Confocal images of soft-tissue extending from the ankle to the

knee containing the PLVs were reconstructed in 3D and stitched together (Figs. 2d-h). The high-

power images confirmed the presence of tdT$^+$ cell arrays adjacent to PLVs that contained CD34$^+$

cells with a mast cell morphology. The low-resolution images revealed the networks of tdT$^+$ cells

that paralleled the vessels from the ankle synovium to cLV.

Telocyte Function in the Synovial Lymphatic System

Previous investigations of lymphatic dysfunction revealed an enrichment of mast cells

around lymphatic vessels (*31*), which prompted us to examine these distinct tdT$^+$ cells and their

interactions within the peri-PLV tissue more closely. Scrutiny of transmission electron

microscopy (TEM) images of PLVs revealed elongated cells in tamoxifen-treated *Efhd1*-

CreER$^{+/-}$x Ai9$^{+/-}$ and *Myoc*-CreER$^{+/-}$ x Ai9$^{+/-}$ mice with morphological characteristics consistent

with the tdT$^+$ cells identified by WMIFM (Fig. 3a). Moreover, these cells have characteristic

ultrastructural features of telocytes (*35*) (i.e. spindle-shaped polygonal cells with a large single

nucleus and scant cytoplasm and thinly attenuated cytoplasmic extensions). These cells were

found to form intimate contacts with mast cells in what appears to be specialized junctional

structures at their plasma membrane interfaces suggesting active intercellular communication

(Fig. 3a). To validate these observations and better characterize the spatial relationships between

telocytes and mast cells, we conducted 3D immunofluorescent confocal microscopy. This

analysis demonstrated that mast cells form direct contacts with the two tdT$^+$ telocyte-like

populations (Fig. 3b and Supplemental Video 1). The spatial distribution and interaction patterns

observed through 3D confocal imaging precisely matched our WMIFM and TEM observations,

confirming that both mural and peri-PLV telocyte populations engage in direct cellular

interactions with mast cells. Taken together, these findings suggest a previously unrecognized

cellular partnership between telocytes and mast cells along lymphatic vessels, potentially

representing a novel signaling axis in cLV regulation.

To provide additional confirmation that the tdT$^+$ cells are telocytes, we reanalyzed our

published scRNAseq data on murine PLV-associated cells (*32*) to identify the *Pdgfra$^+$/Cd34$^+$*

population known to contain telocytes and adventitial fibroblasts (*9*), which produced a uniform

manifold approximation and projection (UMAP) of these closely related populations (Fig. 4a).

Telocyte Function in the Synovial Lymphatic System

To characterize their transcriptome distinctions and similarities, we generated UMAPs of known

marker genes selectively expressed in either one cell type or both telocytes and fibroblasts. The

results confirmed that *Myoc, Efhd1, Pla1a, Dpp4, Pcsk6,* and *Ppp2r2b* genes (*9*) are

preferentially expressed in PLV-associated telocytes (Fig. 4b). The results also confirmed that: i)

the pan marker genes *CD34, Pdgfra,* and *Pdpn* have equivocal expression in both PLV-

associated telocytes and fibroblasts, ii) both cell types produce EMC gene products including

Col1a1, Fn1, and *Lama4*, and iii) *Comp, Dkk3* and *Angpt2* are specific marker genes for PLV-

associated fibroblasts. We also performed a cellular trajectory analysis of the scRNAseq data

from WT and TNF-tg PLVs using Monocle 3, which predicts that telocytes have the potential to

differentiate into fibroblasts (Fig. 4c top). Moreover, we found that there was a 47.7% decrease

in the proportion of telocytes relative to fibroblasts in TNF-tg mice relative to their WT

littermates (Fig. 4c bottom).

To elucidate synovial peri-lymphatic telocyte loss in TNF-tg mice, we performed

scanning electron microscopy (SEM) of PLVs and immunohistochemistry (IHC) on knee

synovium. SEM of WT PLVs revealed apparent telocyte cell bodies embedded within a very rich

ECM structure, and associated mast cells on top of this ECM (Fig. 4d). In contrast, SEM of

TNF-tg PLVs displayed varying degrees of degeneration from partial ECM loss and exposed

LMCs (Fig. 4e) to extensive ECM and LMC loss (Fig. 4f), which is consistent with their

contractility defects observed *ex vivo* (*25*). Consistent with the loss of CD31$^-$/CD34$^+$ telocytes

observed in RA synovium determined by IHC (*37*), we also observed a significant decrease in

CD31$^-$/CD34$^+$ telocyte-like cells in the synovium of TNF-tg mice receiving placebo treatment

compared to WT, and their partially recover from the TNF-tg mice receiving 6 weeks of anti-

TNF treatment (*26*) (Fig. 4g).

Telocyte Function in the Synovial Lymphatic System

To confirm that the CD31$^-$/CD34$^+$ synoviocytes are tdT$^+$ telocytes in our transgenic

mouse models, we performed flow cytometry, scRNAseq, and immunohistochemistry on lower

limb joint tissues from tamoxifen-treated *Efhd1*-CreER$^{+/-}$ x Ai9$^{+/-}$ and *Myoc*-CreER$^{+/-}$ x Ai9$^{+/-}$

mice (Fig. 5a). Flow cytometry of knee and ankle synoviocytes revealed consistent profiles of

tdT$^-$ and tdT$^+$ cells (Supplemental Fig. 2). The scRNAseq analysis of the flow sorted tdT$^+$

synoviocytes identified 13 distinct cell clusters classified by marker gene expression

(Supplemental Fig. 3a-b). To better characterize telocytes in the synovial cells population we re-

clustered the mesenchymal and stromal cell compartments (Supplemental Fig. 5c-e). The clusters

in the synovium closely resembled the telocyte and fibroblast clusters in lymphatics (Fig. 5b).

Specifically, only cells in the upper portion of the UMAP express *Efhd1* and *Pla1a* together with

the conventional non-specific telocyte markers *Cd34, Pdgfra, and Pdpn* (*37, 38*), which are

broadly expressed throughout the UMAP (Fig. 5b). In contrast to its selective expression in PLV-

associated telocytes, *Myoc* is broadly expressed in tdT$^+$ synoviocytes with other ECM genes

including: *Col1a1, Fn1,* and *Lama4* (Fig. 5b). Interestingly, cells in the lower portion of this

UMAP express markers associated with RA fibroblast-like synoviocytes (FLS) (*39, 40*) such as,

Cdh11, Comp, and *Dkk3* (Fig. 5b). Consistent with our observation that the *Efhd1*$^+$/*Pla1a*$^+$ tdT+

synoviocytes are telocytes, these cells are also negative for the endothelial marker gene

CD31/PECAM1 (*41*), and the mesenchymal stromal cell marker *Cd90/Thy-1* (*42*) (Fig. 5b and

Supplemental Fig. 4). However, they are also negative for *FoxL1* (*43*) and *Ano1* (*44*) (Fig. 5b),

previously reported telocyte genes identified in gastrointestinal interstitial cells of Cajal (ICC)

(*35*).

Based on these scRNAseq results, we hypothesized the existence of tissue-specific

telocyte subsets and performed IHC studies on various tissues from tamoxifen-treated *Efhd1*-

CreER$^{+/-}$x Ai9$^{+/-}$ and *Myoc*-CreER$^{+/-}$ x Ai9$^{+/-}$ mice. The results from knee and ankle cryosections

confirmed that tdT$^+$ synoviocytes are CD31$^-$/CD34$^+$ cells with telocyte morphology and are

distinct from Comp$^+$ FLS and CD31$^+$/CD34$^+$ endothelial cells (Fig. 5c). Given that telocytes are

also known to express c-Kit (*35*), a critical gene for mast cells, we examined if c-Kit is also

required for telocyte development and survival. Thus, we also included joints from mast cell

deficient *Kit$^{W-sh/W-sh}$* (cKit$^{-/-}$) mice and confirmed that they have similar levels of synovial CD31$^-$

/CD34$^+$ telocyte-like cells compared to WT mice (Supplemental Fig. 5), which affirms that

lymphatic disfunction in these animals (*31*) is likely due to the loss of mast cells and not loss of

telocytes that also express c-Kit. Additionally, we found tdT$^+$ telocyte-like cells in tendon, skin

hair follicles, and within the ganglion cell layer of the retina, but no tdT$^+$ cells were found among

the c-Kit$^+$ ICC in intestine (Supplemental Fig. 6). Therefore, despite the absence of definitive

markers or functional assays to prove telocyte identity, the preponderance of our ultrastructural,

histologic, and gene expression data show that the *Efhd1*-CreER^{T2} and *Myoc*-CreER^{T2} mice

target tamoxifen-induced gene expression to telocyte subtypes in some but not all tissue types.

***In vivo* Depletion of Telocytes Decreases Lymphatic Function and Exacerbates**

Inflammatory-Erosive Arthritis

To test the hypothesis that in vivo depletion of *Myoc* and *Efhd1* expressing telocytes in the

synovial lymphatic system are critical for lower limb lymphatic function in mice, we generated

an inducible depletion model by crossing the telocyte-specific CreER mice with DTA^{floxed} mice,

followed by tamoxifen treatment to induce diphtheria toxin-α *in vivo* depletion of the gene

targeted cells as outlined in Figure 6a. To estimate *in vivo* depletion efficiency in these models,

we performed WMIFM on PLVs and IHC knee joints from tamoxifen-treated double and single

(control) transgenic mice and assessed CD31^{-}/CD34^{+} cells. WMIFM of PLVs demonstrated a

dramatic DTA elimination of telocytes on the vessel surface and the arrayed telocytes in the

adjacent adipose tissue (Fig. 6b). IHC analysis of the knees confirmed a ~60% reduction in the

number of CD31^{-}/CD34^{+} synoviocytes (Figs. 6c-d). We repeated this experiment to assess

telocyte depletion effects on lymphatic drainage. Cohorts of single and double transgenic mice

underwent NIR-ICG imaging prior to tamoxifen treatment and 6hrs following the last tamoxifen

injection. We observed no tamoxifen effects on DTA$^{f/-}$ mice contrasted with a significant

reduction of ICG clearance in the tamoxifen-treated double transgenic mice (Figs. 6e-f). These

findings support our hypothesis that synovial lymphatic system telocytes regulate synovial

lymphatic clearance, although these results are confounded by our inability to assess systemic

effects and DTA deletion of non-telocytes (e.g. fibroblasts that differentiated from telocytes after

tamoxifen treatment).

To assess the effects of lymphatic dysfunction following telocyte deletion on acute joint

inflammation, we utilized the zymosan-induced arthritis (ZIA) model (*45*) as outlined in Figure

7a. Longitudinal ultrasound confirmed a 10-fold increase in synovial volume following knee

injection with zymosan vs. PBS controls (Fig. 7b-d). Moreover, zymosan-induced 3.2-fold

greater synovitis in telocyte depleted mice, and while this acute synovitis commenced resolution

in WT mice after 7 days (Fig. 7c), the increased synovial volume was unabated at day 14 in

telocyte depleted mice (Fig. 7d). Histology of the knee joints corroborated the imaging results

demonstrating increased pannus tissue in knees with ZIA from mice with depleted telocytes

(Figs. 7e-f). Remarkably, the histology also revealed resorbed cortical bone surfaces in DTA and

zymosan groups (Fig. 7e). To assess telocyte depletion and ZIA effects on focal erosions and

osteoclast numbers, we performed histomorphometry and micro-CT analyses on these femurs.

Telocyte Function in the Synovial Lymphatic System

The results showed increased numbers of tartrate-resistant acid phosphatase (TRAP) positive

osteoclasts on resorbed cortical bone surfaces in some of the femurs from zymosan treated mice

(Supplemental Fig. 7). Surprisingly, we also observed large numbers of osteoclasts in some

femurs from telocyte-depleted mice injected with PBS. Micro-CT further demonstrated that

acute ZIA failed to induce focal erosions in the knees of WT mice within 14 days, whereas it

induced extensive erosions in the patellar groove of telocyte-depleted mice at this time point

(Fig. 7g-h). Together with our lymphatic readouts, these data demonstrate that telocyte depletion

reduces lymphatic function, and upon challenge with acute inflammatory arthritis, leads to more

persistent inflammation, exacerbated synovitis, and rapid focal cortical bone loss.

Synovial Lymphatic System Telocyte Function and Differentiation

To investigate the potential functions of telocytes within the synovial lymphatic system,

we developed an in vitro model in which PLVs from tamoxifen treated *Efhd1*-CreER^{T2} x Ai9 and

Myoc-CreER^{T2} x Ai9 mice were cultured to generate a mixture of growing tdT$^+$ telocytes and

non-fluorescent fibroblasts as described in Figure 8a. As telocytes could be the cells that sense

osmotic pressure in the joint from excess interstitial fluid, and intracellular Ca^{++} signaling by

telocytes has been studied (*46*), we assessed intracellular release of Ca^{++} following exposure to

sucrose and demonstrated that telocytes are more sensitive to osmotic shock than adjacent

fibroblasts (Fig. 8b), suggesting that signaling in response to hypertonic interstitial fluid may be

one of their physiological functions. To further demonstrate that telocytes are functionally

distinct from adjacent fibroblasts, we examined the ability of these primary telocyte and

fibroblast mixed cultures to migrate through Matrigel. The results demonstrated that while both

cell types have a chemotactic response to TGF-β, fibroblasts are more invasive than telocytes

(Fig. 8c).

Lastly, we tested the hypothesis that telocytes differentiate into robust ECM producing

fibroblasts, as predicted by our scRNA-seq trajectory analysis (Fig. 4c) and SEM of PLVs (Fig.

4d). These experiments utilized FACS purified cultures of tdT$^+$ telocytes derived from PLVs

harvested from tamoxifen-treated *Efhd1*-CreER^{T2} x Ai9 and *Myoc*-CreER^{T2} x Ai9 mice, which

retained their telocyte phenotype as evidenced by their long telopodes and their CD34 and Myoc

protein expression (Supplemental Fig. 8). Early passage telocytes were cultured on ECM of

varying stiffness (0.2 kPa (low), 8 kPa (intermediate), and 64 kPa (high), and harvested for bulk

RNAseq. The results demonstrated that telocytes retain their *Efhd1* and *Myoc* expression on low

stiffness ECM but differentiate into other types of mesenchymal cells, including fibroblasts and

myocytes, on stiff matrix (Fig. 9).

DISCUSSION

RA is a complex immune-mediated inflammatory disorder with divergent mechanisms of

disease progression (*47, 48*). Recent high-dimensional studies of RA tissues revealed multiple

cell types and factors that contribute to chronic inflammation in affected joints (*49, 50*), and this

knowledge catalyzed promising treatment strategies for RA patients (*51*). Despite these

therapeutic advances, major unmet clinical needs remain for a large fraction of RA patients that

fail to achieve remission and ~30% who are refractory to all current therapies (*47, 48*). This

treatment shortfall underscores the need to find novel disease mechanisms and treatment

strategies. With this objective in mind, we focused on the autoimmune independent pathologies

of inflammatory-erosive arthritis observed in TNF-tg mice (*52*). This research led to the

discovery of the synovial lymphatic system (*53*), an interactive network critical for joint

homeostasis (*54*), which is dysfunctional in RA and osteoarthritis, and this altered function

correlates with disease progression and flare (*13, 14*). Thus, elucidation of the mechanisms that

regulate the synovial lymphatic system and its dysfunction during chronic arthritis opens new

avenues of investigation for the development of new therapies. Here we demonstrate that DTA-

mediated telocyte deletion inhibits normal lymphatic function (Fig. 6) and exacerbates

inflammatory-erosive arthritis (Fig. 7), which raises the possibility that these cells might serve as

a novel target for treating RA patients who are refractory to all current therapies.

In contrast to the circulatory system, which is responsible for blood flow, and where

development, regulation, and repair have been defined in great cellular and molecular detail,

many fundamental questions about lymphatics remain unanswered, including the origins of

LMCs and cLV maintenance and repair (*3, 55*). Critical questions pertaining to the mechanisms

that control cLV contractions also remain unanswered. To address lymphatic contraction

mechanisms, Zawieja et al utilized conditional-inducible CreER^{T2} transgenic mice to assess the

innate pacemaking potential of mast cells, telocytes, pericytes, fibroblasts and LMCs in cLVs ex

vivo (*9*). These studies failed to identify Ca^{++} events in PDGFRα$^+$ or c-Kit$^+$ adventitial cells that

were in phase with cLV contractile activity. These investigators were unable to consistently elicit

contractions following optogenetic stimulation via a light-activated cation channel rhodopsin2

(ChR2), indicating that adventitial cells (e.g. telocytes) are not spontaneous pacemakers.

However, they did find that MYH11$^+$ LMCs exhibited asynchronous diastolic Ca^{++} events that

were dynamically modulated by pressure, and these cells also propagated contraction in response

to ChR2 photo-stimulated depolarization (*9*). Taken together with other studies (*5, 56*), these

results support a long-standing model of LMCs as the intrinsic pacemaker of cLVs (*57*). While

these studies largely explain the mechanisms responsible for intraluminal pressure-dependent

spontaneous cLV contractions, they do not provide information on the initiating signal for distal

tissue LMC contractions in response to edema (e.g. joint effusions).

In retrospect, our finding that ablation of peri-PLV telocytes is associated with lymphatic

dysfunction is not surprising, as pacemaking in smooth muscle-invested organs is known to be

controlled by this cell type (*6*). Indeed, pioneering studies of gastrointestinal smooth muscle

found that pacemaking is initiated by ICC (*58, 59*), and subsequent studies of the lower urinary

tract (*60*), gallbladder (*61*), and oviduct (*62*) found similar cells, now generically referred to as

telocytes (*35, 36*). Additionally, cells characteristic of telocytes have been identified in LVs (*63*),

and prior studies identified methylene blue staining peri-LV $CD34^+/c-Kit^+$ cells with telocyte

morphology and telopode-like structures extending to LMCs (*9, 63*).

It also reasons that telocytes are involved in cLV contractions, as LMCs are akin to

smooth muscle cells that lack the ionic mechanisms required for action potential regeneration

and electrical signals that efficiently propagate between muscle cells (*64*). Thus, LMCs likely

rely on a telocyte network for signal conduction. Consistently, telocytes mediate rhythmic

electrical activity in other tissues (*65*), which is believed to depend on ion channels for

generating pacemaking currents (*66, 67*). Interestingly, electrically coupled cells are typically

connected by gap junctions (*63*), and evidence demonstrating this physical link between

telocytes and LMCs does not exist (*6*). Studies have also failed to demonstrate functional

electrical communication between telocytes and LMCs or a telocyte Ca^{++} clock to drive the

rhythmic cLV contractions observed *ex vivo* (*9*). However, it is conceivable, based on our

findings, that an intermediary cell provides the depolarizing signals to LMCs to initiate

pacemaking, which we posit to be peri-cLV mast cells integrated into telocytes (Fig. 3). Mast

cells produce, store, and release various inflammatory and vasoactive mediators (*27, 68*). These

mediators influence lymphatic pumping by affecting LMC contractility (*69*), and may be

responsible for directly providing the initiating pacemaking signal to LMCs in joint-draining

cLV secondary to a Ca^{++} flux through gap junctions from attached telocytes. The Ca^{++} flux

generated from telocyte responses to osmotic stress (Figure 8b) and Ehfd1 may play a direct role

in calcium signaling via inhibition of the mitochondrial calcium uniporter (*70*), followed by

shunting of intracellular Ca^{++} through the telocyte network into mast cells.

Our finding that *Myoc* is a peri-PLV telocyte marker gene is also not surprising given that

the extensive ECM (Fig. 4d), and primary functions of this protein are cross-linking ECM

proteins and regulating MMPs (*71*). It is highly probable that *Myoc* is not a telocyte-specific

gene in the joint (Fig. 5b) since the fibroblasts in these musculoskeletal tissues play a major role

in ECM genesis and remodeling. Therefore, since *Efhd1*-CreER[T2] mice have more selective gene

targeting, we conclude that they are superior model for telocyte research. However, mutations in

the *Myoc* are known to be the greatest genetic risk factor for glaucoma. Thus, the *Myoc*-CreER[T2]

mice may be a useful model to better understand this disease.

Here we also report the first evidence of tissue-specific telocytes, which includes our

failure to detect *Efhd1* and *Myoc* expression in ICC of the gut (Supplemental Fig. 5a). This

observation also distinguishes our *Efhd1* and *Myoc* telocyte-targeted CreER[T2] models from the

established *FoxL1*-Cre targeted models (*43, 72*), although all of these mice have transgene

targeting in subepithelial telocytes, suggesting overlaps in the tissue specificity of distinct

telocytes. Additionally, our scRNAseq data demonstrates that synovial and peri-PLV telocytes

do not express *Ano1* (Fig. 5c), which is considered to be the canonical Ca^{++} activated chloride

channel in ICC required for pacemaker activity (*66, 73*). Thus, Ca^{++} signaling in telocytes of the

synovial lymphatic system must involve different pathways, which might include mitochondria,

considering that EFHD1 regulates mitoflash activation that mediates intercellular signaling (*74*)

and physically binds to the mitochondrial calcium uniporter to inhibit the Hippo/YAP pathway

and cellular invasives (*70*). Taken together with our findings that *Efhd1* expression in telocytes is

lost during telocyte to fibroblast/myofibroblast differentiation (Fig. 9), EFHD1 may be a target to

stabilize telocytes in arthritis and other emerging fields that propose telocyte transplantation (*75*).

In terms of the well-known fibrosis that is associated with RA and lymphedema, our *in*

vitro findings demonstrating that relatively non-motile *Efhd1*$^{+}$ telocytes differentiate into

invasive fibroblasts in response to TGF-β and stiff ECM (Figs. 8 & 9) may provide etiologic

insights. Thus, prospective lineage tracing studies with tamoxifen-treated *Efhd1*-CreER^{T2} x Ai9

and *Myoc*-CreER^{T2} x Ai9 mice are warranted to see if this telocyte to myofibroblast

differentiation also occurs in vivo, especially in disease models.

We present the first report of telocytes within the synovial lymphatic system, coupled

with several preliminary findings that must be addressed in future studies. Notable among these

limitations are potential off-target effects of our transgenic mouse models when performing

systemic tamoxifen delivery experiments. As our models may target cells in tissues that were not

studied exhaustively and we confirmed targeting to non-telocytes known to express Efhd1 (e.g.

renal tubule cells (*70*) Supplemental Fig. 9), conclusions about telocyte-specific effects need to

be tempered. We also report some preliminary results that need to be follow up in future studies.

The first is our variable findings of increased osteoclasts in femurs of mice with DTA-deleted

target cells (Supplemental Fig. 7), which warrants prospective studies to determine if these

preliminary observations are significant. We also conclude that our highly variable NIR-ICG

results to assess lymphatic drainage of PBS vs. zymosan injected knees were confounded by

effects of multiple intraarticular injections (data not shown), and that faithful measurement of

lymphatic drainage from the knee will likely require technical innovations (e.g. assessing large

molecule translocation from the joint to iliac lymph nodes).

In summary, we find that telocytes embedded in the ECM around PLVs are lost in TNF-

tg mice with inflammatory arthritis, and telocyte networks adjacent to PLVs are integrated into

mast cells. Furthermore, telocyte deletion decreases lymphatic function and exacerbates ZIA.

Based on these discoveries, we propose a unifying model to explain the functions of telocytes

within the synovial lymphatic system (Fig. 10). This model posits three testable hypotheses that

are the focus of our future directions. The first is the existence of connected telocyte networks

that parallel lymphatic vessels from the synovial lymphatic capillaries to mast cell elaborated

cLV, which is supported by our WMIFM images (Fig. 2h) and can be further tested by high

resolution light-sheet microscopy to confirm the tdT$^+$ cells in the array are physically associated.

The second is that telocytes sense osmotic pressure in joints (e.g. ankle) and signal through the

telocyte network to the peri-PLV mast cells that degranulate and initiate LMC contractions in

response. Experiments to test this will require generation of double transgenic mice in which the

Efhd1-CreER^{T2} are crossed to mice with telocyte or mast cell specific Ca^{++} reporter probes for

intravital microscopy studies following ankle stimulations. Finally, we propose that RA-FLS are

derived from telocytes that differentiate in response to chronic inflammation and adherence to

rigid ECM in inflamed joints. This could be tested via lineage tracing studies with tamoxifen-

treated *Efhd1*-CreER^{T2} x Ai9 mice, and genetic EFHD1 gain and loss of function studies, which

could provide the field with a novel molecular target to treat pauci-immune RA and lymphedema

that is refractory to all known treatments.

Methods

Mouse models. All animal research was conducted with approval by the University of Rochester

Institutional Animal Care and Use Committee. Three transgenic mouse strains were generated

with P2A-CreER^{T2} cassettes, which were knocked into C57BL/6NTac mice via CRISPR/Cas-

mediated genome engineering (Taconic Biosciences, Germantown, NY). Cas9 and gRNA were

co-injected into fertilized eggs with the targeting vector, and the resulting pups were genotyped

by PCR followed by Sanger sequencing analysis to confirm the precise insertion loci (for primers

see Table 1). For the Efhd1-CreER^{T2} model, the TAG stop codon in the mouse *Efhd1* gene

(NCBI Reference Sequence: NM_028889.3) was replaced by a "P2A-CreER^{T2}" cassette

(Transcript: 201-ENSMUST00000027472). A synonymous mutation p. R232= (CGG to AGA)

was also introduced to prevent the binding and re-cutting of the sequence by gRNA after

homology-directed repair. For the Myoc-CreER^{T2} model, the TGA stop codon in the mouse

Myoc gene (NCBI Reference Sequence: NM_010865.3) was replaced by a "P2A-CreER^{T2}"

cassette (Transcript: 201-ENSMUST00000028020). For the Pla1a-CreER^{T2} model, the TGA stop

codon in the mouse *Pla1a* gene (NCBI Reference Sequence: NM_134102.4) was replaced by a

"P2A-CreER^{T2}" cassette (Transcript: 201-ENSMUST00000002926).

B6.Cg-*Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze}$*/J (Ai9)(*33*), B6.129P2-

Gt(ROSA)26Sor$^{tm1(DTA)Lky}$/J (DTA)(*34*) and B6.Cg-*Kit^{W-sh}*/HNihrJaeBsmJ (*Kit$^{W-sh/W-sh}$* or cKit$^{-/-}$)

mice were purchased from The Jackson Laboratory (Bar Harbor, ME, Ai9 Strain #:007909, DTA

Strain #:009669, Kit Strain #: 030764) and maintained at the University of Rochester vivarium.

For crossing between the strains, CreER^{T2} conditional inducible mice were used as male

breeders, and Ai9 or DTA mice as female breeders. For all *in vivo* longitudinal outcome

measures, mice were anesthetized with 1–3% isoflurane. All mice were euthanized with a lethal

dose of ketamine/xylazine cocktail (intraperitoneal) followed by cervical dislocation.

Whole mount immunofluorescent microscopy (WMIFM). PLVs were harvested and fixed as

previous described(*76*). Briefly, fixed PLVs were blocked with 5% normal goat serum (NGS;

ThermoFisher Scientific Cat# 50062Z)/1 × TBS/0.3% Triton X-100 for 1 hr at RT, and then

incubated with the primary antibodies diluted in 5% NGS/1 × TBS / 0.3% Triton X-100

overnight at 4 °C. For αSMA labeling, primary antibodies include mouse anti-αSMA antibody

(AlexaFluor 488 conjugate; ThermoFisher Scientific Cat# 53-9760-82, diluted at 1:100). To

validate the mast cell-telocyte interaction, the following primary antibodies targeting mast cells

were used in this study: Rabbit anti-human Mast Cell Tryptase IgG (mouse reactivity; Bioss

Antibodies Cat# BSM-52533R; 1:100 dilution). To prevent non-specific labeling of Fc receptors

(FcR), exclusively secondary antibodies below featuring F(ab) fragments were employed: Anti-

rabbit IgG (H+L), F(ab')2 Fragment (Alexa Fluor® 647 Conjugate; Cell Signaling Technology

Cat#4414; 1:500 dilution). After 3 × 10 min washes in 1 × TBS/0.1% Triton X-100 following

antibody incubation, the PLVs were mounted on a microscope slide with one drop of both

ProLong Gold Antifade Mountant (ThermoFisher Scientific Cat# P36930) and NucBlue Fixed

ReadyProbes Reagent (DAPI formulation; ThermoFisher Scientific Cat# R37606). The PLVs

were then imaged using a VS120 Slide Scanner for αSMA coverage analysis and Nikon A1R HD

confocal microscopy as previously described (*31, 76*).

To validate the mast cell-telocyte interaction, the following primary antibodies targeting

mast cell were used in this study: Rabbit anti-human Mast Cell Tryptase IgG (mouse reactivity;

Bioss Antibodies Cat# BSM-52533R; 1:100 dilution). To prevent non-specific labeling of Fc

receptors (FcR), exclusively secondary antibodies below featuring F(ab) fragments were

employed: Anti-rabbit IgG (H+L), F(ab')2 Fragment (Alexa Fluor® 647 Conjugate; Cell

Signaling Technology Cat#4414; 1:500 dilution). After 3×10 min washes in $1 \times$ TBS/0.1%

Triton X-100 following antibody incubation, the PLVs were mounted on a microscope slide with

one drop of both ProLong Gold Antifade Mountant (ThermoFisher Scientific Cat# P36930) and

NucBlue Fixed ReadyProbes Reagent (DAPI formulation; ThermoFisher Scientific Cat#

R37606). The PLVs were then imaged using a VS120 Slide Scanner for αSMA coverage

analysis and Nikon A1R HD confocal microscopy as previously described(*26*).

To confirm telocyte identity and characterize marker expression, whole-mounted

lymphatic vessels, OCT-frozen cryosections, or paraffin-embedded tissue sections were

subjected to immunofluorescence staining. The following primary antibodies were used: anti-

CD34 antibody (Abcam Cat# ab81289; 1:50 dilution), CD31 Monoclonal Antibody (Thermo

Fisher Cat# MA1-40074; 1:50), anti-COMP (Abcam Cat# ab231977; 1:125), CD117 (c-Kit)

Monoclonal Antibody (Thermo Fisher Cat# 14-1172-81; 1:100), and anti-Myocilin (Abcam Cat#

ab41552; 1:50). To prevent non-specific Fc receptor binding, only secondary antibodies with

F(ab')2 fragments were used: Alexa Fluor® 647–conjugated anti-rabbit IgG (Cell Signaling

Technology Cat# 4414; 1:500), Alexa Fluor® 555–conjugated anti-rabbit IgG (Cell Signaling

Technology Cat# 4413; 1:1000), FITC-conjugated goat anti-rat IgG (Jackson ImmunoResearch

Cat# 112-096-003; 1:200), Alexa Fluor® 647–conjugated donkey anti-rat IgG (Jackson

ImmunoResearch Cat# 712-606-153; 1:500), and FITC-conjugated goat anti-rabbit IgG (Jackson

ImmunoResearch Cat# 111-096-047; 1:100). Washing, mounting, and imaging were performed

as described above.

Cryosection preparation and Immunohistochemistry (IHC). Mice were anesthetized with

ketamine (100 mg/kg) and xylazine (10 mg/kg) prior to cardiac perfusion with PBS followed by

4% paraformaldehyde (PFA). Tissues of interest were harvested and fixed in 10% neutral

buffered formalin (NBF) for 3 days at room temperature. After fixation, tissues were washed

three times in PBS and equilibrated in 40% sucrose at 4 °C overnight. Tissues were then

infiltrated with optimal control temperature (OCT) embedding medium for 30 minutes at room

temperature, transferred to fresh OCT, and snap-frozen for cryosectioning. Frozen blocks were

sectioned at 10 μm thickness and collected onto glass slides. Sections were used directly to

assess tdT$^+$ endogenous expression via fluorescent microscopy or processed for IHC using the

same antibodies described above for WHIFM.

Histochemistry and Histomorphometry. For histochemistry of ankles and knees and

quantification of synovitis and osteoclasts, lower limbs of the mice were processed for paraffin

embedded demineralized hematoxylin and eosin (H & E) as previously described (22). All slides

were scanned with an Olympus VS120 (full slide scans available upon request), and

histomorphometry for synovial area was quantified with Visiopharm (Hoersholm, Denmark) as

previously described (77).

Electron microscopy. Transmission electron microscopy (TEM) was performed as previously

described (20). The afferent lymphatic vessel to the PLN was identified by injecting Evan's blue

in the footpad and was then excised and fixed overnight at 4°C using a combination fixative of

2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M sodium cacodylate buffer for 24 hours.

The specimens were rinsed in 0.1M sodium cacodylate buffer and post-fixed with buffered 1%

osmium tetroxide. The tissue was dehydrated in a graded series of ethanol to 100%, transitioned

into propylene oxide, infiltrated with EPON/Araldite epoxy resin, followed by embedment in fresh

resin and polymerization for 2 days at 70°C. To identify the lymphatic vessel in the specimen, the

epoxy embedded block was cut serially into 1μm sections and stained with Toluidine blue. Then

the specimen block was trimmed of excess surrounding tissue and thin sectioned at 70 nm with a

diamond knife using an ultramicrotome. The thin sections were placed onto 150 mesh carbon

coated nickel slot grids and stained with uranyl acetate and lead citrate. A Hitachi 7650

Transmission Electron Microscope with a Gatan 11-megapixel Erlangshen digital camera was used

to image the grids. Scanning electron microscopy (SEM) was performed, as previously

described.(*26*) Briefly, PLVs were fixed in 2.5% glutaraldehyde/4% paraformaldehyde/0.1M

cacodylate overnight, post-fixed in buffered 1% osmium tetroxide, dehydrated, critically point

dried, mounted onto aluminum stubs and sputter coated with gold prior to imaging using a Zeiss

Auriga FE SEM. Three SEM micrographs per sample group were randomly chosen for descriptive

analysis.

Single-cell RNA sequencing analyses. *In silico* analyses of previously published scRNAseq

datasets of WT and TNF-tg PLV (GEO; accession number GSE190999) (*32*) were performed

using Seurat (v5.0.1) to generate UMAPs of the mesenchymal cells. Individual gene expression

within this UMAP was performed with FeaturePlot in Seurat. Pseudotime predictions analysis of

the scRNAseq data from WT and TNF-tg PLV was performed with Monocle 3 to generate UMAPs

with heatmaps of the hypothesized cellular differentiation and the telocytes:fibroblasts ratio. For

de novo scRNAseq of synoviocytes, knee tissue from tamoxifen-treated *Efhd1*CreER x Ai9 mice

was digested into a single-cell suspension and processed for scRNAseq as previously

described(*32*). RNA-seq libraries were prepared using the NEBNext® Ultra™ II RNA Library

Prep Kit plus tdT and sequenced on an Illumina NovaSeq X Plus platform. Reads were aligned to

the Mus musculus reference genome (GRCm39) using STAR. UMAPs were generated from the

sequence data (GEO; accession number GSE303999) to identify unique cell clusters and

expression of marker genes.

Bulk RNAseq of purified telocyte cultures on stiffness-defined substrates. Telocytes were

isolated from the ankle synovium of tamoxifen-treated *Efhd1*CreER x Ai9 mice, and tdT$^+$ cells

were FACS purified and plated onto collagen type I–coated CytoSoft® 24-well plates (Advanced

BioMatrix, Cat#5183, #5186, #5189) with defined stiffness levels of 0.2, 8, or 64 kPa. After 24

hours of culture, cells were lysed directly in Buffer RLT Plus (Qiagen, Cat#1053393), and total

RNA was extracted using the RNeasy Plus Micro Kit (Qiagen, Cat# 74134) following the

manufacturer's instructions. RNA-seq libraries were prepared using the NEBNext® Ultra™ II

RNA Library Prep Kit and sequenced on an Illumina NovaSeq X Plus platform. Reads were

aligned to the Mus musculus reference genome (GRCm39) using STAR (GEO; accession number

GSE303624). Differential expression analysis was performed using DESeq2, and visualization of

gene expression patterns was conducted using the Seurat package (v5.0.1).

Near infrared-indocyanine (NIR-ICG) imaging and quantification of lymphatic function.

The subcutaneous injection of 10 µl of a 0.1 mg/ml ICG solution in water was administered to

both hind footpads of the mouse. The fluorescence of the footpad was measured in an IVIS Live

Animal Imaging System (Caliper Life Sciences Inc.) one hour after injection (baseline) to

capture the maximum fluorescence signal intensity, and again at six hours to quantify the

remaining fluorescence signals that were not drained away by lymphatic flow. The ratio of the

fluorescence difference and baseline fluorescence was used to define lymphatic clearance

function, which reflects the lymphatic vessel's ability to remove extra fluid from the footpad.

Zymosan-induced arthritis (ZIA) model. Female 8-weeks-old *Efhd1*-CreER$^{+/-}$ x DTA$^{f/-}$ mice

and wild-type littermate controls received intraperitoneal injections of tamoxifen (0.1 mg/g body

weight) for 5 consecutive days to induce *Efhd1*-driven DTA expression and telocyte depletion.

Two weeks after the initiation of tamoxifen treatment, mice were anesthetized and received a

single intra-articular injection of 180 µg zymosan (Sigma-Aldrich, St. Louis, MO, USA) in 6 µL

PBS into the knee joint cavity. Control mice received PBS alone. Knee joint inflammation was

monitored by high-resolution ultrasound imaging (Vevo 3100, VisualSonics) performed before

zymosan injection (baseline) and on days 2, 7, and 14 post-injection. Analysis of knee joint

inflammation was performed using Amira to measure synovial volume, as described before. (*78*)

On days 13–15, mice were euthanized, and knee joints were harvested, fixed in 4%

paraformaldehyde, micro-CT scanned, and processed for histological analysis.

Micro-CT analysis. Micro-CT scanning and 3D reconstruction were performed as previously

described(*79*), using identical imaging settings and parameters. Following reconstruction, femurs

were digitally segmented from adjacent skeletal structures to improve visualization. To assess

articular erosions, 2D projections were generated from the articular surface of the femoral trochlea,

which articulates with the tibia. This approach maximized the visualization of surface erosion in a

flattened plane and minimized distortion-related quantification errors. Erosive lesions were

manually annotated using QuPath (v0.5.0), and the cumulative erosion area was normalized to the

total projected area of the distal femur (ROI). The erosion percentage (erosion area/ROI) was used

as a quantitative measure of joint damage.

In vitro cellular invasion assays with 3D confocal microscopy and image reconstruction.

Matrigel invasion assays were performed using transwell inserts containing cells suspended in

serum-free medium. Cells migrated toward a bottom chamber containing 10% fetal bovine serum

(FBS), with or without 5 ng/mL TGF-β, for 24 hours. Cultures were then stained with Hoechst to

label nuclei and imaged using an inverted Nikon Ti2-E confocal fluorescence microscopy.

Representative 3D reconstructions were generated and analyzed using Imaris (version 10.2 Oxford

Instruments). For quantitative analysis, the centroid of each Hoechst-stained nucleus was used to

define the position of individual cells. Cells were classified as $Efhd1^+$ telocytes if their nuclei were

spatially enclosed within the tdTomato signal, or as $Efhd1^-$ fibroblasts if there was no overlap with

tdTomato fluorescence. The shallowest nucleus, typically corresponding to a non-migratory or

dead cell, was defined as the reference surface level. The vertical distance between each nucleus

and this reference plane was calculated as the invasion depth. Distributions of invasion depth were

quantified across three independent experiments.

***Ex vivo* co-culture and assays for osmotic pressure response and migration.** PLVs with

surrounding adipose tissue were harvested from *Myoc*-CreER x Ai9 or *Efhd1*-CreER x Ai9

reporter mice and cultured in DMEM + 10% FBS (Sigma-Aldrich, St. Louis, MO, USA) to

establish monolayer co-cultures of tdT$^+$ telocytes and unlabeled tdT$^-$ fibroblasts. For

immunostaining, cells were seeded into the Nunc™ Lab-Tek™ II Chamber Slide™ System

(Thermo Fisher Scientific, Cat# 154534PK), fixed with 10% NBF, and the plastic chamber was

removed prior to staining. Subsequent staining procedures followed the same protocol as for

tissue cryosections. For osmotic shock studies, the primary cells were loaded with Hoechst

nuclear stain (Invitrogen, Waltham, MA) and 10 µM Fluo-4 (Invitrogen, Waltham, MA) as

previously described (*80*). After 30 minutes of incubation, cells were supplemented with 10 mM

Ca^{++} and stimulated with varying concentrations of sucrose (6.25-800 µM) to induce

hyperosmotic pressure changes. Real-time fluorescent microscopy was performed to track Ca^{++}

influx responses. The percentage of responsive cells was calculated by normalizing the number

of Fluo-4$^+$ tdT$^+$ Hoechst$^+$ cells to total tdT$^+$ Hoechst$^+$ cells for telocytes, and Fluo-4$^+$ tdT$^-$

Hoechst$^+$ cells to total tdT$^-$ Hoechst$^+$ cells for fibroblasts.

Statistics. All statistical analyses were performed using GraphPad Prism (version 9.1.1,

GraphPad Software, Boston, MA, USA) and R (version 4.3.2). Additional analyses and data

visualization, including quantification of osmotic shock assay, invasion assays and ZIA model

readouts were conducted using the R packages ggpubr (version 0.6.0). Two-group comparisons

were assessed using unpaired two-tailed Student's t-tests, and multiple group comparisons were

performed using one-way or two-way ANOVA as appropriate. A *p*-value less than 0.05 was

considered statistically significant.

Data availability

The data that support the findings of this study are available upon request. Source data are

available for Figs. 3c & g, 4b, 5d & f, 6c-f, 9b-c, 10a. The RNA-sequencing datasets have been

deposited in GEO database under accession code GSE303999 (scRNA-seq of telocytes in

synovial tissue) and GSE303624 (bulk RNA-seq of telocytes cultured under different matrix

stiffness conditions).

Acknowledgements

We would like to thank the faculty and staff at the Genomics Research Center and Histology,

Biochemistry, Electron Microscopy Resource in the Center for Advanced Resource

Technologies, and Molecular Imaging Core at the University of Rochester Medical Center. This

work was supported by funding from the National Institutes of Health: F30 AG069329 (HMK),

T32GM007356 (HMK), R01AR063650, R01AG059775 (LX), R01AR069000 (CTR),

R01AR056702 (EMS), and P30 AR069655 (EMS).

References

1. B. J. Mehrara *et al.*, The emerging importance of lymphatics in health and disease: an NIH
workshop report. *J. Clin. Invest.* **133**, (2023).
2. D. Negrini, Morphological, Mechanical and Hydrodynamic Aspects of Diaphragmatic Lymphatics.
Biology (Basel) **11**, (2022).
3. J. P. Scallan, S. D. Zawieja, J. A. Castorena-Gonzalez, M. J. Davis, Lymphatic pumping: mechanics,
mechanisms and malfunction. *J. Physiol.* **594**, 5749-5768 (2016).
4. A. R. Hargens, B. W. Zweifach, Contractile stimuli in collecting lymph vessels. *Am. J. Physiol.* **233**,
H57-65 (1977).
5. S. D. Zawieja *et al.*, Ano1 mediates pressure-sensitive contraction frequency changes in mouse
lymphatic collecting vessels. *J. Gen. Physiol.* **151**, 532-554 (2019).
6. M. J. Davis, S. D. Zawieja, Pacemaking in the lymphatic system. *J. Physiol.*, (2024).
7. Q. Zhou, R. Wood, E. M. Schwarz, Y. J. Wang, L. Xing, Near-infrared lymphatic imaging
demonstrates the dynamics of lymph flow and lymphangiogenesis during the acute versus
chronic phases of arthritis in mice. *Arthritis Rheum.* **62**, 1881-1889 (2010).
8. J. P. Scallan *et al.*, Independent and interactive effects of preload and afterload on the pump
function of the isolated lymphangion. *Am. J. Physiol. Heart Circ. Physiol.* **303**, H809-824 (2012).
9. S. D. Zawieja *et al.*, Characterization of the cellular components of mouse collecting lymphatic
vessels reveals that lymphatic muscle cells are the innate pacemaker cells regulating lymphatic
contractions. *eLife*, https://doi.org/10.7554/eLife.90679.90672 (2024).
10. M. J. Davis, C. D. Bertram, Control of lymphatic pacemaking and pumping by mechanobiological
signals. *J. Physiol.*, (2025).
11. M. Muthuchamy, A. Gashev, N. Boswell, N. Dawson, D. Zawieja, Molecular and functional
analyses of the contractile apparatus in lymphatic muscle. *FASEB J.* **17**, 920-922 (2003).
12. S. H. Azhar, H. Y. Lim, B. K. Tan, V. Angeli, The Unresolved Pathophysiology of Lymphedema.
Front. Physiol. **11**, 137 (2020).
13. E. M. Bouta *et al.*, Targeting lymphatic function as a novel therapeutic intervention for
rheumatoid arthritis. *Nat Rev Rheumatol* **14**, 94-106 (2018).
14. M. Cao, M. T. Y. Ong, P. S. H. Yung, R. S. Tuan, Y. Jiang, Role of synovial lymphatic function in
osteoarthritis. *Osteoarthritis Cartilage* **30**, 1186-1197 (2022).
15. Q. Zhang *et al.*, VEGF-C, a Lymphatic Growth Factor, Is a RANKL Target Gene in Osteoclasts That
Enhances Osteoclastic Bone Resorption through an Autocrine Mechanism. *J. Biol. Chem.* **283**,
13491-13499 (2008).
16. R. Guo *et al.*, Inhibition of lymphangiogenesis and lymphatic drainage via vascular endothelial
growth factor receptor 3 blockade increases the severity of inflammation in a mouse model of
chronic inflammatory arthritis. *Arthritis Rheum.* **60**, 2666-2676 (2009).
17. Q. Zhou *et al.*, Vascular endothelial growth factor C attenuates joint damage in chronic
inflammatory arthritis by accelerating local lymphatic drainage in mice. *Arthritis Rheum.* **63**,
2318-2328 (2011).
18. S. T. Proulx *et al.*, MRI and quantification of draining lymph node function in inflammatory
arthritis. *Ann. N. Y. Acad. Sci.* **1117**, 106-123 (2007).
19. J. Li *et al.*, Expanded CD23(+)/CD21(hi) B cells in inflamed lymph nodes are associated with the
onset of inflammatory-erosive arthritis in TNF-transgenic mice and are targets of anti-CD20
therapy. *J Immunol* **184**, 6142-6150 (2010).

20.   E. M. Bouta *et al.*, Treatment of Tumor Necrosis Factor-Transgenic Mice With Anti-Tumor
Necrosis Factor Restores Lymphatic Contractions, Repairs Lymphatic Vessels, and May Increase
Monocyte/Macrophage Egress. *Arthritis Rheumatol* **69**, 1187-1193 (2017).
21.   J. Li *et al.*, Efficacy of B cell depletion therapy for murine joint arthritis flare is associated with
increased lymphatic flow. *Arthritis Rheum.* **65**, 130-138 (2012).
22.   R. D. Bell *et al.*, iNOS dependent and independent phases of lymph node expansion in mice with
TNF-induced inflammatory-erosive arthritis. *Arthritis Res Ther* **21**, 240 (2019).
23.   H. M. Kenney *et al.*, Multi-omics analysis identifies IgG2b class-switching with ALCAM-CD6 co-
stimulation in joint-draining lymph nodes during advanced inflammatory-erosive arthritis. *Front*
Immunol **14**, 1237498 (2023).
24.   R. D. Bell *et al.*, Altered Lymphatic Vessel Anatomy and Markedly Diminished Lymph Clearance
in Affected Hands of Patients With Active Rheumatoid Arthritis. *Arthritis Rheumatol* **72**, 1447-
1455 (2020).
25.   J. P. Scallan *et al.*, Ex vivo Demonstration of Functional Deficiencies in Popliteal Lymphatic
Vessels From TNF-Transgenic Mice With Inflammatory Arthritis. *Front. Physiol.* **12**, 745096
(2021).
26.   H. M. Kenney *et al.*, Persistent popliteal lymphatic muscle cell coverage defects despite
amelioration of arthritis and recovery of popliteal lymphatic vessel function in TNF-Tg mice
following anti-TNF therapy. *Scientific Reports* **12**, 12751 (2022).
27.   S. Pal, S. Nath, C. J. Meininger, A. A. Gashev, Emerging Roles of Mast Cells in the Regulation of
Lymphatic Immuno-Physiology. *Front. Immunol.* **11**, 1234 (2020).
28.   P. A. Nigrovic, D. M. Lee, Mast cells in inflammatory arthritis. *Arthritis Res Ther* **7**, 1 (2004).
29.   F. Rivellese *et al.*, Mast cells in early rheumatoid arthritis associate with disease severity and
support B cell autoantibody production. *Ann Rheum Dis* **77**, 1773-1781 (2018).
30.   J. Keffer *et al.*, Transgenic mice expressing human tumour necrosis factor: a predictive genetic
model of arthritis. *EMBO J.* **10**, 4025-4031 (1991).
31.   Y. Peng *et al.*, Distinct mast cell subpopulations within and around lymphatic vessels regulate
lymph flow and progression of inflammatory-erosive arthritis in TNF-transgenic mice. *Front.*
Immunol. **14**, 1275871 (2023).
32.   H. M. Kenney *et al.*, Single-cell transcriptomics of popliteal lymphatic vessels and peripheral
veins reveals altered lymphatic muscle and immune cell populations in the TNF-Tg arthritis
model. *Arthritis Res Ther* **24**, 64 (2022).
33.   L. Madisen *et al.*, A robust and high-throughput Cre reporting and characterization system for
the whole mouse brain. *Nat. Neurosci.* **13**, 133-140 (2010).
34.   D. Voehringer, H. E. Liang, R. M. Locksley, Homeostasis and effector function of lymphopenia-
induced "memory-like" T cells in constitutively T cell-depleted mice. *J. Immunol.* **180**, 4742-4753
(2008).
35.   L. M. Popescu, M. S. Faussone-Pellegrini, TELOCYTES - a case of serendipity: the winding way
from Interstitial Cells of Cajal (ICC), via Interstitial Cajal-Like Cells (ICLC) to TELOCYTES. *J Cell Mol*
Med **14**, 729-740 (2010).
36.   A. Kondo, K. H. Kaestner, Emerging diverse roles of telocytes. *Development* **146**, (2019).
37.   I. Rosa *et al.*, Impairment in the telocyte/CD34(+) stromal cell network in human rheumatoid
arthritis synovium. *J Cell Mol Med* **25**, 2274-2278 (2021).
38.   S. I. Paerregaard *et al.*, The small and large intestine contain related mesenchymal subsets that
derive from embryonic Gli1(+) precursors. *Nat Commun* **14**, 2307 (2023).
39.   S. Onuora, Rheumatoid arthritis: RA synovium harbours distinct fibroblast subsets. *Nat Rev*
Rheumatol **14**, 250 (2018).

40. F. Mizoguchi *et al.*, Functionally distinct disease-associated fibroblast subsets in rheumatoid
arthritis. *Nat Commun* **9**, 789 (2018).
41. P. Lertkiatmongkol, D. Liao, H. Mei, Y. Hu, P. J. Newman, Endothelial functions of
platelet/endothelial cell adhesion molecule-1 (CD31). *Curr. Opin. Hematol.* **23**, 253-259 (2016).
42. A. Saalbach, U. Anderegg, Thy-1: more than a marker for mesenchymal stromal cells. *FASEB J.*
33, 6689-6696 (2019).
43. M. Shoshkes-Carmel *et al.*, Subepithelial telocytes are an important source of Wnts that
supports intestinal crypts. *Nature* **557**, 242-246 (2018).
44. P. J. Gomez-Pinilla *et al.*, Ano1 is a selective marker of interstitial cells of Cajal in the human and
mouse gastrointestinal tract. *Am. J. Physiol. Gastrointest. Liver Physiol.* **296**, G1370-1381 (2009).
45. E. Bernotiene *et al.*, Delayed resolution of acute inflammation during zymosan-induced arthritis
in leptin-deficient mice. *Arthritis Res. Ther.* **6**, R256-263 (2004).
46. B. M. Radu *et al.*, Calcium Signaling in Interstitial Cells: Focus on Telocytes. *Int J Mol Sci* **18**,
(2017).
47. E. M. Gravallese, G. S. Firestein, Rheumatoid Arthritis - Common Origins, Divergent Mechanisms.
N. Engl. J. Med. **388**, 529-542 (2023).
48. S. Alivernini, G. S. Firestein, I. B. McInnes, The pathogenesis of rheumatoid arthritis. *Immunity*
55, 2255-2270 (2022).
49. F. Zhang *et al.*, Defining inflammatory cell states in rheumatoid arthritis joint synovial tissues by
integrating single-cell transcriptomics and mass cytometry. *Nat. Immunol.* **20**, 928-942 (2019).
50. K. Wei *et al.*, Notch signalling drives synovial fibroblast identity and arthritis pathology. *Nature*
582, 259-264 (2020).
51. E. C. Keystone, J. Smolen, P. van Riel, Developing an effective treatment algorithm for
rheumatoid arthritis. *Rheumatology (Oxford)* **51 Suppl 5**, v48-54 (2012).
52. P. Li, E. M. Schwarz, The TNF-alpha transgenic mouse model of inflammatory arthritis. *Springer*
Semin. Immunopathol. **25**, 19-33 (2003).
53. Q. Liang *et al.*, Lymphatic muscle cells contribute to dysfunction of the synovial lymphatic
system in inflammatory arthritis in mice. *Arthritis Res. Ther.* **23**, 58 (2021).
54. E. M. Bouta *et al.*, The role of the lymphatic system in inflammatory-erosive arthritis. *Semin Cell*
Dev Biol **38**, 90-97 (2015).
55. H. M. Kenney *et al.*, The Enigmas of Lymphatic Muscle Cells: Where Do They Come From, How
Are They Maintained, and Can They Regenerate? *Curr Rheumatol Rev*, (2023).
56. J. A. Castorena-Gonzalez, J. P. Scallan, M. J. Davis, Methods for Assessing the Contractile
Function of Mouse Lymphatic Vessels Ex Vivo. *Methods Mol. Biol.* **1846**, 229-248 (2018).
57. D. F. Van Helden, J. Zhao, Lymphatic vasomotion. *Clin. Exp. Pharmacol. Physiol.* **27**, 1014-1018
(2000).
58. G. D. Hirst, S. M. Ward, Interstitial cells: involvement in rhythmicity and neural control of gut
smooth muscle. *J. Physiol.* **550**, 337-346 (2003).
59. Y. Kito, S. M. Ward, K. M. Sanders, Pacemaker potentials generated by interstitial cells of Cajal in
the murine intestine. *Am. J. Physiol. Cell Physiol.* **288**, C710-720 (2005).
60. B. H. Koh *et al.*, Platelet-derived growth factor receptor-alpha cells in mouse urinary bladder: a
new class of interstitial cells. *J. Cell. Mol. Med.* **16**, 691-700 (2012).
61. B. Lavoie, O. B. Balemba, M. T. Nelson, S. M. Ward, G. M. Mawe, Morphological and
physiological evidence for interstitial cell of Cajal-like cells in the guinea pig gallbladder. *J.*
Physiol. **579**, 487-501 (2007).
62. R. E. Dixon *et al.*, Electrical slow waves in the mouse oviduct are dependent upon a calcium
activated chloride conductance encoded by Tmem16a. *Biol. Reprod.* **86**, 1-7 (2012).

63. D. Briggs Boedtkjer *et al.*, Identification of interstitial Cajal-like cells in the human thoracic duct.
Cells Tissues Organs **197**, 145-158 (2013).
64. K. M. Sanders, L. F. Santana, S. A. Baker, Interstitial cells of Cajal - pacemakers of the
gastrointestinal tract. *J. Physiol.*, (2023).
65. R. E. Dixon *et al.*, Chlamydia infection causes loss of pacemaker cells and inhibits oocyte
transport in the mouse oviduct. *Biol. Reprod.* **80**, 665-673 (2009).
66. S. J. Hwang *et al.*, Expression of anoctamin 1/TMEM16A by interstitial cells of Cajal is
fundamental for slow wave activity in gastrointestinal muscles. *J. Physiol.* **587**, 4887-4904
(2009).
67. S. A. Baker *et al.*, Role of TREK-1 potassium channel in bladder overactivity after partial bladder
outlet obstruction in mouse. *J. Urol.* **183**, 793-800 (2010).
68. J. P. Scallan, M. J. Davis, Itching for answers: how histamine relaxes lymphatic vessels.
Microcirculation **21**, 575-577 (2014).
69. K. J. Plaku, P. Y. von der Weid, Mast cell degranulation alters lymphatic contractile activity
through action of histamine. *Microcirculation* **13**, 219-227 (2006).
70. K. Meng *et al.*, EFHD1, a novel mitochondrial regulator of tumor metastasis in clear cell renal cell
carcinoma. *Cancer Sci* **114**, 2029-2040 (2023).
71. H. Wang *et al.*, Physiological function of myocilin and its role in the pathogenesis of glaucoma in
the trabecular meshwork (Review). *Int. J. Mol. Med.* **43**, 671-681 (2019).
72. H. M. Kolev *et al.*, A FoxL1-CreERT-2A-tdTomato Mouse Labels Subepithelial Telocytes. *Cell Mol*
Gastroenterol Hepatol **12**, 1155-1158 e1154 (2021).
73. R. D. Singh *et al.*, Ano1, a Ca2+-activated Cl- channel, coordinates contractility in mouse
intestine by Ca2+ transient coordination between interstitial cells of Cajal. *J. Physiol.* **592**, 4051-
4068 (2014).
74. T. Hou *et al.*, Identification of EFHD1 as a novel Ca(2+) sensor for mitoflash activation. *Cell*
Calcium **59**, 262-270 (2016).
75. M. Klein, M. Csobonyeiova, S. Ziaran, L. Danisovic, I. Varga, Cardiac Telocytes 16 Years on-What
Have We Learned So Far, and How Close Are We to Routine Application of the Knowledge in
Cardiovascular Regenerative Medicine? *Int J Mol Sci* **22**, (2021).
76. H. M. Kenney *et al.*, Lineage tracing reveals evidence of a popliteal lymphatic muscle progenitor
cell that is distinct from skeletal and vascular muscle progenitors. *Sci Rep* **10**, 18088 (2020).
77. R. D. Bell *et al.*, Selective Sexual Dimorphisms in Musculoskeletal and Cardiopulmonary
Pathologic Manifestations and Mortality Incidence in the Tumor Necrosis Factor-Transgenic
Mouse Model of Rheumatoid Arthritis. *Arthritis Rheumatol* **71**, 1512-1523 (2019).
78. E. M. Bouta *et al.*, Power Doppler ultrasound phenotyping of expanding versus collapsed
popliteal lymph nodes in murine inflammatory arthritis. *PLoS One* **8**, e73766 (2013).
79. H. M. Kenney *et al.*, A high-throughput semi-automated bone segmentation workflow for
murine hindpaw micro-CT datasets. *Bone Rep* **16**, 101167 (2022).
80. M. Vrecl *et al.*, Effect of the ostreolysin A/pleurotolysin B pore-forming complex on intracellular
Ca2+ activity in the vascular smooth muscle cell line A10. *Toxicol In Vitro* **29**, 2015-2021 (2015).

Figure Legends

Figure 1. Generation and validation of CreER^{T2} transgenic mice with tamoxifen-inducible

gene expression in PLV adventitial cells. a Schematic illustration of the P2A-CreER^{T2}

targeting vectors that were used to generate the knock-in transgenic mice as described in

Methods. **b** Schematic illustration of tdT reporter studies performed to assess tissue-specific

tamoxifen-inducible gene expression in the CreER^{T2} knock-in transgenic mice in which founder

lines were crossed with B6.Cg-*Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze}$*/J (Ai9) mice to generate *Efhd1*,

Myoc and *Pla1a* CreER$^{+/-}$ x Ai9$^{+/-}$ test mice and single-transgenic Ai9$^{+/-}$ littermate controls. **c** An

illustration of the tamoxifen dosing regimen (daily intraperitoneal injections on postnatal days

10–13) to the 3 lines of male and female mice (n ≥ 5) is presented with the standard time of

sacrifice at postnatal day 21. **d** WMIFM was performed on PLV immunostained with FITC-

conjugated anti-αSMA antibodies from the indicated tamoxifen-treated mice. Representative 4x

images of tdT^{+} cells (red) surrounding αSMA^{+} (green) in Efhd1CreER$^{+/-}$ x Ai9$^{+/-}$ and

MyocCreER$^{+/-}$ x Ai9$^{+/-}$ mice after tamoxifen induction, but not in CreER negative controls.

Figure 2. Efhd1-CreER^{T2} targets two populations of PLV adventitial cells that are adjacent

to mast cells. a WMIFM of PLV from tamoxifen-treated Efhd1-CreER$^{+/-}$x Ai9$^{+/-}$ mice was

performed as described in Figure 1 and a 4x image is show with 10x images of the boxed ROIs

highlighting **b** a tdT^{+} telocyte-like cell with its telopode (arrow) aligned along the surface of the

PLV, and **c** a network of tdT^{+} telocyte-like cells in the adipose tissue adjacent to the vessels. **d**

Evan's blue dye was injected into the footpad of a tamoxifen-treated *Efhd1*-CreER^{T2} x Ai9

mouse to image the PLVs for dissection, and the tissue was immunostained with fluorescent

antibodies against CD34 and aSMA, and counterstained with DAPI. **e-g** Low power and high-

power confocal microscopy images of the ROI (box) confirm immunostaining of the mast cells

elaborating the tdT$^+$ network along the PLV (green). **h** Eight fluorescent low power images of an

immunostained full-length PLV with associated tissues were stitched together to imaging

telocyte networks that extends from capillaries in the ankle synovium to cLV near the knee. The

box indicates the ROI shown in **e-g**.

Figure 3. TEM and WMIFM imaging of pre-PLV mast cells physically integrated into

telocyte arrays. a TEM was performed on the PLVs described in Fig. 2 and a representative

x1000 image is shown highlighting the endothelium (yellow) and smooth muscle (green) layers,

cells with telocyte ultrastructural morphology (red), and a mast cell (magenta). A high power

x5,000 image of the boxed region is shown highlighting the telocyte-like cell (red arrow)

physically integrated into a mast cell (purple arrow), and remnants of a network of telocyte-like

cells (yellow arrow). **b** PLV with tdT$^+$ telocyte-like cells (red) were labelled with fluorescent

antibodies against αSMA (green) and mast cell tryptase (MCT; magenta) for WMIF confocal

microscopy, and a representative 3D rendering is shown. Note several mast cells are tethered to

the tdT$^+$ telocyte-like cell network (**Supplemental Video 1**). The magnified image of the boxed

ROI illustrates the physical integration, as the distance between the mast cell and tdT$^+$ telocyte-

like cell is less than 500nm.

Figure 4. Transcriptome confirmation of PLV-associated telocytes and the loss of PLV

extracellular matrix (ECM) and telocytes in TNF-tg mice. a Published scRNAseq data on

PLV-associated cells from mice(*32*) were reanalyzed to identify the Pdgfrα$^+$/Cd34$^+$ population

known to contain telocytes and adventitial fibroblasts(*9*), and a color-coated UMAP of these

populations associated with WT PLV is presented to demonstrate their proportions. **b** UMAPs of

known marker genes selectively expressed in one cell type or both telocytes and fibroblasts are

shown to further define these populations. **c** Monocle-3 analysis of the scRNAseq data was

performed to generate UMAPs with pseudotime predictions (heat maps), hypothesized telocyte

differentiation into fibroblasts (lines within the UMAPs), and the 52.3% decrease in the

telocytes:fibroblasts ratio in TNF-tg mice relative to their WT littermates ($p>10-15$ via Fisher's

test). **d** Representative x5000 SEM image of a PLV from a WT mouse with intact ECM

structure, associated mast cell (magenta arrow), and an apparent telocyte cell body (red arrow)

embedded in the ECM. Representative x5000 SEM images of PLVs from TNF-tg mice, **e** one

with partial ECM loss and exposed LMCs (green arrows), and **f** one with extensive ECM and

LMC loss. **g** Immunohistochemistry was performed on knees sections from WT, TNF-tg mice

receiving Placebo or Anti-TNF treatment from a prior study(*26*), and representative fluorescent

microscopy images are shown with quantification of the $CD31^-/CD34^+$ cells as a percentage of

synoviocytes ($n=3$; $p<0.001$ t-test).

Figure 5. Similarities between synovial and PLV-associated tdT$^+$ cells with a telocyte

phenotype and their distinctions from fibroblasts. a Experimental workflow showing

tamoxifen treatment of Efhd1-CreER$^{+/-}$x Ai9$^{+/-}$ mice followed by tissue harvest, cell isolation,

and subsequent analyses. **b** Flow cytometric analysis of knee and ankle synovial tissues revealing

distinct tdT^{high} and tdT^{low} populations (red arrows). **c** UMAPs showing expression of telocyte

and fibroblast marker genes in knee synovial cells, matching the expression patterns observed in

Figure 3b. **d** Representative immunofluorescence images of knee and ankle sections from

tamoxifen-treated Efhd1-CreER$^{+/-}$ x Ai9$^{+/-}$ mice stained for Comp (green) and DAPI (blue),

demonstrating distinct spatial localization of telocytes near the meniscus attachment sites while

fibroblasts (Comp$^+$) occupy peripheral regions. Images at 1x magnification with 5x

magnification of boxed region are shown with parallel H&E bright field images.

Figure 6. In vivo depletion of telocytes decreases lymphatic function. **a** Schematic illustration

of the tamoxifen-induced telocyte deletion model generated by crossing Efhd1- CreER^{T2} mice

DTA-floxed mice. The efficiency of in vivo telocyte deletion following tamoxifen treatment was

assessed via **b** WMIFM of PLVs and **c** IHC of knee synovium with quantification of the relative

telocyte percentage as described in Fig. 3, from the double transgenic mice and their DTA$^{f/-}$

littermates. **d** Histomorphometric quantification of relative telocyte percentage (n=3; $p < 0.5$* or

0.001*** vs. Placebo treated TNF-tg via t-test). **e** Lymphatic clearance in tamoxifen-treated

Efhd1- CreER$^{+/-}$ x DTA$^{f/-}$ mice and their DTA$^{f/-}$ littermate controls was assessed via NIR-ICG

imaging, and representative IVIS images of the ICG injected into the footpads at the time of

injection (Baseline) and 6hrs later are shown with the heatmap of the signal intensity and %

lymphatic clearance for each foot. **f** Data on the % lymphatic clearance of the cohort before and

after tamoxifen injection are presented for each mouse with the mean for the group (n$\geq$6; $p<0.05$

via t-test). Similar results were obtained in experiments with Myoc- CreER$^{+/-}$ x DTA$^{f/-}$ mice (data

not shown).

Figure 7. Telocyte depletion increases severity and duration of synovial inflammation from

zymosan induced arthritis (ZIA). **a** Experimental design and workflow in which 8-week-old

female WT and Efhd1-CreER$^{+/-}$ x DTA$^{f/-}$ mice received tamoxifen treatment to introduce telocyte

depletion in DTA double-transgenics. Knee injections of zymosan or PBS were given at 10-

weeks of age followed by knee ultrasound pre-knee injection and on days- 2, 7, and 14 post-knee

injection. Mice were euthanized on day-15 and knee joints were harvested for histology. **b**

Representative day-14 knee ultrasound images of the synovial volume between the femur (f) and

tibia (t) that were quantified longitudinally (**c** AUC values with mean +/- SD; p<0.01, ANOVA)

and prior to sacrifice on day-14 time (**d**; **p $\leq$ 0.01 via ANOVA). **e** Representative H&E-stained

joint histology obtained at 10x with magnified regions of interest (box) highlighting non-

resorbed cortical bone surfaces in PBS WT (white arrows) vs. resorbed cortical bone surface in

all other groups (yellow arrows), and the ZIA pannus tissue (#). **f** Quantification of the synovial

area is reported as the percentage of ROI, as described in Methods. Data are presented with the

mean +/- SD (**p<0.01via ANOVA). Similar results were obtained in experiments with Myoc-

CreER$^{+/-}$ x DTA$^{f/-}$ mice (data not shown). **g** 3D renderings of the distal femur micro-CT scans,

with quantification of the focal erosions (**h**; *p $\leq$ 0.05 via ANOVA). Data are presented for each

femur with mean +/- SD.

Fig. 8 Telocytes exhibit greater sensitivity to osmotic pressure changes and are less invasive

than fibroblasts. a Schematic illustration of ex vivo culture of PLVs from tamoxifen treated

Myoc-CreER$^{+/-}$x Ai9$^{+/-}$ and Efhd1-CreER$^{+/-}$x Ai9$^{+/-}$ mice, which produce ~1:1 cocultures of tdT^{+}

telocytes and tdT^{-} fibroblasts. For osmotic shock study the cells were preloaded with Hoechst,

Ca^{++}, and Fluo-4, prior to osmotic shock with various concentrations of sucrose (data not shown)

and analyzed via real-time fluorescent microscopy. **b** Representative 10x fluorescent images of a

coculture post-sucrose stimulation are shown to illustrate Ca^{++} responsiveness seen as green

fluorescence (center). The ROI (boxes) are magnified to illustrate four cell types: non-responsive

telocytes (red), responsive telocytes (yellow), non-responsive fibroblasts (blue), and responsive

fibroblasts (green) with (top) and without (bottom) the superimposed tdT channel. The response

to 100 µM (424.5 mOsm) sucrose was quantified as the % Fluo-4 positive telocytes or

fibroblasts, and the data are plotted with the mean +/- SD (** $p<0.01$, t-test, each data point

represents the mean value of 4 images from a single chamber from an independent experiment).

c Matrigel invasion assays were performed with transwell insert containing cells in serum-free

media that invaded towards 10% FBS containing media -/+ 5ng/ml TGF-beta in the bottom

chamber for 24hr. The culture was Hoechst-stained prior to fluorescent confocal microscopy, and

representative 3D reconstructed images are shown with top and side views illustrating the red

telocytes on top of the blue nuclei within the fibroblasts. The migration depth of each nucleus in

the Matrigel was calculated and presented in the histograms, which are representative of three

independent experiments.

Figure 9. PLV-associated telocytes differentiate into fibroblasts on stiff ECM. Primary

cultures of the FACS purified tdT$^+$ cells described in Fig. 4 were cultured on collagen-coated

CytoSoft® Imaging 24-well plates with varying stiffness levels: 0.2 kPa (soft), 8 kPa

(intermediate), and 64 kPa (stiff). After 24hrs of culture the cells were harvested for bulk RNA

sequencing and gene expression analyses. **a** Normalized expression of telocyte (*Efhd1* and *Myoc*)

and myofibroblast (*Acta2* and *Tagln*) maker genes are graphed as a function of the ECM

stiffness. **b** The quantified change in the indicated telocyte, myofibroblast, and fibroblast marker

gene expression is presented with changes as indicated by the key. Also presented are the in

significantly expressed *TIMP*, *Collagen* and *MMP* genes. All indicated changes are significant

(p>0.05).

Figure 10. Proposed telocyte networks within the synovial lymphatic system of the lower

limb and their regulation of joint homeostasis and lymphatic function. a Schematic model of

telocyte regulation of PLV stiffness. Telocytes with telopods integrated along PLV monitor

stiffness and remodel the ECM. When the PLV is too flaccid the telocytes produce myocilin to

crosslink the collagen and fibronectin in the ECM (black arrow). When the PLV becomes too

rigid from ECM accumulation, telocytes release matrix metalloproteases (MMPs) and

downregulate tissue inhibitors of MMPs (TIMPs) to degrade the ECM (white arrow). In the

absence of telocytes in TNF-tg mice, chronic inflammation perpetuates MMP breakdown of the

ECM and dysfunction of flaccid PLV (red arrow). **b** Schematic model of telocyte network

regulation of PLV contraction in the mouse lower limb. Telocytes in the synovium monitor

extracellular osmotic pressure in joint fluid and are activated upon sensing an excessive

threshold (left). Upon activation, an intercellular signal is transmitted through the network of

telopode-to-telopode connected telocytes and is ultimately delivered to mast cells adjacent to the

joint-draining cLV. This signal triggers mast cell degranulation and release of vasoconstricting

factors (e.g. histamine) that initiate LMC contractions in the resting PLV. Subsequent diastole

assisted by ECM tensile force generates vacuum suction to fill the PLV with lymph from the

joint (middle), and the intraluminal shear force triggers intrinsic PLV contractions by LMCs,

which then clears the excess interstitial fluid in the joint returning it to normal osmotic pressure

(right).

Fig. 1 Generation and validation of CreERT2 transgenic mice with tamoxifen-inducible gene expression in PLV adventitial cells. a Schematic illustration of the P2A-CreERT2 targeting vectors that were used to generate the knock-in transgenic mice as described in Methods. **b** Schematic illustration of tdT reporter studies performed to assess tissue-specific tamoxifen-inducible gene expression in the CreERT2 knock-in transgenic mice in which founder lines were crossed with B6.Cg-*Gt(ROSA)26Sor^tm9(CAG-tdTomato)Hze*/J (Ai9) mice to generate *Efhd1*, *Myoc* and *Pla1a* CreERT2[+/-] x Ai9[+/-] test mice and single-transgenic Ai9[+/-] littermate controls. **c** An illustration of the tamoxifen dosing regimen (daily intraperitoneal injections on postnatal days 10–13) to the 3 lines of male and female mice (n ≥ 5) is presented with the standard time of sacrifice at postnatal day 21. **d** WMIFM was performed on PLV immunostained with FITC-conjugated anti-αSMA antibodies from the indicated tamoxifen-treated mice. Representative 4x images of tdT[+] cells (red) surrounding αSMA[+] (green) in Efhd1CreER[+/-] x Ai9[+/-] and MyocCreER[+/-] x Ai9[+/-] mice after tamoxifen induction, but not in CreER negative controls.

Fig. 2 Efhd1-CreERT2 targets two populations of PLV adventitial cells that are adjacent to mast cells. a WMIFM of PLV from tamoxifen-treated Efhd1-CreER[+/-]x Ai9[+/-] mice was performed as described in Figure 1 and a 4x image is show with 10x images of the boxed ROIs highlighting **b** a tdT[+] telocyte-like cell with its telopode (arrow) aligned along the surface of the PLV, and **c** a network of tdT[+] telocyte-like cells in the adipose tissue adjacent to the vessels. **d** Evan's blue dye was injected into the footpad of a tamoxifen-treated *Efhd1*-CreERT2 x Ai9 mouse to image the PLVs for dissection, and the tissue was immunostained with fluorescent antibodies against CD34 and αSMA, and counterstained with DAPI. **e-g** Low power and high-power confocal microscopy images of the ROI (box) confirm immunostaining of the mast cells elaborating the tdT[+] network along the PLV (green). **h** Eight fluorescent low power images of an immunostained full-length PLV with associated tissues were stitched together to imaging telocyte networks that extends from capillaries in the ankle synovium to cLV near the knee.

Fig. 3 TEM and WMIFM imaging of pre-PLV mast cells physically integrated into telocyte arrays. a TEM was performed on the PLVs described in Fig. 2 and a representative x1000 image is shown highlighting the endothelium (yellow) and smooth muscle (green) layers, cells with telocyte ultrastructural morphology (red), and a mast cell (magenta). A high power x5,000 image of the boxed region is shown highlighting the telocyte-like cell (red arrow) physically integrated into a mast cell (purple arrow), and remnants of a network of telocyte-like cells (yellow arrow). **b** PLV with tdT+ telocyte-like cells (red) were labelled with fluorescent antibodies against αSMA (green) and mast cell tryptase (MCT; magenta) for WMIF confocal microscopy, and a representative 3D rendering is shown. Note several mast cells are tethered to the tdT+ telocyte-like cell network (**Supplemental Video 1**). The magnified image of the boxed ROI illustrates the physical integration, as the distance between the mast cell and tdT+ telocyte-like cell is less than 500nm.

Fig. 4 Transcriptome confirmation of PLV-associated telocytes and the loss of PLV extracellular matrix (ECM) and telocytes in TNF-tg mice. a Published scRNAseq data on PLV-associated cells from mice[47] were reanalyzed to identify the Pdgfrα+/Cd34+ population known to contain telocytes and adventitial fibroblasts[13], and a color-coated UMAP of these populations associated with WT PLV is presented to demonstrate their proportions. **b** UMAPs of known marker genes selectively expressed in one cell type or both telocytes and fibroblasts are shown to further define these populations. **c** Monocle-3 analysis of the scRNAseq data was performed to generate UMAPs with pseudotime predictions (heat maps), hypothesized telocyte differentiation into fibroblasts (lines within the UMAPs), and the 52.3% decrease in the telocytes:fibroblasts ratio in TNF-tg mice relative to their WT littermates (p>10-15 via Fisher's test). **d** Representative x5000 SEM image of a PLV from a WT mouse with intact ECM structure, associated mast cell (magenta arrow), and an apparent telocyte cell body (red arrow) embedded in the ECM. Representative x5000 SEM images of PLVs from TNF-tg mice, **e** one with partial ECM loss and exposed LMCs (green arrows), and **f** one with extensive ECM and LMC loss. **g** Immunohistochemistry was performed on knees sections from WT, TNF-tg mice receiving Placebo or Anti-TNF treatment from a prior study[26], and representative fluorescent microscopy images are shown with quantification of the CD31-/CD34+ cells as a percentage of synoviocytes (n=3; p<0.001 t-test).

Fig. 5 Similarities between synovial and PLV-associated tdT⁺ cells with a telocyte phenotype and their distinctions from fibroblasts. a Experimental workflow showing tamoxifen treatment of Efhd1-CreER$^{+/-}$x Ai9$^{+/-}$ mice followed by tissue harvest, cell isolation, and subsequent analyses. **b** Flow cytometric analysis of knee and ankle synovial tissues revealing distinct tdT^{high} and tdT^{low} populations (red arrows). **c** UMAPs showing expression of telocyte and fibroblast marker genes in knee synovial cells, matching the expression patterns observed in Figure 3b. **d** Representative immunofluorescence images of knee and ankle sections from tamoxifen-treated Efhd1-CreER$^{+/-}$ x Ai9$^{+/-}$ mice stained for Comp (green) and DAPI (blue), demonstrating distinct spatial localization of telocytes near the meniscus attachment sites while fibroblasts (Comp$^+$) occupy peripheral regions. Images at 1x magnification with 5x magnification of boxed region are shown with parallel H&E bright field images.

Fig. 6 In vivo depletion of telocytes decreases lymphatic function. **a** Schematic illustration of the tamoxifen-induced telocyte deletion model generated by crossing Efhd1-CreERT2 mice DTA-floxed mice. The efficiency of in vivo telocyte deletion following tamoxifen treatment was assessed via **b** WMIFM of PLVs and **c** IHC of knee synovium with quantification of the relative telocyte percentage as described in Fig. 3, from the double transgenic mice and their DTA$^{f/-}$ littermates. **d** Histomorphometric quantification of relative telocyte percentage (n=3; p < 0.5* or 0.001*** vs. Placebo treated TNF-tg via t-test). **e** Lymphatic clearance in tamoxifen-treated Efhd1-CreERT2$^{+/-}$ x DTA$^{f/-}$ mice and their DTA$^{f/-}$ littermate controls was assessed via NIR-ICG imaging, and representative IVIS images of the ICG injected into the footpads at the time of injection (Baseline) and 6hrs later are shown with the heatmap of the signal intensity and % lymphatic clearance for each foot. **f** Data on the % lymphatic clearance of the cohort before and after tamoxifen injection are presented for each mouse with the mean for the group (n≥6; p<0.05 via t-test). Similar results were obtained in experiments with Myoc-CreERT2$^{+/-}$ x DTA$^{f/-}$ mice (data not shown).

a Tamoxifen treatment · Baseline measurement · Zymosan · Endpoint measurement · Endpoint dissection

Ultrasound at day 0 · Ultrasound at day 2, 7, 14 · µCT scanning of knee joint · Paraffin embedding for histology

Efhd1 CreER · loxP DTA loxP

8 weeks · i.p. injection of 0.1mg/g Tamoxifen · TMX work time · Intra-articular injection of 180µg zymosan · Measurement

Day -12 · Day -7 · Day 0 · Day13-15

b WT · Efhd1CreER$^{+/-}$ x DTA$^{f/-}$

PBS · Zymosan

c Treatment ● PBS ● Zymosan Genotype ✕ DTA ● WT

Synovial Volume Change (%) · Day 0 2 7 14

Area under the curve (AUC) · ** **

d Synovial Volume Change (%) · * ** · Day 14

e WT · Efhd1CreER$^{+/-}$ x DTA$^{f/-}$

PBS · Zymosan · 500µm

f Treatment ● PBS ● Zymosan Genotype ✕ DTA ● WT

Synovial Area (% of ROI) · *** **

g WT · Efhd1CreER$^{+/-}$ x DTA$^{f/-}$

PBS · Zymosan

h Erosio/ROI (%) · * *

Treatment ● PBS ● Zymosan Genotype ✕ DTA ● WT

Fig. 7 Telocyte depletion increases severity and duration of synovial inflammation from zymosan induced arthritis (ZIA). **a** Experimental design and workflow in which 8-week-old female WT and Efhd1-CreERT2$^{+/-}$ x DTA$^{f/-}$ mice received tamoxifen treatment to introduce telocyte depletion in DTA double-transgenics. Knee injections of zymosan or PBS were given at 10-weeks of age followed by knee ultrasound pre-knee injection and on days- 2, 7, and 14 post-knee injection. Mice were euthanized on day-15 and knee joints were harvested for histology. **b** Representative day-14 knee ultrasound images of the synovial volume between the femur (f) and tibia (t) that were quantified longitudinally (**c** AUC values with mean +/- SD; p<0.01, ANOVA) and prior to sacrifice on day-14 time (**d**; **p ≤ 0.01 via ANOVA). **e** Representative H&E-stained joint histology obtained at 10x with magnified regions of interest (box) highlighting non-resorbed cortical bone surfaces in PBS WT (white arrows) vs. resorbed cortical bone surface in all other groups (yellow arrows), and the ZIA pannus tissue (#). **f** Quantification of the synovial area is reported as the percentage of ROI, as described in Methods. Data are presented with the mean +/- SD (**p<0.01via ANOVA). Similar results were obtained in experiments with Myoc-CreERT2$^{+/-}$ x DTA$^{f/-}$ mice (data not shown). **g** 3D renderings of the distal femur micro-CT scans, with quantification of the focal erosions(**h**; *p ≤ 0.05 via ANOVA). Data are presented for each femur with mean +/- SD.

Fig. 8 Telocytes exhibit greater sensitivity to osmotic pressure changes and are less invasive than fibroblasts.
a Schematic illustration of ex vivo culture of PLVs from tamoxifen treated Myoc-CreER[+/-]x Ai9[+/-] and Efhd1-
CreER[+/-]x Ai9[+/-] mice, which produce ~1:1 cocultures of tdT[+] telocytes and tdT[-] fibroblasts. For osmotic shock
study the cells were preloaded with Hoechst, Ca[++], and Fluo-4, prior to osmotic shock with various concentrations of
sucrose (data not shown), and analyzed via real-time fluorescent microscopy. **b** Representative 10x fluorescent
images of a coculture post-sucrose stimulation are shown to illustrate Ca[++] responsiveness seen as green fluorescence
(center). The ROI(boxes) are magnified to illustrate four cell types: non-responsive telocytes (red), responsive
telocytes (yellow), non-responsive fibroblasts (blue), and responsive fibroblasts (green) with (top) and without
(bottom) the superimposed tdT channel. The response to 100 μM (424.5 mOsm) sucrose was quantified as the %
Fluo-4 positive telocytes or fibroblasts, and the data are plotted with the mean +/- SD (** p<0.01, t-test, each data
point represents the mean value of 4 images from a single chamber from an independent experiment). **c** Matrigel
invasion assays were performed with transwell insert containing cells in serum-free media that invaded towards 10%
FBS containing media -/+ 5ng/ml TGF-beta in the bottom chamber for 24hr. The culture was Hoechst-stained prior
to fluorescent confocal microscopy, and representative 3D reconstructed images are shown with top and side views
illustrating the red telocytes on top of the blue nuclei within the fibroblasts. The migration depth of each nuclei in the
Matrigel was calculated and presented in the histograms, which are representative of three independent experiments.

Fig. 9 PLV-associated telocytes differentiate into fibroblasts on stiff ECM. Primary cultures of the FACS purified tdT+ cells described in Fig. 4 were cultured on collagen-coated CytoSoft® Imaging 24-well plates with varying stiffness levels: 0.2 kPa (soft), 8 kPa (intermediate), and 64 kPa (stiff). After 24hrs of culture the cells were harvested for bulk RNA sequencing and gene expression analyses. **a** Normalized expression of telocyte (*Efhd1* and *Myoc*) and myofibroblast (*Acta2* and *Tagln*) maker genes are graphed as a function of the ECM stiffness. **b** The quantified change in the indicated telocyte, myofibroblast, and fibroblast marker gene expression is presented with changes as indicated by the key. Also presented are the in significantly expressed *TIMP*, *Collagen* and *MMP* genes. All indicated changes are significant (p>0.05).

Fig. 10 Proposed telocyte networks within the synovial lymphatic system of the lower limb and their regulation of joint homeostasis and lymphatic function. a Schematic model of telocyte regulation of PLV stiffness. Telocytes with telopods integrated along PLV monitor stiffness and remodel the ECM. When the PLV is too flaccid the telocytes produce myocilin to crosslink the collagen and fibronectin in the ECM (black arrow). When the PLV becomes too rigid from ECM accumulation, telocytes release matrix metalloproteases (MMPs) and downregulate tissue inhibitors of MMPs (TIMPs) to degrade the ECM (white arrow). In the absence of telocytes in TNF-tg mice, chronic inflammation perpetuates MMP breakdown of the ECM and dysfunction of flaccid PLV (red arrow). **b** Schematic model of telocyte network regulation of PLV contraction in the mouse lower limb. Telocytes in the synovium monitor extracellular osmotic pressure in joint fluid and are activated upon sensing an excessive threshold (left). Upon activation, an intercellular signal is transmitted through the network of telopod-to-telopod connected telocytes and is ultimately delivered to mast cells adjacent to the joint-draining cLV. This signal triggers mast cell degranulation and release of vasoconstricting factors (e.g. histamine) that initiate LMC contractions in the resting PLV. Subsequent diastole assisted by ECM tensile force generates vacuum suction to fill the PLV with lymph from the joint (middle), and the intraluminal shear force triggers intrinsic PLV contractions by LMCs, which then clears the excess interstitial fluid in the joint returning it to normal osmotic pressure (right).