Title

Coordination of membrane synthesis with cell growth: the *Escherichia coli* phospholipid synthesis

enzyme PlsB responds to membrane abundance in a manner consistent with filamentation-

mediated inhibition

Author List

Jaïrus Beije, Desi Fierlier, Margot Guurink, Amba Stapert, Adja Zoumaro-Djayoon, Gregory
Bokinsky

Affiliations

Department of Bionanoscience, Kavli Institute of Nanoscience, Delft University of Technology, Delft,
The Netherlands.

Materials and correspondence: g.e.bokinsky@tudelft.nl

Abstract

Cell viability demands tight coordination between growth and the synthesis of new membrane. The mechanisms coordinating cell growth with membrane synthesis in any organism are unclear. In *Escherichia coli,* initiation of phospholipid synthesis by the glycerol-3-phosphate acyltransferase PlsB is regulated by cell growth via an unknown allosteric mechanism. Previous studies have established that PlsB assembles into enzymatically inactive, membrane-bound filaments when overexpressed. We propose that PlsB filamentation regulates PlsB activity and coordinates membrane synthesis with growth. Here, we test our hypothesis by observing the localization of fluorescently labelled PlsB using live-cell fluorescence microscopy. During growth, PlsB localizes as discrete foci, consistent with formation of inactive filaments. Reducing cellular membrane content eliminates PlsB foci and delocalizes PlsB into the cytoplasm. Restoring membrane synthesis causes PlsB foci to reform after a delay. These results are consistent with our hypothesis and suggest a model in which PlsB reversibly assembles into inactive filaments in response to phospholipid abundance. This mechanism establishes a negative feedback loop that controls initiation of phospholipid synthesis by PlsB and effectively coordinates membrane synthesis with growth.

Introduction

How do cells grow without breaking their membranes? Cells must tightly coordinate cell envelope

synthesis with growth to maintain viability. This ability is essential for all cells and yet is not

understood in any organism. The membrane synthesis pathway of the model prokaryote

Escherichia coli is perhaps the best-characterized to date (1). *E. coli* is a Gram-negative bacterial

species whose inner membrane is composed of phospholipids while the outer membrane is an

asymmetric bilayer featuring a phospholipid inner leaflet and a lipopolysaccharide outer leaflet. The

synthesis of membrane phospholipids is initiated in the cytoplasm by the peripheral membrane

enzyme PlsB, which transfers fatty acids from acyl thioesters to glycerol-3-phosphate (G3P) to

generate lysophosphatidic acid (LPA) (**Fig. 1A**). LPA is further acylated by PlsC to generate

phosphatidic acid (PA), the universal precursor of glycerophospholipids (2). Subsequent reactions

add headgroups to the G3P backbone, yielding membrane phospholipids phosphatidylglycerol (PG)

and phosphatidylethanolamine (PE). While LPA is also generated via PlsX and PlsY, this pathway is

not essential in *E. coli,* indicating that most LPA is synthesized by PlsB (3). Synthesized

phospholipids partition into the inner membrane and are subsequently transferred to the inner leaflet

of the outer membrane by several redundant protein bridges (4).

As the first enzyme of the phospholipid synthesis pathway, PlsB activity determines the rate

of membrane phospholipid synthesis. Thus, PlsB activity must be synchronized with growth.

Phospholipid synthesis flux during steady-state growth is not controlled by concentrations of PlsB

substrates (G3P and fatty acyl thioesters) or by PlsB concentration (5–8). Furthermore, PlsB

overexpression does not increase phospholipid abundance (9, 10). Thus, PlsB activity must be

regulated by an allosteric mechanism that activates and deactivates PlsB according to an allosteric

signal. Despite extensive biochemical studies, potential allosteric mechanisms controlling PlsB

activity have remained elusive (11–15). Intriguingly, an early study discovered that PlsB assembles

into protein filaments that are readily detectable if PlsB is overexpressed (10). PlsB filaments consist

of a repeating unit (likely PlsB dimers) helically arranged around a phospholipid core (16). While

filamentous PlsB is enzymatically inactive, dissolution of PlsB filaments with a mild detergent

restores activity. Whether filamentation coordinates PlsB activity with cell growth has remained

unexplored.

Recently, filamentation is increasingly appreciated as a mechanism for regulating metabolic

enzymes (17, 18). High-throughput imaging screens and biochemical experiments have found that

many metabolic enzymes reversibly assemble into filaments (19, 20). Often, filamentation and

inhibition are triggered by the accumulation of a product metabolite (21, 22). In these cases,

filamentation enables negative feedback control over enzyme activity in response to product

demand. Maintaining a fraction of the enzyme population in such an inactive form (described as

"excess capacity" or "enzyme storage") ensures that biosynthetic activity can be rapidly increased

when needed (23).

How might filamentation play a role in regulating PlsB? By analogy to other filament-forming

enzymes, PlsB might assemble into inactive filaments due to accumulation of excess membrane

phospholipids. The fraction of cellular PlsB that is sequestered into inactive filaments might vary in

proportion with the density of phospholipids within the inner membrane: a higher surface density of

phospholipids would tend to increase the fraction of PlsB within a filament, decreasing overall PlsB

activity, while decreased phospholipid content would liberate PlsB from filaments and increase the

active fraction (**Fig. 1B**). This mechanism would establish a direct link between membrane

phospholipid abundance and PlsB activity that acts as a classic negative feedback loop (**Fig. 1C**)

and coordinates phospholipid synthesis with growth (**Fig. 1D**). Intriguingly, *in vitro* experiments

suggest that excess phospholipids (generated by membrane compression) are extruded from

membranes as small lipid filaments (24). Such lipid filaments might nucleate the formation of PlsB

filaments, which have been found to contain an inner phospholipid core (25, 26)

Several lines of evidence are consistent with this model. As already stated, PlsB filaments

are enzymatically inactive. Second, *E. coli* maintains a phospholipid surplus: *E. coli* protoplasts can

be osmotically expanded before lysis occurs (27, 28), and *E. coli* continues to grow for a short time

after phospholipid synthesis is inhibited (29). Third, PlsB is also present in excess: the maximum

activity of PlsB far exceeds activity needed to provide *E. coli* with phospholipids (8). Finally, when

phospholipid synthesis inhibition is relieved, the rate of phospholipid synthesis is transiently faster

than the rate observed during steady-state growth, consistent with a brief period of uninhibited

phospholipid synthesis in the absence of a phospholipid surplus (30).

However, direct evidence for the PlsB filamentation hypothesis is lacking. PlsB filaments

have never been observed in wild-type cells. This is likely due to the comparatively low abundance

of PlsB in *E. coli* (~1000 PlsB/cell (31)). As protein filaments typically assemble in a cooperative

manner (*i.e.* the energetic barrier for nucleating a new filament is higher than for extending an

existing filament (32)), each cell might have at most 1-2 filaments. From these assumptions and

estimates of PlsB filament geometry (26), a filament containing half the total PlsB population would

be small (~180 nm) and thus difficult to observe *in situ.* Here, we use fluorescent tagging to

determine PlsB localization in wild-type cells. Our observations are consistent with our PlsB

filamentation hypothesis: specifically, accumulation of membrane phospholipids triggers PlsB

filamentation and inactivation, enabling negative feedback regulation of phospholipid synthesis and

coordinating phospholipid synthesis with growth.

Results

PlsB localizes as punctate foci during growth.

The ability of proteins to assemble into filaments *in vivo* has been identified in several studies by examining libraries of strains expressing fluorescently tagged proteins (33, 34). To enable PlsB localization, we fused a monomeric GFP variant (msGFP2) to the C-terminus of PlsB (35). As protein fusions may perturb localization and protein activity (36), we first investigated whether our fusion influenced PlsB activity. As PlsB is an essential enzyme, any detrimental effects of protein fusion on PlsB activity are expected to be reflected in the growth rate. Furthermore, localization artefacts may also arise when proteins are expressed above native concentrations. To ensure that PlsB-msGFP2 is expressed at native levels and that any localization observed is not a consequence of overexpression, we integrated the *plsB-msGFP2* gene into the chromosome of wild-type *E. coli* (NCM3722) at the *plsB* locus, replacing unlabelled PlsB while retaining transcriptional control by the *plsB* promoter. *E. coli plsB-msGFP2* grew normally in all tested media (**Fig. 2A**). To determine whether our fusion construct perturbs PlsB function or regulation, we quantified both membrane phospholipids and phospholipid intermediates using liquid chromatography coupled with mass spectroscopy (LCMS). The phospholipid composition of *plsB-msGFP2* cultures (determined by phospholipids/OD) and levels of phospholipid synthesis intermediates was identical to the parent NCM3722 strain in multiple conditions (**Fig. 2B**). Next, we quantified PlsB-msGFP2 using LCMS-based targeted proteomics. The abundance of PlsB peptides in *plsB-msGFP2* closely matched those measured in the parent NCM3722 strain, confirming that PlsB-msGFP2 is expressed from the chromosomal *plsB* locus at native levels (**Fig. 2C**). To confirm that msGFP2 labelling does not disrupt mechanisms that prevent overexpressed PlsB from increasing phospholipid content, we compared phospholipid abundance in our *plsB-msGFP2* strain in which we overexpressed PlsB-msGFP2 from a plasmid with NCM3722 overexpressing unmodified PlsB. While both PlsB and PlsB-msGFP2 overexpression slightly decreased total phospholipid abundance, the phospholipid content of the strain overexpressing PlsB-msGFP2 did not substantially differ from the strain overexpressing PlsB (**Fig. 2D**).

To determine how PlsB localizes during steady-state growth, we imaged exponential phase

plsB-msGFP2 cells from a defined rich medium culture (MOPS / 0.2% glucose / 0.1% CAS amino

acids). Cells were transferred to an agar pad supplemented with the same medium and imaged

using live-cell epifluorescence microscopy. Fluorescent foci were found within 95% of the

population, with diffuse fluorescence appearing within the cytoplasm (**Fig. 3A&B**). Most PlsB foci

localized at the cell poles (**Fig. 3C**). PlsB foci were present at comparable levels in all growth

medium tested, indicating that foci are not an artefact of a specific condition. To determine whether

foci localization is an artefact caused by the msGFP2 fluorescent tag, we also imaged PlsB

localization using a C-terminal HaloTag fusion. PlsB-HaloTag also localized as punctate foci at a

similar frequency as PlsB-msGFP2 (**Fig. 3D**). Finally, we exchanged msGFP2 with the HA epitope

and imaged PlsB using immunofluorescence. PlsB-HA was expressed from a plasmid in wild-type

NCM3722 cells that were subsequently fixed and imaged with anti-HA antibodies. Overexpressed

PlsB-HA also localized as punctate foci in wild-type NCM3722 (**Fig. 3E**). Thus, PlsB foci are not an

artefact of specific tags or overexpression.

Fluorescence intensity of PlsB foci increases with PlsB expression.

The observation that fluorescently tagged proteins localize as foci is consistent with

assembly into filaments or other structures such as chemotaxis receptor arrays (37). In many cases,

the degree of oligomerization is dependent upon protein concentration, with increased

oligomerization correlating with increased protein concentrations. In contrast, if the number or size

of foci do not correlate with protein expression, any foci detected may not indicate protein

oligomers. For instance, the primary phospholipid synthesis enzyme in *Bacillus subtilis,* PlsX,

localizes as foci in some conditions, but foci number did not increase in parallel with PlsX

expression (38). To determine whether the PlsB foci represent protein oligomers, we expressed the

PlsB-msGFP fusion from a plasmid-borne inducible promoter in our *plsB-msGFP2* strain. Inducing

expression of PlsB-msGFP2 did not increase the number of fluorescent foci observed but

dramatically increased the foci brightness (**Fig. 3F**), consistent with PlsB foci indicating formation of

an oligomeric complex. Thus, the effects of increased PlsB-msGFP2 expression are also consistent

with assembly of multimeric complexes.

C-terminal tagging may perturb formation of long filament structures.

The foci observed following overexpression of PlsB-msGFP2 did not appear as elongated filaments

as expected from previous studies (**Fig. 3F**). To determine whether the GFP fusion affects the

formation of filaments, we overexpressed unlabeled PlsB in the *plsB-msGFP2* strain and imaged

our cells using confocal microscopy. In contrast to the foci detected within cells overexpressing

PlsB-msGFP2 alone, we observed linear fluorescent structures more consistent with filaments (**Fig.**

3G). This implies that the presence of a C-terminal tag may adversely affect either the assembly of

PlsB into an orderly filamentous array or the parallel packing of multiple PlsB filaments. However,

immunofluorescence imaging of overexpressed PlsB-HA also appear as foci. Thus, modification of

PlsB on the C-terminus may affect the assembly of PlsB filaments.

Inhibition of fatty acid synthesis delocalizes PlsB.

Despite the unclear effect of C-terminal tagging on the formation of orderly PlsB filaments, the

consistent foci-forming behaviour of PlsB is consistent with assembly into some oligomeric

structure. However, the functional relevance of the foci is unclear. Fluorescent tagging often

introduces artefacts that do not reflect the native location of untagged proteins (39, 40). One

approach for testing connections between foci formation and enzyme function is to determine

whether foci formation is responsive to metabolic perturbations linked with the function of the

enzyme (41). Specifically, if the PlsB foci we observe reflect oligomerization in response to

phospholipid membrane abundance, perturbations to membrane abundance or phospholipid

metabolism are expected to affect localization. Conversely, if the PlsB foci instead are the

consequence of a labelling artefact (e.g. fusion-induced aggregation), the PlsB foci are expected to

be unresponsive to perturbations in membrane metabolism.

We tested whether PlsB localization is relevant to its physiological function by inhibiting fatty

acid synthesis. Triclosan is a fatty acid synthesis inhibitor that acts on the 2*E*-enoyl-acyl-ACP

reductase FabI (42). Triclosan was added to cells growing in liquid culture, which were

subsequently transferred to an agar pad also containing growth medium and triclosan for live-cell

imaging. Triclosan greatly reduced the number of PlsB foci and delocalized fluorescence throughout

the cytoplasm (**Fig. 4A**). A similar result was obtained using the fatty acid synthesis inhibitor

cerulenin, which inactivates the β-keto-acyl-ACP synthases FabB and FabF (43). To determine

whether PlsB foci respond to milder perturbations of fatty acid synthesis, we overexpressed the *E.*

coli thioesterase TesA without its periplasmic trafficking signal. When localized to the cytoplasm,

TesA hydrolyses long-chain fatty acid thioesters (including acyl-ACP PlsB substrates) (44) and

reduces the abundance of membrane phospholipids (45) while allowing growth to continue.

Overexpression of TesA in *plsB-msGFP2* dispersed PlsB-msGFP2 and reduced the foci count (**Fig.**

4A). To determine whether PlsB foci dispersal is caused by generic cell growth arrest rather than a

specific consequence of fatty acid synthesis inhibition, we imaged *plsB-msGFP2* after adding

protein translation inhibitors chloramphenicol and mupirocin. Cells treated with either translation

inhibitor ceased to grow but retained PlsB foci (**Fig. 4B**), indicating that PlsB delocalization is not a

generic response to growth arrest but is a consequence of fatty acid synthesis inhibition. Thus, PlsB

consistently delocalizes in response to perturbations in fatty acid and phospholipid synthesis.

The cytoplasmic localization of PlsB observed when phospholipid synthesis is inhibited is

unexpected. While PlsB is a peripheral membrane protein and not integrated into the membrane, it

is generally regarded as associating stably with the membrane. To further investigate PlsB

localization with higher spatial resolution, we imaged triclosan-treated *plsB-msGFP2* using confocal

microscopy. Consistent with observations from our wide-field images, PlsB-msGFP2 in triclosan-

treated cells is also cytoplasmic and not membrane-associated (**Fig. 4C**). We note that the

phosphatidylserine synthesis enzyme PssA is also a peripheral membrane enzyme that reversibly

associates with the membrane in response to electrostatic charge and substrate availability (46).

PlsB foci gradually reform after phospholipid synthesis is restored.

We next tested whether resumption of fatty acid synthesis might cause PlsB foci to reform.

Triclosan-treated cells in liquid culture were pelleted, resuspended in fresh medium, and plated on

agar containing growth medium and imaged over time. Growth resumed immediately upon triclosan

removal, while PlsB remained dispersed at the beginning of imaging. However, after ~30 minutes of

growth, foci reappeared, coincident with the time of first cell division (**Fig. 4D**). Foci reappearance is

highly synchronized across the population. Thus, both formation and dispersion of PlsB are

reversible and responsive to phospholipid metabolism.

To further investigate PlsB foci formation, we restarted phospholipid synthesis in triclosan-

inhibited cells by feeding exogenous fatty acids palmitate and *cis*-vaccenate. Exogenous fatty acids

are converted by the fatty acid-CoA ligase FadD to acyl-CoA, which are substrates for phospholipid

synthesis by PlsB and PlsC. Fatty acid-fed cells resumed growth at a much slower rate;

nevertheless, PlsB foci steadily accumulated in the population, although less synchronized than

when phospholipid synthesis is restarted by triclosan removal (**Fig. 4E**). We speculate that the

slower growth rate and delayed appearance of foci is due to the inability to synthesize

lipopolysaccharides, which require lipid precursors that cannot be supplied or obtained from

exogenous fatty acids. Thus, PlsB consistently localizes as foci when membrane synthesis is

unperturbed during steady-state growth but disperses when phospholipid abundance is reduced.

Discussion

Indirect evidence from previous studies is consistent with our hypothesis that PlsB activity (and thus

phospholipid synthesis) is regulated by reversible filamentation. First, PlsB concentrations in wild-

type cells are far above levels required to provide sufficient phospholipids for steady-state growth

(1, 8, 47); second, overexpressed PlsB forms enzymatically inactive filaments (25, 26); finally, many

metabolic proteins assemble into enzymatically inactive filaments in the presence of their products

(32, 48). Our observations from live-cell imaging experiments support our hypothesis: PlsB localizes

as a combination of foci and distributed fluorescence, consistent with the formation of a non-

diffusive, membrane-associated state such as a filament. Furthermore, PlsB foci intensity varies in

proportion with PlsB expression, consistent with increased PlsB oligomerization, and contrasts with

results obtained from fluorescently labelled PlsX in *B. subtilis* cells (38). However, while fluorescent

foci are consistent with our hypothesis, many other mechanisms (natural and artefactual) also cause

labelled proteins to appear as foci (36). Thus, our strongest evidence is that PlsB localization

responds reversibly to membrane abundance in a manner predicted by our hypothesis. Specifically,

PlsB foci disappear when membrane phospholipids are depleted and reappear when phospholipid

limitation is alleviated.

While our live-cell microscopy data support our hypothesis, some questions remain. First,

the foci that we observe may not necessarily indicate the formation of PlsB filaments. For instance,

the foci may simply reflect increased localization of PlsB monomers or dimers rather than an

oligomeric structure. This may be due to a preference for binding in regions of anionic phospholipids

or for the curved membrane at the cell poles. Issues also arise from our use of protein fusions to

localize PlsB in live cells. C-terminal fusions appear to compromise filament formation. This may be

caused by the C-terminal extensions disrupting contacts or tweaking the alignment of monomers

and preventing helical packing observed in untagged PlsB. Alternatively, C-terminal tagging may

disrupt the arrangement of multiple PlsB filaments into parallel bundles. Unfortunately, solutions

such as appending tags to the N-terminus, or inserting a tag in an unstructured loop (an approach

successfully used to study MreB localization (49)) are not straightforward to apply to PlsB: the N-

terminus lies at the interface with the membrane, and the Alphafold-predicted structure of PlsB does

not reveal any obvious unstructured loops that might allow the insertion of a bulky fluorescent label

(although a smaller label may be tolerated). Interestingly, although formation of orderly filaments

appears to be disrupted, C-terminal tagging does not dramatically affect PlsB function or regulation:

growth rate and phospholipid abundance of *plsB-msGFP2* are unaffected. This may even be

interpreted as an indication that PlsB and phospholipid homeostasis do *not* in fact rely upon PlsB

filamentation. However, multiple studies that have established the role of filamentation in regulation

of other metabolic enzymes found that disabling filamentation caused surprisingly subtle effects

upon fitness: for instance, disrupting glucokinase filamentation lowered fitness only during growth

transitions, not during steady-state growth (41).

To directly test whether filamentation is required for membrane synthesis regulation, PlsB

mutants that are unable to form filaments should be created and tested *in vivo*. Unfortunately, a

single helical turn of a PlsB filament requires more PlsB monomers (25, 26) than current oligomer

structure prediction tools allow (50). Thus, PlsB filament structures must be experimentally

determined at sufficient resolution to identify residues that stabilize filaments. Expression of

filamentation-deficient PlsB mutants would be expected to increase the phospholipid/protein ratio

well above wild-type levels would provide additional direct support for our hypothesis. Furthermore,

experiments with lipid bilayers supported by a flexible surface could determine whether PlsB

filaments are nucleated by extruded lipid filaments, an approach used to study membrane

remodelling by BAR proteins (51).

Methods

Strains and growth conditions.

Escherichia coli K-12 strain NCM3722 (CGSC #12355) was used for all experiments. Plasmids and

primers used are listed in Tables 1 and 2, respectively, while the amino acid sequences of protein

fusions created are in Table 3. *plsB-msGFP2* was cloned into BglBrick plasmid pBbE5k, sub-cloned

into pDHL940 to fuse with a kanamycin resistance gene, and subsequently inserted together with

the resistance cassette into the NCM3722 chromosome by lambda-red recombination (52). The

kanamycin resistance cassette was subsequently removed. The *plsB-HaloTag* strain was

constructed in a similar manner. Cells were cultured on MOPS minimal media (53) with 0.2% w/v

carbon source (glucose, glycerol, or acetate) or with 0.2% glucose and 0.1% CAS amino acids.

Growth was monitored with optical density measurements (Ultrospec 10 Cell Density Meter, GE

Healthcare). For microscopy experiments, 5 mL cultures were incubated at 37°C in while rotating at

400 rpm. For cultures prepared for experiments requiring larger volumes (growth rate

measurements and LCMS analysis), cells were grown in Erlenmeyer flasks containing a magnetic

stir bar. Culture flasks were partially submerged in in a Grant Instruments Sub Aqua Pro water bath

maintained at 37 °C and stirred (1200 rpm) with submerged 2mag MIXdrive 1 Eco and MIXcontrol

20 stir plates. PlsB expression plasmids (PlsB, PlsB-msGFP2, and PlsB-HA) were constructed by

inserting the PlsB or PlsB fusion gene into a pBbE5k backbone (54).

Culture sampling and LCMS analysis.

291   1 mL samples were rapidly removed from exponential-phase cultures and quenched by pipetting

292   immediately into ice-cooled tubes containing 0.25 mL 10% trichloroacetic acid. Quenched cells were

293   pelleted and stored at -80 °C until analysis. LCMS methods used for phospholipid and PlsB

294   quantification are fully described in reference (8).

Live-cell fluorescence microscopy.

The cultures used for microscopy were prepared at outline in the "culture conditions" section. To

ensure the bacteria grow during the imaging they were placed on agar pads. To make these pads,

first a larger batch of agarose was prepared using 1.5 g low gelling point agarose dissolved in 50

mL milli-Q water for a final concentration of 3%. This agar was then autoclaved to ensure sterility.

Afterwards, the full volume was divided into 0.5 mL aliquots which were stored at -20°C. For each

experiment, one of these aliquots was used and remelted at 80 °C in a heating block. Once melted,

0.5 mL of a MOPS medium prepared at 2x concentration is added. The solution is mixed through

vortexing and briefly (less than 5 min) placed back in the heating block to ensure it does not gel in

the tube. Three 15 x 15 mm glass coverslips are place on a flat piece of parafilm and 0.3 mL of agar

is pipetted on each of these. A second coverslip is then gently placed on top of the agar droplet,

spreading it out to a (near) uniform shape. These are covered and left to gel, after they have

solidified they are placed at 37 °C to pre-heat.

Microscopy slides were prepared in two different ways, depending on the microscope used.

For widefield fluorescence microscopy, a prepared and pre-heated agar pad is taken and one of the

coverslips peeled off. A small amount of culture (1 – 2 μL) is then placed onto the agar pad, which is

then placed agar side down onto a larger 22 x 50 mm coverslip. If used for a timelapse, the sides

are sealed to prevent drying of the agar pad during the acquisition. For confocal microscopy there is

a slight variation on the above protocol, instead of placing the agar onto a 22 x 50 mm coverslip,

they are instead placed onto a microscopy dish with a lid and a small amount of wet tissue paper

inside to prevent dessication.

Widefield microscopy was done initially on an Inverted Olympus IX81 equipped with

a Andor Luca R camera (EM-CCD, 104x1002 pixel chip size, 8x8μm pixel size, 14-bit dynamic

range, 65% quantum efficiency at 600nm), CoolLED pE-4000 fluorescence excitation source, 100x

1.45 NA oil objective (0.13 mm WB), GFP filter cube (ET-EGFP, 470/40 excitation filter, T515lp

dichroic filter cut-off, 525/50 emission filter, reference: C229863), phase ring and heating box. The

microscope was heated to 37°C before the start of the experiment, at least 30 minutes in advance

to ensure a stable temperature. At the start of acquisition the 10x or 20x objective was used to

check alignment of the phase ring and make adjustments if needed. Positions were chosen keeping

in mind a sufficient number of cells in the field of view and no or minimal non-cell debris. Images

were taken in two channels, one in phase contrast (50 ms exposure) and one for GFP

(1000 ms exposure, 490 nm, 40% intensity). Phase contrast imaged were taken through the GFP

filter cube to save time switching blocks. Where relevant, Z-stacks were taken using a 0.2 µm

interval between slices. For timelapses, multiple positions are marked and imaged at 2.5

minute intervals, generally for at least 4 hours but up to 8 hours.

Confocal microscopy is done on a Nikon Eclipse Ti microscope with a 488 nm laser

(confocal-CW, 16 mW), a 100x SR Apo TIRF 1.49 NA oil objective (0.12 mm WB), a 482/35

fluorescence emission filter, a EM-CCD Andor iXon X3 DU897 detection device (512x512 pixel chip

size, 16x16 µm pixel size, 14 bit dynamic range, >90% quantum efficiency at 600nm), and Tokai Hit

sample incubator chamber. The sample is incubated at 37°C during imaging. Suitable locations are

found by looking at the sample using the Resonant scanning. Once a position is found a high

resolution image is taken using Galvano scanning.

HaloTag imaging.

338   Cultures of *plsB-HaloTag* grown to OD600 of ~0.6, incubated with 250 nM JF635 dye for 1 hour at

339   37 °C, and washed three times with medium before imaging.

Immunofluorescence.

An NCM3722 pBbE5K-PlsB-HA culture was induced with 100 µM IPTG and grown for

approximately 1 hour until OD600 of 0.3 – 0.4. Following a gentle fixation protocol (55), the cells

were subsequently fixed for 15 minutes with formaldehyde and glutaraldehyde (final concentrations

2.8% and 0.04%, respectively), washed twice with PBS and concentrated 100-fold by centrifugation

and resuspension in PBS. The dense cell suspension was incubated for 45 min with 1% Triton X-

100 and washed three times with 150 µL PBS. Next, the cells are incubated 45 min in 150 µL PBS

containing 100 µg/mL lysozyme and 5 mM EDTA, followed by three washes with 150 µL PBS and

stored at -20°C until ready for labelling. For antibody labelling, cells were incubated for 1 hour at 37

°C with 0.5 mg/mL antibody (anti-HA rabbit recombinant monoclonal antibody labeled with Alexa

Fluor 488, Abcam EPR22919-101). Cells are washed three times with 150 µL PBS containing

0.05% Tween-20, resuspended in 150 µL PBS and analyzed using fluorescence microscopy.

Image analysis.

To augment the qualitative analysis of microscopy images we developed an image analysis pipeline. The primary goal of this pipeline is to count the number of foci within each cell. This requires detection of individual cells and the subsequent detection of the foci within each cell. The first step is to create image stacks of both the phase contrast and fluorescent images separately. The phase contrast images are loaded into Fiji. The MicrobeJ plugin (56) is then used to segment the cells. The first segmentation is automatic, and this is then followed by manual refinement where necessary. Manual refinement is almost exclusively reserved for eliminating false positives or for separating touching cells. Cells touching the edge are discarded and for all remaining cells the coordinates of their contours are exported. Next, both these contours and the fluorescent images stack are loaded into a custom-made Matlab script. Here the contours are used to find each individual cell in the fluorescent images. All other pixels not defined as cell are considered background and the average background value is subtracted across the image. Next, each cell is analyzed individually. First the cytoplasmic background is estimated by taking the median intensity of cell. This value is then multiplied with a set factor (1.7) to create a threshold. All pixels higher than this threshold are located and further filtered on cluster size. Only clusters of more than 5 pixels are considered a foci. From this we can then collect the number of foci per cell, the number of pixels (i.e. size) and the total intensity of the foci. Next, the centroid of the cell is found as well as the centroid of each foci to determine where in the cell they are located. Besides the exact location of each foci, they are also classified as polar or non-polar. For this determination, foci are considered to be located at the poles when located at the outer 10% of the cell length on either end. Finally, the cell length and width are also measured. All this data is then collected and exported for further analysis.

Acknowledgements

We thank Mehmet Berkmen for sharing HaloTag plasmid pDHL940, Carlas Smith and Michal Shemesh for expertise and assistance with microscopy, Jeremie Capoulade for performing pilot experiments, and Tanneke den Blaauwen and Jolanda Verheul for assistance with cell fixation and

379 immunofluorescence. We thank Ariane Briegel and Arjen Jakobi for encouragement and insightful

380 suggestions, and the entire Bokinsky Lab for stimulating discussions and support.

381 Author Contributions

382 JB: designed and performed experiments, built image analysis pipeline, evaluated data. DF:

383 collected samples for LCMS experiments. MG: constructed *plsB-HaloTag* strain and performed

384 HaloTag imaging experiments. AS: performed initial experiments confirming TesA-driven PlsB

385 delocalization. AZ-D: LCMS analysis. GB: supervised project, constructed *plsB-msGFP2* strain,

386 evaluated data. JB and GB wrote the manuscript. All authors approved the final manuscript.

References

1. Dowhan W. 2013. A retrospective: use of Escherichia coli as a vehicle to study phospholipid synthesis and function. Biochim Biophys Acta 1831:471–494.

2. Yao J, Rock CO. 2013. Phosphatidic acid synthesis in bacteria. Biochim Biophys Acta Mol Cell Biol Lipids https://doi.org/10.1016/j.bbalip.2012.08.018.

3. Yoshimura M, Oshima T, Ogasawara N. 2007. Involvement of the YneS/YgiH and PlsX proteins in phospholipid biosynthesis in both Bacillus subtilis and Escherichia coli. BMC Microbiol 7:69.

4. Ruiz N, Davis RM, Kumar S. 2021. YhdP, TamB, and YdbH Are Redundant but Essential for Growth and Lipid Homeostasis of the Gram-Negative Outer Membrane. mBio 12.

5. Goelz SE, Cronan Jr. JE. 1980. The positional distribution of fatty acids in Escherichia coli phospholipids is not regulated by sn-glycerol 3-phosphate levels. J Bacteriol 144:462–464.

6. Davis MS, Solbiati J, Cronan JE. 2000. Overproduction of acetyl-CoA carboxylase activity increases the rate of fatty acid biosynthesis in Escherichia coli. Journal of Biological Chemistry 275:28593–28598.

7. Heath RJ, Jackowski S, Rock CO. 1994. Guanosine Tetraphosphate Inhibition of Fatty-Acid and Phospholipid-Synthesis in Escherichia coli Is Relieved by Overexpression of Glycerol-3-Phosphate Acyltransferase (PlsB). Journal of Biological Chemistry 269:26584–26590.

8. Noga MJ, Büke F, van den Broek NJF, Imholz NCE, Scherer N, Yang F, Bokinsky G. 2020. Posttranslational Control of PlsB Is Sufficient To Coordinate Membrane Synthesis with Growth in Escherichia coli. mBio 11:e02703-19.

9. Snider MD. 1979. Control of membrane lipid synthesis in Escherichia coli during growth and during the stringent response. J Biol Chem 254:7202–7917.

10. Wilkison WO, Walsh JP, Corless JM, Bell RM. 1986. Crystalline arrays of the Escherichia coli sn-glycerol-3-phosphate acyltransferase, an integral membrane protein. Journal of Biological Chemistry 261:9951–9958.

11. Rock CO, Goelz SE, Cronan JE. 1981. Phospholipid synthesis in Escherichia coli. Characteristics of fatty acid transfer from acyl-acyl carrier protein to sn-glycerol 3-phosphate. Journal of Biological Chemistry 256.

12. Green PR, Merrill Jr. AH, Bell RM. 1981. Membrane phospholipid synthesis in Escherichia coli. Purification, reconstitution, and characterization of sn-glycerol-3-phosphate acyltransferase. Journal of Biological Chemistry 256:11151–11159.

13. Green PR, Bell RM. 1984. Asymmetric reconstitution of homogenous Escherichia coli sn-glycerol-3-phosphate acyltransferase into phospholipid vesicles. Journal of Biological Chemistry 259:14688–14694.

440   14.   Scheideler MA, Bell RM. 1989. Phospholipid dependence of homogeneous,
441   reconstituted sn-glycerol-3-phosphate acetyltransferase of Escherichia coli. Journal of
442   Biological Chemistry.
443

444   15.   Scheideler MA, Bell RM. 1991. Characterization of active and latent forms of the
445   membrane-associated sn-glycerol-3-phosphate acyltransferase of Escherichia coli. Journal of
446   Biological Chemistry.
447

448   16.   Wilkison WO, Bell RM, Taylor KA, Costello MJ. 1992. Structural characterization of
449   ordered arrays of sn-glycerol-3-phosphate acyltransferase from Escherichia coli. J Bacteriol
450   174:6608–6616.
451

452   17.   Simonet JC, Burrell AL, Kollman JM, Peterson JR. 2020. Freedom of assembly:
453   Metabolic enzymes come together. Mol Biol Cell https://doi.org/10.1091/MBC.E18-10-0675.
454

455   18.   Park CK, Horton NC. 2019. Structures, functions, and mechanisms of filament
456   forming enzymes: a renaissance of enzyme filamentation. Biophys Rev
457   https://doi.org/10.1007/s12551-019-00602-6.
458

459   19.   Narayanaswamy R, Levy M, Tschansky M, Stovall GM, O'Connell JD, Mirrielees J,
460   Ellington AD, Marcotte EM. 2009. Widespread reorganization of metabolic enzymes into
461   reversible assemblies upon nutrient starvation. Proc Natl Acad Sci U S A 106.
462

463   20.   Noree C, Begovich K, Samilo D, Broyer R, Monfort E, Wilhelm JE. 2019. A
464   quantitative screen for metabolic enzyme structures reveals patterns of assembly across the
465   yeast metabolic network. Mol Biol Cell 30.
466

467   21.   Barry RM, Bitbol AF, Lorestani A, Charles EJ, Habrian CH, Hansen JM, Li HJ,
468   Baldwin EP, Wingreen NS, Kollman JM, Gitai Z. 2014. Large-scale filament formation inhibits
469   the activity of CTP synthetase. Elife 3:1–19.
470

471   22.   Stoddard PR, Lynch EM, Farrell DP, dosey AM, diMaio F, Williams TA, Kollman JM,
472   Murray AW, Garner EC. 2020. Polymerization in the actin ATPase clan regulates hexokinase
473   activity in yeast. Science (1979) 367:1039–1042.
474

475   23.   Davidi D, Milo R. 2017. Lessons on enzyme kinetics from quantitative proteomics.
476   Curr Opin Biotechnol https://doi.org/10.1016/j.copbio.2017.02.007.
477

478   24.   Staykova M, Holmes DP, Read C, Stone HA. 2011. Mechanics of surface area
479   regulation in cells examined with confined lipid membranes. Proc Natl Acad Sci U S A
480   108:9084–9088.
481

482   25.   Wilkison WO, Walsh JP, Corless JM, Bell RM. 1986. Crystalline arrays of the
483   Escherichia coli sn-glycerol-3-phosphate acyltransferase, an integral membrane protein.
484   Journal of Biological Chemistry 261:9951–9958.
485

486   26.   Wilkison WO, Bell RM, Taylor KA, Costello MJ. 1992. Structural characterization of
487   ordered arrays of sn-glycerol-3-phosphate acyltransferase from Escherichia coli. J Bacteriol
488   174:6608–6616.
489

490   27.   Sun Y, Sun TL, Huang HW. 2014. Physical properties of Escherichia coli spheroplast
491   membranes. Biophys J 107:2082–2090.
492

28.   Buda R, Liu Y, Yang J, Hegde S, Stevenson K, Bai F, Pilizota T. 2016. Dynamics of Escherichia coli's passive response to a sudden decrease in external osmolarity. Proc Natl Acad Sci U S A 113:E5838–E5846.

29.   Bell RM. 1974. Mutants of Escherichia coli defective in membrane phospholipid synthesis: macromolecular synthesis in an sn glycerol 3 phosphate acyltransferase K(m) mutant. J Bacteriol 117.

30.   McIntyre TM, Chamberlain BK, Webster RE, Bell RM. 1977. Mutants of Escherichia coli defective in membrane phospholipid synthesis. Effects of cessation and reinitiation of phospholipid synthesis on macromolecular synthesis and phospholipid turnover. Journal of Biological Chemistry 252:4487–4493.

31.   Li GW, Burkhardt D, Gross C, Weissman JS. 2014. Quantifying absolute protein synthesis rates reveals principles underlying allocation of cellular resources. Cell 157:624–635.

32.   Barry RM, Bitbol AF, Lorestani A, Charles EJ, Habrian CH, Hansen JM, Li HJ, Baldwin EP, Wingreen NS, Kollman JM, Gitai Z. 2014. Large-scale filament formation inhibits the activity of CTP synthetase. Elife 3:1–19.

33.   Narayanaswamy R, Levy M, Tsechansky M, Stovall GM, O'Connell JD, Mirrielees J, Ellington AD, Marcotte EM. 2009. Widespread reorganization of metabolic enzymes into reversible assemblies upon nutrient starvation. Proc Natl Acad Sci U S A 106.

34.   Noree C, Begovich K, Samilo D, Broyer R, Monfort E, Wilhelm JE. 2019. A quantitative screen for metabolic enzyme structures reveals patterns of assembly across the yeast metabolic network. Mol Biol Cell 30.

35.   Valbuena FM, Fitzgerald I, Strack RL, Andruska N, Smith L, Glick BS. 2020. A photostable monomeric superfolder green fluorescent protein. Traffic 21:534–544.

36.   Margolin W. 2012. The price of tags in protein localization studies. J Bacteriol https://doi.org/10.1128/JB.01640-12.

37.   Thiem S, Sourjik V. 2008. Stochastic assembly of chemoreceptor clusters in Escherichia coli. Mol Microbiol 68.

38.   Sastre DE, Bisson-Filho A, de Mendoza D, Gueiros-Filho FJ. 2016. Revisiting the cell biology of the acyl-ACP: Phosphate transacylase PlsX suggests that the phospholipid synthesis and cell division machineries are not coupled in Bacillus subtilis. Mol Microbiol 100:621–634.

39.   Ghodke H, Caldas VEA, Punter CM, van Oijen AM, Robinson A. 2016. Single-Molecule Specific Mislocalization of Red Fluorescent Proteins in Live Escherichia coli. Biophys J 111.

40.   Swulius MT, Jensen GJ. 2012. The helical mreb cytoskeleton in Escherichia coli MC1000/pLE7 is an artifact of the N-terminal yellow fluorescent protein tag. J Bacteriol 194:6382–6386.

41.   Stoddard PR, Lynch EM, Farrell DP, dosey AM, diMaio F, Williams TA, Kollman JM, Murray AW, Garner EC. 2020. Polymerization in the actin ATPase clan regulates hexokinase activity in yeast. Science (1979) 367:1039–1042.

548 42.   Heath RJ, Rubin JR, Holland DR, Zhang E, Snow ME, Rock CO. 1999. Mechanism
549 of triclosan inhibition of bacterial fatty acid synthesis. Journal of Biological Chemistry
550 274:11110–11114.
551
552 43.   D'Agnolo G, Rosenfeld IS, Awaya J, Omura S, Vagelos PR. 1973. Inhibition of fatty
553 acid synthesis by the antibiotic cerulenin. Specific inactivation of beta-ketoacyl-acyl carrier
554 protein synthetase. Biochim Biophys Acta 326:155–156.
555
556 44.   Bonner WM, Bloch K. 1972. Purification and properties of fatty acyl thioesterase I
557 from Escherichia coli. Journal of Biological Chemistry 247.
558
559 45.   Cho H, Cronan JE. 1995. Defective export of a periplasmic enzyme disrupts
560 regulation of fatty acid synthesis. Journal of Biological Chemistry
561 https://doi.org/10.1074/jbc.270.9.4216.
562
563 46.   Louie K, Chen YC, Dowhan W. 1986. Substrate-induced membrane association of
564 phosphatidylserine synthase from Escherichia coli. J Bacteriol
565 https://doi.org/10.1128/jb.165.3.805-812.1986.
566
567 47.   Wilkison WO, Bell RM. 1997. sn-Glycerol-3-phosphate acyltransferase from
568 Escherichia coli. Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism 1348:3–
569 9.
570
571 48.   Park CK, Horton NC. 2019. Structures, functions, and mechanisms of filament
572 forming enzymes: a renaissance of enzyme filamentation. Biophys Rev
573 https://doi.org/10.1007/s12551-019-00602-6.
574
575 49.   Bendezú FO, Hale CA, Bernhardt TG, De Boer PAJ. 2009. RodZ (YfgA) is required
576 for proper assembly of the MreB actin cytoskeleton and cell shape in E. coli. EMBO Journal
577 28.
578
50.   Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O,
Willmore L, Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M,
Reiman D, Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O,
Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB,
Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S, Stecula A,
Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst V, Kohli P,
Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of biomolecular
interactions with AlphaFold 3. Nature 630:493–500.
587
588 51.   Le Roux AL, Tozzi C, Walani N, Quiroga X, Zalvidea D, Trepat X, Staykova M,
589 Arroyo M, Roca-Cusachs P. 2021. Dynamic mechanochemical feedback between curved
590 membranes and BAR protein self-organization. Nat Commun 12.
591
592 52.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
593 Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640–6645.
594
595 53.   Neidhardt FC, Bloch PL, Smith DF. 1974. Culture Medium for Enterobacteria. J
596 Bacteriol 119:736–747.
597
54.   Lee T, Krupa RA, Zhang F, Hajimorad M, Holtz WJ, Prasad N, Lee S, Keasling JD,
Khosla C, Keasling J, Lee S, Chou H, Ham T, Lee T, Keasling J, Atsumi S, Liao J, Raab R,
Tyo K, Stephanopoulos G, Pieper D, Reineke W, Martin V, Pitera D, Withers S, Newman J,
Keasling J, Pitera D, Paddon C, Newman J, Keasling J, Alper H, Fischer C, Nevoigt E,
Stephanopoulos G, Ajikumar P, Xiao W, Tyo K, Wang Y, Simeon F, Leonard E, Mucha O,

Phon T, Pfeifer B, Stephanopoulos G, Dejong J, Liu Y, Bollon A, Long R, Jennewein S,
Williams D, Croteau R, Yan Y, Kohli A, Koffas M, Li M, Elledge S, Shao Z, Zhao H, Zhao H,
Gibson D, Young L, Chuang R, Venter J, Hutchison C, Smith H, Keasling J, Lee S, Kim H,
Park J, Park J, Kim T, Martin C, Nielsen D, Solomon K, Prather K, Canton B, Labno A, Endy
D, Knight T, Anderson J, Dueber J, Leguia M, Wu G, Goler J, Arkin A, Keasling J, Shetty R,
Endy D, Knight T, Tao L, Jackson R, Cheng Q, Anthony J, Anthony L, Nowroozi F, Kwon G,
Newman J, Keasling J, Lee S, Newman J, Keasling J, Lee S, Keasling J, Lee S, Keasling J,
Guzman L, Belin D, Carson M, Beckwith J, Hajimorad M, Gray P, Keasling J, Lee S, Chou H,
Pfleger B, Newman J, Yoshikuni Y, Keasling J, Gorke B, Stulke J, Skulj M, Okrslar V, Jalen
S, Jevsevar S, Slanc P, Strukelj B, Menart V, Carapuca E, Azzoni A, Prazeres D, Monteiro G,
Mergulhao F, Lee C, Ow D, Oh S, Lee C, Kim J, Shin S, Hwang S, Tao L, Jackson R,
Rouviere P, Cheng Q, Lee C, Lee S, Shin S, Hwang S, Friehs K, Antoine R, Locht C, Lutz R,
Bujard H, Manen D, Xia G, Caro L, Grabherr R, Bayer K, Glass R, Kues U, Stahl U, Redding-
Johanson A, Batth T, Chan R, Krupa R, Szmidt H, Adams P, Keasling J, Lee T,
Mukhopadhyay A, Petzold C, Dunlop M, Dossani Z, Szmidt H, Chu H, Lee T, Keasling J,
Hadi M, Mukhopadhyay A, Leguia M, Brophy J, Densmore D, Anderson J, Horton R, Cai Z,
Ho S, Pease L, Pushnova E, Geier M, Zhu Y, Mason G, Provero P, Vaira A, Accotto G. 2011.
BglBrick vectors and datasheets: A synthetic biology platform for gene expression. J Biol Eng
5:12.

55.   Buddelmeijer N, Aarsman M, den Blaauwen T. 2013. Immunolabeling of Proteins in
situ in Escherichia coli K12 Strains. Bio Protoc 3.

56.   Ducret A, Quardokus EM, Brun Y V. 2016. MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. Nat Microbiol 1.

57.   Ke N, Landgraf D, Paulsson J, Berkmen M. 2016. Visualization of periplasmic and
cytoplasmic proteins with a self-labeling protein tag. J Bacteriol 198.

Figures and Tables

Figure 1. Hypothesized mechanism coordinating membrane phospholipid synthesis with cell growth
in *Escherichia coli*. **A.** The role of PlsB in the *E. coli* phospholipid synthesis pathway. **B.** Our
hypothesis predicts that during growth, the PlsB population is partitioned between an active form
(either monomers or dimers) and inactive protein filaments. We propose that the degree of PlsB
filamentation (and thus the level of PlsB activity and the rate of phospholipid synthesis) is
determined by the density of phospholipids within the inner membrane: surplus phospholipids
increase PlsB filamentation, while decreased phospholipid density (i.e. membrane stretching)
decreases PlsB filamentation. This mechanism establishes an effective negative feedback loop that
controls PlsB activity **(C)** and that maintains phospholipid density at a homeostatic level **(D)**.

Figure 2. A msGFP2 fusion with PlsB retains activity. **A.** Growth rates of *E. coli* NCM3722 (wild-type) and NCM3722 *plsB-msGFP2* (PlsB-msGFP2 expressed from the chromosomal *plsB* operon) obtained in three defined media. Two biological replicates of each strain in each condition are shown. **B.** Abundance of phospholipid synthesis intermediates (LPA, PA, and PS) and membrane phospholipids (PG and PE) are unaffected by expression of PlsB-msGFP2. Data obtained from one biological replicate in each condition. **C.** Cellular concentration of PlsB-msGFP2 is equivalent to concentration of PlsB in NCM3722. Triplicate data obtained using LCMS-based peptide quantification from two biological replicates of each strain are shown. **D.** Overexpression of PlsB in NCM3722 and PlsB-msGFP2 in *plsB-msGFP2* generate similar membrane phospholipid profiles. Duplicate data from one biological replicate of each condition is depicted.

Figure 3. PlsB localizes as foci during steady-state growth. **A.** Live-cell fluorescent microscopy
images of *E. coli* NCM3722 *plsB-msGFP2* on agar pads supplemented with defined medium. **B.**
PlsB foci form at similar frequencies across conditions and usually localize at cell poles **(C)**. **D.** PlsB
foci are also observed with different protein fusions, and in fixed cells **(E)**. **F.** Plasmid-based
overexpression of PlsB-msGFP2 does not affect the number of PlsB foci detected per cell but
increases the brightness of the foci. **G.** Overexpression of unlabelled PlsB in *plsB-msGFP2*
background allows resolution of fluorescent structures resembling filaments. For all panels, the
numbers of analysed cells are provided in the figure legends.

Figure 4. PlsB foci respond dynamically to phospholipid abundance. **A.** PlsB-msGFP2 delocalizes into the cytoplasm after membrane synthesis inhibition. Images of *plsB-msGFP2* cells grown in MOPS/0.2% glucose 0.1% CAS amino acids treated with triclosan, cerulenin, or expressing the thioesterase TesA and transferred to agar pads supplemented with identical medium and antibiotics. **B.** PlsB-msGFP2 foci do not substantially respond to translation inhibitors, indicating that delocalization is not a generic response to growth arrest. **C.** A confocal image of triclosan-treated *plsB-msGFP* suggests that PlsB-msGFP2 delocalizes into the cytoplasm and does not associate with the membrane. **D&E.** PlsB-msGFP2 foci reappear when membrane synthesis is restored by either removing triclosan (**D**) or by adding free fatty acids palmitate and *cis*-vaccenate (**E**). For all panels, the numbers of analysed cells are provided in the figure legends.

Table 1. Plasmids in this study.		
Plasmid	**Role**	**Source**
pQE-60NA-msGFP2	msGFP2 sequence	AddGene (plasmid #135301)
pDHL940	HaloTag sequence and antibiotic cassette for recombination	Mehmet Berkmen (57)
pBbE5k	BglBrick vector (lacUV5 promoter, ColE1 origin, kanamycin resistance marker)	AddGene (plasmid #35279)
pBbA2a	BglBrick vector (pTet promoter, p15a origin, ampicillin resistance marker)	AddGene (plasmid #35325)
pBbE5k-PlsB-msGFP2	PlsB-msGFP2 overexpression	This study
pBbA5a-TesA	Cytoplasmic thioesterase expression	This study
pBbE5k-PlsB	PlsB overexpression	This study
pBbE5k-PlsB-HA	HA-tagged PlsB ovreexpression	This study

Plasmid or DNA fragment	primer name	sequence
pBbE5k-PlsB	GB121517-PlsB.f	GCGGATAACAATTTCAGAATTCAAAAGATCTCCAGAGGCTTTACATCGTTTatgTCC
	GB121517-PlsB.r	agatccttactcgagtttggatccttaCCCTTCGCCCTGCGT
	GB121517-plasB.f	ACGCAGGGCGAAGGtaa ggatccaaactcgagtaaggatct
	GB121517-plasB.r	GGAcatAAACGATGTAAAGCCTCTGGAGATCTTTTGAATTCTGAAATTGTTATCCGC
pBbE5k-PlsB-HaloTag	MG_100818-HaloPlsB_Fpls	ctactctggagatttccggttaaggatccaaactcgagtaaggat
	MG_100818-HaloPlsB_Rpls	ccaccgccaccgctCCCTTCGCCCTGCGT
	MG_100818-HaloPlsB_Fphalo	ACGCAGGGCGAAGGagcggtggcggtgg
	MG_100818-HaloPlsB_Rphalo	atccttactcgagtttggatccttaaccggaaatctccagagtag
pBbE5k-PlsB-msGFP2	GB20201012-pPlsB-halo.f	GAGGCAGTGGCGGTTCTtaaggatccaaactcgagtaaggatctcc
	GB20201012-pPlsB-halo.r	GTGAACAGGCTCTCAGTGCTATCatcgttactgccaccgc
	GB20201012-msGFPins.f	gcggtggcagtaacgatGATAGCACTGAGAGCCTGTTCAC
	GB20201012-msGFPins.r	ggagatccttactcgagtttggatccttaAGAACCGCCACTGCCTC
pBbE5k-PlsB-HA	GB101022-plsB.BglII.f	AAAAGATCTCCAGAGGCTTTACATCGTTTATG
	GB221014-plsB.HA-L.XhoI.r	TTACTCGAGTTAAGCGTAATCTGGAACATCGTATGGGTAATCGTTACTGCCACCGCCACC
PlsB-HaloTag chromosomal insertion	MG_091118_HaloTagplsb_F	CGTGCGTTTGACGATTGAGAGTGCGACGCAGGGCGAAGGGagcggtggcggtggcagtaa
	MG_091118_HaloTagplsb_R	CGGATAAGGCGTTTTCGCCGCATCCGGCAATTCTCTCTGAattccggggatccgtcgacc
	MG_101918-SOEplsb.F1	ACCTTCTGGTTGTTGAGTGC
	MG_101918-SOEplsb.R1	CTGCGTCGCACTCTCAAT
	MG_101918-SOEplsb.F2	CTGCAAATTCAATAAATTGCGATG
	MG_101918-SOEplsb.R2	GGTGAGTTAAATGGCTTAGGC
PlsB-msGFP2 chromosomal insertion	GB201110-DHL940.f	GAGGCAGTGGCGGTTCTtaatgtaggctggagctgcttc
	GB201110-msGFP2.r	gaagcagctccagcctacattaAGAACCGCCACTGCCTC

	GB201110-msGFP2.f	ttcgctattacgccagctggatgTCCGGCTGGCCACG
	GB201110-DHL940.r	CGTGGCCAGCCGGAcatccagctggcgtaatagcgaa
	MG_101918-SOEplsb.F2	CTGCAAATTCAATAAATTGCGATG
	GB201127-PlsBint.ubiA.rseq	GGCACTGATGGCGATCATC
	GB201127-PlsBint.50del.r	CATCGCAATTTATTGAATTTGCAGattccgggggatccgtcgacc
	GB201023.plsBint.f	CGCGAAACGCTGCAACG
pBbA5a-TesA	GB20150313-TesA.f	CAGAATTCAAAAGATCTAGGAGGAAAAAAATGGACACGTTATTGATTCTGGGTGATAG
	GB20150313-TesA.r	gagatccttactcgagtttggatccttaTGAGTCATGATTTACTAAAGGCTGC
	GB20150313-TesA-plas.f	GCAGCCTTTAGTAAATCATGACTCAtaaggatccaaactcgagtaaggatctc
	GB20150313-TesA-plas.r	CTATCACCCAGAATCAATAACGTGTCCATTTTTTTCCTCCTAGATCTTTTGAATTCTG

Table 3. Amino acid sequences of PlsB fusions. Linker sequences are underlined while fusion tags are in bold text.

Protein fusion	Amino acid sequence
PlsB-msGFP2	MSGWPRIYYKLLNLPLSILVKSKSIPADPAPELGLDTSRPIMYVLPYNSK ADLLTLRAQCLAHDLPDPLEPLEIDGTLLPRYVFIHGGPRVFTYYTPKE ESIKLFHDYLDLHRSNPNLDVQMVPVSVMFGRAPGREKGEVNPPLR MLNGVQKFFAVLWLGRDSFVRFSPSVSLRRMADEHGTDKTIAQKLAR VARMHFARQRLAAVGPRLPARQDLFNKLLASRAIAKAVEDEARSKKIS HEKAQQNAIALMEEIAANFSYEMIRLTDRILGFTWNRLYQGINVHNAER VRQLAHDGHELVYVPCHRSHMDYLLLSYVLYHQGLVPPHIAAGINLNF WPAGPIFRRLGAFFIRRTFKGNKLYSTVFREYLGELFSRGYSVEYFVE GGRSRTGRLLDPKTGTLSMTIQAMLRGGTRPITLIPIYIGYEHVMEVGT YAKELRGATKEKESLPQMLRGLSKLRNLGQGYVNFGEPMPLMTYLNQ HVPDWRESIDPIEAVRPAWLTPTVNNIAADLMVRINNAGAANAMNLCC TALLASRQRSLTREQLTEQLNCYLDLMRNVPYSTDSTVPSASASELID HALQMNKFEVEKDTIGDIIILPREQAVLMTYYRNNIAHMLVLPSLMAAIV TQHRHISRDVLMEHVNVLYPMLKAELFLRWDRDELPDVIDALANEMQ RQGLITLQDDELHINPAHSRTLQLLAAGARETLQRYAITFWLLSANPSI NRGTLEKESRTVAQRLSVLHGINAPEFFDKAVFSSLVLTLRDEGYISDS GDAEPAETMKVYQLLAELITSDVRLTIESATQGEG<u>SGGGGSND</u>**DSTES LFTGVVPILVELDGDVNGHKFSVRGEGEGDATNGKLTLKFICTTGKLP VPWPTLVTTLTYGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTITFK DDGTYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSH NVYITADKQKNGIKANFKIRHNVEDGSVQLADHYQQNTPIGDGPVLL PDNHYLSTQSKLSKDPNEKRDHMVLLEFVTAAGITGGSGGS**
PlsB-HA	MSGWPRIYYKLLNLPLSILVKSKSIPADPAPELGLDTSRPIMYVLPYNSK ADLLTLRAQCLAHDLPDPLEPLEIDGTLLPRYVFIHGGPRVFTYYTPKE ESIKLFHDYLDLHRSNPNLDVQMVPVSVMFGRAPGREKGEVNPPLR MLNGVQKFFAVLWLGRDSFVRFSPSVSLRRMADEHGTDKTIAQKLAR VARMHFARQRLAAVGPRLPARQDLFNKLLASRAIAKAVEDEARSKKIS HEKAQQNAIALMEEIAANFSYEMIRLTDRILGFTWNRLYQGINVHNAER VRQLAHDGHELVYVPCHRSHMDYLLLSYVLYHQGLVPPHIAAGINLNF WPAGPIFRRLGAFFIRRTFKGNKLYSTVFREYLGELFSRGYSVEYFVE GGRSRTGRLLDPKTGTLSMTIQAMLRGGTRPITLIPIYIGYEHVMEVGT YAKELRGATKEKESLPQMLRGLSKLRNLGQGYVNFGEPMPLMTYLNQ HVPDWRESIDPIEAVRPAWLTPTVNNIAADLMVRINNAGAANAMNLCC TALLASRQRSLTREQLTEQLNCYLDLMRNVPYSTDSTVPSASASELID HALQMNKFEVEKDTIGDIIILPREQAVLMTYYRNNIAHMLVLPSLMAAIV TQHRHISRDVLMEHVNVLYPMLKAELFLRWDRDELPDVIDALANEMQ RQGLITLQDDELHINPAHSRTLQLLAAGARETLQRYAITFWLLSANPSI NRGTLEKESRTVAQRLSVLHGINAPEFFDKAVFSSLVLTLRDEGYISDS GDAEPAETMKVYQLLAELITSDVRLTIESATQGEG<u>SGGGGSND</u>**YPYDV PDYA**
PlsB-HaloTag	MSGWPRIYYKLLNLPLSILVKSKSIPADPAPELGLDTSRPIMYVLPYNSK ADLLTLRAQCLAHDLPDPLEPLEIDGTLLPRYVFIHGGPRVFTYYTPKE ESIKLFHDYLDLHRSNPNLDVQMVPVSVMFGRAPGREKGEVNPPLR MLNGVQKFFAVLWLGRDSFVRFSPSVSLRRMADEHGTDKTIAQKLAR VARMHFARQRLAAVGPRLPARQDLFNKLLASRAIAKAVEDEARSKKIS HEKAQQNAIALMEEIAANFSYEMIRLTDRILGFTWNRLYQGINVHNAER VRQLAHDGHELVYVPCHRSHMDYLLLSYVLYHQGLVPPHIAAGINLNF WPAGPIFRRLGAFFIRRTFKGNKLYSTVFREYLGELFSRGYSVEYFVE GGRSRTGRLLDPKTGTLSMTIQAMLRGGTRPITLIPIYIGYEHVMEVGT YAKELRGATKEKESLPQMLRGLSKLRNLGQGYVNFGEPMPLMTYLNQ HVPDWRESIDPIEAVRPAWLTPTVNNIAADLMVRINNAGAANAMNLCC

TALLASRQRSLTREQLTEQLNCYLDLMRNVPYSTDSTVPSASASELID
HALQMNKFEVEKDTIGDIIILPREQAVLMTYYRNNIAHMLVLPSLMAAIV
TQHRHISRDVLMEHVNVLYPMLKAELFLRWDRDELPDVIDALANEMQ
RQGLITLQDDELHINPAHSRTLQLLAAGARETLQRYAITFWLLSANPSI
NRGTLEKESRTVAQRLSVLHGINAPEFFDKAVFSSLVLTLRDEGYISDS
GDAEPAETMKVYQLLAELITSDVRLTIESATQGEGS<u>GGGGSND</u>**GSEIG**
TGFPFDPHYVEVLGERMHYVDVGPRDGTPVLFLHGNPTSSYVWRNII
PHVAPTHRCIAPDLIGMGKSDKPDLGYFFDDHVRFMDAFIEALGLEE
VVLVIHDWGSALGFHWAKRNPERVKGIAFMEFIRPIPTWDEWPEFAR
ETFQAFRTTDVGRKLIIDQNVFIEGTLPMGVVRPLTEVEMDHYREPFL
NPVDREPLWRFPNELPIAGEPANIVALVEEYMDWLHQSPVPKLLFWG
TPGVLIPPAEAARLAKSLPNCKAVDIGPGLNLLQEDNPDLIGSEIARW
LSTLEISG