Multi-ancestry Transcriptome-Wide Association Study Reveals Shared and Population-Specific Genetic Effects in Alzheimer's Disease

Xinyu Sun[1,3], Makaela Mews[2,3], Nicholas R. Wheeler[1,3], Penelope Benchek[1,3], Tianjie Gu[14,15], Lissette Gomez[14,15], Nicholas Ray[4,5,6], Christiane Reitz[4,5,6], Adam C. Naj[7,8], Jennifer Elizabeth Below[9], Giuseppe Tosto[4,5,6], Mario Cornejo-Olivas[10,11], Goldie S. Byrd[12], Briseida E. Feliciano-Astacio[13], Katrina Celis[14,15], Farid Rajabli[14,15], Brian W. Kunkle[14,15], Margaret A. Pericak-Vance[14,15], Jonathan L. Haines[1,3], Anthony J. Griswold[14,15], and William S. Bush[1,3,□]

[1] Department of Population and Quantitative Health Sciences, School of Medicine, Case Western Reserve University, Cleveland, Ohio, USA.

[2] System Biology & Bioinformatics; Department of Nutrition, School of Medicine, Case Western Reserve University, Cleveland, Ohio, USA.

[3] Cleveland Institute for Computational Biology, Case Western Reserve University, Cleveland, Ohio, USA.

[4] Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Vagelos College of Physicians and Surgeons, Columbia University, New York, New York, USA.

[5] The Gertrude H. Sergievsky Center, Vagelos College of Physicians and Surgeons, Columbia University, New York, New York, USA.

[6] Department of Neurology, Vagelos College of Physicians and Surgeons, Columbia University, and the New York Presbyterian Hospital, New York, New York, USA.

[7] Department of Biostatistics and Epidemiology, University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA.

[8] Penn Neurodegeneration Genomics Center, Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA.

[9] Vanderbilt University Medical Center

[10] Neurogenetics Working Group, Universidad Cientifica del Sur, Lima, Peru.

[11] Neurogenetics Research Center, Instituto Nacional de Ciencias Neurologicas, Lima, Peru.

[12] Maya Angelou Center for Health Equity, Wake Forest University, Winston-Salem, North Carolina, USA.

[13] Universidad Central del Caribe, Bayamón, Puerto Rico.

[14] John P. Hussman Institute for Human Genomics, Miller School of Medicine, University of Miami, Miami, Florida, USA

[15] Dr. John T. Macdonald Foundation Department of Human Genetics, University of Miami Miller School of Medicine, Miami, Florida, USA

[□] Correspondence: William S. Bush <wsb36@case.edu>

Abstract

Alzheimer's disease (AD) risk differs across ancestral populations, yet most genetic studies have focused on Non-Hispanic White (NHW) cohorts. We conducted a multi-population transcriptome-wide association study (TWAS) using whole-blood RNA-seq and genotype data from reported NHW (n=235), African American (AA; n=224), and Hispanic (HISP; n=292) participants in MAGENTA. Using SuShiE for multi-population fine-mapping, we identified credible sets of eQTLs for 8,748 genes and improved fine-mapping precision relative to analyses using fewer populations. eQTL effects were largely shared across populations, with population-specific regulation for a subset of genes. Population-stratified TWAS and sample size–weighted meta-analysis (FUSION + MAFOCUS) prioritized and and fine-mapped nine genes (FDR<0.05, PIP>0.8), including established AD loci (*BIN1*, *PTK2B*, *DMPK*) with consistent effects across populations. Importantly, at *BIN1* we fine-mapped regulatory variants associated with gene expression and AD risk beyond the GWAS index SNP—most notably rs11682128, which is only in modest LD with rs6733839 ($r^2 \approx 0.34$)—demonstrating that multi-population TWAS can implicate additional functional variants not captured by single-SNP GWAS signals. We also discovered a novel association between COG4 expression and AD in NHW, implicating Golgi apparatus function. Using independent SuShiE-derived models from TOPMed MESA (PBMC), several associations replicated directionally across ancestries, with statistical significance most evident in NHW. Our results show that multi-population fine-mapping improves eQTL resolution and TWAS interpretability, reveals regulatory variants beyond GWAS index SNPs, and underscores the need to expand non-European AD cohorts to resolve shared and population-specific mechanisms.

Introduction

Alzheimer's Disease (AD) is the most common form of dementia and affects millions worldwide, with its prevalence expected to increase significantly as global populations age[1]. AD is heritable, though AD risk varies substantially across ancestral populations, with African Americans having approximately

double the risk and Hispanics having about one and one-half times the risk

compared to non-Hispanic Whites[1,2]. Recent and ongoing ascertainment efforts

have increased participation of African American and Admixed Hispanic

individuals in genetic studies, which have led to multiple ancestry-specific

GWAS[3], and a recent multi-ancestry GWAS meta-analysis[4]. These studies

(along with detailed analyses of APOE-□4 effects[5]) have revealed substantial

heterogeneity in effect sizes of established AD risk variants[4], and different

genetic architectures in AD risk.

Growing recognition that AD involves coordinated dysregulation of both

peripheral and central immune systems has shifted the field toward viewing AD

as a systemic disorder. As highlighted by Bettcher et al.[6], mounting clinical and

mechanistic evidence reveals dynamic crosstalk between the peripheral immune

system and the central nervous system, suggesting that immune perturbations in

blood may not only reflect, but also influence, AD pathogenesis. This systems-

level perspective underscores the value of studying whole blood as an

informative, accessible tissue for understanding disease mechanisms across

diverse populations.

Multiple transcriptome-wide association studies (TWAS) of AD have been

conducted to extend GWAS results to functional effects in brain and blood[7–17]. In

principle, TWAS accelerates the advancement of interventions by jointly

modeling associations from multiple (typically non-coding) variants to estimate

the effect of changes in gene expression on a phenotype, which provides more

biologically interpretable relationships than single variant associations alone.

Various multi-variable modeling strategies (such as Elastic net[18] and LASSO[19])

have demonstrated gene expression is predominantly driven by a sparse set of

genetic variants[20–22]. Recent methodological advances in Bayesian

fine☐mapping[23,24] have also led to approaches that use fine-mapped cis-eQTLs

as variables for gene expression prediction[21,25]. Bayesian fine-mapping methods

such as SuSIE quantify uncertainty by defining credible sets of highly correlated

variants, which enables more robust interpretation in the presence of complex

linkage disequilibrium (LD) structures[23]. Notably, all of these approaches require

population-specific eQTL reference panels[26] that have distinct eQTL effects

specific to a particular population and also matches the LD structure of the target

GWAS dataset. Due to the scarcity of non-European eQTL datasets, nearly all

TWAS have been conducted in Non-Hispanic white populations only.

In this work, we perform multi-ancestry fine-mapping of whole-blood eQTL effects

and conduct the first multi-ancestry TWAS of AD within reported African

American (AA), Non-Hispanic White (NHW), and Hispanic (HISP) participants in

the MAGENTA study. We first build population-specific gene expression

prediction models; then perform population-stratified TWAS, integrate these

associations via sample-size–weighted meta-analysis, and fine-map putative

causal genes. Finally, we compare dense (all cis-SNPs) versus sparse (credible

set–based) models. We report novel association signals and further dissect the

blood-based functional genetic architecture of known AD association signals.

Methods

eQTL Reference Panel Study Participants

The *Multi-Ancestry Genomics, Epigenomics, and Transcriptomics of Alzheimer's*

(MAGENTA) study accessed existing sample collections that were part of

multiple previous NIA-funded research projects. Some legacy samples were

drawn from studies that ascertained White or African American participants. All

other samples were drawn from more recent studies including the Puerto Rican

Alzheimer's Disease Initiative, the Cuban American Alzheimer's Disease

Initiative, the Peru Alzheimer's Disease Initiative, and the Research in African-

American Alzheimer's Disease Initiative. All participants and/or their consenting

proxy provided written informed consent as part of the study protocols approved

by the site-specific Institutional Review Boards.

Population Descriptors and Study Group Definitions

Consistent with recent recommendations[27], we define our rationale and definition

for the population descriptors in this study. MAGENTA participants were recruited

across multiple studies that differed in catchment geography (U.S., Peru), terms

used for study recruitment and outreach, and cultural context. As a result,

demographic fields and participant descriptors varied across individuals.

To enable cross-group comparisons while acknowledging differences in genetic

ancestry and admixture, we used a combination of geographic and self-identity

terms as follows:

- "Non-Hispanic White" (NHW): individuals ascertained in North Carolina, Tennessee, and South Florida where participants self-identified as Non-Hispanic White or were categorized as not Hispanic in legacy datasets where recontact is impossible.

- "African American" (AA): individuals ascertained in Northeast Ohio, North Carolina, and South Florida using descriptors intended to reach self-identified Black/African American participants.
- "Caribbean Hispanic": individuals ascertained in the U.S. using descriptors intended to reach individuals with Cuban or Puerto Rican heritage, reflecting similar admixture patterns and environmental exposures.
- "Peruvian Hispanic": individuals ascertained in Lima, Peru and surrounding areas.

For consistency with prior published GWAS studies, we combined the Caribbean Hispanic and Peruvian Hispanic into a single "Hispanic" group to facilitate our TWAS analyses. These labels reflect recruitment context and self-identification; downstream genetic analyses incorporate ancestry-informative covariates to account for population structure.

Genotyping and Quality Control

All samples were genotyped using either the Illumina MEGA v2 or v3 multi-population array and subsequently imputed using TOPMed reference panels v2 (build 38) corresponding with self-identified ancestry and/or ascertainment criteria. We restricted our analysis to variants directly genotyped by the array and variants with an imputation quality control score of R^2 > 0.8. Genotyping data was processed using the ADSP xqtl-protocol release 0.1.1. All analyses were performed stratified by group yielding a total of three analyses. Relatedness was assessed using kinship analysis generated by the KING software. Subsequently, we removed samples to eliminate relatedness (kinship coefficient $\geq$ 0.0625) and maintained samples by prioritizing inclusion of cognitively normal individuals with advanced age, diagnosed AD cases, and control samples with rarer APOE genotypes (ε3/4 or ε4/4). We concluded our genotyping quality control using PLINK2.0 to filter variants with minor allele counts (MAC) of 10, Hardy-Weinberg

equilibrium of $1 * 10^{-8}$, variant-level missingness of 0.1, and sample-level

missingness of 0.1.

Post-QC, 224 AA, 235 NHW, 292 HISP samples from MAGENTA, with both

whole-blood expression and genotype data, were analyzed. The gene expression

was measured in whole blood using RNA-seq. We included the common protein-

coding genes that were available in all three populations. In total, we retained

14,436 genes. Major Histocompatibility Complex (MHC) region from

chromosome 6p (~24–34 Mbp) was excluded from the study, due to the complex

LD structure. In the end, we analyzed 14,211 protein-coding genes.

RNA Processing

Sample preparation and RNA sequencing of whole blood samples have been

previously described[28]. Briefly, RNA was extracted from previously frozen whole

blood samples, quality controlled with RNA integrity (RIN) scores >= 5, prepared

using the NuGEN Universal Plus mRNA-Seq with globin and ribosomal depletion,

and sequenced using 125bp paired end Illumina HiSeq3000. RNA sequencing

FASTQ files were processed using the STAR aligner (v.2.7.10a), quality

controlled using Picard, and quantified using RSEM (v.1.3.3) using the following

reference files:

GRCh38_full_analysis_set_plus_decoy_hla.noALT_noHLA_noDecoy_ERCC.fast

a, Homo_sapiens.GRCh38.103.chr.reformatted.collapse_only.gene.ERCC.gtf,

and /hg38_GENCODE.v38.bed (see ADSP xqtl-protocol 0.1.1 for full details).

Genes with a TPM expression level of 10% or less in over 20% of samples

and/or genes with read counts less than 6 among fewer than 20% of samples

were excluded from the analysis. In addition, samples detected as outliers using

relative log expression (RLE), hierarchical clustering, and D-statistics were

removed. Finally, TPM data was normalized using the Trimmed Mean of M-value

(TMM) method.

182   Bulk RNA-seq Deconvolution and Hidden Factor Detection

As we used bulk whole blood RNA data for our analysis, we performed

CIBERSORTx deconvolution using the European-based LM22 reference panel to

estimate relative cell-type proportions in each sample[29]. We performed this

analysis using all samples combined. We included estimated cell-type

proportions for 14 estimated cell types as covariates in our analysis based on an

identified significant difference in relative cell-type proportions within each

population or across AD status (Kruskal-Wallis Test $p < 0.05$). The 14 estimated

cell-type proportions are Neutrophils, CD4+ T cells-Memory resting, B cells –

Naïve, B cells – Memory, Macrophages – M1, Mast cells – activated, T cells –

follicular helper, Plasma cells, T cells – CD4 naive, Regulatory T cells – Tregs,

Macrophages – M0, NK cells – activated, Monocytes, and NK cells - resting.

We also performed Hidden Factor Analysis using the Marchenko-Pastur PCA

approach as described in the ADSP xqtl-protocol 0.1.1. The approach applies the

Marchenko-Pastur distribution to determine the optimal number of principal

components to retain by modeling the eigenvalues of random covariance

matrices. Components with eigenvalues above the Marchenko-Pastur threshold

are considered to represent meaningful biological variation, while those below it

are attributed to random noise. This technique ensures that only components

capturing substantive biological variation are retained for downstream analyses,

avoiding overfitting while enhancing the interpretability of confounder correction

in our QTL mapping.

AD GWAS Summary Statistics

We accessed the largest available clinically diagnosed AD GWAS summary

statistics for each population. All summary statistics were derived from models

adjusted for age, sex, and population substructure principal components. We

lifted coordinates from GRCh37/hg19 using UCSC `liftOver`[30] as needed.

We accessed summary statistics from the Ray et al. African American GWAS,

which included 2,784 AD cases and 5,222 controls (total n=9,168)[31]. We also

accessed summary statistics from the Hispanic subset of Rajabli et al.,

comprising 3,005 AD cases and 5,894 controls (total n=8,899)[4]. For the Non-

Hispanic White population, we used Stage I data from Kunkle et al., which

included 21,982 AD cases and 44,944 controls (total n=63,926)[32].

Harmonization of GWAS and Expression Prediction Reference

To ensure consistency and improve the integration of GWAS summary statistics

and our gene expression prediction reference panel, we performed several

harmonization procedures. First, we ensured the reference and alternative alleles

of each variant were defined consistently across the GWAS and the reference

panel. If the reference and alternative alleles were flipped between the GWAS

and the reference panel, we inverted the GWAS Z-score so that its sign matched

the alternative allele in the reference panel. Second, since the prediction model

prioritizes weights mostly to eQTL variants within credible sets, we performed

summary statistics imputation to maximize the power and SNP coverage. We

followed the PredictDB GTEx v8 gwas imputation pipeline to impute the missing

variant z-scores in GWAS summary statistics, according to the MAGENTA

genotype reference panel for each population. 1,252,104, 1,253,727, and

1,081,547 variants are imputed for AA, EUR, and HISP GWAS respectively,

which represented 19.07% of total variants on average. Third, only unambiguous

SNPs, that do not involve substitutions of complementary base pairs, were kept.

Statistical methods

SuShiE

SuShiE[33] incorporated individual level gene expression and genotype data from

all three populations to infer credible sets and population-specific weights. We

followed empirical convention[23,24,33] and selected the expected maximum number

of credible sets, L, as 10 for each protein-coding gene. We specified posterior

inclusion probability (PIP) threshold for SNPs to be included in the credible sets

as 0.9. In some cases, credible sets may contain a few lead variants in tight LD

and low LD to others in a region. By definition, each credible set captures a

homogeneous and distinct signal. To refine the final inference results, credible

sets with lowest absolute pairwise correlation, which is defined as *purity*, and

weighted by sample sizes across all populations, are removed, if the purity is less

than 0.5. For the remaining parameters, default settings are used (Table S2).

SuShiE produces population-specific cis-eQTL weights to predict gene

expression (GReX). SNPs with large magnitude weights were identified by

SuShiE as members of credible sets. Such variable selection approaches reduce

TWAS estimation errors by reducing the potential of weak gene expression

prediction instruments[34]. SuShiE also computed cis-SNP heritability using the

`limix` python package for each gene within each population.

Cis-eQTL weighted prediction modeling was performed for each protein-coding

gene using SNPs within a $\pm$ 500kb region, adjusting for age, sex, PCs, cell-type

proportions and hidden factors.

FUSION
254

FUSION estimates the association between genetically regulated gene

expression (GReX) and Alzheimer's disease using GWAS summary statistics.

The TWAS test statistic (Z_{TWAS}) is computed as a weighted combination of

GWAS SNP-level Z-scores (Z_{gwas}), where the weights (W_{snp}) are derived from

the SNPs' effects on gene expression. To account for correlation structure

among SNPs, the test statistic is normalized by the square root of the weighted

linkage disequilibrium (LD) matrix:

$$Z_{TWAS} = \frac{W_{snp} * Z_{gwas}}{\sqrt{W_{snp} * LD * W_{snp}}}$$

We followed the FUSION[35] pipeline and used MAGENTA genotype data as the

LD reference panel for each population-stratified TWAS analysis. There are

14,127, 14,116, and 13,742 gene models available for NHW, AA, HISP

respectively. There are 13,731 common models shared across all populations.

SuShiE-derived population-specific cis-eQTL effect sizes were used to predict

the gene expression for each gene, and identify potential AD-related genes for

each population.

Meta analysis of population-stratified TWAS associations

Effective sample size weighting (SSW) meta-analysis was performed on three

population-stratified TWAS associations to share TWAS associations across all

populations. While inverse-variance weighting (IVW)[26] is generally preferred,

summary-level TWAS methods like FUSION cannot produce the effect sizes

(beta) and their standard errors required for this approach.

Due to the limited correlations and concordances of TWAS associations between

pairs of two populations, that was similarly observed in prior studies[26], we set the

FDR's q-value threshold to 0.05 ($q_c < 0.05$), in order to identify shared AD-

related gene candidates.

TWAS Fine-mapping

TWAS would nominate multiple genes, while some genes could be non-causal

due to sharing of eQTLs or co-regulated genes. We applied MAFOCUS[36] to

prioritize putative causal genes by correcting for correlation structures induced by

linkage disequilibrium (LD) and prediction weights in our TWAS results.

Population-specific genotype data from NHW, AA, and HISP groups in the

MAGENTA study served as LD reference panels. We performed fine-mapping on

LD blocks containing at least one significant gene ($q_c < 0.05$) using default

parameters. For each LD block, MAFOCUS calculated posterior inclusion

probabilities (PIPs) and derived 80% credible sets (ρ = 0.8) of putative causal

genes, both for individual populations and across all three populations. To

prioritize genes with the strongest evidence of causality, we established PIP

thresholds based on the MAFOCUS methodology; we classified a significant

TWAS association as a high-confidence causal gene if it achieved either a

population-specific $PIP_{pop} > 0.8$ in its corresponding population-stratified

analysis, or a multi-population $PIP_{ME} > 0.8$ in the sample-size weighted meta-

analysis that integrated evidence across all populations.

Independent Multi-ancestry Expression Reference Panel

In our analyses, we re-evaluate the significant TWAS associations from our

discovery dataset (MAGENTA) using the publicly released SuShiE-derived

weights from the TOPMed MESA visit-1 mRNA dataset[33,37,38]. As we did not have

access to the individual level genotype of the MESA samples, we used high-

coverage 1000 Genomes data to generate a LD reference for each ancestry[39].

The MESA dataset contained expression measurements from 956 participants

(402 European American, 175 African American, and 277 Hispanic American

individuals) with corresponding whole genome sequencing data. The mRNA

expression was quantified in peripheral blood mononuclear cells (PBMCs),

covering 21,747 protein-coding genes. The original analysis adjusted for 15 gene

expression principal components, 10 genotype principal components, age, sex,

and assay lab as covariates. These SuShiE-derived prediction models were

accessed at https://zenodo.org/records/10963034.

Results

MAGENTA Study Overview

We used the Multi-Ancestry Genomics, Epigenomics, and Transcriptomics of

Alzheimer's (MAGENTA) cohort as the eQTL reference, comprised of African

American (AA, n=224), Non-Hispanic White (NHW, n=235), and a combined

Hispanic group (HISP, n=292). The Hispanic group contains Caribbean and

Peruvian sub-cohorts to align with available AD GWAS summary statistics.

Case/control status was approximately balanced across groups (~50% each),

and age distributions were comparable (means ~76.6–77.5 years), supporting

cross-population comparability for downstream fine-mapping and TWAS

analyses (Table 1).

Table 1: **Sample characteristics of MAGENTA study participants.**

	Subcategory	All	AA	NHW	HISP
Sample Size (N)		751	224	235	292
Sex	Male	267 (35.6%)	64 (28.6%)	110 (46.8%)	93 (31.8%)
	Female	484 (64.4%)	160 (71.4%)	125 (53.2%)	199 (68.2%)
Age (years)	Mean ± SD	77.0 ± 7.7	76.6 ± 7.8	77.5 ± 7.4	76.9 ± 7.9
	Median [IQR]	77.0 [71.0-83.0]	76.5 [70.0-83.0]	78.0 [72.0-83.0]	77.0 [71.0-83.0]
	Range	53.0-98.0	63.0-97.0	65.0-97.0	53.0-98.0
Status	Case	378 (50.3%)	111 (49.6%)	118 (50.2%)	149 (51.0%)
	Control	373 (49.7%)	113 (50.4%)	117 (49.8%)	143 (49.0%)

*The table presents demographic and clinical characteristics of the MAGENTA study across all
participants and stratified by population groups (AA: African American, NHW: Non-Hispanic
White, HISP: Hispanic). For each characteristic, counts and percentages are shown for
categorical variables, while mean ± standard deviation, median [interquartile range], and range
are provided for continuous variables. Sample sizes, sex distribution, age statistics, and
Alzheimer's Disease status are reported for the total cohort and each population group.

330   We found that cis-eQTL effects were largely consistent whether or not

331   Alzheimer's disease (AD) status was included as a covariate in the MAGENTA

332   datasets (Figure S3, Table S3, Figure S4), consistent with previous studies[40–42].

From multi-ancestry fine-mapping to interpretable TWAS: workflow overview

*Figure 1: **Multi-population TWAS analytical workflow for Alzheimer's Disease.** The workflow integrates MAGENTA genotype and expression data (n=751; AA: 224, NHW: 235, HISP: 292) with published AD GWAS summary statistics across three population groups: Hispanic (Rajabli et al. 2023), African American (Ray et al. 2022), and Non-Hispanic White (Kunkle et al. 2019). SuShiE is applied for multi-population eQTL fine-mapping to create population-specific GReX prediction models, which are then used in FUSION analysis alongside GWAS summary statistics and population-matched reference panels. Population-stratified TWAS results are combined through sample size weighted (SSW) meta-analysis to produce multi-population TWAS associations. Fine-mapping of these results using MA-FOCUS identifies the most likely causal genes.*

Our multi-population workflow (Figure 1) integrates MAGENTA expression and genotype data with population-matched AD GWAS to generate interpretable TWAS associations. SuShiE multi-ancestry fine-mapping yields population-specific GReX models that we test with FUSION within each ancestry and meta-analyzed across populations, followed by MAFOCUS gene-level fine-mapping of TWAS hits to identify putative causal genes. A detailed dense versus sparse comparison demonstrating that credible-set variants recapitulate dense-model TWAS signals is presented in the "Sparse Models Using Fine-mapped eQTLs" subsection. Overall, We showed this workflow increases eQTL resolution and enables variant-resolved interpretation of TWAS signals.

Multi-population fine-mapping improves credible set precision and identifies shared eQTL effects

Using MAGENTA whole-blood RNA-seq and genotypes from AA, NHW, and HISP (14,436 shared protein-coding genes), we applied SuShiE multi-ancestry fine-mapping and identified credible sets (CS) for 8,748 genes. The mean number of CS per gene was 1.27 (median 1), with very few genes requiring more than seven CSs, supporting our choice of $L = 10$. Credible sets were compact (median 4 SNPs; Figure S1), consistent with a sparse cis-regulatory architecture[20–22,43,44].

369 *Table 2:* **Improved fine-mapping precision of shared cis-eQTLs through**
370 **multi-population analysis.**

Metric	Average of Any Two Populations	Three Populations (NHW+AA+HISP)	Improvement
Median SNPs per credible set	6	4	33% reduction
Mean credible sets per gene	0.52	1.27	144% increase
Percent of eQTL had PIP > 0.9	0.018	0.024	32% increase

*This table demonstrates the benefits of incorporating three diverse populations (Non-Hispanic
White, African American, and Hispanic) in SuShiE analysis compared to using only two
populations. The tri-population approach resulted in more precise credible sets (33% fewer SNPs
per set), identified more credible sets per gene (144% increase), and increased the proportion of
SNPs with high posterior inclusion probability (32% increase in SNPs with $PIP > 0.9$). These
findings support the sparse genetic architecture of gene expression regulation and highlight the
value of diverse population inclusion in fine-mapping studies.

Including all three populations improved fine-mapping precision relative to any

two-population analysis (NHW–AA, NHW–HISP, and AA–HISP) (Table 2);

median SNPs per CS decreased from 6 to 4 (–33%), consistent with reduced LD-

induced ambiguity across ancestries, the mean number of CS per gene

increased from 0.52 to 1.27 (+144%), and the fraction of SNPs with $PIP > 0.9$

rose from 0.018 to 0.024 (+32%), indicating improved confidence in variant-level

fine-mapping. These findings are consistent with the benefits of multi-ancestry

fine-mapping results reported by the PAGE consortium[45].

*Figure 2: **Cross-population analysis of eQTL effects in the first credible set (CS1).** (a) Density plots showing the distribution of effect size correlations across population pairs (NHW-AA, NHW-HISP, and AA-HISP) for all genes (n=8,748) (left) and genes with significant heritability (right, p < 0.05). The bimodal distributions indicate predominantly consistent effects across populations (~70% positive correlations) with some population-specific patterns. (b) Venn diagram showing overlap of genes with strong negative correlations (r < -0.5) between population pairs. The substantial pairwise overlaps (1223-1708 genes) suggests that genetic effects typically differ in one population while remaining consistent between the other two, with HISP showing the most distinct patterns (1708 overlapping genes with other pairs).*

Following the approach of Lu et al.[33], we evaluated cross-ancestry consistency of

effects for first SuShiE reported single shared effect (CS1), which represented

the largest effects on gene expressions. Effect size correlations were highly

consistent for roughly 70% of genes, indicating broadly shared regulation

(Figure 2) across groups. A subset of genes (~30%) showed anticorrelation (r < -

0.5), with 2,509 (AA–NHW), 2,990 (AA–HISP), and 2,954 (NHW–HISP) genes,

and substantial pairwise overlap (1,223–1,708 genes) where genes have a

different direction of effect in one population compared to the other two. These

findings highlight population-specific signals with HISP showing the most distinct

patterns.

We then assessed posterior eQTL weight concordance of variants in CS1 across

populations. For all genes, correlations were moderate (AA–NHW r=0.526; AA–

HISP r=0.379; NHW–HISP r=0.441), and increased when restricting to genes

with significant cis-heritability (AA–NHW r=0.689; AA–HISP r=0.484; NHW–HISP

r=0.540) (Figure S2). These results support substantial sharing of cis-regulatory

mechanisms across ancestries, with clearer concordance in genes under

stronger genetic control though notable population-specific effects remain.

Population-stratified TWAS reveals both shared and group-specific AD gene associations

We applied MAGENTA-derived, population-specific weights in FUSION to AD

GWAS from Kunkle et al. (NHW), Ray et al. (AA), and Rajabli et al. (HISP). Test

calibration was acceptable (λ_{GC}: NHW = 1.080, AA = 0.909, HISP = 0.979). At

$q_c < 0.05$ (NHW: $P < 1.4 * 10^{-4}$; AA: $P < 2.8 * 10^{-6}$; HISP: $P < 3.8 * 10^{-6}$), we

identified 40, 4, and 3 significant genes, respectively (Figure 3 (a–c); Table S6).

Using MAFOCUS to prioritize putative causal genes among the significant hits

$(q_c < 0.05)$, we identified high-confidence genes with $PIP > 0.8$: 12 in NHW and 2

each in AA and HISP.

*Figure 3: **population-stratified and meta-analyzed TWAS associations for***
***Alzheimer's Disease across diverse populations.** The figure displays TWAS*
associations as Manhattan plots with -log10(P-value) on the Y-axis, with positive
associations shown in the top panel and negative associations in the bottom
panel. The gray dashed lines indicate the population-stratified FDR threshold of
0.05, corresponding to $P_{NHW} < 1.4 * 10^{-4}$; $P_{AA} < 2.8 * 10^{-6}$; $P_{HISP} < 3.8 * 10^{-6}$;
$P_{META} < 1.4 * 10^{-4}$ for each respective population. Red dots highlight genes that
meet both statistical significance ($FDR < 0.05$) and fine-mapping criteria ($PIP >$
0.8), with gene symbols labeled for these prioritized associations. Each panel
represents different population-stratified or meta-analyzed associations. a) A
Miami plot of NHW population-stratified TWAS associations; b) A Miami plot of
AA ancestry-population TWAS associations; c) A Miami plot of HISP population-
stratified TWAS associations; d) A Miami plot of SSW meta-analyzed TWAS
associations

We first highlight within-population signals ($q_c < 0.05$, population-specific

$PIP > 0.8$) that did not reach significance in the multi-population meta-analysis.

In NHW, two genes outside known AD loci were significant: *CEBPZOS* and

COG4 (Figure 3 (a)). *CEBPZOS* showed a negative association ($Z_{NHW} = -4.076$,

$P = 4.58 * 10^{-5}$). SuShiE fine-mapping identified 11 SNPs in 5 credible sets;

CS1–CS2 exhibited near-perfect cross-population effect concordance ($\overline{\rho} \approx$

0.999), whereas CS3–CS5 were population-specific (anticorrelated). In particular,

NHW's effect size correlation from CS4-5 exhibited strong anticorrelation than

HISP and AA (Figure S5). Expression was strongly predictable in NHW ($R^2 =$

0.225) and modestly but significantly in AA ($R^2 = 0.0246$, $P = 0.0106$). Using

independent SuShiE models from TOPMed MESA, the NHW effect direction

validated ($Z_{NHW} = -2.513$). No genome-wide significant GWAS SNPs were

present within ±1 Mb ($P < 5 * 10^{-8}$; the most significant association is from

rs1072218 $P = 5.50 * 10^{-5}$). The direction agrees with prior NHW TWAS[46]

($OR = 0.97$; 0.96–0.99).

COG4 showed a significant association with AD risk in NHW ($q_c < 0.05$, $Z_{NHW} = $

4.601, $P = 4.20 * 10^{-6}$; population-specific $PIP > 0.8$). SuShiE fine-mapping

identified 9 SNPs in two credible sets, with strong anticorrelation of effects

between NHW and the other populations ($\overline{\rho} \approx -0.934$). Expression was

genetically predictable in both NHW ($R^2 = 0.0487$, $P < 0.05$) and AA ($R^2 = 0.123$,

$P < 0.05$), yet AA showed no TWAS association ($Z_{AA} = -0.00317$, $P = 0.997$). No

genome-wide significant GWAS SNPs were present within ±1 Mb ($P < 5 * 10^{-8}$;

the most significant association is from rs3752786 $P = 3.98 * 10^{-6}$). Using

independent SuShiE models from TOPMed MESA, the NHW direction replicated

but was not significant ($Z_{NHW} = 0.281$, $P = 0.779$}); attenuation is plausibly due to

reduced cis-SNP coverage and differing fine-mapped eQTLs in MESA (±500 kb;

MAGENTA $N_{snp} = 1358$ vs. MESA $N_{snp} = 783$, −42.3%) and potential age/AD-

context differences.

In HISP TWAS, one gene reached significance at $q_c < 0.05$: *GARIN2* (also

known as *FAM71D*) on 14q23.3 ($Z_{HISP} = -4.622$, $P = 3.80 * 10^{-6}$) (Figure 3 (c)).

No SuShiE credible sets were identified and 5-fold CV was not significant

($R^2 = -0.00345$, $P = 0.991$); SuShiE posterior weights in HISP were uniformly

small (max magnitude $\approx 1 * 10^{-5}$), indicating the TWAS signal is unlikely to be

mediated by predicted expression. The result may instead reflect aggregated

weak SNP effects within HISP that may tag other variants in the region

(Figure S8).

In AA TWAS, *MARK4* was significant ($Z_{AA} = -4.710, P = 2.50 \times 10^{-6}$) within the

APOE region (~ 170,000 bp from *APOE*). SuShiE fine-mapping identified 13

SNPs across five credible sets; CS1 showed near-perfect cross-ancestry

concordance ($\overline{\rho}_1 \approx 1$), while averaging across all sets indicated heterogeneity

($\overline{\rho}_{AA\text{-}NHW} \approx 0.200; \overline{\rho}_{AA\text{-}HISP} \approx 0.599; \overline{\rho}_{NHW\text{-}HISP} \approx -0.199$). Predictive performance

was not significant in AA (5-fold CV R^2 not significant ($R^2 = 0.0033; P =$

$0.190 =$)), but was significant in NHW and HISP ($R^2_{NHW} = 0.308; P = 1.35 * 10^{-20}$,

$R^2_{HISP} = 0.102; P = 1.52 * 10^{-8}$). In TOPMed MESA, the AA effect direction was

consistent but not significant ($Z_{AA} = -1.662, P = 9.66 * 10^{-2}$). Given regional

complexity and proximity to *APOE* (Figure S7), this association should be

interpreted with caution.

We explored shared TWAS associations using Sample Size Weighted (SSW)

Meta-analysis. There are 39 statistically significant ($q_c < 0.05; P_{META} < 1.4 *$

10^{-4}) genes, and 9 of them had MAFOCUS's $PIP > 0.8$ (Figure 3 (d)). All fine-

mapped meta-analysis associations were driven mostly by the NHW population,

due to its larger sample size. Relative to the NHW TWAS, no new associations

became significant after meta-analysis. However, the sign concordance of the 9

identified genes between NHW and the other two populations is 66.67%, which

demonstrates moderate consistency and some heterogeneity across populations.

*Table 3: **Multi-population TWAS significant genes identified by SSW meta-***
analysis.

CHR	SBP	EBP	Gene Symbol	Meta Z	Meta P	Z AA	Z NHW	Z HISP
2	126656133	126696667	GYPC	-4.555	5.242e-06	-1.213	-4.433	-0.292

CHR	SBP	EBP	Gene Symbol	Meta Z	Meta P	Z AA	Z NHW	Z HISP
2	127048027	127107288	BIN1	5.176	2.269e-07	2.056	4.734	1.856
8	27311482	27459391	PTK2B	4.482	7.380e-06	0.453	4.560	-0.406
11	47331406	47352702	MYBPC3	6.162	7.169e-10	-1.183	6.283	1.148
11	60200270	60243137	MS4A4E	7.311	2.651e-13	-1.438	7.730	-0.610
11	85657742	85682908	CREBZF	-5.788	7.140e-09	1.103	-5.861	-1.356
19	44890569	44903689	TOMM40	47.363	1.000e-50	10.800	47.700	-6.752
19	45178784	45216933	BLOC1S3	21.455	1.000e-50	2.760	21.200	2.017
19	45769709	45782552	DMPK	5.565	2.621e-08	0.099	5.500	1.130

*SSW meta-analysis ($q_c < 0.05$) and MAFOCUS ($PIP > 0.8$). chromosomal location (CHR), start
base position (SBP), end base position (EBP), population-stratified TWAS z-scores (Z AA, Z
NHW, Z HISP).

In total, we identified multiple significant TWAS associations across

chromosomes 2, 8, 11, and 19, with variable patterns of genetic regulation

across populations (Table Table 3).

Chromosome 2 – *BIN1* and *GYPC*

BIN1 showed highly consistent TWAS associations across AA, NHW, and HISP

populations, with fine-mapping implicating three SNPs in two credible sets (mean

cross-population effect size correlation $\overline{\rho} \approx 0.956$). The lead eQTL rs11682128

(previously linked to *BIN1* expression in blood and brain) was in moderate LD

($r^2 \approx 0.34$; 53,336 bp away) with the top GWAS SNP rs6733839. 5-fold cross-

validation R^2 (0.116–0.263) was significant in all populations.

GYPC, located near *BIN1*, also showed consistent TWAS associations but had

significant R^2 only in the HISP population.

Chromosome 8 – *PTK2B*

PTK2B was significant in the meta-analysis and fine-mapping identified 28 SNPs

in three credible sets with high effect size concordance ($\bar{\rho} \approx 0.995$). Notably, the

28 SNPs are in minimal LD with the GWAS index SNP, rs2741342, with the

highest LD in the first credible set ($r^2 \approx 0.1$). Predictive performance ($R^2 =$

$0.040 - 0.120$) were significant in all populations, though TWAS significance was

observed mainly in NHW, with consistent direction across datasets.

Chromosome 11 – *MYBPC3*, *MS4A4E*, and *CREBZF*

MYBPC3 showed the strongest signal in NHW, with high cross-population effect

size correlation ($\bar{\rho} \approx 0.999$). R^2 was significant in only NHW ($h^2 = 0.052$).

MS4A4E was driven by the NHW signal, with weak or negative effect size

correlation with AA and HISP (AA-NHW: $\rho \approx 0.004$; AA-HISP: $\rho \approx -0.004$) and

no significant predictive performance in any population. *CREBZF* showed no

significant R^2, or fine-mapped variants, and was likely influenced by nearby

strong GWAS signals ($P < 1 * 10^{-8}$ from 86Mb to 86.2Mb).

Chromosome 19 – *TOMM40*, *DMPK*, and *BLOC1S3*

Among the three genes near APOE locus, only *TOMM40*'s SuShiE TWAS

prediction model contained APOE-ε4 allele, rs429358 in its first CS.

TOMM40 exhibited strong but heterogeneous associations, with significant R^2

only in NHW and variable effect directions across populations. The learned prior

effect size correlations showed substantial heterogeneity across populations

especially in HISP, with moderate positive correlation between AA and NHW

($\overline{\rho} \approx 0.599$), but negative correlations between AA-HISP ($\overline{\rho} \approx -0.492$) and NHW-

HISP ($\overline{\rho} \approx -0.190$). *DMPK* showed consistent positive direction across

populations, moderate cross-population correlations ($\overline{\rho}_{AA-HISP} \approx 0.661$;

$\overline{\rho}_{AA-NHW} \approx 0.382$; $\overline{\rho}_{NHW-HISP} \approx 0.053$), and significant predictive performance

($R^2 = 0.026 - 0.146$) in all groups. *BLOC1S3* had highly consistent effect sizes

across populations ($\overline{\rho} \approx 0.998$) but significant R^2 only in NHW ($R^2 = 0.031$;

$h^2 = 0.048$).

Across loci, patterns emerged where some genes (*BIN1*, *PTK2B*, *BLOC1S3*) had

consistent cross-population genetic regulation, whereas others (*MYBPC3*,

TOMM40, *MS4A4E*) displayed marked population-specific architectures in one

group. Using direction as the primary criterion, we observed replicated effects in

TOPMed MESA for several discovery genes in at least two ancestries—notably

BIN1, *PTK2B*, *DMPK*, *TOMM40*, and *MYBPC3*, and *MS4A4E*—with significance

most evident in NHW and generally insignificant in AA and HISP, consistent with

GWAS sample size differences. Detailed association results, cis-h^2, R^2, and

replication outcomes are presented in Table 3, Table S5, and Table S7.

Sparse models using fine-mapped eQTLs capture key TWAS associations and implicate variants beyond GWAS index SNPs

Across fine-mapped TWAS hits (MAFOCUS PIP>0.8), credible sets typically did

not include the GWAS index SNPs. At *BIN1*, the top CS SNP rs11682128 was

only modestly correlated with the NHW index SNP rs6733839 ($r^2 \approx 0.34$),

indicating additional regulatory variants beyond the GWAS lead. Similar patterns

were observed at *PTK2B* and *DMPK*, where rs2741342 and rs429358 are not in

the credible sets for each gene respectively. However, in the APOE region

TOMM40's CS1 included the ε4-defining rs429358 and drove heterogeneous

associations across populations. Dense-versus-sparse comparisons showed that

restricting models to credible-set variants largely preserved association

magnitudes (Table 4), supporting a fine-mapped variant-level regulatory basis for

these TWAS signals.

To confirm that fine-mapped eQTLs in credible sets provide adequate models for

TWAS associations, we compared the TWAS Z-scores between two approaches:

a dense model using SuShiE expression estimates for all SNPs within the cis-

region (±500kb), and a sparse model using only SNPs identified in credible sets.

Our analysis focused on significant genes identified in the dense model

approach, examining the consistency and magnitude of associations across both

modeling strategies. A detailed summary of the credible sets for these significant

genes, including the number of variants, posterior inclusion probabilities, and

other fine-mapping statistics, is provided in the Supplementary Material

(Table S4).

The comparison revealed remarkable consistency between dense and sparse

models for most genes (Table 4). Key AD-associated genes like *BIN1*, *TOMM40*,

COG4, and *MARK4* showed particularly stable associations across both

approaches. For instance, *BIN1* maintained consistent negative associations

across all populations in dense and sparse models (NHW: 4.73 vs 4.34; AA: 2.06

vs 2.00; HISP: 1.86 vs 1.60), with only slight attenuation in the sparse model.

Similarly, *TOMM40* preserved its strong population-specific pattern, showing

nearly identical Z-scores in both approaches for all populations (NHW: 47.7 vs

48.3; AA: 10.8 vs 10.8; HISP: -6.75 vs -6.88). However, the strong and

heterogeneous associations of *TOMM40* are likely driven by the APOE effect, as

the APOE ε4 allele, rs429358, is present in the first CS.

The novel association we identified, *COG4*, also demonstrated stability between

dense and sparse models (NHW: 4.6 vs 3.88; AA: 0 vs 0.29; HISP: -0.93 vs -

0.42), suggesting that its association with AD is primarily driven by the fine-

mapped eQTLs.

However, some genes showed notable differences between the two approaches.

For example, *BLOC1S3* in the APOE region showed different magnitudes of

association in the dense versus sparse models in the NHW population (21.2 vs

3.47). This difference could be explained by the complex LD structure and APOE

signals in the region, where many variants could contribute to the association

signal through LD with known AD risk variants. Similarly, *GYPC* showed marked

differences in the NHW population (-4.43 vs 0.06), suggesting that some of its

association might be driven by SNPs outside the credible sets. Since *GYPC* is in

close proximity to *BIN1*, the strong TWAS association of *GYPC* in NHW, is likely

to be driven by rs6733839 (not in CSs of *GYPC*), which is an index SNP in the

region of NHW AD GWAS.

Overall, our comparison demonstrates that fine-mapped eQTLs in credible sets

can largely explain the TWAS associations for most genes, particularly those with

strong and consistent effects across populations. The stability of associations

between dense and sparse models provides additional support for the sparse

genetic architecture of gene expression regulation and validates our fine-

mapping approach.

*Table 4: **Comparison of Dense and Sparse TWAS Z-scores for Dense Model-**
Identified Significant Genes*

Chr	Gene	NHW dense	NHW sparse	AA dense	AA sparse	HISP dense	HISP sparse
2	BIN1	4.73	4.34	2.06	2	1.86	1.6
2	GYPC	-4.43	0.06	-1.21	-1.98	-0.29	0.3
8	PTK2B	4.56	4.39	0.45	0.42	-0.41	0.15
11	MS4A4E	7.73	7.74	-1.44	-1.48	-0.61	-0.51
11	MYBPC3	6.28	6.29	-1.18	-1.18	1.15	0.96
16	COG4	4.6	3.88	0	0.29	-0.93	-0.42
19	BLOC1S3	21.2	3.47	2.76	2.16	2.02	2.53
19	DMPK	5.5	5.31	0.1	0.08	1.13	1.01
19	MARK4	10.9	11.5	-4.71	-4.45	-1.17	-2.19
19	TOMM40	47.7	48.3	10.8	10.8	-6.75	-6.88

*TWAS Z-scores comparing dense and sparse modeling approaches across three populations
(Non-Hispanic White (NHW), African American (AA), and Hispanic (HISP)). Chr: Chromosome.

As illustrated in Figure 4, *BIN1*'s variant (rs11682128) in the first credible set

(CS1) exhibited consistently positive eQTL weights across all three populations,

with magnitudes ranging from 0.44 to 0.64. The visualization further revealed

population-specific patterns in the corresponding GWAS signals, with the

strongest Z-scores observed in the NHW population (~4.5), followed by more

moderate signals in AA and Hispanic populations (1-2), which may explain the

magnitude of TWAS associations across ancestries. The difference could be due

to the sample size differences in GWAS studies, or real effect size differences

across populations as we observe with APOE[5]. Despite these differences in

GWAS signal intensity, the direction and relative magnitude of the TWAS

associations remain consistent (Table 4), suggesting that the identified credible

set variants captured the core regulatory mechanisms influencing BIN1

expression in relation to AD risk. The genomic context provided in the bottom

panel demonstrates that the two credible sets are located in the intronic region of

BIN1. Collectively, this integrated view of BIN1 regulation supports our finding

that sparse models using only credible set variants can effectively recapitulate

the signals detected in dense models, while providing greater interpretability

regarding the specific variants driving these associations.

*Figure 4: **Fine-mapped eQTLs and GWAS associations for the BIN1
locus.** The figure displays the relationship between eQTL weights (top panel),
GWAS Z-scores (middle panel), and gene structure (bottom panel) for BIN1 on
chromosome 2. In the top and middle panel, eQTL weights and GWAS
associations are shown across three ancestral populations (African
American (AA, circles), Hispanic (HISP, squares), and Non-Hispanic White
(NHW, triangles)), with variants colored by credible set membership. Point sizes*

indicate posterior inclusion probability (PIP) values. The bottom panel illustrates
the genomic structure of BIN1 at chromosome 2. In CS1, there is only one
variant, rs11682128; In CS2, there are two variants, rs6714626 and rs6710752

COG4 is identified as a novel AD-related gene from our main analysis. When we

focused on the sparse model, which included only 9 variants in 2 credible sets,

we observed a similar TWAS association ($Z_{NHW} = 3.88$) in the NHW population,

while the other two populations did not show significant associations (Table 4).

After analyzing the SNP weights and GWAS associations for these fine-mapped

SNPs (Figure 5), we noticed substantial differences in the NHW population,

particularly in the second credible set. For example, rs11639579, the fine-

mapped SNP in the second credible set with the largest PIP ($PIP = 0.709$),

exhibited the largest difference in NHW's GWAS association and SNP weight

compared to the other two populations (Table 4). We observed that this variant is

also located near *IL34*, but the SNP is not fine-mapped in *IL34*'s credible set, and

the TWAS association of *IL34* is not significant in the dense model.

*Figure 5: **Fine-mapped eQTLs and GWAS associations for the COG4***
***locus.** The figure displays the relationship between eQTL weights (top panel),*
GWAS Z-scores (middle panel), and gene structure (bottom panel) for TOMM40
on chromosome 19. In the top and middle panels, eQTL weights and GWAS
associations are shown across three ancestral populations (African American
(AA, circles), Hispanic (HISP, squares), and Non-Hispanic White (NHW,
triangles)), with variants colored by credible set membership. Point sizes indicate

665 *posterior inclusion probability (PIP) values. The bottom panel illustrates*
666 *the genomic structure of COG4 at chromosome 16.*

Discussion
667

In this multi-ancestry TWAS of AD, we show that jointly modeling cis-eQTLs

across African American, Non-Hispanic White, and Hispanic groups improves

fine-mapping resolution and yields more interpretable, variant-resolved

associations. Through the lens of whole-blood gene expression, we implicate

regulatory variants beyond GWAS index SNPs at established loci at BIN1,

PTK2B, and DMPK. We further nominate COG4 as a novel candidate gene for

AD.

Importantly, we demonstrate that sparse credible-set models from our fine-

mapping recapitulate dense-model associations, concentrating evidence onto

compact variant sets without a significant loss of association signal. Our multi-

ancestry fine-mapping of eQTL results decreased the number of SNPs in each

credible set while also increasing the overall number of credible sets. These

findings are consistent with the results reported in Lu et al.[33]. and provide

additional empirical evidence supporting the sparse genetic architecture of gene

expression regulation previously reported in multiple studies[20–22]. Our results also

align with prior studies showing that while differences in LD and allele

frequencies across ancestral groups might lead to heterogeneous eQTL effects,

the correlation is largely consistent across populations[33,47–49].

Our multi-population TWAS analysis revealed both known and novel associations

with AD risk. Importantly, many of the TWAS associations appear to be driven

not by GWAS index SNPs, but by fine-mapped regulatory variants highlighted in

credible sets, suggesting that the detected AD TWAS associations are driven by

subsets of gene expression regulatory variants. We identified regulatory variants

for *BIN1*, *PTK2B*, *DMPK*, and *GYPC*, all mostly independent of established

GWAS signals at these loci. *BIN1* demonstrated remarkable consistency in

TWAS associations that were successfully confirmed in the TOPMed MESA

dataset across three groups. Importantly, our results confirmed the eQTL effect

of rs11682128 is shared across three populations, which was previously reported

by Zhu et al[50]. Lastly, two additional SNPs were identified in the second credible

set, which implied more regulatory variants in BIN1.

We also identified *CEBPZOS* (which has been reported from prior TWAS[50]) and

COG4, neither of which have nearby genome-wide significant GWAS signals. To

the best of our knowledge, *COG4* had not been identified as an AD-related gene

previously. Notably, single-ancestry fine-mapping in NHW alone identified no

credible sets, whereas including all three populations yielded two credible sets,

underscoring the advantage of multi-ancestry fine-mapping for variable selection.

Specifically, four variants in the first credible set had higher allele frequency in

AA than the other two populations Table S8. *COG4* encodes a component of an

oligomeric protein complex that plays a crucial role in the structure and function

of the Golgi apparatus, which is essential for processing and distributing proteins

necessary for neuronal function. Notably, rs11639579, the SNP in the second

credible set with the largest PIP ($PIP = 0.709$), exhibited substantial differences

in GWAS association and SNP weight in the NHW population compared to AA

and HISP populations. According to Agora[51], an AD knowledge portal, *COG4* has

meaningful expression across brain regions. The differential expression, between

AD cases and controls, is significant in cerebellum ($\log_2 FC = -0.149$, $P = 1.44 *$

10^{-3}) and parahippocampal gyrus ($\log_2 FC = 0.0827$, $P = 1.44 * 10^{-2}$).

Furthermore, Golgi fragmentation occurs in many neurodegenerative diseases.

In AD, Golgi fragmentation results in enhanced APP trafficking and Aβ

production[52,53]. The significant association of genetically predicted *COG4*

expression in blood with AD suggests a potentially underexplored link between

systemic vesicle trafficking pathways and disease risk.

Our study faces several important limitations. First, while whole blood is readily

accessible and often used in large-scale genomic studies, it may not optimally

reflect the tissue-specific regulatory mechanisms occurring in the brain.

Currently, the scarcity of brain-specific eQTL data from diverse populations

constrains our ability to conduct optimal TWAS analysis and validate these

findings in the most relevant tissue. However, several previous literatures

observed largely concordant AD TWAS effect size directions and eQTL effect

sizes between brain and peripheral tissues[54,55]. Additionally, Huseby et al. found

eight fundamental cell biological functions that are altered in whole blood

samples from six neurodegenerative diseases[56], suggesting AD risk can be

modulated by multiple systemic factors[57]. Second, the substantial disparity in

GWAS sample sizes between populations significantly impacted our ability to

detect population-specific associations. This imbalance is reflected in our meta-

analysis results, where the significant associations were predominantly driven by

NHW signals, as evidenced by the stronger Z-scores in NHW population across

most identified genes (Table 3). Nevertheless, despite these power limitations,

we observed consistent effect directions across populations for several key

genes. Third, a practical limitation of multi-ancestry fine-mapping (including

SuShiE) is the requirement to analyze only SNPs shared across populations to

ensure matched LD and comparability; such filtering can exclude ancestry-

unique variants, biasing credible sets toward shared signals and potentially

underestimating truly population-specific regulatory variants, especially in AA and

admixed HISP where allele frequencies and LD differ. We attempted to mitigate

extreme sparsity by applying a per-population MAC threshold (MAC ≥ 10). Future

work should explicitly accommodate ancestry-unique variants in multi-ancestry

fine-mapping method developments, and expand recruitment of

underrepresented populations to enable adequately powered ancestry-specific

fine-mapping and TWAS that are comparable to EUR-based studies. Forth, we

acknowledge that reliance on the European-derived LM22 panel may reduce cell

type deconvolution accuracy and introduce population-specific bias in our AA and

HISP cohorts, so these cell-type covariates should be interpreted cautiously and

revisited with ancestry-matched or single-cell–derived references as they

become available.

Several key future directions could build upon our findings. First, as brain-specific

multi-population eQTL datasets become available, replicating our analyses in

neural tissue would be valuable for validating these associations and potentially

uncovering additional AD-related genes. More critically, ongoing efforts to

expand AD GWAS in non-European populations will be crucial for achieving

more balanced statistical power across populations. This enhanced power would

not only improve our ability to detect population-specific effects but also provide

more robust evidence for shared genetic mechanisms across populations. Such

advancements would help address the current European bias in genetic studies

and contribute to a more comprehensive understanding of AD's genetic

architecture across diverse populations.

Acknowledgement

765 This work was supported by grants from the National Institute on Aging

766 R01AG070935 (Bush, Griswold), U19AG074865 (Pericak-Vance, Byrd, Bush,

767 Haines, Kunkle, Reitz, Vance), and RF1AG061351 (Below, Bush, Naj). This work

768 made use of the High Performance Computing Resource in the Core Facility for

769 Advanced Research Computing at Case Western Reserve University.

References

1. (2024). 2024 Alzheimer's disease facts and figures. Alzheimer's & Dementia
20, 3708–3821.

2. Daniel, E.V., Kleiman, M.J., and Galvin, J.E. (2023). Exploring Reasons for
Differential Vulnerability and Alzheimer's Disease Risk in Racial and Ethnic
Minorities. Journal of Alzheimer's Disease : JAD *91*, 495.

3. Kunkle, B.W., Schmidt, M., Klein, H.-U., Naj, A.C., Hamilton-Nelson, K.L.,
Larson, E.B., Evans, D.A., De Jager, P.L., Crane, P.K., Buxbaum, J.D., et al.
(2021). Novel Alzheimer Disease Risk Loci and Pathways in African American
Individuals Using the African Genome Resources Panel: A Meta-analysis. JAMA
Neurology *78*, 102–113.

4. Rajabli, F., Benchek, P., Tosto, G., Kushch, N., Sha, J., Bazemore, K., Zhu,
C., Lee, W.-P., Haut, J., Hamilton-Nelson, K.L., et al. (2025). Multi-ancestry
genome-wide meta-analysis of 56,241 individuals identifies known and novel

cross-population and ancestry-specific associations as novel risk loci for
Alzheimer's disease. Genome Biology *26*, 210.

5. Belloy, M.E., Andrews, S.J., Le Guen, Y., Cuccaro, M., Farrer, L.A., Napolioni,
V., and Greicius, M.D. (2023). APOE Genotype and Alzheimer Disease Risk
Across Age, Sex, and Population Ancestry. JAMA Neurology *80*, 1284–1294.

6. Bettcher, B.M., Tansey, M.G., Dorothée, G., and Heneka, M.T. (2021).
Peripheral and central immune system crosstalk in Alzheimer disease — a
research prospectus. Nature Reviews Neurology *17*, 689–701.

7. Mews, M.A., Naj, A.C., Griswold, A.J., Alzheimer's Disease Genetics
Consortium, Below, J.E., and Bush, W.S. (2025). Brain and blood transcriptome-
wide association studies identify five novel genes associated with Alzheimer's
disease. Journal of Alzheimer's Disease: JAD *105*, 228–244.

8. Zhou, D., Jiang, Y., Zhong, X., Cox, N.J., Liu, C., and Gamazon, E.R. (2020).
A unified framework for joint-tissue transcriptome-wide association and
Mendelian randomization analysis. Nature Genetics *52*, 1239–1246.

9. Liu, W., Li, M., Zhang, W., Zhou, G., Wu, X., Wang, J., Lu, Q., and Zhao, H.
(2020). Leveraging functional annotation to identify genes associated with
complex diseases. PLoS Computational Biology *16*, e1008315.

10. Gockley, J., Montgomery, K.S., Poehlman, W.L., Wiley, J.C., Liu, Y.,
Gerasimov, E., Greenwood, A.K., Sieberts, S.K., Wingo, A.P., Wingo, T.S., et al.
(2021). Multi-tissue neocortical transcriptome-wide association study implicates 8
genes across 6 genomic loci in Alzheimer's disease. Genome Medicine *13*, 76.

11. Chen, H.-H., Petty, L.E., Sha, J., Zhao, Y., Kuzma, A., Valladares, O.,
Alzheimer's Disease Genetics Consortium, International Genomics of
Alzheimer's Project, Bush, W., Naj, A.C., Gamazon, E.R., et al. (2021).
Genetically regulated expression in late-onset Alzheimer's disease implicates risk
genes within known and novel loci. Translational Psychiatry *11*, 618.

12. Wang, Y.H., Luo, P.P., Geng, A.Y., Li, X., Liu, T.-H., He, Y.J., Huang, L., and
Tang, Y.Q. (2023). Identification of highly reliable risk genes for Alzheimer's
disease through joint-tissue integrative analysis. Frontiers in Aging Neuroscience
15, 1183119.

13. Bae, Y.E., Wu, L., and Wu, C. (2021). InTACT: An adaptive and powerful
framework for joint-tissue transcriptome-wide association studies. Genetic
Epidemiology *45*, 848–859.

14. Hu, Y., Li, M., Lu, Q., Weng, H., Wang, J., Zekavat, S.M., Yu, Z., Li, B., Gu,
J., Muchnik, S., et al. (2019). A statistical framework for cross-tissue
transcriptome-wide association analysis. Nature Genetics *51*, 568–576.

15. Sun, Y., Zhu, J., Zhou, D., Canchi, S., Wu, C., Cox, N.J., Rissman, R.A.,
Gamazon, E.R., and Wu, L. (2021). A transcriptome-wide association study of
Alzheimer's disease using prediction models of relevant tissues identifies novel
candidate susceptibility genes. Genome Medicine *13*, 141.

16. Tang, S., Buchman, A.S., De Jager, P.L., Bennett, D.A., Epstein, M.P., and
Yang, J. (2021). Novel Variance-Component TWAS method for studying complex
human diseases with applications to Alzheimer's dementia. PLoS Genetics *17*,
e1009482.

17. Luningham, J.M., Chen, J., Tang, S., De Jager, P.L., Bennett, D.A.,
Buchman, A.S., and Yang, J. (2020). Bayesian Genome-wide TWAS Method to
Leverage both cis- and trans-eQTL Information through Summary Statistics.
American Journal of Human Genetics *107*, 714–726.

18. Zou, H., and Hastie, T. (2005). Regularization and Variable Selection Via the
Elastic Net. Journal of the Royal Statistical Society Series B: Statistical
Methodology *67*, 301–320.

19. Tibshirani, R. (1996). Regression Shrinkage and Selection via the Lasso.
Journal of the Royal Statistical Society. Series B (Methodological) *58*, 267–288.

20. Wheeler, H.E., Shah, K.P., Brenner, J., Garcia, T., Aquino-Michaels, K., Cox,
N.J., Nicolae, D.L., and Im, H.K. (2016). Survey of the Heritability and Sparse
Architecture of Gene Expression Traits across Human Tissues. PLoS Genetics
12, e1006423.

21. Barbeira, A.N., Melia, O.J., Liang, Y., Bonazzola, R., Wang, G., Wheeler,
H.E., Aguet, F., Ardlie, K.G., Wen, X., and Im, H.K. (2020). Fine-mapping and
QTL tissue-sharing information improves the reliability of causal gene
identification. Genetic Epidemiology *44*, 854–867.

22. Consortium, T.G. (2020). The GTEx Consortium atlas of genetic regulatory
effects across human tissues. Science.

23. Zou, Y., Carbonetto, P., Wang, G., and Stephens, M. (2022). Fine-mapping
from summary data with the "Sum of Single Effects" model. PLoS Genetics *18*,
e1010299.

24. Wang, G., Sarkar, A., Carbonetto, P., and Stephens, M. (2020). A Simple
New Approach to Variable Selection in Regression, with Application to Genetic
Fine Mapping. Journal of the Royal Statistical Society Series B: Statistical
Methodology *82*, 1273–1300.

25. Zhao, S., Crouse, W., Qian, S., Luo, K., Stephens, M., and He, X. (2024).
Adjusting for genetic confounders in transcriptome-wide association studies
improves discovery of risk genes of complex traits. Nature Genetics 1–12.

26. Bhattacharya, A., Hirbo, J.B., Zhou, D., Zhou, W., Zheng, J., Kanai, M.,
Global Biobank Meta-analysis Initiative, Pasaniuc, B., Gamazon, E.R., and Cox,
N.J. (2022). Best practices for multi-ancestry, meta-analytic transcriptome-wide
association studies: Lessons from the Global Biobank Meta-analysis Initiative.
Cell Genomics *2*, 100180.

27. (2023). Using Population Descriptors in Genetics and Genomics Research: A
New Framework for an Evolving Field
(https://pubmed.ncbi.nlm.nih.gov/36989389/).

28. Griswold, A.J., Sivasankaran, S.K., Van Booven, D., Gardner, O.K., Rajabli,
F., Whitehead, P.L., Hamilton-Nelson, K.L., Adams, L.D., Scott, A.M., Hofmann,
N.K., et al. (2020). Immune and Inflammatory Pathways Implicated by Whole
Blood Transcriptomic Analysis in a Diverse Ancestry Alzheimer's Disease
Cohort. Journal of Alzheimer's Disease *76*, 1047–1060.

29. Steen, C.B., Liu, C.L., Alizadeh, A.A., and Newman, A.M. (2020). Profiling
Cell Type Abundance and Expression in Bulk Tissues with CIBERSORTx.
Methods in Molecular Biology (Clifton, N.J.) *2117*, 135.

30. Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G.,
Clawson, H., Diekhans, M., Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The
UCSC Genome Browser Database: Update 2006. Nucleic Acids Research *34*,
D590–D598.

31. Extended genome-wide association study employing the African Genome
Resources Panel identifies novel susceptibility loci for Alzheimer's Disease in
individuals of African ancestry - Ray - 2023 - Alzheimer's & Dementia - Wiley
Online Library (https://alz-
journals.onlinelibrary.wiley.com/doi/10.1002/alz.078904).

32. Kunkle, B.W., Grenier-Boley, B., Sims, R., Bis, J.C., Damotte, V., Naj, A.C.,
Boland, A., Vronskaya, M., van der Lee, S.J., Amlie-Wolf, A., et al. (2019).
Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci
and implicates Aβ, tau, immunity and lipid processing. Nature Genetics *51*, 414–
430.

33. Lu, Z., Wang, X., Carr, M., Kim, A., Gazal, S., Mohammadi, P., Wu, L.,
Pirruccello, J., Kachuri, L., Gusev, A., et al. (2025). Improved multiancestry fine-
mapping identifies cis-regulatory variants underlying molecular traits and disease
risk. Nature Genetics *57*, 1881–1889.

34. Lorincz-Comi, N., Yang, Y., Li, G., and Zhu, X. (2024). MRBEE: A bias-
corrected multivariable Mendelian randomization method. HGG Advances *5*,
100290.

35. Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W.J.H., Jansen,
R., de Geus, E.J.C., Boomsma, D.I., Wright, F.A., et al. (2016). Integrative

approaches for large-scale transcriptome-wide association studies. Nature
Genetics *48*, 245–252.

36. Lu, Z., Gopalan, S., Yuan, D., Conti, D.V., Pasaniuc, B., Gusev, A., and
Mancuso, N. (2022). Multi-ancestry fine-mapping improves precision to identify
causal genes in transcriptome-wide association studies. The American Journal of
Human Genetics *109*, 1388–1404.

37. Bild, D.E., Detrano, R., Peterson, D., Guerci, A., Liu, K., Shahar, E., Ouyang,
P., Jackson, S., and Saad, M.F. (2005). Ethnic Differences in Coronary
Calcification. Circulation *111*, 1313–1320.

38. Taliun, D., Harris, D.N., Kessler, M.D., Carlson, J., Szpiech, Z.A., Torres, R.,
Taliun, S.A.G., Corvelo, A., Gogarten, S.M., Kang, H.M., et al. (2021).
Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program.
Nature *590*, 290–299.

39. Byrska-Bishop, M., Evani, U.S., Zhao, X., Basile, A.O., Abel, H.J., Regier,
A.A., Corvelo, A., Clarke, W.E., Musunuri, R., Nagulapalli, K., et al. (2022). High-
coverage whole-genome sequencing of the expanded 1000 Genomes Project
cohort including 602 trios. Cell *185*, 3426–3440.e19.

40. Sieberts, S.K., Perumal, T.M., Carrasquillo, M.M., Allen, M., Reddy, J.S.,
Hoffman, G.E., Dang, K.K., Calley, J., Ebert, P.J., Eddy, J., et al. (2020). Large
eQTL meta-analysis reveals differing patterns between cerebral cortical and
cerebellar brain regions. Scientific Data *7*, 340.

41. Fromer, M., Roussos, P., Sieberts, S.K., Johnson, J.S., Kavanagh, D.H.,
Perumal, T.M., Ruderfer, D.M., Oh, E.C., Topol, A., Shah, H.R., et al. (2016).
Gene expression elucidates functional impact of polygenic risk for schizophrenia.
Nature Neuroscience *19*, 1442–1453.

42. Ng, B., White, C.C., Klein, H.-U., Sieberts, S.K., McCabe, C., Patrick, E., Xu,
J., Yu, L., Gaiteri, C., Bennett, D.A., et al. (2017). An xQTL map integrates the
genetic architecture of the human brain's transcriptome and epigenome. Nature
Neuroscience *20*, 1418–1426.

43. Shang, L., Smith, J.A., Zhao, W., Kho, M., Turner, S.T., Mosley, T.H., Kardia,
S.L.R., and Zhou, X. (2020). Genetic Architecture of Gene Expression in
European and African Americans: An eQTL Mapping Study in GENOA. The
American Journal of Human Genetics *106*, 496–512.

44. Mostafavi, H., Spence, J.P., Naqvi, S., and Pritchard, J.K. (2023). Systematic
differences in discovery of genetic effects on gene expression and complex traits.
Nature Genetics *55*, 1866–1875.

45. Wojcik, G.L., Graff, M., Nishimura, K.K., Tao, R., Haessler, J., Gignoux, C.R.,
Highland, H.M., Patel, Y.M., Sorokin, E.P., Avery, C.L., et al. (2019). Genetic

analyses of diverse populations improves discovery for complex traits. Nature
570, 514–518.

46. Sun, Y., Zhou, D., Rahman, M.R., Zhu, J., Ghoneim, D., Cox, N.J., Beach,
T.G., Wu, C., Gamazon, E.R., and Wu, L. (2021). A transcriptome-wide
association study identifies novel blood-based gene biomarker candidates for
Alzheimer's disease risk. Human Molecular Genetics *31*, 289–299.

47. Mogil, L.S., Andaleon, A., Badalamenti, A., Dickinson, S.P., Guo, X., Rotter,
J.I., Johnson, W.C., Im, H.K., Liu, Y., and Wheeler, H.E. (2018). Genetic
architecture of gene expression traits across diverse populations. PLoS Genetics
14, e1007586.

48. Chen, Y., Liu, S., Ren, Z., Wang, F., Liang, Q., Jiang, Y., Dai, R., Duan, F.,
Han, C., Ning, Z., et al. (2024). Cross-ancestry analysis of brain QTLs enhances
interpretation of schizophrenia genome-wide association studies. American
Journal of Human Genetics *111*, 2444–2457.

49. Chen, M.-H., Raffield, L.M., Mousas, A., Sakaue, S., Huffman, J.E., Moscati,
A., Trivedi, B., Jiang, T., Akbari, P., Vuckovic, D., et al. (2020). Trans-ethnic and
ancestry-specific blood-cell genetics in 746,667 individuals from 5 global
populations. Cell *182*, 1198–1213.e14.

50. Zhu, J., Liu, X., Yin, H., Gao, Y., and Yu, H. (2021). Convergent lines of
evidence support BIN1 as a risk gene of Alzheimer's disease. Human Genomics
15, 9.

51. Greenwood, A.K., Gockley, J., Daily, K., Aluthgamage, D., Leanza, Z.,
Sieberts, S.K., Woo, K.H., Simon, S., Gendel, A., Thyer, T., et al. (2020). Agora:
An open platform for exploration of Alzheimer's disease evidence:
Genetics/omics and systems biology. Alzheimer's & Dementia *16*, e046129.

52. Joshi, G., Chi, Y., Huang, Z., and Wang, Y. (2014). Aβ-induced Golgi
fragmentation in Alzheimer's disease enhances Aβ production. Proceedings of
the National Academy of Sciences of the United States of America *111*, E1230–
E1239.

53. Joshi, G., Bekier, M.E., and Wang, Y. (2015). Golgi fragmentation in
Alzheimer's disease. Frontiers in Neuroscience *9*, 340.

54. Gerring, Z.F., Lupton, M.K., Edey, D., Gamazon, E.R., and Derks, E.M.
(2020). An analysis of genetically regulated gene expression across multiple
tissues implicates novel gene candidates in Alzheimer's disease. Alzheimer's
Research & Therapy *12*, 43.

55. Gamazon, E.R., Segrè, A.V., van de Bunt, M., Wen, X., Xi, H.S.,
Hormozdiari, F., Ongen, H., Konkashbaev, A., Derks, E.M., Aguet, F., et al.
(2018). Using an atlas of gene regulation across 44 human tissues to inform
complex disease- and trait-associated variation. Nature Genetics *50*, 956–967.

56. Huseby, C.J., Delvaux, E., Brokaw, D.L., and Coleman, P.D. (2023). Blood
RNA transcripts reveal similar and differential alterations in fundamental cellular
processes in Alzheimer's disease and other neurodegenerative diseases.
Alzheimer's & Dementia *19*, 2618–2632.

57. Zhong, H., Zhou, X., Uhm, H., Jiang, Y., Cao, H., Chen, Y., Mak, T.T.W., Lo,
R.M.N., Wong, B.W.Y., Cheng, E.Y.L., et al. (2024). Using blood transcriptome
analysis for Alzheimer's disease diagnosis and patient stratification. Alzheimer's
& Dementia *20*, 2469.