Phospho-eIF2Bε links stress response-driven translation to amyloidogenesis in Alzheimer's disease

Hugo Fanlo-Ucar,[1] Clara Planas-Clopes,[1] Oriol Bagudanch,[1] Carlos Pardo-Pastor,[1] Scarlett Troncoso-Cotal,[2] Alejandra R. Alvarez,[2] Baldo Oliva[3] and Francisco J. Muñoz[1]

Abstract

A central aetiopathological event in Alzheimer's disease is the accumulation of amyloid β-peptide (Aβ) derived from the amyloidogenic processing of the amyloid precursor protein, a pathway initiated by BACE1. Chronic activation of the Integrated Stress Response is linked to Alzheimer's disease, typically through eIF2α phosphorylation, which selectively enhances translation of key stress-responsive mRNAs like ATF4 and BACE1. We investigated a novel regulatory mechanism mediated by Glycogen Synthase Kinase-3β (GSK-3β) a hyperactive kinase in Alzheimer's disease, on the translational factor eIF2B, the guanine nucleotide exchange factor (GEF) for eIF2. We treated human neuroblastoma cells (SH-SY5Y) with a GSK-3β inhibitor (NP031112) or overexpressed a constitutively active GSK-3β mutant (GSK-3β-S9A) on HEK cells to modulate GSK-3β activity. GSK-3β inhibition significantly reduced ATF4 and BACE1 protein levels in SH-SY5Y cells in a dose-dependent manner as we tested by western blot ($P<0.001$), independent of eIF2α phosphorylation. This effect was translational, as the inhibitor still reduced BACE1 levels when transcription was blocked. Crucially, NP031112 reversed stress-induced eIF2Bε phosphorylation at Ser540 ($P<0.001$). Furthermore, H_2O_2-induced $A\beta_{1-42}$ secretion by SH-SY5Y cells was significantly reversed by GSK-3β inhibition ($P<0.001$). Luciferase reporter assays using the BACE1 5' untranslated region (5' UTR) confirmed that both eIF2Bε silencing and a phosphomimetic mutant eIF2Bε-S540E increased BACE1 5' UTR-driven translation ($P<0.001$), demonstrating that reduced functional eIF2B enhances BACE1 translation. Finally, immunohistofluorescence and western blot analysis of human AD hippocampi confirmed the translational link, showing a significant increase in pS540-eIF2Bε, ATF4, and BACE1 levels in AD patients compared to non-demented controls ($P<0.001$ for all three). Our findings establish a novel GSK-3β-eIF2Bε-BACE1 axis that links stress response dysregulation to amyloidogenesis, independent of canonical eIF2α phosphorylation. The phosphorylation of eIF2Bε functionally mimics the effects of eIF2α phosphorylation by reducing the available functional eIF2B pool,

thereby reprogramming translation to favour the production of stress proteins like ATF4 and BACE1. Based on our results, we proposed that eIF2Bε phosphorylation is a key etiological mechanism in Alzheimer's disease.

Author affiliations:

1 Laboratory of Molecular Physiology, Department of Medicine and Life Sciences, Faculty of Medicine and Life Sciences, Universitat Pompeu Fabra, 08003 Barcelona, Spain

2 Cell Signaling Laboratory, Biological Sciences Faculty, Millennium Institute on Immunology and Immunotherapy, Pontificia Universidad Católica de Chile, 8331150 Santiago de Chile, Chile

3 Laboratory of Structural Bioinformatics (GRIB), Department of Medicine and Life Sciences, Faculty of Medicine and Life Sciences, Universitat Pompeu Fabra, 08003 Barcelona, Spain

Correspondence to: Francisco J. Muñoz

MELIS-Universitat Pompeu Fabra

Carrer del Doctor Aiguader, 88, 08003 Barcelona

E-mail: paco.munoz@upf.edu

Keywords: Amyloid β-peptide; BACE1; ATF4; eIF2B; GSK-3β; eIF2α.

Running title: eIF2Bε phosphorylation links stress and amyloid

Introduction

A hallmark of Alzheimer's disease is the accumulation of misfolded amyloid β-peptide (Aβ) peptides, produced through amyloidogenic processing of the amyloid precursor protein (APP). In this pathway, BACE1 initiates APP cleavage,[1] followed by γ-

secretase,[2,3] which determines Aβ length. These peptides aggregate into toxic oligomers and fibrils, impairing synaptic function and neuronal viability.[4,5] However, how amyloidogenic processing becomes activated with age, the major Alzheimer's disease risk factor, remains unclear.

In Alzheimer's disease and other neurodegenerative diseases, chronic integrated stress response (ISR) activation disrupts proteostasis, synaptic function and survival.[6–8] Dysregulation of ISR components, particularly eIF2α phosphorylation and ATF4-mediated transcription, is linked to protein misfolding,[9,10] oxidative stress,[11] and neuroinflammation.[12] ATF4 regulates genes controlling adaptation, metabolism and apoptosis,[13] but under prolonged stress it promotes neuronal death.[14] The key regulatory event is eIF2α phosphorylation, which reduces general translation but enhances synthesis of uORF-containing mRNAs,[15] such as ATF4 and BACE1.[16–20] By lowering ternary complex availability, phosphorylated eIF2α slows scanning and reinitiation, increasing translation of ATF4 and BACE1, the latter contributing to Aβ overproduction.[13,17]

eIF2B, the GEF of eIF2, is central in this process.[21] Normally, it replenishes eIF2-GTP to sustain translation. Under stress, phosphorylated eIF2α binds eIF2B with high affinity, sequestering it in an inactive state and reducing protein synthesis.[13] This allows global suppression while maintaining selective translation of stress-response proteins. eIF2B activity is further regulated by GSK-3β,[22,23] a kinase hyperactive in Alzheimer's disease due to Aβ accumulation.[24,25] GSK-3β promotes tau hyperphosphorylation, neuronal dysfunction and possibly ISR modulation. It phosphorylates the ε subunit of eIF2B at Ser540, reducing GEF activity by ~50%.[22,23] Although GSK-3β has been linked to BACE1 transcriptional regulation via NF-κB,[26,27] its role in ISR-driven translational control of ATF4 and BACE1 remains unknown.

In this work we hypothesized that eIF2Bε phosphorylation significantly decreases active eIF2B, enhancing ATF4 and BACE1 expression under stress. We aim to unveil a novel translational regulatory mechanism contributing to neurodegeneration in Alzheimer's disease.

Materials and methods

Reagents

GSK-3β inhibitor (NP031112) was purchased from SelleckChem (S2823). DRB was purchased from Sigma (D1916-50MG). Hydrogen peroxide solution was purchased from Sigma (216763-500ML-M). MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) was purchased from Sigma (M5655).

Cell lines

Human neuroblastoma cells (SH-SY5Y) were grown with Ham's F12 medium + GlutaMAX (Gibco) supplemented with 15% FBS (Biowest) and 1% penicillin/streptomycin (Gibco). Human embryonic kidney 293 cells (HEK-293) were grown with DMEM medium (Gibco) supplemented with 10% foetal bovine serum and 1% penicillin/streptomycin. Cells were incubated at 37°C in a humidified atmosphere of 5% CO_2.

SH-SY5Y were plated on 96-well plates (15.000 cells/well) for MTT viability assays, on 6-well plates (400.000 cells/well) for WB studies and on 90 mm Petri dishes (1.500.000 cells/plate) for the ELISA assay. HEK-293 cells were seeded 96-well plates (16.000 cells/well) for luciferase assays and in 6-well plates (300.000 cells/well) for transfection and WB studies.

Human hippocampal samples

Human hippocampal samples were supplied by the Neurological Tissue Bank of the Biobank Hospital Clínic-IDIBAPS, Barcelona, Spain. The procedure was carried out according to the rules of the Helsinki Declaration and to the Ethics Committee of the Institut Municipal d'Investigacions Mèdiques-Universitat Pompeu Fabra (EC-IMIM-UPF). Samples were obtained from six non-demented controls and six Alzheimer's disease patients.

MTT viability assay

SH-SY5Y cells seeded in a 96-well plate were treated with vehicle (DMSO) and GSK-3β inhibitor at rising concentrations both in FBS-free Ham's F12 medium for 24 h. Then, cells were incubated with 0.5 mg/mL MTT solution for 2 h. Formazan crystals were

solubilized in 100 µL DMSO and cell viability was determined by absorbance at 595 nm by VICTOR Nivo Multimade Plate Reader (PerkinElmer).

Immunoblotting

SH-SY5Y cells in 6-well plates kept/treated with FBS-containing Ham's F12, FBS-free Ham's F12 or drugs (in FBS-free Ham's F12) for 24 h; HEK-293 cells in 6-well plates transfected with 2.5 µg of GSK-3β-S9A/GSK-3β-WT constructs for 24 h; and hippocampal samples from control and Alzheimer's disease patients, were lysed on ice with 50 µL of lysis buffer (150 mM Sodium Chloride, 1.0% Triton X-100, 0.5% Sodium Deoxycholate, 0.1% sodium dodecyl sulphate (SDS), 50 mM Tris, pH 8.0) supplemented with phosphatase and protease inhibitors tablets. Samples were centrifuged at 15,000 g for 10 min at 4ºC and then loaded into 10% acrylamide protein gels. Next, proteins were transferred onto 0.2 µm pore nitrocellulose membranes. Membranes were blocked for 1 h at room temperature (RT) with either 5% bovine serum albumin (BSA) for phosphoproteins or 5% skimmed milk for total proteins in Tween 20-Tris buffer solution (TTBS: 100 mM Tris-HCl, 150 mM NaCl, pH 7.5 plus 0.1% Tween-20). Then, membranes were incubated overnight (o.n.) at 4ºC with anti-BACE1 (1:1,000; ThermoFisher, PA1-757), anti-ATF4 (1:1,000; Cell Signaling, 11815S), anti-pS9-GSK-3β (1:1,000; Cell Signaling, 9323S), anti-GSK-3β (1:1,000; Cell Signaling, 9832), anti-pS51-eIF2α (1:1,000; Cell Signaling, 3597S), anti-eIF2α (1:1,000; Cell Signaling, 2103), anti-pS540-eIF2Bε (1:1,000; Invitrogen, 44-530G), anti-eIF2Bε (1:1,000; Cell Signaling, 3595S) or anti-GAPDH (1:2,000; Abcam, ab8245) antibodies (Abs). Horseradish peroxidase-conjugated donkey anti-rabbit and anti-mouse Abs (GE Healthcare) were used as secondary Abs at 1:2,000 for 1 h at RT. Primary and secondary Abs were diluted in blocking solution. Bands were visualized with Clarity Western ECL Substrate (BioRad) and analysed with ImageJ software.

Human Aβ$_{1-42}$ ELISA assay

Approximately 5 mL of supernatants were collected from SH-SY5Y cells cultured in 90 mm dishes and treated with vehicle, 10 µM NP031112, 10 µM H_2O_2 or both. These samples were centrifuged at 1200 rpm for 5 min to remove any floating cells, and concentrated 60x using the AMICON 3k filters (Sigma). Assay was performed with the

commercial Ultrasensitive Amyloid beta 42 Human ELISA Kit from Invitrogen following their protocol. Final Aβ concentrations were relativized to total protein concentration.

Structural modelling of eIF2-eIF2B

Structural models of eIF2 in complex with selected eIF2B subunits were generated using AlphaFold3. Each prediction included the eIF2 subunits α, β and γ, and the γ and ε subunits of eIF2B; the remaining eIF2B subunits were excluded from the modelling. The complete eIF2B structure was reconstructed by superimposing the modelled components onto the eIF2B structure from PDB entry 6K71. A GTP molecule was included in AlphaFold3 modelling which position it in the nucleotide-binding pocket of eIF2γ. To mimic regulatory phosphorylation, serine residues S525, S535 and S540 in eIF2Bε were substituted with glutamate prior to model generation (thus, two models were obtained, one phosphorylated and another unphosphorylated).

To assess the potential interaction between tRNA and the modelled eIF2-eIF2B complex, a tRNA molecule was incorporated by aligning the modelled eIF2 structure with the eIF2-tRNA complex from PDB entry 3JAP using the Matchmaker tool in ChimeraX. Conformational states of eIF2β and its interactions with GTP, the HEAT domain of eIF2Bε, and tRNA were analysed in ChimeraX. Electrostatic interactions were inferred based on the spatial proximity of acidic and basic side chains.

Cloning of BACE1 5′-untranslated region

Total RNA was extracted from SH-SY5Y cells, and one-step RT-PCR was carried out using a One-Step PCR kit (Qiagen) with primers designed to amplify *BACE1* 5′ UTR: FW 5′- TGGTACCACAAGTCTTTCCGCCTCCCCA-3′, RV 5′-GAAGCTTGGTGGG CCCCGGCCTTC-3′. The PCR product, a single band matching the molecular weight of *BACE1* 5′ UTR (~460 nt), was isolated and purified from an agarose gel using the Illustra™ GFX™ PCR DNA and Gel Band Purification kit (GE Healthcare). The 5′ UTR DNA fragment was then inserted into a modified pGL4.10-*luc2* vector from Promega containing the SV40 promoter, between Kpn1 and HindIII restriction enzyme sites.

Transient DNA transfection of HEK-293 cells

For the overexpression studies, HEK-293 cells seeded in 6-well plates and transfected with 2.5 µg of either pcDNA3 or GSK-3β-S9A/GSK-3β-WT constructs for 24 h using the

Lipofectamine 3000 transfection reagent (ThermoFisher). GSK-3β constructs were gently provided by Félix Hernández Pérez (Universidad Autónoma de Madrid, Madrid).

For the luciferase assays, HEK-293 cells were seeded in 96-well plates at a density of 12,000 cells per well and grown for 24 h with DMEM plus 10% FBS and at the same time transfected with either siCtrl or sieIF2Bε (100 pmol) using lipofectamine RNAiMAX transfection reagent (ThermoFisher). After 24 h, a total of 100 ng of DNA were transfected into each well, adjusting to the following conditions: 33 ng of either pGL4.10-*luc2* or pGL4.10-5' UTR·*BACE1*, plus 33 ng of pcDNA3, eIF2Bε-WT or eIF2Bε-S540E, plus 33 ng of Renilla luciferase. Cells were transfected using Lipofectamine 3000 transfection reagent (ThermoFisher) for 24 h.

Luciferase assay

Luciferase and Renilla activities of previously transfected HEK-293 cells were measured by using the Dual-Glo™ Luciferase Assay System (Promega) following the manufacturer instructions, and luminescence was read using a luminometer (VICTOR Nivo, PerkinElmer).

Immunohistofluorescence assays

Human hippocampal sections were treated with 0.15 M glycine and 10 mg/mL $NaBH_4$ in PBS for 30 min. Then they were washed in deionized water, permeabilized with 0.3% Triton X-100 in PBS for 1 h at 4ºC, and then blocked with 5% FBS + 1% BSA diluted in 0.1% Triton X-100 in PBS for 2 h. The following step was incubation o.n. at 4 ºC with anti-pS9-GSK-3β (1:200; Cell Signaling, 9323S) or anti-pY216-GSK-3β (1:200; Abcam, ab75745), and anti-GSK-3β (1:250; Cell Signaling, 9832) Abs. Samples were incubated the next day with Alexa Fluor 488 donkey anti-rabbit Ab (1:2000; Invitrogen, A32790), Alexa Fluor 555 donkey anti-mouse Ab (1:2000; Invitrogen, A32773) and TO-PRO-3 (1:1000) for 1 h at RT. They were then treated for 10 min with 0.3% Sudan Black (Sigma) in ethanol that was previously mixed o.n. at 4ºC, centrifuged at 3000 g for 20 min and filtered using common filter paper. Sections were washed with deionized water and cover slips were mounted using Fluoromont G. Digital images were taken with a Leica TCS SP5 confocal microscope and analysed with ImageJ software.

Statistical analyses

Data are expressed as mean ± SD of n experiments, as indicated in the corresponding figures. All statistics were performed with GraphPad software (9.0.1). Statistical tests to assure normality of the data were performed first, including Anderson-Darling, D'Agostino-Pearson omnibus, Shapiro-Wilk and Kolmogorov-Smirnov. Then, statistical analyses were performed using two-tailed Student's t-test and one-way or two-way ANOVA, depending on the variables analysed, followed by either Dunnett's or Tukey's post hoc test, as appropriate for the specific experimental design and comparisons required.

Results

The ISR up-regulates ATF4 and BACE1 protein levels via GSK-3β activation

Serum deprivation is known to activate the ISR.[28] To evaluate this effect in SH-SY5Y cells, we examined the expression levels of ATF4, since it is the major ISR downstream effector, and BACE1, a key enzyme in Alzheimer's disease aetiology, after 24 h of serum starvation. A significant increase in the expression of both proteins was observed under serum-free conditions ($P < 0.001$) (Fig. 1A). Furthermore, GSK-3β, another key enzyme in Alzheimer's disease etiopathology, is tightly regulated by phosphorylation: it becomes enzymatically enhanced via autophosphorylation at Tyr216,[29,30] and is inhibited by phosphorylation at Ser9 by mainly Akt.[31] Interestingly, it has been reported that serum deprivation activates GSK-3β in cortical neurons [32]. Therefore, we investigated the role of GSK-3β in the upregulation of ATF4 and BACE1 under serum deprived conditions, using NP031112, a non-ATP competitive and irreversible GSK-3β inhibitor.[33] First, we assessed its cytotoxicity after 24 h of treatment and found that concentrations up to 10 μM were non-toxic (Fig. 1B), which were subsequently used for further analyses on SH-SY5Y cells. Western blot (WB) assays were performed to evaluate the protein levels of ATF4, BACE1 and pS9-GSK-3β in the presence of NP031112. As expected, we observed a dose-dependent reduction in ATF4 and BACE1 protein levels, reaching a maximum decrease of over 50% at the highest NP031112 concentration tested ($P < 0.001$) (Fig. 1C). Similar results were obtained with AR·A014418, another GSK-3β inhibitor (Supplementary Fig. 1). Moreover, pS9-GSK-3β levels increased proportionally to the

inhibitor concentration, becoming significant ($P < 0.001$) at 10 µM (Fig. 1C). This observation supports the established mechanism where GSK-3β sustains its active state through the inhibitory phosphorylation of I-2, a PP1 inhibitor, consequently allowing PP1 to dephosphorylate inactive pS9-GSK-3β.[34] Notably, NP031112 appears to interrupt this positive feedback loop by directly inhibiting GSK-3β, offering a potential explanation for the observed elevation in pS9-GSK-3β. This phenomenon is cohesive with the PI3K/Akt-independent increase in Ser9 phosphorylation observed in cells treated with LiCl.[34] However, at the lowest inhibitor concentrations tested (1 and 5 µM), the difference was not statistically significant, likely due to being insufficient to prevent PP1 activity (Fig. 1C).

To confirm the specific effect of GSK-3β on the results obtained in Fig. 1A and 1C, we transfected HEK-293 cells with a constitutively active non-phosphorylatable GSK-3β mutant, GSK-3β-S9A, encoding vector. In agreement with prior results, ATF4 and BACE1 levels were significantly increased ($P < 0.05$ and $P < 0.001$, respectively) in cells overexpressing the mutant kinase and kept in FBS-containing media (Fig. 1D). Cohesively, overexpressing the wild-type enzyme also resulted in higher levels of ATF4 and BACE1 ($P < 0.001$ and $P < 0.01$, respectively) (Supplementary Fig. 2); however, these levels were lower than those obtained with GSK-3β-S9A, as expected.

GSK-3β modulates BACE1 translation and promotes eIF2Bε phosphorylation

To determine whether the increase in BACE1 expression is due to transcriptional regulation, we used 5,6-dichloro-1-beta-D-ribofuranosylbenzimidazole (DRB), a general transcription inhibitor that blocks activating phosphorylation of the RNA polymerase II by inhibiting CDK9, and, therefore, transcript elongation.[35] Given the high toxicity associated with transcriptional inhibition,[36] and since DRB acts rapidly (within 5 min)[37] and does not induce cell death at 5 h, we shortened the treatment duration from 24 to 5 h, also because our results show that GSK-β inhibition decreases BACE1 at 3-5 h (Supplementary Fig. 3). As expected, WB results revealed that both NP031112 and DRB independently reduce BACE1 levels ($P < 0.001$) (Fig. 2A). Notably, a more pronounced reduction in BACE1 levels was observed when both treatments were combined ($P <$

0.001), suggesting that NP031112 still manages to modulate BACE1 expression even when transcription is blocked (Fig. 2A). This observation supports a transcription-independent effect of GSK-3β on BACE1 expression.

eIF2α is one of the key regulators of the ISR, and its phosphorylation at Serine 51 occurs when one of the four ISR kinases is activated.[13] Although GSK-3β has not been reported to be an eIF2α kinase, we aimed to determine whether GSK-3β inhibition might influence eIF2α phosphorylation, and therefore explain the typical ISR outcomes we have observed. WB analysis showed a significant increase in eIF2α phosphorylation under serum deprivation ($P < 0.05$). Importantly, GSK-3β inhibition did not revert this, proving that blocking GSK-3β does not affect eIF2α phosphorylation triggered by serum deprivation (Fig. 2B, left panel). Therefore, the increase in pS51-eIF2α observed after serum starvation is independent of GSK-3β overactivation, but rather to GCN2 activation, as GCN2 is the eIF2α kinase that responds to amino acid deprivation.[38] While no direct link between eIF2α and GSK-3β has been reported, previous studies have shown that GSK-3β phosphorylates eIF2B at Serine 540 of the ε subunit in humans, reducing its activity by approximately 50%.[22,23] After investigating whether the GSK-3β inhibitor affected eIF2Bε phosphorylation, we found that 10 μM NP031112 significantly reduced pS540-eIF2Bε levels elevated by serum deprivation ($P < 0.001$) (Fig. 2B, right panel). This indicates that the decrease in both ATF4 and BACE1 levels in human neuroblastoma cells observed upon treatment with the GSK-3β inhibitor (Fig. 1) are not caused by a reduction of the canonical eIF2α phosphorylation. Overexpression of the GSK-3β-S9A mutant in HEK-293 cells also increased eIF2Bε phosphorylation ($P < 0.05$), further confirming our findings (Fig. 2C).

Finally, we addressed the pathophysiological relevance of our hypothesis by measuring secreted Aβ$_{1-42}$ (Fig. 2D), the most amyloidogenic Aβ species,[39,40] in the supernatant of cultured SH-SY5Y cells. To induce Aβ$_{1-42}$ production, we treated human neuroblastoma cells with H_2O_2, a well-known GSK-3β activator.[41] In serum-free conditions, 10 μM NP031112 reduced Aβ$_{1-42}$ levels by approximately 20% ($P < 0.01$). Cells challenged with 10 μM H_2O_2 exhibited a 23% increase in Aβ$_{1-42}$ production compared to controls ($P < 0.01$), which was reversed by co-treatment with NP031112 ($P < 0.001$). These findings are consistent with the previous results, showing that less BACE1 implies less Aβ$_{1-42}$ production, and confirms that GSK-3β inhibition contributes to reduced Aβ generation, a result previously observed in mice models.[42,43]

While GSK-3β induces Aβ$_{1-42}$ generation by increasing BACE1 transcription,[27] indirectly enhancing γ-secretase activity,[44] and phosphorylating APP and increasing its processing ,[45,46] no modulation of BACE1 translation by GSK-3β has been established yet. Taken together, our results suggest a model in which GSK-3β modulates eIF2Bε phosphorylation and activates a non-canonical ISR, which at the same time upregulates BACE1 translation, in a situation analogous to that caused by eIF2alpha phosphorylation, which also culminates in a reduction of available eIF2B for protein translation (Fig. 2E). GSK-3β inhibition reduces eIF2Bε phosphorylation, relieving its suppressive effect on global translation initiation, thereby decreasing ATF4 and BACE1 synthesis. This mechanism supports a translational control axis of BACE1 expression and amyloid pathology involving GSK-3β-eIF2Bε-BACE1-Aβ$_{1-42}$.

An interaction model reveals that eIF2Bε phosphorylation disrupts eIF2 and eIF2B dynamics

To study possible effects on eIF2 and eIF2B dynamics mediated by eIF2Bε phosphorylation, we decided to model their interaction. The structural dynamics of the eIF2-eIF2B interaction are central to translational control via guanine nucleotide exchange mediated by the eIF2B GEF activity. The active structure of eIF2 bound to tRNA (PDB: 3JAP) reveals that GTP is primarily coordinated by the γ subunit, while the β subunit forms a closed conformation over the bound tRNA. The α subunit also participates in stabilizing the tRNA interaction, together forming a ternary complex that is competent for ribosomal recruitment and translation initiation (Supplementary Fig. 4A). Upon GTP hydrolysis, GDP remains tightly bound, likely due to the continued closed conformation of the β subunit, thereby preventing spontaneous nucleotide exchange (Fig. 3A, upper panel).

The GEF activity of eIF2B is required for GDP release. Structural insights from the eIF2B-eIF2 complex (PDB: 6K71) show that the α subunit of eIF2 adopts an open conformation, while the β subunit is unresolved, suggesting structural flexibility. Superposition of the eIF2-tRNA complex (3JAP) onto the eIF2B-eIF2 complex (6K71) indicates that tRNA could still be accommodated in the complex during nucleotide

exchange. The ε and γ subunits of eIF2B mediate the core interaction with eIF2 (Fig. 3A, upper panel).

Using AlphaFold3,[47] we modelled partial eIF2-eIF2B complexes comprising only the γ and ε subunits of eIF2B, and GDP. Two conformations emerged: (i) a closed β subunit of eIF2 retaining GDP within the active site, with the HEAT domain of the ε subunit distant; and (ii) an open conformation, with the β subunit displaced toward the γ subunit of eIF2B and in proximity to the HEAT domain of eIF2Bε, exposing the GDP-binding site. AlphaFold3 predictions suggest that the interaction favours the open β conformation, facilitating GDP release and subsequent GTP/tRNA binding (Supplementary Fig. 4B).

This supports a model in which eIF2B binds the GDP-bound form of eIF2 and induces opening of the β subunit to enable GDP release. Subsequent GTP loading and tRNA binding promote re-closure of the β subunit, stabilizing the ternary complex. This conformational change is likely mediated by electrostatic interactions, particularly involving positively charged residues on eIF2β (e.g., R261, R296, K215, K234, K259, K315), with R325 and R329 involved in GTP coordination. The α subunit then locks the tRNA in place, and the active ternary complex is released from eIF2B (Supplementary Fig. 4B).

To examine the impact of regulatory phosphorylation, we modelled mutations in the ε subunit of eIF2B that mimic phosphorylation (S525E, S535E and S539E). These mutations induced local disorder in the region connecting the HEAT domain to the core of the ε subunit, as reflected by low pLDDT scores. Despite the disorder, all AlphaFold models predicted an open conformation of the eIF2 β subunit, exposing the GDP-binding site. This conformation appears to result from electrostatic interactions between negatively charged glutamates from eIF2Bε (mimicking the effects of serine phosphorylation) and positively charged residues on eIF2β (notably K215, K259, R261, K324, R325 and R329) (Fig. 3A, lower panel).

Functionally, this phosphorylation mimic mutants disrupts the normal closing of the β subunit, thereby impairing GTP binding and subsequent tRNA capture. The negative charges introduced by phosphorylation compete with the tRNA own phosphate backbone for binding to key basic residues, preventing the formation of a stable ternary complex (Supplementary Fig. 4C). As a result, phosphorylated eIF2 remains in an inactive, GDP-

bound state and is unable to initiate translation effectively. These results explain a potential mechanism on how GSK-3β-mediated eIF2Bε phosphorylation achieves ISR activation.

Inhibition of eIF2Bε enhances BACE1 translation

Next, we studied the effect of eIF2Bε inhibition on BACE1 levels to directly demonstrate a link between eIF2Bε and BACE1 translation and to show the functional relationship between the two. We cloned *BACE1* 5' UTR into a pGL4.10-*luc2* vector containing the firefly luciferase gene under the control of the SV40 promoter (Fig. 3B). Moreover, to modulate eIF2Bε activity, we used the eIF2Bε-S540E mutant, where the serine at position 540 was replaced by a glutamic acid to structurally mimic the phosphorylation induced by GSK-3β, and a siRNA proven to repress eIF2Bε expression (Supplementary Fig. 5). Compared to the control vector, the insertion of the 5' UTR reduced the expression of luciferase, as expected ($P < 0.001$) (Fig. 3C). Silencing eIF2Bε, or overexpressing the eIF2Bε-S540E phosphomimetic mutant, rescued luciferase expression in 5' UTR-expressing cells ($P < 0.001$) (Fig. 3C). These results demonstrate that a reduction of functional eIF2Bε levels induces BACE1 translation, similar to what occurs when eIF2α is phosphorylated.

Phosphorylated eIF2Bε is increased in Alzheimer's disease hippocampal samples

To assess whether the molecular alterations observed in culture are relevant for Alzheimer's disease etiopathogenesis, we analysed post-mortem hippocampal samples from Alzheimer's disease patients and age-matched non-demented controls. Our study focused on evaluating GSK-3β activation status by analysing its inhibitory phosphorylation at Ser9 (pS9-GSK-3β) and activatory phosphorylation at Tyr216 (pY216-GSK-3β), as well as the expression of downstream targets related to the ISR.

Immunohistofluorescence studies performed on human hippocampal slices revealed a significantly lower pS9-GSK-3β/GSK-3β ratio in Alzheimer's disease hippocampi compared to controls ($P < 0.001$). In parallel, the pY216-GSK-3β/GSK-3β ratio was found to be significantly higher in Alzheimer's disease hippocampi ($P < 0.001$),

supporting increased GSK-3β activation. These results were confirmed by quantitative analysis of mean fluorescence intensities for both total and phosphorylated forms of GSK-3β (Fig. 4A).

To explore the downstream consequences of increased GSK-3β activity, we next analysed the protein levels of key ISR-related components by WB. Hippocampal lysates from Alzheimer's disease and non-demented individuals were used to quantify pS540-eIF2Bε, ATF4 and BACE1. Consistent with enhanced ISR signalling, we observed a significant increase in ATF4, BACE1 and pS540-eIF2Bε levels in Alzheimer's disease hippocampi compared to controls ($P < 0.01$, $P < 0.001$ and $P < 0.001$, respectively) (Fig. 4B). These data highlight the relevance of the phosphorylation of eIF2Bε and its potential role in promoting pathological translation of stress-related proteins in Alzheimer's disease.

Together, these findings provide strong evidence that dysregulation of GSK-3β activity and the ISR axis, particularly via eIF2Bε phosphorylation, contributes to the molecular pathology in Alzheimer's disease.

Discussion

Our study uncovers a previously unrecognized role of the ISR through phosphorylation of the eIF2Bε subunit, establishing a novel translational control mechanism that links cellular stress with amyloidogenic processing. Crucially, this pathway converges on BACE1, the rate-limiting enzyme for Aβ generation, placing it at the centre of Alzheimer's disease aetiology and progression.

We demonstrate that GSK-3β activity increases under stress-like conditions and is both necessary and sufficient to drive a sustained ISR-like state characterized by elevated ATF4 and BACE1. Importantly, this regulation occurs independently of eIF2α phosphorylation, a canonical ISR hallmark, and instead involves direct phosphorylation of eIF2Bε at Ser540, a modification that selectively impairs global translation while favouring translation of stress-responsive mRNAs, such as ATF4 and BACE1. This alternative mechanism mimics the downstream effects of ISR activation but bypasses traditional upstream sensors, suggesting that chronic GSK-3β activation can keep the cell in a "maladaptive translational state".

The pathological relevance of this mechanism is underscored by our findings in human Alzheimer's disease hippocampi, where we observed increased GSK-3β activity alongside elevated p-eIF2Bε, ATF4 and BACE1 levels in the hippocampus of Alzheimer's disease patients compared to controls. These data support the notion that chronic activation of GSK-3β in the aging or diseased brain perpetuates an active ISR state, driving BACE1 expression and consequently enhancing Aβ production, a central event in Alzheimer's disease pathogenesis.[48]

This mechanism has several implications for our understanding of Alzheimer's disease. First, it expands the role of GSK-3β beyond insulin resistance, Wnt signalling and tau phosphorylation,[25] establishing it as a direct modulator of the translational machinery in neurons. Second, it identifies eIF2Bε phosphorylation as a critical node linking stress, translation and amyloid pathology.

While some GSK-3β inhibitors have shown promise in preclinical Alzheimer's disease models,[49] translating them to clinical approval has proven challenging, as they are tested in Alzheimer's disease patients at advanced stages of the disease.[50,51] Our study, however, offers a crucial perspective by highlighting that the GSK-3β effect on eIF2Bε phosphorylation represent a fundamental mechanism directly impacting protein translation and Alzheimer's disease aetiology. Specifically, we demonstrate that eIF2Bε phosphorylation leads to increased BACE1 levels and, consequently, enhanced amyloid production. These findings open a promising path for further studies on the regulation of eIF2B activity.

While traditionally viewed as a downstream target inhibited by phosphorylated eIF2α during stress, our research demonstrates that eIF2B itself can act as a trigger of ISR-like responses when inhibited by GSK-3β-mediated phosphorylation. This occurs because the reduced availability of functional eIF2B alters translation dynamics, slowing ribosomal activity in a manner similar to the effects of eIF2α phosphorylation. These results are consistent with recently published evidence that shows that deletion of eIF2B alone promotes the activation of an alternative ISR, inducing a metabolic reprogramming to maintain energy homeostasis upon eIF2B attenuation.[52] These discoveries shift the perspective on eIF2B from being merely a passive target to an active component in stress sensing and response. Its relevance is further underscored by the fact that mutations in eIF2B subunits cause Vanishing White Matter (VWM) disease, a leukodystrophy marked

by white matter degeneration and high vulnerability to physiological stress, particularly in glial cells like astrocytes, which significantly shortens the life expectancy of patients, who have severe neurological symptoms.[53–55] These mutations often disrupt eIF2B guanine nucleotide exchange activity,[56] which resembles the phosphorylation effect we observed in our interaction model. On the other hand, the discovery of molecules such as ISRIB, which stabilizes eIF2B and restores its function even under stress,[57] has opened up new therapeutic possibilities not only for neurodegenerative diseases, but also for disorders directly linked to eIF2B dysfunction.

In summary, our findings suggest that a decline in functional eIF2Bε levels triggers ISR-associated protein translation, paralleling the mechanism activated by eIF2α phosphorylation. It establishes a novel GSK3β-eIF2Bε-BACE1 axis as a mechanistic link between stress response dysregulation of translation and Alzheimer's disease pathology. By elucidating how ISR reprograms translation in neurons to favour amyloidogenic pathways, we reveal a central pathogenic mechanism in Alzheimer's disease aetiology.

Data availability

The data that support the findings of this study are available on reasonable request from the corresponding author.

Acknowledgements

We thank Dr. Miguel A. Valverde (Universitat Pompeu Fabra, Barcelona) for his critical review of the manuscript.

Funding

This work was supported by the Spanish Ministry of Science and Innovation and Agencia Estatal de Investigación plus FEDER Funds through grants PID2023-149767OB-I00 (FJM) and PID2023-150068OB-I00 (BO) funded by MICIU/AEI/10.13039/501100011033 and by 'ERDF A way of making Europe'. This

work was also funded by MICIU/AEI/10.13039/501100011033 as 'Unidad de Excelencia María de Maeztu' CEX2024-001431-M.

Competing interests

The authors report no competing interests.

Supplementary material

Supplementary material is available at *XXXX* online.

References

1. Vassar R, Bennett BD, Babu-Khan S, et al. β-Secretase Cleavage of Alzheimer's Amyloid Precursor Protein by the Transmembrane Aspartic Protease BACE. *Science*. 1999;286(5440):735-741. doi:10.1126/science.286.5440.735

2. Yu G, Nishimura M, Arawaka S, et al. Nicastrin modulates presenilin-mediated notch/glp-1 signal transduction and βAPP processing. *Nature*. 2000;407(6800):48-54. doi:10.1038/35024009

3. Francis R, McGrath G, Zhang J, et al. aph-1 and pen-2 Are Required for Notch Pathway Signaling, γ-Secretase Cleavage of βAPP, and Presenilin Protein Accumulation. *Dev Cell*. 2002;3(1):85-97. doi:10.1016/S1534-5807(02)00189-2

4. Lambert MP, Barlow AK, Chromy BA, et al. Diffusible, nonfibrillar ligands derived from Aβ1–42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci*. 1998;95(11):6448-6453. doi:10.1073/pnas.95.11.6448

5. Dahlgren KN, Manelli AM, Stine WB, Baker LK, Krafft GA, LaDu MJ. Oligomeric and Fibrillar Species of Amyloid-β Peptides Differentially Affect Neuronal Viability. *J Biol Chem*. 2002;277(35):32046-32053. doi:10.1074/jbc.M201750200

6. Flury A, Aljayousi L, Park HJ, et al. A neurodegenerative cellular stress response linked to dark microglia and toxic lipid secretion. *Neuron*. 2025;113(4):554-571.e14. doi:10.1016/j.neuron.2024.11.018

7. Hu Z, Yu P, Zhang Y, et al. Inhibition of the ISR abrogates mGluR5-dependent long-term depression and spatial memory deficits in a rat model of Alzheimer's disease. *Transl Psychiatry*. 2022;12(1):96. doi:10.1038/s41398-022-01862-9

8. Costa-Mattioli M, Gobert D, Stern E, et al. eIF2α Phosphorylation Bidirectionally Regulates the Switch from Short- to Long-Term Synaptic Plasticity and Memory. *Cell*. 2007;129(1):195-206. doi:10.1016/j.cell.2007.01.050

9. Ho YS, Yang X, Lau JCF, et al. Endoplasmic Reticulum Stress Induces Tau Pathology and Forms a Vicious Cycle: Implication in Alzheimer's Disease Pathogenesis. *J Alzheimers Dis*. 2012;28(4):839-854. doi:10.3233/JAD-2011-111037

10. Radford H, Moreno JA, Verity N, Halliday M, Mallucci GR. PERK inhibition prevents tau-mediated neurodegeneration in a mouse model of frontotemporal dementia. *Acta Neuropathol (Berl)*. 2015;130(5):633-642. doi:10.1007/s00401-015-1487-z

11. Batenburg KL, Kasri NN, Heine VM, Scheper W. Intraneuronal tau aggregation induces the integrated stress response in astrocytes. Luo ZG, ed. *J Mol Cell Biol*. 2023;14(10):mjac071. doi:10.1093/jmcb/mjac071

12. Huang T chao, Luo L, Jiang S hai, et al. Targeting integrated stress response regulates microglial M1/M2 polarization and attenuates neuroinflammation following surgical brain injury in rat. *Cell Signal*. 2021;85:110048. doi:10.1016/j.cellsig.2021.110048

13. Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM. The integrated stress response. *EMBO Rep*. 2016;17(10):1374-1395. doi:10.15252/embr.201642195

14. Lockshin ER, Calakos N. The integrated stress response in brain diseases: A double-edged sword for proteostasis and synapses. *Curr Opin Neurobiol*. 2024;87:102886. doi:10.1016/j.conb.2024.102886

15. Herrera-Fernández V, Fanlo-Ucar H, Gohl P, et al. Predicting cellular adaptation proteins dependent on eIF2α regulation under stress conditions: Physiological and pathophysiological implications in neuronal function. *Comput Struct Biotechnol J*. 2025;27:3127-3136. doi:10.1016/j.csbj.2025.07.015

16. Lu PD, Harding HP, Ron D. Translation reinitiation at alternative open reading frames regulates gene expression in an integrated stress response. *J Cell Biol*. 2004;167(1):27-33. doi:10.1083/jcb.200408003

17. O'Connor T, Sadleir KR, Maus E, et al. Phosphorylation of the Translation Initiation Factor eIF2α Increases BACE1 Levels and Promotes Amyloidogenesis. *Neuron*. 2008;60(6):988-1009. doi:10.1016/j.neuron.2008.10.047

18. Ill-Raga G, Palomer E, Wozniak MA, et al. Activation of PKR causes amyloid ß-peptide accumulation via de-repression of BACE1 expression. *PloS One*. 2011;6(6):e21456. doi:10.1371/journal.pone.0021456

19. Ill-Raga G, Tajes M, Busquets-García A, et al. Physiological Control of Nitric Oxide in Neuronal BACE1 Translation by Heme-Regulated eIF2α Kinase HRI Induces Synaptogenesis. *Antioxid Redox Signal*. 2015;22(15):1295-1307. doi:10.1089/ars.2014.6080

20. Picón-Pagès P, Gutiérrez DA, Barranco-Almohalla A, et al. Amyloid Beta-Peptide Increases BACE1 Translation through the Phosphorylation of the Eukaryotic Initiation Factor-2α. *Oxid Med Cell Longev*. 2020;2020:2739459. doi:10.1155/2020/2739459

21. Pavitt GD, Ramaiah KVA, Kimball SR, Hinnebusch AG. eIF2 independently binds two distinct eIF2B subcomplexes that catalyze and regulate guanine-nucleotide exchange. *Genes Dev*. 1998;12(4):514-526. doi:10.1101/gad.12.4.514

22. Welsh GI, Miller CM, Loughlin AJ, Price NT. Regulation of eukaryotic initiation factor eIF2B: glycogen synthase kinase-3 phosphorylates a conserved serine which undergoes dephosphorylation in response to insulin. *FEBS Lett*. 1998;421(2):125-130. doi:10.1016/S0014-5793(97)01548-2

23. Pap M, Cooper GM. Role of Translation Initiation Factor 2B in Control of Cell Survival by the Phosphatidylinositol 3-Kinase/Akt/Glycogen Synthase Kinase 3β Signaling Pathway. *Mol Cell Biol*. 2002;22(2):578-586. doi:10.1128/MCB.22.2.578-586.2002

24. Leroy K, Yilmaz Z, Brion J -P. Increased level of active GSK-3β in Alzheimer's disease and accumulation in argyrophilic grains and in neurones at different stages of neurofibrillary degeneration. *Neuropathol Appl Neurobiol*. 2007;33(1):43-55. doi:10.1111/j.1365-2990.2006.00795.x

25. Lauretti E, Dincer O, Praticò D. Glycogen synthase kinase-3 signaling in Alzheimer's disease. *Biochim Biophys Acta BBA - Mol Cell Res*. 2020;1867(5):118664. doi:10.1016/j.bbamcr.2020.118664

26. Chen CH, Zhou W, Liu S, et al. Increased NF-κB signalling up-regulates BACE1 expression and its therapeutic potential in Alzheimer's disease. *Int J Neuropsychopharmacol*. 2012;15(01):77-90. doi:10.1017/S1461145711000149

27. Ly PTT, Wu Y, Zou H, et al. Inhibition of GSK3β-mediated BACE1 expression reduces Alzheimer-associated phenotypes. *J Clin Invest*. 2013;123(1):224-235. doi:10.1172/JCI64516

28. Cerqua M, Foiani M, Boccaccio C, Comoglio PM, Altintas DM. The integrated stress response drives MET oncogene overexpression in cancers. *EMBO J*. 2025;44(4):1107-1130. doi:10.1038/s44318-024-00338-4

29. Hughes K, Nikolakaki E, Plyte SE, Totty NF, Woodgett JR. Modulation of the glycogen synthase kinase-3 family by tyrosine phosphorylation. *EMBO J*. 1993;12(2):803-808. doi:10.1002/j.1460-2075.1993.tb05715.x

30. Cole A, Frame S, Cohen P. Further evidence that the tyrosine phosphorylation of glycogen synthase kinase-3 (GSK3) in mammalian cells is an autophosphorylation event. *Biochem J*. 2004;377(1):249-255. doi:10.1042/bj20031259

31. Moule SK, Welsh GI, Edgell NJ, Foulstone EJ, Proud CG, Denton RM. Regulation of Protein Kinase B and Glycogen Synthase Kinase-3 by Insulin and β-Adrenergic Agonists in Rat Epididymal Fat Cells. *J Biol Chem*. 1997;272(12):7713-7719. doi:10.1074/jbc.272.12.7713

32. Hetman M, Cavanaugh JE, Kimelman D, Xia Z. Role of Glycogen Synthase Kinase-3β in Neuronal Apoptosis Induced by Trophic Withdrawal. *J Neurosci*. 2000;20(7):2567-2574. doi:10.1523/JNEUROSCI.20-07-02567.2000

33. Luna-Medina R, Cortes-Canteli M, Sanchez-Galiano S, et al. NP031112, a Thiadiazolidinone Compound, Prevents Inflammation and Neurodegeneration under Excitotoxic Conditions: Potential Therapeutic Role in Brain Disorders. *J Neurosci*. 2007;27(21):5766-5776. doi:10.1523/jneurosci.1004-07.2007

34. Zhang F, Phiel CJ, Spece L, Gurvich N, Klein PS. Inhibitory Phosphorylation of Glycogen Synthase Kinase-3 (GSK-3) in Response to Lithium. *J Biol Chem*. 2003;278(35):33067-33077. doi:10.1074/jbc.M212635200

35. Kuo YH, Lai TC, Chang CH, Hsieh HC, Yang FM, Hu MC. 5,6-Dichloro-1-β-D-ribofuranosylbenzimidazole (DRB) induces apoptosis in breast cancer cells through inhibiting of Mcl-1 expression. *Sci Rep*. 2023;13(1):12621. doi:10.1038/s41598-023-39340-x

36. Eciroglu-Sarban H, Altin-Celik P, Kelicen-Ugur P, Donmez-Altuntas H. Neuroprotective Effects of Hesperidin and CK2 Inhibitor DRB on Aβ1-42-Induced Neurotoxicity in Differentiated SH-SY5Y Cells. *Mol Neurobiol*. Published online May 30, 2025. doi:10.1007/s12035-025-05082-2

37. Clement JQ, Wilkinson MF. Rapid induction of nuclear transcripts and inhibition of intron decay in response to the polymerase II inhibitor DRB. *J Mol Biol*. 2000;299(5):1179-1191. doi:10.1006/jmbi.2000.3745

38. Misra J, Carlson KR, Spandau DF, Wek RC. Multiple mechanisms activate GCN2 eIF2 kinase in response to diverse stress conditions. *Nucleic Acids Res*. 2024;52(4):1830-1846. doi:10.1093/nar/gkae006

39. Vandersteen A, Broersen K, Schymkowitz J, Rousseau F. Amyloid Beta-peptide Length Strongly Affects Aggregation Kinetics. *Alzheimers Dement*. 2009;5(4S_Part_17). doi:10.1016/j.jalz.2009.07.071

40. Gu L, Guo Z. Alzheimer's Aβ42 and Aβ40 peptides form interlaced amyloid fibrils. *J Neurochem*. 2013;126(3):305-311. doi:10.1111/jnc.12202

41. Quintanilla RA, Muñoz FJ, Metcalfe MJ, et al. Trolox and 17β-Estradiol Protect against Amyloid β-Peptide Neurotoxicity by a Mechanism That Involves Modulation of the Wnt Signaling Pathway. *J Biol Chem*. 2005;280(12):11615-11625. doi:10.1074/jbc.M411936200

42. Avrahami L, Farfara D, Shaham-Kol M, Vassar R, Frenkel D, Eldar-Finkelman H. Inhibition of glycogen synthase kinase-3 ameliorates β-amyloid pathology and restores lysosomal acidification and mammalian target of rapamycin activity in the Alzheimer disease mouse model: in vivo and in vitro studies. *J Biol Chem*. 2013;288(2):1295-1306. doi:10.1074/jbc.M112.409250

43. Rockenstein E, Torrance M, Adame A, et al. Neuroprotective effects of regulators of the glycogen synthase kinase-3beta signaling pathway in a transgenic model of Alzheimer's disease are associated with reduced amyloid precursor protein

phosphorylation. *J Neurosci Off J Soc Neurosci*. 2007;27(8):1981-1991. doi:10.1523/JNEUROSCI.4321-06.2007

44. Uemura K, Kuzuya A, Shimozono Y, et al. GSK3β Activity Modifies the Localization and Function of Presenilin 1. *J Biol Chem*. 2007;282(21):15823-15832. doi:10.1074/jbc.M610708200

45. Lee MS, Kao SC, Lemere CA, et al. APP processing is regulated by cytoplasmic phosphorylation. *J Cell Biol*. 2003;163(1):83-95. doi:10.1083/jcb.200301115

46. Aplin AE, Gibb GM, Jacobsen JS, Gallo J, Anderton BH. In Vitro Phosphorylation of the Cytoplasmic Domain of the Amyloid Precursor Protein by Glycogen Synthase Kinase-3β. *J Neurochem*. 1996;67(2):699-707. doi:10.1046/j.1471-4159.1996.67020699.x

47. Abramson J, Adler J, Dunger J, et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*. 2024;630(8016):493-500. doi:10.1038/s41586-024-07487-w

48. Hampel H, Hardy J, Blennow K, et al. The Amyloid-β Pathway in Alzheimer's Disease. *Mol Psychiatry*. 2021;26(10):5481-5503. doi:10.1038/s41380-021-01249-0

49. Serenó L, Coma M, Rodríguez M, et al. A novel GSK-3β inhibitor reduces Alzheimer's pathology and rescues neuronal loss in vivo. *Neurobiol Dis*. 2009;35(3):359-367. doi:10.1016/j.nbd.2009.05.025

50. Arciniegas Ruiz SM, Eldar-Finkelman H. Glycogen Synthase Kinase-3 Inhibitors: Preclinical and Clinical Focus on CNS-A Decade Onward. *Front Mol Neurosci*. 2022;14:792364. doi:10.3389/fnmol.2021.792364

51. ARGO investigators, Lovestone S, Boada M, et al. A Phase II Trial of Tideglusib in Alzheimer's Disease. *J Alzheimers Dis*. 2015;45(1):75-88. doi:10.3233/JAD-141959

52. Chen CW, Papadopoli D, Szkop KJ, et al. Plasticity of the mammalian integrated stress response. *Nature*. 2025;641(8065):1319-1328. doi:10.1038/s41586-025-08794-6

53. Van Der Knaap MS, Leegwater PAJ, Könst AAM, et al. Mutations in each of the five subunits of translation initiation factor eIF2B can cause leukoencephalopathy with vanishing white matter. *Ann Neurol*. 2002;51(2):264-270. doi:10.1002/ana.10112

54. Dooves S, Bugiani M, Postma NL, et al. Astrocytes are central in the pathomechanisms of vanishing white matter. *J Clin Invest*. 2016;126(4):1512-1524. doi:10.1172/jci83908

55. Leferink PS, Dooves S, Hillen AEJ, et al. Astrocyte Subtype Vulnerability in Stem Cell Models of Vanishing White Matter. *Ann Neurol*. 2019;86(5):780-792. doi:10.1002/ana.25585

56. Fogli A, Schiffmann R, Hugendubler L, et al. Decreased guanine nucleotide exchange factor activity in eIF2B-mutated patients. *Eur J Hum Genet*. 2004;12(7):561-566. doi:10.1038/sj.ejhg.5201189

57. Wong YL, LeBon L, Edalji R, Lim HB, Sun C, Sidrauski C. The small molecule ISRIB rescues the stability and activity of Vanishing White Matter Disease eIF2B mutant complexes. *eLife*. 2018;7. doi:10.7554/elife.32733

58. Pettersen EF, Goddard TD, Huang CC, et al. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci Publ Protein Soc*. 2021;30(1):70-82. doi:10.1002/pro.3943

Figures

Figure 1 ISR increases ATF4 and BACE1 protein expression through GSK-3β. (**A**) Human neuroblastoma cells (SH-SY5Y cells) were grown for 24 h in the presence/absence of FBS. ATF4 and BACE1 expression was assayed by WB. Data are the mean ± SD of three independent experiments. *** $P < 0.001$ vs. control by Student's t-test. (**B**) SH-SY5Y cells were treated with increasing concentrations of the GSK-3β inhibitor NP031112 in FBS-free medium for 24 h and survival was assayed by MTT reduction. Data are the mean ± SD of three independent experiments. ** $P < 0.01$ and *** $P < 0.001$ vs. control by one-way ANOVA plus Dunnett as post hoc test. (**C**) SH-

SY5Y cells were treated with increasing concentrations of NP031112 in FBS-free medium for 24 h and the expression of ATF4, BACE1 and pS9-GSK-3β were assayed by WB. Data are the mean ± SD of five-seven independent experiments, respectively. * $P <$ 0.05, ** $P < 0.01$ and *** $P < 0.001$ vs. control by one-way ANOVA plus Dunnett as post hoc test. (**D**) HEK-293 cells were transfected with empty (pcDNA3) or GSK-3β-S9A-encoding plasmids and BACE1 and ATF4 protein levels were analysed by WB after 24 h. Data are the mean ± SD of four and three independent experiments, respectively. * $P < 0.05$ and *** $P < 0.001$ vs. control by Student's t-test.

Figure 2 BACE1 up-regulation depends on eIF2Bε phosphorylation. (**A**) SH-SY5Y cells were treated with either 10 μM NP031112, 50 μM DRB, or a combination of both in FBS-free medium for 5 h. The expression of BACE1 was assayed by WB. Data are the mean ± SD of six independent experiments. *** $P < 0.001$ vs. untreated controls and ## $P < 0.01$ vs. DRB plus vehicle by two way-ANOVA plus Tukey as post hoc test. (**B**) SH-SY5Y cells were cultured in FBS-containing medium, serum-starved (Ctrl), or treated with 10 μM NP031112 in FBS-free medium for 24 h and the phosphorylation of eIF2α and eIF2Bε was assayed by WB. Data are the mean ± SD of four independent experiments, respectively. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ and non-significant (n.s.) vs. control by one way-ANOVA plus Dunnett as post hoc test. (**C**) HEK-293 cells were transfected with pcDNA3 and GSK-3β-S9A plasmid and the level of pS540-eIF2Bε was analysed by WB 24 h after transfection. Data are the mean ± SD of three independent experiments. * $P < 0.05$ vs. control by Student's t-test. (**D**) SH-SY5Y cells were treated with 10 μM NP031112, 10 μM H_2O_2, or a combination of both in FBS-free medium for 24 h. Aβ$_{1-42}$ concentration in the supernatant was measured by ELISA. Data are the mean ± SD of seven independent experiments. ** $P < 0.01$ vs. control and ### $P < 0.001$ vs. H_2O_2 plus vehicle by two way-ANOVA plus Tukey as post hoc test. (**E**) Schematic representation of the hypothesis: Phosphorylation of the eIF2Bε subunit at Ser540 by GSK-3β mimics the effects of p-eIF2α, due to reduced levels of active eIF2B.

Figure 3 eIF2Bε modelling and mutations support its role on BACE1 translation.
(**A**) Ribbon representation of the structural model of eIF2 (subunits α, β, γ) in complex with eIF2B in the closed (up) and open (down) conformations using ChimeraX.[58] eIF2 subunits α and γ are shown in green; subunit β is shown in cyan in the upper panel and blue in the lower panel. eIF2B subunits γ and ε are shown in dark and light brown, respectively. The remaining eIF2B subunits are shown in beige. The HEAT domain and the GTP molecule are indicated, with the latter displayed in surface representation and coloured by heteroatom. In the upper panel, a tRNA molecule is included to indicate its ability to bind the complex. In the lower panel, the region spanning residues 525-540, enriched in negatively charged residues (including glutamate substitutions for phosphorylated serines), is highlighted in red with side-chain atoms shown. Side-chain atoms of selected positively charged residues (Arg and Lys) are indicated in both panels. (**B**) Schematic representation of the pGL4.10-*luc2* vectors used. (**C**) HEK-293 cells were transfected with vectors pGL4.10-*luc2* or pGL4.10-*luc2*-5' UTR-*BACE1*; pRL; eIF2Bε-WT or eIF2Bε-S540E; siCtrl or sieIF2Bε. Then, firefly and renilla luciferase activities were analysed with a Dual-Glo luciferase kit and luminescence was recorded with a luminometer. Data are the mean ± SD of 15-25 independent experiments. *** $P < 0.001$ vs. pGL4.10-*luc2*-5' UTR-*BACE1* or ### $P < 0.001$ vs. eIF2Bε-WT by one-way ANOVA plus Tukey as post hoc test.

Figure 4 BACE1 and eIF2Bε are increased in the Alzheimer's disease hippocampal samples. (**A**) Immunohistofluorescence studies of hippocampal slices of Alzheimer's disease and non-demented individuals were used to investigate GSK-3β activity. The panels on the right show enlarged views of the regions within the yellow-outlined squares in the images on the left panels. Scale bars represent 50 μm in both left and right panels. pS9-GSK-3β and pY216-GSK-3β fluorescence intensities were measured and relativized to the total GSK-3β to create ratios. Data are the mean ± SD of five independent

individuals per group. *** $P < 0.001$ vs. control by Student's t-test. Images were obtained using a Leica SP8 confocal microscope and analysed with ImageJ. (**B**) Hippocampal tissue of Alzheimer's disease and non-demented individuals was homogenized and levels of ATF4, BACE1 and pS540-eIF2Bε were analysed by WB. Data are the mean ± SD of six independent individuals per group. ** $P < 0.01$ and *** $P < 0.001$ vs. control by Student's t-test.

SUPPLEMENTARY MATERIALS

eIF2Bε phosphorylation links stress response-driven translation to amyloidogenesis in Alzheimer's disease

Supplementary Figure 1 AR·A014418 reduces ATF4 and BACE1 levels. (**A**) SH-SY5Y cells were treated for 24 h with increasing concentrations of AR·A014418. Then, a MTT assay was performed to analyse cell survival. Data are the mean ± SEM of 3 independent experiments, respectively. *** $P < 0.001$ vs. control by one-way ANOVA plus Dunnett as post hoc test. (**B**) SH-SY5Y cells were treated for 24 h with increasing subtoxic concentrations of AR·A014418. Then, WB were performed to analyse ATF4 and BACE1 protein levels. Data are the mean ± SEM of 3 independent experiments, respectively. *** $P < 0.001$, ** $P < 0.01$ and * $P < 0.05$ vs control by one-way ANOVA plus Dunnett as post hoc test.

Supplementary Figure 2 Overexpression of GSK-3β-WT upregulates stress-responsive signalling. HEK-293 cells were transfected with empty (pcDNA3) or GSK-3β-encoding plasmids and BACE1 and ATF4 protein levels were analysed by WB after 24 h. Data are the mean ± SD of four and three independent experiments, respectively. * $P < 0.05$ and *** $P < 0.001$ vs. control by Student's t-test.

Supplementary Figure 3 NP031112 and AR·A014418 reduce ATF4 and BACE1 levels. SH-SY5Y cells were treated for 0 (Ctrl + FBS and Ctrl), 0.5, 1, 3 or 5 h with 10 μM NP031112 (**A**) or 1 μM AR·A014418 (**B**). Then, a WB was performed to analyse BACE1 protein levels, and the bands were quantified.

A

eIF2α   eIF2γ

tRNA   eIF2β

GTP

B

eIF2γ

eIF2α   eIF2Bε

Closed
eIF2β (1)

Open
eIF2β (2)

eIF2Bγ

eIF2B

Closed   Open

+GTP, + tRNA

Rotation
(closing)

tRNA   R325   GTP
R296
K315   R329

C

Fragment negatively charged
(S525E, S535E, S540E)
eIF2Bε

Fragment positively charged
(K324, R325, R329, R261)
eIF2β

Supplementary Figure 4 Phosphorylation of eIF2Bε disrupts proper formation of the translation ternary complex. (**A**) Ribbon representation of the structural model of eIF2 (subunits α, β, γ) in complex with a GTP molecule and a tRNA (in pink) using ChimeraX. eIF2 subunits α and γ are shown in green; subunit β is shown in cyan in the upper panel and blue in the lower panel. (**B**) Binding of eIF2B to eIF2 predicts two possible β subunit conformations: 1) a closed conformation shown in cyan; and 2) an open conformation shown in dark blue. The electrostatic interaction between the positively charged residues R296, K315, R325, and R329 on eIF2β and the GTP and tRNA molecules is depicted. eIF2B γ subunit is shown in brown/maroon and ε subunit is shown in red/translucent red. The rest of eIF2B subunits are partially showed in beige. (**C**) Representation of the interaction between eIF2 and eIF2B containing the phosphomimetic glutamate residues in eIF2Bε. eIF2β is represented in dark blue, with positively charged fragment highlighted in cyan; eIF2Bε subunit is shown in translucent red, with negatively charged phosphomimetic residues highlighted in red.

Supplementary Figure 5 Silencing eIF2Bε reduces its mRNA and protein levels. (**A**) HEK-293 cells were transfected for 24 h with four different siRNAs targeting different

sequences of the *EIF2B5* mRNA. Then, a RT-qPCR was performed to study levels of *EIF2B5* mRNA. (**B**) A western blot was also performed to study the protein levels of eIF2Bε after silencing.