# scFPC-DE: Robust Differential Expression Analysis Along Single Cell Trajectories via Functional Principal Component Analysis

Ricardo J. López Candelaria[1], Yu Qian[2], Fang Chen[3], Mansun Law[3], Yun Zhang[4, *] and Xing Qiu[1,*]

## Affiliations

[1]Department of Biostatistics and Computational Biology, University of Rochester, Rochester, NY 14642, USA. [2]Department of Informatics, J. Craig Venter Institute, La Jolla, CA 92037, USA. [3]Department of Immunology and Microbiology, Scripps Research, La Jolla, CA 92037, USA. [4]Division of Intramural Research, National Library of Medicine, Bethesda, MD 20894, USA.

*To whom correspondence should be addressed.

## Abstract

**Motivation:** Identifying temporally differentially expressed genes (TDEGs) along pseudotime trajectories from single cell RNA sequencing (scRNA-seq) data helps characterize the cellular states that underlie the dynamic process of cellular development. However, existing tests based on generalized additive models (GAMs) suffer from increased false positive rates under zero inflation caused by high dropouts, a ubiquitous technical artifact of scRNA-seq data. Furthermore, by testing each gene independently, existing tests ignore the variance-covariance structure shared across genes along the trajectory, leading to suboptimal power and reduced interpretability.

**Results:** We present scFPC-DE, a trajectory-based differential expression analysis (TDEA) method based on functional data analysis (FDA). It models the gene expression as a function of pseudotime in the $L^2$ space and represents the covariance structure of these functions by eigenfunctions derived from functional principal component (FPC) analysis. This approach effectively captures informative gene expression patterns along the trajectory, while mitigating the influence of zero inflation in both simulation and real data analysis. In simulations, scFPC-DE exhibited superior control of type I error and achieved the highest ROC-AUC among competing methods. When applied to an scRNA-seq dataset of B cell subtypes, scFPC-DE uniquely identified TDEGs enriched for B cell differentiation pathways, outperforming existing methods in biological relevance. These results show that scFPC-DE effectively captures the shared gene expression variation and pseudo-temporal structure along the single cell trajectory for TDEG identification.

**Availability:** R package and code vignettes are publicly available at https://github.com/LopezRicardo1/scFPCDE.

**Contact:** xing_qiu@urmc.rochester.edu; yun.zhang@nih.gov.

37  **Supplementary information:** Supplementary data are available at *Bioinformatics* online

## Introduction

Single cell RNA sequencing (scRNA-seq) has revolutionized transcriptomics by enabling the

measurement of gene expression at the resolution of individual cells, capturing cellular

heterogeneity and revealing complex biological processes such as cell differentiation [1].

scRNA-seq experiments generate large-scale datasets composed of thousands of cells and

genes, posing substantial statistical and computational challenges for data processing and

analysis. Gene expression signals in single cells are much lower than those in bulk

transcriptomics. Robust statistical methods are required to effectively capture the characteristics

of cells and extract meaningful biological insights from scRNA-seq data.   An emerging, rapidly

growing area of scRNA-seq analysis is single cell trajectory inference, which aims to reconstruct

the continuous development of cells through dynamic biological processes such as cell

differentiation and cell activation [1]. This is typically achieved by estimating a latent variable

known as pseudotime using methods such as *Monocle3* [2], which orders cells along a trajectory

that reflects the biological states of the cells in the process. Trajectory-based differential

expression (DE) analysis aims to identify temporally differentially expressed genes (TDEGs)

along a cell trajectory inferred via pseudotime. Methods for trajectory-based DE analysis, such

as the "regression analysis" approach in the DE module in *Monocle3* [3] and *tradeSeq* [4], use

generalized additive models (GAMs) of smooth functions for fitting gene expression values along

the estimated pseudotime. There is also a "graph-autocorrelation analysis" approach for DE

analysis in *Monocle3* that uses Moran's I [5] to detect genes with expression patterns spatially

correlated with the reconstructed cell graph/trajectory. While this test is sensitive to localized or

non-monotonic patterns, it does not provide effect estimates or easy covariate adjustment

(including experimental designs). In addition, its graph reconstruction, gene-wise Moran's I

calculations, and Monte-Carlo resampling for $p$-value calculation scale very poorly with cell and

gene counts. PseudotimeDE [6] accounts for uncertainties in pseudotime estimation and applies

a permutation-based inference framework with the GAM (Gaussian or negative-binomial)

approach to produce recalibrated $p$-values. Because it corrects for pseudotime variability and

has outperformed alternatives such as *Monocle3*-DE and *tradeSeq* in recent benchmarks,

PseudotimeDE is widely regarded as the current state-of-the-art for TDEG detection.

Most of these modeling approaches rely on parametric assumptions, such as negative

binomial or zero-inflated negative binomial distributions for the scRNA-seq count data. However,

parametric approaches are vulnerable to deviations from the underlying distribution

assumptions. An alternative is to model the gene expression changes along pseudotime using

functional principal component analysis (FPCA), a well-known technique in functional data

analysis (FDA), [7, 8]. An FPCA-based F-statistic was introduced for DE testing in time-course

bulk RNA-seq studies [9], which outperformed alternative time-course methods, particularly

when the number of time points is sparse or irregular. FPCA modeling was also used for robust

gene set enrichment analysis (GSEA) in time-course bulk RNA-seq studies [10]. A major strength

of FPCA lies in its ability to learn a data-driven basis in the $L^2$ space (eigenfunctions of the

covariance operator), which provides a representation of the dominant temporal patterns of gene

expressions along time or pseudotime. This stands in contrast to approaches like GAM, which

typically rely on fixed, predefined basis functions. By the celebrated Karhunen–Loève theorem

[11-13], top $L$ eigenfunctions estimated by FPCA capture more of the total $L^2$ variance than any

other $L$-dimensional basis, making them the most efficient parsimonious low-rank representation

of temporal data. In the scRNA-seq context, where each cell provides a pseudo-temporal

observation, FPCA naturally captures the *shared temporal patterns* across genes and

pseudotime points (i.e., cells along the trajectory), overcoming the time point sparsity in the bulk

RNA-seq experiments that often involve a limited number of discrete sampling times.

Unlike bulk RNA-seq, scRNA-seq data are known for high zero inflation. Zero counts in

scRNA-seq data may arise from sampling zero, technical zero, and biological zero [14]. The

dropout technical artifact due to the low amount of mRNA in scRNA-seq experiments contributes

to excessive zero counts in scRNA-seq data [15]. High zero inflation presents a non-trivial

challenge for downstream analysis because many standard approaches assume continuous,

well-sampled observations. Consequently, single cell trajectory analysis requires methods that

remain robust when a large proportion of measurements are zeros.

Motivated by these considerations, we introduce scFPC-DE, a trajectory-based DE analysis

(TDEA) testing method grounded in the FDA hypothesis testing principles [16]. The traditional

functional F-test (FPC-F) summarizes trajectory variability by the ratio of between- to

within-trajectory variance. When zeros dominate a gene's profile, the denominator can be very

small, producing spuriously large F-statistic value driven by just a few nonzero points. In

contrast, scFPC-DE avoids this issue by using a distance-based test statistic ($L^2$ distance in the

subspace spanned by eigenfunctions) which is numerically stable even when expression profiles

contain excessive zeros. Secondly, scFPC-DE constructs its functional basis (eigenfunctions)

from a selected subset of trajectory-informative genes (TIGs): Well-expressed genes that exhibit

sufficient variation relevant to the pseudotime. As a result, our trajectory model reflects

biologically meaningful variation, leading to more robust inference and substantially tighter

false-positive control in zero-inflated scRNA-seq data. Our strategy addresses the limitations of

existing methods and provides a powerful framework for cell trajectory downstream analysis.

In our simulation studies designed to reflect the high zero inflation nature of scRNA-seq

data, scFPC-DE stringently controls the type I error at the nominal level, while competing

methods exhibited considerable inflation of type I error. scFPC-DE also achieved comparable or

higher power than existing approaches. Consequently, scFPC-DE attained the highest area

under the curve (AUC) in Receiver Operating Characteristic (ROC) analysis among the methods

tested.

We further evaluate method performance using a real scRNA-seq dataset of phenotypically

sorted B cell subtypes [17]. scFPC-DE and PseudotimeDE were applied to identify TDEGs

associated with the B cell differentiation trajectory of transitional, naïve, and classical memory B

cell subtypes. We apply *ReactomePA* (for pathway enrichment analyses) [18], and the

*Enrichr-KG* online tool (for phenotype/network queries) [19], to evaluate biological relevance of

the TDEGs identified by both methods. Compared with PseudotimeDE, the TDEGs identified by

scFPC-DE form a more concise list that exhibits greater biological relevance and stronger

enrichment for pathways related to B cell development and immune system processes. This

more stringent and focused DEG detection using scFPC-DE is consistent with the tight type I

error control observed in simulations, which supports more targeted and biologically interpretable

downstream analyses.

The remainder of this manuscript is organized as follows. Section 2 develops the core

statistical properties of the FPC model for trajectory gene expression and introduces the

distance-based test statistic in the FPC latent space. Section 3 presents results from

zero-inflated simulation studies and compares the performance of scFPC-DE with previously

established FPC-F and PseudotimeDE methods, using the latter as the current benchmark for

trajectory-based differential expression analysis (TDEA). This section also includes results from

the real B cell scRNA-seq dataset and enrichment analysis of the identified TDEGs. Section 4

concludes with a discussion of the contributions. The scFPC-DE R package is publicly available

at GitHub: https://github.com/LopezRicardo1/scFPCDE.

## Methods

Functional principal component analysis (FPCA) [7, 20] is an extension of standard PCA for

continuous functions. Here, we propose to model the gene expression changes along the

trajectory as a function of pseudotime and use FPCA to capture the main sources of variations

across the gene expression functions, providing a data-driven framework for modeling the gene

expression changes along the pseudotime trajectory.

**2.1 Modeling Cell Trajectories with FPCA**

Let $Y_i(t_j)$ denote the log-transformed, centered gene expression of gene $i$ at pseudotime point

$j$. Using FPCA, we can express them as

$$Y_i(t_j) = \sum_{k=1}^{L} \xi_{ik} \Phi_k(t_j) + \epsilon_i(t_j). \#(1)$$

141 Here $\Phi_k(t)$, $k = 1, \dots L$, are the top $L$ eigenfunctions that capture the dominant temporal

142 expression modes, and $\xi_{ik}$ are the corresponding functional principal component (FPC) scores.

143 The residual term $\epsilon_i(t_j)$ models random noise and accounts for the remaining variability. FPC

144 scores are obtained by projecting each gene trajectory onto the linear space spanned by

145 eigenfunctions learned non-parametrically satisfying

$$\int C(s,t)\Phi_k(s)ds = \lambda_k \Phi_k(t) \#(2)$$

where $C(s,t)$ and $\lambda_k$ denotes the pseudotime covariance and $k$-th eigenvalue, respectively

(see Supplementary Section 1.1). Genes with large FPC scores have larger pseudo-temporal

variations that align with the dynamic behavior represented by eigenfunctions, while small scores

indicate flat or noise-driven profiles. We performed FPCA using a standard procedure based on

roughness penalized B-spline [7, 21], implemented in the *fda* R package [21]

**2.2 Trajectory-based Differential Expression FPC Test**

Under the null hypothesis of no pseudo-temporal variation, FPC scores remain centered at zero

with their variability driven only by the measurement noise, yielding small magnitudes of FPC

scores relative to the alternative hypothesis. Formally, for each gene $i$, the null and alternative

hypotheses can be stated as:

$$H_{0i}: \xi_{ik} = 0, \forall k \text{ vs. } H_{1i}: \exists k \text{ such that } \xi_{ik} \neq 0.$$

156 We use the squared Euclidean norm of FPC scores as the test statistics:

$$D_i = \sum_{k=1}^{L} \xi_{ik}^2 . \#(3)$$

157   Mathematically, $D_i$ is the squared $L^2$-distance between the constant zero function and the $i$th

158   trajectory projected to the linear subspace spanned by the eigenfunctions. A larger $D_i$ indicates

159   stronger evidence of the pseudotime dynamics. To obtain a null distribution of $D_i$, we randomly

160   permute the pseudotime labels across cells and recompute $D_i$ from these permutations. The

161   permutation-based empirical p-value is defined as:

$$p_i = \frac{1}{B \cdot G} \sum_{b=1}^{B} \sum_{g=1}^{G} I\left\{ D_g^{(b)} \geq D_i \right\}, \#(4)$$

where $D_g^{(b)}$ is the test statistic for gene $g$ under the $b$-th permutation, and $B$, $G$ are the

number of permutations and genes, respectively. A significant gene identified by this procedure

is called a temporally differentially expressed gene (TDEG).

An outline of the scFPC-DE workflow is provided in Figure 1. We first apply an established

trajectory analysis tool (e.g., *Monocle3*) to cluster cells and estimate pseudotime from properly

processed scRNA-seq data. Next, we model each gene's log-transformed, mean-centered

expression as a smooth function of pseudotime using roughness-penalized B-splines and

perform FPCA to extract top eigenfunctions and corresponding FPC scores. For each gene, we

compute the distance statistic $D_i$ with and without random pseudotime permutations. The

resulting null distribution of $D_i$ enables us to derive empirical $p$-values and set a decision

boundary in the FPC score space to identify TDEGs. Smoothed gene expression trajectories

produced by scFPC-DE can be visualized in the FPC space overlaid with a directional field from

their pseudo-temporal derivatives. Finally, we perform gene-set and pathway enrichment

analyses on the identified TDEGs to understand the underlying biological processes.

### 2.3 FPC Refinement with Trajectory-Informative Genes

Data from scRNA-seq experiments are highly zero-inflated due to the technical artifact of dropouts [14, 15]. To mitigate the excessive sparsity in the data that may dilute the estimation of pseudo-temporal expression patterns, we adopt a strategy to estimate the eigenfunctions by focusing on a subset of trajectory-informative genes (TIGs) that are likely to exhibit non-flat, informative pseudo-temporal trends.

Selecting highly variable genes is a common practice in the clustering-based scRNA-seq data analysis pipeline (e.g., *Seurat* [22]). However, standard approaches based on dispersion metrics do not necessarily capture useful variability along the pseudotime trajectory. To ensure our FPCA basis represents genuinely dynamic genes, we implement a two-stage procedure focusing on TIGs. First, we fit an initial FPCA model to all genes and compute the distance statistic $D_i$ for each gene, which captures its overall pseudo-temporal variation. Empirical evidence suggests the eigenfunctions obtained in this step are overly smoothed, likely due to the "dilution effects" of non-informative genes and zero inflation (see Supplementary Section 3.2; Figure 8). Next, we select top 20~30% genes ranked by $D_i$ as TIGs and re-estimate the covariance surface, eigenfunctions, and FPC scores using only TIGs. The result is a stable functional basis that reflects major temporal dynamics of informative genes. This retraining procedure improves interpretability and the statistical power of our DE test, particularly under high sparsity.

### 2.4 Simulation Study

We use *scDesign3* [23] to generate synthetic data that reflect realistic characteristics of scRNA-seq data. Simulated dataset is designed to consist of 4,000 genes, including 500 true TDEGs along 500 pseudotime points (cells) equally spaced in a unit interval.

200        To model the dynamic expression patterns, we design the simulation model for the centered

201    and log-transformed mean gene expression curves using three basis functions with distinct

202    dynamic patterns (see Supplementary Figure 1):

$$\phi_1(t) = e^{-t/2}\cos(2\pi t), \phi_2(t) = e^{-t/2}\sin(2\pi t), \quad \phi_3(t) = e^{-2t}. \#(5)$$

204    FPC scores for each TDEG are sampled from a standard normal distribution $\xi_{ik} \sim \mathcal{N}(0,1)$,

205    scaled by eigenvalues prioritizing dominant expression trends: $\lambda_1 = 1$, $\lambda_2 = 0.5$, and $\lambda_3 = 0.25$.

206    For each TDEG, the true mean expression at each pseudotime point is defined as:

$$\mu_i(t_j) = \exp\left(\sum_{k=1}^{3} \sqrt{\lambda_k}\, \xi_{ik}\phi_k(t_j)\right). \#(6)$$

Non-TDEGs have constant mean values along pseudotime, sampled uniformly between 0 and

the median expression of all simulated TDEGs.

Gene expression counts are first generated using a programmatic random number generation

from a zero-inflated negative binomial (ZINB) distribution defined as $W_i(t_j) = W_i^0(t_j)Z_i(t_j)$,

where $W_i^0(t_j) \sim \mathrm{NB}(\mu_i(t_j), \theta = 1)$, and $Z_i(t_j) \sim \mathrm{Bernoulli}(p = 0.25)$ as a dropout indicator

introducing high zero inflation at the same level as observed in the real data analysis. Using the

programmatically generated count data as input, we then use *scDesign3* to further introduce

stochastic components that reflect the realistic characteristics of real scRNA-seq data. The

*scDesign3* simulated data no longer follows the classical distributional assumptions, thus limiting

the validity of methods based on the GAM approach. The high zero inflation dataset will serve as

the primary simulation dataset in this manuscript.

In addition, we generated three supplementary simulation datasets with low to moderate zero

inflation in Supplementary Section 2.1, and present the results in Supplementary Section 2.2,

which show the progressive impact of zero inflation on the method performance.

***2.5 Real Data Analysis***

We applied scFPC-DE to a publicly available real scRNA-seq dataset of B cell subtypes [17]

downloaded          from          the          EMLB-EBI          ArrayExpress          data          portal

(https://www.ebi.ac.uk/biostudies/arrayexpress) with Array Express accession: E-MTAB-9544.

We preprocessed the raw counts with the Seurat workflow [24] and then inferred cell trajectories

and pseudotime ordering with *Monocle3* [3]. The resulting pseudotime trajectories are consistent

with those reported in [17]. For trajectory-based differential expression analyses, we focused on

the developmental trajectory of transitional (Trans), naïve (Naïve), and classical memory

(C-mem1) B cell clusters. Genes with exceptionally low expression or high sparsity across these

subsets were removed (see Supplementary Figure 7). The filtered data still comprises 93.9%

overall sparsity (Supplementary Table 3), mirroring the sparsity of our primary simulated data.

Using scFPC-DE, we identified consistent TDEGs including the canonical markers found in

the original study [17] and B-cell specific pathways. All analyses can be reproduced using scripts

available at: https://github.com/LopezRicardo1/scFPCDE/tree/main/vignettes.

We performed gene set enrichment analysis using the *ReactomePA* [18] package and

Enrichr Knowledge Graph online platform for querying phenotype/network features from multiple

libraries (https://maayanlab.cloud/enrichr-kg) [19]. Reactome pathway analysis was conducted

via Over-Representation Analysis (ORA), using the sets of TDEGs identified by scFPC-DE and

PseudotimeDE as input, against a common background gene universe [18]. *ReactomePA* was

used for Reactome pathway enrichment, while Enrichr-KG was employed to analyze Mammalian

Phenotype Level 4 ontology (2021 release) and construct phenotype-level interaction networks.

Pathways and phenotype categories with FDR-adjusted p-values below 0.05 were considered

significantly enriched.

## Results

### *3.1 Benchmarking Performance of scFPC-DE in Simulations*

PseudotimeDE using a negative binomial generalized additive model (NB-GAM) is considered

the state-of-the-art method for TDEG identification. To benchmark the method performance in

simulations, we compared scFPC-DE with PseudotimeDE using NB-GAM. We also compared

scFPC-DE with FPC-F, which serves as a baseline for a generic FPC-based functional test.

Additional comparisons with the Gaussian and ZINB-GAM variants of PseudotimeDE are

provided in Supplementary Section 2.2.

In the simulation benchmarking, we used the oracle pseudotime simulated in the synthetic

data, which provides a fair performance evaluation of the TDEG identification itself and removes

the uncertainty from pseudotime estimation. Table 1 summarizes the type I error rate and

statistical power of the three benchmarked methods using the high zero inflation simulated

dataset. scFPC-DE consistently showed the best type I error control while maintaining

competitive power compared to the other two methods. Figure 2 presents the systematic

evaluation of the simulation results in the ROC analysis. For type I error control, scFPC-DE

always had the closest observed false positive rate as the expected nominal α level (Figure 2a).

The ROC analysis also confirmed that scFPC-DE outperformed the competing methods by

achieving the highest AUC statistic (Figure 2b), indicating the overall improved performance of

sensitivity and specificity. For the supplementary simulated data with moderate-to-low zero

inflation, the performance gap between scFPC-DE and benchmarked methods are smaller

(Supplementary Section 2.2), suggesting that the differences in type I error control are tightly

linked to the dropout rate in the data [14].

To further illustrate from where scFPC-DE gained the overall improvement, we projected the

simulated data in the FPC score space (Figure 2c). By employing FPCA to capture the global

covariance patterns across genes [9], scFPC-DE is able to define a more precise and stable null

boundary in the FPC score space than the gene-wise GAM approaches [4, 25]. That is the main

reason scFPC-DE achieves superior discrimination between true signal and noise (Figure 2c).

In the simulation studies, we also compared the computing time needed by each method.

scFPC-DE demonstrated substantial computational gains, requiring only 4.82 seconds on

average per dataset—more than ten times faster than the computing time of PseudotimeDE

(49.62 seconds; Supplementary Table 2).

***3.2 scFPC-DE Performance in Real Data Analysis***

In the scRNA-seq dataset of B cell subtypes   [17], we reconstructed the pseudotime trajectory

using *Monocle3* (Figure 3a–b), which reproduces the results of Figure 2A in the original study.

We captured the cell trajectory of the development pathway of Trans, Naïve, and C-mem1 B cell

clusters, which recapitulates the biological cell states along the pseudotime (Figure 3c). By

applying scFPC-DE, gene expression changes along the trajectory are modeled as smooth

functions of pseudotime. In Figure 3d, we visualized selected genes for the Trans, Naïve, and

C-mem 1 trajectory from the original study using our scFPC-DE framework. As expected, all

selected genes were identified as TDEG using scFPC-DE.

In the FDA-based FPC modeling framework, FPCA-derived eigenfunctions (i.e., FPCs) can

effectively capture the principal modes of gene expression variations along the pseudotime

(Figure 4a). The functional variance-covariance structure, $C(t,s)$, shared among genes indicate

that there are increased gene-gene interactions among neighboring pseudotime points (Figure

4b). The linear structure of eigenfunctions also permits efficient computation of time derivatives,

which may suggest the directions of cell development in the directional vectors in the FPC score

space (Figure 4c). These vectors, derived via straight forward calculations involving

eigenfunctions derivatives and FPC scores pseudo inverse [26-28], offer a geometric perspective

on local developmental progression and complement the global pseudotime covariance captured

in the FDA model (see Supplementary Section 1.3 for details). Projecting cells into the FPC space

showed coherent cluster structure and cell state transitions as expected by the biological cell

developmental trajectory and pseudotime estimated by *Monocle3*, providing a strong internal

validation of the useful information captured in top FPC space and the cell trajectory derived by

the pseudotime analysis (Figure 4c–d).

We also compared the results by using scFPC-DE and PseudotimeDE in the real dataset.

Out of 2980 total genes, scFPC-DE identified 474 TDEGs and PseudotimeDE identified 1795

TDEGs along the Trans, Naïve, and C-mem1 trajectory. In Figure 5a, we compare the

distributions of FDR-corrected p-values obtained from scFPC-DE and PseudotimeDE. The

p-value histogram from scFPC-DE shows a relatively uniform distribution with a modest

concentration near zero, reflecting conservative type I error control and a more stringent DEG

selection. In contrast, PseudotimeDE yields a sharp spike near zero with minimal values

elsewhere, indicating potential inflation of false positives. In Figure 5b, the TDEG results are

projected in the FPC score space. The scFPC-DE result led to a clear separation between null

genes and TDEGs, proving a stable decision boundary. This stable separation is achieved by

using the projected $L^2$ distance ($D_i$) as the test statistic, which quantifies the deviation from a

constant mean gene expression function in FPCA. This distance measure provides a meaningful

ranking of genes in the FPC score space after the effective variance decomposition through

FPCA. In Figure 6, we visualized the top 10 TDEGs identified by scFPC-DE. The smoothed

functions of the top TDEGs exhibit pronounced dynamic expression patterns across pseudotime,

including the predicted downregulation of IgD, a known hallmark of B cell maturation.

**3.3 Pathway enrichment analysis**

We    used    the    *ReactomePA*    R    package    [18]    and    the    *Enrichr-KG*    online    tool

([https://maayanlab.cloud/enrichr-kg](https://maayanlab.cloud/enrichr-kg)) [19] to conduct pathway enrichment analysis for the TDEG

lists identified by scFPC-DE and PseudotimeDE. In addition to the Reactome pathways, by using

*Enrichr-KG*, we included more biologically curated pathway and phenotype ontologies, such as

the KEGG pathways, the Mammalian Phenotype Ontology [29], the Human U133A/GNF1H

Gene Atlas [30] and the HuBMAP ASCT+B tables for single cell-specific annotations [31]

Despite identifying substantially fewer TDEGs (Figure 7a), scFPC-DE uncovered 43 uniquely

enriched pathways that are associated with B-cell immune functions and lineage progression in

the Reactome pathway analysis (Figure 7b; Supplementary Table 4). While both scFPC-DE and

PseudotimeDE identified 25 common pathways (Figure 7b), the four enriched pathways unique to

PseudotimeDE reflected broad cellular functions with limited immune relevance (Supplementary

Table 5). This contrast indicates higher biological specificity of the scFPC-DE result by precisely

modeling the functionally coherent gene programs and avoiding false discoveries. These results

are consistent with our simulation observations, where scFPC-DE maintained sufficient power

while substantially reducing false positives, especially for data with high zero inflation. Both

methods recovered infection-related pathways (Supplementary Table 6) in which B cells play

important roles, however scFPC-DE achieved this with only a quarter of the input TDEGs

compared with PseudotimeDE. Notably most immune-specific pathways were uniquely identified

by scFPC-DE. This demonstrates that scFPC-DE can identify a more compact set of TDEGs with

higher signal-to-noise ratio and biologically more coherent. We additionally examined the 25

Reactome pathways shared across both methods. Enrichment p-values were consistently and

substantially smaller when using the scFPC-DE gene set, reflecting stronger statistical support for

true pathway-level associations. Using Over-Representation Analysis (ORA) summary statistics

from *ReactomePA*, the most striking result highlighting the advantage of scFPC-DE is portrayed

in Figure 7c, which compares the enrichment strength between scFPC-DE and PseudotimeDE

using gene-to-background ratio plots. scFPC-DE identified immune and developmental pathways

with higher gene ratios and greater statistical significance, while PseudotimeDE primarily

highlighted broad categories such as mRNA splicing with weaker enrichment signals. Notably, the

largest gene-to-background ratio was observed for the Immune System pathway, reinforcing the

biological context of this analysis and underscoring the relevance of scFPC-DE in capturing

immune-specific gene programs. Overall, scFPC-DE yields consistently stronger pathway

enrichment signals in both effect size and statistical significance compared to PseudotimeDE. In

summary, scFPC-DE identifies a more concise and biologically informative set of TDEGs for

pathway analysis.

To further assess the biological relevance of the TDEG lists from the two DE methods, we

used the convenient *Enrichr-KG* online tool to query more biological entity databases. Figure 7d-e

show the top 5 terms from the selected ontologies. Using the single cell-specific knowledge

curated in the HuBMAP ASCT+B tables (green), the scFPC-DE list correctly annotated the B cells

from blood as PBMC samples were used in the original study (Stewart, et al., 2021). The thoracic

thymus is a critical hub for developing central immune tolerance; thus, it would be expected to

contain similar populations of Naïve B cell. On the contrary, the PseudotimeDE list annotated two

B cell populations from spleen – an irrelevant organ to the real data, therefore the TDEG list is

likely to contain misinformation from false discoveries. Both methods recognized specific terms

related to developmental and mature B cell phenotypes and functions, such as the CD19+ B cells

(purple), humoral immune response (orange), and B cell receptor signaling pathway (grey), with

higher ranking in the scFPC-DE output ordered by the combined score of *Enrichr-KG* than in the

PseudotimeDE output. Supplementary Figure 9 also shows the knowledge graph visualization of

biological phenotypes and connected genes derived from the TDEG lists, revealing gene-specific

contributions to the phenotypes.

Together, these results illustrate that scFPC-DE can effectively detect a more interpretable,

statistically robust, and biologically meaningful enrichment profile than existing method such as

PseudotimeDE.

## Discussion

In this study, we introduced scFPC-DE, a novel method based on functional data analysis

techniques for trajectory-based differential expression analysis using scRNA-seq data. By

leveraging FPCA, scFPC-DE provides a continuous, data-driven representation of gene

expression dynamics along the cell trajectory reconstructed by pseudotime. The FPCA modeling

framework captures the dominant variation across genes while preserving the temporal pattern

inherent to the underlying cellular progression in the cell trajectory, enabling a more integrated

and interpretable model of gene expression changes.

Simulation studies, designed to reflect the high zero inflation phenomenon in the real

scRNA-seq data, demonstrate that scFPC-DE offers substantial advantages over existing

approaches. It consistently achieved the best control of type I error while attaining the highest

AUC in the ROC analysis. The improved overall performance is largely attributed from its ability

to borrow information across genes and across pseudotime, in contrast to the GAM-based

methods that model each gene independently and lack global covariance modeling.

While PseudotimeDE and other GAM-based approaches have advanced the modeling of

nonlinear trajectories, they fail to incorporate inter-gene dependencies. In contrast, scFPC-DE's

FPCA-based approach effectively captures the global inter-gene variance-covariance structure

in the top FPC space and uses functional projection to distinguish null from alternative

hypotheses in the FPC space with a robust null boundary that leads to a tighter control of type I

error, particularly for scRNA-seq data with high zero inflation. Application to a well-characterized

B cell subtypes dataset further illustrates the advantages of scFPC-DE. With a more concise list

of DEGs identified by scFPC-DE than by PseudotimeDE, the scFPC-DE identified DEG list

revealed specific immune pathways related to B cell functions in Reactome enrichment analysis, indicating a stringent control of false positive DEGs by scFPC-DE. The real data analysis suggests that scFPC-DE can effectively avoid those noises associated with a large number of false positive genes, yielding a cleaner and more biologically coherent enrichment analysis.

One limitation in the current scFPC-DE pipeline is that it takes in the estimated pseudotime as input data, which may potentially impact the inference. In this study, our results indicate that our FPCA-based framework remains robust to variations in the input pseudotime, as shown by its consistent performance across simulations and real data. As a natural extension, we plan to develop an end-to-end FDA-based pipeline that integrates pseudotime inference and differential expression testing into a unified functional framework. We also plan to extend scFPC-DE to support complex branching trajectories and to include functional pseudotime estimation using the same FDA framework.

In conclusion, scFPC-DE defines a statistically rigorous and biologically interpretable framework for trajectory-based differential expression analysis. Its ability to model smooth gene expression variation across pseudotime, incorporate inter-gene covariance, and control false discoveries marks a significant advancement over existing methods.

## R Package Availability

The scFPC-DE R package implementing all methods described in this study is publicly available at: https://github.com/LopezRicardo1/scFPCDE.

## Funding

Research reported in this publication was partially supported by the National Institute of Environmental Health Sciences of the National Institutes of Health (NIH) under award number T32ES007271, National Institute of Allergy and Infectious Diseases of the National Institutes of Health (NIH) under award numbers 1R01AI184931, 1U01AI187057 and 1R01AI168917. This

research was supported in part by the Intramural Research Program of the National Institutes of

Health (NIH). The contributions of the NIH author(s) are considered Works of the United States

Government. The findings and conclusions presented in this paper are those of the author(s) and

do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human

Services.

*Conflict of Interest:* none declared.

**Tables and Figure Legend**

**Table 1. Type I error rate (TIER) and statistical power (PWR) across varying significance**

**levels ($\alpha$) under a high zero inflation setting (93.9% sparsity).** Simulation benchmarking

results compare the performance of scFPC-DE, FPC-F, and PseudotimeDE for identifying the

true TDEGs. scFPC-DE consistently exhibits the lowest TIER across all α thresholds, indicating

superior type I error control. Notably, at α = 0.01, scFPC-DE reduces the TIER to 0.017,

outperforming both FPC-F (0.039) and PseudotimeDE (0.049). At more lenient thresholds ($\alpha$ =

0.05 and 0.10), the FPRs of competing methods increase substantially, while scFPC-DE

maintains markedly better calibration. Despite this advantage in error control, scFPC-DE does not

compromise on statistical power. It attains power values that are comparable or superior across

all tested levels—demonstrating robust sensitivity to true signal even under extreme dropout

conditions. The best results for each significance level are shown in boldface.

**Fig. 1. scFPC-DE workflow.** (1) Data preprocessing of the input scRNA-seq dataset, including

normalization, log-transformation, dimensionality reduction, finding neighbors, UMAP/tSNE

embedding, and clustering. Note that the scFPC-DE workflow uses the log-transformed gene

expression data as the input data for the downstream analysis workflow. (2) Trajectory analysis

for estimating the pseudotime of cells in a trajectory of interest. (3) Modeling the log-transformed

gene expression as a smoothed function of pseudotime centered at functional mean expression

of 0 for each gene. Examples of a temporally DE gene (left) and a temporally non-DE gene

(right). (4) Identifying the top eigenfunctions $\Phi_k(t)$ from FPCA and construction of the FPC

latent space. (5) Visualization of the reconstructed cell trajectory in the FPC space. (6) Projecting

each gene onto the FPC space allows calculation the distance statistic $D_i$ for hypothesis testing.

Genes are represented as coordinates with respect to the eigenfunctions, with the significance

decision boundary is denoted by dashed ellipsoid; genes positioned farther away are classified

as TDEGs/TIGs. (7-8) Enrichment analysis of DEGs and knowledge network visualization using

online tools.

**Fig. 2. Simulation results for scFPC-DE and benchmarking methods**. (a) Observed versus

expected type I error curves demonstrate that scFPC-DE aligns closely with the diagonal line,

demonstrating its accurate type I error control across significance thresholds. In contrast, both

FPC-F and PseudotimeDE deviate above the diagonal line, indicating inflated error rates, with

PseudotimeDE showing the most severe inflation. (b) ROC curve comparison among scFPC-DE

(AUC = 0.867), FPC-F (AUC = 0.798), and PseudotimeDE (AUC = 0.795) shows that scFPC-DE

achieves the best performance in sensitivity and specificity. (c) In the FPC projected score space,

null genes (grey) cluster near the origin, while true DEGs (black) separate outward. Controlling at

the 5% significance level, scFPC-DE produces fewer false positives (245 out of 4000; red), mostly

located at the boundary between null and alternative regions, compared to the substantially

higher false positive misclassifications observed in both FPC-F (430) and PseudotimeDE (501).

**Fig. 3. Pseudotime reconstruction and smoothed gene expression functions using real**

**data.** (a) 3-D UMAP (UMAP 1 and UMAP 3) as referenced in Stewart et al. (2021), with

reconstructed *Monocle3* cell trajectory. (b) Pseudotime-colored UMAP shows consistency with

the results shown in referenced study. (c) Cell type ordering and pseudotime distribution for the

Trans, Naive, and CMem1 subtypes annotated by the referenced study. (d) Gene expression

functions for the biologically relevant genes identified in Stewart et al. (2021), Figure 2C. The

smoothed gene expression functions were modeled by the FPC framework. scFPC-DE also

identified these genes as TDEGs after false discovery rate (FDR) correction.

**Fig. 4. FPC space reconstruction in real data analysis.** (a) Top 3 FPC eigenfunctions explain

92.6% of the total variation along the cell trajectory between Trans, Naïve, and C-mem1 clusters,

representing the major pseudo-temporal gene expression patterns. (b) Pseudotime covariance

contour structure shows regions of high gene expression interactions along cell pseudotime

progression. (c) Projecting cells onto the FPC space deciphers the cell development trajectory,

with directional vectors indicating cell direction along the cell trajectory. (d) Cells projected in the

FPC space with pseudotime estimated by *Monocle3*.

**Fig. 5. scFPC-DE and PseudotimeDE comparison on real data**. (a) FPC gene score

projection identifies highly differentially expressed genes, showing a well-defined boundary

between the null genes and TDEGs identified by scFPC-DE. In contrast, PseudotimeDE lacks a

clearly defined boundary. (b) The distribution of FDR-corrected p-values between scFPC-DE and

PseudotimeDE.

**Fig. 6. Gene expression patterns of the top TDEGs identified by scFPC-DE**. Top 10 TDEGs

ranked by the scFPC-DE projected distance statistic allows further inspection for TIGs along cell

pseudotime progression. Violin plots grouped by the discrete clusters as well as the smoothed

gene expression functions are plotted for each top TDEG. The functional data representation

offers a dynamic view of the continuous gene expression changes across the cell states.

**Fig. 7. Pathway and phenotype-level enrichment analysis results for scFPC-DE and**

**PseudotimeDE.** (a) Number of method-specific and common TDEGs identified by scFPC-DE

and PseudotimeDE. scFPC-DE identified a smaller, more targeted set of TDEGs—approximately

one-fourth the size of that returned by PseudotimeDE. (b) Number of method-specific and

common pathways identified in the Reactome pathway analysis. scFPC-DE uniquely identified 43

pathways, PseudotimeDE identified 4 unique pathways, and 25 pathways were shared between

the two methods. Even when sharing most of the same DEGs, PseudotimeDE did not identify

these unique pathways when following the same GSEA protocol. (c) The gene-to-background

ratio plots show that the enrichment signals from scFPC-DE are stronger for both shared and

unique pathways. (d–e) The *Enrichr-KG* results for pathway- and phenotype-level enrichment

analysis using multiple ontologies (MGI Mammalian Phenotype Level 4, HuBMAP ASCT+B 2022,

Human Gene Atlas, and KEGG 2021).

# References

495
496
1.  Saelens W, Cannoodt R, Todorov H, Saeys Y: **A comparison of single-cell trajectory inference methods**. *Nature biotechnology* 2019, **37**(5):547-554.

497
498
499
2.  Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen TS, Rinn JL: **The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells**. *Nature biotechnology* 2014, **32**(4):381-386.

500
501
3.  Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, Zhang F, Mundlos S, Christiansen L, Steemers FJ: **The single-cell transcriptional landscape of mammalian organogenesis**. *Nature* 2019, **566**(7745):496-502.

502
503
504
4.  Van den Berge K, Roux de Bézieux H, Street K, Saelens W, Cannoodt R, Saeys Y, Dudoit S, Clement L: **Trajectory-based differential expression analysis for single-cell sequencing data**. *Nature communications* 2020, **11**(1):1201.

505
5.  Moran PA: **Notes on continuous stochastic phenomena**. *Biometrika* 1950, **37**(1/2):17-23.

506
507
6.  Song D, Li JJ: **PseudotimeDE: inference of differential gene expression along cell pseudotime with well-calibrated p-values from single-cell RNA sequencing data**. *Genome biology* 2021, **22**(1):124.

508
509
7.  Ramsay J, Hooker G, Graves S, Ramsay J, Hooker G, Graves S: **Introduction to functional data analysis**. *Functional data analysis with R and MATLAB* 2009:1-19.

510
8.  Ramsay JO, Silverman BW: **Applied functional data analysis: methods and case studies**: Springer; 2002.

511
512
9.  Wu S, Wu H: **More powerful significant testing for time course gene expression data using functional principal component analysis approaches**. *BMC bioinformatics* 2013, **14**:1-13.

513
514
10. Zhang Y, Topham DJ, Thakar J, Qiu X: **FUNNEL-GSEA: FUNctioNal ELastic-net regression in time-course gene set enrichment analysis**. *Bioinformatics* 2017, **33**(13):1944-1952.

515
11. Ramsay JO, Silverman BW: **Functional data analysis**: Springer; 2005.

516
517
12. Payen R: **Fonctions aléatoires du second ordre à valeurs dans un espace de Hilbert**. In: *Annales de l'institut Henri Poincaré Section B Calcul des probabilités et statistiques: 1967*. 323-396.

518
519
13. Karhunen K: **Über lineare Methoden in der Wahrscheinlichkeitsrechnung**. *Ann Acad Sci Fennicae* 1947, **37**:1.

14. Silverman JD, Roche K, Mukherjee S, David LA: **Naught all zeros in sequence count data are the same**.
*Computational and structural biotechnology journal* 2020, **18**:2789-2798.

15. Jiang R, Sun T, Song D, Li JJ: **Statistics or biology: the zero-inflation controversy about scRNA-seq data**.
*Genome biology* 2022, **23**(1):31.

16. Su Y-R, Di C-Z, Hsu L: **Hypothesis testing in functional linear models**. *Biometrics* 2017, **73**(2):551-561.

17. Stewart A, Ng JC-F, Wallis G, Tsioligka V, Fraternali F, Dunn-Walters DK: **Single-cell transcriptomic analyses**
**define distinct peripheral B cell subsets and discrete development pathways**. *Frontiers in immunology*
2021, **12**:602539.

18. Yu G, He Q-Y: **ReactomePA: an R/Bioconductor package for reactome pathway analysis and visualization**.
*Molecular BioSystems* 2016, **12**(2):477-479.

19. Evangelista JE, Xie Z, Marino GB, Nguyen N, Clarke DJ, Ma'ayan A: **Enrichr-KG: bridging enrichment**
**analysis across multiple libraries**. *Nucleic acids research* 2023, **51**(W1):W168-W179.

20. Ramsay JO, Silverman BW: **Principal components analysis for functional data**. In: *Functional data analysis.*
Springer; 2005: 85-109.

21. Ramsay J, Wickham H, Ramsay MJ, deSolve S: **Package 'fda'**. *R Project, July* 2018.

22. Satija R, Farrell JA, Gennert D, Schier AF, Regev A: **Spatial reconstruction of single-cell gene expression**
**data**. *Nature biotechnology* 2015, **33**(5):495-502.

23. Song D, Wang Q, Yan G, Liu T, Sun T, Li JJ: **scDesign3 generates realistic in silico data for multimodal**
**single-cell and spatial omics**. *Nature Biotechnology* 2024, **42**(2):247-252.

24. Gribov A, Sill M, Lück S, Rücker F, Döhner K, Bullinger L, Benner A, Unwin A: **SEURAT: visual analytics for**
**the integrated analysis of microarray data**. *BMC medical genomics* 2010, **3**:1-6.

25. Hastie TJ: **Generalized additive models**. *Statistical models in S* 2017:249-307.

26. MacAusland R: **The moore-penrose inverse and least squares**. *Math 420: Advanced Topics in Linear*
*Algebra* 2014:1-10.

27. Prasad KM, Bapat R: **The generalized moore-penrose inverse**. *Linear Algebra and its Applications* 1992,
**165**:59-69.

28. Shang HL: **A survey of functional principal component analysis**. *AStA Advances in Statistical Analysis* 2014,
**98**:121-142.

29. Smith CL, Goldsmith C-AW, Eppig JT: **The Mammalian Phenotype Ontology as a tool for annotating,**
**analyzing and comparing phenotypic information**. *Genome biology* 2004, **6**(1):R7.

30. Su AI, Wiltshire T, Batalov S, Lapp H, Ching KA, Block D, Zhang J, Soden R, Hayakawa M, Kreiman G: **A gene**
**atlas of the mouse and human protein-encoding transcriptomes**. *Proceedings of the National Academy of*
*Sciences* 2004, **101**(16):6062-6067.

31. Börner K, Blood PD, Silverstein JC, Ruffalo M, Satija R, Teichmann SA, Pryhuber GJ, Misra RS, Purkerson JM,
Fan J: **Human biomolecular atlas program (hubmap): 3D human reference atlas construction and usage**.
*Nature methods* 2025:1-16.

| $\alpha$ | scFPC-DE | | FPC-F | | PseudotimeDE | |
| --- | --- | --- | --- | --- | --- | --- |
| | TIER | PWR | TIER | PWR | TIER | PWR |
| 0.01 | **0.017** | **0.434** | 0.039 | 0.406 | 0.049 | 0.426 |
| 0.05 | **0.070** | 0.602 | 0.123 | 0.552 | 0.143 | **0.604** |
| 0.1 | **0.130** | **0.694** | 0.193 | 0.632 | 0.217 | 0.674 |

# scFPC-DE Workflow

**(1) Pre-processing**

**(2) Cell Trajectory Pseudotime**

**(3) Centered Gene Expression**
Smooth Function of Pseudotime

DE Gene | Non-DE Gene

**(4) Learn Functional Modes**
FPC Reconstruction

Eigenfunction — Phi1 — Phi2

**(5) Cell Projected in FPC Space**

Cluster: 1, 2, 3

**(6) Project Genes in FPC Score Space**
Run scFPC-DE Test

**(7) Top Enriched Pathways**

**(8) Gene–Pathway Knowledge Network**

# Simulation Sensitivity Analysis Results

**(a)** **Type-I Error Control**
Observed vs Expected False Positive

**(b)** **ROC - AUC by Test**

AUC = 0.867-scFPC_DE
AUC = 0.798-FPC_F
AUC = 0.795-PseudotimeDE

**(c) Simulation Gene FPC Score Projection**

Test Result

False Negative
False Positive
True Negative
True Positive

**Eigenfunctions, Expression Covariance and FPC Space Cell Projection**

**(a) FPC Eigen Functions**

PEV = 92.6 %

Eigenfunction
— FPC1
– – FPC2
⋯ FPC3

**(b) Expression Covariance Countour**

Covariance

**(c) Cell Projected in FPC Space**

Colored by Cell Cluster

Cluster
● Trans
● Naive
● C-mem1

**(d) Cell Projected in FPC Space**

Colored by Pseudotime

Pseudotime

**UMAP with Cell Clusters and Pseudotime Trajectory**

**(a)**

**(b)**

**(c) Ordered Violin of Pseudotime by Cluster**

**(d) FPC-Smoothed Gene Expression**

# Real Data FPC Gene Scores and TDEA Test p-Values

**(a)** Gene Scores Colored by Test Desicion

**(b)** FDR Corrected p-Values

# FPC Smoothed Expression Curves Along Pseudotime

**Reactome Over Representation Analysis and EnrichR KG Network**

(a) DEGs Sets By Method

scFPC_DE 9 | 474 | PseudotimeDE 1795

(b) Enriched Reactome Pathways

scFPC_DE 43 | 25 | PseudotimeDE 4

(c) scFPC-DE - Gene vs BG Ratio

Immune System
Developmental Biology
Cellular responses to stimuli
Cellular responses to stress

-log10(q.val): 4, 8, 12, 16

Shared: Common, Unique

PseudotimeDE - Gene vs BG Ratio

HIV Infection mRNA Splicing
mRNA Splicing - Major Pathway

(d) scFPC-DE - KG Network

HuBMAP_ASCTplusB_augmented_2022 — Transitional Stage B cell - Blood
HuBMAP_ASCTplusB_augmented_2022 — Immature B cell - Blood
HuBMAP_ASCTplusB_augmented_2022 — Naive B cell - Blood
HuBMAP_ASCTplusB_augmented_2022 — Naive B cell - Thoracic Thymus
HuBMAP_ASCTplusB_augmented_2022 — Memory B cell - Blood
Human_Gene_Atlas — CD19+ BCells(neg. sel.)
KEGG_2021_Human — Coronavirus disease
MGI_Mammalian_Phenotype_Level_4_2021 — abnormal B cell physiology MP:0002459
KEGG_2021_Human — Ribosome
MGI_Mammalian_Phenotype_Level_4_2021 — abnormal B cell differentiation MP:0002144
MGI_Mammalian_Phenotype_Level_4_2021 — decreased IgG1 level MP:0008495
MGI_Mammalian_Phenotype_Level_4_2021 — abnormal humoral immune response MP:0001800
MGI_Mammalian_Phenotype_Level_4_2021 — increased pro-B cell number MP:0008186
KEGG_2021_Human — B cell receptor signaling pathway
Human_Gene_Atlas — BDCA4+ DendriticCells
Human_Gene_Atlas — CD8+ Tcells
Human_Gene_Atlas — Lymphoma burkitts(Raji)
Human_Gene_Atlas — CD56+ NKCells
KEGG_2021_Human — Leishmaniasis
KEGG_2021_Human — Hematopoietic cell lineage

(e) PseudotimeDE - KG Network

HuBMAP_ASCTplusB_augmented_2022 — Transitional Stage B cell - Blood
HuBMAP_ASCTplusB_augmented_2022 — Naive B cell - Spleen
HuBMAP_ASCTplusB_augmented_2022 — Immature B cell - Blood
HuBMAP_ASCTplusB_augmented_2022 — Naive B cell - Blood
HuBMAP_ASCTplusB_augmented_2022 — Mantle Zone B Cell - Spleen
Human_Gene_Atlas — CD8+ Tcells
Human_Gene_Atlas — CD19+ BCells(neg. sel.)
Human_Gene_Atlas — CD56+ NKCells
MGI_Mammalian_Phenotype_Level_4_2021 — abnormal B cell physiology MP:0002459
MGI_Mammalian_Phenotype_Level_4_2021 — decreased B-1a cell number MP:0008168
MGI_Mammalian_Phenotype_Level_4_2021 — decreased B cell proliferation MP:0005093
MGI_Mammalian_Phenotype_Level_4_2021 — abnormal humoral immune response MP:0001800
MGI_Mammalian_Phenotype_Level_4_2021 — decreased IgG1 level MP:0008495
Human_Gene_Atlas — CD4+ Tcells
Human_Gene_Atlas — BDCA4+ DendriticCells
KEGG_2021_Human — Epstein-Barr virus infection
KEGG_2021_Human — Coronavirus disease
KEGG_2021_Human — Measles
KEGG_2021_Human — Colorectal cancer
KEGG_2021_Human — B cell receptor signaling pathway