Rapamycin can revert CCM phenotype in brain endothelial cells and ameliorates chronic cavernoma development in combination with lapatinib

Mar García-Colomer[1], José E Martínez[1], Luis Diaz-Gomez[2], Miriam Sartages[1], Eva M. Esquinas-Román[3], Cristina Riobello[3], David Martínez-Delgado[1], Diego González-Pérez[3], Aurora Gómez-Durán[3], Miguel Fidalgo[1], Marta Varela-Rey[3], Celia M Pombo[1*], Juan Zalvide[1*]

[1]Department of Physiology, Centro Singular de Medicina Molecular e Enfermedades Crónicas (CiMUS) and Instituto Sanitario de Santiago de Compostela (IDIS), Universidade de Santiago de Compostela, (USC), Santiago de Compostela, 15782, Spain.

[2]Department of Pharmacology, Pharmacy, and Pharmaceutical Technology, Instituto de Materiales (iMATUS), and IDIS, Universidade de Santiago de Compostela, Santiago de Compostela, 15782 Spain.

[3]Department of Biochemistry and Molecular Biology, CiMUS and IDIS, USC, Spain.

*Corresponding authors: Celia M. Pombo: Tel: +34-881815413. E-mail: celiamaria.pombo@usc.es; Juan Zalvide, Tel: +34-881815414. Email juan.zalvide@usc.es.

Short title: Rapamycin, CCM phenotype and cavernomas

Word count: 7591.

Abstract

Objective

This study investigates the impact of rapamycin and propranolol on cerebral cavernous malformations (CCMs). Employing an unbiased transcriptomic analysis, we aimed to comprehensively elucidate the molecular mechanisms underlying these drug effects. We also sought to determine whether rapamycin, especially in combination with the tyrosine kinase inhibitor lapatinib, can reduce lesion volume even after lesion growth has occurred.

Approach and results

Mouse Brain Microvascular Endothelial Cells (mBMEC) deficient in *Ccm3* were treated with propranolol (10 µM) or rapamycin (100 nM) and were analyzed by RNA-seq and immunofluorescence. *Ccm3*-null mice in which cavernoma development had been induced were treated with rapamycin alone or combined with lapatinib, assessing lesion volume using micro-CT imaging.

While propranolol shows limited efficacy in modulating the CCM transcriptomic phenotype in mBMEC, rapamycin demonstrates a significant impact. Rapamycin reverses gene expression changes induced by *Ccm3* deficiency, restoring *Klf2/4*-dependent genes like *Nos3*, *Adamts1*, and *Thbs1*. Notably, we observed a reduction in KLF2 protein levels in *Ccm3* KO cells treated with rapamycin. Critically, *in vivo* experiments demonstrate that a combination of rapamycin and lapatinib effectively reduces lesion volume in a chronic CCM model.

Conclusion

Rapamycin normalizes the expression of key genes implicated in cavernoma pathogenesis in *Ccm3* KO endothelial cells, offering new insights into its potential therapeutic mechanisms. In contrast, the observed lack of efficacy of propranolol in brain endothelial cells challenges current understanding of its mechanism of action in CCM treatment and warrants further investigation. Significantly, rapamycin, when combined with lapatinib, exhibits promising efficacy *in vivo* by reducing the volume of established CCM lesions. This finding is particularly noteworthy as it suggests a potential treatment strategy for existing lesions. Our work has identified a path for the application of drug combination in the treatment of CCMs.

Graphic abstract

Non-standard abbreviations and acronyms

CCMs: Cerebral cavernous malformations.

mBMEC: Mouse Brain Microvascular Endothelial Cells.

fCCM: familial cavernomatosis.

Introduction

Cerebral cavernous malformations (CCMs) are vascular lesions, predominantly occurring in the central nervous system and associated with recurrent hemorrhages and seizures(1, 2). CCMs are among the most common cerebral vascular malformations, especially in young people (it is estimated that 0.5% of the population has at least one CCM)(3, 4). Most CCMs are sporadic, but they can also have a genetic basis caused by inactivating mutations in one of the three CCM genes: *KRIT1/CCM1*, *OSM/CCM2* and *PDCD10/CCM3* (4). Patients with this familial cavernomatosis (fCCM) have multiple CCMs, which increases the severity of their condition. The products of these CCM genes form a ternary complex which binds to MEKK3, resulting in the inhibition of the MEKK3/MEK5/ERK5 axis and downregulation of the *KLF2* and *KLF4* transcription factors. Dysregulation of *KLF2* and *KLF4* in CCM mutant cells is essential for cavernoma development (5).

The current treatment for CCMs relies on surgical intervention for accessible lesions in the brain and spinal cord. However, for deep-seated CCM lesions and for fCCMs, when many lesions may appear, novel pharmacological interventions are clearly needed, and their development is a long-standing goal of translational CCM research. As a result of the studies on the function of CCM genes and cavernoma pathogenesis, several possible drug treatments have been proposed, among them statins, known to inhibit Rho activity, which is deregulated in CCM-deficient cells, especially in *Ccm1-* and *Ccm2*-deficient endothelium (6, 7); sulindac, which inhibits transcriptional β-catenin activity(8); the multikinase inhibitor sorafenib, which inhibits excessive angiogenesis (9); the free radical scavenger tempol (10); intestinal microbiome modification to alter Toll-mediated

signaling (11); vitamin D and indirubin-3-monoxime, identified in a systematic pharmacological screen (12); lapatinib, because of the increase of EGFR in *Ccm3*-deficient endothelial cells (13); or rapamycin, based on the stimulated mTOR activity in CCM lesions (14). All these treatments diminish one or several of the effects of CCM deficiency *in vitro*, and some of them diminish the number of lesions that arise in animal models of cavernoma development, as long as they are given before their appearance or growth. Despite promising preclinical results, no clinically significant effect has been found in the two trials performed so far, a pilot study studying possible effects of by simvastatin on lesion permeability (15) and a blinded study on the effects of atorvastatin on lesion bleeding (16). Also, the phase 2 trial using the drug REC-99 (NCT05085561) has recently been discontinued because of lack of clinical benefits.

Interestingly, the largest number of clinical trials proposed for CCMs involve the non-selective β-blocker propranolol (trials NCT03589014, NCT03474614 and NCT03523650), which has been selected due to its known effect on infantile hemangiomas (17). Another treatment that has been proposed recently is rapamycin. This was initially based on the seminal observation that *Pik3ca* gain of function (GOF) mutations fueled growth of cavernoma lesions and was reinforced by the fact that CCM-deficient endothelial cells without *PIK3CA* mutations also had increased phosphorylation of the mTOR target S6 (14).

Propranolol treatment has been shown to reduce lesion burden in chronic murine models of cavernoma development and restore barrier function when initiated before or during lesion growth (18). Furthermore, while meta-analyses of clinical studies have not demonstrated a statistically significant protective effect of β-blockers (including propranolol) in preventing intracerebral hemorrhage or focal neurologic deficits in individuals with CCM, a phase II clinical trial suggested a potential benefit in preventing

cavernoma development, although the study was not powered to achieve statistical significance (19). Despite these indications of activity, the clinical efficacy of propranolol remains unclear. Addressing this uncertainty requires further research, as the cellular mechanism of propranolol's effect on CCM development is unknown, including the specific cell type in which it exerts its effects. Several studies have described the effects of propranolol on CCM-deficient endothelial cells, but at a concentration of 100 μM (20), which is significantly higher than the estimated *in vivo* concentration after propranolol treatment, even in the brain (no more than 10 μM). Moreover, treatment of mice with 15 mg/kg.day of propranolol did not affect lesion size or hemorrhage in already established cavernomas, at least under the treatment regimen used (20).

As stated above, the combination of the detection of *Pik3ca* GOF mutations in large cavernomas and enhanced mTOR activity in CCM-deficient cells makes mTOR-inhibiting drugs such as rapamycin good candidates for pharmacological treatment of cavernomas. Enhanced activity of mTOR has been shown in *CCM1*-deficient HuVECs (14) and in progenitor endothelial cells in *in vivo* lesions (21). Its inhibition with rapamycin reverts some of the effects of CCM loss in the latter, without affecting the mRNA overexpression of *Klf2* or *Klf4*, two genes that are important for cavernoma pathogenesis. It is unclear through what mechanism rapamycin affects the CCM phenotype, although it is believed that it acts specifically on progenitor endothelial cells (21).

In agreement with *in vitro* effects, treatment with rapamycin *in vivo* inhibits cavernoma development in several preclinical models (14, 21) However, the effects of rapamycin are not clear when a chronic model of CCM development and lower doses of the drug are used (22), suggesting that our understanding of the effects of rapamycin on CCM-deficient endothelial cells are not complete.

In this study, we investigated the effects of propranolol and rapamycin on *Ccm3*-deficient mBMEC. While neither treatment reversed known effects of CCM deficiency on protein membrane distribution, our unbiased transcriptomic analysis of drug effects revealed a key finding: rapamycin reverses the expression of genes dysregulated by *Ccm3* loss. In stark contrast, propranolol failed to demonstrate any significant effect in reversing the *Ccm3*-deficient phenotype, even under unbiased RNA-seq analysis. This observation suggests that if propranolol exhibits therapeutic effects in CCMs, it is unlikely to be through directly reversing the CCM phenotype in endothelial cells.

Importantly, we also present novel evidence that rapamycin, in combination with the tyrosine kinase inhibitor lapatinib, significantly reduces lesion volume in a chronic model of cavernomas. This finding underscores the potential of drug combinations as a promising therapeutic strategy for cavernomas, warranting further investigation.

Methods

Protein immunoblotting and immunofluorescence were performed as detailed in supplementary methods.

Mouse experiments

All procedures were performed in accordance with the European Parliament and of the Council Directive 2010/63/EU and Spanish legislation (RD 53/2013) and were approved by the Ethics Committee on Animal Welfare of University of Santiago de Compostela (reference 15010/19/003). For analysis of cavernomas or isolation of mBMEC, animals were euthanized by CO_2 inhalation followed by cervical dislocation. All experiments involving animals were performed following the ARRIVE guidelines.

All mice used in this study, both for the isolation of brain endothelial cells and for *in vivo* treatments, were maintained in a C57BL/6J background from those developed by

Elisabetta Dejana laboratory crossing *Pdcd10*^{fl/fl} mice (8) with *cdh5*(PAC)-CreERT2 mice (23), as described in (8). Endothelial-specific *Ccm3*-null mice were obtained after treatment with tamoxifen (Merck Life Sciences #T5648) as specified in supplemental methods. Number of mice used in each experimental group are outlined in figure legends.

Procedures for cavernoma induction, drug administration and assessment of lesions are detailed in supplemental methods.

Isolation of mBMEC and treatments

mBMEC were isolated from adult 8–10-week *Cdh5* (PAC)-CreERT2/*Pdcd10*^{fl/fl} mice. Mice were euthanized and brain tissue harvest and tissue homogenization and digestion performed as described by Czupalla (24). Then cells were resuspended and cultured as described by Assmann with slight modifications as described in supplementary methods (25).

RNA extraction, sequencing and data analysis

RNA was handled, and the bioinformatic analysis of RNA-seq performed, as described in detail in supplemental methods.

Data and materials availability:

The data underlying this article will be shared at reasonable request to the corresponding authors. The RNA-seq data is available in the NCBI GEO repository (accession GSE298723).

Please see the Major Resources Table in the Supplemental Materials

Sex as a biological variable

Male and female mice were used in the experiments, as no sex differences have been reported in the development of cavernomas.

Statistics

Statistical analysis was performed using GraphPad software (https://www.graphpad.com/), version 7.0, San Diego, CA, applying the t-Test when 2 groups were compared, and ANOVA analysis with a Tukey's multiple comparison test when comparing >2 groups. For cavernoma volumes in figures 5 and 6, Welch's ANOVA test followed by a Dunnett´s T3 multiple comparisons test was used, based on unequal variances of the different groups. A value of $p<0.05$ was considered significant. All graphs represent the mean ± SEM with dots indicating individual values.

Results

To investigate the effects of drugs proposed for cavernoma treatment, we used mouse brain microvascular endothelial cells (mBMEC), a cellular model that closely mimics the capillary/venous endothelium from which these lesions originate. We employed mice carrying a floxed *Ccm3* allele (26) that express a tamoxifen-inducible cre recombinase (*cdh5*-creERT2) (23). To generate *Ccm3*-deficient mBMEC, we inactivated *Ccm3* in endothelial cells at passage 1 by treating the cultures with 4-OH-tamoxifen (4-OHT) for 4 days (27), which significantly reduced levels of *Ccm3* mRNA and protein compared to untreated cells (figure S1A and B). These *Ccm3*-deficient cells displayed characteristic effects of CCM deficiency, including elevated *Klf2* and *Klf4* mRNA levels (figure S1C) and reduced junctional levels of VE-cadherin and β-catenin (figure S1D). The presence of these established CCM-related phenotypes in our *Ccm3*-deficient mBMEC validates the suitability of this model for further investigating the effects of propranolol and rapamycin on CCM pathogenesis.

To further characterize the effects of *Ccm3* deficiency in mBMEC, we subjected cells to RNA-seq analysis. *Ccm3* inactivation deregulated 168 genes (figure S2A and table S1). Analysis of the differentially expressed genes (figure S2B) showed enrichment in biological processes that have been reported to be related to cavernomas, such as angiogenesis, cell adhesion, cell migration or extracellular matrix organization (28), which confirms that mBMEC are a good model to study the effects of CCM deficiency.

We sought to elucidate the specific effects of propranolol and rapamycin within this defined cellular model of CCM. We treated cells for 24 hours with 10 µM propranolol, based on its plasma levels in treated hypertensive patients and the distribution of the drug in brain tissue (29, 30). For rapamycin, we treated cells for 24 hours at a 100 nM concentration, based on the peak and trough levels attained in plasma after mice dosing (31-33). First, we analyzed the localization of VE-cadherin and β-catenin. VE-cadherin distribution was not affected by treatments in *Ccm3* KO cells, while a 24-hour exposure to rapamycin diminished its junctional localization in WT cells (figure 1A). β-catenin distribution to the cell junctions was also not affected by treatments in *Ccm3* KO cells, but, surprisingly, both drugs reduced its amount in wild-type cells after a 24-hour exposure, making them more similar to *Ccm3*-deficient cells at least in this regard (figure 1B). We concluded that neither rapamycin nor propranolol recovered the abnormal distribution of VE cadherin or β -catenin in *Ccm3*-deficient mBMEC.

To study the effect of both treatments in an unbiased manner, we performed an RNA-seq analysis of *Ccm3* KO cells after treatment and analyzed their effect on the 168 genes that were differentially expressed in KO vs WT cells, which reflected the effect of *Ccm3* loss on brain endothelial cells. Treatment with propranolol affected the expression of several of these genes (figure 2A). However, when a principal component analysis (PCA) was performed comparing WT, untreated and treated KO cells, the effect of propranolol was

shown not to revert the effects of *Ccm3* deficiency but acted along a different component (figure 2B). In contrast, rapamycin partially recovered the displacement induced by the loss of *Ccm3*, suggesting a normalization of gene expression and thus a possible therapeutic effect in these cells (figure 2D). This is also reflected in the heatmap of the DEGs, where KO cells treated with rapamycin cluster apart from untreated KO cells (figure 2C).

In-depth analysis of the genes recovered by rapamycin showed that they included some of the genes known to be dependent on *Klf2/4* transcription factors, which are deregulated in CCM deficient cells. This included genes such as *Adamts1*, overexpressed in CCM-deficient cells (34, 35), or *Thbs1*, downregulated in CCM-deficient deficient cells (28). To confirm this effect, we performed RT-qPCRs in cell preparations different from those used for RNA-seq. *Adamts1* was confirmed as overexpressed and inhibited by rapamycin in *Ccm3* KO mBMEC (figure 3A), while *Thbs1* was downregulated in KO cells, with rapamycin recovering it (figure 3B). *Nos3*, a gene regulated by *Klf2/4* and important in CCM pathogenesis(36), was also overexpressed in KO cells and inhibited by rapamycin when checked by RT-qPCR (figure 3C). On the contrary, *Ccm3*, *Klf2* and *Klf4* were not affected by rapamycin treatment in *Ccm3* KO cells (figures 3D, E and F), which is consistent with its described effects in other CCM-deficient endothelial cells(14). Contrary to the effects of rapamycin, and in accordance to the RNA-seq results, propranolol did not influence the expression of any of these genes.

Overexpression of *Nos3* is one of the hallmarks of cavernous malformations, and it is important for lesion development through a feedback loop involving NO production by cavernoma endothelial cells and the ensuing secretion of VEGF by astrocytes that contributes to lesion enlargement (36). Thus, we wanted to see if the effect of rapamycin on *Nos3* mRNA was reflected at the protein level. As seen in figure 4A and B, both eNOS

protein levels, and its phosphorylation in serine 1177 were stimulated in *Ccm3*-deficient cells, and this was reverted when they were treated with rapamycin for 24 hours. The same western blot shows that *Ccm3*-deficient cells had a tendency to have higher levels of S6 phosphorylation with no change in Akt phosphorylation in S473, which, together with a higher S6K1 phosphorylation in Serine 371 (figure 4D), is in agreement with the effects of *Ccm1* KO in brain endothelial cells and HuVECs, as reported by Ren et al (14). Rapamycin acted as expected both in WT and KO cells, inhibiting S6 phosphorylation after 3 and 24 hours of treatment, and diminishing AKT phosphorylation in S473 only after 24 hours.

Another important functional consequence of *Ccm3* deficiency in endothelial cells is the abnormal dispersion of the Golgi apparatus, which is linked to their impaired ability to polarize during cell migration (37). Notably, rapamycin partially reversed this dispersion in *Ccm3* KO mBMEC (figure 4C).

The effect of rapamycin on known transcriptional targets of *Klf2* and *Klf4*, together with its lack of effect on the expression of these transcription factors at the mRNA level, hints at a possible post-transcriptional effect of the drug on *Klf2* and/or *Klf4*, probably through the known effects of rapamycin on translation. However, there is no apparent effect on *Klf4*, as its protein levels have been shown not to change with rapamycin in *Ccm1*-deficient endothelial cells, both *in vivo* and *in vitro* (14). Intriguingly, in a different setting, Wang et al have shown that rapamycin inhibits *Klf2* by post-transcriptional mechanisms in endothelial cells (38). When we checked KLF2 protein levels in *Ccm3* KO cells, we saw that rapamycin treatment could diminish KLF2 protein (figure 4D), even if it did not inhibit its mRNA. This shows that rapamycin inhibits KLF2 even without affecting its mRNA levels, and points to a direct mechanism for the effect of the drug on KLF2/4 targets in *Ccm3* KO cells.

The effects of rapamycin in the reversal of alterations induced by the lack of *Ccm3* reinforce the idea that it could be used for treatment both of growing and mature cavernoma lesions, either alone or in combination with other agents. Rapamycin has been shown to prevent the development of cavernomas using animal models of the disease, both in lesions with only mutations in a CCM gene and in those harboring also *Pik3ca* mutations (14). However, concerns have been raised on these results based on the high doses of rapamycin used, which may have achieved supratherapeutic levels that cannot be safely tolerated in humans (33). The trough levels of rapamycin that are usually targeted in plasma for benign conditions are 5-15 ng/mL (33, 39). Thus, we decided to induce cavernoma in neonatal *Cdh5* (PAC)-CreERT2/*Pdcd10*$^{fl/fl}$ mice (23) and administer 1.5 mg/Kg rapamycin intraperitoneally, a lower dose than the 2 mg/Kg used by Bishu *et al* to attain trough levels of 13.8 ng/mL (39) (figure 5A). Under this regime, rapamycin could inhibit the development of cavernoma lesions, although it did not completely prevent them, as happens when higher doses are used (14, 21, 40) (figure 5B and 5C).

We then considered the possibility of using rapamycin in combination with other agents for the treatment of cavernomas. Since propranolol had no apparent effect in our cellular system, we turned to the HER2/EGFR inhibitor lapatinib, as we had previously shown that *Ccm3*-deficient endothelial cells have higher levels of EGF receptor, and are more susceptible to apoptosis after its inhibition with lapatinib *in vitro* (13) . Further, it is known that high levels of PI3K activity in tumors make them resistant to EGF receptor inhibition and rapamycin enhances the effect of EGFR inhibition in breast cancer cells (41). Treatment with 300 mg/Kg lapatinib alone did not significantly prevent cavernoma development. However, the combined rapamycin/lapatinib treatment resulted in a trend towards lower volume of the cerebellum occupied with lesions as compared with the already low volume of cavernomas in mice treated with rapamycin alone (figure 5B and

5C). This hinted at the possibility that it could be a good strategy to inhibit the EGF receptor family in combination with rapamycin treatment for the handling of cavernomatosis, and that it could have a greater effect in situations where rapamycin alone is less effective, such as cavernomas that have already developed.

Being able to inhibit lesions that have already developed would be an optimal strategy for treatment of cavernomas. Unfortunately, treatment of chronic lesions with pharmacologically relevant doses of rapamycin does not result in lesion improvement (22). We wanted to see if the addition of lapatinib could help reduce lesion volume of more mature lesions. We used a model of chronic cavernoma developed for the same animals used for acute cavernoma development by Malinverno *et al* (42), and treated them with an injection of rapamycin alone or in combination with lapatinib every 24 hours from postnatal day 30 (P30) to P39, measuring cavernoma volume at P40 (figure 6A). Consistent with recent reports (22), rapamycin alone did not affect the evolution of lesions in this model but, surprisingly, combined treatment of rapamycin and lapatinib had an effect on the volume occupied by lesions in these animals (figure 6B), suggesting that the combination of rapamycin with inhibition of tyrosine kinase receptors may be especially useful to treat established cavernomas.

Discussion

We have investigated the effects of two proposed cavernoma treatments, rapamycin and propranolol, on the biology of *Ccm3*-deficient brain endothelial cells. To ensure an unbiased approach, we analyzed the impact of both drugs on the transcriptomic signature of *Ccm3* KO brain endothelial cells, assessing their ability to revert it toward that of WT cells. Using this strategy, we discovered that propranolol exhibits no discernible effect on the transcriptional phenotype induced by *Ccm3* loss in our model with the doses and

treatment time used. In contrast, rapamycin significantly reverses the transcriptomic signature triggered by *Ccm3* deficiency.

Rapamycin was first proposed as a possible treatment for cavernomas by Ren et al (14), based on their groundbreaking discovery of activating *Pik3ca* mutations in cavernomas and the observation that *Ccm1* loss or *KLF4* overexpression increased mTORC1 activity in cavernoma lesions and HuVECs, an effect which has been described as mediated by caveolin 1(21). Our cellular model effectively recapitulates the effects of CCM deficiency on mTOR activity in endothelial cells. Importantly, despite the absence of *Pik3ca* activating mutations, we observed signs of enhanced mTOR activity in *Ccm3*-deficient mBMEC compared to *Ccm3* WT cells, as evidenced by S6K1 phosphorylation and a trend towards S6 phosphorylation, while AKT phosphorylation remained unaffected. These effects mirror those resulting from *Ccm1* loss (14).

Using this model, we show that rapamycin significantly impacts the expression of genes known to be upregulated in CCM-deficient endothelial cells, such as *Nos3*, *Adamts1* and *Thbs1*. Moreover, the deregulated expression of these genes in KO cells is known to be dependent on the transcription factors *Klf2* and *Klf4*. Importantly, we also show that rapamycin, despite not affecting its mRNA levels, diminishes KLF2 protein. This suggests a mechanism for the restoration of expression of *Klf2/4* transcriptional targets in *Ccm3* KO cells. This specific effect on KLF2 is particularly noteworthy, as rapamycin is known not to affect KLF4 protein levels (14). Interestingly, while rapamycin treatment influences several pathways altered in cavernomas, it does not affect them all. Specifically, rapamycin does not revert the altered distribution of β-catenin or VE cadherin in the membrane of *Ccm3*-deficient cells, suggesting these alterations are not dependent on KLF2 overexpression.

The transcriptional effects of rapamycin have important functional consequences. *Nos3* overexpression and the ensuing overproduction of NO in CCM-deficient endothelial cells is known to be important in cavernoma pathogenesis (36), whereas inhibition of *Thbs1* also favors cavernoma development (28). We show that rapamycin can diminish the amount of the *Nos3* gene product, eNOS and its phosphorylation, and can revert Golgi dispersion in CCM-deficient cells. This underscores the potential of rapamycin as a component of a pharmacological strategy to treat cavernomas, even in the absence of *PIK3CA* mutations.

Importantly, in this study we provide compelling evidence that rapamycin, when combined with a drug possessing a distinct mechanism of action such as lapatinib, can effectively improve chronic cavernoma lesions. With the cavernoma model developed by Malinverno et al. (42), we can begin treatment at postnatal day 30, 28 days after lesion induction, when cavernomas are known to be already detectable. This suggests that the combination of rapamycin and lapatinib, but not rapamycin alone, may improve cavernomas after they have already formed. A common characteristic of all proposed cavernoma treatments that have failed in clinical trials to date is their efficacy in preventing cavernoma development, rather than in improving existing cavernomas. However, in clinical settings, cavernomas are typically already detectable upon diagnosis. The activity of the rapamycin/lapatinib combination in improving cavernomas after 30 days of development makes this strategy worthy of further pursuit, including refinement in preclinical models and incorporating MRI exams in animals before and after treatment.

In contrast to rapamycin, propranolol exhibits no activity in reverting the phenotype of *Ccm3*-deficient cells in our model, both when analyzed in a directed manner (e.g., β-catenin or VE-cadherin distribution) and in an unbiased manner (RNA-seq analysis). These results diverge from reports demonstrating phenotypic effects of propranolol in

CCM-deficient endothelial cells (18, 20, 43). However, in those experiments, the concentration of propranolol used was significantly higher (50 to 100 µM) and not readily achievable in clinical settings. One possibility is that propranolol has beneficial effects on CCM-deficient endothelium that do not involve reversing the CCM phenotype. Alternatively, the reported effects of propranolol on intracranial hemorrhage or focal neurological deficits in patients and animal models may be mediated by its actions in non-endothelial cells or on a specific subset of endothelial cells within a cavernoma that are not reflected in the mBMEC used in this study.

In summary, our study demonstrates that rapamycin effectively reverses key aspects of CCM3 deficiency in brain endothelial cells and, when combined with lapatinib, significantly reduces lesion volume in a chronic model. This highlights the potential of rapamycin-based combination therapies as a promising strategy for treating established CCM lesions, addressing a critical unmet need in cavernoma therapy.

Acknowledgements

The authors thank Dr Ralf Adams for generously sharing the (*Cdh5*) CreERT2 allele and Elisabetta Dejana for generously sharing the *Pdcd10*^fl/fl allele and the protocol for mBMEC isolation and culture. We also thank the Galicia Supercomputing Centre (CESGA) for the availability of computational resources.

The graphical abstract and schemes in figures 5 and 6 were created with Biorender.com.

Sources of funding

This work was supported by projects PID2021-123365OB-I00, financed by MCIN/AEI/10.13039/501100011033/ and by "ERDF A way of making Europe", and ED431C 2023/10 financed by the Consellería de Cultura, Educación e Ordenación Universitaria, Xunta de Galicia (to C-M-P. and J.Z); and projects PID2020 119486RB-

100 funded by MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe" and PID2023-152685OB-I00 funded by MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe", by the "European Union" (to M.V.-R). M. García -Colomer was a predoctoral fellow from Xunta de Galicia. The fellowship to E.M.E.R is supported by PRE2021-099788 funded by MCIN/AEI/ 10.13039/501100011033 and "ESF Investing in your future", to C.R.S MU-21-UP2021-03071902373A.

Disclosures

The authors declare they have no conflict of interest.

Supplemental materials

Figures S1 and S2.
Table S1
Supplemental methods.
Major resources table.

References

1. Clatterbuck RE, Eberhart CG, Crain BJ, Rigamonti D. Ultrastructural and immunocytochemical evidence that an incompetent blood-brain barrier is related to the pathophysiology of cavernous malformations. Journal of Neurology, Neurosurgery & Psychiatry. 2001;71(2):188–192.

2. Fischer A, Zalvide J, Faurobert E, Albiges-Rizo C, Tournier-Lasserve E. Cerebral cavernous malformations: From CCM genes to endothelial cell homeostasis. Trends in Molecular Medicine. 2013;19(5):302–308.

3. Maraire JN, Awad IA. Intracranial cavernous malformations: Lesion behavior and management strategies. Neurosurgery;37(4):591–605.

4. Labauge P, Denier C, Bergametti F, Tournier-Lasserve E. Genetics of cavernous angiomas. Lancet Neurol. 2007;6:237–244.

5. Zhou Z, Tang AT, Wong W, Bamezai S, Goddard LM, Shenkar R, Zhou S, Yang J, Wright AC, Foley M, Arthur JSC, Whitehead KJ, Awad IA, Li DY, Zheng X, Kahn ML.

Cerebral cavernous malformations arise from endothelial gain of MEKK3–KLF2/4 signalling. Nature. 2016;532(7597):122–126.

6. Nishimura S, Mishra-Gorur K, Park J, Surovtseva YV, Sebti SM, Levchenko A, Louvi A, Gunel M. Combined HMG-COA reductase and prenylation inhibition in treatment of CCM. Proc Natl Acad Sci U S A. 2017;114(21):5503–5508.

7. Shenkar R, Shi C, Austin C, Moore T, Lightle R, Cao Y, Zhang L, Wu M, Zeineddine HA, Girard R, Mcdonald DA, Rorrer A, Gallione C, Pytel P, Liao JK, Marchuk DA, Awad IA. RhoA kinase inhibition with fasudil versus simvastatin in murine models of cerebral cavernous malformations. Stroke. 2017;48(1):184–194.

8. Bravi L, Rudini N, Cuttano R, Giampietro C, Maddaluno L, Ferrarini L, Adams RH, Corada M, Boulday G, Tournier-Lasserve E, Dejana E, Lampugnani MG. Sulindac metabolites decrease cerebrovascular malformations in CCM3-knockout mice. Proc Natl Acad Sci U S A . 2015;112(27):8421–8426.

9. Wüstehube J, Bartol A, Liebler SS, Brütsch R, Zhu Y, Felbor U, Sure U, Augustin HG, Fischer A. Cerebral cavernous malformation protein CCM1 inhibits sprouting angiogenesis by activating DELTA-NOTCH signaling. Proc Natl Acad Sci U S A. 2010;107(28):12640–12645.

10. Alfa R, Considine T, Virani S, Pfeiffer M, Donato A, Dickerson D, Shuster D, Ellis J, Rushton K, Wei H, Gibson C. Clinical pharmacology and tolerability of REC-994, a redox-cycling nitroxide compound, in randomized phase 1 dose-finding studies. Pharmacology Res & Perspec. 2024;12(3):e1200.

11. Tang AT, Choi JP, Kotzin JJ, Yang Y, Hong CC, Hobson N, Girard R, Zeineddine HA, Lightle R, Moore T, Cao Y, Shenkar R, Chen M, Mericko P, Yang J, Li L, Tanes C, Kobuley D, Võsa U, Whitehead KJ, Li DY, Franke L, Hart B, Schwaninger M, Henao-Mejia J, Morrison L, Kim H, Awad IA, Zheng X, Kahn ML. Endothelial TLR4 and the microbiome drive cerebral cavernous malformations. Nature. 2017;545(7654):305–310.

12. Otten C, Knox J, Boulday G, Eymery M, Haniszewski M, Neuenschwander M, Radetzki S, Vogt I, Hähn K, De Luca C, Cardoso C, Hamad S, Igual Gil C, Roy P, Albiges-rizo C, Faurobert E, Von Kries JP, Campillos M, Tournier-lasserve E, Derry WB, Abdelilah-seyfried S. Systematic pharmacological screens uncover novel pathways involved in cerebral cavernous malformations. EMBO Mol Med. 2018;10(10):e9155.

13. Sartages M, Floridia E, García-Colomer M, Iglesias C, Macía M, Peñas P, Couraud P, Romero IA, Weksler B, Pombo CM, Zalvide J. High levels of receptor tyrosine kinases in CCM3-deficient cells increase their susceptibility to tyrosine kinase inhibition. Biomedicines. 2020;8(12):624.

14. Ren AA, Snellings DA, Su YS, Hong CC, Castro M, Tang AT, Detter MR, Hobson N, Girard R, Romanos S, Lightle R, Moore T, Shenkar R, Benavides C, Beaman MM, Müller-Fielitz H, Chen M, Mericko P, Yang J, Sung DC, Lawton MT, Ruppert JM, Schwaninger M, Körbelin J, Potente M, Awad IA, Marchuk DA, Kahn ML. PIK3CA and CCM mutations fuel cavernomas through a cancer-like mechanism. Nature. 2021;594(7862):271–276.

15. Mabray MC, Caprihan A, Nelson J, McCulloch CE, Zafar A, Kim H, Hart BL, Morrison L. Effect of simvastatin on permeability in cerebral cavernous malformation type 1 patients: Results from a pilot small randomized controlled clinical trial. Transl Stroke Res. 2020;11(3):319–321.

16. Awad IA, Alcazar-Felix RJ, Stadnik A, Kinkade S, Jhaveri A, Lee J, Hage S, Iqbal J, Polster SP, Shenkar R, Treine K, Mcbee N, Ostapkovich N, Lane K, Liao JK, Sorrentino M, Lee C, Flemming KD, Girard R, Carroll TJ, Thompson RE, Hanley DF, Alcazar-Felix J, Ms S, Bs J, Bsn JL, Ba T, Mph M, Ccrp L, Hanley F, Md F. Safety and efficacy of atorvastatin for rebleeding in cerebral cavernous malformations (AT CASH EPOC): A phase 1/2a, randomised placebo-controlled trial. The Lancet Neurology. 2025;24(4):295–304.

17. Léauté-Labrèze C, Dumas de la Roque E, Hubiche T, Boralevi F, Thambo J, Taïeb A. Propranolol for severe hemangiomas of infancy. N Engl J Med. 2008;358(24):2649–2651.

18. Li W, Shenkar R, Detter MR, Moore T, Benavides C, Lightle R, Girard R, Hobson N, Cao Y, Li Y, Griffin E, Gallione C, Zabramski JM, Ginsberg MH, Marchuk DA, Awad IA. Propranolol inhibits cavernous vascular malformations by β1 adrenergic receptor antagonism in animal models. Journal of Clinical Investigation. 2021;131(3):e154909.

19. Lanfranconi S, Scola E, Meessen JMTA, Pallini R, Bertani GA, Al-Shahi Salman R, Dejana E, Latini R, Abete Fornara G, Agnelli NM, Albanese A, Awad I, Bagnati R, Balconi G, Ballabio E, Beghi E, Bernasconi R, Besana S, Blanda A, Bossi C, Bresolin N, Buratti MG, Calabrese R, Carriero MR, Castori M, Ciceri EF, Ciurleo R, Comi GP, Contarino V, Conte G, Dagruma L, alessandris GQ, De Grazia U, Di Bonaventura R, orio P, Farago G, Foresta A, Fusco C, Gaudino C, Lampugnani MG, Lanno A, Lazzaroni F, Lee C, Locatelli M, Maggioni AP, Magnusson P, Malinverno M, Mangiavacchi M, Mangraviti A, Marino S, Mazzola S, Nicolis EB, Novelli D, Ojeda Fernandez ML, Petracca A, Pignotti F, Pogliani S, Poloni M, Prelle A, Raggi P, Raucci F, Regna-Gladin C, Ronchi D, Scelzo E, Seyfried S, Simeone A, Sturiale CL, Tassi L, Tettamanti M, Torri V, Tournier-Lasserve E, Treglia R, Triulzi FM, Ungaro C, Ursi E, Valcamonica G, Vasami', Antonella, Zarino B. Safety and efficacy of propranolol for treatment of familial cerebral cavernous malformations (Treat_CCM): A randomised, open-label, blinded-endpoint, phase 2 pilot trial. The Lancet Neurology. 2022;22(1):35–44.

20. Oldenburg J, Malinverno M, Globisch MA, Maderna C, Corada M, Orsenigo F, Conze LL, Rorsman C, Sundell V, Arce M, Smith RO, Yau ACY, Billström GH, Mägi CÖ, Beznoussenko GV, Mironov AA, Fernando D, Daniel G, Olivari D, Fumagalli F, Lampugnani MG, Dejana E, Magnusson PU. Propranolol reduces the development of lesions and rescues barrier function in cerebral cavernous malformations. Stroke. 2021;52(4):1418–1427.

21. Min W, Qin L, Zhang H, López-Giráldez F, Jiang N, Kim Y, Mohan VK, Su M, Murray KN, Grutzendler J, Zhou JH. mTORC1 signaling in brain endothelial progenitors contributes to CCM pathogenesis. Circulation Research. 2024;135(4):e94–e113.

22. Alcazar-Felix RJ, Shenkar R, Benavides CR, Bindal A, Srinath A, Li Y, Kinkade S, Terranova T, Debose-Scarlett E, Lightle R, Debiasse D, Almazroue H, Cruz DV,

Romanos S, Jhaveri A, Koskimäki J, Hage S, Bennett C, Girard R, Marchuk DA, Awad IA. Except for robust outliers, rapamycin increases lesion burden in a murine model of cerebral cavernous malformations. Transl Stroke Res. 2024;16(3):859–867.

23. Wang Y, Nakayama M, Pitulescu ME, Schmidt TS, Bochenek ML, Sakakibara A, Adams S, Davy A, Deutsch U, Lüthi U, Barberis A, Benjamin LE, Mäkinen T, Nobes CD, Adams RH. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. Nature. 2010;465(7297):483–486.

24. Czupalla CJ, Yousef H, Wyss-Coray T, Butcher EC. Collagenase-based single cell isolation of primary murine brain endothelial cells using flow cytometry. BIO-PROTOCOL. 2019;8(22):e3092.

25. Assmann JC, Müller K, Wenzel J, Walther T, Brands J, Thornton P, Allan SM, Schwaninger M. Isolation and cultivation of primary brain endothelial cells from adult mice. Bio-protocol. 2017 May 20;7(10):1–16.

26. Maddaluno L, Rudini N, Cuttano R, Bravi L, Giampietro C, Corada M, Ferrarini L, Orsenigo F, Papa E, Boulday G, Tournier-Lasserve E, Chapon F, Richichi C, Retta SF, Lampugnani MG, Dejana E. EndMT contributes to the onset and progression of cerebral cavernous malformations. Nature. 2013;498(7455):492–496.

27. Sartages M, García-Colomer M, Iglesias C, Howell BW, Macía M, Peña P, Pombo CM, Zalvide J. GCKIII (germinal center kinase III) kinases STK24 and STK25 (serine/threonine kinase 24 and 25) inhibit cavernoma development. Stroke. 2022;53(3):976–986.

28. Lopez-Ramirez MA, Fonseca G, Zeineddine HA, Girard R, Moore T, Pham A, Cao Y, Shenkar R, De Kreuk B, Lagarrigue F, Lawler J, Glass CK, Awad IA, Ginsberg MH. Thrombospondin1 (TSP1) replacement prevents cerebral cavernous malformations. Journal of Experimental Medicine. 2017;214(11):3331–3346.

29. Wong L, Nation RL, Chiou WL, Mehta PK. Plasma concentrations of propranolol and 4-hydroxypropranolol during chronic oral propranolol therapy. Br J Clin Pharmacol. 2015;8(2):163–167.

30. Myers MG, Lewis PJ, Reid JL, Dollery CT. Brain concentration of propranolol in relation to hypotensive effect in the rabbit with observations on brain propranolol levels in man. J Pharmacol Exp Ther. 1975;192(2):327–335.

31. Granville CA, Warfel N, Tsurutani J, Hollander MC, Robertson M, Fox SD, Veenstra TD, Issaq HJ, Linnoila RI, Dennis PA. Identification of a highly effective rapamycin schedule that markedly reduces the size, multiplicity, and phenotypic progression of tobacco Carcinogen–Induced murine lung tumors. Clinical Cancer Research. 2007;13(7):2281–2289.

32. Banerjee S, Gianino SM, Gao F, Christians U, Gutmann DH. Interpreting mammalian target of rapamycin and cell growth inhibition in a genetically engineered mouse model of Nf1-deficient astrocytes. Molecular Cancer Therapeutics. 2011;10(2):279–291.

33. Hagan MJ, Shenkar R, Srinath A, Romanos SG, Stadnik A, Kahn ML, Marchuk DA, Girard R, Awad IA. Rapamycin in cerebral cavernous malformations: What doses to test in mice and humans. ACS Pharmacol Transl Sci. 2022;5(5):266–277.

34. Choi JP, Wang R, Yang X, Wang X, Wang L, Ting KK, Foley M, Cogger V, Yang Z, Liu F, Han Z, Liu R, Baell J, Zheng X. Ponatinib (AP24534) inhibits MEKK3-KLF signaling and prevents formation and progression of cerebral cavernous malformations. Sci Adv. 2018;4(11):eaau0731.

35. Yang X, Wu S, Gao R, Wang R, Wang Y, Dong Z, Wang L, Qi C, Wang X, Schmitz ML, Liu R, Han Z, Zheng X. Release of STK24/25 suppression on MEKK3 signaling in endothelial cells confers cerebral cavernous malformation. JCI Insight. 2023;8(5):e160372.

36. Lopez-Ramirez MA, Lai CC, Soliman SI, Hale P, Pham A, Estrada EJ, Mccurdy S, Girard R, Verma R, Moore T, Lightle R, Hobson N, Shenkar R, Poulsen O, Haddad GG, Daneman R, Gongol B, Sun H, Lagarrigue F, Awad IA, Ginsberg MH. Astrocytes propel neurovascular dysfunction during cerebral cavernous malformation lesion formation. Journal of Clinical Investigation. 2021;131(13):e139570.

37. Fidalgo M, Fraile M, Pires A, Force T, Pombo C, Zalvide J. CCM3/PDCD10 stabilizes GCKIII proteins to promote golgi assembly and cell orientation. Journal of Cell Science. 2010;123(8):1274–1284.

38. Wang Y, Li Q, Zhang Z, Peng K, Zhang D, Yang Q, Passerini AG, Simon SI, Sun C. mTOR contributes to endothelium-dependent vasorelaxation by promoting eNOS expression and preventing eNOS uncoupling. Commun Biol. 2022;5(1):726.

39. Bishu K, Ogut O, Kushwaha S, Mohammed SF, Ohtani T, Xu X, Brozovich FV, Redfield MM. Anti-remodeling effects of rapamycin in experimental heart failure: Dose response and interaction with angiotensin receptor blockade. PLoS ONE. 2013;8(12):e81325.

40. Li L, Ren AA, Gao S, Su YS, Yang J, Bockman J, Mericko-Ishizuka P, Griffin J, Shenkar R, Alcazar R, Moore T, Lightle R, Debiasse D, Awad IA, Marchuk DA, Kahn ML, Burkhardt J. mTORC1 inhibitor rapamycin inhibits growth of cerebral cavernous malformation in adult mice. Stroke. 2023;54(11):2906–2917.

41. Liu Q, Yu S, Zhao W, Qin S, Chu Q, Wu K. EGFR-TKIs resistance via EGFR-independent signaling pathways. Mol Cancer. 2018;17(1):53.

42. Malinverno M, Maderna C, Abu Taha A, Corada M, Orsenigo F, Valentino M, Pisati F, Fusco C, Graziano P, Giannotta M, Yu QC, Zeng YA, Lampugnani MG, Magnusson PU, Dejana E. Endothelial cell clonal expansion in the development of cerebral cavernous malformations. Nat Commun. 2019;10(1):2761.

43. Lopes-coelho F, Nunes S, Martins F, Hipólito A, Gouveia-fernandes S, Domingues G, Melo BF, Sacramento JF, Conde SV, Pereira SA, Vinhais S, Salgado D, Serpa J. Propranolol therapy for cerebral cavernous malformations. World Acad Sci J. 2022;4(3):1–9.

Highlights

- Rapamycin can partially revert the transcriptomic phenotype of CCM3-deficient brain endothelial cells.

- Rapamycin inhibits Klf2 protein levels in CCM3-deficient endothelial cells.

- Combination of rapamycin and lapatinib can reduce lesion volume in a chronic model of cerebral cavernomas.

Figure 1. Rapamycin or propranolol do not recover abnormal distribution of VE-cadherin or β-catenin in *Ccm3* KO mBMEC. **A**. Distribution of VE-cadherin. WT or KO mBMEC were either left untreated or treated with 10 μM propranolol for 24 h, 100 nM rapamycin for 24 h, or 100 nM rapamycin for 3 h, and an immunofluorescence performed for VE-cadherin, counterstaining with DAPI. Upper panels: representative photographs of the immunofluorescence with staining for VE-cadherin (green) and DNA (blue). Scale bar is 50 μm. Graph shows the ratio of junctional/total VE-cadherin after each treatment, referenced to untreated WT mBMEC, of >100 cells per treatment from 4 independent biological replicates (cell cultures). Each cell culture was obtained from a pool of 3 to 4 brains. P-values are from ANOVA analysis with a Tukey's multiple comparison test. **B**. Distribution of β-catenin in WT and KO mBMEC. Upper panels: representative photographs of the immunofluorescence with staining for β-catenin (green) and DNA (blue). Scale bar is 50 μm. Graph shows the ratio of junctional/total β-catenin after each treatment, referenced to untreated WT mBMEC, of >100 cells per treatment from 4 independent biological replicates (cell cultures). Each cell culture was obtained from a pool of 3 to 4 brains. P-values are from ANOVA analysis with a Tukey's multiple comparison test.

Figure 2. Rapamycin, but not propranolol, can partially revert the transcriptional effect of *Ccm3* inactivation in mBMEC. WT and KO mBMEC were left untreated or treated with 10 µM propranolol for 24 h (KO+P) or 100 nM rapamycin for 24 h (KO+R). RNAseq analysis was performed, and their expression profile was compared using genes differentially expressed in KO vs WT mBMEC. Three independent biological replicates (cell cultures) were analyzed in all cases. Each cell cultures was obtained from a pool of 3 to 4 brains. **A.** Heatmap analysis of WT, KO, and KO+P, showing that KO+ P cells cluster with KO cells. **B.** Principal component analysis of WT, KO, and KO+P mBMEC, showing a displacement of KO vs WT mBMEC in the horizontal axis that is not reverted by treatment with propranolol. **C.** Heatmap analysis of WT, KO, and KO+R, showing that treated KO+R mBMEC no longer cluster with KO mBMEC. **D.** Principal component analysis of WT, KO, and KO+R mBMEC, showing a displacement of KO WT mBMEC in the horizontal axis that is partially reverted by treatment with rapamycin.

Figure 3. Rapamycin, but not propranolol, reverts the expression of genes dependent on *Klf2/4*, without affecting *Klf2* or *Klf4* mRNA expression. WT or KO mBMEC were either left untreated or treated with 10 µM propranolol for 24 h or 100 nM rapamycin for 24 h. RNA was extracted and RT-qPCR was performed using GAPDH as a reference gene. Each graph shows the average and SEM of mRNA relative expression and the individual values of six independent biological replicates (cell cultures). Each cell culture was obtained from a pool of 3 to 4 brains. P-values are from ANOVA analysis with a Tukey's multiple comparison test. **A.** Expression of *Nos3*, **B.** Expression of *Adamts1*, **C.** Expression of *Thbs1*, **D.** Expression of *Ccm3*, **E.** Expression of *Klf2*, **F.** Expression of *Klf4*.

Figure 4. Effect of rapamycin on eNOS, Golgi dispersion and KLF2 protein levels. **A.** Western blot of mBMEC proteins after treatments. Extracts were prepared from WT and KO mBMEC treated with 100 nM rapamycin for 3 or 24 hours, or with 10 μM propranolol for 24 hours, and western blots performed for total eNOS, eNOS phosphorylated in Ser 1177 (peNOS), phosphorylated S6 (pS6), AKT phosphorylated in Ser 473 (pAKT), total AKT, *Ccm3*, with GAPDH as a loading control. **B.** Average and SEM of the intensity of peNOS, eNOS, pAKT/AKT ratio and pS6 and from four independent biological replicates (cell cultures). Each cell culture was obtained from a pool of 3 to 4 brains. P-values are from ANOVA analysis with a Tukey's multiple comparison test. **C.** Rapamycin treatment reverts Golgi dispersion in *Ccm3*-deficient mBMEC. WT and KO mBMEC were untreated or treated with 100 nM rapamycin for 24 hours and stained for

GM130 to visualize Golgi. Left panels: Photographs of untreated WT cells, and KO cells untreated or treated with rapamycin. Scale bar is 20 μm. Right graph: Golgi fragments per cell of >100 cells per treatment from 3 independent biological replicates (cell cultures). Each cell culture was obtained from a pool of 3 to 4 brains. P-values are from ANOVA analysis with a Tukey's multiple comparison test. **D.** Rapamycin diminishes KLF2 protein levels in *Ccm3* deficient cells. KO and WT mBMEC were left untreated or treated with 100 nM rapamycin for 24 hours. Western blots were performed for KLF2, vinculin and pS6K1. Left panel, representative western blot. Right graph, average and SEM of corrected intensity of KLF2 of western blots from four biological replicates (cell cultures). Each cell culture was obtained from a pool of 3 to 4 brains. P-values are from ANOVA analysis with a Tukey's multiple comparison test.

Figure 5. Rapamycin, but not Lapatinib, can inhibit acute cavernoma development. **A.** Scheme showing the induction and treatment of cavernomas. 50 µg tamoxifen were administered to newborn Cdh5 (PAC)-CreERT2/PDCD10$^{fl/fl}$ mice at P1. Rapamycin and/or Lapatinib were injected at days P2 to P6, and mice were euthanized at P8, brains extracted and analyzed by micro-CT. **B.** Micro-CT images of brains of P8 mice with induced cavernomas untreated, or treated with Lapatinib, Rapamycin, or Lapatinib + Rapamycin. **C.** Volume of cavernomas untreated mice (n=19 mice), or mice treated with Lapatinib (n=10 mice), Rapamycin (n=10 mice), or Lapatinib + Rapamycin (n=10 mice). Total volumes of cavernomas per brain were calculated from micro-CT images. P-values are from Welch's ANOVA test followed by a Dunnett´s T3 multiple comparisons test. Panel A was created with biorender.com.

Figure 6. Rapamycin/Lapatinib combination can diminish cavernoma volume when used after cavernoma development. **A.** Scheme showing the induction and treatment of cavernomas. 2.5 µg tamoxifen were administered to Cdh5 (PAC)-CreERT2/PDCD10$^{fl/fl}$ mice at P2. Rapamycin or Rapamycin + Lapatinib were injected at days P30 to P39, and mice were euthanized at P40, brains extracted and analyzed by micro-CT. **B.** Micro-CT images of brains of P30 mice with induced cavernomas untreated, or treated with Rapamycin or Lapatinib + Rapamycin. **C.** Volume of cavernomas in untreated mice (n= 13 mice), or mice treated with Rapamycin (n=14 mice) or Lapatinib + Rapamycin (n=8 mice). Total volumes of cavernomas per brain were calculated from micro-CT images. P-values are from Welch's ANOVA test followed by a Dunnett´s T3 multiple comparisons test. Panel A was created with biorender.com.