Sexually antagonistic co-evolution at the molecular level: patterns of genetic co-variation predict phenotypic outcomes of mating interactions.

Alma Thorson[1], Rhonda R. Snook[1,+], R. Axel W. Wiberg[1,*,+]

1. Department of Zoology, Stockholm University, Svante Arrhenius väg 18B, Stockholm, 106 91, Sweden.

* Correspondence: Department of Zoology, Stockholm University, Svante Arrhenius väg 18B, Stockholm, 106 91, Sweden. **e-mail:** raw.wiberg@protonmail.com

+ RAWW and RRS should be considered shared senior authors.

Running title: Sexually antagonistic co-evolution at the molecular level

Keywords: population genomics, sexual selection, sexual conflict, mating behaviour

Abstract

Sexually antagonistic co-evolution (SAC) is a potent driver of phenotypic change, with males and females engaged in cycles of co-evolution in which sex-specific trait expression enhances their own fitness at the expense of the other. The cyclical process results in transient mismatches between male and female traits wherin the relative exaggeration of sex-specific phenotypes predicts which sex has the upper-hand during mating interactions. Parallel genetic signatures of exaggeration of SAC traits should be observed, but due to a poor knowledge of genes underlying SAC traits this remains untested, impeding understanding of how SAC shapes genetic variation and the phenotypic consequences. Here we show the first evidence that SAC theory readily extends to the molecular level and classic patterns of phenotypic SAC are recapitulated. We exploit the iconic sex-peptide (SP) network in *Drosophila melanogaster*, which comprises male seminal fluid proteins and female reproductive tract proteins that interact to influence sexually antagonistic female post-mating behaviours. Using genomic data from ~150 populations, we show that genetic differentiation at loci of the male- and female-components of the SP-network loci co-vary as expected under SAC. We use relative levels of genetic differentiation between male- and female- components of the SP-network to infer which sex has the advantage in mating interactions. Mating and survival assays using populations established from wild-derived isofemale lines show populations with inferred male-advantage have reduced female remating and earlier peaks of egg production compared to populations with inferred female-advantage. Thus, relative levels of genetic differentiation in the classic SP-network with known SAC phenotypes accurately predicts outcomes of mating interactions within populations. Our findings shed light how SAC operates at the molecular level, and sets a standard for future investigations.

Introduction

Sexual conflict arises when the fitness optima of a trait shared by the sexes are not aligned (Parker, 1970, 1979; Arnqvist & Rowe, 2005; Perry & Rowe, 2015). Interlocus sexual conflict occurs when different loci in males and females influence the phenotypic value of a trait that depends on contributions from both partners. A classic trait influenced by interlocus sexual conflict is the female remating rate (Parker, 1970, 2020). Males benefit when they can prevent future inseminations by rival males, for example by reducing female receptivity to further matings. In contrast, females may gain fitness, either *via* direct or indirect benefits, from controlling when, how often, and with whom to remate (Arnqvist & Nilsson, 2000; Rundle et al., 2007; Priest, Galloway, et al., 2008; Priest, Roach, et al., 2008; Fromonteil et al., 2023). This does not preclude strong selection on males to court females aggressively, if the opportunity exists for them to accrue benefits from such courting, and thereby increase the overall female mating rate (Arnqvist & Rowe, 2002a; Perry & Rowe, 2012). Such conflicts can result in sexually antagonistic co-evolution (SAC), an escalating arms race between the sexes leading to novel and increasingly exaggerated persistence and resistance traits in males and females (Holland & Rice, 1998; Arnqvist & Rowe, 2002a, 2005; Perry & Rowe, 2015). On average, males and females should be evenly matched in this arms race, making conflicts difficult to detect (Arnqvist & Rowe, 2002a). However, a central prediction of SAC theory is that due to the cyclical nature of SAC, transient mismatches can occasionally result wherein for some populations (or species) either males or females have the upper-hand. Which sex has the advantage can be detected from patterns of relative phenotypic exaggeration of male and female traits (Arnqvist & Rowe, 2002a, 2005).

The most well-documented example of SAC influencing phenotypic traits is among *Gerris* water striders (Arnqvist & Rowe, 2002a, 2002b). Intense mating struggles between the sexes generates strong selection for male persistence and female resistance

morphologies and behaviours (Arnqvist & Rowe, 2002a, 2002b). Across both species and

populations, the relative level of exaggeration between male and female morphologies

predicts the outcome of mating interactions, reflecting which sex has an advantage, in

agreement with SAC theory (Arnqvist & Rowe, 2002a; Perry & Rowe, 2012). Relatively

more exaggerated male morphologies result in, for example, shorter pre-mating struggles,

and higher male mating success (Arnqvist & Rowe, 2002a; Perry & Rowe, 2012). Patterns

of co-variation in antagonistic male and female traits in other taxa also likely reflect SAC

(Bergsten et al., 2001; Bergsten & Miller, 2007; Rönn et al., 2007; Tatarnic & Cassis, 2010;

Perry & Rowe, 2015), making this a compelling hypothesis explaining phenotypic diversity

across several systems. As genetic evolution underlies phenotypic evolution, the logic of

SAC should also extend to the genetic level at loci underlying male and female conflict

traits. That is, these loci (e.g. Swanson & Vacquier, 2002, and reviewed in Sirot et al.,

2015; Rowe et al., 2018) should show patterns of co-varying "molecular exaggeration"

between male- and female- beneficial loci across populations and species. More critically,

the relative degree of "molecular exaggeration" of genes under interlocus sexual conflict

should predict which sex has the upper-hand and therefore fitness-related outcomes of

mating interactions, just as for phenotypic traits.

However, to-date no studies have tested these predictions. This is because the loci

underlying most conflict traits remain unknown (but see Khila et al., 2009; Crumière &

Khila, 2019; Pruvôt et al., 2025). Meanwhile, classes of loci proposed to be associated

with sexual conflicts and SAC, such as genes encoding male seminal fluid proteins

(SFPs), show rapid evolutionary change across species (Swanson & Vacquier, 2002;

Haerty et al., 2007). These SFPs, transferred to females during mating, mediate female

reproductive physiology, manipulating female traits such as receptivity to further mating

attempts, egg production, and survival (Sirot et al., 2015). While SFPs are relatively well-

described genetically with many having known phenotypic functions, their co-evolution with

loci underlying female resistance traits is unexplored and how genetic (co-)variation at these loci impacts fitness outcomes of male-female interactions is unknown. Thus, unlike the progress of the "genomics revolution" in understanding the genetics of ecological adaptations over the past quarter century, studies of the genomics of sexually selected and sexual conflict traits have lagged behind (Wilkinson et al., 2015; Kasimatis et al., 2017; Rowe et al., 2018). Empirical tests of SAC theory require integrating genetic (co-)variation at known SAC loci and the consequence of the co-variation for outcomes of mating interactions across multiple populations and species. Such tests are critical for understanding how many loci, and what proportion of those underlying a given trait, are involved in this ubiquitous process in sexually reproducing taxa. And also how sex-specific selection and co-evolution between males and females shapes patterns of genetic variation across populations and species, and what the fitness consequences of this genetic (co-)variation are (Wilkinson et al., 2015; Kasimatis et al., 2017; Rowe et al., 2018).

We capitalize on the iconic "sex peptide" (SP) network of *Drosophila melanogaster* to explicitly test SAC theory in a molecular framework for the first time. The SP-network includes a male component, the SP protein and a suite of other SFPs, that interact with a female component, "sex-peptide receptor" (SPR) and other proteins expressed in the female reproductive tract (FRT), to influence a wide range of female post-mating behaviours (Kubli & Bopp, 2012; Findlay et al., 2014; Singh et al., 2018; Hopkins & Perry, 2022). SP has been shown to depress the female remating rate, induce a higher ovulation rate, shorten female lifespan, and reduce reproductive success (reviewed in Kubli & Bopp, 2012; Hopkins & Perry, 2022). These traits are likely under sexual conflict and therefore also potentially subject to SAC in *D. melanogaster* (Parker, 1970; Rundle et al., 2007; Priest, Galloway, et al., 2008; Priest, Roach, et al., 2008; Smith et al., 2017; Parker, 2020). However, whether the evolution of *SP* and *SPR* is driven by SAC is controversial (Hopkins

& Perry, 2022; Hopkins et al., 2024). Controversy arises because studies are mixed in

whether SP receipt consistently (across studies) results in reductions in lifespan or lifetime

reproductive success for females (Hopkins & Perry, 2022). Additionally, across species,

molecular evolution of *SP* and *SPR* seems uncorrelated (Findlay et al., 2014; Hopkins et

al., 2024) contrary to SAC theory (Parker, 1970; Arnqvist & Rowe, 2005). However, other

traits influenced by SP, such as female remating rate and oviposition patterns, may be

equally important sources of sexual conflict in natural populations (e.g. Wensing & Fricke,

2018). These traits are frequently ignored in assessment of the role of sexual conflict and

SAC in SP and SPR co-evolution, as are the other loci of the male-and female-

components of the SP-network.

Leveraging the known loci of the male- and female- components of the SP-network,

we use population genomic data across ~150 natural populations of *D. melanogaster*

(Kapun et al., 2021; Nunez et al., 2025) to test co-evolutionary patterns within and

between these components. Our analysis determines the relative molecular exaggeration

of the male- and female- components in natural populations, which we then use to infer the

relative advantage of the two sexes in each population. If SAC is a driver of molecular

evolution at these loci, then the relative molecular advantage of each sex should

predictably influence female post-mating behaviour. We then test this critical prediction

using lab populations established from a resource of isofemale lines established from

natural populations that vary continuously in the degree of predicted advantage to one or

the other sex (Durmaz Mitchell et al., 2025).

Methods

Genetic data

Population genomic data were obtained from the publicly available DEST dataset

which comprises a total of 530 pool-seq samples of ~50 flies from >170 populations world-

wide sampled across years (https://dest.bio/; (Kapun et al., 2021; Nunez et al., 2025). We

ascertained that individual samples from each population can serve as representatives on

the basis of hierarchical Fst analysis of a neutral set of SNPs with poolfstat (Hivert et al.,

2018; Gautier et al., 2024; see "1.1. Hierarchichal Fst analysis" in the Supplementary

Materials). Samples from the same population showed overall low within-population Fst

(Fsg = 0.01, [0.0096, 0.0104]), compared to between-population Fst (Fgt = 0.04 [0.038,

0.045]), in agreement with previous results (Nunez et al., 2025). Thus, we chose a single

pool-seq sample per population for further analysis, favouring samples with highest

coverage, to ensure the highest quality data and estimates. Populations from the ancestral

range in Africa were excluded due to very few samples, different sequencing strategies,

and a large impact on overall diversity and differentiation so that the final set of samples

represented 149 populations (Table S1).

A set of interacting loci that make up the SP-network have been described from

several studies. Here we use the union of loci described in Findlay et al., (2014) and Singh

et al., (2018), because these represent the current most complete understanding of the

system (see also McGeary & Findlay, 2020). Several loci are poorly characterised in this

context, but all have been shown to be necessary for the observation of SP-network

mediated post-mating behaviours (such as the reduced female receptivity to re-mating,

oviposition rates), as well as longevity.

These loci can be divided into "male"- and "female"-components of the SP-network

based on sex-specific reproductive tissue expression (Tables S2 and S3) although this

does not necessarily preclude expression of those loci in the other sex. For example, SPR

is expressed only in the female reproductive tract (FRT) but is expressed in the nervous

system of both sexes (Table S3). We consider two different definitions of the SP-network

loci, depending on the analysis. Where only coding sites are analysed ($\pi N/\pi S$), we limit

ourselves to the coding sequence (CDS) of each gene. In other analyses (Fst, π, and

Tajima's D) we include introns and sites 2,000 bp up-/down-stream of the locus. This is to strike a balance between a focus on the loci of interest while ensuring enough data for accurate population genetic estimates and inclusion of potential regulatory elements. To obtain genomic background estimates as a comparison, we also sampled 100 randomly selected loci from the *D. melanogaster* annotation.

To investigate patterns of genetic diversity and evaluate the role of selection at SP-network loci, population genetic summary statistics were computed (π, Tajima's D, and $\pi N/\pi S$), with npstat (v. 1.0; Ferretti et al., 2013) and SNPGenie (v. 2019.10.31; Nelson et al., 2015). For $\pi N/\pi S$, we summed the mean number of pairwise (non-)synonymous differences and the number of (non-)synonymous sites, across all populations for each gene. We then computed overall $\pi N/\pi S$ for each locus using these sums. To derive confidence intervals we bootstrap sampled populations with replacement and re-computed $\pi N/\pi S$ 100 times. We also computed population-specific Fst values (see "1.2. Definition of population-specific Fst" in the Supplementary Materials) for each locus in the SP-network with poolfstat (v. 2.1.2; Hivert et al., 2018).

Analysis of molecular co-evolution

Population-specific Fst estimates were analysed to characterise patterns of co-variation in Fst between male- and female- components of the SP-network. We use 2-Block Partial Least Squares (2B-PLS) to identify male and female axes that describe the largest amount of co-variance in Fst between male- and female- components of the SP-network. 2B-PLS is well suited for analyses of co-evolution across species or populations (Rohlf & Corti, 2000) and has been employed to study complex multivariate morphologies in the context of SAC (e.g. Arnqvist & Rowe, 2002a, 2002b; Perry & Rowe, 2012). 2B-PLS produces two main reduced axes that describe co-variation between the two blocks, as well as individual loading scores for the components of those blocks on the axes. A

permutation test of rows within the two blocks tests whether the degree of co-variation is greater than expected by chance. This permutation test conserves relationships between the variables (in this case the loci of the SP-network) within blocks, but should break any signal of co-variation across blocks.

Populations are not independent but genetically related to varying degrees. Such relatedness structure could produce correlations between loci due purely to demographic processes, and should therefore be accounted for in analyses (Stone et al., 2011; Castillo, 2017; Chelini et al., 2021; Raas & Dutheil, 2024). 2B-PLS also integrates phylogenetic or relatedness information using a phylogenetic tree (Adams & Felice, 2014). We first computed a relatedness matrix for all populations using BayPass (v. 2.4; (Gautier et al., 2013); see "1.3. Computation of the relatedness matrix" in the Supplementary Materials) and then converted the summarised omega matrix of BayPass to a dendrogram using the neighbour-joining method in the "ape" R package (v. 5.8; (Paradis et al., 2004). Phylogenetically corrected 2B-PLS was used to test for and describe the main axes of co-variation between male and female components of the SP-network. 2B-PLS analysis was performed with the "geomorph" R package (v. 4.0.8; Adams et al., n.d.; Baken et al., 2021). Because the sign of the strongest loadings on both axes was negative (Table S4), we switched the sign of all values on both axes for ease of interpretation and plotting, such that higher values on the 2B-PLS axes indicated higher differentiation at the loci with strongest loadings on these axes. The main axes of co-variation from the 2B-PLS analysis can be further processed into two principle components by PCA, where the first PC describes the absolute degree of genetic differentiation across both male- and female-components of the SP-network. Meanwhile, the second PC axis describes the *relative* degree of genetic differentiation and therefore provides a continuous measure of the degree of mismatch between males and female (Arnqvist & Rowe, 2002a, 2002b).

Behavioural experiments

We set up stock population cages using isofemale line resources available through the DrosEU consortium (Durmaz Mitchell et al., 2025; see "1.4. Isofemale line resources and stock population cages" in the Supplementary Materials). Briefly, we created three replicate population cages using isofemale lines from each of six natural populations (18 cages in total). For each population, we took 15 males and 15 females of each isofemale line, and split them across the three replicate cages. Populations were allowed to mate freely and recombine for 4-9 generations prior to the experiment. Experimental flies were obtained from each population cage using egg laying plates consisting of a molasses, agar, and water mix (in 1L dH_2O: 16.7g agar, 159.7g molasses) set in a small Petri dish lid (4 cm in diameter) and topped with a small amount of yeast paste (in 10mL dH_2O: 3g dried yeast) placed in each cage overnight. 100 eggs from each cage were collected and distributed across 5 vials (20 eggs / vial) containing standard fly medium (see Supplementary Materials). These were left to develop for 9-11 days in an incubator (Panasonic MIR-154-PE, Panasonic Healthcare Co. Ltd., 25º C, 12h:12h L:D). On the days of eclosion, virgin flies were collected from each vial throughout the day, ensuring collected flies were no older than 2 hours post-eclosion and thus virgin. Flies were kept in single-sex groups of up to 10 flies in vials containing standard fly medium for 3 days prior to mating and re-mating assays. Collections of eggs and virgins for mating and re-mating assays were staggered in order to obtain, as far as possible, males of controlled ages. All males were 2-5 days old at the time of mating and re-mating assays.

On the day of the mating assay, a female and male from the same population were loaded into individual vials. We recorded the time at which pairs entered the vial, and both the start and end time of copulation. The assay was run for 2 hours from the time of the last pair entering the vial. Flies that did not mate in that time were discarded (~3-12% of all flies across populations). There was no relationship between the percentage of "failed"

pairings and the relative degree of genetic differentiation (Pearson's correlation: cor = -0.17, t = -0.34, d.f. 4, p = 0.75). Flies that were observed to mate later than 2 hours after pair formation were censored and discarded. Mated males were removed from their vial and females were left to lay eggs for 2 days. A female re-mating assay was performed the second day after the mating assay in which each female was transferred to a new vial and paired with a new male from the same population of approximately equal age to the initial male to isolate the effect of re-mating. We again recorded the time at which pairs entered the vial, and, if re-mated, copulation start and end times. To account for female re-mating refraction, we ran the assay for 3 hours from the time of the last pair entering the vial, pairs that were observed to mate later than 3 hours after pair formation were censored and discarded.

We collected data on two fitness related traits for each female – mortality and offspring production. Mortality was assessed for each female up to 30 days after the first mating day. All vials were checked every day (except weekends) and the date of death recorded for each female. Females that died due to handling error or escaped were censored as were all females that survived past the 30-day cut-off. Throughout the experiment, females were provided with fresh vials of fly medium on days 2, 4, 7, 11, 17, and 24 after the mating assay, producing 7 vials of offspring in total. Total reproductive success was scored by counting offspring from each vial after 14 days, to allow ample time for the majority of offspring to hatch, pupate, and eclose as adults. Total reproductive success, was measured as either the total offspring in vials 2 and 3, or the total offspring across all 7 vials. This was done to keep a maximum amount of data, as some females died during the experiment and did not have an opportunity to lay eggs in later vials. We also measured timing of peak offspring production computing the time, in days, until the vial with the maximum number of offspring. The full experiment was conducted over five

batches to obtain suitable sample sizes and batch was included as a random factor in

analyses.

In a second mortality experiment, experimental flies were collected from each cage

using egg laying plates as above. Upon collection of emerged adults, virgin flies were

distributed across vials in groups of 10 females and 10 males per vial. Vials were then kept

in a controlled temperature room (25º C, 12h:12h L:D) for the duration of the experiment.

Survival of females was checked every day, except weekends, until less than 10% of flies

from each cage were remaining.

See the Supplementary Materials for the details of all statistical analyses.

Results and Discussion

To assess molecular evidence for SAC in the male- and female- components of the

SP-network across *D. melanogaster* populations we first evaluated the overall population

genetic signature of selection at the loci of the SP-network. Comparative work has found

rapid evolution of *Drosophila* SFPs generally, and co-variation in the evolutionary rates

across species among the loci of the SP-network in particular (Haerty et al., 2007; Findlay

et al., 2014; Hopkins et al., 2024). However, *SP* and *SPR* seem to have uncoupled

evolutionary rates across species (Findlay et al., 2014; Hopkins et al., 2024). These

patterns are consistent with either relaxed- or positive- selection, and have not been

evaluated across populations. Selection (whether positive, in favour of new beneficial

alleles, or purifying, against new deleterious mutations) should leave a characteristic signal

of reduced non-synonymous diversity (πN) relative to synonymous diversity (πS),

comparable to other protein-coding genes subject to selection. We find that nucleotide

diversity (π) for SP-network loci is no different from the set of randomly selection loci

(Figure 1A) and $\pi N/\pi S$ values are well below 1 (Figure 1A), patterns of strong evidence of

selection. Combined with the observed rapid evolution across species at SP and other

SFPs (Haerty et al., 2007; Findlay et al., 2014; Hopkins et al., 2024), our results support a process of recurrent selective sweeps. Such a dynamic genomic signature is expected for SAC traits, in which alleles beneficial to one sex rapidly spread, but which are then countered by new alleles beneficial to the other sex.

We next found strong evidence for co-variation in population genetic differentiation between the male- and female- components of the SP-network from analysis of population-specific Fst estimates of each locus (Phylogenetically corrected 2-Block Partial Least Squares (2B-PLS) analysis; D. C. Adams & Felice, 2014; Rohlf & Corti, 2000; r-PLS = 0.44, Z = 4.17, p = 0.001; Figure 1B). The male axis of this co-variation loads most heavily for SP network genes, *SP* and *Sems*, while the female axis loads most heavily for *Esp* and *hdly* (Table S4). There is relatively low loading of *SPR* on the female axis, which is consistent with the weak patterns of covariation between *SP* and *SPR* observed across species (Findlay et al., 2014; Hopkins et al., 2024) (Table S4). In summary, across 149 natural *D. melanogaster* populations, our results suggest ongoing selection at SP-network loci, and natural genetic co-evolution between male- and female- components of the network consistent with SAC. The pattern of co-variation is weak between the most directly interacting *SP* and *SPR*, and strongest between *SP*, *Sems* and other indirectly interacting loci of the female component of the SP-network (*Esp* and *hdly*). These also show the highest correlations in Fst in simple pairwise tests (Figure S1). Our results highlight that the most obvious candidate loci underlying conflict traits may not always be the strongest drivers of SAC, and that a only a subset of the loci underlying a suite of sexually antagonistic traits may be contributing to SAC. By including multiple interacting loci of SP-network, rather than focusing on the most intuitive one from each sex, we are able to detect molecular signatures of co-evolution that may otherwise be elusive.

A central prediction of SAC is that relatively more exaggeration in male- compared to female- traits infer a fitness advantage in sexual conflicts to males and *vice versa*

(Arnqvist & Rowe, 2002a, 2005). We extend this prediction to the molecular level, defining the relative degree of "molecular exaggeration" as the difference in genetic differentiation between male- and female- components of the SP-network, expressed as a continuous measure using the second principle component (PC2) of the male- and female- axes of co-variation derived from the 2B-PLS (Arnqvist & Rowe, 2002a, 2002b; Perry & Rowe, 2012; Figure 1B; Table S5). Higher scores indicate more exaggeration, or greater genetic differentiation in allele frequencies from the overall average, in the male- compared to the female- component, and thus a more derived state, and an inferred advantage to males. If these patterns of genetic co-variation between components of the SP-network, with previously described phenotypic effects (Kubli & Bopp, 2012, but see Hopkins & Perry, 2022), reflect a process of SAC, then populations with an inferred male-advantage should have: lower female remating rates and an earlier peak in offspring production (faster initial rate of offspring production) than populations with an inferred female-advantage. If SP harms females by reducing lifespan, then we also expect shorter total lifespan and lower survival in populations with inferred male advantage.

We tested these predictions using six populations along the axis of relative genetic differentiation (PC2; Figure 1B), representing a continuum from strong inferred male-advantage to strong inferred female-advantage, for which isofemale line resources are available; these represent a small proportion of the original 149 populations (Durmaz Mitchell et al., 2025; Figure 1B, Table S5). As our aim was to investigate population-level phenotypic effects, we reconstructed each of the six populations by combining individuals from all isofemale lines for each population in population cages, with three replicate cages per population, and assayed female remating rates, survival, offspring production per vial, and total offspring number.

Remating rates varied across populations, from 14% (Uman, Ukraine) to 40% (Munich, Germany) (Figure 2A; Table S6; total N = 765, n= 113-140 females per

population). Critically, the variation in remating rates confirmed the predictions from SAC

theory: populations with relatively more genetic differentiation in the male component of

the SP-network, i.e. greater inferred male-advantage, had lower female remating rates

(Figure 2A and 2B). Over 94% of the posterior distribution of the relationship between

relative genetic differentiation and female remating rates was < 0, indicating a clearly

negative relationship, as predicted by SAC theory (Figure 2B). The model in which relative

genetic differentiation (PC2) as the explanatory variable was represented by the degree of

male- and female- advantage was preferred over a model that did not include this

explanatory variable ($ELPD_{w/ term}$ = -399.3 ±14.7, $ELPD_{w/o term}$ = -399.4 ±14.6, ELPD

difference = -0.1 ±0.7). These effects are even stronger when contrasting female remating

rates from populations with the greatest inferred male- and female- advantage, with 99%

of the posterior distribution supporting a lower remating rate for populations with an

inferred male advantage (Figure S2).

Molecular exaggeration of SP-network loci was also associated with patterns of

oviposition as predicted by SAC theory. Male-advantaged populations had peak

oviposition earlier than female-advantaged populations (Figure 3; Figure S3). This resulted

in females from populations with inferred female advantage producing 68% of their total

offspring within the first 7 days after mating, on average, compared to 71% for females

from populations with inferred male advantage. Male induction of faster egg-laying rates

and earlier peaks of offspring production bias paternity to the most recently mated male

and may be counter to female interests insofar as they override female-beneficial

determination of the timing of reproductive investment (i.e. cryptic female choice;

Eberhard, 1996; Crudgington et al., 2005; Arnqvist & Rowe, 2005; Edward et al., 2010).

Contrary to SAC predictions, we found no relationship between molecular

exaggeration at SP network loci and total number of offspring (Figure S4). Similarly, while

there is strong evidence that survival differs between populations (Cox regression survival

analysis; X^2 = 27.03, d.f. = 5, p < 0.001; Figure S5A), there was no evidence that this

varied based on which sex was advantaged (HR = 1.23, Z-ratio = 1.04, p = 0.3). Moreover,

there was no evidence that either female remating status (X^2 = 0.36, d.f. = 1, p = 0.55) or

the interaction between remating status and population (X^2 = 4.66, d.f. = 5, p = 0.46)

influenced survival. Survival results were similar in the second experiment tracking survival

over ~80 days, while also allowing for continuous mating in groups of 10 males and 10

females (Figure S5B).

By using patterns of natural genetic (co-)variation of loci of multiple male- and

female- SP network components across multiple populations, we explicitly test and support

the role of SAC. We show that both the female remating rate and egg production timing

are predictable based on relative molecular exaggeration of SP-network genes. Our

results are in complete agreement with predictions from SAC theory for these traits and

extend SAC theory to the molecular level for the first time. Although Interest in *Drosophila*

SFPs as mediators of sexual conflict arose because of their apparent toxic effects to

females, observed as reduced lifetime reproductive success and survival the role of sexual

conflict in the (co-)evolution of SP and SPR has been questioned (Hopkins & Perry, 2022;

Kubli & Bopp, 2012). This is due mainly inconsistent observations for the expected

reductions in female longevity or lifetime reproductive success, leading to speculation that

these costs may be highly condition or environment dependent, or that these traits are not

sexually antagonistic at all (Hopkins & Perry, 2022). We found no relationship between this

exaggeration on total offspring production or lifespan, suggesting that these traits, in fact,

are not driven by SAC at these loci. However, fewer studies have focussed on female

remating rates and the timing of offspring production, despite these traits likely also being

under sexual conflict and highly relevant in nature (Eberhard, 1996; Arnqvist & Nilsson,

2000; Arnqvist & Rowe, 2005).

Our study addresses critical shortcomings in previous work and illuminates seemingly contradictory results. SAC theory predicts that sex-specific effects will be likely be highly dependent on genotype-by-genotype interactions (Rowe et al., 2003; Wensing & Fricke, 2018), so mixed results across studies using either different populations or inter-population crosses in different ways is unsurprising (e.g. Smith et al., 2009; Wensing & Fricke, 2018). Our study, by taking an explicitly comparative approach and using within-population crosses, avoids this unpredictability, and instead focusses on patterns of co-evolution between males and females, across populations. Additionally, our study also highlights the perils of a specific focus on genetic variation in, and the direct interactions of, only *SP* and *SPR;* the SP-network consists of several other male and interacting female proteins that may play a more important role in the co-evolution between males and females. We show that effects of molecular exaggeration at SPR is relatively small and that, instead, other loci of the female component of the SP-network is driving female resistance evolution. Not accounting for the effects of variation at these loci will miss much of the relevant biology.

Conclusions

In sum, we extend SAC to the molecular level for the first time. We predict and test outcomes of mating interactions by documenting patterns of natural genetic co-evolution and molecular exaggeration at interacting loci of the classic *Drosophila* SP-network. Our results show that relatively few interacting loci can drive SAC, and predict sexually antagonistic phenotypes, which, coupled with strong selection, may explain the rapid pace of this evolutionary process, at least for some traits and genes. This also offers encouragement to other studies of sexually antagonistic complex behavioural and morphological traits in that, while good knowledge of the relevant interacting loci is needed, not all loci underlying these traits need be accounted for. Our approach offers a

standard that can be extended to ongoing work identifying loci underlying conflict traits in other systems to yield important insights into how SAC can shape patterns of genetic variation (e.g. Khila et al., 2009; Rowe et al., 2018; Crumière & Khila, 2019; Pruvôt et al., 2025).

Endmatter

Data and code availability

Author contributions

R.A.W.W. contributed with the conception of the study, funding acquisition, data collection and analysis, and led the writing of the manuscript. A.T. contributed with experimental design, collection and analysis of data, R.R.S. contributed with the conception of the study, funding acquisition, and writing of the manuscript.

Funding

This work was funded by a Starting Grant from the Swedish Research Council (Vetenskapsrådet) [grant no. 2022-03701], and a European Commission Marie Sklodowska-Curie Actions Independent Postdoctoral Fellowship [grant no. 101061664] to RAWW, as well as a Project Grant from the Swedish Research Council (Vetenskapsrådet) to RRS [2018-04598].

Conflicts of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgements

We are grateful to the DrosEU and DrosRTEC consortia for their heroic efforts of sampling, sequencing, and establishing of isofemale line resources of Drosophila melanogaster, and also for helpful feedback. In particular we are grateful to Thomas Flatt, Pau Carazo, Mathieu Gautier for advice in the planning of experiments; to Élio Sucena for maintaining and shipping us isofemale lines. We are also grateful to Nadeem Naber, Caja Samsen, and Matilde Stella for help in the lab during experiments and to Luc Bussiere and Mike Ritchie for comments that improved the manuscript. The computations and data handling was enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS), partially funded by the Swedish Research Council through grant agreement no. 2022-06725.

References

Adams, D. C., & Felice, R. N. (2014). Assessing Trait Covariation and Morphological Integration on Phylogenies Using Evolutionary Covariance Matrices. *PLoS ONE*, *9*, e94335. https://doi.org/10.1371/journal.pone.0094335

Adams, D., Collyer, M., Kaliontzopoulou, A., & Baken, E. (n.d.). *Geomorph: Software for geometric morphometric analyses* (Version 4.0.7) [R]. https://cran.r-project.org/package=geomorph

Arnqvist, G., & Nilsson, T. (2000). The evolution of polyandry: Multiple mating and female fitness in insects. *Animal Behaviour*, *60*, 145–164. https://doi.org/10.1006/anbe.2000.1446

Arnqvist, G., & Rowe, L. (2002a). Antagonistic coevolution between the sexes in a group of insects. *Nature*, *415*, 787–789. https://doi.org/10.1038/415787a

Arnqvist, G., & Rowe, L. (2002b). Correlated evolution of male and female morphologies in water striders. *Evolution*, *56*, 936–947. https://doi.org/10.1111/j.0014-3820.2002.tb01406.x

Arnqvist, G., & Rowe, L. (2005). *Sexual conflict*. Princeton University Press.

Baken, E. K., Collyer, M. L., Kaliontzopoulou, A., & Adams, D. C. (2021). geomorph v4.0 and gmShiny: Enhanced analytics and a new graphical interface for a comprehensive morphometric experience. *Methods in Ecology and Evolution*, *12*, 2355–2363. https://doi.org/10.1111/2041-210X.13723

Bergsten, J., & Miller, K. B. (2007). Phylogeny of diving beetles reveals a coevolutionary arms race between the sexes. *PLoS ONE*, *2*, e522. https://doi.org/10.1371/journal.pone.0000522

Bergsten, J., Töyrä, A., & Nilsson, A. N. (2001). Intraspecific variation and intersexual correlation in secondary sexual characters of three diving beetles (Coleoptera: Dytiscidae). *Biological Journal of the Linnean Society*, *73*, 221–232. https://doi.org/10.1111/j.1095-8312.2001.tb01359.x

Castillo, D. M. (2017). Factors contributing to the accumulation of reproductive isolation: A mixed model approach. *Ecology and Evolution*, *7*, 5808–5820. https://doi.org/10.1002/ece3.3093

Chelini, M. C., Brock, K., Yeager, J., & Edwards, D. L. (2021). Environmental drivers of sexual dimorphism in a lizard with alternative mating strategies. *Journal of Evolutionary Biology*, *34*, 1241–1255. https://doi.org/10.1111/jeb.13881

Crudgington, H. S., Beckerman, A. P., Brüstle, L., Green, K., & Snook, R. R. (2005). Experimental Removal and Elevation of Sexual Selection: Does Sexual Selection Generate Manipulative Males and Resistant Females? *American Naturalist*, *165*, S72–S87.

Crumière, A. J. J., & Khila, A. (2019). Hox genes mediate the escalation of sexually antagonistic traits in water striders. *Biology Letters*, *15*, 20180720. http://dx.doi.org/10.1098/rsbl.2018.0720

Durmaz Mitchell, E., Kerdaffrec, E., Harney, E., Paulo, T. F., Veselinovic, M. S., Tanaskovic, M., Tyukmaeva, V., de Arcaya, T. A. F., Aksoy, C., Argyridou, E., Bailly, T. P. M., Can, D., Cobanoglu, E., Cook, N., Coşkun, S., Davidovic, S., Demir, E., Dias, T., Rasouli-Dogaheh, S., … Flatt, T. (2025). Continent-wide differentiation of

fitness traits and patterns of climate adaptation among European populations of *Drosophila melanogaster*. *Evolution Letters*, *9*, 473–490. https://doi.org/10.1093/evlett/qraf014

Eberhard, W. G. (1996). *Female Control: Sexual Selection by Cryptic Female Choice*. Princeton University Press.

Edward, D. A., Fricke, C., & Chapman, T. (2010). Adaptations to sexual selection and sexual conflict: Insights from experimental evolution and artificial selection. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*, 2541–2548. https://doi.org/10.1098/rstb.2010.0027

Ferretti, L., Ramos-Onsins, S. E., & Pérez-Enciso, M. (2013). Population genomics from pool sequencing. *Molecular Ecology*, *22*, 5561–5576. https://doi.org/10.1111/mec.12522

Findlay, G. D., Sitnik, J. L., Wang, W., Aquadro, C. F., Clark, N. L., & Wolfner, M. F. (2014). Evolutionary Rate Covariation Identifies New Members of a Protein Network Required for Drosophila melanogaster Female Post-Mating Responses. *PLoS Genetics*, *10*, e1004108. https://doi.org/10.1371/journal.pgen.1004108

Fromonteil, S., Marie-Orleach, L., Winkler, L., & Janicke, T. (2023). Sexual selection in females and the evolution of polyandry. *PLOS Biology*, *21*, e3001916. https://doi.org/10.1371/journal.pbio.3001916

Gautier, M., Coronado-Zamora, M., & Vitalis, R. (2024). *Estimating hierarchical F – statistics from Pool–Seq data*. bioRxiv. https://doi.org/10.1101/2024.11.22.624688

Gautier, M., Foucaud, J., Gharbi, K., Cézard, T., Galan, M., Loiseau, A., Thomson, M., Pudlo, P., Kerdelhué, C., & Estoup, A. (2013). Estimation of population allele frequencies from next-generation sequencing data: Pool-versus individual-based genotyping. *Molecular Ecology*, *22*, 3766–3779. https://doi.org/10.1111/mec.12360

Haerty, W., Jagadeeshan, S., Kulathinal, R. J., Wong, A., Ravi Ram, K., Sirot, L. K., Levesque, L., Artieri, C. G., Wolfner, M. F., Civetta, A., & Singh, R. S. (2007). Evolution in the Fast Lane: Rapidly Evolving Sex-Related Genes in Drosophila. *Genetics*, *177*, 1321–1335. https://doi.org/10.1534/genetics.107.078865

Hivert, V., Leblois, R., Petit, E. J., Gautier, M., & Vitalis, R. (2018). Measuring Genetic Differentiation from Pool-seq Data. *Genetics*, *210*, 315–330.

Holland, B., & Rice, W. R. (1998). Perspective: Chase-Away Sexual Selection: Antagonistic Seduction Versus Resistance. *Evolution*, *52*, 1–7.

Hopkins, B. R., Angus-Henry, A., Kim, B. Y., Carlisle, J. A., Thompson, A., & Kopp, A. (2024). Decoupled evolution of the *Sex Peptide* gene family and *Sex Peptide Receptor* in Drosophilidae. *Proceedings of the National Academy of Sciences*, *121*(3), e2312380120. https://doi.org/10.1073/pnas.2312380120

Hopkins, B. R., & Perry, J. C. (2022). The evolution of sex peptide: Sexual conflict, cooperation, and coevolution. *Biological Reviews*, *97*, 1426–1448. https://doi.org/10.1111/brv.12849

Kapun, M., Nunez, J. C. B., Bogaerts-Marquez, M., Murga-Moreno, J., Paris, M., Outten, J., Coronado-Zamora, M., Tern, C., Rota-Stabelli, O., Guerreiro, M. P. G., Casillas, S., Orengo, D. J., Puerma, E., Kankare, M., Ometto, L., Loeschcke, V., Onder, B. S., Abbott, J. K., Schaeffer, S. W., … Bergland, A. O. (2021). *Drosophila Evolution over Space and Time (DEST): A New Population Genomics Resource*. *38*, 5782–5805.

Kasimatis, K. R., Nelson, T. C., & Phillips, P. C. (2017). Genomic Signatures of Sexual Conflict. *Journal of Heredity*, *108*, 780–790. https://doi.org/10.1093/jhered/esx080

Khila, A., Abouheif, E., & Rowe, L. (2009). Evolution of a Novel Appendage Ground Plan in Water Striders Is Driven by Changes in the Hox Gene Ultrabithorax. *PLoS Genetics*, *5*, e1000583. https://doi.org/10.1371/journal.pgen.1000583

Kubli, E., & Bopp, D. (2012). Sexual Behavior: How Sex Peptide Flips the Postmating Switch of Female Flies. *Current Biology*, *22*, R520–R522. https://doi.org/10.1016/j.cub.2012.04.058

McGeary, M. K., & Findlay, G. D. (2020). Molecular evolution of the sex peptide network in *Drosophila*. *Journal of Evolutionary Biology*, *33*, 629–641. https://doi.org/10.1111/jeb.13597

Nelson, C. W., Moncla, L. H., & Hughes, A. L. (2015). SNPGenie: Estimating evolutionary parameters to detect natural selection using pooled next-generation sequencing data. *Bioinformatics*, *31*, 3709–3711. https://doi.org/10.1093/bioinformatics/btv449

Nunez, J. C. B., Coronado-Zamora, M., Gautier, M., Kapun, M., Steindl, S., Ometto, L., Hoedjes, K. M., Beets, J., Wiberg, R. A. W., Mazzeo, G. R., Bass, D. J., Radionov, D., Kozeretska, I., Zinchenko, M., Protsenko, O., Serga, S., Amor-Jimenez, C., Casillas, S., Sanchez-Gracia, A., … Gonzalez, J. (2025). Footprints of worldwide adaptation in structured populations of *D. melanogaster* through the expanded DEST 2.0 genomic resource. *Molecular Biology and Evolution*, *42*, msaf132. https://doi.org/10.1101/2024.11.10.622744

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics*, *20*, 289–290. https://doi.org/10.1093/bioinformatics/btg412

Parker, G. A. (1970). Sperm competition and its evolutionary consequences in the insects. *Biological Reviews*, *45*, 525–567. https://doi.org/10.1111/j.1469-185X.1970.tb01176.x

Parker, G. A. (1979). Sexual selection and sexual conflict. In *Sexual Selection and Reproductive Competition in Insects* (pp. 123–166). Academic Press. https://doi.org/10.1016/B978-0-12-108750-0.50010-0

Parker, G. A. (2020). Conceptual developments in sperm competition: A very brief synopsis. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*, 20200061. https://doi.org/10.1098/rstb.2020.0061

Perry, J. C., & Rowe, L. (2012). Sexual conflict and antagonistic coevolution across water strider populations. *Evolution*, *66*, 544–557. https://doi.org/10.1111/j.1558-5646.2011.01464.x

Perry, J. C., & Rowe, L. (2015). The evolution of sexually antagonistic phenotypes. *Cold Spring Harbor Perspectives in Biology*, *7*, a017558. https://doi.org/10.1101/cshperspect.a017558

Priest, N. K., Galloway, L. F., & Roach, D. A. (2008). Mating Frequency and Inclusive Fitness in *Drosophila melanogaster*. *The American Naturalist*, *171*, 10–21. https://doi.org/10.1086/523944

Priest, N. K., Roach, D. A., & Galloway, L. F. (2008). Cross-generational fitness benefits of mating and male seminal fluid. *Biology Letters*, *4*, 6–8. https://doi.org/10.1098/rsbl.2007.0473

Pruvôt, C., Armisén, D., Roux, P., Arnqvist, G., Rowe, L., Husby, A., & Khila, A. (2025). A shared developmental genetic basis for sexually antagonistic male and female adaptations in the toothed water strider. *Evolution Letters*, *9*, 13–23. https://doi.org/10.1093/evlett/qrae056

Raas, M. W. D., & Dutheil, J. Y. (2024). The rate of adaptive molecular evolution in wild and domesticated *Saccharomyces cerevisiae* populations. *Molecular Ecology*, *33*(10), e16980. https://doi.org/10.1111/mec.16980

Rohlf, F. J., & Corti, M. (2000). Use of Two-Block Partial Least-Squares to Study Covariation in Shape. *Systematic Biology*, *49*, 740–753. https://doi.org/10.1080/106351500750049806

Rönn, J., Katvala, M., & Arnqvist, G. (2007). Coevolution between harmful male genitalia and female resistance in seed beetles. *Proceedings of the National Academy of Sciences of the United States of America*, *104*, 10921–10925. https://doi.org/10.1073/pnas.0701170104

Rowe, L., Cameron, E., & Day, T. (2003). Detecting sexually antagonistic coevolution with population crosses. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *270*, 2009–2016. https://doi.org/10.1098/rspb.2003.2453

Rowe, L., Chenoweth, S. F., & Agrawal, A. F. (2018). The Genomics of Sexual Conflict. *The American Naturalist*, *192*, 274–286. https://doi.org/10.1086/698198

Rundle, H. D., Ödeen, A., & Mooers, A. Ø. (2007). An experimental test for indirect benefits in Drosophila melanogaster. *BMC Evolutionary Biology, 7*, 36. https://doi.org/10.1186/1471-2148-7-36

Singh, A., Buehner, N. A., Lin, H., Baranowski, K. J., Findlay, G. D., & Wolfner, M. F.
(2018). Long-term interaction between Drosophila sperm and sex peptide is
mediated by other seminal proteins that bind only transiently to sperm. *Insect
Biochemistry and Molecular Biology*, *102*, 43–51.
https://doi.org/10.1016/j.ibmb.2018.09.004

Sirot, L. K., Wong, A., Chapman, T., & Wolfner, M. F. (2015). Sexual Conflict and Seminal
Fluid Proteins: A Dynamic Landscape of Sexual Interactions. *Cold Spring Harbor
Perspectives in Biology*, *7*, a017533. https://doi.org/10.1101/cshperspect.a017533

Smith, D. T., Clarke, N. V. E., Boone, J. M., Fricke, C., & Chapman, T. (2017). Sexual
conflict over remating interval is modulated by the *sex peptide* pathway.
Proceedings of the Royal Society B: Biological Sciences, *284*, 20162394.
https://doi.org/10.1098/rspb.2016.2394

Smith, D. T., Hosken, D. J., Ffrench-Constant, R. H., & Wedell, N. (2009). Variation in sex
peptide expression in *D. melanogaster*. *Genetics Research*, *91*, 237–242.
https://doi.org/10.1017/S0016672309000226

Stone, G. N., Nee, S., & Felsenstein, J. (2011). Controlling for non-independence in
comparative analysis of patterns across populations within species. *Philosophical
Transactions of the Royal Society B: Biological Sciences*, *366*(1569), 1410–1424.
https://doi.org/10.1098/rstb.2010.0311

Swanson, W. J., & Vacquier, V. D. (2002). The rapid evolution of reproductive proteins.
Nature Reviews Genetics, *3*, 137–144. https://doi.org/10.1038/nrg733

Tatarnic, N. J., & Cassis, G. (2010). Sexual coevolution in the traumatically inseminating
plant bug genus *Coridromius*. *Journal of Evolutionary Biology*, *23*, 1321–1326.
https://doi.org/10.1111/j.1420-9101.2010.01991.x

Wensing, K. U., & Fricke, C. (2018). Divergence in sex peptide-mediated female post-
mating responses in *Drosophila melanogaster*. *Proceedings of the Royal Society B:
Biological Sciences*, *285*, 20181563. https://doi.org/10.1098/rspb.2018.1563

Wilkinson, G. S., Breden, F., Mank, J. E., Ritchie, M. G., Higginson, A. D., Radwan, J.,
Jaquiery, J., Salzburger, W., Arriero, E., Barribeau, S. M., Phillips, P. C., Renn, S. C.

P., & Rowe, L. (2015). The locus of sexual selection: Moving sexual selection studies into the post-genomics era. *Journal of Evolutionary Biology*, *28*, 739–755. https://doi.org/10.1111/jeb.12621

Figure 1. A). Estimates of π (top), Tajima's D (middle), and πN/πS (bottom) for loci in the

male- and female-components of the SP-network. Red dashed lines show the maximum

and minimum values obtained for a set of 100 randomly sampled genes from the rest of

the genome. **B)** Male and female axes of co-variation in Fst at SP-network loci. Points are

coloured by their continent of origin. Populations chosen for behavioural experiments, and

the sex with inferred advantage are highlighted with coloured circles. The dashed lines

labelled "PC1" and "PC2" illustrate the first two principle components from a PCA of the

male and female axes of co-variation.

Figure 2. A) Female re-mating rates as a function of a continuous measure of relative

genetic differentiation at female- and male- components of the SP-network. Negative

values indicate relatively more genetic differentiation in the female- versus the male-

component. Each individual point is a replicate population cage, points are coloured by

which sex is inferred to have an advantage in sexual conflicts (red – female-advantage,

blue – male advantage, black - intermediate). The slope and error ribbons represent the

posterior mean and credible intervals for the slope. **B)** Posterior distribution of the slope

estimate. The area shaded in red indicates the proportion of the posterior distribution with

slope estimates < 0, that supports a negative relationship between relative genetic

differentiation and the female re-mating rate, the grey box and bar indicate the posterior

mean and 89% HDPI respectively.

Figure 3. A) The log(number of days) until the peak of offspring production as a function of

a continuous measure of relative genetic differentiation at female- and male- components

of the SP-network. Negative values indicate relatively more genetic differentiation in the

female- *versus* the male-component. Each individual point is a replicate population cage,

points are coloured by which sex is inferred to have an advantage in sexual conflicts (red –

female-advantage, blue – male advantage, black - intermediate). The slope and error

ribbons represent the posterior mean and credible intervals for the slope. **B)** Posterior

distribution of the slope estimate. The area shaded in red indicates the proportion of the

posterior distribution with slope estimates < 0, that supports a negative relationship

between relative genetic differentiation and the peak of offspring production, the grey box

and bar indicate the posterior mean and 89% HDPI respectively.