**Hemocyte Viability Assay as an Alternative Method for Testing Bacterial Pathogenicity in**

**Bivalves**

Jaypee S. Samson[1,2], David C. Rowley[3], and Marta Gomez-Chiarri[1]

[1] Department of Fisheries, Animal and Veterinary Science, University of Rhode Island, Kingston, Rhode Island, USA

[2] College of Fisheries and Freshwater Aquaculture Center, Central Luzon State University, Science City of Muñoz, Nueva Ecija, Philippines

[3] Department of Biomedical and Pharmaceutical Sciences, University of Rhode Island, Kingston, Rhode Island, USA

## ABSTRACT

Bacterial pathogens cause disease outbreaks in hatcheries, resulting to larval mortality and significant economic losses. One of the major challenges in studying bacterial pathogenesis in larvae is the limited availability of healthy larvae throughout the year. This seasonal constraint delays testing and impedes progress in understanding bacterial infections, disease management, and the development of effective treatments for bivalve aquaculture. To address this challenge, a hemocyte viability assay based on the resazurin assay was developed as an alternative for testing bacterial pathogenesis. The assay was optimized using eastern oyster hemocytes, and the lethal concentration $(LC)_{50}$ values of a well-known bacterial pathogen of larvae (*Vibrio coralliilyticus* RE22) ($107 \pm 22$ multiplicity of infection (MOI)), a probiont (*Phaeobacter inhibens* S4) ($2157 \pm 167$ MOI), and several bacterial isolates from shellfish hatcheries were determined. The most virulent isolates had an $LC_{50}$ ranging from 53 to 195 MOI, indicating high pathogenicity. To validate the assay, the isolates were tested on oyster larvae, confirming that bacterial isolates with low $LC_{50}$ (<200 MOI) caused significant larval mortality, while other isolates exhibiting $LC_{50}$ values ranging from 690 to 25389 MOI showed no adverse effect on the oyster larvae. This high-throughput hemocyte viability assay provides a reliable and less time-consuming alternative to larval assays, particularly during the off-season, facilitating year-round research on bacterial pathogenicity in bivalves.

Key words: Bivalve, Larvae, Hemocytes, Resazurin, Oyster, Pathogenicity, Viability

## INTRODUCTION

Bivalves are important in fisheries, aquaculture, and ecosystem structure and function. Because of these critical roles, significant research has focused on their immunological responses to infectious diseases, suggesting that they could potentially be used as a model system to study human health (Fernández Robledo et al., 2019). Like other invertebrates, bivalves have a diverse range of immune cells, genes, and proteins. Although mucosal tissues in contact with the environment play an important role in host defense, hemocytes are the backbone of the bivalve immune system (Allam and Raftos, 2015).

Among the various threats to bivalve health includes the bacterial pathogens, which are a leading cause of disease outbreaks in bivalve larvae, resulting in significant mortality and economic losses. Most bacterial diseases of bivalves are caused by a large range of *Vibrio* spp., *Pseudomonas,* and *Aeromonas* (Zannella et al., 2017). Due to the filter-feeding behavior of bivalves, they concentrate a rich and diverse bacterial commensal microbiota, comprising several species belonging to distinct genera including *Vibrio, Pseudomonas, Acinetobacter, Photobacterium, Moraxella, Aeromonas, Micrococcus* and *Bacillus* (Kueh and Chan, 1985; Zannella et al., 2017); some of them may be harmful in larval rearing systems but not in the wild. These bacteria can rapidly proliferate in aquaculture environments, often exploiting stress or environmental fluctuations to infect and overwhelm bivalve hosts. *Vibrio* spp. are particularly well-characterized, producing virulence factors like hemolysins and ciliostatic toxins which cause deciliation, loss of velar epithelial, and abnormal swimming behavior (Paillard et al., 2004).

A major challenge in studying the pathogenesis of bacterial isolates and their effects on the larvae is the limited availability of healthy bivalve larvae throughout the year. This seasonal

limitation impedes the ability to consistently assess the pathogenicity of bacterial isolates and their

interaction with host organisms under varying environmental conditions. As larvae are often only

available during specific times of the year, researchers are left with a narrow window for

experimental testing, which may limit the robustness and reproducibility of findings. This

restriction can delay research into bacterial infections, disease management, and the development

of effective treatments for bivalve aquaculture. To address the challenges of studying bacterial

pathogenicity and virulence in bivalve larvae, alternative high-throughput methods that are more

consistent, accessible,  repeatable, and faster are essential. The Alamar Blue assay, also known as

the resazurin assay, has emerged as a valuable tool for assessing hemocyte viability and metabolic

activity. This assay relies on the reduction of resazurin to resorufin, a fluorescent compound that

serves as an indicator of cellular metabolic activity (O'Brien et al., 2000). This assay has been

widely used to measure cell viability and cytotoxicity in a range of biological and environmental

systems and several cell types, including bacteria, yeast, fungi, protozoa, and cultured mammalian

and piscine cells (O'Brien et al., 2000; Petiti et al., 2025; Rampersad, 2012).

Bivalves have an open circulatory system in which hemolymph, containing hemocytes,

bathes all the organs before returning to the heart via sinuses and respiratory structures. These

circulating hemocytes are primarily responsible for defending against pathogens through

mechanisms such as phagocytosis and encapsulation (Canesi et al., 2002; Fisher, 1986).

Additionally, antibacterial effectors, opsonins, nonspecific hydrolytic enzymes, and toxic oxygen

intermediates found in bivalve hemolymph coordinate the immune response (Chong, 2022; Gerdol

et al., 2018; Song et al., 2010). The classification of bivalve hemocytes has been a subject of

debate, though they are generally categorized into hyalinocytes and granulocytes (de la Ballina et

al., 2022; López et al., 1997). The latter can be further subdivided into eosinophilic granular

hemocytes and basophilic granular hemocytes (López et al., 1997; Wootton et al., 2003). Other

hemocyte types, such as morula-like cells and blast-like cells, have also been reported in some

species (Donaghy et al., 2009; Wootton et al., 2003). Among these, granular hemocytes are the

most numerous and function as the primary phagocytic cells in bivalves(Allam et al., 2001;

Cajaraville and Pal, 1995). Hemocytes can recognize, locate, ingest, transport, and digest foreign

particles, playing a central role in phagocytosis, encapsulation, inflammation, and wound repair

(Fisher, 1986). Mobility is a critical characteristic of hemocytes, enabling them to migrate toward

pathogens and perform immune functions efficiently. Studies on *Crassostrea virginica* and

*Mercenaria mercenaria* have demonstrated that granulocytes are the predominant phagocytes

responsible for host defense (Cheng et al., 1975; Foley and Cheng, 1975). While granulocytes are

generally associated with pathogen clearance, some evidence suggests that agranulocytes may also

contribute to immune responses (Sokolnikova et al., 2023).

Hemocyte viability is a measure of the proportion of alive cells to evaluate the overall

hemocyte health. Viability assays can be assessed based on cellular metabolism, enzyme activity,

or cell membrane integrity. The widely used approach was double fluorescence staining with

membrane-permeable nuclear dyes, such as SYBR, and the membrane-impermeable dyes, such as

propidium iodide (PI), and detected by use of fluorescence microscopy or flow cytometry (Allam

et al., 2001). In contrast, the resazurin assay offers a high-throughput, plate-based method that is

both rapid and quantifiable.

The goal of this work is to develop an alternative and high-throughput viability assay that

can be used to assess bacterial infections and virulence without relying on the seasonal availability

of bivalve larvae. Hemocytes, as key immune cells in mollusks, can serve as an alternative to

assess the impact of bacterial infections, enabling the testing of bacterial virulence in the absence

of larvae. This assay can provide valuable insights into host-pathogen interactions, mechanisms of

pathogenesis, immune responses, and virulence quantification of bacterial isolates. This could

enhance the efficiency and flexibility of pathogen characterization, ultimately contributing to

better disease management and prevention strategies in bivalve aquaculture.

## MATERIALS AND METHODS

### Bacteria Isolation and Identification

Bacteria were isolated from bivalve (eastern oyster *Crassostrea virginica* and hard clam *Mercenaria mercenaria*) larvae samples from different hatcheries (n = 4) in the Atlantic Coast of the US (hatchery identity not provided per non-disclosure agreement with the hatcheries). Upon samples receipt, larvae were washed and placed in filtered sterile seawater (FSSW) overnight at room temperature (20 to 22 °C) with aeration before bacterial isolation. To isolate bacterial strains, larvae were filtered through a 40-µm mesh and washed with FSSW. A 100 µL aliquot of the concentrated larvae was combined with 100 µL FSSW and macerated using a sterile grinder or pipette tip until the sample became homogeneous. A 100 µL aliquot was then taken for serial dilution. Then, 20 µL was plated on various media, including YP30 (yeast extract (0.5 %), peptone (0.1 %), sea salts (3 %) and agar (1.5%), pH 7.6) (Karim et al., 2013), marine agar (MilliporeSigma, USA), seawater tryptone agar (SWT, tryptone (0.5%), yeast extract (0.3 %), glycerol (0.3%), agar (1.5%), pH 7.6), and nutrient-free agar plates, and incubated for 2-5 days at 27 °C. After incubation, distinct colonies were selected and streaked onto fresh sterile plates until single colonies were isolated.

The DNA from isolated bacteria were extraction following a modified protocol (Li et al., 2023). Briefly, overnight culture (500 µL) of each isolate was harvested and resuspended in DNase-free water with 0.1 g of 0.1 mm glass beads (Sigma-Aldrich, USA). The cells were heated at 80 °C at 500 rpm for 10 minutes, followed by RNase treatment (Qiagen, Germany). Bacterial isolates were identified by 16S rRNA sequencing using universal primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTTACCTTGTTACGACTT-3') (Weisburg et al., 1991). PCR reactions were prepared in a total volume of 25 µL, comprising 5 µL

of extracted DNA, 12.5 µL of PCR Mastermix, 1 µL each of forward (27F) and reverse (1492R)

primers, and 5.5 µL of nuclease-free water. Amplification was performed using the following

thermal cycling conditions: initial denaturation at 95°C for 3 minutes; 30 cycles of denaturation at

95°C for 30 seconds, annealing at 55°C for 30 seconds, and extension at 72°C for 90 seconds;

followed by a final extension at 72°C for 5 minutes. PCR products were purified using a Qiagen

PCR purification kit following the manufacturer's protocol and sequenced at the URI-INBRE

facility. Sequence identification was conducted by comparing obtained sequences to the NCBI 16S

database using BLAST (Camacho et al., 2009).

**Preparation of Bacterial Dilutions**

Bacterial isolates (Table 1) were cultured overnight on Yeast Peptone 30 agar plates at

27°C for 18-24 hours. After incubation, individual colonies were picked and transferred to liquid

YP30 medium, where they were incubated at 27°C with shaking (150 rpm) for 18-24 hours.

Following the incubation, the bacterial cultures were centrifuged at 13,000 rpm for 5 minutes to

pellet the cells. The supernatant was discarded, and the bacterial pellet was washed twice with

filtered seawater (FSSW) to remove the residual medium. The cell pellet was then resuspended in

FSSW, and the optical density ($OD_{600}$) was adjusted to the desired concentration ($10^8$ CFU mL$^{-1}$).

To prepare for the assay, a series of two-fold bacterial dilutions was made in a 96-well

plate. The final bacterial concentration in each well was verified by plating appropriate dilutions

onto YP30 agar plates and determining colony counts to confirm the accuracy of the dilution series

(Karim et al., 2013).

**Preparation of Oyster Hemocytes**

Adult eastern oysters (*Crassostrea virginica*) from Rhode Island farms were collected for

hemocyte isolation, and all procedures were conducted under aseptic conditions. The shells of the

oysters were scrubbed clean and rinsed in freshwater and sprayed with 70% ethanol.  For each

oyster, a small hole was made near the adductor muscle using flame-sterilized pliers. Then, a sterile

1.5" 21 G needle, attached to a 3 mL syringe, was primed with 100 µL of ice-cold, filtered sterile

seawater (FSSW, salinity of 28 psu), ensuring that air bubbles were removed from the syringe.

The primed needle was inserted into the adductor muscle notch, and approximately 1 mL of

hemolymph was extracted from each oyster. The needle was rotated gently during extraction to

help remove tissue adhering to the needle. Hemolymph samples were passed through a 40 µm

mesh filter into a 50 mL Falcon tube kept on ice to remove large tissue debris.

The pooled hemolymph (6-12 oysters) was then centrifuged at 2500 rpm for 3 minutes at

4°C to collect the hemocytes. The supernatant was discarded, and the hemocytes were washed

twice with FSSW and resuspended in the same solution. Hemocyte concentration and viability

were evaluated using a hemocytometer, and concentration was adjusted to between $1.0 \times 10^6$ and

$2.0 \times 10^6$ cells $mL^{-1}$. For heat-killed hemocyte preparation, half of the total hemocyte suspension

volume was placed in a water bath at 90°C for 10 minutes (Fig. 1A). Live and heat-killed (HK)

hemocyte solutions were kept on ice for use in subsequent assays.

**Hemocyte Viability Assay Development**

To assess the impact of bacterial pathogens and probiotics on hemocytes, we developed a

hemocyte viability assay based on the reduction of resazurin dye. This approach quantifies the

hemocyte metabolic activity as a proxy for cell viability following bacterial exposure. The assay

workflow is illustrated in Fig. 1, outlining the key steps: collection and preparation of alive and

heat-killed hemocytes, bacterial serial dilutions, bacterial-hemocyte interaction, antibiotic

addition, a 20-hr incubation period, and finally, measurement of cellular metabolic activity via

fluorescence.

175   The hemocyte viability was assessed resazurin-based dye CellTiter-Blue™ (Promega). The

176 assay was carefully optimized to determine the appropriate hemocyte concentration, antibiotic

177 concentration, and incubation durations for each phase: hemocyte–bacteria interaction, antibiotic

178 treatment, and resazurin metabolism. For the assay, black clear-bottom 96-well plates (Corning,

179 USA) were used. Hemocytes Fifty microliters of either live and heat-killed hemocytes

180 (background control) were added to designated wells of the plate (Fig.1B). A known concentration

181 of pathogenic *Vibrio coralliilyticus* RE22 (Estes et al., 2004) and probiotic *Phaeobacter inhibens*

182 S4 (Karim et al., 2013) were prepared by dilution and spot plating to determine the actual bacterial

183 concentration required for calculating the multiplicity of infection (MOI), defined as the ratio of

184 infectious agents (e.g., bacteria or viruses) to target host cells. Fifty microliters of the bacterial

185 suspension were added to each well containing the hemocytes, each replicated four times. The

186 plates were incubated for 1 hour at room temperature to allow bacterial interaction with hemocytes.

187   After incubation, chloramphenicol was added to the wells to a final concentration of 200

188 $\mu g\ mL^{-1}$ to prevent further bacterial proliferation during the assay. The plates were incubated for

189 an additional 3 hours. Next, 10 µL of resazurin reagent was added to each well, and the plate was

190 shaken gently for 10 seconds. The plate was then incubated for 20 hours at room temperature to

191 allow the reduction of resazurin. After the incubation period, the plate was shaken again for 10

192 seconds, and fluorescence was measured using a fluorometer (BioTek, USA) at an excitation

193 wavelength of 560 nm and an emission wavelength of 590 nm (Fig. 1C).

194 **Data Calculation and Analysis**

195   To calculate hemocyte viability, fluorescence from heat-killed hemocyte wells was

196 subtracted from the fluorescence readings of live hemocyte wells (no treatment control) to account

197 for background fluorescence. The heat-killed hemocyte control consists of hemocytes that have

been inactivated by heat treatment, rendering them metabolically inactive. This control helps

account for background fluorescence as well as any potential effects caused by bacteria utilizing

nutrients released from dead hemocytes, which could influence bacterial metabolism and interfere

with assay readings. The relative viability of the hemocytes was determined by normalizing the

fluorescence of the treated hemocyte samples to that of the control wells. Relative viability was

calculated using the following formula:

$$\text{Viability (\%)} = \frac{F_{\text{Experimental}} - F_{\text{Blank}}}{F_{\text{Control}} - F_{\text{Background}}} \times 100$$

Where:

$F_{\text{Experimental}}$ = fluorescence reading of the well containing hemocytes and the pathogen

$F_{\text{Blank}}$ = fluorescence reading of the well containing heat-killed hemocytes and the pathogen

$F_{\text{Control}}$ = fluorescence reading of the well containing the hemocytes without the pathogen

$F_{\text{Background}}$ = fluorescence reading of the well containing the heat-killed hemocytes without the

pathogen

The MOI was calculated by dividing the bacterial concentration (CFU mL$^{-1}$) by the

hemocyte concentration (cells mL$^{-1}$). Then, a linear regression analysis was performed to plot the

relationship between MOI and relative viability. Using the regression equation, the lethal

concentration for 50% viability (LC$_{50}$) was determined.

**Larval Assay**

To compare the result of the hemocyte viability assay with the larval assay, the effect of

the bacterial isolates on larval viability was assessed using a previously described protocol

(Gómez-León et al., 2008) with modifications.  Briefly, 6-8-day-old healthy eastern oyster and

hard clam larvae (veligers) were added to 24-well plates at 75-100 larvae mL$^{-1}$.  Then, bacterial

isolates were added at a final concentration of $10^5$ CFU mL$^{-1}$ and incubated for 16-20 h at room

temperature. At this point, larval mortality was evaluated by visual examination under a

microscope (Leica, 400x magnification). Larvae mortality was calculated as follows:

$$\text{Mortality (\%)} = \frac{\text{Number of dead larvae}}{\text{Number of dead larvae} + \text{Number of alive larvae}} \times 100$$

**RESULTS**

**Assay Optimization**

To eliminate bacterial growth during the incubation with the resazurin dye, antibiotics

(chloramphenicol) were added to the wells after hemocytes were incubated with the live bacteria

to allow for bacterial effect on live hemocytes. Different concentrations of chloramphenicol were

added in wells of *V. coralliilyticus* RE22 ($10^8$ CFU mL$^{-1}$) and hemocytes ($10^6$ cells mL$^{-1}$),

respectively, to determine the minimum antibiotic concentration that affects the growth of the

bacteria but not the viability of the hemocytes. The minimum concentration of chloramphenicol

that significantly decreased or inhibited the growth of *V. coralliilyticus* RE22 was 200 µg mL$^{-1}$.

On the other hand, hemocyte viability remained even at higher antibiotic concentrations (Table 2).

This approach helps prevent bacterial proliferation, which could otherwise artificially increase the

reduction rate of the dye. It was determined that a hemocyte density of 1 to 2 × $10^6$ cells/mL is

optimal for reliable assay performance. Furthermore, fluorescence readings stabilized after 20

hours of incubation with the resazurin dye, indicating sufficient time for metabolic conversion and

accurate viability measurement. Overall, the assay lasts for 24 h, including the 1 h bacteria-

hemocyte interaction and 3 h antibiotic treatment (Fig. 1C).

**Hemocyte Viability Assay**

To test the capability of the assay in quantifying bacterial virulence, well-known pathogenic (*V. coralliilyticus* RE22) and probiotic (*Phaeobacter inhibens* S4) bacteria were used. A known pathogenic isolate, *V. coralliilyticus* RE22, had a computed $LC_{50}$ value of $107 \pm 22$, while the probiotic *P. inhibens* S4 had $2157 \pm 167$ (Fig. 2). The assay was repeated three times to get the standard deviation. The coefficient of variation was calculated for inter assay accuracy and has 7.7% and 20.5% for S4 and RE22.

The assay was further applied to evaluate the virulence of several bacterial isolates newly isolated from oyster and clam larvae. (Table 1). The $LC_{50}$ values of the bacterial isolates ranged from 53 to 25,389 MOI (Table 3). Among these, the most virulent isolate to oyster larvae was *V. toranzoniae* CH7, with an $LC_{50}$ of 53 MOI, followed by *V. rotiferianus* CH3 (81 MOI), *V. fortis* CH6 (86 MOI), *V. neptunius* DEN11 (129 MOI), *V. chagasii* CH4 (190 MOI), and *C. lytica* CH30 (195 MOI). In contrast, the highest $LC_{50}$ values among the isolates corresponded to *V. chagasii* CLAM14 (25,389 MOI) and *T. ascidiaceicola* CLAM15 (20,315 MOI), suggesting these are non-pathogenic isolates on oyster.

**Larval Assay**

To confirm the result of the hemocyte viability assay, the tested isolates were used to challenge eastern oyster and hard clam larvae for 16 – 20 h at a standardized concentration of $10^5$ CFU mL$^{-1}$ (Fig. 3) . Oyster larvae challenged with the known pathogen *V. coralliilyticus* RE22 had almost 100% mortality. Similarly, potential pathogenic isolates *V. fortis* CH6 and *V. neptunius* DEN11 also had 100% mortality. In addition, *V. chagasii* CH4, *C. lytica* CH30, *V. rotiferianus* CH3, *V. toranzoniae* CH7, and *V. tubiashii* DEN41 had larval survival rates below 50%, confirming their pathogenicity towards the larvae. Conversely, larvae treated with the potential

probiotic isolates exhibited a relative survival rate of 96-100%, suggesting these isolates are

generally non-pathogenic and safe for larvae. In general, bacterial isolates that showed $LC_{50}$ values

below 200 MOI (*i.e., V. toranzoniae* CH7, *V. rotiferianus* CH3, *V. fortis* CH6, *V. neptunius*

DEN11, *V. chagasii* CH4, and *C. lytica* CH30) demonstrated significant pathogenicity towards

oyster larvae.

In clam larvae, highest mortality was observed in *V. coralliilyticus* RE22, *V. tubiashii*

(DEN41 and DEN12), and *V. neptunius* DEN11. Two isolates *V. chagasii* CLAM14, *T.*

*ascidiaceicola* CLAM15 that did not cause significant mortality in oyster larvae, exhibited notably

higher mortality rates in the clam larvae from which they were originally isolated.

**DISCUSSION**

The hemocyte viability assay offers a rapid, high-throughput alternative to traditional larval

assays for assessing bacterial pathogenicity in larvae. Unlike larval-based methods, which are

limited by the seasonal availability of healthy larvae and can be time-consuming and variable, this

assay provides a consistent and quantifiable measure of bacterial effects on hemocytes. To our

knowledge, this is the first report of using resazurin-based viability measurement in bivalve

hemocytes to test bacterial pathogenicity. By using resazurin reduction as a measure for hemocyte

viability, the assay enables sensitive detection of pathogenic impacts in a controlled, reproducible

format. One major challenge encountered during the assay development was the potential

interference of bacterial growth with the resazurin, as metabolically active bacteria can also reduce

the dye, confounding results. This issue was addressed by the addition of antibiotics after initial

bacterial-hemocyte interaction, effectively halting bacterial proliferation during the assay period

and ensuring fluorescence readings primarily reflect hemocyte viability. We found that hemocytes

remain capable of reducing resazurin even in the presence of antibiotics, allowing effective

inhibition of bacterial growth during incubation without compromising the assay. Similar to

previous reports, hemocyte viability was not affected by the addition of antibiotics (Lacaze et al.,

2015; Matozzo et al., 2016).

Potential limitations include the fact that this in vitro assay may not fully capture the

complexity of whole-organism responses in larvae, such as tissue-specific effects or systemic

immune responses. Additionally, some bacterial species may exhibit variable sensitivity to

antibiotics or resazurin, potentially affecting assay accuracy, thus other antibiotics or even

antibiotic cocktail can be optimized for future use. Also, the use of hemocytes from other bivalves

can be explored to test the host specificity of bacterial isolates being tested. Nonetheless, the assay

represents a valuable tool for rapid screening of bacterial virulence and host-pathogen interactions,

complementing more traditional larval infection studies.

***Hemocyte viability assay corresponds with the larval assay***

In this study, we assess the result of the hemocyte assay using the same test isolates we

tested on oyster larvae. To evaluate the pathogenicity, a 20% mortality threshold was arbitrarily

set to identify the pathogenic isolates. The correlation between mortality and $LC_{50}$ of the

pathogenic isolates was strong, with a Pearson correlation coefficient of 0.84 ($P = 0.0022$),

indicating a significant relationship (Fig. 4). Similar to the results of the hemocyte assay, *V.*

*toranzoniae* CH7, *V. rotiferianus* CH3, *V. fortis* CH6, *V. neptunius* DEN11, *V. chagasii* CH4, and

*C. lytica* CH30, which showed less than 200 MOI $LC_{50}$ values, caused more than 20% mortality

in the oyster larvae. In addition, *V. tubiashii* (DEN12 and DEN41) and *V. alginolyticus* DEN30

also had significant mortality on the larvae, although they showed higher $LC_{50}$. The tested potential

pathogens *V. tasmaniensis* CLAM18*, V. cyclitrophicus* CLAM19*, V. chagasii* CLAM14, *T.*

*ascidiaceicola* CLAM15, and *V. alginolyticus* DEN30, which showed high $LC_{50}$ values, did not

affect the larval survival significantly. It is important to note that four of these isolates, except the

latter, were isolated from clams; and when tested against clam larvae (*M. mercenaria*), they

resulted in mortalities of 47%, 17%, 31%, and 51%, respectively. This host-specific pathogenicity

is a well-known occurrence of bacterial pathogens. Because different hosts occupy diverse

environmental niches, a pathogen's fitness can vary depending on the host species. Pathogens that

are well-adapted to certain hosts can evolve into unique strains, a phenomenon known as

polymorphism. Alternatively, certain pathogens may evolve into generalists capable of surviving

and replicating in multiple host species via a variety of methods, a process known as polyphenism

(Bäumler and Fang, 2013).

### 323      *Pathogenic bacteria require less concentration to cause hemocyte mortality*

In this study, several pathogenic bacterial isolates (*i.e., V. toranzoniae* CH7, *V. rotiferianus*

CH3, *V. fortis* CH6, *V. neptunius* DEN11, *V. chagasii* CH4, and *C. lytica* CH30) showed $LC_{50}$

values below 200 MOI. This indicates that these isolates might produce toxins that degrade the

hemocytes. These results are consistent with previous reports indicating pathogenicity for these

species. *V. toranzoniae* and several other *Vibrio* spp. (*V. lentus*, *V. crassostreae*, and *V.*

*cyclitrophicus*) caused significantly higher hemocyte cytotoxic rates in the scallop, *Argopecten*

*purpuratus* (González et al., 2025). Similarly, *V. rotiferianus* has also been reported to show

pathogenicity in insect infection models (Harrison et al., 2022). *V. fortis* has been implicated in

the heat stress-induced mortality of *C. gigas* (Green et al., 2019). *V. neptunius* is a natural part of

the mollusk microbiota and an opportunistic pathogen that causes disease outbreaks in oysters and

clams. It is known to produce siderophores, piscibactin, and amphibactin, which contribute

significantly to its virulence (Galvis et al., 2021). In addition, *V. chagasii* was pointed out as the

causative agent of the massive larval mortalities of the farmed scallop (*A. purpuratus*) in Chile

(Urtubia et al., 2023). *C. lytica* has been isolated in the eroded mantle epithelium of juvenile paua

(*Haliotis iris*) in New Zealand during an outbreak of bacterial disease of the epithelium of the

mantle, epipodium, and foot (Diggles, 2005).

Other *Vibrio* strains tested, including *V. tubiashii* DEN12, DEN41, *V. tasmaniensis*

CLAM18*, V. cyclitrophicus* CLAM19*, V. chagasii* CLAM14, and *V. alginolyticus* DEN30,

showed high $LC_{50,}$ indicating low to non-pathogenicity for hemocytes. Vibrios are gram-negative,

curved rod-shaped, motile, heterotrophic, rapid-growing, and highly adaptable bacteria comprising

more than 140 species that are ubiquitous in aquatic environments. Most are non-pathogenic. They

are often associated with phytoplankton, zooplankton, crustaceans, and bivalves, showing free-

living, symbiotic, and, in some cases, pathogenic life strategies. Pathogenicity is determined at the

strain level, with most species exhibiting a diversity of nonpathogenic and pathogenic strains

(Gómez-Chiarri et al., 2025). Furthermore, pathogenicity depends on the host species, their life

stage, the number of bacteria, and environmental factors (Zannella et al., 2017).

**CONCLUSIONS**

This study aimed to develop an alternative assay for assessing bacterial pathogenicity in

bivalves, utilizing adult oyster hemocytes as a substitute for oyster larvae to overcome the issue of

larval shortages during the off-season. The results demonstrated that the developed hemocyte

viability assay is an effective tool for testing the pathogenicity of bacterial isolates in oysters.

Importantly, this assay can be further adapted to assess the pathogenicity of bacterial isolates in

hemocytes from other bivalve species, such as clams, scallops, and mussels. This flexibility makes

it a valuable tool for investigating host-specific pathogens across a broader range of bivalve

species.

**Conflict of Interests**

The authors declare that they have no competing interests

**Acknowledgements**

We are grateful to all the research and commercial hatcheries involved in the Bivalve Hatchery

Health Consortium, current and previous members of the Gomez-Chiarri lab for their assistance

during this study, and all the members of the Probiotics Working Group at the University of Rhode

Island. We thank Janet Atoyan at the URI-INBRE for the 16S sequencing support.

**Fundings**

This work was supported by USDA AFRI Aquaculture Research Award #2019-70007-

30146 awarded to MGC and DCR.

**REFERENCES**

Allam, B., Ashton-Alcox, K.A., Ford, S.E., 2001. Haemocyte parameters associated with resistance to brown ring disease in *Ruditapes* spp. clams. Dev. Comp. Immunol. 25, 365–375. https://doi.org/10.1016/S0145-305X(00)00072-0

Allam, B., Raftos, D., 2015. Immune responses to infectious diseases in bivalves. J. Invertebr. Pathol., Pathogens and Disease Processes in Marine Molluscs 131, 121–136. https://doi.org/10.1016/j.jip.2015.05.005

Bäumler, A., Fang, F.C., 2013. Host Specificity of Bacterial Pathogens. Cold Spring Harb. Perspect. Med. 3, a010041. https://doi.org/10.1101/cshperspect.a010041

Cajaraville, M.P., Pal, S.G., 1995. Morphofunctional Study of the Haemocytes of the Bivalve Mollusc *Mytilus galloprovincialis* with Emphasis on the Endolysosomal Compartment. Cell Struct. Funct. 20, 355–367. https://doi.org/10.1247/csf.20.355

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: architecture and applications. BMC Bioinformatics 10, 421. https://doi.org/10.1186/1471-2105-10-421

Canesi, L., Gallo, G., Gavioli, M., Pruzzo, C., 2002. Bacteria–hemocyte interactions and phagocytosis in marine bivalves. Microsc. Res. Tech. 57, 469–476. https://doi.org/10.1002/jemt.10100

Cheng, T.C., Rodrick, G.E., Foley, D.A., Koehler, S.A., 1975. Release of lysozyme from hemolymph cells of *Mercenaria mercenaria* during phagocytosis. J. Invertebr. Pathol. 25, 261–265. https://doi.org/10.1016/0022-2011(75)90076-2

Chong, R.S.-M., 2022. Chapter 57 - Molluscan immunology, in: Kibenge, F.S.B., Baldisserotto, B., Chong, R.S.-M. (Eds.), Aquaculture Pathophysiology. Academic Press, pp. 383–392. https://doi.org/10.1016/B978-0-323-95434-1.00057-7

de la Ballina, N.R., Maresca, F., Cao, A., Villalba, A., 2022. Bivalve Haemocyte Subpopulations: A Review. Front. Immunol. 13, 826255. https://doi.org/10.3389/fimmu.2022.826255

Diggles, B., 2005. Diseases of Cultured Paua (*Haliotis iris*) in New Zealand.

Donaghy, L., Kim, B.-K., Hong, H.-K., Park, H.-S., Choi, K.-S., 2009. Flow cytometry studies on the populations and immune parameters of the hemocytes of the Suminoe oyster, *Crassostrea ariakensis*. Fish Shellfish Immunol. 27, 296–301. https://doi.org/10.1016/j.fsi.2009.05.010

Estes, R.M., Friedman, C.S., Elston, R.A., Herwig, R.P., 2004. Pathogenicity testing of shellfish hatchery bacterial isolates on Pacific oyster *Crassostrea gigas* larvae. Dis. Aquat. Organ. 58, 223–230. https://doi.org/10.3354/dao058223

Fernández Robledo, J.A., Yadavalli, R., Allam, B., Pales Espinosa, E., Gerdol, M., Greco, S., Stevick, R.J., Gómez-Chiarri, M., Zhang, Y., Heil, C.A., Tracy, A.N., Bishop-Bailey, D., Metzger, M.J., 2019. From the raw bar to the bench: Bivalves as models for human health. Dev. Comp. Immunol. 92, 260–282. https://doi.org/10.1016/j.dci.2018.11.020

Fisher, W.S., 1986. Structure and Functions of Oyster Hemocytes. Springer, Berlin, Heidelberg.

Foley, D.A., Cheng, T.C., 1975. A quantitative study of phagocytosis by hemolymph cells of the pelecypods *Crassostrea virginica* and *Mercenaria mercenaria*. J. Invertebr. Pathol. 25, 189–197. https://doi.org/10.1016/0022-2011(75)90068-3

Galvis, F., Ageitos, L., Rodríguez, J., Jiménez, C., Barja, J.L., Lemos, M.L., Balado, M., 2021. *Vibrio neptunius* Produces Piscibactin and Amphibactin and Both Siderophores Contribute

Significantly to Virulence for Clams. Front. Cell. Infect. Microbiol. 11.

https://doi.org/10.3389/fcimb.2021.750567

Gerdol, M., Gomez-Chiarri, M., Castillo, M.G., Figueras, A., Fiorito, G., Moreira, R., Novoa, B.,

Pallavicini, A., Ponte, G., Roumbedakis, K., Venier, P., Vasta, G.R., 2018. Immunity in

Molluscs: Recognition and Effector Mechanisms, with a Focus on Bivalvia, in: Cooper,

E.L. (Ed.), Advances in Comparative Immunology. Springer International Publishing,

Cham, pp. 225–341. https://doi.org/10.1007/978-3-319-76768-0_11

Gómez-Chiarri, M., Andrews, J.S., Coppersmith, J., Guidry, M.E., Houtz, A., Mills, B., Murphy,

M., Nelson, D.R., Rowley, D.C., 2025. 11 - Vibriosis of bivalves, in: Smolowitz, R. (Ed.),

Diseases of Bivalves. Academic Press, pp. 143–162. https://doi.org/10.1016/B978-0-12-

820339-2.00005-X

Gómez-León, J., Villamil, L., Salger, S.A., Sallum, R.H., Remacha-Triviño, A., Leavitt, D.F.,

Gómez-Chiarri, M., 2008. Survival of eastern oysters *Crassostrea virginica* from three

lines following experimental challenge with bacterial pathogens. Dis. Aquat. Organ. 79,

95–105. https://doi.org/10.3354/dao01902

González, R., Peña-Jaña, M., Rojas, R., Oyanedel, D., Henríquez-Castillo, C., Astudillo, O.,

Schmitt, P., Brokordt, K., 2025. Pathogenic impact of *Vibrio* spp. in scallop hemolymph:

Implications for health in aquaculture. Aquaculture 742507.

https://doi.org/10.1016/j.aquaculture.2025.742507

Green, T.J., Siboni, N., King, W.L., Labbate, M., Seymour, J.R., Raftos, D., 2019. Simulated

Marine Heat Wave Alters Abundance and Structure of Vibrio Populations Associated with

the Pacific Oyster Resulting in a Mass Mortality Event. Microb. Ecol. 77, 736–747.

https://doi.org/10.1007/s00248-018-1242-9

Harrison, J., Nelson, K., Morcrette, H., Morcrette, C., Preston, J., Helmer, L., Titball, R.W., Butler,

C.S., Wagley, S., 2022. The increased prevalence of *Vibrio* species and the first reporting

of *Vibrio jasicida* and *Vibrio rotiferianus* at UK shellfish sites. Water Res. 211, 117942.

https://doi.org/10.1016/j.watres.2021.117942

Karim, M., Zhao, W., Rowley, D., Nelson, D., Gomez-Chiarri, M., 2013. Probiotic Strains for

Shellfish Aquaculture: Protection of Eastern Oyster, *Crassostrea virginica*, Larvae and

Juveniles Against Bacterial Challenge. J. Shellfish Res. 32, 401–408.

https://doi.org/10.2983/035.032.0220

Kueh, C.S., Chan, K.Y., 1985. Bacteria in bivalve shellfish with special reference to the oyster. J.

Appl. Bacteriol. 59, 41–47. https://doi.org/10.1111/j.1365-2672.1985.tb01773.x

Lacaze, E., Pédelucq, J., Fortier, M., Brousseau, P., Auffret, M., Budzinski, H., Fournier, M., 2015.

Genotoxic and immunotoxic potential effects of selected psychotropic drugs and

antibiotics on blue mussel (*Mytilus edulis*) hemocytes. Environ. Pollut. 202, 177–186.

https://doi.org/10.1016/j.envpol.2015.03.025

Li, S., Liu, X., Li, Z., Liu, H., Hu, D., 2023. Combination of direct boiling and glass beads

increases the purity and accuracy of bacterial DNA extraction. Biotechnol. J. 18, 2300135.

https://doi.org/10.1002/biot.202300135

López, C., Carballal, M.J., Azevedo, C., Villalba, A., 1997. Morphological Characterization of the

Hemocytes of the Clam, *Ruditapes decussatus* (Mollusca: Bivalvia). J. Invertebr. Pathol.

69, 51–57. https://doi.org/10.1006/jipa.1996.4639

Matozzo, V., Bertin, V., Battistara, M., Guidolin, A., Masiero, L., Marisa, I., Orsetti, A., 2016.

Does the antibiotic amoxicillin affect haemocyte parameters in non-target aquatic

invertebrates? The clam *Ruditapes philippinarum* and the mussel *Mytilus galloprovincialis*

as       model       organisms.      Mar.      Environ.      Res.      119,      51–58.

https://doi.org/10.1016/j.marenvres.2016.05.017

O'Brien, J., Wilson, I., Orton, T., Pognan, F., 2000. Investigation of the Alamar Blue (resazurin)

fluorescent dye for the assessment of mammalian cell cytotoxicity. Eur. J. Biochem. 267,

5421–5426. https://doi.org/10.1046/j.1432-1327.2000.01606.x

Paillard, C., Roux, F.L., Borrego, J.J., 2004. Bacterial disease in marine bivalves, a review of

recent studies: Trends and evolution. Aquat. Living Resour. 17, 477–498.

https://doi.org/10.1051/alr:2004054

Petiti, J., Revel, L., Divieto, C., 2025. Optimizing resazurin-based viability assays for P-

MSC/TER308 cell line to enhance results reliability. BMC Res. Notes 18, 228.

https://doi.org/10.1186/s13104-025-07298-w

Rampersad, S.N., 2012. Multiple Applications of Alamar Blue as an Indicator of Metabolic

Function and Cellular Health in Cell Viability Bioassays. Sensors 12, 12347–12360.

https://doi.org/10.3390/s120912347

Sokolnikova, Y., Mokrina, M., Magarlamov, T., Grinchenko, A., Kumeiko, V., 2023.

Specification of hemocyte subpopulations based on immune-related activities and the

production of the agglutinin MkC1qDC in the bivalve *Modiolus kurilensis*. Heliyon 9,

e15577. https://doi.org/10.1016/j.heliyon.2023.e15577

Song, L., Wang, L., Qiu, L., Zhang, H., 2010. Bivalve Immunity, in: Söderhäll, K. (Ed.),

Invertebrate Immunity, Advances in Experimental Medicine and Biology. Springer US,

Boston, MA, pp. 44–65. https://doi.org/10.1007/978-1-4419-8059-5_3

Urtubia, R., Miranda, C.D., Rodríguez, S., Dubert, J., Barja, J.L., Rojas, R., 2023. First Report,

Characterization and Pathogenicity of Vibrio chagasii Isolated from Diseased Reared

Larvae of Chilean Scallop, *Argopecten purpuratus* (Lamarck, 1819). Pathogens 12, 183.

https://doi.org/10.3390/pathogens12020183

Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA amplification

for phylogenetic study. J. Bacteriol. 173, 697–703. https://doi.org/10.1128/jb.173.2.697-

703.1991

Wootton, E.C., Dyrynda, E.A., Ratcliffe, N.A., 2003. Bivalve immunity: comparisons between the

marine mussel (*Mytilus edulis*), the edible cockle (*Cerastoderma edule*) and the razor-shell

(*Ensis siliqua*). Fish Shellfish Immunol. 15, 195–210. https://doi.org/10.1016/S1050-

4648(02)00161-4

Zannella, C., Mosca, F., Mariani, F., Franci, G., Folliero, V., Galdiero, Marilena, Tiscar, P.G.,

Galdiero, Massimiliano, 2017. Microbial Diseases of Bivalve Mollusks: Infections,

Immunology and Antimicrobial Defense. Mar. Drugs 15, 182.

https://doi.org/10.3390/md15060182

Table 1. List of bacterial isolates used in the study.

| ISOLATE TESTED | ISOLATION SOURCE | REFERENCE |
|---|---|---|
| *Phaeobacter inhibens* S4 | *C. virginica* | Karim et al., 2013 |
| *Vibrio coralliilyticus* RE22 | *C. gigas* | Estes et al., 2004 |
| *Vibrio toranzoniae* CH7 | *C. virginica* | This study |
| *Vibrio rotiferianus* CH3 | *C. virginica* | This study |
| *Vibrio fortis* CH6 | *C. virginica* | This study |
| *Vibrio neptunius* DEN11 | *C. virginica* | This study |
| *Vibrio chagasii* CH4 | *C. virginica* | This study |
| *Cellulophaga lytica* CH30 | *C. virginica* | This study |
| *Vibrio tubiashii* DEN41 | *C. virginica* | This study |
| *Vibrio tasmaniensis* CLAM18 | *M. mercenaria* | This study |
| *Vibrio cyclitrophicus* CLAM19 | *M. mercenaria* | This study |
| *Vibrio tubiashii* DEN12 | *C. virginica* | This study |
| *Vibrio alginolyticus* DEN30 | *C. virginica* | This study |
| *Tenacibaculum ascidiaceicola* CLAM15 | *M. mercenaria* | This study |
| *Vibrio chagasii* CLAM14 | *M. mercenaria* | This study |
| *Pseudooceanicola nitratireducens* NEH7 | *C. virginica* | This study |
| *Marinomonas gallaica* CLAM9 | *M. mercenaria* | This study |
| *Algoriphagus yeomjeoni* DEN5 | *C. virginica* | This study |
| *Glutamicibacter soli* CLAM16 | *M. mercenaria* | This study |

502
503

504     Table 2. Antibiotic concentration to prevent or reduce bacterial growth.

| Chloramphenicol (ug mL$^{-1}$) | *V. coralliilyticus* RE22 | Hemocyte viability |
|:---:|:---:|:---:|
| 5 | + | + |
| 10 | + | + |
| 20 | + | + |
| 40 | + | + |
| 60 | + | + |
| 80 | + | + |
| 100 | + | + |
| 150 | +/- | + |
| 200 | - | + |
| 300 | - | + |
| 400 | - | + |
| 500 | - | + |

505    + = resazurin dye turned into pinkish
506    - = resazurin dye remained bluish
507

509 Table 3. Computed $LC_{50}$ value of bacteria isolated from bivalves using the developed hemocyte
510         viability assay.

| ISOLATE TESTED | $LC_{50}$ (MOI) |
|---|---|
| *Phaeobacter inhibens* S4 | $2157 \pm 167$ |
| *Vibrio coralliilyticus* RE22 | $107 \pm 22$ |
| *Vibrio toranzoniae* CH7 | 53 |
| *Vibrio rotiferianus* CH3 | 81 |
| *Vibrio fortis* CH6 | 84 |
| *Vibrio neptunius* DEN11 | 129 |
| *Vibrio chagasii* CH4 | 190 |
| *Cellulophaga lytica* CH30 | 195 |
| *Vibrio tubiashii* DEN41 | 848 |
| *Vibrio tasmaniensis* CLAM18 | 877 |
| *Vibrio cyclitrophicus* CLAM19 | 927 |
| *Vibrio tubiashii* DEN12 | 4160 |
| *Vibrio alginolyticus* DEN30 | 12204 |
| *Tenacibaculum ascidiaceicola* CLAM15 | 20315 |
| *Vibrio chagasii* CLAM14 | 25389 |
| *Pseudooceanicola nitratireducens* NEH7 | 690 |
| *Marinomonas gallaica* CLAM9 | 704 |
| *Algoriphagus yeomjeoni* DEN5 | 935 |
| *Glutamicibacter soli* CLAM16 | 1913 |

Figure 1. Overview of the hemocyte viability assay including the (A) isolation, preparation, and

quantification; (B) plate layout; and (C) the procedure of the hemocyte viability assay using the

resazurin dye. Figure created with BioRender.com.

**Figure 2.** Representative example of linear regression analysis of *V. coralliilyticus* RE22 and *P. inhibens* S4. Relationship between multiplicity of infection (MOI) and hemocyte viability measured by resazurin reduction. Linear regression analysis demonstrates the negative correlation between increasing MOI and hemocyte metabolic activity, indicating reduced viability at higher bacterial loads. Data points represent mean values from replicate wells, with fitted regression lines, corresponding equations, and $R^2$ values.

527
528 **Figure 3.** Relative percent survival of eastern oyster (upper)  and hard clam (lower) larvae

529 challenge and treated with pathogenic and probiotic isolates. Data are presented as mean ± standard

530 error of the mean (SEM). Red dash line indicates 80% survival threshold. ** Known pathogen and

531 probiotics. *Potential probiotics.

**Figure 4**. Correlation of the survival data from pathogenic isolates causing >20% mortalities and their corresponding $LC_{50}$ values based on multiplicity of infection (MOI). The survival percentage is plotted against the $LC_{50}$ values on a logarithmic scale. Survival data points represent mean values. Pearson correlation and linear regression with 95% confidence intervals are shown.

538