1   **Models for the architecture of the human inner kinetochore CCAN complex**
2   **on centromeric α-satellite CENP-A nucleosome arrays**
3

4   Cong Yu[1], Kyle W. Muir[1,3], Jing Yang[1], Ziguo Zhang[1], Stephen H. McLaughlin[1] and David
5   Barford[1,2]
6

7   [1]MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge, CB2 0QH, UK
8

9   [2]Corresponding author: dbarford@mrc-lmb.cam.ac.uk
10

11   [3]Present address: Institute of Cell Biology, University of Edinburgh, Edinburgh, UK.
12

13   **Abstract**
14

Human kinetochores assemble onto centromeric DNA defined by tandem copies of
thousands of the 171 bp α-satellite repeat sequence. The centromere-specific CENP-A
nucleosome (CENP-A[Nuc]) recruits the inner kinetochore constitutive centromere associated
network (CCAN) complex. A previous cryo-EM structure of CCAN bound to a CENP-A[Nuc]
reconstituted with a 171 bp α-satellite repeat sequence showed how extranucleosomal
linker DNA threads through an internal tunnel in the CCAN complex. To understand the
higher-order architecture of the inner kinetochore assembled onto α-satellite repeat arrays
we have determined cryo-EM structures of CCAN with longer DNA sequences. These include
free DNA and single and dimeric α-satellite repeats with CENP-A nucleosomes. We show
from the structures of CCAN bound to both free DNA and monomeric CENP-A[Nuc] that CCAN
engages 65-70 bp of DNA comprising 30-35 bp of an upstream α-satellite repeat. This
upstream DNA interacts with the histone fold domain subunits of the CENP-TWSX module in
a manner resembling how DNA is wrapped in nucleosomes. A complex of CCAN assembled
onto a dimeric α-satellite repeat with two CENP-A nucleosomes showed that CCAN can only
be accommodated on the linker DNA by unwrapping DNA from the CENP-TWSX module
together with 25 bp of DNA from the upstream nucleosome. We discuss the implications of
these results for models of CCAN assembly on arrays of α-satellite chromatin containing
CENP-A[Nuc].
33

34   **Introduction**
35

The faithful inheritance of genetic information is crucial for life. Central to this process in
eukaryotes are the roles fulfilled by kinetochores – large protein complexes that assemble

at centromeric chromatin. Kinetochores function to attach sister chromatids to the mitotic

spindle, and couple the work of microtubule depolymerization to chromatid segregation [1-3].

Kinetochores also act as signalling hubs for the error correction and spindle assembly

checkpoints that together coordinate the correct attachment of sister chromatids to the

mitotic spindle with the initiation of sister chromatid segregation [4,5].

Human regional centromeres are megabase-sized structures, comprising thousands of

contiguous 171 bp α-satellite repeat sequences. Identical α-satellite repeats are organized

into higher order repeat (HOR) arrays of typically four, seven and eleven repeats [6]. Only a

small fraction of α-satellite repeats is associated with the approximately four-hundred

CENP-A nucleosomes per centromere [7]. CENP-A nucleosomes are restricted to the hypo-

methylated centromere dip region (CDR), the most recently evolved segment of

centromeres [6,8,9], and are interspersed with canonical H3 nucleosomes [10]. Assuming at most

147 bp of DNA wraps a histone octamer [11], at least 24 bp of α-satellite repeat sequence

forms extranucleosomal linker DNA. Both *in vivo* ChIP-seq mapping and in *vitro*

reconstitution of CENP-A nucleosomes on α-satellite sequences indicate that the CENP-A

nucleosome is installed precisely towards the 3' end of the repeat, with the centre of the α-

satellite repeat palindrome coinciding with the nucleosome dyad axis [12,13]. Human

centromeres, except for the Y chromosome centromere, contain the 17 bp B-box element.

The majority of B-boxes are arranged on alternating α-satellite repeats [14] positioned such

that the B-box on linker DNA partially overlaps a nucleosome entry site [15]. CENP-B

engagement by the B-box facilitates unwrapping of the DNA gyre from the CENP-A histone

octamer [16], whereas the engagement of the dimeric CENP-B to two separate B-boxes

induces looping of centromeric DNA [17].

The centromere-kinetochore assembly of vertebrates comprises large macromolecular

complexes at the constriction point of chromosomes visible by electron microscopy in

prometaphase cells as plate-like structures ~200 nm in diameter, and attached to 20-30

microtubule filaments [18]. Kinetochores are composed of two main components: the inner

kinetochore centromere-binding CCAN complex, and the microtubule-binding KMN complex

of the outer kinetochore [19]. The 16-subunit CCAN complex is organized into five modules:

CENP-C and CENP-OPQUR, and the DNA-binding modules, CENP-LN, CENP-HIKM and CENP-

TWSX. The latter module comprises four histone fold domain proteins [20]. Cryo-EM

structures of human and budding yeast CCAN have elucidated mechanisms of centromere

and CENP-A nucleosome recognition [12,21-25]. CCAN encloses extranucleosomal linker DNA

within a topologically-enclosed chamber [12,22], a mechanism recently shown to be conserved

in the CENP-A-independent holocentric kinetochores of *Lepidoptera* [26]. In vertebrates,

interactions with the CENP-A nucleosome, that confer CCAN specificity for CENP-A rather

than H3 nucleosomes, are mediated primarily by the CENP-C subunit through its short linear

interaction motifs that bridge the CENP-A histone of CENP-A[Nuc] with the CCAN modules

CENP-LN and CENP-HIKM [12,21,23,27-30]. Human CENP-C incorporates two CENP-A-binding

motifs, generating four potential CENP-A-binding sites in the CENP-C dimer [31].

In our previous cryo-EM structures of CCAN:CENP-A[Nuc] and CCAN:DNA complexes, a DNA

duplex was observed threading through a 50 Å-long central tunnel running the length of the

CCAN molecule [12]. This DNA-binding tunnel is formed by the CENP-LN channel, and CENP-I

and CENP-TW subunits. As the DNA exits the tunnel it curves around the CENP-TWSX

module, interacting with the histone fold domains of CENP-TW. The DNA molecules used in

the earlier study (the 171 bp α-satellite DNA used to reconstitute the CENP-A[Nuc] for the

CCAN:CENP-A[Nuc] complex, and the 54 bp DNA used in the CCAN:DNA complex) were of

insufficient length to extend beyond CENP-TW, and so in these complexes DNA did not

contact CENP-SX of the histone-fold CENP-TWSX module. Previous cell biology and

biochemical studies however, had identified CENP-SX as DNA-binding proteins at the

centromere [32]. We were therefore missing important details regarding how CCAN interacts

with DNA at the centromere, and this information is required for understanding the higher-

order architecture of kinetochores arranged on α-satellite repeat arrays. Furthermore, the

repeating CCAN:CENP-A nucleosome unit at centromeres is not defined. Thus, the aims of

this study are two-fold. First, we aim to obtain a more complete understanding of how

CCAN interacts with DNA, and second we seek to provide models for how multiple copies of

CCAN and CENP-A nucleosomes assemble onto contiguous α-satellite repeat sequences at

human centromeres.

Results

<u>CCAN binds 70 bp of DNA in a loop configuration that wraps the CENP-TWSX module</u>

To understand how centromeric DNA interacts with CCAN, including all of the histone fold

domains (HFDs) of the CENP-TWSX module, we reconstituted CCAN with a longer DNA

duplex (human α-satellite DNA: 171 bp) and used mild cross-linking to stabilize the complex.

The cryo-EM structure, resolved to 3.3 Å resolution, showed clearly defined cryo-EM density

for ~70 bp of DNA that wraps around CCAN in a loop-like structure (**Fig. 1 and Extended**

Data Fig. 1 and Extended Data Table 1). As observed in our previous CCAN:CENP-A[Nuc] and

CCAN:DNA complexes [12], DNA threads through the enclosed CCAN chamber, contacting

CENP-TW as it exits the tunnel (**Fig. 1**). However, in the new structure, we observed an

additional 20 bp of DNA exiting the DNA-binding tunnel, so that in total 40 bp of DNA

interacts with the entire CENP-TWSX module. Additionally, a short segment of DNA (~5 bp)
extends beyond the CENP-TWSX module and approaches CENP-QU.
115

116
Fig. 1. CCAN binds 70 bp of DNA in a loop configuration. **a** and **b**, Orthogonal views of the
CCAN:DNA complex. 41 bp of DNA interacts with the CENP-TWSX module. Insert: Schematic of the
CCAN:DNA complex. **c**, Cryo-EM map of the CCAN:DNA complex with DNA coloured in gold and
CCAN subunits in grey. **d**, Electrostatic surface of CCAN as viewed in panel B, with DNA shown as a
ribbons representation.

The HFDs of the CENP-TWSX module assemble into a heterotetramer that structurally
resembles the H3:H4 tetramer of canonical nucleosomes (**Fig. 2a, b**) [11,32]. In the CCAN:DNA

complex, DNA wraps around CENP-TWSX comparably to how H3:H4 tetramers wrap DNA in
the context of an octameric nucleosome [11]. In both structures DNA has a similar curvature
and adopts a left-handed helical geometry. The CENP-TWSX module creates a positively-
charged ridge that binds the DNA phosphate backbone (**Fig. 1d**). Strikingly, DNA wraps the
H3:H4 and CENP-TWSX tetramers with identical registers such that the DNA major groove
projects towards both the dyad axis of the H3:H4 tetramer, and the related pseudo-dyad
axis of CENP-TWSX (**Fig. 2a, b**). Consequently, in both structures, at superhelical locations
(SHLs) ±0.5, ±1.5, ±2.5, the DNA minor groove projects towards and contacts equivalent
residues in the histone fold domains of the H3:H4 tetramer and CENP-TWSX module. Our
structure shows that DNA interacts with conserved basic residues of all four HFD domains of
CENP-TWSX (**Fig. 2c, d**), consistent with prior mutagenesis data demonstrating the role of
basic CENP-TWSX amino acids in DNA binding *in vitro* and centromere localization *in vivo*
[20,32].

Fig. 2. The CENP-TWSX module binds DNA analogous to the H3:H4 histone tetramer of canonical
nucleosomes. a, CENP-TWSX and CCAN-associated 70 bp DNA, 40 bp interacts with the CENP-TWSX

module. **b**, H3:H4 tetramer of a nucleosome with almost with 67 bp of DNA. Super helical locations
(SHLs) are indicated. The pseudo dyad axes of CENP-TWSX and the H3:H4 tetramer are shown in A
and B (PDB: [11]). **c**, Details of basic residues from the HFDs of CENP-S and CENP-X and CENP-S C-
terminal α-helix interacting with the DNA phosphate backbone. **d**, Details of basic residues from the
HFDs of CENP-T and CENP-W, and CENP-T N-terminal extension interacting with the DNA phosphate
backbone. In (C) and (D) residues previously shown to contribute to CENP-TWSX – DNA binding are
underlined [32]. **e**, Multiple sequence alignment of the CENP-S C-terminal α-helix. Alignment created
using Jalview [33].

Additional protein-DNA contacts stabilize CENP-TWSX – DNA interactions. These involve an

extended basic segment immediately N-terminal of the CENP-T HFD that inserts into the

DNA minor groove, and a C-terminus α-helix of CENP-S that engages the DNA major groove

(**Fig. 2a, c and d**). This CENP-S α-helix becomes ordered on binding DNA, such that a

conserved stretch of basic residues stabilize its interactions with the DNA backbone (**Fig. 2c,**

e), as similarly observed in the crystal structure of a CENP-SX:DNA complex [34]. Our structural

data are consistent with previous studies showing that deletion of the C-terminus of CENP-

S, including this basic region, abolished CENP-S binding to DNA *in vitro* [32]. Similar

observations regarding the interaction of linker DNA with the CENP-TWSX module were

obtained by Musacchio and colleagues from a cryo-EM structure of a CCAN:DNA complex

(M. Pesenti, I. Vetter and A. Musacchio, unpublished data).

<u>DNA of CENP-A^{Nuc} reconstituted with longer α-satellite repeat DNA wraps CENP-TWSX</u>

We next assessed whether CENP-TWSX wraps linker DNA in the context of a CENP-A

nucleosome. We assembled CCAN onto a CENP-A^{Nuc} reconstituted with 211 bp DNA that

comprised the 171 bp α-satellite repeat and 40 bp from the 5' upstream α-satellite repeat,

and determined its structure using cryo-EM to an overall resolution of 4.5 Å (**Extended Data**

Fig. 2). Both individual CCAN:DNA and CENP-A^{Nuc} modules were resolved in cryo-EM density,

showing relative flexibility (**Extended Data Fig. 2d**). A total of 200 bp DNA was visible in the

cryo-EM map. Similar to our previous CCAN:CENP-A^{Nuc} structure with 171 bp DNA [12], CENP-

A^{Nuc} is positioned at the 3' end of the α-satellite repeat (**Fig. 3a, c**). The 5' terminus of the

CENP-A^{Nuc} DNA gyre is unwrapped by 10 bp. This unwrapped DNA then leads into

extranucleosomal linker DNA that threads through the CCAN DNA-binding tunnel. On exiting

the tunnel, DNA wraps around the entire CENP-TWSX module, as seen in the CCAN:DNA

complex described above. This DNA comprises linker DNA of the 3' downstream α-satellite

repeat together with 30 bp of the 5' upstream α-satellite repeat. Thus, a total of ~65 bp of

free DNA 5' of CENP-A^{Nuc} interacts with CCAN (**Fig. 3c**). An unexpected finding was that

compared with our previous CCAN:CENP-A^{Nuc} cryo-EM structure [12], CENP-A^{Nuc} adopts a

different orientation relative to CCAN. In our new structure, CENP-A^{Nuc} is rotated by 216°

and shifted by six bp away from CCAN. A shift of register while maintaining the same DNA

phosphate backbone interactions with the CCAN DNA-binding tunnel causes the DNA duplex

(and hence CENP-A^{Nuc}) to rotate 36° per bp, rotating 216° overall (**Fig. 3b and**

Supplementary Video 1). The reasons for the differences in CENP-A^{Nuc} position between the

two CCAN:CENP-A^{Nuc} complexes are not entirely clear. A possible explanation is that the

longer extranucleosomal DNA linker used in this study optimizes interactions with DNA-

binding sites on CCAN, resulting in displacement of CENP-A^{Nuc}. Cryo-EM density for CENP-C

is observed at the CENP-C-binding sites on CENP-A^{Nuc}, CENP-LN and CENP-HIKM, however,

due to the displaced CENP-A^{Nuc}, there are no direct contacts between CENP-A^{Nuc} and the

back-side of CCAN.

190
191
Fig. 3. Cryo-EM structure of CCAN:CENP-A^{Nuc} showing CENP-TWSX binding DNA. a, Ribbons

representation of the CCAN:CENP-A^{Nuc} complex. DNA coloured in red indicates DNA of the 5'

upstream α-satellite repeat. **b**, Superimposition of our new CCAN:CENP-A^{Nuc} complex (with 211 bp

DNA) (coloured as in panel A) onto the previous model of CCAN:CENP-A^{Nuc} with a single 171 bp α-

satellite repeat DNA (coloured grey) [12]. This shows the difference in position and orientation of the 3'

CENP-A^{Nuc}. The two nucleosomes differ in orientation by 216° due to a 6 bp translation of the repeat

sequence through the tunnel (**Supplementary Video 1**). **c**, Schematic of the CCAN:CENP-A^{Nuc} on an

α-satellite repeat dimer. A total of 200-210 bp of DNA is embedded by the complex including 30 bp

from the 5' upstream α-satellite repeat. A B-box accessible for CENP-B binding in the upstream

repeat is shown in blue.

Although the cryo-EM density for the DNA phosphate backbone is well resolved, density for

individual DNA bases is less well defined such that purine and pyrimidine bases are not

distinguished. We therefore do not observe sequence-specific protein-DNA interactions. In

conclusion, these new structures of CCAN with both free DNA and a CENP-A^{Nuc} with an

extended extranucleosomal DNA linker are consistent with earlier proposals that the CENP-
TWSX module of the kinetochore functions as a nucleosome-like particle at the centromere
[20,32].

Implications for CCAN binding to α-satellite repeat dimers

Our new CCAN:DNA and CCAN:CENP-A[Nuc] structures showing that CCAN itself interacts with
65 bp of duplex DNA has implications for understanding the higher-order organization of
CCAN on repetitive α-satellite repeat CENP-A nucleosome arrays. The 65 bp DNA interacting
with CCAN comprises 30 bp from the 5' upstream α-satellite repeat and 35 bp of the 3'
downstream α-satellite repeat: 24 bp of extranucleosomal linker DNA and 10 bp of
unwrapped CENP-A[Nuc] DNA (**Fig. 3c**). The 30 bp DNA interacting with CCAN from the 5'
upstream α-satellite repeat is DNA of the CENP-A nucleosome gyre when CENP-A
nucleosomes are positioned at the 3' end of the α-satellite repeat [12,13]. This suggests that in
the context of a CCAN:CENP-A[Nuc] complex, the upstream α-satellite repeat might be devoid
of a nucleosome (either CENP-A or canonical H3). In alternative non-exclusive scenarios, to
accommodate CCAN bound between two CENP-A nucleosomes on adjacent α-satellite
repeats, a CENP-A nucleosome positioned on the upstream 5' α-satellite repeat might
unwrap by 30 bp, or the upstream CENP-A[Nuc] may force DNA to unwrap from the CENP-
TWSX module. A third possibility is that CCAN repositions CENP-A[Nuc] by causing DNA to slide
around the histone octamer. However, the defined position of CENP-A[Nuc] at the 3' of α-
satellite repeats *in vivo* [13] does not support this scenario.

Substantial evidence indicates that the DNA termini of CENP-A nucleosomes are loosely
wrapped and flexible. First, nuclease digestion sensitivity assays showed that CENP-A
nucleosomes protect only 100-120 bp DNA, compared to 140-150 bp for canonical H3
nucleosomes [13,24,35,36]. Second, FRET data indicated increased flexibility of terminal DNA
segments in CENP-A[Nuc] relative to canonical nucleosomes [16]. Third, both crystal and cryo-EM
structures of CENP-A[Nuc] revealed unwrapping of terminal DNA segments [12,16,24,37-41], in some
instances as much as 24-36 bp of flexible DNA at one end [42]. Finally, CENP-B and CENP-C
have been implicated in promoting DNA unwrapping from CENP-A[Nuc] (Refs [16,43,44]).

CCAN binds a dimeric α-satellite repeat with two CENP-A nucleosomes

To investigate the feasibility of CCAN binding to linker DNA connecting two adjacent CENP-A
nucleosomes, we reconstituted two CENP-A nucleosomes on an α-satellite repeat dimer.
The α-satellite repeat dimer was designed to include only linker DNA connecting the two

CENP-A nucleosomes: upstream and downstream α-satellite repeats were 147 bp and 171

bp, respectively (**Extended Data Fig. 3a**). To stabilize the two nucleosome positions on this

324 bp DNA, we replaced 92 bp of native α-satellite repeat DNA at the centre of each CENP-

A[Nuc] site with Widom601 DNA. We observed that CCAN associated with this di-CENP-A[Nuc]

array, as assessed by size exclusion chromatography (SEC) (**Extended Data Fig. 3b, c**).

Analysis of the peak SEC fractions using interferometric scattering microscopy (iSCAT)

revealed two species: the major 1,054 kDa complex corresponded to a single CCAN

promoter bound to the di-CENP-A[Nuc] array (CCAN:di-CENP-A[Nuc]), although with a lower mass

than the expected 1,282 kDa. The underestimate of molecular mass using ISCAT may be due

to the DNA component which scatters less than the globular proteins used for calibration. A

445 kDa component corresponded to a di-CENP-A[Nuc] array without CCAN (expected

molecular mass of 431 kDa) (**Extended Data Fig. 3d**). The partial dissociation of CCAN from

di-CENP-A[Nuc] is likely due to the low sample concentrations required for single molecule

counting in iSCAT.

CCAN and CENP-B binding to the same linker DNA segment is mutually exclusive

Our α-satellite dimer incorporated a B-box within the linker DNA of the downstream α-

satellite repeat connecting the two CENP-A nucleosomes (**Extended Data Fig. 4a**). SEC data

showed that the CENP-B protein recognized the B-box specifically because CENP-B bound to

the di-CENP-A[Nuc] array with an intact B-box, but not to a di-CENP-A[Nuc] array with a mutant B-

box (**Extended Data Fig. 4b, c**). We next tested whether CCAN and CENP-B bind the di-CENP-

A[Nuc] array simultaneously. We observed that CCAN displaced CENP-B from the array,

showing that CCAN and CENP-B engagement of linker DNA is mutually exclusive (**Extended

Data Fig. 4d, e**). This is consistent with our model of CCAN:CENP-A[Nuc] (Ref. [12]), that with

CCAN bound to linker DNA, the B-box is enclosed by the CCAN DNA-binding tunnel and

therefore inaccessible to CENP-B.

<u>CCAN binds to linker DNA in the dimeric α-satellite repeat with two CENP-A nucleosomes</u>

To understand the structural basis of how CCAN interacts with linker DNA connecting

adjacent CENP-A nucleosomes, we prepared cryo-EM grids with the CCAN:di-CENP-A[Nuc]

complex. Despite cross-linking and the use of additives to reduce sample denaturation at

the air-water interface, CCAN:di-CENP-A[Nuc] particles were not visible on either cryo-electron

micrographs or subsequent 2D classes of the cryo-EM dataset. We instead observed a

heterogeneous mixture of free DNA, CENP-A[Nuc] and protein complexes (data not shown),

indicative of CCAN:di-CENP-A[Nuc] denaturation on cryo-EM grids. To stabilize the CCAN:di-

CENP-A[Nuc] complex for cryo-EM analysis, we combined it with the CENP-A nucleosome

stabilizing-single chain antibody fragment (ScFv) [40], omitting CENP-C due to its binding site

overlapping that of the ScFv (**Extended Data Fig. 5)**, and collected a large cryo-EM dataset

on this ScFv-stabilized CCAN:di-CENP-A^{Nuc} complex (**Extended Data Figs 6 and 7 and**

Extended Data Table 1).

2D classification of the selected particles and subsequent 3D reconstruction and

classification revealed a highly heterogenous mixture of molecular states (**Extended Data**

Figs 6b-d and 7). The majority of 2D classes from nearly nine million picked particles

comprised either free mono-CENP-A nucleosome or CCAN:DNA with no obvious associated

CENP-A nucleosome. *Ab initio* reconstruction and 3D refinement resulted in high resolution

maps of CCAN:DNA and a CENP-A mono-nucleosome with bound ScFv (2.9 Å and 2.7 Å,

respectively; **Extended Data Fig. 7**). The CCAN reconstruction showed DNA enclosed within

the DNA-binding tunnel. We re-extracted CCAN particles in RELION [45] using a larger box and

performed further 3D classification (**Extended Data Fig. 7**). This procedure revealed 3D

classes with diffuse, poorly-defined EM densities at both 3' and 5' entrances to the CCAN-

DNA tunnel that likely represented flexible CENP-A nucleosomes. To determine whether the

density at the 3' entrance corresponded to a 3' CENP-A nucleosome, we generated a wide

mask covering this density and performed signal-subtraction of density outside the mask.

Subsequent 3D reconstruction and classification of the mask-generated recentred particles

recovered a CENP-A nucleosome, confirming that the diffuse density observed in 2D classes

as CENP-A^{Nuc}. Using the corresponding unmasked particle set, we then calculated a new 3D

reconstruction at 9.5 Å resolution that revealed both CCAN and 3' CENP-A^{Nuc}. Consistent

with this reconstruction, 2D classes calculated from this particle set showed density features

for CCAN and 3' CENP-A^{Nuc}. A similar approach was used to analyse EM density at the 5'

entrance to the CCAN-DNA tunnel, resulting in a 3D reconstruction with CCAN and

associated 5' CENP-A^{Nuc} at 15 Å resolution (**Extended Data Fig. 7**). In the 2D class galleries

corresponding to both the 3' CCAN:CENP-A^{Nuc} and 5' CCAN:CENP-A^{Nuc} complexes, we

observed diffuse densities at the opposite end of the CCAN DNA-tunnel (i.e. 5' and 3',

respectively), for example **Extended Data Fig. 7i**. These particles are consistent with CCAN

bound to the linker DNA of the di-CENP-A^{Nuc} array. However, the small number of the

particles precluded a 3D reconstruction. We therefore generated a composite cryo-EM map

of a CCAN:di-CENP-A^{Nuc} complex from the individual 3' CCAN:CENP-A^{Nuc} and 5' CCAN:CENP-

A^{Nuc} maps (**Extended Data Fig. 7i** and **Fig. 4b**). One of its projected views matches an

experimental 2D class of a CCAN:di-CENP-A^{Nuc} , validating our composite approach

(**Extended Data Fig. 7i**).

319
320
Fig. 4. Structure of the CCAN:di-CENP-A^{Nuc} complex. **a**, Ribbons representation of CCAN:di-CENP-
A^{Nuc} complex. Insert: Schematic of the CCAN:di-CENP-A^{Nuc} on a dimer of α-satellite repeats. 20-25 bp
of the upstream CENP-A nucleosome is unwrapped. A total of 300-310 bp of DNA is embedded by
the complex. A B-box accessible for CENP-B binding in the upstream repeat is shown in blue. **b**, Cryo-
EM map of CCAN:di-CENP-A^{Nuc} complex. **C**. Superimposition of the new CCAN:CENP-A^{Nuc} and
CCAN:di-CENP-A^{Nuc} complexes showing the identical position of the 3' CENP-A^{Nuc}.

CCAN bound to linker DNA forces DNA unwrapping from CENP-TWSX and 5' CENP-A^{Nuc}

5' CCAN:CENP-A^{Nuc} complex: Despite the limited resolution of this map, reliable alignment of

CENP-A^{Nuc} into cryo-EM density was facilitated by the ScFv associated with both sides of the

nucleosome (**Fig. 4b**). Analysis of this structure indicated that a CENP-A nucleosome

positioned 5' to CCAN is incompatible with DNA wrapping around the CENP-TWSX module

(**Fig. 4a, b)**. Analogous to our previous CCAN:CENP-A[Nuc] and CCAN:DNA complexes [12], we

observe that as DNA exits the CCAN-DNA tunnel, it runs along the CENP-I-CENP-TW channel,

although unlike our new CCAN:DNA and CCAN:CENP-A[Nuc] complexes (this study), does not

continue to bend around CENP-SX, but instead projects in a straighter trajectory to join the

DNA of the 5' CENP-A nucleosome. The DNA gyre of the 5' CENP-A nucleosome connected

to linker DNA is itself unwrapped by 20-25 bp. Thus, even without DNA wrapping around

CENP-SX, the 5' CENP-A nucleosome is forced to unwrap to allow its positioning onto an

upstream α-satellite repeat. Our fitting of the 5' CENP-A[Nuc] to the cryo-EM density map is

consistent with the α-satellite repeat length of 171 bp such that the DNA length from the

end of the downstream (3') α-satellite repeat sequence to the dyad axis of the 5' CENP-A[Nuc]

is 74 bp (**Fig. 4a**).

3' CCAN:CENP-A[Nuc] complex: The position and orientation of the 3' CENP-A[Nuc] in the

CCAN:di-CENP-A[Nuc] complex is the same as our new CCAN:mono-CENP-A[Nuc] complex with

211 bp DNA (this study), differing from that in the previously published CCAN:CENP-A[Nuc]

complex [12] (**Fig. 4a, c**). Thus, the ScFv used for the CCAN:di-CENP-A[Nuc] structure

determination does not affect the orientation of the 3' CENP-A[Nuc], consistent with modelling

showing that ScFv binding to CENP-A[Nuc], when in the mono-CENP-A[Nuc] position, is possible

without a steric clash with CCAN.

Discussion

In this study we investigated the molecular basis for how human CCAN interacts with CENP-

A nucleosomes and DNA of α-satellite repeat arrays, revealing that CCAN engages as much

as 70 bp of DNA. As described previously [12], DNA threads through the enclosed tunnel

generated by the CENP-LN module, CENP-I and CENP-TW subunits. The longer DNA

sequences used in this study showed additional DNA interactions with the CENP-SX subunits

of the CENP-TWSX HFD module. The CENP-TWSX subunits interact with DNA analogously to

how the H3:H4 heterotetramer, in the context of octameric nucleosomes, wraps its DNA

gyre. The specific details of the interactions of DNA with CENP-SX, together with CENP-TW,

observed in our structure, are in agreement with biochemical and cell biology data,

confirming the proposal that human CCAN incorporates a nucleosome-like particle at the

centromere [20,32]. Consistent with this, CENP-TWSX augments the affinity of CCAN for a

CENP-A nucleosome three-fold [12]. Thus, this study indicates that CCAN creates a tight

attachment to centromeric DNA by both topological entrapment of DNA through the CCAN

DNA-binding tunnel (CENP-LN, CENP-I), and by wrapping DNA around a HFD tetramer

(CENP-TWSX), mimicking the mechanism by which nucleosomes package DNA.

The finding that CCAN alone engages 65-70 bp of DNA has implications for the higher-order

architecture of CCAN and CENP-A nucleosomes on repetitive α-satellite sequences. The 65

bp of extranucleosomal DNA bound by CCAN greatly exceeds the 24 bp linker DNA

connecting two adjacent and fully wrapped nucleosomes (147 bp) on tandem α-satellite

repeats. In our CCAN:CENP-A^{Nuc} structures, 10 bp of DNA are unwrapped from the 3' CENP-

A nucleosome gyre as it enters the CCAN DNA-binding tunnel ([12] and this study). Thus, to

accommodate 65 bp of DNA interacting with CCAN requires ~30 bp DNA of the upstream α-

satellite repeat, in addition to the 24 bp of α-satellite extranucleosomal linker DNA and 10

bp of unwrapped DNA from the downstream CENP-A nucleosome (**Fig. 3c**). Since CENP-A

nucleosomes are positioned exactly on α-satellite repeats such that the 3' end of the repeat

defines the start of the CENP-A nucleosome DNA gyre [12,13], the 30 bp of upstream α-satellite

repeat DNA would require the upstream CENP-A nucleosome to unwrap 30 bp of DNA. Such

unwrapping is potentially feasible as demonstrated by the extensive levels of DNA

unwrapping from CENP-A nucleosomes documented previously [12,16,24,37-42]. To test whether

CCAN associates with DNA connecting two adjacent CENP-A nucleosomes, we reconstituted

a dimeric α-satellite repeat with CENP-A nucleosomes. SEC and iSCAT data confirmed

interactions of CCAN with a dimeric α-satellite repeat with two CENP-A nucleosomes

(**Extended Data Fig. 3**). That this interaction required the free linker DNA connecting the

two nucleosomes was confirmed by the ability of CCAN to displace CENP-B from its B-box

binding site located within the linker DNA (**Extended Data Fig. 4d, e**). Our cryo-EM structure

of the CCAN:di-CENP-A^{Nuc} complex revealed CCAN bound to DNA linking two CENP-A

nucleosomes (**Fig. 4**). However, accommodating CCAN at this site involved not only 20-25 bp

of DNA to be unwrapped from the 5' upstream CENP-A nucleosome, but also dissociation of

DNA from CENP-SX. Thus, although the upstream CENP-A nucleosome is unwrapped, the

available free DNA duplex is insufficient to bind CENP-SX, presumably because of steric

hindrance between the upstream CENP-A nucleosome and CCAN.

Whether such an arrangement of tandem CENP-A nucleosomes and central CCAN that we

observe in our CCAN:di-CENP-A^{Nuc} complex occurs at native human centromeres is unclear.

The dissociation of DNA from CENP-SX is inconsistent with the role of conserved basic

residues of CENP-SX in binding DNA revealed by our CCAN:DNA and CCAN:mono-CENP-A^{Nuc}

structures and from prior *in vitro* binding assays [32]. Furthermore, the extensive unwrapping

of DNA from a CENP-A nucleosome, although observed for a small population of particles in

a cryo-EM study [42], is not consistent with CENP-A nucleosomes defining a 120 bp footprint *in*

vivo [13,46], and the size of the nuclease-resistant core observed *in vitro*. Although our cryo-EM

structure of the CCAN:di-CENP-A^{Nuc} complex embeds a total of 300-310 bp (**Fig. 4a**), in

agreement with the 250-300 bp centromere core-selective footprint identified by single

molecule chromatin fibre sequencing (Fiber-seq) [46], we propose two alternative scenarios

for the arrangement of CCAN and CENP-A nucleosomes on α-satellite arrays. In one

scenario, the kinetochore unit would constitute a monomeric CCAN:CENP-A^{Nuc} complex in

which DNA from the upstream α-satellite repeat engages CENP-TWSX such that a total of

200-210 bp of DNA is bound by the CCAN:CENP-A^{Nuc} complex (**Fig. 3**). In a second related

scenario, a central CENP-A nucleosome would interact with two flanking CCAN protomers in

a pseudo-symmetric arrangement of the mono-CCAN:CENP-A^{Nuc} complex (di-CCAN:CENP-

A^{Nuc}) (**Fig. 5**). This proposed assembly, with the central CENP-A nucleosome and associated

CCAN protomers spanning three α-satellite repeats, embeds 260-270 bp. The 210 bp bound

by mono-CCAN:CENP-A^{Nuc} is shorter than the 250-300 bp centromere core-selective

footprint [46], however the 260-270 bp bound to the proposed di-CCAN:CENP-A^{Nuc} complexes

matches the reported 250-300 bp centromere footprint. We note that Dubocanin et al.,

(2025) also identified centromere-associated footprints of 210 bp in size [46]. Although these

were assigned as di-nucleosomes, it is also plausible that these methylation-free footprints

correspond to mono-CCAN:CENP-A^{Nuc} assemblies. Consistent with this, a recent cryo-ET

study of human centromeres found that the majority of putative kinetochore particles were

consistent in size with a mono-CCAN complex closely associated with a single CENP-A

nucleosome [47]. Additionally, larger globular densities containing multiple nucleosomes and

CCAN complexes were also observed, although whether these constitute tandem CCAN and

CENP-A^{Nuc} complexes was not defined.

Fig. 5. Model of a di-CCAN:CENP-A^Nuc complex. The model was generated from the CCAN:mono-CENP-A^Nuc structure by superimposing on CENP-A^Nuc. The two CCANs together with a single CENP-A^Nuc overlap three α-satellite repeats. In the model, DNA from the central α-satellite sequence is coloured gold. DNA bound to the two CCANs from the 5' and 3' α-satellite repeats (repeats 1 and 3, respectively) is coloured red. Insert: Schematic of the model with DNA coloured red for the 5' and 3' repeats as in the model. A B-box accessible for CENP-B binding in the upstream repeat is shown in blue. B-boxes in the two upstream repeats are occluded by CCAN. 260-270 bp of DNA are bound by the di-CCAN:CENP-A^Nuc complex.

Finally, in this study we show that the association of CCAN and CENP-B to the same extranucleosomal linker DNA is mutually exclusive (**Extended Data Fig. 4**). As indicated in Figures 3 to 5, for all three models of CCAN:CENP-A^Nuc complexes discussed above, a B-box in the upstream α-satellite repeat is accessible for CENP-B binding.

In summary this work, combined with recent 2D and 3D mapping of CCAN, CENP-A nucleosomes and DNA-binding proteins at native centromeres [46,47], suggest potentially heterogenous structures of CCAN and CENP-A nucleosomes assembled on α-satellite repeat arrays including mono-CCAN:CENP-A^Nuc, di-CCAN:CENP-A^Nuc and CCAN:di-CENP-A^Nuc complexes.

Supplementary Video 1. Video illustrates the difference in position of the 3' CENP-A nucleosome between our previous CCAN:CENP-A^Nuc complex with 171 bp DNA (grey cartoon) [12] and the new structure of CCAN:CENP-A^Nuc with 211 bp DNA (ASW6) (colour-coded according to subunits as in Fig. 1). Relative to the previous CCAN:CENP-A^Nuc structure, in the new model, the DNA register of the α-satellite repeat is shifted 6 bp (translation of 20

Å) away from CCAN (towards the 3' end). Because the DNA phosphate backbone contacts

with CCAN are preserved, this results in a 216° rotation of CENP-A^{Nuc}.

Methods

<u>DNA cloning</u>

Protein complexes for CCAN and CENP-A histone octamer were prepared as previously

described [12], except CENP-Q was modified to CENP-Q$^{55-268}$, CENP-U was modified to CENP-

U$^{235-418}$, CENP-I was changed to isoform 1 from previously isoform 2, and CENP-B full length

was cloned into pOPINS plasmid, with a double strep II-sumo-tev tag at their N-termini. The

DNA fragment ASW14 was designed as two repeats from α-satellite DNA of chromosome 2

with 91 base pairs centred on their dyad positions replaced with Widom601 sequence

aligned at the dyad base. A B-box sequence is in the linker region. Six ASW14 fragments

flanked by GATATC (EcoRV recognition site) were assembled into the pMK-RQ plasmid. The

plasmid containing ASW14 fragments was produced by using Plasmid Plus Giga Kit

(QIAGEN). The DNA fragment was released by EcoRV-HF (NEB) at 37C over night. The

digested sample was diluted five-fold with PBS and loaded onto a pre-equlibrated (PBS) 1 mL

Resource Q Column (GE 17-1177-01, Merck). The column was washed with 10 mL PBS

(buffer A), followed by a wash with 20 mL of 50% of buffer B (PBS with 850 mM NaCl). The

DNA fragment was eluted using a gradient to 70% of buffer B at 1 mL.min^{-1} over 120 min.

The target fragment eluted around 60% buffer B. The fractions containing DNA were checked

on an agarose gel, and DNA was precipitated using 1:1 of isopropanol at 20 °C for 30 min at

15,000g. The DNA pellet was dissolved in the buffer needed for its following application.

The sequence of the generated ASW14 fragment is:

482
```
483  1     ATCAGGAAAT ATCTTCACAT AAAAACTAGA CAATTGGTCG TAGACAGCTC TAGCACCGCT
484  61    TAAACGCACG TACGCGCTGT CCCCCGCGTT TTAACCGCCA AGGGGATTAC TCCCTAGTCT
485  121   CCTTGTAGTA TCTGCAAGTG GACATTCGGA GCACTTTGTG GCCTTCGTTG GAAACGGGAA
486  181   TATGTTCACA TAAAAACTAG ACAATTGGTC GTAGACAGCT CTAGCACCGC TTAAACGCAC
487  241   GTACGCGCTG TCCCCCGCGT TTTAACCGCC AAGGGGATTA CTCCCTAGTC TCCTTGTAGT
488  301   ATCTGGAATT GGACATTTGG AGAT
```
489

The Widom601 sequence underlined. Dyad base of the two CENP-A^{Nuc} indicated in italics.

ASW14 with mutant B-box: The B-box of CTTCGTTGGAAACGGGA in bold of ASW14 was

mutated to CAAAATTGGTAATTTTA in the B-box mutant α-satelitte repeat dimer.

The 211 bp DNA fragment for the CCAN:CENP-A^{Nuc} complex (ASW6) has an additional 40 bp

5' of an α-satellite repeat sequence (AS2) with a central 92 bp Widom601 sequence. ASW6

sequence is:

497
1   CATACTCTTT TTGTAGTATC TGCAAGTGGA CATTCGGAGC ACTTTGTGGC CTTCGTTGGA
61   AACGGGAATA TGTTCACATA AAAACTAGAC AATTGGTCGT AGACAGCTCT AGCACCGCTT
121   AAACGCACGT ACGCGCTGTC CCCCGCGTTT TAACCGCCAA GGGGATTACT CCCTAGTCTC
181   CTTGTAGTAT CTGGAATTGG ACATTTGGAG AT

<u>CCAN:DNA complex assembly</u>

CCAN:DNA complexes were assembled in CCAN buffer (300 mM NaCl, 20 mM Hepes pH 7.8,

0.5 mM TCEP) as follows. 6 µM CCAN was incubated at 22 °C for 30 mins before addition of

an equal volume of 6 µM 171 bp (α-satellite repeat) DNA (AS2 described in [12]), and

incubated for a further 30 mins at at 22 °C, resulting in a final CCAN:DNA concentration of 3

µM and a final volume of 100 µL. The CCAN:DNA complex was then centrifuged at 10 °C in a

table-top centrifuge for 10 min. To stabilize complexes for cryo-EM analysis, on-column

crosslinking was performed.

<u>On-column crosslinking of CCAN:DNA</u>

An Agilent 500 column coupled to a MicroAkta was pre-equilibrated in CCAN buffer. 100 µL

2.5 mM BS(PEG)$_5$ (ThermoFisherScientific) was injected into the system for 1 mL at a flow

rate of 0.07 µL/min. The CCAN:DNA complex was then injected into the column and eluted

at the same flow rate into a 96-well plate in 50 µL fractions. Each plate well was

supplemented with 5 µL 1 M Tris.HCl buffer pH 7.4 to quench the crosslinker during elution.

Fractions containing CCAN:DNA were pooled, buffer exchanged into 150 mM NaCl, 20 mM

Hepes pH 7.8, 0.5 mM TCEP and concentrated to 1.5 mg.mL^{-1} (Amicon 100 kDa, 0.5 mL).

Following concentration, CCAN:DNA complexes were again centrifuged at 10 °C in a table-

top centrifuge for 10 min.

<u>CENP-A mono-nucleosome (ASW6) and CENP-A di-nucleosome (ASW14) reconstitution</u>

The CENP-A histone octamer was prepared as described previously [12]. Either ASW6 or

ASW14 DNA were desalted into buffer B (20 mM HEPES pH 7.5, 2 M KCl). DNA at 8 µM was

mixed with the CENP-A histone octamer at molar ratios of 1.1:1 protein:DNA for mono-

CENP-A^{Nuc} and 2.1:1 protein:DNA for di-CENP-A^{Nuc}. The histone octamers were then wrapped

with DNA using the gradient dialysis method starting from 2 M KCl (in 20 mM HEPES pH 7.5,

1 mM EDTA) to 300 mM KCl over 16 h at 22 °C. The mixture was then dialysed against 20

mM HEPES pH 7.5, 300 mM KCl, 1 mM EDTA for a further 5 h. The wrapped mono-CENP-A

and di-CENP-A nucleosomes were assessed by agarose gel and stored at 4 °C.

Assembly of the CCAN:CENP-A^{Nuc} (ASW6 DNA) complex

CENP-A^{Nuc} and CCAN components (CENP-C^N, CENP-LN, CENP-HIKM, CENP-TWSX, and CENP-OPQUR$^{\Delta N}$) were mixed at a 1:1 molar ratio at 5 μM in PBS buffer and TCEP. The sample was injected into an Agilent Bio SEC-5 1000A 4.6x300 column in CCAN gel filtration buffer (20 mM HEPES pH 7.8, 300 mM NaCl, 1 mM TCEP). The peak fractions were analysed using a 4-12% SDS-PAGE gel, stained with Instant Blue Coomassie Stain. To prepare the complex for structural analysis, on-column cross-linking using BS(PEG)$_5$ (ThermoFisherScientific) was performed. 100 μl of 2.5 mM BS(PEG)$_5$ was injected into an Agilent Bio SEC-5 1000A 4.6x300 column (equilbrated in PBS buffer) at a flow rate set of 0.07 μL/min, then 0.2 mL buffer, after which the CCAN:CENP-A^{Nuc} sample in (100 μl) was applied and eluted in PBS buffer. The sample was eluted into a 96-well plate in 50 μL fractions. Each plate well was supplemented with 5 μL 1 M Tris.HCl buffer pH 7.4 to quench the crosslinker during elution. The peak fractions of the CCAN:CENP-A^{Nuc} samples were pooled and concentrated to 1.5 mg.mL^{-1} for cryo-EM grid preparation.

Assembly of the CCAN:di-CENP-A^{Nuc} (ASW14 DNA) complex

For assembly of the CCAN:di-CENP-A^{Nuc} (ASW14 DNA) complex, we used the same procedure as for the CCAN:mono-CENP-A^{Nuc} (ASW6 DNA) complex except we used 20 mM HEPES pH 7.8, 300 mM NaCl, 1 mM TCEP buffer in the cross-linking final gel filtration step.

Assembly of the CCAN$^{\Delta CENP-C}$:di-CENP-A^{Nuc} (ScFv) (ASW14 DNA) complex

 1. *Cross-linking di-CENP-A^{Nuc} and ScFv (di-CENP-A^{Nuc}:ScFv)*
Di-CENP-A^{Nuc} and ScFv were combined at a 1:2.3 molar ratio and dialysed into a buffer of 20 mM HEPES pH 7.5, 50 mM NaCl. Cross-linking of the di-CENP-A^{Nuc}:ScFv complex was performed using a 24 mL Superose 6 column by injecting 500 μL of 2.5 mM BS(PEG)$_5$ at a flow rate of 0.25 mL.min^{-1} for 7 mL before loading the CENP-A^{Nuc}:ScFv complex onto the column at the same flow rate. The cross-linking was assessed using SDS-PAGE gel. The cross-linked fractions were pooled and concentrated to 10 μM.

 2. *Assembly of cross-linked di-CENP-A^{Nuc}:ScFv with CCAN*
All CCAN components (omitting CENP-C^N) were without DTT. The components were at a 1:1:1:1:1 molar ratio (CENP-OPQUR:CENP-LN:CENP-HIKM:CENP-TWSX:di-CENP-A^{Nuc}:ScFv) with CCAN components at a concentration of 5 μM. The resultant complex was cross-linked

on an Agilent Bio SEC-5 1000A 4.6x300 column. Collected fractions were concentrated to

1.5-2 mg.mL^{-1} for cryo-EM grid preparation.

CCAN displaces CENP-B from di-CENP-A^{Nuc}-B-box

Di-CENP-A^{Nuc}-CENP-B binding test. Di-CENP-A^{Nuc} and CENP-B were mixed at a 1:1 ratio at 5

µM final concentration and injected into a Superose 6 Increase 10/300 column (Cytiva).

Appropriate fractions were collected and concentrated.

Di-CENP-A^{Nuc}-CENP-B-CCAN binding test. An equal molar ratio of CCAN$^{\Delta CENP-C}$ components

(CENP-LN, CENP-HIKM, CENP-TWSX, and CENP-OPQUR$^{\Delta N}$) was added to the di-CENP-

A^{Nuc}:CENP-B complex at a final concentration of 5 µM and then loaded onto an Agilent Bio

SEC-5 1000A 4.6x300 column. The eluted fractions were visualized using a 4-12% SDS-PAGE

gel, stained with InstantBlue Coomassie Stain.

Mass photometry

To assess the oligomeric state of CCAN:di-CENP-A^{Nuc} and CCAN$^{\Delta CENP-C}$:di-CENP-A^{Nuc}, 18 µL of

sample at approximately 100 nM was applied to wells in silicone gaskets on clean coverslips

(Refeyn, UK) on a OneMP instrument (Refeyn). Movies were imaged for 60 s using the

AcquireMP software (Refeyn) and the results were processed using the DiscoverMP

software. The data were analysed by using automatic Gaussian fitting. The contrast

intensities of landing events due to interference scattering were converted to masses using a

calibration performed with a standard set of protein oligomers (P1, Refeyn) under similar

buffer conditions. Data events were exported and plotted in Prism (GraphPad).

Cryo-EM grid preparation and data collection

CCAN:DNA sample vitrification

Immediately prior to plunge-freezing, CCAN:DNA samples were supplemented with n-octyl-

β-D-glucopyranoside (Glycon-biochem) to a final concentration of 0.05%. Quantifoil 300

mesh copper R1.2/1.3 grids (Quantifoil Micro Tools) were rendered hydrophilic by glow

discharge in a Edwards S150B glow discharger for 1 min 15 sec, setting 6, 30-35 mA, 1.2 kV,

0.2 mBar (0.15 Torr). Subsequently, 3 µL sample was applied to the grids, which were then

mounted in a Vitrobot IV (Thermo Fisher Scientific), blotted with Whatman Filter paper 1 (0-

0.5 s waiting time, 2 s blotting time, -7 blotting force), and plunge frozen in liquid ethane

maintained at a temperature of -170 °C.

CCAN:DNA data aquisition

10,628 micrograph movies were collected on a Titan Krios (ThermoFisherScientific) electron

microscope, equipped with a K3 direct electron detector (Gatan) and energy filter (Gatan)

with a slit width of 20 eV, operating at 300 keV with a flux of 25 e⁻/pixel/s, at a nominal

magnification of 105k, yielding a pixel size of 0.725 Å/px and a total dose of 40e/Å^2.

Automated data acquisition was performed using EPU (ThermoFisherScientific) with

aberration-free image shift enabled, and a defocus range of -0.8 to -2.2 µm in 0.2 µm

increments.

CCAN:CENP-A^{Nuc} complex sample vitrification and cryo-EM data acquistion

To prepare the CCAN:CENP-A^{Nuc} complex for cryo-EM, 3.5 µL of sample at a concentration of

2 mg.mL^{-1} was applied to glow-discharged Quantifoil R2/1 300 mesh copper grids (Quantifoil

Micro Tools). Grids were blotted for 2 s at 4°C , 100% humidity, and a blotting force of -10,

then plunge-frozen in liquid ethane using a Vitrobot Mark IV (Thermo Fisher Scientific). Data

were collected on a Titan Krios microscope (Thermo Fisher Scientific) operated at 300 kV,

equipped with a K3 direct electron detector (Gatan) and an energy filter (Gatan) with an

energy slit of 20 eV. Images were acquired at a nominal magnification of 81,000×,

corresponding to a pixel size of 1.072 Å. Movies were recorded at a dose rate of 15 e⁻/pixel/s

with an exposure time of 3 s over 40 frames. Automated data collection was performed

using EPU (Thermo Fisher Scientific) with aberration-free image shift (AFIS) enabled. The

defocus range was set from -1.0 to -2.0 µm in 0.2 µm increments. A total of 50,781 movie

frames were collected.

CCAN:di-CENP-A^{Nuc} complex sample vitrification and cryo-EM data acquistion

To prepare the CCAN:di-CENP-A^{Nuc} complex for cryo-EM, 3.5 µL of sample at a concentration

of 2 mg.mL^{-1} was applied to glow-discharged Quantifoil R1.2/1.3 300 mesh copper grids

(Quantifoil Micro Tools). Grids were blotted for 2 s at 4°C , 100% humidity, and a blotting

force of -7, then plunge-frozen in liquid ethane using a Vitrobot Mark IV (Thermo Fisher

Scientific). Data were collected on a Titan Krios microscope (Thermo Fisher Scientific)

operated at 300 kV, equipped with a K3 direct electron detector (Gatan) and an energy filter

(Gatan) with an energy slit of 20 eV. Images were acquired at a nominal magnification of

81,000×, corresponding to a pixel size of 0.928 Å. Movies were recorded at a dose rate of

25 e⁻/pixel/s with an exposure time of 1.3 s over 82 frames. Automated data collection was

performed using EPU (Thermo Fisher Scientific) with aberration-free image shift (AFIS)

enabled. The defocus range was set from -1.0 to -2.0 µm in 0.2 µm increments. A total of

28,703 movie frames were collected.

<u>Cryo-EM data processing</u>

CCAN:DNA complex

Micrograph movies were imported into RELION [45] and motion-corrected using RELION's own

implementation of MotionCor2 [48], followed by contrast transfer function estimation using

the RELION wrapper for CTFFIND4 [49]. Particle picking was performed using the crYOLO

general model and particles were extracted in RELION with a box size of 150 pixels. Particles

were subsequently exported to cryoSPARC and subjected to 2D classification. Particles

exhibiting clear secondary structure features were used to generate an *ab initio*

reconstruction. The initial particle set was then subjected to three rounds of heterogeneous

refinement against one true volume (the CCAN:DNA *ab initio* reconstruction) and three

noise decoy volumes arising from *ab initio* reconstruction of particles with no obvious

proteo-typic features. Particles accumulating in the CCAN:DNA class were selected for each

subsequent round of refinement until few or no particles reclassified into the noise decoy

classes. The cleaned particle set was then subjected to homogeneous refinement, followed

by 3D variability analysis. Particles exhibiting the most continuous DNA and CCAN subunit

density were re-extracted without binning in RELION and subjected to particle polishing and

CTF refinement. Finally, polished and CTF-corrected particles were re-imported into

cryoSPARC where a final round of homogeneous refinement was performed, yielding a 3D

reconstruction of CCAN:DNA to an overall resolution of 3.54 Å. The resulting map was

filtered according to local resolution, as estimated in cryoSPARC, and then used for structural

model building (in COOT) [50] and real-space refinement in PHENIX [51].

CCAN:CENP-A^{Nuc} complex

For the CCAN:CENP-A^{Nuc} complex, movie frames were motion-corrected, followed by

estimation of Contrast transfer function (CTF) parameters using the patch-based method in

cryoSPARC [52]. Particle picking was performed with crYOLO [53] using the genaral model, and

particles were extracted with a box size of 400 pixels. Reference-free 2D classfication was

then conducted to clean the particle dataset. Classes exhibting clear secondary structure

features were selected for *ab-initio* reconstruction and heterogeneous refinement in

cryoSPARC. One class with 1,525,520 particles showing nucleosome features was selected

for non-uniform refinement, yielding a reconstruction of the CENP-A^{Nuc}:CENP-C^{N} at 2.4 Å

resolution. Another dominant classs containing 3,815,026 particles was import into RELION

for further 3D classification. Three classes displaying CCAN feature were selected for particle

subtraction using a mask covering 3' CENP-A^{Nuc} region. After an additional round of 2D and

3D classifion on the recentered particle dataset, 325,201 well-aligned nucleosome particles

were reverted to their original, un-subtracted particles for further 3D classification. This

yielded a subset of 39,036 particles corresponding to the CENP-A^{Nuc}:CCAN complex, which

was refined to 4.5 Å resolution using Refine3D followed by Post-process.

CCAN:di-CENP-A^{Nuc} complex

For the CCAN:di-CENP-A^{Nuc} complex, all movie frames were motion-corrected, followed by

estimation of Contrast transfer function (CTF) parameters using the patch-based method in

cryoSPARC. Particle picking was performed with crYOLO using the genaral model, and the

particles were extracted with a box size of 400 pixels. Reference-free 2D classfication was

then conducted to clean the particle dataset, and classes exhibting clear secondary structure

features were selected for *ab-initio* reconstruction and heterogeneous refinement in

cryoSPARC. Three well-populated classes with clear structural features were further selected

for non-uniform refinement, CTF refinement and local refinement, resulting three high-

resolution maps: CENP-A^{Nuc}:ScFv (1:1), CENP-A^{Nuc}:ScFv (1:2) and CCAN:DNA, resolved at 2.7

Å, 2.7 Å and 2.9 Å, respectively.

To dissect the CENP-A^{Nuc} association with CCAN, the CCAN:DNA data was exported to

RELION [45] for 3D classification, which revealed diffuse density at both the 3' and 5' end of

the DNA. To recover CENP-A^{Nuc} at each side, two masks were genarated and to subtract the

density outside either the 3' or 5' CENP-A^{Nuc} regions. After re-centering the particle on the 3'

or 5' CENP-A^{Nuc}, nucleosome density was resolved at both side, as confirmed by 2D

classfication and 3D classfication showing dominant classes with clear nucleosome views.

These dominant classes were reverted to their original un-subtracted particles for further 3D

classification, yielding two subsets: 9,661 particles of CCAN:3' CENP-A^{Nuc} and 5,396 particles

of CCAN:5' CENP-A^{Nuc}. These were refined to 9.5 Å and 15.5 Å respectively. Subsequent

multibody refinement and post-process modestly improved the map quality of both the

CCAN:DNA and CENP-A^{Nuc} modules. Finally, the maps of CCAN with 3' and 5' CENP-A^{Nuc},

were combined to gerarate a composite map of the CCAN di-CENP-A^{Nuc} complex.

<u>Cryo-EM model building and refinement</u>

For the CCAN:DNA structure, the PDB entry 7R5S was used as the initial model. The model

was fitted into the cryo-EM map of CCAN:DNA, and the DNA wrapping around CENP-

TWSX was guided by a predicted CCAN:DNA model generated using AlphaFold3, followed

by manual correction in COOT and real-space refinement in PHENIX. For the CCAN:CENP-A^{Nuc}

structure, the newly refined CCAN:DNA complex model was docked into the corresponding

map, followed by docking of a 3' CENP-A^{Nuc} using PDB entry 7YWX. The models were then

combined in COOT and real-space refined in PHENIX. For the CCAN:di-CENP-A^{Nuc} model,

a composite map combining CCAN:3' CENP-A^{Nuc} and CCAN:5' CENP-A^{Nuc} was used. The

CCAN:DNA model was first fitted into the map, followed by docking of CENP-A^{Nuc} models

(PDB: 6E0P) to the 3' and 5' nucleosomes, respectively. All models were then combined

in COOT, and the linker DNA region was manually built and refined. Additional manual

corrections were also performed in COOT. Figures were generated using ChimeraX [54].

Data availability

Atomic models and cryo-EM maps have been deposited in the Protein Data Bank and the

Electron Microscopy Data Bank, respectively. Accession numbers are listed in Extended Data

Table 1.

Acknowledgements

We are grateful to the LMB and eBIC EM Facilities for help with the EM data collection; J.

Grimmett, T. Darling and I. Clayson for scientific computing; K. Turton for help with insect cell

expression. We thank Andrea Musacchio, Marion Pesenti and Ingrid Vetter for sharing

unpublished experimental observations. For the purpose of open access, the authors have

applied a CC BY public copyright license to any Author Accepted Manuscript version arising.

Funding

UKRI/Medical Research Council MC_UP_1201/6 (D.B.)

Cancer Research UK C576/A25675 (D.B.)

References

1. Ariyoshi, M. & Fukagawa, T. An updated view of the kinetochore architecture. *Trends*
Genet **39**, 941–953 (2023).
2. McAinsh, A.D. & Marston, A.L. The Four Causes: The Functional Architecture of
Centromeres and Kinetochores. *Annu Rev Genet* **56**, 279–314 (2022).
3. Musacchio, A. & Desai, A. A Molecular View of Kinetochore Assembly and Function.
Biology (Basel) **6**(2017).
4. McAinsh, A.D. & Kops, G. Principles and dynamics of spindle assembly checkpoint
signalling. *Nat Rev Mol Cell Biol* **24**, 543–559 (2023).
5. Fischer, E.S. Kinetochore-catalyzed MCC formation: A structural perspective. *IUBMB*
Life **75**, 289–310 (2023).
6. Altemose, N. et al. Complete genomic and epigenetic maps of human centromeres.
Science **376**, eabl4178 (2022).
7. Bodor, D.L. et al. The quantitative architecture of centromeric chromatin. *Elife* **3**,
e02137 (2014).
8. Logsdon, G.A. et al. The structure, function and evolution of a complete human
chromosome 8. *Nature* **593**, 101–107 (2021).

9.   Gershman, A. et al. Epigenetic patterns in a complete human genome. *Science* **376**,
eabj5089 (2022).

10.   Blower, M.D., Sullivan, B.A. & Karpen, G.H. Conserved organization of centromeric
chromatin in flies and humans. *Dev Cell* **2**, 319–30 (2002).

11.   Luger, K., Mader, A.W., Richmond, R.K., Sargent, D.F. & Richmond, T.J. Crystal
structure of the nucleosome core particle at 2.8 A resolution. *Nature* **389**, 251–60
(1997).

12.   Yatskevich, S. et al. Structure of the human inner kinetochore bound to a centromeric
CENP-A nucleosome. *Science* **376**, 844–852 (2022).

13.   Hasson, D. et al. The octamer is the major form of CENP-A nucleosomes at human
centromeres. *Nat Struct Mol Biol* **20**, 687–95 (2013).

14.   Corda, L. & Giunta, S. Chromosome-specific centromeric patterns define the centeny
map of the human genome. *Science* **389**, eads3484 (2025).

15.   Fujita, R. et al. Stable complex formation of CENP-B with the CENP-A nucleosome.
Nucleic Acids Res **43**, 4909–22 (2015).

16.   Nagpal, H. et al. CENP-A and CENP-B collaborate to create an open centromeric
chromatin state. *Nat Commun* **14**, 8227 (2023).

17.   Chardon, F. et al. CENP-B-mediated DNA loops regulate activity and stability of
human centromeres. *Mol Cell* **82**, 1751–1767 e8 (2022).

18.   Magidson, V. et al. Adaptive changes in the kinetochore architecture facilitate proper
spindle assembly. *Nat Cell Biol* **17**, 1134–44 (2015).

19.   Yatskevich, S., Barford, D. & Muir, K.W. Conserved and divergent mechanisms of inner
kinetochore assembly onto centromeric chromatin. *Curr Opin Struct Biol* **81**, 102638
(2023).

20.   Hori, T. et al. CCAN makes multiple contacts with centromeric DNA to provide distinct
pathways to the outer kinetochore. *Cell* **135**, 1039–52 (2008).

21.   Pesenti, M.E. et al. Structure of the human inner kinetochore CCAN complex and its
significance for human centromere organization. *Mol Cell* **82**, 2113–2131 e8 (2022).

22.   Dendooven, T. et al. Cryo-EM structure of the complete inner kinetochore of the
budding yeast point centromere. *Sci Adv* **9**, eadg7480 (2023).

23.   Tian, T. et al. Structural insights into human CCAN complex assembled onto DNA. *Cell*
Discov **8**, 90 (2022).

24.   Yan, K. et al. Structure of the inner kinetochore CCAN complex assembled onto a
centromeric nucleosome. *Nature* **574**, 278–282 (2019).

25.   Hinshaw, S.M. & Harrison, S.C. The structure of the Ctf19c/CCAN from budding yeast.
Elife **8**(2019).

26.   Yu, C.W.H. et al. Architecture and Function of Holocentric CENP-A-Independent
Kinetochores. *BioRxiv* (2025).

27.   Kato, H. et al. A conserved mechanism for centromeric nucleosome recognition by
centromere protein CENP-C. *Science* **340**, 1110–3 (2013).

28.   Klare, K. et al. CENP-C is a blueprint for constitutive centromere-associated network
assembly within human kinetochores. *J Cell Biol* **210**, 11–22 (2015).

29.   McKinley, K.L. et al. The CENP-L-N Complex Forms a Critical Node in an Integrated
Meshwork of Interactions at the Centromere-Kinetochore Interface. *Mol Cell* **60**,
886–98 (2015).

30.   Pentakota, S. et al. Decoding the centromeric nucleosome through CENP-N. *Elife*
6(2017).

31. Walstein, K. et al. Assembly principles and stoichiometry of a complete human
kinetochore module. *Sci Adv* **7**(2021).

32. Nishino, T. et al. CENP-T-W-S-X forms a unique centromeric chromatin structure with
a histone-like fold. *Cell* **148**, 487–501 (2012).

33. Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M. & Barton, G.J. Jalview
Version 2--a multiple sequence alignment editor and analysis workbench.
Bioinformatics **25**, 1189–91 (2009).

34. Zhao, Q. et al. The MHF complex senses branched DNA by binding a pair of crossover
DNA duplexes. *Nat Commun* **5**, 2987 (2014).

35. Dechassa, M.L. et al. Structure and Scm3-mediated assembly of budding yeast
centromeric nucleosomes. *Nat Commun* **2**, 313 (2011).

36. Kingston, I.J., Yung, J.S. & Singleton, M.R. Biophysical characterization of the
centromere-specific nucleosome from budding yeast. *J Biol Chem* **286**, 4021–6
(2011).

37. Migl, D. et al. Cryoelectron Microscopy Structure of a Yeast Centromeric Nucleosome
at 2.7 A Resolution. *Structure* **28**, 363–370 e3 (2020).

38. Tachiwana, H. et al. Crystal structure of the human centromeric nucleosome
containing CENP-A. *Nature* **476**, 232–5 (2011).

39. Ariyoshi, M. et al. Cryo-EM structure of the CENP-A nucleosome in complex with
phosphorylated CENP-C. *EMBO J* **40**, e105671 (2021).

40. Zhou, B.R. et al. Atomic resolution cryo-EM structure of a native-like CENP-A
nucleosome aided by an antibody fragment. *Nat Commun* **10**, 2301 (2019).

41. Roulland, Y. et al. The Flexible Ends of CENP-A Nucleosome Are Required for Mitotic
Fidelity. *Mol Cell* **63**, 674–685 (2016).

42. Boopathi, R. et al. Phase-plate cryo-EM structure of the Widom 601 CENP-A
nucleosome core particle reveals differential flexibility of the DNA ends. *Nucleic Acids*
Res **48**, 5735–5748 (2020).

43. Falk, S.J. et al. Chromosomes. CENP-C reshapes and stabilizes CENP-A nucleosomes at
the centromere. *Science* **348**, 699–703 (2015).

44. Ali-Ahmad, A., Bilokapic, S., Schafer, I.B., Halic, M. & Sekulic, N. CENP-C unwraps the
human CENP-A nucleosome through the H2A C-terminal tail. *EMBO Rep* **20**, e48913
(2019).

45. Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S.H.W. New tools for
automated cryo-EM single-particle analysis in RELION-4.0. *Biochem J* **478**, 4169–4185
(2021).

46. Dubocanin, D. et al. Conservation of dichromatin organization along regional
centromeres. *Cell Genom* **5**, 100819 (2025).

47. Kixmoeller, K., Tarasovetc, E.V., Mer, E., Chang, Y.W. & Black, B.E. Centromeric
chromatin clearings demarcate the site of kinetochore formation. *Cell* **188**, 1280–
1296 e19 (2025).

48. Zheng, S.Q. et al. MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331–332 (2017).

49. Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. *J Struct Biol* **192**, 216–21 (2015).

50. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486–501 (2010).

51.   Afonine, P.V. et al. Towards automated crystallographic structure refinement with
856   phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352–67 (2012).

857   52.   Punjani, A., Zhang, H. & Fleet, D.J. Non-uniform refinement: adaptive regularization
858   improves single-particle cryo-EM reconstruction. *Nat Methods* **17**, 1214–1221
859   (2020).

860   53.   Wagner, T. et al. SPHIRE-crYOLO is a fast and accurate fully automated particle picker
861   for cryo-EM. *Commun Biol* **2**, 218 (2019).

862   54.   Goddard, T.D. et al. UCSF ChimeraX: Meeting modern challenges in visualization and
863   analysis. *Protein Sci* **27**, 14–25 (2018).

Extended Data Table 1. Cryo-EM data collection, refinement, and validation statistics

Complex	CCAN:DNA PDB XXXX EMD-XXXXX	CCAN:mono-CENP-A PDB XXXX EMD-XXXXX	CCAN:di-CENP-A PDB XXXX EMD-XXXXX	
Data collection and Processing				
Microscope	Titan Krios	Titan Krios	Titan Krios	Titan Krios
Voltage (keV)	300	300	300	300
Camera	K3	K3	K3	K3
Magnification	105,000	81,000	81,000	81,000
Pixel size at detector (Å/pixel)	0.725	1.072	0.928	0.928
Total electron exposure (e^-/Å^2)	40	40	40	40
Exposure rate (e-/pixel/sec)	25	15	18	18
Number of frames	40	40	82	82
Defocus range (µm)	0.8-2.2	1-2	1.0-2.0	1.0-2.0
Automation software	EPU	EPU	EPU	EPU
Energy filter slit width (eV)	20	20	20	20
Micrographs collected (no.)	10,628	50,781	28,703	28,703
Total extracted particles (no.)	1,963,587	21,223,394	8,973,622	8,973,622
For each reconstruction:			CCAN/3'-CENP-A	CCAN/5'-CENP-A
Final particles (no.)	80,236	39,306	9,661	5,396
Point-group	C1	C1	C1	C1
Resolution (global, Å)	3.54	4.5	9.5	15.5
Resolution range (local, Å)	3.54-20	4.5-18	9.5-20	15.5-20
Map sharpening B factor (Å^2)	-73	-20	-50	-50
Map sharpening methods	cryoSPARC	RELION5	RELION5	RELION5
Model composition				
Protein (residues)	3,165	3,962	5,600	
RNA/DNA (nucleotides)	162	398	636	
Refinement				
Refinement package	PHENIX 1.20.1	PHENIX 1.20.1	PHENIX 1.21	
real or reciprocal space	real	real	real	
Initial model used (Alphafold2)	7R5S/AlphaFold3	7YWX	7R5S/6E0P	
Model-Map scores				
CC Volume/Mask	0.66/0.67	0.48/0.46	0.26/0.36	
B factors (Å^2)				
Protein residues	2.50/63.61/30.63	111.54	80.43	
RNA/DNA	46.81/99.32/79.68	48.18	50	
R.m.s. deviations from ideal values				
Bond lengths (Å)	0.006	0.012	0.012	
Bond angles (°)	1.003	1.160	1.203	
Validation				
MolProbity score	1.56	1.94	1.59	
CaBLAM outliers	1.22	1.38	0.04	
Clashscore	7.26	20.98	6.74	
Poor rotamers (%)	0.14	0	0.29	
C-beta deviations	0	0	0	
EMRinger score	1.26	0.25	*n.d.	
Ramachandran plot				
Favored (%)	97.13	97.37	96.63	
Allowed (%)	2.81	2.48	3.25	
Outliers (%)	0.06	0.15	0.13	

*Not determined (n.d.)

Extended Data Fig. 1. Cryo-EM data and processing workflow for the CCAN:DNA complex. **a**, 2D class averages and FSC curves for CCAN:DNA. **b**, Cryo-EM processing workflow. **c**, 3D reconstruction. **d**, FSC curves.

Extended Data Fig. 2. Cryo-EM images and processing workflow for CCAN:CENP-A^Nuc complex. **a**, Cryo-EM processing workflow. **b**, Cryo-EM images. **c**, 2D class average gallery. **d**, Cryo-EM reconstruction colour-coded according to local resolution. **e**, FSC curves.

Extended Data Fig. 3. Reconstitution of CCAN:di-CENP-A^Nuc complex. a, Schematic of CCAN:di-CENP-A^Nuc complex showing α-satellite repeat dimer used in this study. The 3' (downstream) repeat is 171 bp whereas the 5' upstream repeat is 147 bp, i.e. without linker DNA. This design ensures only one CCAN-binding linker DNA per α-satellite repeat dimer. **b** and **c**, Size exclusion chromatogram (b) and associated SDS PAGE gel (c) of purified CCAN:di-CENP-A^Nuc complex. **d**, iSCAT data for CCAN:di-CENP-A^Nuc complex showing a mixture of di-CENP-A^Nuc alone (445 kDa) and the CCAN:di-CENP-A^Nuc complex (1,054 kDa).

Extended Data Fig. 4. CCAN association with linker DNA displaces CENP-B bound to linker DNA B-box. a, Schematic of the CCAN:di-CENP-A^{Nuc} complex showing a-satellite repeat dimer used in this study with B-box in blue. **b**, Di-CENP-A^{Nuc}:CENP-B complex (with wild type B-box). **c**, CENP-B does not bind to di-CENP-A^{Nuc} with a mutated B-box, demonstrating linker DNA binding. **d**, and **e**, Size exclusion chromatogram (d) and associated SDS PAGE gel (e) showing that CCAN binding to di-CENP-A^{Nuc} displaces CENP-B.

Extended Data Fig. 5. Reconstitution of CCAN$^{DCENP-C}$:di-CENP-A^{Nuc} complex for cryo-EM. a and **b**, Size exclusion chromatogram (A) and associated SDS PAGE gel (B) of purified CCAN$^{\Delta CENP-C}$:di-CENP-A^{Nuc}:ScFv complex. **c**, iSCAT data for CCAN$^{DCENP-C}$:di-CENP-A^{Nuc}:ScFv complex.

Extended Data Fig. 6. Cryo-EM data for the CCAN:di-CENP-A^{Nuc} complex.
a, Representative cryo-electron micrograph. **b**, 2D class averages of CENP-A^{Nuc}.
c, 2D class averages of CCAN with 3' CENP-A^{Nuc}. **d**, 2D class averages of CCAN with 5' CENP-A^{Nuc}.

Extended Data Fig. 7. Cryo-EM workflow for CCAN$^{DCENP-C}$:di-CENP-A^{Nuc}:ScFv complex reconstruction. a, Cryo-EM processing workflow. **b**, FSC curves.