Title: High-resolution mapping of a rapidly evolving complex trait reveals genotype-phenotype stability

and an unpredictable genetic architecture of adaptation

Smiley-Rhodes J.A*[1], Bitter M.C.*[1], Berardi S.[2], Oken H.[2], Beltz J.[2], Petrov D.A.[#1,3], and Schmidt P.[#2]

* = equal contribution

# = equal contribution

1 – Department of Biology, Stanford University, Stanford CA 94305

2 – Department of Biology, University of Pennsylvania, Philadelphia, PA 19104

3 – Chan Zuckerberg Biohub, San Francisco, CA 94158, USA

Correspondance: dpetrov@stanford.edu, schmidtp@sas.upenn.edu

Abstract

The extent to which adaptation can be predicted, particularly for traits with complex genetic bases, is

unknown. Here, we leveraged a model complex trait, model species, and high-powered longitudinal

sampling design to test the efficacy of genomic prediction of complex trait variation and evolution in

ecologically relevant settings. We monitored genome-wide allele frequencies and pigmentation variation

in genetically diverse populations of *Drosophila melanogaster* across seven generations of evolution in

both field mesocosms exposed to natural environmental fluctuations, as well as mesocosms housed in a

controlled, lab-based setting. At two time points throughout trait evolution, we conducted a high-

powered, tail-based mapping of pigmentation. We produced a well-resolved genotype-phenotype map

that reaffirms canonical pigmentation genes and unveils novel loci underpinning a substantial portion of

trait variation. While we were able to use this map to correctly infer the direction of pigmentation

evolution in both the field and lab mesocosms, the particular loci responding to selection, and thus

architecture of adaptation itself, were largely unpredictable. We suggest this unpredictability to be a result

of pleiotropic constraint, which was more pronounced in the field, relative to the lab-based, environment.

Finally, we quantified a striking stability of the genotype-phenotype map across genetically diverged

populations, demonstrating that shifting epistatic landscapes associated with the evolutionary process

itself does not alter trait architecture to an extent that precludes phenotypic prediction, provided the

mapping is sufficiently powered. In concert our results highlight both the promise and limitations of

genomic prediction, and exemplify the challenges of applying lab-based studies of complex traits to their

evolutionary dynamics in the wild.

Introduction

Whether or not adaptation can be predictable is an outstanding question in evolutionary biology (Blount et al., 2018; Wortel et al., 2022). This fundamental question permeates numerous issues of societal concern: from the preservation of species in the face of global change to the management of disease outbreaks. Two distinct ways in which adaptation can be predicted are: i) correctly inferring whether or not a focal trait has evolved, given patterns of genomic variation alone; or ii) correctly inferring the presence and direction of selection on genomic loci, given observations of phenotypic change alone. In order to employ either method of prediction, researchers must first quantify the molecular genetic basis of phenotypic variation, and then understand how such phenotypic variation is translated into fitness-relevant differences among genotypes.

Instances in which adaptation has proven to be predictable are almost exclusively in the context of traits exhibiting simple genetic bases, whereby trait variation is predominantly explained by one or a few large effect loci (Colosimo et al., 2005; Enattah et al., 2008; Hoekstra et al., 2006; Hof et al., 2016; Kunte et al., 2014; Nadeau et al., 2016; Reed et al., 2011). While such case studies provide fundamental insight into the evolutionary process, the vast majority of fitness-relevant traits instead exhibit complex genetic bases, whereby phenotypic differences are due to allelic variation at tens to thousands of loci spread throughout the genome (Sella and Barton, 2019; Visscher et al., 2017). Quantifying, and ultimately predicting, how these traits evolve remains a challenge.

Two facets of quantitative trait variation that may preclude evolutionary predictability are pleiotropy and epistasis. Pleiotropy occurs when a single genomic locus impacts phenotypic variation in multiple traits, a phenomenon expected to be pervasive among the large-effect quantitative trait loci that impact fitness in natural populations (Mackay and Anholt, 2024; Walsh and Blows, 2009). As populations adapt, correlated selection across fitness-relevant traits can then impact the evolution of pleiotropic loci, constraining their effect on the evolution of the trait in question (Svensson et al., 2021). In such a scenario, patterns of genetic variation at a mapped locus may behave as effectively neutral during the evolution of the trait itself, or even indicate trait shifts in the opposite direction of how it in fact evolved through time (Albecker et al., 2025; Conover and Schultz, 1995; Sella and Barton, 2019; Walsh and Blows, 2009).

Epistasis is the interaction of allelic effects at multiple loci and, in the context of predicting trait adaptation, may shift the genotype-phenotype map throughout the evolutionary process (Duncan et al., 2019; Huang et al., 2012; Phillips, 2008; Privé et al., 2022). Specifically, epistasis can cause the marginal effect of trait-associated loci to change as a function of genetic background, which varies across populations with distinct demographic histories and patterns of selection (Duncan et al., 2019; Durvasula and Lohmueller, 2019; Guo et al., 2014; Huang et al., 2012; Kosheleva and Desai, 2018; Lohmueller,

2014; Martin et al., 2017; Novembre and Barton, 2018; Patel et al., 2022; Privé et al., 2022; Yair and

Coop, 2021). Similarly, context-dependent dominance may alter the phenotypic effect of alleles as a

function of shifting environments (Grieshop et al., 2021). It is therefore unclear whether loci mapped

within a particular population and at a particular point in time will be those that underpin phenotypic

variation when the trait mean shifts, as is often assumed by studies that use contemporary mapping data to

infer past trait evolution using genomic data (Abdellaoui et al., 2023; Akbari et al., 2024; Cox et al.,

2019; Halldorsson et al., 2022; Mathieson et al., 2015).

In concert, epistasis and pleiotropy lead to a genetic architecture of adaptation that may be much

less predictable based on the genotype-phenotype mapping data alone. As these processes and their

effects manifest through time, inferring them using static sequencing data can be challenging. Thus,

longitudinal studies, which directly examine selection on individual traits and loci throughout the genome

over the progression of evolution itself represent a powerful method to study pleiotropy, epistasis, and the

predictability of complex trait evolution (Chiu et al., 2016; Kerner et al., 2009; Lenski, 2017; Stroud and

Ratcliff, 2025; Wu et al., 2014). Such studies can probe the predictability of adaptation in a systematic

manner, answering fundamental questions, such as: are the loci underpinning phenotypic variation in a

complex trait the same that respond to selection and underpin trait evolution in nature? How stable is the

trait genotype-to-phenotype map throughout evolution and across different environments associated with

distinct selection pressures?

Our study comprised a longitudinal analysis of abdominal pigmentation evolution in *Drosophila*

melanogaster within an ecologically realistic context. Pigmentation is one of the most conspicuous

complex traits associated with phenotypic diversity across the tree of life, demonstrating adaptive

evolution across a range of taxa (Kronforst et al., 2012). In *D. melanogaster*, abdominal pigmentation

exhibits latitudinal clines across widespread geographic populations, and rapidly and cyclically evolves in

response to seasonally varying selection pressures (Berardi et al., 2025). Importantly, abdominal

pigmentation has long served as a model complex trait, whereby the biosynthetic pathway of the two

main molecules, darkly pigmented melanin and light-colored sclerotin, have been well-characterized

(Massey and Wittkopp, 2016; Wright, 1987). More recent genome wide associations studies have

pinpointed three core genes in the melanin/sclerotin biosynthetic pathway that explain the majority of

intra- and inter-population pigmentation variation: *bric-a-brac* (*bab*), *tan*, and *ebony* (Bastide et al., 2016,

2013; Dembeck et al., 2015). Interestingly, clines in pigmentation across space and time do not appear to

be explained by expected signatures of selection at these pigmentation genes of large effect (Berardi et al.,

2025), a pattern suggestive of underlying pleiotropy and/or epistasis but challenging to attribute to these

factors directly in the face of confounding population demography and linkage.

Here, using a single genetically diverse founding population we quantified pigmentation
evolution across seven generations in two environments with distinct selective pressures: a simple, lab-
based environment and a dynamic, semi-natural field mesocosm. We quantified evidence of selection on
pigmentation, as well as genome-wide single nucleotide polymorphisms (SNPs) as our focal trait itself
evolved rapidly over seven generations in each environment. At two points throughout this process, we
conducted a tail-based mapping of replicate populations to independently identify trait-associated loci in
each of four total mapping populations (Fig. 1). Our genotype-phenotype mapping both validated the role
of previously identified candidate pigmentation genes, and also identified novel loci underpinning
pigmentation variation in the species. We found that this map could then be used to correctly infer
pigmentation evolution through time in each environment given patterns of allele frequency data alone.
Interestingly, however, the loci displaying signatures of selection as pigmentation evolved (i.e., the
genetic architecture of adaptation) did not always behave in a manner as predicted by mapping, a dynamic
that was exacerbated in the complex, field setting. This pattern argues for the strong effect of pleiotropy
affecting the genomic predictability, and underlying genetic architecture, of phenotypic evolution in
nature. Finally, we observed genotype-phenotype mapping to be remarkably stable and transferable across
mapping populations, indicating that marginal effects of mapped loci remain detectable even amidst
shifting epistatic landscapes associated with the evolutionary process itself.
119
Results
Study scheme and power
Our study population was derived via outcrossing a panel of 76 inbred lines originally collected
in local Pennsylvania orchards (Behrman et al. 2018). Several key advantages arise from this
methodological decision. By allowing evolution to proceed from a single founding population we could
evaluate the effect of epistasis on genotype-phenotype map portability directly and in a manner that is not
confounded by population-specific demographic histories and patterns of linkage (Lohmueller, 2014;
Novembre and Barton, 2018). Furthermore, the pigmentation and fitness-relevant variation subject to
selection in our study was evaluated in the same local region from which it originated and is maintained
naturally, and in which we have documented repeatable seasonal evolution of lighter pigmentation from
late spring through summer (Berardi et al., 2025). Finally, the haplotype-informed allele frequency
estimates we are able to generate from the reconstituted outbred populations are of the accuracy produced
using standard pool-seq methods and sequencing depths of roughly 1000x (Tilk et al., 2019). Such
haplotype-imputed methods of mapping have been recently shown to be significantly greater powered for
complex trait mapping than more standard SNP-by-SNP approaches (Long et al. 2025).

The outbred population was allowed to recombine for four generations, before being used to seed

each of 10 large indoor cage replicates maintained in controlled, lab-based conditions (Bitter et al.,

2024b), as well as 10 replicate outdoor field cages comprising a semi-natural mesocosm exposed to

natural environmental fluctuations in Philadelphia, PA (Fig. 1) (Bitter et al., 2024a, 2024b; Rudman et al.,

2022). Populations evolving within the indoor environment were subject to increasing population

densities, but in the absence of the suite of fluctuating abiotic conditions imposed upon the populations

evolved in the outdoor mesocosm. We allowed the replicate populations to expand and evolve for eight

weeks (seven generations) in each environment, during which time we monitored patterns of

pigmentation variation and genome-wide allele frequencies at 1.9 M single nucleotide polymorphisms

(SNPs). At weeks 2 and 8 of the experiment (generations 2 and 7), we conducted tail-based mapping

whereby we selectively isolated and sequenced pools of 75 female flies per cage from the 15% lightest,

15% darkest, and representative midpoint of the trait value distribution (hereafter referred to as 'light',

'dark', and 'midpoint' trait value groups, see Methods) from each of the ten replicate populations of the

lab and field mesocosms. Whole genome sequencing reads derived from each sample were then subject to

a bioinformatic pipeline that leverages knowledge of the haplotype structure from a founding set of

individuals (here, our inbred reference panel) to compute genome-wide haplotype and allele frequency

data in evolved, pooled samples (Kessner et al., 2013; Tilk et al., 2019). This resulted in, for each sample,

estimates of haplotype frequencies within locally unrecombined windows (quantified using the

recombination rate and the number of generations since cage founding) and allele frequencies at each of

the 1.9 M SNPs we previously identified in our inbred reference panel.

Figure 1 | Experimental design. Ten cages each in an indoor laboratory and in an outdoor mesocosm were seeded with an outbred population of *D. melanogaster*. The replicate populations were allowed to evolve from late June to late October, during which time significant pigmentation evolution was observed (see text). At weeks 2 and 8 of the experiment (generations 2 and 7) we conducted tail-based mapping of pigmentation. Specifically, following two generations of common garden rearing, flies were segregated into color fractions based on the population's distribution of abdominal pigmentation scores (fractions comprised 75 female individuals from the darkest 15%, the lightest 15%, and a representative sample from the trait midpoint) across each of ten replicates per time point and environment. Each color fraction from each replicate and time point was then used in whole-genome sequencing and estimation of high-accuracy, haplotype-informed allele frequencies at 1.9 M SNPs.

Phenotypic and genomic variation across environments and throughout evolution

We first assessed the degree of pigmentation variation in our study population and whether or not it was affected by natural selection (whether direct or indirect) in either the lab or field mesocosm. Specifically, we quantified pigmentation independently for each inbred reference line, across individuals

in the derived outbred population (hereafter, 'baseline'), and among replicate populations at weeks two

and eight of evolution in each environment. As expected, our baseline population exhibited variation in

pigmentation that was representative of the variation observed across the 76 inbred lines from which it

was derived (Fig. 2A). Pigmentation after two generations of evolution resulted in significantly lighter

pigmentation across replicates in both the lab and field environments, relative to the baseline (pairwise t-

tests, week 2 outdoor $p \ll 0.001$, week 2 indoor $p \ll 0.001$). Also as expected, the replicate populations

then evolved to become significantly lighter during the six additional weeks of evolution in both the

outdoor and indoor environments (pairwise t-tests; week 2 vs. 8 outdoor $p \ll 0.001$, week 2 vs. 8 indoor

$= 0.0245$). Notably, a significant treatment by time interaction on variation in pigmentation throughout

the experiment was observed (two-way ANOVA; $F = 17.03$, $p \ll 0.001$), driven by the accelerated rate

and magnitude of pigmentation evolution towards lighter pigmentation observed in the outdoor replicates

(Fig. 2A).

Next, we quantified whether there existed evidence of environment-specific patterns of selection

that would, in effect, induce unique epistatic landscapes across lab and field-based mapping populations.

Specifically, we ran principal component analysis (PCA; *prcomp* function, R), on the normalized allele

frequency data of all 1.9 M SNPs, and then projected samples onto the first two principal components

(Fig. 2C-D). Strikingly, the first principal component (6.4% variance explained) segregated samples in

accordance with trait group, and along the second principal component (4.8% variance explained) in

accordance with environment of selection (Fig. S1). When segregated by chromosome, we observed the

same correlation between PC1 and color fraction on all chromosomes except X, and all chromosomes

except 2R demonstrated the same relationship between PC2 and treatment (Fig. S2). These patterns were

not disrupted when the major cosmopolitan inversions were excluded (Fig. S3). This analysis thus

demonstrates the pronounced impact of both phenotypic group and environment of selection on genome

wide patterns of variation, while also suggesting that a shared set of loci underpinned variation in

pigmentation across mapping populations.

Figure 2 | Phenotypic and genomic variation throughout study period. (A) Distribution of the total pigmentation scores (whereby higher scores indicate increased pigmentation) across samples. "Founder" refers to our 76 inbred lines that we outbred for four generations to generate our baseline (week 0) population. Grey dots each represent the average pigmentation score of N = 3 females from each inbred line. The baseline pigmentation distribution represents the scores of N = 90 females from the initial outbred population which was used to seed our indoor and outdoor cages. From the samples collected during the experiment at weeks 2 and 8 in both environments, we scored 30 females per replicate cage (N = 300 in total for each timepoint/treatment combination) and found a significant shift towards lighter pigmentation in both outdoor and indoor populations (paired t-test; p-values << 0.001). (B-C) Projection of samples used in trait mapping onto the first two principal components generated via PCA of allele frequency data from all 1.9 M SNPs. In (B) samples are colored by the color fraction from which they originate (dark, midpoint, or light) and the shape of the points corresponds to the timepoint when the sample was collected. PC1 value and color fraction identity are significantly associated (linear regression p-value << 0.001). In (C) samples are colored by treatment. PC2 value and treatment are significantly associated (ANOVA p-value << 0.001).

Ancestry-informed mapping provides resolved architecture of abdominal pigmentation

Given the substantial variation in pigmentation in our study population (Fig. 2B), we next aimed

to map the underlying genetic architecture of this trait. We initially focused mapping on samples collected

at week 8 in the outdoor mesocosm, as these replicates had undergone evolution in a semi-natural

environment and also experienced more generations of recombination since initial outbreeding than the

week 2 samples, and were thus predicted to generate better resolved genetic maps. Our general approach

to mapping was to identify SNPs harboring alleles with systematic frequency changes across the light,

midpoint, and dark trait value groups. However, given the finite number of haplotypes from the founding

inbred lines used to derive our outbred population, a significant association between a SNP and trait value

group could indicate either its linkage to a locally causal allele and/or its broader inbred-line ancestry.

Accordingly, to infer the magnitude of the inbred-line ancestry genome-wide, we quantified the average

contribution of haplotypes coming from each inbred line to each trait value group and tested for the

association of this quantity with the pigmentation score of that inbred line. The results of this are depicted

in Figure 3A, where we observe that the light trait value group contains a disproportionate amount of

ancestry from the lightest inbred lines and, likewise, the dark trait value group contains a similarly

disproportionate amount of ancestry from the darkest inbred lines ($R^2 = 0.032$; $p < 0.01$). To further

quantify this ancestry effect on pigmentation, we a ran a simple regression (GLM; logistic link function

and quasibinomial error variance) of allele frequencies on trait value group. We took the top 1% of the

SNPs (based on nominal p-value from the GLM) and calculated their expected frequency difference

between light and dark color fractions based on estimates of average contribution of each inbred line to

that sample, as well as the association between that SNP and its presence in the inbred line (Fig 3B).

Indeed, we see that the GLM-identified light-associated SNPs tend to be associated with the lighter

pigmented lines, making it hard to disentangle the SNP's local association with a causal locus from its

association with global line ancestry.

To account for the strong ancestry effects on SNP-pigmentation association (Fig. 3A-B), we ran

an ancestry-informed GLM. Specifically, allele frequencies were regressed both on pigmentation trait

value and the SNP's association with each inbred line and its genome-wide relative contribution to each

trait value group (light, midpoint, or dark). As expected, in Fig. 3B we see that the distribution of

expected light-dark frequency differences for SNPs identified by this model is now centered around zero,

indicating that the global bias produced by our trait-only GLM is eliminated. We further validated that

SNPs identified via this approach were not exclusive to the lightest nor darkest founding inbred lines

(colored points, Fig. 2A), but rather present at appreciable frequency across all founding haplotypes (Fig.

S4). We then defined a robust set of pigmentation-associated SNPs using permutations of the trait value

data and re-running our ancestry-informed GLM (N = 100 permutations). A SNP was considered

significant if its nominal p-value from the empirical GLM fell in the top 0.05% of permuted p-values

(corresponding to an empirical FDR threshold < 0.0005; Fig. S5). Even at this conservative threshold, we

see that trait-associated SNPs are pervasive throughout the genome and distributed across all

chromosomal arms (Fig 3C). Hereafter, we refer to pigmentation-associated SNPs identified via this

approach as either light or dark-associated given their relative frequency in the light and dark color

fractions: the light-associated allele for each mapped SNP is that occurring at higher relative frequency in

the light color fraction.

We next used our set of pigmentation-associated SNPs throughout the genome to identify the

number and location of unlinked loci underpinning variation in the trait. To do so, we first identified the

most significant SNP on each chromosomal arm, as well as its surrounding, locally un-recombined region

(defined by the recombination rate and number of generations since cage founding, see Methods). We

then eliminated all trait-associated SNPs on the chromosome whose relative presence in the light and dark

fractions can be explained by haplotype frequencies in the locally un-recombined focal region, in effect

eliminating all signal due to linkage at the focal SNP location. From the set of remaining SNPs, we

identified the SNP with the most significant nominal p-value and repeated the procedure. This was

iterated until we exhausted all trait-associated SNPs on the chromosome and resulted in a total of 76

genomic loci, the midpoint of which was the most significant, locus-defining SNP. Merging loci with

overlapping start and endpoints, we obtained a final list of 42 loci, ranging in size from 700 Kb to 3 Mb,

and distributed across all chromosomal arms (Supplementary Data File 1). This indicates that the genetic

architecture of our focal, ecologically relevant complex trait, is not underpinned solely by a single or few

large-effect loci, nor exclusively by a very large number of weak-effect loci (e.g., 1000's). Rather, we

infer the presence of a moderate number of large to moderate-effect loci underpinning trait variation,

while acknowledging there may be further loci of small effect that our approach lacks sufficient power to

detect. We once again validated that pigmentation-associated SNPs within the largest-effect loci were not

exclusive to those lightest/darkest haplotypes enriched in the trait tails (Fig. 3A), but rather shared across

all founding inbred lines (Fig. S6).

We quantified the extent to which our identified loci validated previously described, and

potentially identified new, candidate genes associated with pigmentation variation in the species.

Assuredly, the top loci on chromosomal arms 3L, 3R, and X contained the previously described core

genes impacting pigmentation variation within the species: *bab*, *ebony*, and *tan* (Bickel et al., 2011; Jeong

et al., 2008; Massey and Wittkopp, 2016; Pool and Aquadro, 2007). A comparison of the association

signal between loci identified here and two recent, well-powered association studies of the trait (Bastide

et al., 2013; Dembeck et al., 2015) indicated significantly more overlap than expected by chance

(permutation p < 0.001, see Methods). However, this signal is eliminated when we no longer consider

SNPs within our mapped loci containing *bab*, *ebony*, and *tan* (p = 0.994) suggesting little overlap outside

of these three core genes. Thus, the extensive enrichment of pigmentation-associated variation outside

these candidate loci we observe in our data suggests the presence of novel candidate pigmentation loci

identified in our study, which we discuss in the supplement (Supplementary Discussion; Table S1-2).

Pleiotropy and predictions of complex trait evolution

A potential of application of a well-resolved genetic map is to predict whether or not a trait has

evolved using genomic data alone. However, this effort could be obscured by underlying pleiotropy at

quantitative trait loci, which may constrain their impact on trait evolution and leave the architecture of

trait adaptation largely unpredictable. Such pleiotropic interactions may be particularly pronounced in

complex environments where selection across multiple fitness-relevant traits may induce correlated

selection on pleiotropic loci. Our longitudinal study design afforded the unique opportunity to test this

directly by combining our well-resolved genetic mapping data (Fig. 2C-D) with our time-series allele

frequency data collected throughout phenotypic evolution of the trait (Fig. 1; 2A). Accordingly, for each

of the 42 independent, unlinked loci identified above, we quantified the frequency shifts of the light-

associated alleles as the population itself evolved to become significantly lighter in the outdoor mesocosm

(Fig. 2A). We then inferred whether observed allele frequency movement within each locus indicated

lighter pigmentation evolution by comparing its behavior to a distribution derived from a set of matched

control SNPs (see Methods). We found that out of the 42 loci, 18 exhibited an overall increase in light-

associated allele frequency and thus a behavior concordant with expectations based on the evolution of

lighter pigmentation (i.e., overall increase in frequency of light-associated alleles). For 17 pigmentation-

associated loci, however, we did not detect any systematic shift indicative of trait evolution, while 7 loci

showed behavior in the direction of darker pigmentation (Fig. 3D). Thus, in sum, while our genome-wide

inference correctly predicted evolution of lighter pigmentation in the outdoor mesocosm (p << 0.001; Fig.

4A) we observed that at particular loci, and even for entire chromosomal arms (Fig. S7), the genetic basis

of pigmentation evolution was unpredictable based on mapping data alone. This likely reflects the role of

pleiotropy and constraint during complex trait adaptation in natural settings characterized by a high

diversity of selection pressures, such as our outdoor mesocosm (Fig. 1, Fig. 3D. 3D). This unpredictable

genetic architecture of adaptation was also evident at the known loci of major effect: allele frequency

shifts within the *ebony* locus, as well as the largest effect locus on 2R, predicted evolution of lighter

pigmentation; while the locus containing *bab* predicted evolution of darker pigmentation and the *tan* locus

predicted that pigmentation would not evolve (arrows in Fig. 3D).

Figure 3 | Genetic architecture of female abdominal pigmentation. (A) The pigmentation score of the founding inbred lines compared with the light vs. dark frequency differences across all samples for the haplotypes originating from these lines. Inbred line pigmentation is highly correlated to the haplotype's light vs. dark enrichment (linear regression p << 0.001). Colored points indicate inbred lines that drive the significance of relationship between pigmentation score and enrichment in color fractions (gold points correspond to inbred lines enriched in light color fraction and dark brown to lines enriched in dark color fractions) (B) Distribution of the difference between average expected light frequencies and average expected dark frequencies for the top 1% of SNPs from a model only considering the impact of trait value on allele frequency variation (Trait-Only), versus a model with an additional term that represents the expected frequency in the sample based on the SNPs' ancestry (Ancestry). (C) Manhattan plot depicting nominal p-values derived from an ancestry-informed mapping of SNP association with pigmentation variation. The dashed horizontal line corresponds to an empirical FDR threshold of 0.0005, derived via permutations. (D) Genomic distribution of SNPs confined within N = 42 unlinked, independent loci underpinning pigmentation variation. SNPs are plotted as a function of their genomic coordinate and regression-derived nominal p-value (as in (C)). Shaded regions denote the stop and end points of each of the identified unlinked, independent loci. Shaded region color then denotes the predominant behavior of light-associated SNPs within each locus through time: red shading indicates loci with light-associated alleles favored throughout evolution of lighter pigmentation (18 total loci; binomial test, FDR < 0.05), blue shading indicates loci in which the light-associated alleles are predominately selected against (7 total loci; binomial test, FDR < 0.05), and grey shading indicates loci with no signature of selection on light-associated alleles (17 total loci; binomial test, FDR > 0.05). The location of the canonical pigmentation genes, *bric-a-brac (bab), ebony,* and *tan*, are indicated on the Manhattan plots. Potential candidate genes underlying the additional loci observed throughout the genome are discussed in the Supplement.

The effect of pleiotropy on genomic prediction differs across complex and simple environments

The pleiotropy we quantified above manifested in a field setting where replicate populations were exposed to a dynamic suite of fluctuating selection pressures. This begs the question of whether the magnitude of pleiotropy would be subdued in a simpler, lab-based environment with fewer selective pressures. Our study design provided a unique opportunity to test this hypothesis via comparing genomic predictions of trait evolution for the replicate populations evolved in the field mesocosm relative to those that evolved within the indoor, lab-based mesocosm (Figs. 1-2). We first quantified whether light-associated alleles were favored during the evolution of lighter pigmentation, finding that their genome-wide frequencies increased significantly throughout trait evolution in each environment, relative to background allele frequency change (paired t-test p-values << 0.001; Fig. 4A; also see Fig. S8-S9). Next, we used the relative effect sizes produced by our mapping data, as well as their magnitude of allele frequency during trait evolution, to provide a genome-wide summary statistic of pigmentation evolution in each environment (Fig. 2A, Methods). This analysis is analogous to the polygenic score-based predictions of trait evolution that are pervasive in studies of human adaptation (Akbari et al., 2024; Cox et

al., 2019; Mathieson et al., 2015), and correctly inferred that lighter pigmentation evolved in both the

indoor and outdoor environment (Fig. 4B). Interestingly, however, the relative predicted magnitude of

phenotypic change was greater in the indoor environment, despite our phenotypic assays showing that

replicate populations in this environment evolved to a lesser degree (Fig 2A). Such a disconnect between

observed patterns of phenotypic evolution and our genomic inference of the process provides strong

evidence of a context-dependent pleiotropy across environments – the more complex environmental

conditions in the outdoor mesocosm induces selection on a greater number of traits overall, leading to

more pleiotropic constraint at pigmentation-associated loci and poorer genomic prediction.

We investigated context-specific pleiotropy further with a focus on the *bab* locus, a canonical

pigmentation gene identified by our mapping and for which raw allele frequency trajectories showed

evidence of pleiotropic constraint (Fig. 3D). By excluding variation in this locus from our genomic

prediction, we observed that the inferred relative magnitude of pigmentation evolution increased in both

environments and, strikingly, to a much greater degree in the outdoor environment (Fig 4C). This analysis

thus provides a key example of how pleiotropy at a single locus can skew genomic predictions of

phenotypic evolution. Furthermore, it elucidates how studying such processes in a natural setting may

reveal properties of segregating variation that are masked in simpler, lab-based environments.

Interestingly, regenerating these predictions using variation within the canonical pigmentation loci alone

led to a spurious prediction of darker evolution in the outdoor environment (Fig. S10), while excluding

variation in these loci from our prediction recovered the correct inference of trait evolution (Fig. S11).

Thus, the key takeaway of these results is that accurate genomic prediction of trait mapping hinges upon a

comprehensive, well-resolved genetic map that includes both the largest-effect loci, as well as those of

more moderate or minor impact on trait variation.

As our genomic predictions of trait evolution were generated from mapping results derived from

samples evolved in the outdoor mesocosm, we next quantified if and how these predictions differed when

derived from mapping of the indoor populations sampled at week 8. Using the same mapping procedure,

we found that while this independent set of mapped loci correctly predicts trait evolution in the indoor

environment, it fails to predict evolution of lighter pigmentation in the outdoor environment (Fig. 4D).

This discrepancy in genomic prediction of trait evolution in the outdoor environment (i.e., comparing

magenta distributions in Fig. 4B vs. Fig. 4C) potentially indicates epistasis-driven differences in the

genotype-phenotype map that arise between environments of selection. Strikingly, but also consistent

with those genomic predictions generated using the outdoor mapping data (Fig. 4B-C), we found that the

exclusion of *bab* locus variation allowed us to correctly predict trait evolution in the outdoor environment

using the indoor mapping data (Fig. 4E). This further exemplifies the pleiotropic constraints of this large

effect locus on trait evolution and the consequences therein on genomic prediction.

Figure 4 | **Prediction of phenotypic evolution in simple and complex environments**. (A) Genome-wide allele

frequency change at pigmentation-associated SNPs identified in the outdoor, week 8 mapping population and

quantified throughout trait evolution in the outdoor (purple) and indoor (blue) mesocosm, relative to matched

control shifts (grey distributions). (B) Distribution of Z-scores for bootstrapped estimates of genomic predictions for

pigmentation evolution in the outdoor (magenta) and indoor (turquoise) mesocosm using SNPs identified in the

outdoor, week 8 mesocosm. Positive Z-scores indicate net lighter pigmentation evolution while negative values

would indicate net darker evolution. Z-scores were computed by comparing inferred relative phenotypic change

using mapped SNPs to a that computed using a set of matched control SNPs (grey distributions). (C) Same as in

panel (D), but genomic predictions discard pigmentation-associated SNPs within the *bab* locus. (E-F) Same as (D-E)

but for pigmentation-associated SNPs identified via mapping the week 8 indoor populations.

Stable genotype-phenotype mapping throughout evolution and across environments of selection

How well do genotype-phenotype maps derived from a single population, at a single point in time,

transfer, and allow phenotypic prediction, across genetically diverged populations? This represents

another fundamental question in complex trait genetics and evolution that we could address within the

context of our longitudinal, multi-environment study. Specifically, we observed a pronounced signature

of genome-wide divergence associated between our lab-based, indoor environment and the outdoor, field

mesocosm (Fig. 2B), and previously reported substantial evolutionary change within each of these

environments throughout the observation period monitored in this study (Bitter et al., 2024a, 2024b). One

implication of this environmental and temporal genomic divergence is that it may induce shifts in the

epistatic landscape and/or elicit context dependent-dominance of pigmentation-associated loci. In effect,

these processes may alter the marginal effect of pigmentation-associated loci and generate distinct

genotype-phenotype maps that preclude phenotypic prediction across mapping populations (Novembre

and Barton, 2018).

We evaluated the stability of the genotype-phenotype map by re-implementing our mapping

approach, as described above, on the replicate populations sampled in outdoor mesocosms at week 2, as

well as the indoor populations sampled at weeks 2 and 8. The number of loci underpinning variation

across all mapping populations was comparable (ranging from 30-43 independent loci, Supplementary

Data File 1; Table S2). Furthermore, as expected, the total genomic distance spanned by these loci

declined from weeks 2 to 8 (from ~1 Mb at week 2 to ~700 Kb at week 8) due to additional

recombination events that allowed for more localized mapping.

The similarities in architecture inferred in the lab and field environments at week 8 are clearly

apparent in the mirrored Manhattan plots presented in Fig. 5A, where regions of heightened pigmentation-

associated variation are replicable across mapping environments. Notably, loci identified across all four

mapping populations contained the canonical pigmentation genes *tan*, *ebony*, and *bab* (Fig. S12),

demonstrating consistency of mapping as well as the pronounced effect these genes have on pigmentation

variation in this species. We next took two approaches to rigorously quantify, genome-wide, the similarity

in loci harboring pigmentation-associated variation across the four independent mapping populations.

First, we assessed whether the start and end points of loci identified in each independent mapping

population were closer than expected by chance: for each set of loci identified within a particular mapping

population, we selected the locus-defining, midpoint SNP (i.e., that with the most significant p-value) and

then quantified the number of times it fell within a locus in other three sets of mapped loci (each locus

could therefore overlap 0 to 3 times with the remaining mapping populations). The number of overlaps

across all loci for a particular mapping population were then summed and compared to an empirical null

distribution derived by iterating this procedure $N = 2000$ times using a set of randomly selected SNPs (see

Methods). Across sets of mapped loci, midpoint SNPs were far more likely to fall within a locus of an

independent mapping than expected by chance (Fig. 5B), demonstrating that the regions harboring

pigmentation-associated variation were conserved across mapping populations. To probe this consistency

further, we quantified the distance between locus midpoints among mapping populations. As expected,

these distances were significantly closer than expected based on an empirical null (Fig. 5C, see Methods),

further indicating the conservation of pigmentation associated loci across mapping populations.

Given the pronounced mapping stability at the locus-level, we next interrogated whether

pigmentation-associated SNPs were also shared across mapping populations. To do so, we reidentified

trait-associated SNPs using our ancestry-informed GLM in groups of 5 (of the 10 total) replicates per

mapping population. This then allowed for a balanced comparison of the overlap in trait-associated SNPs

within subsets of the same mapping population (e.g., overlap in trait associated SNPs between subsets of

five replicates from the outdoor week 8 samples), and between mapping populations (e.g., overlap in trait

associated SNPs between subsets of five outdoor week 8 and five indoor week 2 replicates). Strikingly,

we found that the amount of overlap in SNPs identified between subsets of the same mapping population

did not exceed that between mapping populations, either from the same environment and a different time

point or even between environments (Fig. 5D; Fig. S13). In the supplement, we further quantified this

SNP-level parallelism using a leave-one-out cross validation scheme (Fig. S14).

While our quantification of overlap in pigmentation-associated loci across mapping populations

indicated a strong stability of the genotype-phenotype map (Fig. 5B-D), we observed several conspicuous

instances in which genomic regions harboring pigmentation-relevant variation for one mapping

population appeared to lack pigmentation-associated variation in an independent mapping population.

Several of such instances can be observed in the mirrored Manhattan plot in Fig. 5A, where the start and

end points of pigmentation loci (indicated by shaded regions) within one environment exhibit non-

overlapping start and endpoints with any of the loci identified in the second environment. We suspect

these environment-specific loci to potentially indicate epistatic interactions driving differences in the

marginal effect of pigmentation-associated SNPs across mapping populations. In the Supplement, we

show that such instances of environment-specific loci are not solely due to technical noise, but rather

treatment-specific marginal effects of pigmentation-associated loci and variation (Fig. S15 & S16), as

first indicated by our genomic prediction results presented above (Fig. 4). Nonetheless, the vast majority

of loci are overlapping across populations (e.g., 30 of 43 loci identified in the week 8 outdoor population

overlapped with at least one of the 42 loci in the week 8 indoor population), and thus the dominant signal

in the data is one of strong parallelism. Indeed, when we exclude treatment-specific loci from our

calculations of predicted phenotypic evolution, our ability to correctly predict trait change across

environments is unaffected (Fig. S17).

Finally, given the overall stability of SNP and locus mapping, we queried our ability to predict

relative phenotypic values across independent populations using allele frequency data alone. Specifically,

we inferred the relative pigmentation of phenotypic groups (i.e., the light, midpoint, and dark color

fractions) both within and across mapping populations using our mapping data (see Methods). Strikingly,

we found that we predicted relative pigmentation scores with statistically indistinguishable accuracy

between, as well as within, mapping populations (Fig 5E; Fig. S18). In concert, these analyses suggest

that epistasis or dominance shifts caused by the genetic evolutionary change in our experiment were

insufficient to preclude genotype-phenotype map portability and phenotypic prediction. While seemingly

at odds with expectations and empirical evidence of limited portability of mapping results across

independent populations (de Bakker et al., 2006; Mostafavi et al., 2020; Privé et al., 2022), it bolsters

more recent research that have demonstrated increased portability under instances in which mapping is

conducted in a high-powered manner and the loci underpinning trait variation are fine mapped (Hu et al.,

2025; Wu et al., 2025)

Figure 5 | Epistasis does not preclude portability of trait mapping. (A) Mirrored Manhattan plots of genetic

architecture quantified at week 8 in the outdoor (top) and indoor (bottom) environments. Only SNPs with p-value

corresponding to empirical FDR < 0.0005 p-value are depicted. For each mapping, shaded regions correspond to

unlinked loci underpinning pigmentation-associated variation (B) Observed (diamonds) vs expected (grey violin

distributions) overlap between pigmentation associated loci across mapping populations. (C) Observed (diamonds)

vs expected (grey violin distributions) distances between midpoint SNPs of trait associated loci between mapping

populations. (D) Observed percent overlap of trait-associated SNPs identified within and between mapping

populations. (E) Predicted relative phenotypic values for dark, midpoint, and light color fractions using genotype-

phenotype mapping conducted in the outdoor mesocosm at week 8 where positive and negative values indicate

lighter or darker pigmentation relative to the baseline population, respectively. Predictions (diamonds) were

conducted for each trait value group and replicate cage. Colored violins represent 1000 bootstrapped predictions of

each predicted, relative phenotypic value.

Discussion

505   The central goal of this study was to rigorously evaluate the predictability of complex trait

adaptation. Specifically, we aimed to quantify the extent to which we could correctly infer complex trait

evolution given genomic data alone, and also predict which loci came under selection during trait

evolution. Leveraging a model complex trait (abdominal pigmentation), a model species (*Drosophila*

melanogaster), and a highly powered, longitudinal study design, we showed that well-resolved genotype-

phenotype maps can be used to correctly predict if, and in what direction, a complex trait evolves.

However, the specific genetic architecture of adaptation (i.e. which loci show signatures of selection

during trait evolution) was largely unpredictable based on mapping data alone. Specifically, despite a

genome-wide increase in the frequency of light-associated alleles, pigmentation-associated variation in

several loci was observed to be net-neutral, or even selected against in a 'counter-gradient' manner,

throughout the evolution of lighter pigmentation (Albecker et al., 2025; Paaby et al., 2014). This dynamic

exemplifies underlying pleiotropy in complex trait variation which, under a regime of correlated

selection, can ultimately constrain such variation's impact on trait evolution (Walsh and Blows, 2009). It

is unsurprising that this dynamic emerged in our outdoor mesocosm, as we have shown that multiple

fitness-relevant traits to evolve across seasons in this system (Rudman et al., 2022).

520   A key additional component of this study was to assess if, and how, pleiotropic constraint

differed across distinct environments of selection. Specifically, we paired our outdoor mesocosm with an

indoor mesocosm in which the same founding population evolved concurrently in the context of the same

general biotic shifts (e.g., increases in population density), but the absence of the multivariate abiotic

fluctuations that characterize the outdoor habitat (Fig. 1). In effect, we showed that the pleiotropic

constraint of several large-effect pigmentation-associated loci (e.g., *bab*) was alleviated in the simpler,

lab-based environment, leading to better overall genomic predictions of trait evolution. This finding thus

exemplifies a context-dependence of pleiotropy, whereby more complex environments elicit an increase

in pleiotropic constraint. Furthermore, from a practical sense, this implies that generic principles of the

evolutionary process itself may be obscured in simplified, lab-based environments and bolster recent calls

to study the ecology and evolution of natural populations in realistic settings (Bergelson et al., 2021).

While the impact of pleiotropy was pronounced, we found epistasis to have a less detrimental

effect on the prediction of complex trait variation overall. Specifically, we found a striking stability of the

genotype-phenotype map across four independent mapping populations sampled at two points in the

evolutionary process and evolved in two distinct environments. Furthermore, we showed that relative

phenotypic predictions were effectively as accurate across environments of selection, as within.

Therefore, at least within the confines of the magnitude and timescale of evolution manipulated here,

epistasis (and/or context-dependent dominance shifts) may not inhibit a general transferability of mapping

results across genetically diverged populations. While this finding may be at odds with some studies

evaluating the portability of mapping data across diverged populations, it bolsters more recent work

demonstrating increased portability under instances in which trait architecture has been well-resolved

(i.e., fine-mapped) such as in the high-powered mapping conducted here (de Bakker et al., 2006; Hu et al.,

2025; Mostafavi et al., 2020; Privé et al., 2022; Wu et al., 2025)

It warrants note that the unprecedented insight into the predictability of evolution was only made

possible by several key aspects of our methodological approach: our generation of haplotype-informed

allele frequencies at extremely high coverage, in combination with an exceptional amount of biological

replicates per mapping treatment and time point, provided the power to produce well-resolved genetic

maps (Long et al., 2025; Tilk et al., 2019). Furthermore, our longitudinal analysis of trait variation using a

single founding population was the only way to exclude confounding variables such as unique patterns of

demographic history and linkage when assessing the role of pleiotropy and epistasis on complex trait

prediction (Lohmueller, 2014; Novembre and Barton, 2018). In concert, this study design then allowed us

to both reaffirm canonical pigmentation genes, as well as identify new candidates, underpinning variation

in this classic complex trait. Crucially, our accurate genomic prediction of trait evolution hinged upon this

well-resolved genotype-phenotype map, as prediction using canonical gene variation alone would have

led to incorrect inferences of trait evolution. Our study therefore exemplifies the critical roles of study

design and power in the rigorous evaluation of fundamental evolutionary dynamics.

Looking forward, our results are pertinent to numerous applications of genomic data: predicting

phenotype across disparate populations, inferring past selection using time-series and ancient DNA

genomic data, or manipulating contemporary variation to shape future evolutionary trajectories (e.g.,

augmenting population resilience to future global change conditions). The critical contingency of

effectively implementing such approaches, however, will be the procurement of well-resolved genotype-

phenotype maps and knowledge of the focal trait variation's pleiotropy and epistasis, a venture that may

not be feasible across study systems.

Methods

Experimental overview

A genetically diverse population of outbred *D. melanogaster* was generated via outcrossing ten male and female individuals from a total of 76 inbred lines, originally collected from Linvilla Orchards, Media, PA (Behrman et al., 2018). The outbred population was allowed to recombine and expand for 4 generations, after which 500 males and 500 females were used to seed each of 10 outdoor and indoor replicate cages. The outdoor mesocosm consisted of ten replicate, two m^3 mesh cages, each harboring a single dwarf peach tree, and was exposed to natural environmental conditions, insect, and microbial communities (Bitter et al., 2024a). The indoor mesocosm then consisted of ten replicate, ~0.5 m^3 cages housed in a temperature-controlled laboratory at the University of Pennsylvania. Constant food was supplied to the replicates within each environment, whereby 400 ml of Drosophila media was provided in 900 cm^3 aluminum loaf pans within each cage three times per week (Bitter et al., 2024a, 2024b). Monitoring of phenotypic and genomic variation in the replicates began two weeks after cage founding and continued for a period of eight total weeks and six generations of evolution. Generation time estimates were produced using a degree day model (Kamiyama et al., 2020).

Population-wide pigmentation variation was quantified in our panel of inbred lines, the recombinant outbred population after 4 generations of expansion and recombination, and the evolving replicate populations in the outdoor and indoor environments at weeks 2 and 8 (20 July and 1 September, separated by a period of 5 generations). To quantify evolved changes in pigmentation, eggs were collected overnight from all replicate cages (indoor/lab and outdoor/field), after which they were transferred to common garden conditions (25°C, 12L:12D) where they were reared for 2 generations. Following density control of F2 eggs (30 +/-5) in replicate vials, adult female abdominal pigmentation (N = 30 individuals per replicate) was scored as the proportion of pigmentation observed on tergites 5, 6, and 7 (David et al., 1990). The sum of these scores across the three tergites provided the total pigmentation score for the assayed individual.

We quantified genome-wide allele frequencies throughout evolution weekly in the outdoor environment and at weeks 2 and 8 in the indoor environment. At each collection point, eggs were collected overnight from each cage and then transferred to common garden conditions where larval development took place. 100 random females from the eclosed adult cohort were then collected and stored in ethanol at -20°C for later DNA extraction and sequencing (see below).

At the weeks two and eight sampling points, we conducted a tail-based mapping of pigmentation variation for each mapping population. Specifically, we first quantified pigmentation variation within each treatment using a random sample of sorted, 100 F2 females (3-6 days post-eclosure) and the visual scoring methods described above (David et al., 1990). Then, for each replicate across both environments,

we segregated individuals into one of three trait value groups: (1) those exhibiting pigmentation within

the upper 15% of the population-wide phenotypic distribution, (2) those exhibiting pigmentation within

the lower 15% of the trait distribution and (3) a random sample from those representative of the midpoint

(~25th-75th percentile) of the trait distribution. Seventy-five individuals from each trait value group and

replicate were then stored in ethanol and at -20ºC for later DNA extraction and sequencing. In total this

resulted in 120 samples available for mapping: three trait value groups across ten replicate cages sampled

from two environments and at two time points.

Sequencing and allele frequency estimation

Genomic DNA was extracted for both mesocosm time-series and trait mapping samples using

NEB Monarch DNA kit. Following sample QC via nanodrop and qubit, whole genome libraries were

constructed using Illumina DNA prep kit. Each sample was sequenced to an equal coverage of 8x using

Illumina Novaseq 6000. From the resulting sequencing data, we generated haplotype-informed allele

frequencies using previously described methods (Kessner et al., 2013; Tilk et al., 2019). This method

computes allele frequencies at sites previously identified in the genome sequences of each founding

inbred strain using an expectation-maximization algorithm that estimates haplotype frequencies in a

pooled sample of mapped sequencing reads. Haplotype inference was conducted in window sizes that

varied proportionally to the length of un-recombined haplotype blocks expected as a function of the

estimated number of generations since original inbred line outbreeding (Kessner et al., 2013). After

filtering sites an average minor allele frequency >= 0.02 in the founder population, and present in at least

one evolved sample at a minor allele frequency > =0.01, we retained 1.9 M SNPs for analysis.

Genome-wide patterns of variation across environments and throughout evolution

We first assessed genome-wide patterns of variation across all samples used in trait mapping

using principal component analysis (*prcomp* package, R) and normalized allele frequency data across all

1.9 M SNPs. We projected samples onto the first two principal components and visualized the data by

shaping and coloring points according to different underlying variables (e.g., trait value group, treatment)

to infer which variables may dominate variance in allele frequencies across samples. We quantified

whether samples segregated along the first two principal components in accordance with trait value group

or environment using linear regression and ANOVA, respectively.

Ancestry-informed association analysis of pigmentation variation

Our general approach to mapping pigmentation was to identify SNPs harboring alleles with

systematic frequency changes across the light, midpoint, and dark trait value groups. First, however, it

was important to consider that due to the finite number of haplotypes from the founding inbred lines used

to derive our outbred population, a significant association between a SNP and trait value group could

indicate its biological impact on pigmentation variation and/or its inbred-line ancestry. This effect could

appear at the level of entire chromosomes due to long range patterns of linkage, as well as more locally

around causal loci. We quantified the magnitude of this effect genome-wide by computing the average

contribution of haplotypes coming from each inbred line to each trait value group (i.e., the difference in

haplotype frequency between the light and dark tails) and quantified the relationship of this value with the

pigmentation score of each inbred line using linear regression and Pearson's correlation coefficient.

We interrogated the effect of linkage on SNP mapping further by running two generalized linear

models (GLM; logistic link function and quasibinomial error variance). The first model only took into

consideration the impact of trait value group on variation in allele frequencies (model form: allele

frequency ~ trait value; where allele frequencies are normalized by the SNP's effective coverage and the

total number of chromosomes sampled). If the effects of ancestry on SNP association were pervasive,

SNPs identified via this approach would be enriched in those SNPs expected to have elevated light to

dark frequency differences based on their association with particular haplotypes alone. To test this, we

selected the top 1% of the SNPs (based on nominal p-value from the GLM) and calculated their expected

frequency difference between light and dark color fractions based on estimates of the average contribution

of each inbred line to that sample and the association between that SNP and inbred line. We then

compared these results to a separate GLM model in which the haplotypic ancestry was accounted for.

This model then regressed allele frequencies both on pigmentation trait value, as well as each SNPs

association with the line contribution to the sample using average genome-wide haplotype frequencies. In

accordance with the previous, ancestry-agnostic, model we used a logistic link function and

quasibinomial error variance, and allele frequencies were weighted to account for variation in coverage

and the finite number of chromosomes sampled. We then compared the density distributions of expected

light-dark frequency differences for SNPs identified via each of these distinct models using a KS test.

Following GLM model structure selection, we used permutations to define a final set of trait-

associated SNPs. Specifically, we randomly shuffled trait value data and re-ran our regressions N = 100

total times to generate an empirical null distribution. We then considered a SNP significant if its nominal

p-value from the empirical GLM fell in the top 0.05% of its permuted p-values (corresponding to an

empirical FDR threshold < 0.0005).

The set of pigmentation-associated SNPs identified via GLM was used to quantify the number

and distribution of unlinked loci underpinning trait variation. We developed an iterative procedure for

this, whereby we first identified the most significant SNP on each chromosomal arm, as well as its

surrounding, locally un-recombined region (defined by the recombination rate and number of generations

since cage founding; Tilk *et al*. 2019). We then eliminated all trait-associated SNPs on the chromosomal

arm (i.e., those with a significant GLM p-value) whose relative presence in the light and dark fractions

can be explained by haplotype frequencies (i.e., ancestry) in the locally un-recombined focal region alone.

As the significance of these SNPs could be driven either by linkage to the focal top SNP, or variation in

the trait itself, we eliminated these and defined the trait-associated locus as the locally unrecombined

region (as defined using number of generations since founder line outbreeding and genome-wide

recombination rates) around the lowest p-value SNP. The specific threshold used in SNP elimination was

defined via permutations: we iteratively subset the 10 replicates into two equal subsets (N = 5). In one

subset, termed 'real-data', light-dark haplotype frequency differences within the focal region were

computed as above. In the other subset, the dark-light haplotype frequency difference was calculated (i.e.,

termed 'sign-flipped' data). In each subset we sequentially eliminated percentiles of SNPs starting with

the top and bottom 2.5% of expected light-dark frequency differences and selected our threshold as the

percentile at which the number of true significant SNPs eliminated in the real-date matched that of the

sign-flipped data (Fig. S19). From the set of remaining SNPs (those not eliminated in the first iteration),

we identified the SNP with the most significant nominal p-value and repeated the procedure. This was

iterated until we exhausted all trait-associated SNPs on the chromosome. For each chromosomal arm, we

then quantified how the number of trait-associated SNPs decreased as a function of these iterations (Fig.

S20). Finally, we merged all loci with overlapping breakpoints to procure a final set of unlinked and

independent trait-associated loci.

We additionally explored which candidate genes may be underlying the identified loci.

Specifically, we identified genes within each locus with a GO term associated with pigmentation

(excluding eye pigmentation) or cuticle development/molting. For each of the identified genes, we then

quantified whether the number of pigmentation-associated SNPs within the gene (identified via mapping)

was greater than expected based on their genome-wide abundance using a binomial test with a Benjamini-

Yekutieli false discovery rate correction.

Comparison of mapping to previous association studies

We leveraged results reported by two previous genome wide association studies (Bastide et al.,

2013; Dembeck et al., 2015) to quantify the consistency of mapping of pigmentation across disparate

populations and studies. In brief, Bastide et al. (2013) conducted a pooled-GWAS approach using wild

populations of *D. melanogaster* from Austria and Italy, and focused on female pigmentation on tergite 7.

Dembeck et al. (2015) conducted a GWAS using individual *D. melanogaster* lines from the Drosophila

Genetics Reference panel (Mackay et al., 2012), focusing on pigmentation variation of tergites A5 and

A6. We evaluated whether all SNPs identified as trait-associated in the Dembeck study, and the top 100

SNPs from Bastide et al., exhibited more overlap with our unlinked loci than expected by chance.

Specifically, we compared the number of SNPs falling within each locus to a random set of SNPs within

the chromosome of the same test set size as those obtained from the comparison studies. We iterated this

process 1000 times and generated an empirical p-value given the observed overlap, relative to the

randomly generated 1000 overlaps. We then re-conducted this analysis, but excluded those within loci

containing *tan*, *ebony*, and *bab* to infer the extent to which shared signal across studies was driven by

these large effect genes.

Predicting evolution using mapping results

We quantified whether mapped SNPs displayed a behavior during lighter pigmentation evolution

that would be expected based on this mapping data alone. Specifically, we computed the frequency shifts

for all light-associated alleles at pigmentation-associated SNPs between weeks 1 and 8 of evolution in the

outdoor mesocosm, separately for of the unliked, trait-associated loci. We compared the resulting

distribution to one derived via a set of matched control SNPs (matched on chromosome and starting

frequency (+-/ 2.5%), and excluding those SNPs within 500 Kb of the respective target SNP) using a

binomial test. In effect, we quantified whether the proportion of light-associated SNPs within each locus

shifting in a manner concordant with that predicted by mapping (i.e., increasing in frequency as lighter

evolution evolved) was significantly greater than expected based on our matched control distribution.

We then compared how our genomic predictions of trait evolution differed between the semi-

natural, outdoor mesocosms and relatively simple, indoor environment. First, we quantified the trajectory

of trait-associated SNPs (identified in our outdoor, week 8 mapping population) separately in the outdoor

and indoor mesocosm throughout the period of lighter pigmentation evolution (weeks 1 through 8). We

compared each resulting distribution to a set of matched control SNPs using a binomial test to infer

whether the direction and magnitude of light-favored allele frequency shifts exceeded background allele

frequency movement. This analysis was then re-conducted separately for SNPs within each of the top loci

on chromosomal arms 2L and 2R, as well as the loci containing the genes *bab*, *ebony*, and *tan* on

chromosomal arms 3L, 3R, and X, respectively. Furthermore, the distribution of trait-associated SNP

trajectories throughout trait evolution within each environment were compared directly to each other

using a paired t-test.

Finally, we generated genomic predictions of trait evolution using an approach that explicitly

accounted for the observed effect-size of each SNP on trait variation observed via mapping. Specifically,

we summed the frequency shifts of light-associated alleles, this time weighting each allele's shift by the

associated SNP's GLM regression coefficient. This summation then represents an effective shift in the

polygenic score of the population's pigmentation through time, whereby positive values indicate the

evolution of lighter pigmentation and negative values, darker pigmentation. We bootstrapped these

estimates N = 1000 times and their significance was then assessed using Z-scores computed via a

comparison of the estimate to a distribution generated via N = 1000 matched control SNP sets (matched

on chromosome and starting frequency). We considered the empirical, bootstrapped value significant if its

confidence intervals did not overlap with the matched control distribution.

Stability of mapping throughout evolution and across environments of selection

We next sought to determine the stability of the genotype-phenotype map across independent mapping populations (i.e., each mapping environment and timepoint). Accordingly, we re-conducted our mapping procedure, as described above, for the indoor weeks 2 and 8, as well as the outdoor week 2 mapping population. First, we asked if the proximity of loci identified across mapping populations were more similar than expected by chance. To address this, for each of the four genotype-phenotype maps we identified the top SNP of each locus (the SNP with the lowest p-value, and also the locus midpoint) and summed the number of times it fell within a locus identified in one of the independent mapping populations. This then produced a score, ranging from 0-3, for each locus and these scores were summed across all loci to produce an empirical overlap score for each mapping population (note that to increase power of this analysis, we selected focal SNPs using the set of pre-merged loci for each mapping population). To determine if the observed overlaps were greater than expected by chance, we generated an empirical null distribution via random sampling of sets of SNPs for each genotype-phenotype map and re-computing overlap scores with each independent mapping (a procedure iterated N = 2000 times for each of the four genotype-phenotype maps). We subsequently interrogated whether the distance between locus-defining midpoint SNPs across sets of loci were closer than expected by chance. Specifically, as before, we first identified the top SNPs for each set of loci from every genotype-phenotype map, and then computed its distance to the closest top SNP of each of the remaining three mappings. This distance metric was averaged to produce an average proximity score which, as above, was compared to a distribution of scores derived via a random set of SNPs (N = 2000 times).

We next aimed to quantify the extent to which the identity of specific pigmentation-associated SNPs was shared across independent mapping populations. To do so, we compared the overlap in identified SNPs within and between independent mapping populations. To conduct this analysis in a balanced manner (i.e., compare sets of SNPs identified using the same number of replicate cages/populations), we iteratively subset our set of 10 replicates from each mapping population into groups of 5 replicates and identified pigmentation associated SNPs in each subset using the ancestry-informed regression introduced above. This was iterated with each possible combination of subset replicates, yielding 126 total sets of significant SNPs per mapping population. The percentage of shared SNPs was then calculated for subsets of the same mapping population (e.g. overlap between pigmentation-associated SNPs identified using different subsets of 5 replicate populations from the outdoor, week 8 replicates), and compared to those observed between independent mapping populations (e.g. overlap between sets of pigmentation associated SNPs identified using 5 replicate populations from outdoor week 8 replicates and sets of SNPs from independent treatment and/or time point). Whether the

amount of overlap in identified SNPs within vs. between mapping populations differed was evaluated

with a planned contrast using a one-way ANOVA model.

We next sought to quantify whethe there existed mapped loci with environment-specific marginal

effects on pigmentation variation. Accordingly, we first isolated those loci from the outdoor and indoor

week 8 mapping populations that had non-overlapping breakpoints with any of the loci in the opposing

environments (N = 13 total unique outdoor loci, and N = 13 total unique indoor loci). We then identified

all pigmentation-associated SNPs within each unique locus and quantified the light dark-frequency

difference for the light-associated allele for the environment in which it was identified, as well as the

environment that did not contain the locus. Evidence for environment-specific marginal effects of each

locus was then determined by comparing the distribution of light-dark frequency shifts for each locus

between environments (using a Wilcoxon Signed Rank Test), as well as to a distribution derived from an

equally sized set of matched control SNPs (using a Mann Whitney U test).

Finally, we sought to determine if genotype-phenotype maps produced at a particular point in

evolution, and within a particular environment of selection, could correctly predict phenotypes of

individuals sampled at an independent time and environment. To do so, we used those SNPs identified in

our outdoor week 8 mapping and calculated the relative expected phenotypic score for trait value groups

(e.g., light, midpoint, and dark color fractions) within this focal mapping, as well as all remaining,

independent mapping populations. Specifically, for each light-associated allele at all pigmentation-

associated SNPs, we calculated the difference in frequency between the baseline population and each

pooled sample. This allele frequency change was weighted in accordance with the SNPs effect size on

trait variation as computed via mapping (i.e., by multiplying that by the regression coefficient of our

ancestry-informed GLM by the light-favored allele frequency change of that sample). These values were

then summed across all SNPs for each sample, providing a metric of the relative lightness of the sample,

as predicted by mapping (values greater than 0 indicating lighter pigmentation and less than 0 darker). We

then bootstrapped this value 1000 times for each sample. The accuracy of relative predicted phenotypic

scores was then evaluated using three criteria: (i) whether the dark samples had, on average, predicted

phenotypic values below 0, (ii) whether the light samples had, on average, predicted phenotypic values

above 0, and (iii) if there existed a significant linear association between trait value group and relative

predicted phenotypic value. The latter was tested using a simple linear regression with the model trait

value group ~ phenotype prediction, where trait values groups were assigned a numeric value (dark = 0,

midpoint = 1, and light = 2).

Code and data availability: All code necessary to replicate the analyses presented in this manuscript are

available at the following: https://github.com/jrhodes613/PigMap2021 .

Founder line sequences are available at the NCBI with the accession ID PRJNA722305. Upon manuscript

acceptance, sequences generated via pooled sequencing for this study will be available at the NCBI with

the accession ID PRJNA1031645 and phenotypic and allele frequency data will be uploaded to a dryad

repository.

Author contributions:

Conceptualization: MCB, DAP, PS

Experimental methodology: MCB, SB, HO, JB, DAP, PS

Formal analysis: MCB, JASR

Writing of manuscript: MCB, JASR, DAP, PS

Manuscript editing: all authors

Acknowledgements: We thank Lauren McIntyre, Jonathan Pritchard, and all members of the Petrov and

Schmidt lab for helpful feedback of experimental design and analysis and interpretation. We thank

Caitlynn T. Tran and Andy V. Huynh for assistance in pigmentation phenotyping. We thank Heloise

Bastide for an extended list of pigmentation-associated loci and results from Bastide et al 2013 (doi:

10.1371/journal.pgen.1003534).

Funding: We are grateful to our funding organizations: the National Science Foundation (NSF

PRFB 2109407 to M.C.B.), the National Institutes of Health (NIH R35GM118165-07 to D.A.P. and

NIH R01GM137430 to P.S.), and the Chan Zuckerburg Biohub.

References

Abdellaoui, A., Yengo, L., Verweij, K.J.H., Visscher, P.M., 2023. 15 years of GWAS discovery:
Realizing the promise. The American Journal of Human Genetics 110, 179–194.
https://doi.org/10.1016/j.ajhg.2022.12.011

Akbari, A., Barton, A.R., Gazal, S., Li, Z., Kariminejad, M., Perry, A., Zeng, Y., Mittnik, A., Patterson,
N., Mah, M., Zhou, X., Price, A.L., Lander, E.S., Pinhasi, R., Rohland, N., Mallick, S., Reich, D.,
2024. Pervasive findings of directional selection realize the promise of ancient DNA to elucidate
human adaptation. https://doi.org/10.1101/2024.09.14.613021

Albecker, M.A., Bittar, T.B., Trussell, G.C., Lotterhos, K.E., 2025. A Quantitative Survey of Cogradient
and Countergradient Variation in Nature. The American Naturalist 000–000.
https://doi.org/10.1086/737682

Bastide, H., Betancourt, A., Nolte, V., Tobler, R., Stöbe, P., Futschik, A., Schlötterer, C., 2013. A
Genome-Wide, Fine-Scale Map of Natural Pigmentation Variation in Drosophila melanogaster.
PLOS Genetics 9, e1003534. https://doi.org/10.1371/journal.pgen.1003534

Bastide, H., Lange, J.D., Lack, J.B., Yassin, A., Pool, J.E., 2016. A Variable Genetic Architecture of
Melanic Evolution in Drosophila melanogaster. Genetics 204, 1307–1319.
https://doi.org/10.1534/genetics.116.192492

Behrman, E.L., Howick, V.M., Kapun, M., Staubach, F., Bergland, A.O., Petrov, D.A., Lazzaro, B.P.,
Schmidt, P.S., 2018. Rapid seasonal evolution in innate immunity of wild Drosophila
melanogaster. Proceedings of the Royal Society B: Biological Sciences 285, 20172599.
https://doi.org/10.1098/rspb.2017.2599

Berardi, S., Rhodes, J.A., Berner, M.C., Greenblum, S.I., Bitter, M.C., Behrman, E.L., Betancourt, N.J.,
Bergland, A.O., Petrov, D.A., Rajpurohit, S., Schmidt, P., 2025. Drosophila melanogaster
pigmentation demonstrates adaptive phenotypic parallelism over multiple spatiotemporal scales.
Evolution Letters qraf008. https://doi.org/10.1093/evlett/qraf008

Bergelson, J., Kreitman, M., Petrov, D.A., Sanchez, A., Tikhonov, M., 2021. Functional biology in its
natural context: A search for emergent simplicity. eLife 10, e67646.
https://doi.org/10.7554/eLife.67646

Bickel, R.D., Kopp, A., Nuzhdin, S.V., 2011. Composite Effects of Polymorphisms near Multiple
Regulatory Elements Create a Major-Effect QTL. PLOS Genetics 7, e1001275.
https://doi.org/10.1371/journal.pgen.1001275

Bitter, M.C., Berardi, S., Oken, H., Huynh, A., Lappo, E., Schmidt, P., Petrov, D.A., 2024a. Continuously
fluctuating selection reveals fine granularity of adaptation. Nature 1–8.
https://doi.org/10.1038/s41586-024-07834-x

Bitter, M.C., Greenblum, S., Rajpurohit, S., Bergland, A.O., Hemker, J.A., Betancourt, N.J., Tilk, S.,
Berardi, S., Oken, H., Schmidt, P., Petrov, D.A., 2024b. Pervasive fitness trade-offs revealed by
rapid adaptation in large experimental populations of Drosophila melanogaster.
https://doi.org/10.1101/2024.10.28.620721

Blount, Z.D., Lenski, R.E., Losos, J.B., 2018. Contingency and determinism in evolution: Replaying
life's tape. Science 362, eaam5979. https://doi.org/10.1126/science.aam5979

Chiu, Y.-F., Justice, A.E., Melton, P.E., 2016. Longitudinal analytical approaches to genetic data. BMC
Genetics 17, S4. https://doi.org/10.1186/s12863-015-0312-y

Colosimo, P.F., Hosemann, K.E., Balabhadra, S., Villarreal, G., Dickson, M., Grimwood, J., Schmutz, J.,
Myers, R.M., Schluter, D., Kingsley, D.M., 2005. Widespread Parallel Evolution in Sticklebacks
by Repeated Fixation of Ectodysplasin Alleles. Science 307, 1928–1933.
https://doi.org/10.1126/science.1107239

Conover, D.O., Schultz, E.T., 1995. Phenotypic similarity and the evolutionary significance of
countergradient variation. Trends in Ecology & Evolution 10, 248–252.
https://doi.org/10.1016/S0169-5347(00)89081-3

Cox, S.L., Ruff, C.B., Maier, R.M., Mathieson, I., 2019. Genetic contributions to variation in human
stature in prehistoric Europe
. PNAS 116, 21484–21492. https://doi.org/10.1073/pnas.1910606116

David, J.R., Capy, P., Gauthier, J.-P., 1990. Abdominal pigmentation and growth temperature in
Drosophila melanogaster: Similarities and differences in the norms of reaction of successive
segments. Journal of Evolutionary Biology 3, 429–445. https://doi.org/10.1046/j.1420-
9101.1990.3050429.x

de Bakker, P.I.W., Burtt, N.P., Graham, R.R., Guiducci, C., Yelensky, R., Drake, J.A., Bersaglieri, T.,
Penney, K.L., Butler, J., Young, S., Onofrio, R.C., Lyon, H.N., Stram, D.O., Haiman, C.A.,
Freedman, M.L., Zhu, X., Cooper, R., Groop, L., Kolonel, L.N., Henderson, B.E., Daly, M.J.,
Hirschhorn, J.N., Altshuler, D., 2006. Transferability of tag SNPs in genetic association studies in
multiple populations. Nat Genet 38, 1298–1303. https://doi.org/10.1038/ng1899

Dembeck, L.M., Huang, W., Magwire, M.M., Lawrence, F., Lyman, R.F., Mackay, T.F.C., 2015. Genetic
Architecture of Abdominal Pigmentation in Drosophila melanogaster. PLOS Genetics 11,
e1005163. https://doi.org/10.1371/journal.pgen.1005163

Duncan, L., Shen, H., Gelaye, B., Meijsen, J., Ressler, K., Feldman, M., Peterson, R., Domingue, B.,
2019. Analysis of polygenic risk score usage and performance in diverse human populations. Nat
Commun 10, 3328. https://doi.org/10.1038/s41467-019-11112-0

Durvasula, A., Lohmueller, K.E., 2019. Negative selection on complex traits limits genetic risk prediction
accuracy between populations. bioRxiv 721936. https://doi.org/10.1101/721936

Enattah, N.S., Jensen, T.G.K., Nielsen, M., Lewinski, R., Kuokkanen, M., Rasinpera, H., El-Shanti, H.,
Seo, J.K., Alifrangis, M., Khalil, I.F., Natah, A., Ali, A., Natah, S., Comas, D., Mehdi, S.Q.,
Groop, L., Vestergaard, E.M., Imtiaz, F., Rashed, M.S., Meyer, B., Troelsen, J., Peltonen, L.,
2008. Independent Introduction of Two Lactase-Persistence Alleles into Human Populations
Reflects Different History of Adaptation to Milk Culture. The American Journal of Human
Genetics 82, 57–72. https://doi.org/10.1016/j.ajhg.2007.09.012

Grieshop, K., Ho, E.K.H., Kasimatis, K.R., 2021. Dominance reversals and the maintenance of genetic
variation. https://doi.org/10.48550/arXiv.2109.01571

Guo, Z., Tucker, D.M., Basten, C.J., Gandhi, H., Ersoz, E., Guo, B., Xu, Z., Wang, D., Gay, G., 2014.
The impact of population structure on genomic prediction in stratified populations. Theor Appl
Genet 127, 749–762. https://doi.org/10.1007/s00122-013-2255-x

Halldorsson, B.V., Eggertsson, H.P., Moore, K.H.S., Hauswedell, H., Eiriksson, O., Ulfarsson, M.O.,
Palsson, G., Hardarson, M.T., Oddsson, A., Jensson, B.O., Kristmundsdottir, S., Sigurpalsdottir,
B.D., Stefansson, O.A., Beyter, D., Holley, G., Tragante, V., Gylfason, A., Olason, P.I., Zink, F.,
Asgeirsdottir, M., Sverrisson, S.T., Sigurdsson, B., Gudjonsson, S.A., Sigurdsson, G.T.,
Halldorsson, G.H., Sveinbjornsson, G., Norland, K., Styrkarsdottir, U., Magnusdottir, D.N.,
Snorradottir, S., Kristinsson, K., Sobech, E., Jonsson, Helgi, Geirsson, A.J., Olafsson, I., Jonsson,
P., Pedersen, O.B., Erikstrup, C., Brunak, S., Ostrowski, S.R., Thorleifsson, G., Jonsson, F.,
Melsted, P., Jonsdottir, I., Rafnar, T., Holm, H., Stefansson, H., Saemundsdottir, J., Gudbjartsson,
D.F., Magnusson, O.T., Masson, G., Thorsteinsdottir, U., Helgason, A., Jonsson, Hakon, Sulem,
P., Stefansson, K., 2022. The sequences of 150,119 genomes in the UK Biobank. Nature 607,
732–740. https://doi.org/10.1038/s41586-022-04965-x

Hoekstra, H.E., Hirschmann, R.J., Bundey, R.A., Insel, P.A., Crossland, J.P., 2006. A Single Amino Acid
Mutation Contributes to Adaptive Beach Mouse Color Pattern. Science 313, 101–104.
https://doi.org/10.1126/science.1126121

Hof, A.E. van't, Campagne, P., Rigden, D.J., Yung, C.J., Lingley, J., Quail, M.A., Hall, N., Darby, A.C.,
Saccheri, I.J., 2016. The industrial melanism mutation in British peppered moths is a transposable
element. Nature 534, 102–105. https://doi.org/10.1038/nature17951

Hu, S., Ferreira, L.A.F., Shi, S., Hellenthal, G., Marchini, J., Lawson, D.J., Myers, S.R., 2025. Fine-scale
population structure and widespread conservation of genetic effect sizes between human groups
across traits. Nat Genet 57, 379–389. https://doi.org/10.1038/s41588-024-02035-8

Huang, W., Richards, S., Carbone, M.A., Zhu, D., Anholt, R.R.H., Ayroles, J.F., Duncan, L., Jordan,
K.W., Lawrence, F., Magwire, M.M., Warner, C.B., Blankenburg, K., Han, Y., Javaid, M.,
Jayaseelan, J., Jhangiani, S.N., Muzny, D., Ongeri, F., Perales, L., Wu, Y.-Q., Zhang, Y., Zou, X.,
Stone, E.A., Gibbs, R.A., Mackay, T.F.C., 2012. Epistasis dominates the genetic architecture of
Drosophila quantitative traits. Proceedings of the National Academy of Sciences 109, 15553–
15559. https://doi.org/10.1073/pnas.1213423109

Jeong, S., Rebeiz, M., Andolfatto, P., Werner, T., True, J., Carroll, S.B., 2008. The Evolution of Gene
Regulation Underlies a Morphological Difference between Two Drosophila Sister Species. Cell
132, 783–793. https://doi.org/10.1016/j.cell.2008.01.014

Kamiyama, M.T., Bradford, B.Z., Groves, R.L., Guédot, C., 2020. Degree day models to forecast the
seasonal phenology of Drosophila suzukii in tart cherry orchards in the Midwest U.S. PLOS ONE
15, e0227726. https://doi.org/10.1371/journal.pone.0227726

Kerner, B., North, K.E., Fallin, M.D., 2009. Use of Longitudinal Data in Genetic Studies in the Genome-
wide Association Studies Era: Summary of Group 14. Genet Epidemiol 33, S93–S98.
https://doi.org/10.1002/gepi.20479

Kessner, D., Turner, T.L., Novembre, J., 2013. Maximum Likelihood Estimation of Frequencies of Known Haplotypes from Pooled Sequence Data. Mol Biol Evol 30, 1145–1158. https://doi.org/10.1093/molbev/mst016

Kosheleva, K., Desai, M.M., 2018. Recombination Alters the Dynamics of Adaptation on Standing Variation in Laboratory Yeast Populations. Mol Biol Evol 35, 180–201. https://doi.org/10.1093/molbev/msx278

Kronforst, M.R., Barsh, G.S., Kopp, A., Mallet, J., Monteiro, A., Mullen, S.P., Protas, M., Rosenblum, E.B., Schneider, C.J., Hoekstra, H.E., 2012. Unraveling the thread of nature's tapestry: the genetics of diversity and convergence in animal pigmentation. Pigment Cell & Melanoma Research 25, 411–433. https://doi.org/10.1111/j.1755-148X.2012.01014.x

Kunte, K., Zhang, W., Tenger-Trolander, A., Palmer, D.H., Martin, A., Reed, R.D., Mullen, S.P., Kronforst, M.R., 2014. doublesex is a mimicry supergene. Nature 507, 229–232. https://doi.org/10.1038/nature13112

Lenski, R.E., 2017. Experimental evolution and the dynamics of adaptation and genome evolution in microbial populations. ISME J 11, 2181–2194. https://doi.org/10.1038/ismej.2017.69

Lohmueller, K.E., 2014. The Impact of Population Demography and Selection on the Genetic Architecture of Complex Traits. PLOS Genetics 10, e1004379. https://doi.org/10.1371/journal.pgen.1004379

Long, A.D., Hanson, K.M., Macdonald, S.J., 2025. The Illusion of Polygenicity in Poolseq studies: Insufficient Power Can Mask Simple Genetic Architectures. https://doi.org/10.1101/2025.07.23.666414

Mackay, T.F.C., Anholt, R.R.H., 2024. Pleiotropy, epistasis and the genetic architecture of quantitative traits. Nat Rev Genet 25, 639–657. https://doi.org/10.1038/s41576-024-00711-3

Mackay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., Casillas, S., Han, Y., Magwire, M.M., Cridland, J.M., Richardson, M.F., Anholt, R.R.H., Barrón, M., Bess, C., Blankenburg, K.P., Carbone, M.A., Castellano, D., Chaboub, L., Duncan, L., Harris, Z., Javaid, M., Jayaseelan, J.C., Jhangiani, S.N., Jordan, K.W., Lara, F., Lawrence, F., Lee, S.L., Librado, P., Linheiro, R.S., Lyman, R.F., Mackey, A.J., Munidasa, M., Muzny, D.M., Nazareth, L., Newsham, I., Perales, L., Pu, L.-L., Qu, C., Ràmia, M., Reid, J.G., Rollmann, S.M., Rozas, J., Saada, N., Turlapati, L., Worley, K.C., Wu, Y.-Q., Yamamoto, A., Zhu, Y., Bergman, C.M., Thornton, K.R., Mittelman, D., Gibbs, R.A., 2012. The Drosophila melanogaster Genetic Reference Panel. Nature 482, 173–178. https://doi.org/10.1038/nature10811

Martin, A.R., Gignoux, C.R., Walters, R.K., Wojcik, G.L., Neale, B.M., Gravel, S., Daly, M.J., Bustamante, C.D., Kenny, E.E., 2017. Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. The American Journal of Human Genetics 100, 635–649. https://doi.org/10.1016/j.ajhg.2017.03.004

Massey, J., Wittkopp, P.J., 2016. The genetic basis of pigmentation differences within and between Drosophila species. Curr Top Dev Biol 119, 27–61. https://doi.org/10.1016/bs.ctdb.2016.03.004

Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg, S.A., Harney, E., Stewardson, K., Fernandes, D., Novak, M., Sirak, K., Gamba, C., Jones, E.R., Llamas, B., Dryomov, S., Pickrell, J., Arsuaga, J.L., de Castro, J.M.B., Carbonell, E., Gerritsen, F., Khokhlov, A., Kuznetsov, P., Lozano, M., Meller, H., Mochalov, O., Moiseyev, V., Guerra, M.A.R., Roodenberg, J., Vergès, J.M., Krause, J., Cooper, A., Alt, K.W., Brown, D., Anthony, D., Lalueza-Fox, C., Haak, W., Pinhasi, R., Reich, D., 2015. Genome-wide patterns of selection in 230 ancient Eurasians. Nature 528, 499–503. https://doi.org/10.1038/nature16152

Mostafavi, H., Harpak, A., Agarwal, I., Conley, D., Pritchard, J.K., Przeworski, M., 2020. Variable prediction accuracy of polygenic scores within an ancestry group. eLife 9, e48376. https://doi.org/10.7554/eLife.48376

Nadeau, N.J., Pardo-Diaz, C., Whibley, A., Supple, M.A., Saenko, S.V., Wallbank, R.W.R., Wu, G.C., Maroja, L., Ferguson, L., Hanly, J.J., Hines, H., Salazar, C., Merrill, R.M., Dowling, A.J., ffrench-Constant, R.H., Llaurens, V., Joron, M., McMillan, W.O., Jiggins, C.D., 2016. The gene

993   cortex controls mimicry and crypsis in butterflies and moths. Nature 534, 106–110.
994   https://doi.org/10.1038/nature17961

995 Novembre, J., Barton, N.H., 2018. Tread Lightly Interpreting Polygenic Tests of Selection. Genetics 208,
996   1351–1355. https://doi.org/10.1534/genetics.118.300786

997 Paaby, A.B., Bergland, A.O., Behrman, E.L., Schmidt, P.S., 2014. A highly pleiotropic amino acid
998   polymorphism in the Drosophila insulin receptor contributes to life-history adaptation. Evol 68,
999   3395–3409. https://doi.org/10.1111/evo.12546

1000 Patel, R.A., Musharoff, S.A., Spence, J.P., Pimentel, H., Tcheandjieu, C., Mostafavi, H., Sinnott-
1001   Armstrong, N., Clarke, S.L., Smith, C.J., Durda, P.P., Taylor, K.D., Tracy, R., Liu, Y., Johnson,
1002   W.C., Aguet, F., Ardlie, K.G., Gabriel, S., Smith, J., Nickerson, D.A., Rich, S.S., Rotter, J.I.,
1003   Tsao, P.S., Assimes, T.L., Pritchard, J.K., 2022. Genetic interactions drive heterogeneity in causal
1004   variant effect sizes for gene expression and complex traits. The American Journal of Human
1005   Genetics 109, 1286–1297. https://doi.org/10.1016/j.ajhg.2022.05.014

1006 Phillips, P.C., 2008. Epistasis — the essential role of gene interactions in the structure and evolution of
1007   genetic systems. Nat Rev Genet 9, 855–867. https://doi.org/10.1038/nrg2452

1008 Pool, J.E., Aquadro, C.F., 2007. The genetic basis of adaptive pigmentation variation in Drosophila
1009   melanogaster. Molecular Ecology 16, 2844–2851. https://doi.org/10.1111/j.1365-
1010   294X.2007.03324.x

1011 Privé, F., Aschard, H., Carmi, S., Folkersen, L., Hoggart, C., O'Reilly, P.F., Vilhjálmsson, B.J., 2022.
1012   Portability of 245 polygenic scores when derived from the UK Biobank and applied to 9 ancestry
1013   groups from the same cohort. The American Journal of Human Genetics 109, 12–23.
1014   https://doi.org/10.1016/j.ajhg.2021.11.008

1015 Reed, R.D., Papa, R., Martin, A., Hines, H.M., Counterman, B.A., Pardo-Diaz, C., Jiggins, C.D.,
1016   Chamberlain, N.L., Kronforst, M.R., Chen, R., Halder, G., Nijhout, H.F., McMillan, W.O., 2011.
1017   optix drives the repeated convergent evolution of butterfly wing pattern mimicry. Science 333,
1018   1137–1141. https://doi.org/10.1126/science.1208227

1019 Rudman, S.M., Greenblum, S.I., Rajpurohit, S., Betancourt, N.J., Hanna, J., Tilk, S., Yokoyama, T.,
1020   Petrov, D.A., Schmidt, P., 2022. Direct observation of adaptive tracking on ecological time scales
1021   in Drosophila. Science 375, eabj7484. https://doi.org/10.1126/science.abj7484

1022 Sella, G., Barton, N.H., 2019. Thinking About the Evolution of Complex Traits in the Era of Genome-
1023   Wide Association Studies. Annual Review of Genomics and Human Genetics 20, 461–493.
1024   https://doi.org/10.1146/annurev-genom-083115-022316

1025 Stroud, J.T., Ratcliff, W.C., 2025. Long-term studies provide unique insights into evolution. Nature 639,
1026   589–601. https://doi.org/10.1038/s41586-025-08597-9

1027 Svensson, E.I., Arnold, S.J., Bürger, R., Csilléry, K., Draghi, J., Henshaw, J.M., Jones, A.G., De Lisle, S.,
1028   Marques, D.A., McGuigan, K., Simon, M.N., Runemark, A., 2021. Correlational selection in the
1029   age of genomics. Nat Ecol Evol 5, 562–573. https://doi.org/10.1038/s41559-021-01413-3

1030 Tilk, S., Bergland, A., Goodman, A., Schmidt, P., Petrov, D., Greenblum, S., 2019. Accurate Allele
1031   Frequencies from Ultra-low Coverage Pool-Seq Samples in Evolve-and-Resequence
1032   Experiments. G3 (Bethesda) 9, 4159–4168. https://doi.org/10.1534/g3.119.400755

1033 Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A., Yang, J., 2017. 10 Years
1034   of GWAS Discovery: Biology, Function, and Translation. Am J Hum Genet 101, 5–22.
1035   https://doi.org/10.1016/j.ajhg.2017.06.005

1036 Walsh, B., Blows, M.W., 2009. Abundant Genetic Variation + Strong Selection = Multivariate Genetic
1037   Constraints: A Geometric View of Adaptation. Annu. Rev. Ecol. Evol. Syst. 40, 41–59.
1038   https://doi.org/10.1146/annurev.ecolsys.110308.120232

1039 Wortel, M.T., Agashe, D., Bailey, S.F., Bank, C., Bisschop, K., Blankers, T., Cairns, J., Colizzi, E.S.,
1040   Cusseddu, D., Desai, M.M., van Dijk, B., Egas, M., Ellers, J., Groot, A.T., Heckel, D.G.,
1041   Johnson, M.L., Kraaijeveld, K., Krug, J., Laan, L., Lässig, M., Lind, P.A., Meijer, J., Noble,
1042   L.M., Okasha, S., Rainey, P.B., Rozen, D.E., Shitut, S., Tans, S.J., Tenaillon, O., Teotónio, H., de
1043   Visser, J.A.G.M., Visser, M.E., Vroomans, R.M.A., Werner, G.D.A., Wertheim, B., Pennings,

P.S., 2022. Towards evolutionary predictions: Current promises and challenges. Evol Appl 16, 3–
21. https://doi.org/10.1111/eva.13513
Wright, T.R.F., 1987. The Genetics Of Biogenic Amine Metabolism, Sclerotization, And Melanization In
Drosophila Melanogaster*, in: Scandalios, J.G., Caspari, E.W. (Eds.), Advances in Genetics,
Molecular Genetics of Development. Academic Press, pp. 127–222.
https://doi.org/10.1016/S0065-2660(08)60008-5
Wu, Y., Zheng, Z., Thibaut, L., Lin, T., Feng, Q., Cheng, H., Yengo, L., Goddard, M.E., Wray, N.R.,
Visscher, P.M., Zeng, J., 2025. Genome-wide fine-mapping improves identification of causal
variants. medRxiv 2024.07.18.24310667. https://doi.org/10.1101/2024.07.18.24310667
Wu, Z., Hu, Y., Melton, P.E., 2014. Longitudinal Data Analysis for Genetic Studies in the Whole-
Genome Sequencing Era. Genetic Epidemiology 38, S74–S80. https://doi.org/10.1002/gepi.21829
Yair, S., Coop, G., 2021. Population differentiation of polygenic score predictions under stabilizing
selection. https://doi.org/10.1101/2021.09.10.459833

Supplementary Figures

Supplementary Figure 1 | Relationship between color fraction and PC1. Samples plotted by color fraction (dark, midpoint, or light) and the coordinate along the first principal component generated via PCA of allele frequency data across all SNPs and samples used in mapping. A linear regression line (line in figure) between trait value and PC1 is significant ($R^2 = 0.762$; $p \ll 0.001$).

Supplementary Figure 2 | PCA by chromosome. Projection of samples used in trait mapping onto the first two principal components generated via PCA of allele frequency data from all SNPs located on each chromosomal arm. In the column on the left, samples are colored by the color fraction from which they originate (dark, midpoint, or light) and the shape of the points corresponds to the timepoint when the sample was collected. In the right column, samples are colored by treatment.

Supplementary Figure 3 | PCA without variation within inversions. Projection of samples used in trait mapping onto the first two principal components generated via PCA of allele frequency data from all SNPs except those located in each of six common inversions found in wild populations. In the column on the left, samples are colored by the color fraction from which they originate (dark, midpoint, or light) and the shape of the points corresponds to the timepoint when the sample was collected. In the right column, samples are colored by treatment.

Supplementary Figure 4 | Distribution of pigmentation-associated SNPs across inbred reference panel. The proportion of all pigmentation-associated, significant SNPs (minor allele, identified via our ancestry-informed GLM conducted on the week 8, outdoor mapping population) present on each of the 76 founding inbred lines. Inbred lines are ordered from lightest to darkest pigmentation, as quantified via manual scoring. As can be observed, those SNPs identified by our ancestry-informed GLM are not exclusive to either the lightest or darkest lines, but rather present at appreciable frequency across all haplotypes.

Supplementary Figure 5 | GLM empirical and permuted p-value cumulative distribution functions. CDFs of log10 p-values of SNPs resulting from a GLM performed with real data (red) and 100 GLMs performed with shuffled, permuted data (grey). The cutoff, determined as the top 0.05% of permuted p-values, is drawn as a blue dotted line for each timepoint and treatment.

Supplementary Figure 6 | Distribution of pigmentation-associated SNPs across inbred reference panel for the largest-effect, independent loci identified on each chromosomal arm. The proportion of all pigmentation-associated SNPs within each of the top independent loci (clusters) on each chromosomal arm (minor allele, identified via our ancestry-informed GLM conducted on the week 8, outdoor mapping population) present on each of the 76 founding inbred lines. Inbred lines are ordered from lightest to darkest pigmentation score as quantified via manual scoring.

Supplementary Figure 7 | Seasonal shifts of trait-associated SNPs in outdoor cages. A) Distribution of light-associated allele frequency shifts at all trait-associated SNPs (identified at Week 8 in outdoor cages) between Weeks 1 and 8 of evolution in the outdoor field environment. Distributions of frequency shifts at trait-associated SNPs are dark magenta, while distributions derived from matched controls are grey. Reported p-values are derived from a comparison of trait-associated and matched control SNPs using a binomial test. B) Same as (A), but segregated by chromosomal arm.

Supplementary Figure 8 | Seasonal shifts of trait-associated SNPs in indoor cages. A) Distribution of light-associated allele frequency shifts at all trait-associated SNPs (identified at Week 8 in outdoor cages) between Weeks 1 and 8 of evolution in the indoor lab environment. Distributions of frequency shifts at trait-associated SNPs are turquoise, while distributions derived from matched controls are grey. Reported p-values are derived from a comparison of trait-associated and matched control SNPs using a binomial test. B) Same as (A), but segregated by chromosomal arm.

Supplementary Figure 9 | Seasonal shifts of trait-associated SNPs in outdoor versus indoor cages. A) Distribution of light-associated allele frequency shifts at all trait-associated SNPs (identified at Week 8 in outdoor cages) between Weeks 1 and 8 of evolution in the outdoor field vs. indoor lab environment across the whole genome. Distributions of frequency shifts at trait-associated SNPs in the outdoor cages are magenta, while distributions in indoor cages are turquoise. The indoor and outdoor distributions were compared using a two-sided t-test. B) Same as (A), but segregated by chromosomal arm.

Supplementary Figure 10 | Phenotypic prediction with only *tan*, *ebony*, **and** *bab*, **using mapping from each timepoint/treatment subset.** Relative, predicted phenotypic change between Weeks 1 and 8 in both indoor and outdoor environments, based on only the three clusters that contain *tan, ebony*, or *bab*. Each quadrant uses a different mapping (for example, from Week 8 in the outdoor cages) and within each quadrant, we use the mapping to predict the phenotypic change in the outdoor environment (magenta) and the indoor environment (turquoise). The predicted change is denoted with diamonds, while the violins indicate the distribution of 1000 bootstrapped values (colored for trait-associated SNPs and grey to matched control SNPs).

Supplementary Figure 11 | Phenotypic prediction excluding *tan*, *ebony*, **and** *bab*. Relative, predicted phenotypic change between Weeks 1 and 8 in both outdoor (magenta) and indoor (turquoise) environments, generated using our trait mapping results derived at weeks 8 in the outdoor (left) and indoor environment, and using variation from every mapped locus excluding those containing *tan, ebony*, or *bab*. Predicted phenotypic change are quantified as Z-scores (computed relative to matched control shifts) whereby positive Z-scores indicate net lighter pigmentation evolution while negative values would indicate net darker evolution. Z-scores were computed by comparing inferred relative phenotypic change using mapped SNPs to a that computed using a set of matched control SNPs (grey distributions).

Supplementary Figure 12 | Manhattan plots produced across all mapping populations. Manhattan plot depicting nominal p-value's derived from an ancestry-informed mapping of SNP association with pigmentation variation, separately for each mapping population (week 2 indoor, week 2 outdoor, week 8 indoor, week 8 outdoor). The dashed horizontal line corresponds to an empirical FDR threshold of 0.0005, derived via permutations. Shaded regions denote the stop and end points of each of the independent loci, while bar color denotes behavior of SNPs within each locus through time as the populations evolved lighter pigmentation within the outdoor mesocosm: red shading indicates loci with light-associated alleles favored throughout evolution of lighter pigmentation, blue shading indicates loci in which the light-associated alleles are predominately selected against and grey shading indicates loci with no signature of selection on light associated alleles. The location of the canonical pigmentation genes, *bric-a-brac (bab)*, *ebony*, and *tan*, are indicated on the Manhattan plots. Potential candidate genes underlying the remaining large effect loci observed throughout the genome are discussed in the Supplement.

Supplementary Figure 13 | Shared significant SNPs across mapping populations. Overlap percentage from each pair of GLMs that each use five of the cages from a specific timepoint and treatment combination. Significant SNP overlap is either considered between GLMs using cages within the same timepoint/treatment (green), the same treatment but the other timepoint (yellow), the same timepoint but the other treatment (red), or cages from both the other timepoint and treatment (blue). The box plots represent the real percentage from each paired comparison, while the colored violins represent our bootstrapping results.

Supplementary Figure 14 | Leave-one-out across the full genome, for all timepoint/treatment combinations. Results from leave-one-out validation of mapping within and across treatments and time points for all timepoint/treatment combinations. Depicted are the median allele frequency difference for each test cage between light and dark samples of trait-associated SNPs (red), as well as a set of matched control SNPs (grey).

Supplementary Figure 15 | Evidence of field-specific pigmentation-associated loci. Distribution of light-dark allele frequencies (boxes denote the 50% interquartile range, dark horizontal lines correspond to the median, and points indicate outliers) for pigmentation associated SNPs located within mapped loci occurring the outdoor, week 8 mapping population that share no overlapping breakpoints with any loci in the indoor, week 8 mapping population. The magenta distribution corresponds to allele frequency differences within the outdoor cages, while the turquoise to the indoor environment. Grey distributions then represent the light-dark allele frequency differences at matched control SNPs for each environment. The light-dark frequency differences are significantly greater than the matched control shifts for the outdoor environment (Mann-Whitney U test p-value = 0.003), and significantly less than matched control SNPs for the indoor environment (Mann-Whitney U test p-value < 0.001). Furthermore, the outdoor and indoor frequency differences are significantly different than each other (Wilcoxon Signed Rank Test p-value < 0.001), indicating that light-associated alleles in outdoor environment are putatively enriched in the dark color fraction of the indoor enviornment for this set of loci.

Supplementary Figure 16 | Evidence of lab-specific pigmentation-associated loci. Distribution of light-dark allele frequencies (boxes denote the 50% interquartile range, dark horizontal lines correspond to the median, and points indicate outliers) for pigmentation associated SNPs located within mapped loci occurring the indoor, week 8 mapping population that share no overlapping breakpoints with any loci in the outdoor, week 8 mapping population. The magenta distribution corresponds to allele frequency differences within the outdoor cages, while the turquoise to the indoor environment. Grey distributions then represent the light-dark allele frequency differences at matched control SNPs for each environment. The light-dark frequency differences are significantly greater than the matched control shifts for the indoor environment (Mann-Whitney U test p-value < 0.001), and significantly less than matched control SNPs for the outdoor environment (Mann-Whitney U test p-value < 0.001). Furthermore, the indoor and outdoor frequency differences are significantly different than each other (Wilcoxon Signed Rank Test p-value < 0.001), indicating that light-associated alleles in indoor environment are putatively enriched in the dark color fraction of the outdoor enviornment for this set of loci.

Supplementary Figure 17 | Phenotypic prediction excluding treatment-specific loci. Relative, predicted phenotypic change between Weeks 1 and 8 in both outdoor (magenta) and indoor (turquoise) environments, generated using our trait mapping results derived at weeks 8 in the outdoor (left) and indoor environment, and using variation from every mapped locus excluding those that are identified only in the outdoor cages at week 8 or identified only in the indoor cages at week 8. Predicted phenotypic change are quantified as Z-scores (computed relative to matched control shifts) whereby positive Z-scores indicate net lighter pigmentation evolution while negative values would indicate net darker evolution. Z-scores were computed by comparing inferred relative phenotypic change using mapped SNPs to a that computed using a set of matched control SNPs (grey distributions).

Supplementary Figure 18 | Phenotypic predictions of color samples across timepoint and treatment.
Using the mapping from each timepoint/treatment subset and considering all significant SNPs, we generate a phenotypic prediction score for each color fraction sample. A negative value indicates an expected phenotypic shift towards dark pigmentation and a positive value a shift towards lighter pigmentation. We consider our prediction "correct" when the majority of dark samples (dark brown) have a negative predicted value, the majority of light samples (gold) have a positive one, and a linear regression (where we encode dark = 0, midpoint = 1, and light = 2) is significant and has a positive slope. We predict the phenotype of all samples from either within the same timepoint/treatment (green), the same treatment but the other timepoint (yellow), the same timepoint but the other treatment (red), or from both the other timepoint and treatment (blue).

Supplementary Figure 19 | Percentage of top SNPs eliminated in first round of unlinked locus identification with different quantile cutoffs. In our clustering method, the SNPs with the highest light-dark difference as predicted by haplotype frequencies in the highest signal region are eliminated sequentially. Here, we look at the percentage of the eliminated SNPs that are significant SNPs in the first round of clustering with cutoffs at different quantiles using real data versus permutations.

Supplementary Figure 20 | Signal elimination throughout clustering. Plot of the number of remaining significant SNPs before each round of clustering for each chromosome, showing the elimination of linked significant SNPs throughout the clustering process.

Supplementary Tables

Table S1 | List of pigmentation (Pig) and cuticle-associated genes located within mapped loci identified in the week 8, outdoor mapping population that are enriched in pigmentation-associated, mapped SNPs (binomial test, FDR < 0.05). Also reported is the locus ID (corresponding to the locus information provided in Supplementary Data File 1) within which each gene resides, the distance of each gene to the locus midpoint SNP, the number of SNPs within or within 10 Kb of the gene, the proportion of SNPs within the gene that are pigmentation associated (Sig. SNP proportion), and the Benjamini-Yekutieli false discovery rate corrected p-value for observed enrichment of pigmentation-associated SNPs.

Locus ID	FBID	Gene Name	Pig GO?	Cuticle GO?	Distance to midpoint SNP (Mb)	Total SNPs in gene ± 10Kb	Sig. SNP Proportion	FDR
TP8.E.2L.3	FBgn0003255	rickets		✓	0.50	956	0.0251	0.0019
TP8.E.2R.1	FBgn0002533	Larval cuticle protein 2		✓	0.12	330	0.0636	3.13×10^{-9}
	FBgn0002534	Larval cuticle protein 3		✓	0.12	322	0.0621	1.27×10^{-8}
	FBgn0002535	Larval cuticle protein 4		✓	0.12	315	0.0603	5.15×10^{-8}
	FBgn0002531	Larval cuticle protein 1		✓	0.13	344	0.0436	1.10×10^{-4}
	FBgn0002567	lightoid	✓		0.62	422	0.0284	0.0383
TP8.E.3L.1	FBgn0004870	bric a brac 1	✓		0.55	1951	0.1630	2.80×10^{-265}
	FBgn0035281	Cuticular protein 62Bc		✓	1.36	601	0.0499	1.44×10^{-10}
	FBgn0035280	Cuticular protein 62Bb		✓	1.35	577	0.0433	1.20×10^{-7}
	FBgn0035279	Cuticular protein 62Ba		✓	1.34	634	0.0426	4.36×10^{-8}
TP8.E.3L.2	FBgn0035398	Cht7		✓	0.43	772	0.0311	6.90×10^{-5}
TP8.E.3L.5	FBgn0035512	Cuticular protein 64Ac		✓	0.10	511	0.1683	1.66×10^{-72}
	FBgn0035511	Cuticular protein 64Ab		✓	0.10	508	0.1673	1.64×10^{-71}
	FBgn0035513	Cuticular protein 64Ad		✓	0.10	478	0.1611	1.62×10^{-63}
	FBgn0035510	Cuticular protein 64Aa		✓	0.10	551	0.1506	5.41×10^{-66}
TP8.E.3R.2	FBgn0000527	ebony	✓	✓	0.00	625	0.1296	3.55×10^{-59}
TP8.E.3R.3	FBgn0038714	Cuticular protein 92A		✓	0.53	578	0.0329	3.36×10^{-4}
TP8.E.X.1	FBgn0086367	tan	✓		0.01	382	0.1545	1.74×10^{-47}

Table S2 | List of pigmentation (Pig) and cuticle-associated genes located within mapped loci identified in the week 8, indoor mapping population that are enriched in pigmentation-associated, mapped SNPs (binomial test, FDR < 0.05). Also reported is the locus ID (corresponding to the locus information provided in Supplementary Data File 1) within which each gene resides, the distance of each gene to the locus midpoint SNP, the number of SNPs within or within 10 Kb of the gene, the proportion of SNPs within the gene that are pigmentation associated (Sig. SNP proportion), and the Benjamini-Yekutieli false discovery rate corrected p-value for observed enrichment of pigmentation-associated SNPs.

Locus ID	FBID	Gene Name	Pig GO?	Cuticle GO?	Distance to midpoint SNP (Mb)	Total SNPs in gene ± 10Kb	Sig. SNP Proportion	FDR
TP8.I.2L.3	FBgn0053302	Cuticular protein 31A		✓	0.03	642	0.0685	1.05×10^{-14}
TP8.I.2R.1	FBgn0002534	Larval cuticle protein 3		✓	0.43	322	0.1770	1.58×10^{-40}
	FBgn0002535	Larval cuticle protein 4		✓	0.44	315	0.1619	3.09×10^{-34}
	FBgn0002533	Larval cuticle protein 2		✓	0.43	330	0.1485	4.15×10^{-31}
	FBgn0002531	Larval cuticle protein 1		✓	0.43	344	0.1279	3.87×10^{-25}
TP8.I.3L.1	FBgn0000543	ecdysoneless		✓	0.69	535	0.0897	1.08×10^{-20}
	FBgn0035279	Cuticular protein 62Ba		✓	0.25	634	0.0757	1.08×10^{-17}
	FBgn0035280	Cuticular protein 62Bb		✓	0.26	577	0.0676	7.67×10^{-13}
	FBgn0022702	Chitinase 2		✓	0.18	350	0.0657	2.17×10^{-7}
	FBgn0035281	Cuticular protein 62Bc		✓	0.27	601	0.0416	1.74×10^{-4}
TP8.I.3L.2	FBgn0035875	Cuticular protein 66Cb		✓	0.30	429	0.0559	1.82×10^{-6}
TP8.I.3L.5	FBgn0004870	bric a brac 1	✓		0.33	1951	0.1978	1.44×10^{-295}
TP8.I.3L.6	FBgn0261797	Dynein heavy chain 64C		✓	0.23	782	0.0665	7.88×10^{-17}
TP8.I.3R.2	FBgn0027508	tankyrase		✓	0.34	537	0.0503	2.38×10^{-6}
TP8.I.3R.3	FBgn0000527	ebony	✓	✓	0.07	625	0.0432	3.97×10^{-5}
	FBgn0038819	Cuticular protein 92F		✓	0.49	464	0.0409	0.0024
TP8.I.3R.6	FBgn0038901	Bursicon	✓	✓	0.36	325	0.0554	8.00×10^{-5}
TP8.I.X.1	FBgn0086367	tan	✓		0.01	382	0.1440	3.10×10^{-34}

Supplementary Discussion

Novel pigmentation-associated loci identified by high-powered, tail-based mapping

Here we describe in further detail genes occurring within our mapped loci that likely play a functional role in pigmentation variation. As nearly 80% of total genes in the genome were located within the start and endpoints of the loci identified across all mapping populations, we concentrate this discussion on those genes with a gene ontology (GO) term associated with pigmentation (excluding eye-specific pigmentation GO terms) and/or with cuticle development and molting, as well as confine our assessment to mapping conducted at week 8. Additionally, for each putative candidate gene, we ensured that the proportion of mapping significant SNPs within or nearby (within 10 Kb) gene breakpoints were significantly greater than expected based on the genome-wide distribution of significant SNPs (binomial test).

As expected, *bric-a-brac* (*bab*), *ebony*, and *tan* each fell within mapped loci and contained a significant enrichment of pigmentation-associated SNPs. *Ebony* and *tan* are opposing enzymes that catalyze dopamine to a precursor of sclerotin (N-β-alanyl-dopamine, NBAD) and NBAD back to dopamine, respectively (Wright, 1987). Thus, together they control the amount of NBAD available to create light pigmented sclerotin instead of darkly pigmented melanin. In contrast, *bab* is a set of two duplicated transcription factors (*bab1* and *bab2*) that repress pigmentation (Couderc et al., 2002). For example, *bab* is expressed in tergites A2 through A6 in female *D. melanogaster,* but only A2 through A4 in males, giving rise to the characteristic, male-only, completely pigmented terminal tergites for which *D. melanogaster* is named (Kopp et al., 2000). Binding sites for *Abdominal-B* (*Abd-B*) and *doublesex* (dsx) are located in the long intron of bab1, and the binding of *doublesex* to *bab1* mediates this sex-specific pigmentation (Williams et al., 2008). Additionally, previous studies have shown that the regulation of both *bab* and *tan* is influenced by temperature, providing a likely explanation for the plasticity of pigmentation across developmental temperatures observed in *D. melanogaster (Castro et al., 2018; Gibert et al., 2016).* For *bab*, the *hox* gene and transcription factor Abd-B has been shown to mediate this temperature-dependent expression (Gibert et al., 2007). We also identify Abd-B in our study which is located within the top locus on chromosomal arm 3R.

On 2R, our largest peak (which shows high consistency across timepoints and treatments), overlies larval cuticle proteins 1, 2, and 3 (*Lcp1*, *Lcp2*, *Lcp3*) – three out of four larval cuticle protein genes that form a set of two pairs of genes (*Lcp1* and *Lcp2*, and *Lcp3* and *Lcp4*) that each act as a separate transcriptional unit and are transcribed at different periods during the third instar stage (Kimbrell et al., 1988). *Lcp3* and *Lcp4* accumulate first, followed by *Lcp1* and *Lcp2*, and this is likely part of a set of changes that shift the third instar soft cuticle to a harder, less flexible one for the wandering stage (Kimbrell et al., 1988). Even though the larval cuticle is not preserved into the adult stage, certain pigmentation genes (namely yellow) that are important for adult pigmentation are also expressed and influence pigmentation in the larval stages (Kornezos and Chia, 1992).

We find further evidence of pigmentation-associated SNPs in loci concentrated in and around *Bursicon* (*Burs*), *rickets* (*rk*), and *Partner of bursicon* (*Pburs*). These three genes are the core genes vital to the cuticular hardening after eclosion. *Burs* and *Pburs* dimerize and bind to rickets to initialize the final tanning or hardening of the cuticle (Baker and Truman, 2002; Dewey et al., 2004). Prior to this, the eclosed fly is entirely white, with very little visible pigmentation.

Tanning then occurs when deposited cuticular proteins are linked together and the pigmentation becomes stark and visible. Despite this direct connection to pigmentation of the cuticle, this mapping study represents the first to directly identify pigmentation-relevant variation in these genes.

In only the outdoor week 8 mapping population, we find more significant SNPs than expected around and within a gene on chromosomal arm 3L that encodes for a transmembrane chitinase, *Chitinase 7* (*Cht7*) (Zhu et al., 2008). It appears that *Cht7* is primarily involved in chitin organization in the developing cuticle, as RNAi knockdown of *Cht7* has shown the old cuticle to be disassembled successfully, but the density and organization of the new cuticle is compromised (Noh et al., 2018). Another chitinase gene, *Chitinase 2* (*Cht2*), is nearby *Cht7* on 3L, but only exhibited a significant enrichment of significant SNPs in the indoor mapping population. *Cht2*, like *Cht7*, seems to play a role in organizing the developing cuticle and not in disassembling the old cuticle (Pesch et al., 2017).

In the indoor week 8 mapping population, we identified loci with high signal in and around *tankyrase*, *Dynein heavy chain 64c* (*Dhc64c*), and *ecdysoneless*. Both *tankyrase* and *Dhc64c* are involved in the proper expression and localization of *Wg* (Feng et al., 2014; Kim et al., 2021). *Wg* and other components in, or affected by, the *Wnt* pathway are known to be associated with pigmentation differences within and between species (Massey and Wittkopp, 2016). *Ecdysoneless*, while it does affect the production of ecdysone as its name implies, actually has a broader role in correctly localizing pre-mRNA processing factor 8 (*Prp8*), a highly conserved splicing factor (Claudius et al., 2014; Erkelenz et al., 2021).

Supplementary References:

Baker, J.D., Truman, J.W., 2002. Mutations in the Drosophila glycoprotein hormone receptor, rickets, eliminate neuropeptide-induced tanning and selectively block a stereotyped behavioral program. J Exp Biol 205, 2555–2565. https://doi.org/10.1242/jeb.205.17.2555

Castro, S.D., Peronnet, F., Gilles, J.-F., Mouchel-Vielh, E., Gibert, J.-M., 2018. bric à brac (bab), a central player in the gene regulatory network that mediates thermal plasticity of pigmentation in Drosophila melanogaster. PLOS Genetics 14, e1007573. https://doi.org/10.1371/journal.pgen.1007573

Claudius, A.-K., Romani, P., Lamkemeyer, T., Jindra, M., Uhlirova, M., 2014. Unexpected Role of the Steroid-Deficiency Protein Ecdysoneless in Pre-mRNA Splicing. PLOS Genetics 10, e1004287. https://doi.org/10.1371/journal.pgen.1004287

Couderc, J.-L., Godt, D., Zollman, S., Chen, J., Li, M., Tiong, S., Cramton, S.E., Sahut-Barnola, I., Laski, F.A., 2002. The bric à brac locus consists of two paralogous genes encoding BTB/POZ domain proteins and acts as a homeotic and morphogenetic regulator of imaginal development in Drosophila. Development 129, 2419–2433. https://doi.org/10.1242/dev.129.10.2419

Dewey, E.M., McNabb, S.L., Ewer, J., Kuo, G.R., Takanishi, C.L., Truman, J.W., Honegger, H.-W., 2004. Identification of the Gene Encoding Bursicon, an Insect Neuropeptide Responsible for

Cuticle Sclerotization and Wing Spreading. Current Biology 14, 1208–1213. https://doi.org/10.1016/j.cub.2004.06.051

Erkelenz, S., Stanković, D., Mundorf, J., Bresser, T., Claudius, A.-K., Boehm, V., Gehring, N.H., Uhlirova, M., 2021. Ecd promotes U5 snRNP maturation and Prp8 stability. Nucleic Acids Res 49, 1688–1707. https://doi.org/10.1093/nar/gkaa1274

Feng, Y., Li, X., Ray, L., Song, H., Qu, J., Lin, S., Lin, X., 2014. The Drosophila tankyrase regulates Wg signaling depending on the concentration of Daxin. Cell Signal 26, 1717–1724. https://doi.org/10.1016/j.cellsig.2014.04.014

Gibert, J.-M., Mouchel-Vielh, E., Castro, S.D., Peronnet, F., 2016. Phenotypic Plasticity through Transcriptional Regulation of the Evolutionary Hotspot Gene tan in Drosophila melanogaster. PLOS Genetics 12, e1006218. https://doi.org/10.1371/journal.pgen.1006218

Gibert, J.-M., Peronnet, F., Schlötterer, C., 2007. Phenotypic Plasticity in Drosophila Pigmentation Caused by Temperature Sensitivity of a Chromatin Regulator Network. PLOS Genetics 3, e30. https://doi.org/10.1371/journal.pgen.0030030

Kim, J.-Y., Tsogtbaatar, O., Cho, K.-O., 2021. Dynein Heavy Chain 64C Differentially Regulates Cell Survival and Proliferation of Wingless-Producing Cells in Drosophila melanogaster. Journal of Developmental Biology 9, 43. https://doi.org/10.3390/jdb9040043

Kimbrell, D.A., Berger, E., King, D.S., Wolfgang, W.J., Fristrom, J.W., 1988. Cuticle protein gene expression during the third instar of *Drosophila melanogaster*. Insect Biochemistry 18, 229–235. https://doi.org/10.1016/0020-1790(88)90087-X

Kopp, A., Duncan, I., Carroll, S.B., 2000. Genetic control and evolution of sexually dimorphic characters in Drosophila. Nature 408, 553–559. https://doi.org/10.1038/35046017

Kornezos, A., Chia, W., 1992. Apical secretion and association of the Drosophila yellow gene product with developing larval cuticle structures during embryogenesis. Molec. Gen. Genet. 235, 397–405. https://doi.org/10.1007/BF00279386

Massey, J., Wittkopp, P.J., 2016. The genetic basis of pigmentation differences within and between Drosophila species. Curr Top Dev Biol 119, 27–61. https://doi.org/10.1016/bs.ctdb.2016.03.004

Noh, M.Y., Muthukrishnan, S., Kramer, K.J., Arakane, Y., 2018. A chitinase with two catalytic domains is required for organization of the cuticular extracellular matrix of a beetle. PLOS Genetics 14, e1007307. https://doi.org/10.1371/journal.pgen.1007307

Pesch, Y.-Y., Riedel, D., Behr, M., 2017. *Drosophila* Chitinase 2 is expressed in chitin producing organs for cuticle formation. Arthropod Structure & Development, Arthropod Cuticle Function: from Molecules to Structures to Ecology 46, 4–12. https://doi.org/10.1016/j.asd.2016.11.002

Williams, T.M., Selegue, J.E., Werner, T., Gompel, N., Kopp, A., Carroll, S.B., 2008. The Regulation and Evolution of a Genetic Switch Controlling Sexually Dimorphic Traits in Drosophila. Cell 134, 610–623. https://doi.org/10.1016/j.cell.2008.06.052

Wright, T.R.F., 1987. The Genetics Of Biogenic Amine Metabolism, Sclerotization, And Melanization In *Drosophila Melanogaster**, in: Scandalios, J.G., Caspari, E.W. (Eds.), Advances in Genetics, Molecular Genetics of Development. Academic Press, pp. 127–222. https://doi.org/10.1016/S0065-2660(08)60008-5

Zhu, Q., Arakane, Y., Banerjee, D., Beeman, R.W., Kramer, K.J., Muthukrishnan, S., 2008. Domain organization and phylogenetic analysis of the chitinase-like family of proteins in three species of insects. Insect Biochemistry and Molecular Biology, Special Issue on the Tribolium Genome 38, 452–466. https://doi.org/10.1016/j.ibmb.2007.06.010